

Surrogate R-spondin agonists potentiate Wnt signaling

Surrogate R-spondin agonists for tissue-specific potentiation of Wnt signaling

Vincent C. Luca^{1,3,+,*}, Yi Miao^{1,+}, Xingnan Li², Michael J. Hollander⁴, Calvin J. Kuo², K. Christopher Garcia^{1,*}

¹Department of Molecular and Cellular Physiology, Howard Hughes Medical Institute, and Department of Structural Biology, Stanford University School of Medicine, Stanford, California 94305, USA;

²Department of Medicine, Division of Hematology, Stanford University School of Medicine, Stanford, California 94305, USA; ³Present Address: Department of Drug Discovery, Moffitt Cancer Center & Research Institute, Tampa, Florida, 33612, USA; ⁴Department of Bioengineering, Stanford University, Stanford, California 94305, USA

Running title: *Surrogate R-spondin agonists potentiate Wnt signaling*

⁺These authors contributed equally

*To whom correspondence should be addressed: Vincent C. Luca (Vince.Luca@moffitt.org) and K. Christopher Garcia (kcgarcia@stanford.edu)

Keywords: Wnt signaling, stem cells, regeneration, receptor endocytosis, protein engineering, R-spondin

ABSTRACT

Secreted R-spondin1-4 proteins (RSPO1-4) orchestrate stem cell renewal and tissue homeostasis by potentiating canonical Wnt/ β -catenin signaling. RSPOs induce the turnover of negative Wnt regulators RNF43 and ZNRF3 through a process that requires RSPO interactions with either Leucine-rich repeat-containing G-protein coupled receptors (LGRs) or heparin sulfate proteoglycans (HSPGs). Here, we describe the engineering of ‘surrogate RSPOs’ that function independently of LGRs and HSPGs to enhance Wnt signaling on cell types expressing a target surface marker. These bispecific proteins were generated by fusing an RNF43- or ZNRF3-specific single chain antibody variable fragment (scFv) to the immune cytokine IL-2. Surrogate RSPOs mimic the function of natural RSPOs by crosslinking the extracellular domain (ECD) of RNF43 or ZNRF3 to the ECD of the IL-2 receptor CD25, which sequesters the complex and results in highly selective amplification of Wnt signaling on CD25+ cells. Furthermore, surrogate RSPOs were able to substitute for wild-type RSPO in a colon organoid growth assay when intestinal stem cells were transduced to express CD25. Our results provide proof-of-concept for a technology that may be adapted for use on a broad range of cell- or tissue-types and will open new avenues

for the development of Wnt-based therapeutics for regenerative medicine.

The Wnt/ β -catenin pathway controls cell fate determination and tissue homeostasis in all metazoans (1). Pleiotropic Wnt signaling exhibits differential effects on a wide array of cell types and is therefore tightly regulated by several host-encoded enhancers and inhibitors. R-spondin proteins (RSPO1-4 in mammals) potentiate Wnt signaling by antagonizing negative regulators of the Wnt receptor Frizzled (2). The effect of RSPO is remarkably potent, and co-administration of Wnts and RSPOs can result in signaling outputs that are several hundred-fold greater than those of Wnt alone (3).

RSPO-mediated enhancement occurs via an indirect mechanism that greatly increases expression levels of the Wnt receptors Frizzled (Fzd) and LRP5 or LRP6 on the cell surface (2). In the absence of RSPO, the transmembrane E3 ligases RNF43 and ZNRF3 (4) ubiquitinate the intracellular regions of Fzd, which results in the internalization and degradation of both Fzd and associated LRP5/6 proteins (2) (Fig 1A). RSPO proteins are comprised of two Furin-like domains (Fu1, Fu2) followed by a thrombospondin domain (TSP) and basic region (BR). The Fu1

Surrogate R-spondin agonists potentiate Wnt signaling

domain of RSPO binds to the extracellular domains (ECDs) of RNF43 or ZNRF3 while the Fu2 domain binds to the ECD of a co-receptor, the Leucine-rich repeat-containing G-protein coupled receptor 4, 5 or 6 (LGR4-6) (5-8). This crosslinking event induces endocytosis of the RSPO-LGR-E3 ligase ternary complex, which sequesters RNF43 or ZNRF3 from Fzd and, in turn, potentiates Wnt signaling (Fig 1B). It was recently determined that RSPO2 and RSPO3 can function independently of LGRs by binding to heparan sulfate proteoglycans (HSPGs) on the cell surface via their TSP and BR domains (9). These data suggest RSPO are modular molecules consisting of an RNF43/ZNRF3 binding domain that is necessary for the potentiation of Wnt signaling and LGR/HSPG-binding domains that facilitate internalization of the complex.

Biomedical applications of recombinant RSPOs have been widely explored in the field of regenerative medicine, especially in the context of the small intestine where the RSPOs function to promote the renewal of stem cells at the base of the crypt (10,11). For example, recombinant RSPO1 protects animals from both experimentally induced colitis and from lethal doses of chemoradiation by stimulating intestinal regeneration (12). Furthermore, the addition of exogenous RSPO is critical for the proliferation of intestinal organoids (10), an in vitro model system for intestinal epithelium studies. Interestingly, it has been recently reported that Wnts and RSPOs play non-equivalent roles in intestinal stem cell (ISC) self-renewal through a feedback loop, in which Wnts drive the expression of LGR5 and RSPOs to induce ISC expansion (13), raising the possibility that targeting the pathways individually or in concert may be associated with unique therapeutic outcomes.

Potential obstacles for the clinical use of Wnt agonists include toxicity, off-target effects, and diminished potency caused by RNF43/ZNRF3-mediated antagonism. Wnts are powerful morphogens and hyperactivation of the Wnt pathway has been implicated in the development of several human cancers (14). RSPOs are promising candidates to selectively boost innate Wnt activity while circumventing off-target effects associated with global Wnt agonism. However, our ability to target RSPOs to specific

tissue or cell types is restricted by the expression profiles of RSPO receptors, since HSPGs are ubiquitously expressed and LGR expression is restricted to subsets of tissue stem cells. To develop precisely targeted RSPO mimetics, we engineered synthetic protein ligands that phenocopy RSPO-mediated Wnt signal enhancement via an LGR- and HSPG-independent mechanism. These “surrogate RSPOs” function by cross-linking the RNF43 or ZNRF3 ECD to an immune cell surface marker, CD25, that is known to undergo internalization upon ligand binding (15), thereby mimicking RSPO-mediated sequestration of RNF43/ZNRF3 (Fig. 1C). Surrogate RSPOs selectively boost Wnt signaling in CD25+ reporter cells and human colon organoids, and their modular design allows them to be adapted to target virtually any cell type, a feature that has major implications for the development of Wnt-based therapeutics.

RESULTS

Isolation of RNF43- and ZNRF3-specific scFvs using yeast surface display.

Our surrogate RSPO design concept involves the fusion of two components: (i) an RNF43- or ZNRF3-binding protein and (ii) a tissue-specific targeting protein that induces endocytosis of its receptor. Administration of the surrogate RSPO protein will cross-link ECD of RNF43 or ZNRF3 to the module's cognate receptor and mimic the function of natural RSPOs by driving membrane clearance of the complex. We isolated the RNF43- or ZNRF3-binding protein components for the surrogate RSPOs by selecting single-chain variable fragments (scFvs) from a published yeast display library of $\sim 1 \times 10^9$ sequences derived from human B-cells (16). The scFvs were obtained by performing two parallel sets of selections against either the RNF43 ECD or the ZNRF3 ECD using magnetic-activated cell sorting (MACS) and fluorescence-activated cell sorting (FACS). After several rounds of selection, we isolated a panel of different RNF43- or ZNRF3-specific scFvs (Fig. S1), and the clones “R5” and “Z6” were chosen for incorporation into surrogate RSPOs based on their ability to robustly bind to RNF43 and ZNRF3, respectively, in a fluorescence-based assay (Fig. 2A).

Surrogate R-spondin agonists potentiate Wnt signaling

Design, expression and purification of surrogate RSPOs.

Membrane clearance of RSPO signaling complexes has been reported to require the internalization of the receptors LGR4-6 (17). We therefore selected the immune cytokine IL-2 to act as a proxy for RSPO-LGR binding in our surrogate RSPO constructs based on its ability to induce endocytosis of its cognate receptor, CD25 (15). We generated two surrogate RSPO constructs, one targeted to RNF43 and one targeted to ZNRF3, by fusing R5 to IL-2 (R5-IL2) and Z6 to IL-2 (Z6-IL2). Single-chain constructs were designed such that R5 or Z6 is connected to the N-terminus of IL-2 via a flexible 5x(Gly-Ser) linker (Fig. 2B), which allows R5/Z6 and IL-2 to independently bind their respective targets. R5-IL2 and Z6-IL2 were expressed in insect cells using Baculovirus and both proteins eluted from a gel filtration column as monodisperse peaks, which indicates favorable biochemical behavior (Fig. S2).

Biophysical characterization of surrogate RSPO-receptor interactions.

We used surface plasmon resonance (SPR) to measure the binding affinity of R5-IL2 and Z6-IL2 for RNF43 and ZNRF3, respectively. We determined that R5-IL2 bound to RNF43 with a dissociation constant (K_d) of 58 nM, and that Z6-IL2 bound to ZNRF3 with a K_d of 80 nM (Fig. 2C and 2D). R5-IL2 did not cross-react with ZNRF3, and Z6-IL2 did not cross-react with RNF43 (Fig. S3). We also used SPR to determine that R5-IL2 bound to CD25 with a K_d of 8.4 nM and that Z6-IL2 bound to CD25 5.7 nM, both of which are similar to the reported K_d values of the wild-type CD25-IL2 interaction (Fig. S4) (18,19). Collectively, our SPR experiments revealed that (i) R5 and Z6 bind to their respective targets with similar affinities, (ii) that the affinities of surrogate RSPOs fall within the range of those reported between wild-type RSPOs and RNF43/ZNRF3 (5 nM to 10,000 nM) (7,8), and (iii) that CD25 binding is retained in the surrogate RSPO format.

Surrogate RSPO-mediated enhancement of Wnt signaling.

We performed a luciferase reporter assay to test the selectivity of R5-IL2 and Z6-IL2 in

potentiating Wnt signaling. We compared the activity of surrogate RSPOs on HEK 293 Super Top Flash (STF) Wnt reporter cells, or on HEK 293 STF cells that had been retrovirally transduced with CD25. In the CD25-expressing cells, R5-IL2 increased Wnt3a reporter activity up to 9.5-fold, and Z6-IL2 increased Wnt3a activity up to 41-fold (Fig. 3A). The greater activity of Z6-IL2 relative to R5-IL2 is consistent with a previous report that ZNRF3 is the dominantly expressed homolog (relative to RNF43) in HEK 293 cells (2). Given that natural RSPOs are cross-reactive, and that sequestration of both RNF43 and ZNRF3 may be required for the potent activity of RSPOs, we posited that the combination of R5-IL2 and Z6-IL2 would have a synergistic effect. We therefore incubated the cells with a 1:1 mixture of R5-IL2 and Z6-IL2, and found that this combination increased Wnt3a activity by 148-fold (Fig. 3A). However, in the untransduced cells, the addition of R5-IL2, Z6-IL2, and R5-IL2+Z6-IL2 led to increases in Wnt3a activity of only 2.7, 3.9 and 5.8-fold, respectively (Fig. 3B), indicating that surrogate RSPO activity is highly selective for cells expressing CD25.

In both the CD25 positive and CD25 negative cells, we found that a mixture of conditioned Wnt3a media and RSPO2 (Furin domains 1 and 2) elicited a response that was 286- to 304-fold greater than that of Wnt3a alone (Fig. 3A, 3B), which is similar to the previously observed levels of RSPO-mediated enhancement (3). As negative controls, we treated cells with Wnt3a conditioned media supplemented with either R5, Z6, or IL-2. The R5 scFv and IL2 cytokine did not substantially potentiate Wnt3a signaling (Fig. 3B), however, the Z6 scFv exhibited mild agonist activity that resulted in 10-fold or 12-fold increases in signaling in the HEK293 STF and CD25-expressing cells, respectively (Fig. 3A, 3B).

Surrogate RSPO-mediated stimulation of intestinal organoid growth.

The growth of primary human colon organoids is presumably driven by LGR5+ stem cells and requires the addition of exogenous RSPOs along with Wnt, EGF, and Noggin (8,9). To determine whether R5-IL2 or Z6-IL2 can substitute for RSPOs in human colon organoid

Surrogate R-spondin agonists potentiate Wnt signaling

culture, we performed an organoid growth assay using organoids that had been retrovirally transduced to express CD25. In this assay, CD25⁺ organoid cells were cultured in a basal medium of Wnt3A, EGF and Noggin, supplemented with one of the following recombinant proteins: RSPO2, IL-2, R5 scFv, Z6 scFv, R5-IL2, Z6-IL2, or a 1:1 mixture of R5-IL2 and Z6-IL2. Subsequently, organoid growth was monitored by fluorescence. Although not as potent as RSPO2 (73-fold), the addition of R5-IL2, Z6-IL2, or the mixture of R5-IL2 and Z6-IL2 each led to significant increases in organoid growth relative to the negative control protein, IL-2 (R5-IL2: 7.3-fold, Z6-IL2: 9.4-fold, mixture: 11-fold) (Fig. 4). However, the addition of R5 scFv or Z6 scFv did not result in a significant increase. Importantly, neither R5-IL2 nor Z6-IL2 stimulated the growth of wild-type organoids (Fig. 4), indicating that surrogate RSPOs specifically affect cells expressing CD25.

DISCUSSION

The successful development of surrogate RSPO proteins indicates that a non-natural receptor, CD25, may be co-opted to facilitate selective amplification of Wnt signaling. Bispecific surrogate RSPOs do not include any components that recognize cognate LGR or HSPG receptors and their Wnt-potentiating activity is likely driven instead by IL-2-mediated endocytosis of CD25. This finding supports the present model that RSPOs function by inducing membrane clearance of RNF43/ZNRF3 and opens new avenues for the engineering surrogate RSPOs that act upon a broad spectrum of receptors that induce endocytosis. For example, tyrosine kinase receptors, G-protein coupled receptors (GPCRs), pattern recognition receptors and several other classes of receptors are internalized upon ligand-binding and may serve as alternatives for the targeting arms of surrogate RSPOs.

The selective amplification of Wnt signaling on CD25⁺ cells by R5-IL2 and Z6-IL2 has the potential to influence immune cell behavior for biomedical applications. CD25 is expressed predominantly on activated T cells, regulatory T cells (T regs), and natural killer cells (NK cells) (20) and Wnt activation is known to modulate the behavior of several T cell subtypes. For example,

Wnt signaling attenuates the suppressive activity of T regs by disrupting Foxp3 transcription activity (21), and autocrine inhibition of Wnt signaling in NK cells mediates tumor immune evasion by downregulating the expression of activating ligands (22). Administration of CD25-targeted surrogate RSPOs alone or in combination with Wnt agonists may therefore augment the effectiveness of T cell- or NK-cell based cancer immunotherapies.

We designed R5-IL2 and Z6-IL2 such that they bind to RNF43 and ZNRF3, respectively, and neither surrogate RSPO is cross-reactive (Fig. 2C&D, S3). In our 293-based Wnt reporter assay, the highest levels of Wnt signal enhancement were achieved by co-administering R5-IL2 and Z6-IL2 (Fig. 3), suggesting that maximal activity requires the antagonism of both RNF43 and ZNRF3. Natural RSPOs, which are cross-reactive with both RNF43 and ZNRF3, were also more potent stimulators of colon organoid growth when compared to R5-IL2 and Z6-IL2 (Fig. 4). Based on these findings, we anticipate that second-generation surrogate RSPOs may be optimized through the incorporation of cross-reactive RNF43/ZNRF3 binding modules, such as the RSPO Fu1 domain (8), in place of the scFv component. Additional optimization strategies may involve the use of scFvs that recognize different epitopes on the RNF43 or ZNRF3 ECDs or the use of scFvs that bind to RNF43 or ZNRF3 with higher affinity. On the other hand, the ability of R5-IL2 and Z6-IL2 to target a given E3 ligase (RNF43 versus ZNRF3) confers an additional degree of selectivity that is not attainable using natural RSPOs, and this property may be advantageous for targeting cell types that express only one of the two E3 ligases. Collectively, our data provide proof-of-concept for surrogate RSPOs whose modular design will allow for the targeted enhancement of Wnt signaling on cells expressing RNF43, ZNRF3 or various other tissue-specific markers.

EXPERIMENTAL PROCEDURES

Protein expression and purification

The ECDs of human RNF43 (amino acids 24-197) and ZNRF3 (amino acids 56-219) were cloned into the pAcGp67A vector and modified to include a C-terminal biotin-acceptor peptide tags (GLNDIFEAQKIEW) followed by a

Surrogate R-spondin agonists potentiate Wnt signaling

hexahistidine tag. Unless otherwise specified, when RNF43 or ZNRF3 are named in the Experimental Procedures section, this refers to the ECD only. The R5 and Z6 human antibody scFv fragments were cloned into pAcGp67A vector and include a C-terminal hexahistidine tag. These human antibody scFv fragments were also cloned into the pAcGp67A vector in frame with a (GS)₅ linker followed by human interleukin-2 (amino acids 21-153) and a hexahistidine tag to generate the R5-IL2 and Z6-IL2 surrogate RSPOs. All proteins were expressed in Hi-Five cells (Invitrogen) from *Trichoplusia ni* using Baculovirus and the cultures were harvested 60 hours after infection. Proteins were purified by nickel affinity chromatography followed by size exclusion chromatography in HBS buffer (10 mM HEPES pH 7.4, 150 mM sodium chloride). RNF43 and ZNRF3 were site-specifically biotinylated by BirA ligase at the C-terminal biotin-acceptor peptide prior to size-exclusion chromatography. All proteins were either used when freshly purified or supplemented with 20% glycerol, snap-frozen in liquid nitrogen, and then thawed for use as necessary.

Selection of RNF43/ZNRF3-binding scFv fragments

The nonimmune yeast surface display library of human antibody scFv fragments was generously provided by Prof. Dane Wittrup (16). A sequential selection strategy involving a series of magnetic bead selections followed by flow-cytometric sorting was used as previously reported (23). Round 1 of the selections was performed by mixing 250 μ L magnetic streptavidin microbeads (Miltenyi) with either biotinylated RNF43 or ZNRF3 to a final concentration of 400 nM. These microbeads were further mixed with 10^{10} yeast cells from the scFv library in 1 X phosphate buffered saline (PBS) containing 0.5% bovine serum albumin and 2mM EDTA (PBE). Yeast were subsequently passed over a Magnetic-Activated cell sorting (MACS) LS separation column (Miltenyi), the flow through was discarded, and the RNF43 or ZNRF3 binders were isolated from the column elution. Round 2 was performed as Round 1 except 10^8 yeast cells from the recovered Round 1 sample were used for the selection. For Round 3, yeast cells were incubated with 200 nM biotinylated

RNF43 or ZNRF3 ECD, washed with PBE, and then stained with 50 nM streptavidin labeled with Alexafluor-647 (SA-647). The RNF43 and ZNRF3 binders were enriched by MACS using anti-647 microbeads (Miltenyi). In Round 4, yeast cells were pre-incubated with 5 nM biotinylated RNF43 or 10 nM biotinylated ZNRF3, washed with PBE, and then double-stained with SA-647 and Alexa Fluor 488-conjugated antibody to the c-Myc epitope (Myc-488, Cell Signaling). High-affinity RNF43 or ZNRF3 binders were isolated by fluorescence-activated cell sorting (FACS).

Plasmids from the final round of selection were isolated using the Zymoprep Yeast Plasmid Miniprep Kit (Zymo Research) and sequenced. These plasmids were electroporated into *S. cerevisiae* EBY100 yeast and recovered in SDCAA selection media, followed by induction in SGCAA induction media (23). The individual scFv-transformed yeast was incubated with increasing concentrations of biotinylated RNF43 or ZNRF3 proteins. These were stained with SA-647 and Myc-488 and fluorescence was monitored by flow cytometry. The data were analyzed by GraphPad Prism 7 and the highest affinity clones encoding full scFv sequences (R5, Z6) were selected for use in subsequent experiments.

Binding of RNF43 and ZNRF3 to yeast surface displayed scFvs

Yeast cells expressing either the R5 or Z6 scFvs were stained with 1 μ M recombinant biotinylated RNF43 or ZNR3 ECDs, respectively, in PBE at 4 °C for 2 hours with slow rotation. Cells were then washed, incubated with SA-647 and Myc-488 for 20 min, washed again, and then analyzed by flow cytometry.

Surface plasmon resonance

All binding measurements were conducted using a BIAcore T100 instrument (GE Healthcare). Biotinylated RNF43, ZNRF3, and CD25 were coupled on an SA sensor chip (GE Healthcare) at low density. An unrelated biotinylated protein was captured at equivalent coupling density to the reference flow cell. Increasing concentrations of R5-IL2 and Z6-IL2 were injected onto the chip in HBS-P (GE Healthcare) at 30 μ L/mL for 120 s or 300 s,

Surrogate R-spondin agonists potentiate Wnt signaling

respectively. Subsequently, the chip was regenerated with 0.5 M MgCl₂ and 25% ethylene glycol in HBS-P buffer for 80 s after each injection. Resonance units were calculated by subtracting the resonance units observed in RNF43, ZNRF3, or CD25-containing flow cells from those of the control flow cell. Curves were fitted to a 1:1 binding model using the accompanying Biacore evaluation software (Biacore/GE Healthcare).

Retrovirus production

Human full-length CD25 was cloned into retroviral vector pMSCV-IRES-YFP (a gift from Dr. Melissa McCracken, Stanford University). Retrovirus was packaged in HEK293T cells as described (24). Briefly, HEK293T cells were plated the day before transfection, cells were transfected with a mixture of the pMSCV-CD25-IRES-GFP vector and the pCL-10A packaging vector using X-tremeGENE HP (Sigma-Aldrich). The supernatant was collected and then filtered using a sterile 0.45 µm filter.

Luciferase signaling assays

HEK293T cells were stably transduced with Firefly Luciferase Reporter under the control of a concatemer of seven LEF/TCF binding sites and Renilla Luciferase Reporter by lentivirus as previously reported from the Garcia laboratory (3). These cells were further stably infected with human CD25 by retrovirus co-expressing YFP. Transduced HEK293T cells were sorted by FACS to enrich for YFP expression. CD25 surface expression was further confirmed by staining the transduced HEK293T cells with an anti-CD25 Brilliant Violet 605 antibody (BioLegend, Clone BC96, #302632). Fluorescence was monitored by flow cytometry and was compared to non-transduced HEK293T cells (data not shown). Luciferase reporter assays were conducted using the Dual Luciferase Assay

kit (Promega) as instructed. Briefly, cells were plated at a density of 10,000 cells per well in a 96-well plate and incubated for 24 hours prior to stimulation. Following the plating step, cells were stimulated with 20% Wnt3A conditioned media (ATCC) supplemented with the Fu1-Fu2 domains of the RSPO2 protein (25 nM) or various concentrations of R5-IL2 or Z6-IL2. Cells were cultured in the presence of signaling proteins for another 24 hours before being washed and lysed as suggested by the instructions in the Dual Luciferase Assay Kit (Promega) manual. Luciferase activity was recorded using SpectraMax Paradigm and analyzed using GraphPad Prism 7.

Organoid growth assays

Fresh human colonic samples were collected at Stanford Hospital with informed consent and grown as primary organoids in Matrigel domes as described (3). The culture was passaged several times in WENR media (Wnt3a, R-spondin, EGF, Noggin) to confirm robust outgrowth of organoids. To evaluate surrogate activity, the organoids were transduced with retrovirus expressing CD25 and FACS sorted for YFP+ (YFP co-expression with CD25) to enrich the CD25+ population. CD25+ cells were then subcultured in basal media containing Wnt3a, EGF and Noggin but lacking RSPO, and then supplemented with test candidate surrogate R-spondins or RSPO2 as in Fig. 4. Organoid morphology was observed and phase contrast pictures were taken using Nikon TS100. To quantify cell growth, organoids were dissociated to single cells and plated at 10,000 cells per well in 96-well format. Cell viability assay was performed 3 days after plating using AlamarBlue cell viability reagent (Thermo Fisher).

Surrogate R-spondin agonists potentiate Wnt signaling

Acknowledgements

We thank Lora Picton for edits and helpful comments. This work was supported by a gift from the Mathers Fund, and the Ludwig Cancer Foundation (K.C.G.); the Howard Hughes Medical Institute (K.C.G.); NIH R01-DK115728 (K.C.G. and C.J.K.) and NIH DK085527 (C.J.K.).

Competing financial interests

K.C.G. and C.J.K. are founders of Surrozen, Inc. K.C.G. and V.C.L. are inventors on a patent applied for by Stanford University describing surrogate RSPOs.

Surrogate R-spondin agonists potentiate Wnt signaling

REFERENCES

1. Nusse, R., and Clevers, H. (2017) Wnt/ β -Catenin Signaling, Disease, and Emerging Therapeutic Modalities. *Cell* **169**, 985-999
2. Hao, H. X., Xie, Y., Zhang, Y., Charlat, O., Oster, E., Avello, M., Lei, H., Mickanin, C., Liu, D., Ruffner, H., Mao, X. H., Ma, Q. C., Zamponi, R., Bouwmeester, T., Finan, P. M., Kirschner, M. W., Porter, J. A., Serluca, F. C., and Cong, F. (2012) ZNRF3 promotes Wnt receptor turnover in an R-spondin-sensitive manner. *Nature* **485**, 195-200
3. Janda, C. Y., Dang, L. T., You, C. J., Chang, J. L., de Lau, W., Zhong, Z. D. A., Yan, K. S., Marecic, O., Siepe, D., Li, X. N., Moody, J. D., Williams, B. O., Clevers, H., Piehler, J., Baker, D., Kuo, C. J., and Garcia, K. C. (2017) Surrogate Wnt agonists that phenocopy canonical Wnt and beta-catenin signalling. *Nature* **545**, 234-237
4. Koo, B. K., Spit, M., Jordens, I., Low, T. Y., Stange, D. E., van de Wetering, M., van Es, J. H., Mohammed, S., Heck, A. J. R., Maurice, M. M., and Clevers, H. (2012) Tumour suppressor RNF43 is a stem-cell E3 ligase that induces endocytosis of Wnt receptors. *Nature* **488**, 665-669
5. de Lau, W., Barker, N., Low, T. Y., Koo, B. K., Li, V. S. W., Teunissen, H., Kujala, P., Haegebarth, A., Peters, P. J., van de Wetering, M., Stange, D. E., van Es, J. E., Guardavaccaro, D., Schasfoort, R. B. M., Mohri, Y., Nishimori, K., Mohammed, S., Heck, A. J. R., and Clevers, H. (2011) Lgr5 homologues associate with Wnt receptors and mediate R-spondin signalling. *Nature* **476**, 293-297
6. Xie, Y., Zamponi, R., Charlat, O., Ramones, M., Swalley, S., Jiang, X. M., Rivera, D., Tschantz, W., Lu, B., Quinn, L., Dimitri, C., Parker, J., Jeffery, D., Wilcox, S. K., Watrobka, M., LeMotte, P., Granda, B., Porter, J. A., Myer, V. E., Loew, A., and Cong, F. (2013) Interaction with both ZNRF3 and LGR4 is required for the signalling activity of R-spondin. *Embo Reports* **14**, 1120-1126
7. Wang, D., Huang, B., Zhang, S., Yu, X., Wu, W., and Wang, X. (2013) Structural basis for R-spondin recognition by LGR4/5/6 receptors. *Genes & development* **27**, 1339-1344
8. Zebisch, M., Xu, Y., Krastev, C., MacDonald, B. T., Chen, M., Gilbert, R. J., He, X., and Jones, E. Y. (2013) Structural and molecular basis of ZNRF3/RNF43 transmembrane ubiquitin ligase inhibition by the Wnt agonist R-spondin. *Nature communications* **4**, 2787
9. Lebensohn, A. M., and Rohatgi, R. (2018) R-spondins can potentiate WNT signaling without LGRs. *eLife* **7**, 18
10. Sato, T., Vries, R. G., Snippert, H. J., van de Wetering, M., Barker, N., Stange, D. E., van Es, J. H., Abo, A., Kujala, P., Peters, P. J., and Clevers, H. (2009) Single Lgr5 stem cells build crypt-villus structures in vitro without a mesenchymal niche. *Nature* **459**, 262-265
11. Ootani, A., Li, X. N., Sangiorgi, E., Ho, Q. T., Ueno, H., Toda, S., Sugihara, H., Fujimoto, K., Weissman, I. L., Capecchi, M. R., and Kuo, C. J. (2009) Sustained in vitro intestinal epithelial culture within a Wnt-dependent stem cell niche. *Nature Medicine* **15**, 701-706
12. Zhou, W.-J., Geng, Z. H., Spence, J. R., and Geng, J.-G. (2013) Induction of intestinal stem cells by R-spondin 1 and Slit2 augments chemoradioprotection. *Nature* **501**, 107-111
13. Yan, K. S., Janda, C. Y., Chang, J. L., Zheng, G. X. Y., Larkin, K. A., Luca, V. C., Chia, L. A., Mah, A. T., Han, A., Terry, J. M., Ootani, A., Roelf, K., Lee, M., Yuan, J., Li, X., Bolen, C. R., Wilhelmy, J., Davies, P. S., Ueno, H., von Furstenberg, R. J., Belgrader, P., Ziraldo, S. B., Ordonez, H., Henning, S. J., Wong, M. H., Snyder, M. P., Weissman, I. L., Hsueh, A. J., Mikkelsen, T. S., Garcia, K. C., and Kuo, C. J. (2017) Non-equivalence of Wnt and R-spondin ligands during Lgr5(+) intestinal stem-cell self-renewal. *Nature* **545**, 238-242
14. Zhan, T., Rindtorff, N., and Boutros, M. (2017) Wnt signaling in cancer. *Oncogene* **36**, 1461
15. Hémar, A., Subtil, A., Lieb, M., Morelon, E., Hellio, R., and Dautry-Varsat, A. (1995) Endocytosis of interleukin 2 receptors in human T lymphocytes: distinct intracellular localization and fate of the receptor alpha, beta, and gamma chains. *The Journal of cell biology* **129**, 55-64
16. Feldhaus, M. J., Siegel, R. W., Opresko, L. K., Coleman, J. R., Feldhaus, J. M. W., Yeung, Y. A., Cochran, J. R., Heinzelman, P., Colby, D., Swers, J., Graff, C., Wiley, H. S., and Wittrup, K. D.

Surrogate R-spondin agonists potentiate Wnt signaling

- (2003) Flow-cytometric isolation of human antibodies from a nonimmune *Saccharomyces cerevisiae* surface display library. *Nature Biotechnology* **21**, 163-170
17. Glinka, A., Dolde, C., Kirsch, N., Huang, Y. L., Kazanskaya, O., Ingelfinger, D., Boutros, M., Cruciat, C. M., and Niehrs, C. (2011) LGR4 and LGR5 are R-spondin receptors mediating Wnt/ β -catenin and Wnt/PCP signalling. *EMBO reports* **12**, 1055-1061
 18. Rickert, M., Boulanger, M. J., Goriatcheva, N., and Garcia, K. C. (2004) Compensatory energetic mechanisms mediating the assembly of signaling complexes between interleukin-2 and its α , β , and γ receptors. *Journal of molecular biology* **339**, 1115-1128
 19. Spangler, J. B., Tomala, J., Luca, V. C., Jude, K. M., Dong, S., Ring, A. M., Votavova, P., Pepper, M., Kovar, M., and Garcia, K. C. (2015) Antibodies to interleukin-2 elicit selective T cell subset potentiation through distinct conformational mechanisms. *Immunity* **42**, 815-825
 20. Liao, W., Lin, J.-X., and Leonard, W. J. (2013) Interleukin-2 at the crossroads of effector responses, tolerance, and immunotherapy. *Immunity* **38**, 13-25
 21. van Loosdregt, J., Fleskens, V., Tiemessen, M. M., Mokry, M., van Boxtel, R., Meering, J., Pals, C., Kurek, D., Baert, M. R. M., Delemarre, E. M., Grone, A., Koerkamp, M., Sijts, A., Nieuwenhuis, E. E. S., Maurice, M. M., van Es, J. H., ten Berge, D., Holstege, F. C., Staal, F. J. T., Zaiss, D. M. W., Prakken, B. J., and Coffey, P. J. (2013) Canonical Wnt Signaling Negatively Modulates Regulatory T Cell Function. *Immunity* **39**, 298-310
 22. Malladi, S., Macalinao, D. G., Jin, X., He, L., Basnet, H., Zou, Y., de Stanchina, E., and Massagué, J. (2016) Metastatic latency and immune evasion through autocrine inhibition of WNT. *Cell* **165**, 45-60
 23. Chao, G., Lau, W. L., Hackel, B. J., Sazinsky, S. L., Lippow, S. M., and Wittrup, K. D. (2006) Isolating and engineering human antibodies using yeast surface display. *Nature protocols* **1**, 755
 24. Naviaux, R. K., Costanzi, E., Haas, M., and Verma, I. M. (1996) The pCL vector system: rapid production of helper-free, high-titer, recombinant retroviruses. *Journal of virology* **70**, 5701-5705
 25. Janda, C. Y., Waghray, D., Levin, A. M., Thomas, C., and Garcia, K. C. (2012) Structural Basis of Wnt Recognition by Frizzled. *Science* **337**, 59-64
 26. Cheng, Z., Biechele, T., Wei, Z., Morrone, S., Moon, R. T., Wang, L., and Xu, W. (2011) Crystal structures of the extracellular domain of LRP6 and its complex with DKK1. *Nature structural & molecular biology* **18**, 1204

Surrogate R-spondin agonists potentiate Wnt signaling




Figure 1. The R-spondin signaling mechanism. In the absence of RSPO, RNF43 and ZNRF3 drive membrane clearance of Fzd receptors to negatively regulate Wnt signaling. RSPO-mediated crosslinking of the ECD of RNF43 or ZNRF3 to the ECD of LGR4, LGR5, or LGR6 sequesters RNF43/ZNRF3 to restore Fzd surface levels, thereby potentiating Wnt activity. Surrogate RSPOs mimic the function of wild type RSPOs by cross-linking RNF43 or ZNRF3 to tissue-specific markers known to undergo endocytosis upon ligand binding. PDB files 5UK5 (7), 4F0A (25), 3S8Z, and 3S94 (26) were used to generate the images above.

Surrogate R-spondin agonists potentiate Wnt signaling




Figure 2. Characterization of RNF43- and ZNR3-specific scFvs. (A) Flow cytometry dot plots depicting the binding of yeast-displayed R5 and Z6 scFvs to RNF43 and ZNR3, respectively. R5- or Z6-expressing yeast were stained with 1 μ M concentrations of RNF43 or ZNR3 and surface expression was detected with an antibody to the c-Myc epitope. (B) Construct design for R5-IL2 and Z6-IL2 surrogate RSPOs. (C) & (D). SPR was used to measure the binding of R5-IL2 or Z6-IL2 to RNF43 and ZNR3, respectively. Cartoons (left) depict the orientation of streptavidin (SA)-coupled RNF43 or ZNR3 ECDs and the R5-IL2 or Z6-IL2 analytes on the sensor chip. Curves (center) indicate a series of injections of R5-IL2 (3-fold dilutions, maximum concentration 3 μ M) or Z6-IL2 (2-fold dilutions, maximum concentration 1 μ M). Dissociation constants were obtained by plotting maximum RU values and fitting to a 1:1 binding model (right).

Surrogate R-spondin agonists potentiate Wnt signaling




Figure 3. Surrogate RSPOs potentiate Wnt signaling. (A) Representative Luciferase reporter assay measuring surrogate RSPo potentiation of Wnt activity in CD25-expressing cells. HEK STF 293T cells retrovirally transduced with CD25 were incubated with various recombinant proteins (final protein concentrations indicated on graph labels) in the presence of 20% Wnt3a conditioned media. Error bars represent the standard deviation between three independent measurements. (B) A reporter assay was performed under the same conditions as in (A) using uninfected (CD25 negative) HEK STF 293T cells.

Surrogate R-spondin agonists potentiate Wnt signaling




Figure 4. Surrogate RSPOs stimulate intestinal organoid growth. Surrogate RSPOs were tested for their ability to stimulate the growth of LGR5+ human colon organoids that had been retrovirally transduced to express CD25. Basal media containing Wnt3a, EGF and Noggin, but lacking RSPO2 was supplemented with 500nM concentrations of the indicated proteins, or with 25nM RSPO2. Microscopy was used to observe the appearance of spherical organoids in each condition (top) and growth was quantitated by fluorescence (bar graphs, bottom). The bar graph on the right is a zoomed panel of the colored region from the graph on the left.