2 Roost selection by male northern long-eared bats (*Myotis*

3 *septentrionalis*) in a managed fire-adapted forest

Jesse M. Alston[a,b*], Ian M. Abernethy[b], Douglas A. Keinath[c], and Jacob R.

Goheen[d]

a. Program in Ecology, Department of Zoology and Physiology, University of Wyoming,

Laramie, WY 82071, USA

b. Wyoming Natural Diversity Database, University of Wyoming, Laramie, WY 82071, USA

c. Wyoming Ecological Services Field Office, U. S. Fish and Wildlife Service, Cheyenne, WY

82009, USA

d. Department of Zoology and Physiology, University of Wyoming, Laramie, WY 82071, USA

*Corresponding author: jalston@uwyo.edu

Abstract: Wildlife conservation in multi-use landscapes requires identifying and conserving

critical resources that may otherwise be destroyed or degraded by human activity. Summer day-

roost sites are critical resources for bats, so conserving roost sites is thus a focus of many bat

conservation plans. Studies quantifying day-roost characteristics typically focus on female bats

due to their relative importance for reproduction, but large areas of species' ranges can be

occupied predominantly by male bats due to sexual segregation. We used VHF telemetry to

identify and characterize summer day-roost selection by male northern long-eared bats (*Myotis

septentrionalis*) in a ponderosa pine (*Pinus ponderosa*) forest in South Dakota, USA. We tracked

18 bats to 43 tree roosts and used an information theoretic approach to determine the relative

importance of tree- and plot-level characteristics on roost site selection. Bats selected roost trees

that were larger in diameter, more decayed, and under denser canopy than other trees available

on the landscape. Much like studies of female northern long-eared bats have shown, protecting

large-diameter snags within intact forest is important for the conservation of male northern long-

eared bats. Unlike female-specific studies, however, many roosts in our study (39.5%) were

located in short (≤ 3 m) snags. Protecting short snags may be a low-risk, high-reward strategy for

conservation of resources important to male northern long-eared bats. Other tree-roosting bat

species in fire-prone forests are likely to benefit from forest management practices that promote

these tree characteristics, particularly in high-elevation areas where populations largely consist of

males.

1. Introduction

Habitat degradation by humans is a leading cause of extinction and population declines of species globally (Dobson et al., 1997; Halpern et al., 2008; Hansen et al., 2013). Less than 15% of Earth's land surface falls within a protected area, and less than half of that area is free from human development, agriculture, livestock grazing, light pollution, and transportation infrastructure (Jones et al., 2018). Even in relatively undisturbed areas, land uses other than conservation of nature—such as wildfire prevention, livestock grazing, recreation, and extraction of timber and other forest products—are the norm rather than the exception. Conservation measures targeting these multi-use landscapes are thus vital for conserving species (Kremen and Merenlender, 2018).

In multi-use landscapes, successful conservation often requires the identification of critical resources for species of conservation concern so that the supply of those critical resources can be maintained or increased. Day-roosts appear to be critical resources for many bats, providing shelter from predators and environmental stressors (Fenton et al., 1994; Solick and Barclay, 2006), communal sites for social interactions (Willis and Brigham, 2004), and secure places to raise young (Kunz, 1982). Bats spend most of their time in day-roosts, alone or in groups of up to millions of individuals, depending on sex, species, and reproductive status. Patterns of bat abundance and distribution are correlated with roost availability (Humphrey, 1975), and declines in reproductive success have been documented when pregnant or lactating bats are experimentally excluded from preferred roosts (Brigham and Fenton, 1986). Because day-roosts are so important for bats, measures to conserve roosts feature prominently in bat conservation plans. Resource managers seeking to conserve bats while managing landscapes for multiple uses benefit from knowledge that promotes bat roost conservation.

We evaluated day-roost selection by male northern long-eared bats (*Myotis*

septentrionalis) in a ponderosa pine forest in the Black Hills of South Dakota, USA. Our study

population inhabits an intensively logged landscape at the western edge of this species' range.

Northern long-eared bats inhabit much of the eastern United States and southern Canada

(Caceres and Barclay, 2000), but are increasingly threatened by white nose syndrome and have

been protected under the Endangered Species Act since 2015. Throughout their range, northern

long-eared bats roost almost exclusively in tree cavities and under sloughing bark within intact

forest (Lacki et al., 2009), and forage within forests or at forest edges (Henderson and Broders,

2008; Owen et al., 2003; Patriquin and Barclay, 2003).

At our study site and other high elevation areas in the Black Hills, male bats are much

more common than females (Choate and Anderson, 1997; Cryan et al., 2000). Sexual segregation

driven by elevation or temperature is widespread among bats, and is believed to be driven by

differences in energy requirements that allow males to inhabit areas that are colder or have less

prey (Barclay, 1991; Ford et al., 2002; Senior et al., 2005). Male northern long-eared bats are

therefore likely to occupy substantially different habitat than females, but range-wide

conservation for the species is informed predominantly by studies focusing on female bats (J.

Alston, unpublished data). Forest managers in male-dominated areas may therefore rely on

incomplete information to conserve the majority of bats within their jurisdictions. Our study

provides managers in such areas information to appropriately guide management in male-

dominated areas and supplement the existing wealth of information on female habitat use.

To evaluate factors driving roost selection, we tracked adult male northern long-eared

bats to day-roosts and quantified characteristics of both used and available roost trees using

variables easily measured by forest and wildlife managers. We evaluated these data using an

information-theoretic approach to select the best models from a suite of candidate models. We

hypothesized that in this intensively logged ecosystem, bats primarily select roost trees with

characteristics that promote cavity formation (e.g., tree size and amount of decay), the number of

nearby roosts (e.g., plot-level tree and snag density), and thermal characteristics suitable for

behavioral thermoregulation (e.g., canopy cover and orientation in relation to sunlight).

2. Methods

2.1. Study Area

We conducted our study during the summers of 2017 and 2018 on Jewel Cave National

Monument (43° 45' N, 103° 45' W) and surrounding areas of Black Hills National Forest, 16 km

west of Custer, South Dakota, USA. In this area, mean monthly summer high temperatures range

between 22 – 27° C and mean monthly summer precipitation ranges between 60 – 80 mm

(Western Regional Climate Center, 2018). Open ponderosa pine (*Pinus ponderosa*) forests

dominate our study site, with Rocky Mountain juniper (*Juniperus scopulorum*) and quaking

aspen (*Populus tremuloides*) occurring locally. In our local study area, forests form a

heterogenous mosaic with northern mixed-grass prairie where a large stand-replacing fire

occurred in 2000. A large cave system and several smaller caves lie underground at our study

site, and there is substantial topographic relief on the landscape in the form of intersecting

canyon systems and rock outcrops.

Forests in this landscape are intensively managed. Black Hills National Forest typically

uses even-aged management techniques other than clear-cutting (e.g., two-step shelterwood

harvest). Stand harvest rotations are 120 years on average, but selective cutting occurs at 10- to

20-year intervals to harvest mature trees and thin the understory. Aside from large severe

wildfires, the forest self-regenerates and does not require planting. Forest management on private

lands generally also follow this formula but thinning intervals vary (B. Phillips, personal

communication). Forests on Jewel Cave National Monument are managed for resource

preservation, primarily using prescribed fire.

2.2. Capture and VHF Telemetry

We used mist nets to capture bats over permanent and semi-permanent water sources

(e.g., springs, stock tanks, and stock ponds). In summer (Jun–Aug) 2017 and 2018, we netted 20

and 49 nights at 9 and 15 water sources, respectively. We opened mist nets at civil sunset and

closed them after five hours and during inclement weather. We affixed VHF transmitters (LB-2X

model .28 g – Holohil Systems Ltd., Carp, ON, Canada; .25 g model – Blackburn, Nacogdoches,

TX, USA) between the scapulae of adult male northern long-eared bats with latex surgical

adhesive (Osto-Bond, Montreal Ostomy, Montreal, QC, Canada). In our study area and others in

the region (Cryan et al. 2000), sex ratios are overwhelmingly male. Because patterns of roost

selection can differ between male and female bats (Boland et al., 2009; Elmore et al., 2004; Hein

et al., 2008; Perry and Thill, 2007), we targeted males specifically. Additionally, the roosting

habits of male bats are less studied than those of females—only 2 of the 14 peer-reviewed studies

on roost selection of northern long-eared bats provide data on males, and 11 out of 111 peer-

reviewed studies on roost selection of cavity-roosting bats in general provide data on males (J.

Alston, unpublished data). All transmitters weighed <5% of the mass of the bat (Aldridge and

Brigham, 1988). We tracked bats to roosts each day transmitters were active. All protocols were

approved by the University of Wyoming and National Park Service Animal Care and Use

Committees and met guidelines approved by the American Society of Mammalogists (Sikes,

2016).

2.3. Roost Characterization

To characterize roosts, we collected data for each roost and randomly sampled available

roost trees in our study area. We identified available roost trees by generating a sample of 200

random points within 2.53 km (the farthest distance we located a bat roosting from its capture

site during our study) of sites where we captured northern long-eared bats and selecting the

nearest available roost tree at a random bearing from each point. We defined available roost trees

as live trees >20 cm in diameter or any dead tree with a visible defect (e.g. sloughing bark or

cavities) sufficiently large for a bat to roost within. For each tree and plot, we measured

characteristics that may influence roost suitability (Table 1). We measured vegetation

characteristics at two spatial scales: 1) individual trees, and 2) a 706.86 m^2 (15 m radius) plot

around the tree. We also measured topographic variables at the plot scale.

2.4. Statistical Analysis

To quantify differences between roost trees used by northern long-eared bats and the 200

randomly sampled available roost trees, we used the R statistical software environment (R Core

Team, 2018) to build binomial-family generalized linear models in a use-availability sampling

design (Manly et al., 2007). We employed an information theoretic approach using Akaike's

Information Criterion adjusted for small sample sizes (AIC_c) to compare competing models

(Burnham and Anderson, 2002) using the 'MuMIn' package (Barton, 2018). We calculated AIC_c

values and model weights (w_i) for all possible combinations of a maximum of 8 predictors (one

variable for each 5 observations) in our set of candidate models to prevent bias and unreliable

confidence interval coverage (Vittinghoff and McCulloch, 2007). Predictors with variance

inflation factors > 10 were removed from consideration in our global model to reduce problems

associated with multicollinearity (Kutner, 2005). We averaged model coefficients for all models

with cumulative w_i > .95 (Burnham and Anderson, 2002) using the full-averaging method.

Finally, we validated our averaged model using area under the receiver operating characteristic

curve (AUC; Manel et al., 2001; Swets, 1988).

3. Results

We located 44 roosts used on 59 days by 18 bats during our study. Aside from one roost

in a rock crevice, bats roosted exclusively in ponderosa pines, either in cavities or under loose

bark. Thirty-six out of 43 tree roosts (83.7%) occurred in dead trees (hereafter termed "snags").

We found 2.4 ± 0.3 (range: 1-5) roost trees per bat. Bats typically roosted in the same patch of

contiguous forest for the active life of the transmitter. Bats roosted 790 ± 90 m (range: 55 –

2,530 m) from the sites at which they were captured.

Our global model distinguishing used roost trees from available roost trees incorporated

DBH, tree height, decay class (Maser et al., 1979), slope, aspect (split into two components—

eastness and southness), percent bark remaining, plot tree density, plot snag density, plot canopy

cover, and interaction terms between slope and eastness and slope and southness. The global

model provided an adequate fit to the data (le Cessie-van Houwelingen-Copas-Hosmer global

goodness of fit test; z = 0.805, p = 0.421). Our averaged model (incorporating 104 models in our

confidence set; Table A.1) indicated that DBH, decay class, and canopy cover were important

variables (Table 2). Significant (p < .05) averaged model coefficients, confidence intervals, and

scaled and unscaled odds ratios are reported in Table 3. Mean differences between used and

available roost trees among our variables of interest are reported in Table 4. Predictive

performance of the averaged model was very high (AUC = 0.924).

Three variables (DBH, decay class, and canopy cover) were positively related to roost

selection (Fig. 1; Table 2). For each 5 cm increase in DBH, odds of selection increased by 61%

(CI: 21-113%). For each 1 unit increase in decay class, odds of selection increased by 111% (CI:

47-203%). For each additional 10% increase in canopy cover, the odds of selection increased by

126% (CI: 55-230%).

4. Discussion

Northern long-eared bats primarily selected roosts in trees with characteristics that

promote cavity formation. At the level of individual trees, northern long-eared bats selected for

large diameter trees with substantial decay. This corroborates previous work on northern long-

eared bats (Jung et al., 2004; Rojas et al., 2017) and is intuitive because large trees with more

decay have more roost structures (i.e., cavities and loose bark) for bats to use (Reynolds et al.,

1985). This is particularly true of ponderosa pines, which can produce large amounts of resin to

defend against physical injury (Kane and Kolb, 2010; Lewinsohn et al., 1991) and therefore tend

to develop cavities only when they are scarred or dead. In intensively logged landscapes like the

Black Hills, cavities are found overwhelmingly in snags because most trees are harvested before

they reach ages at which cavities typically form.

Conservation actions targeting northern long-eared bats should include preservation of

large snags whenever possible. Our study demonstrated that northern long-eared bats select

large-diameter snags (>37 cm), and large diameter snags also tend to remain standing longer than

thinner snags (Bull, 1983; Chambers and Mast, 2014). These snags need not be tall—short (≤ 3

m) snags are important resources for male northern long-eared bats as well. Seventeen of 43

(39.5%) roosts that we located occurred in broken-off snags ≤ 3 m in height. These are important

resources and are likely more vulnerable to loss during forest management activities (particularly

prescribed fire) than other potential roost trees. Snags are often intentionally removed during

forest management activities because of hazards posed to forest management personnel (e.g.,

loggers and firefighters) and the general public. However, these short snags pose less danger to

forest management personnel and the public than taller snags, and their preservation is therefore

a realistic and actionable step toward bat conservation.

Of the variables we considered that may influence thermal characteristics of roosts, only

canopy cover influenced roost selection significantly. Trees were more likely to be used as roosts

as surrounding canopy cover increased, and use was greater than availability at all canopy cover

levels >19%. Although many snags were available at our study site in open areas burned by a

severe wildfire in 2000, northern long-eared bats rarely use those snags, instead selecting snags

in the interior of forest stands with live canopy. Forty out of 43 (93.0%) roosts were within or

immediately bordering intact forest stands with live canopy, and all roosts were within 50 m of

intact forest stands. Bats may prefer these areas because canopy cover creates cooler

environments, but they may also simply prefer to be immediately near forested areas where they

forage (Henderson and Broders, 2008; Owen et al., 2003; Patriquin and Barclay, 2003). Either

way, stand-replacing fire likely poses risks to local populations of northern long-eared bats at the

western edge of its range, where severe wildfire is increasingly prevalent due to climate change

(Westerling et al., 2006). Clearcutting also poses risks to local populations of northern long-

eared bats in these areas, even if snags are retained. Selective logging that leaves some level of

canopy cover remaining would ensure that snag retention is effective for bat roost conservation.

Dynamics of regional disturbance may be important when evaluating local-scale factors

that influence roost selection (O'Keefe and Loeb, 2017). The ponderosa-dominated landscape

where we conducted our research is substantially different than other landscapes (i.e., deciduous

and mixed forests in eastern North America) where roost selection by northern long-eared bats

has been studied. Although many of the factors driving roost selection appear to be similar

among areas, the processes that create roosts may be fundamentally different in different areas.

Snags in ponderosa pine forests are often generated in large pulses by severe wildfire and

mountain pine beetles (*Dendroctonus ponderosae*), but the long-term ramifications of these

resource pulses for bats are not well understood. Severe wildfire appears to create snags that are

largely unused by bats. Mountain pine beetle outbreaks may do the same if beetle-induced

mortality reduces or eliminates canopy cover over large areas, or if outbreaks lead to more severe

fires. Northern long-eared bats may instead depend on snag-generating processes that operate at

more local scales and over longer intervals to create suitable roosts.

Roost selection by bats varies by sex, age class, and reproductive condition (Elmore et

al., 2004; Hein et al., 2008). Studies on roost selection generally focus on females because they

tend to drive reproduction, which is required to sustain populations. However, targeting roost

conservation toward females exclusively may neglect resources that are important for males.

Because sex ratios can be heavily biased in some areas (Cryan et al., 2000), ignoring the needs of

males could leave resources that are important for most individuals inhabiting these areas

unprotected. On the other hand, designing roost conservation measures on studies of males alone

will leave resources that are important for females unprotected. For example, short (≤ 3 m) snags

are important resources for males, but they may not be for females, which aggregate in maternity

colonies that require larger cavities than largely solitary males (Perry and Thill, 2007). Resource

managers seeking to conserve bats should take these sex differences into account when

developing conservation plans and designing studies to inform those plans. In high elevation

areas, males may be more important than females for sustaining local populations because there

are few females in those areas.

5. Conclusions

Forest managers require actionable knowledge to guide conservation, and our results

indicate that conserving large-diameter snags within intact forest stands is one such action that

can be taken to conserve bats in wildfire-prone coniferous forests. Short ($\leq$ 3 m) snags in

particular represent a low-risk, high-reward resource to target for preservation in male-biased,

high elevation populations. For federally threatened northern long-eared bats, conserving these

snags at the western edge of their range may prevent range contraction and local extinction.

Similar patterns are likely to hold true for other cavity-roosting bat species in wildfire-prone

coniferous forests, like those found throughout western North America. Although bats face

danger from many threats unrelated to roosts (e.g., white nose syndrome, wind energy

development, etc.), roost conservation remains an important tool for bat conservation in the face

of such threats.

Acknowledgements

Many thanks to L. Boodoo, C. McFarland, E. Greene, B. Tabor, and B. Phillips for help with

fieldwork; J. Rick for helpful comments on pre-submission versions of this manuscript; R.

Anderson-Sprecher for helpful comments concerning statistical analyses; and P. Ortegon, D.

Licht, M. Wiles, D. Austin, B. Phillips, and E. Thomas for their logistical support of this project.

Research funding was provided by the National Park Service, the Department of Zoology and

Physiology at the University of Wyoming, the Berry Ecology Center, the American Society of

Mammalogists, Prairie Biotic Research, Inc., and the Wyoming Chapter of The Wildlife Society.

The findings and conclusions in this article are those of the authors and do not necessarily

represent the views of the U.S. Fish and Wildlife Service or the National Park Service.

Appendix A: Supplementary Data

Table A.1. Candidate models, ΔAIC values, and model weights (w_i) used to determine model-

averaged coefficients.

Data Availability

*Data and R code used in analysis are available in the supplementary material and will be

archived on the lead author's personal website if this manuscript is accepted for publication.

Literature Cited

Aldridge, H.D.J.N., Brigham, R.M., 1988. Load carrying and maneuverability in an

insectivorous bat: a test of the 5% "rule" of radio-telemetry. Journal of Mammalogy 69,

379–382. https://doi.org/10.2307/1381393

Barclay, R.M.R., 1991. Population structure of temperate zone insectivorous bats in relation to

foraging behaviour and energy demand. Journal of Animal Ecology 60, 165–178.

https://doi.org/10.2307/5452

Barton, K., 2018. MuMIn: multi-model inference.

Boland, J.L., Hayes, J.P., Smith, W.P., Huso, M.M., 2009. Selection of day-roosts by Keen's

myotis (*Myotis keenii*) at multiple spatial scales. Journal of Mammalogy 90, 222–234.

https://doi.org/10.1644/07-MAMM-A-369.1

Brigham, R.M., Fenton, M.B., 1986. The influence of roost closure on the roosting and foraging

behaviour of *Eptesicus fuscus* (Chiroptera: Vespertilionidae). Can. J. Zool. 64, 1128–

1133. https://doi.org/10.1139/z86-169

Bull, E.L., 1983. Longevity of snags and their use by woodpeckers (General Technical Report

No. GTR-RM-99), Proceedings of the symposium: snag habitat management. Rocky

Mountain Research Station, Forest Service, US Department of Agriculture, Fort Collins,

CO.

Burnham, K.P., Anderson, D.R., 2002. Model selection and multimodel inference: a practical

information-theoretic approach, 2nd ed. ed. Springer, New York.

Caceres, M.C., Barclay, R.M.R., 2000. *Myotis septentrionalis*. Mammalian Species 1–4.

https://doi.org/10.1644/1545-1410(2000)634<0001:MS>2.0.CO;2

Chambers, C.L., Mast, J.N., 2014. Snag dynamics and cavity excavation after bark beetle

outbreaks in southwestern ponderosa pine forests. Forest Science 60, 713–723.

https://doi.org/10.5849/forsci.13-018

Choate, J.R., Anderson, J.M., 1997. The bats of Jewel Cave National Monument, South Dakota.

The Prairie Naturalist 29, 38–47.

Cryan, P.M., Bogan, M.A., Altenbach, J.S., 2000. Effect of elevation on distribution of female

bats in the Black Hills, South Dakota. Journal of Mammalogy 81, 719–725.

https://doi.org/10.1644/1545-1542(2000)081<0719:EOEODO>2.3.CO;2

Dobson, A.P., Bradshaw, A.D., Baker, A.J.M., 1997. Hopes for the future: restoration ecology

and conservation biology. Science 277, 515–522.

https://doi.org/10.1126/science.277.5325.515

Elmore, L.W., Miller, D.A., Vilella, F.J., 2004. Selection of diurnal roosts by red bats (*Lasiurus*

borealis) in an intensively managed pine forest in Mississippi. Forest Ecology and

Management 199, 11–20. https://doi.org/10.1016/j.foreco.2004.03.045

Fenton, M.B., Rautenbach, I.L., Smith, S.E., Swanepoel, C.M., Grosell, J., van Jaarsveld, J.,

1994. Raptors and bats: threats and opportunities. Animal Behaviour 48, 9–18.

https://doi.org/10.1006/anbe.1994.1207

Ford, W.M., Menzel, M.A., Menzel, J.M., Welch, D.J., 2002. Influence of summer temperature

on sex ratios in eastern red bats (*Lasiurus borealis*). The American Midland Naturalist

147, 179–184.

Halpern, B.S., Walbridge, S., Selkoe, K.A., Kappel, C.V., Micheli, F., D'Agrosa, C., Bruno, J.F.,

Casey, K.S., Ebert, C., Fox, H.E., Fujita, R., Heinemann, D., Lenihan, H.S., Madin,

E.M.P., Perry, M.T., Selig, E.R., Spalding, M., Steneck, R., Watson, R., 2008. A global

map of human impact on marine ecosystems. Science 319, 948–952.

https://doi.org/10.1126/science.1149345

Hansen, M.C., Potapov, P.V., Moore, R., Hancher, M., Turubanova, S.A., Tyukavina, A., Thau,

D., Stehman, S.V., Goetz, S.J., Loveland, T.R., Kommareddy, A., Egorov, A., Chini, L.,

Justice, C.O., Townshend, J.R.G., 2013. High-resolution global maps of 21st-century

forest cover change. Science 342, 850–853. https://doi.org/10.1126/science.1244693

Hein, C.D., Castleberry, S.B., Miller, K.V., 2008. Sex-specific summer roost-site selection by

336   Seminole bats in response to landscape-level forest management. Journal of Mammalogy

337   89, 964–972. https://doi.org/10.1644/07-MAMM-A-335.1

338   Henderson, L.E., Broders, H.G., 2008. Movements and resource selection of the northern long-

339   eared myotis (*Myotis septentrionalis*) in a forest—agriculture landscape. Journal of

340   Mammalogy 89, 952–963. https://doi.org/10.1644/07-MAMM-A-214.1

341   Humphrey, S.R., 1975. Nursery roosts and community diversity of Nearctic bats. Journal of

342   Mammalogy 56, 321–346. https://doi.org/10.2307/1379364

343   Jones, K.R., Venter, O., Fuller, R.A., Allan, J.R., Maxwell, S.L., Negret, P.J., Watson, J.E.M.,

344   2018. One-third of global protected land is under intense human pressure. Science 360,

345   788–791. https://doi.org/10.1126/science.aap9565

346   Jung, T.S., Thompson, I.D., Titman, R.D., 2004. Roost site selection by forest-dwelling male

347   *Myotis* in central Ontario, Canada. Forest Ecology and Management 202, 325–335.

348   https://doi.org/10.1016/j.foreco.2004.07.043

349   Kane, J.M., Kolb, T.E., 2010. Importance of resin ducts in reducing ponderosa pine mortality

350   from bark beetle attack. Oecologia 164, 601–609. https://doi.org/10.1007/s00442-010-

351   1683-4

352   Kremen, C., Merenlender, A.M., 2018. Landscapes that work for biodiversity and people.

353   Science 362, eaau6020. https://doi.org/10.1126/science.aau6020

354   Kunz, T.H., 1982. Roosting ecology of bats, in: Kunz, T.H. (Ed.), Ecology of Bats. Springer US,

355   Boston, MA, pp. 1–55. https://doi.org/10.1007/978-1-4613-3421-7_1

356   Kutner, M.H. (Ed.), 2005. Applied linear statistical models, 5th ed. ed, The McGraw-Hill/Irwin

357   series operations and decision sciences. McGraw-Hill Irwin, Boston.

Lacki, M.J., Cox, D.R., Dickinson, M.B., 2009. Meta-analysis of summer roosting characteristics

of two species of *Myotis* bats. The American Midland Naturalist 162, 318–326.

Lewinsohn, E., Gijzen, M., Croteau, R., 1991. Defense mechanisms of conifers: differences in

constitutive and wound-induced monoterpene biosynthesis among species. Plant

Physiology 96, 44–49. https://doi.org/10.1104/pp.96.1.44

Manel, S., Williams, H.C., Ormerod, S.J., 2001. Evaluating presence–absence models in

ecology: the need to account for prevalence. Journal of Applied Ecology 38, 921–931.

https://doi.org/10.1046/j.1365-2664.2001.00647.x

Manly, B.F.L., McDonald, L., Thomas, D.L., McDonald, T.L., Erickson, W.P., 2007. Resource

selection by animals: statistical design and analysis for field studies. Springer Science &

Business Media.

Maser, C., Anderson, R.G., Kromack, Jr., K., 1979. Dead and down woody material, in: Wildlife

Habitats in Managed Forests: The Blue Mountains of Oregon and Washington.

Agricultural Handbook 553. U.S. Department of Agriculture, Forest Service,

Washington, D.C., pp. 78–95.

O'Keefe, J.M., Loeb, S.C., 2017. Indiana bats roost in ephemeral, fire-dependent pine snags in

the southern Appalachian Mountains, USA. Forest Ecology and Management 391, 264–

274. https://doi.org/10.1016/j.foreco.2017.01.036

Owen, S.F., Menzel, M.A., Ford, W.M., Chapman, B.R., Miller, K.V., Edwards, J.W., Wood,

P.B., 2003. Home-range size and habitat used by the northern myotis (*Myotis*

septentrionalis). The American Midland Naturalist 150, 352–359.

https://doi.org/10.1674/0003-0031(2003)150[0352:HSAHUB]2.0.CO;2

Patriquin, K.J., Barclay, R.M.R., 2003. Foraging by bats in cleared, thinned and unharvested

boreal forest. Journal of Applied Ecology 40, 646–657. https://doi.org/10.1046/j.1365-

2664.2003.00831.x

Perry, R.W., Thill, R.E., 2007. Roost selection by male and female northern long-eared bats in a

pine-dominated landscape. Forest Ecology and Management 247, 220–226.

https://doi.org/10.1016/j.foreco.2007.04.041

R Core Team, 2018. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria.

Reynolds, R.T., Linkhart, B.D., Jeanson, J., 1985. Characteristics of snags and trees containing

cavities in a Colorado conifer forest (USDA Forest Service Research Note No. RM-455).

Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.

Rojas, V.G., O'Keefe, J.M., Loeb, S.C., 2017. Baseline capture rates and roosting habits of

Myotis septentrionalis (northern long-eared bat) prior to white-nose syndrome detection

in the Southern Appalachians. Southeastern Naturalist 16, 140–148.

https://doi.org/10.1656/058.016.0202

Senior, P., Butlin, R.K., Altringham, J.D., 2005. Sex and segregation in temperate bats. Proc Biol

Sci B 272, 2467–2473. https://doi.org/10.1098/rspb.2005.3237

Sikes, R.S., 2016. 2016 guidelines of the American Society of Mammalogists for the use of wild

mammals in research and education. Journal of Mammalogy 97, 663–688.

https://doi.org/10.1093/jmammal/gyw078

Solick, D.I., Barclay, R.M.R., 2006. Thermoregulation and roosting behaviour of reproductive

and nonreproductive female western long-eared bats (*Myotis evotis*) in the Rocky

Mountains of Alberta. Canadian Journal of Zoology 84, 589–599.

https://doi.org/10.1139/z06-028

Swets, J.A., 1988. Measuring the accuracy of diagnostic systems. Science 240, 1285–1293.

https://doi.org/10.1126/science.3287615

Vittinghoff, E., McCulloch, C.E., 2007. Relaxing the rule of ten events per variable in logistic

and Cox regression. Am J Epidemiol 165, 710–718. https://doi.org/10.1093/aje/kwk052

Westerling, A.L., Hidalgo, H.G., Cayan, D.R., Swetnam, T.W., 2006. Warming and earlier

spring increase western U.S. forest wildfire activity. Science 313, 940–943.

https://doi.org/10.1126/science.1128834

Western Regional Climate Center, 2018. NCDC 1981-2010 monthly normals: Custer, SD. URL

https://wrcc.dri.edu/cgi-bin/cliMAIN.pl?sd2087

Willis, C.K.R., Brigham, R.M., 2004. Roost switching, roost sharing and social cohesion: forest-

dwelling big brown bats, *Eptesicus fuscus*, conform to the fission–fusion model. Animal

Behaviour 68, 495–505. https://doi.org/10.1016/j.anbehav.2003.08.028

 Figure Legends

 Fig. 1. Unscaled odds ratios associated with each variable in the averaged roost selection model.

 Error bars represent 95% confidence intervals.

422   **Table 1**. Variables measured at used and available summer day-roosts of male northern long-

423   eared bats (*Myotis septentrionalis*) in the Black Hills of South Dakota, 2017–2018.

Variable	Definition
DBH	Tree diameter at breast height (cm)
Height	Tree height (m)
Snag	Tree status (live/dead)
Decay Class	Stage of tree decay on ordinal scale
Bark Remaining	Bark remaining on tree stem (%)
Canopy Cover	Average of 4 canopy cover measurements (N/E/S/W) taken 5 m from tree (%)
Slope	Slope of 706.9 m^2 (15 m radius) plot centered at tree (%)
Tree Density	Number of trees in 706.9 m^2 plot centered at tree
Snag Density	Number of snags in 706.9 m^2 plot centered at tree
Eastness	Difference between aspect of 706.9 m^2 plot centered at tree and 90 degrees (˚)
Southness	Difference between aspect of 706.9 m^2 plot centered at tree and 180 degrees (˚)
Slope*Eastness	Interaction term between slope and eastness
Slope*Southness	Interaction term between slope and southness

425   **Table 2**. Coefficient estimates in the averaged model and 95% confidence intervals. Bold variables denote significance at α = .05.

Variable	Estimate	LCL (95%)	UCL (95%)
Height	0.0133	-0.0767	0.1033
DBH	**0.0948**	**0.0382**	**0.1514**
Decay Class	**0.7465**	**0.3835**	**1.1094**
Bark Remaining	0.0033	-0.0113	0.0180
Snag Density	0.1010	-0.0039	0.2059
Tree Density	-0.0182	-0.0653	0.0289
Canopy Cover	**0.0816**	**0.0438**	**0.1195**
Slope	0.0323	-0.0354	0.0999
Eastness	-0.0069	-0.0207	0.0068
Southness	0.0004	-0.0041	0.0050
Slope*Eastness	0.0001	-0.0004	0.0005
Slope*Southness	0.0000	-0.0002	0.0002

428   **Table 3**. Averaged model coefficients, confidence intervals, and scaled and unscaled odds ratios for significant variables.

Variable	Coefficient	Unscaled OR	Scaled OR	Units	Scaled OR LCL (95%)	Scaled OR UCL (95%)
DBH	0.0948	1.0995	1.6065	5 cm	1.2105	2.1321
Decay Class	0.7447	2.1095	2.1095	1 unit	1.4674	3.0327
Canopy Cover	0.0814	1.0850	2.2619	10%	1.5491	3.3025

430 **Table 4**. Means and standard errors for variables of interest among used and available trees.

431 Bold font denotes statistically significant variables in the final averaged model.

	Roost		Available	
Variable	Mean	SE	Mean	SE
Height (m)	8.53	1.11	9.01	0.43
DBH (cm)	**35.69**	**1.57**	**30.33**	**0.69**
Decay Class	**4.95**	**0.33**	**3.72**	**0.18**
Bark Remaining (%)	74.19	4.22	69.73	2.49
Snag Density	4.74	1.03	2.12	0.23
Tree Density	19.84	2.15	10.76	1.12
Canopy Cover (%)	**36.83**	**3.02**	**14.96**	**1.39**
Slope (%)	16.87	1.62	11.66	0.64
Eastness (°)	76.36	8.21	93.35	3.81
Southness (°)	109.48	11.14	96.58	5.48