Dark aerobic sulfide oxidation by anoxygenic phototrophs in anoxic waters

Jasmine S. Berg[1,2]*, Petra Pjevac[3], Tobias Sommer[4], Caroline R.T. Buckner[1], Miriam Philippi[1], Philipp F. Hach[1], Manuel Liebeke[1], Moritz Holtappels[5], Francesco Danza[6,7], Mauro Tonolla[6,7], Anupam Sengupta[8,9], Carsten J. Schubert[4], Jana Milucka[1], Marcel M.M. Kuypers[1]

[1]Department of Biogeochemistry, Max Planck Institute for Marine Microbiology, 28359 Bremen, Germany
[2]Institut de Minéralogie, Physique des Matériaux et Cosmochimie, Université Pierre et Marie Curie, CNRS UMR 7590, 4 Place Jussieu, 75252 Paris Cedex 05, France
[3]Division of Microbial Ecology, Department of Microbiology and Ecosystem Science, University of Vienna, 1090 Vienna, Austria
[4]Eawag, Swiss Federal Institute of Aquatic Science and Technology, Kastanienbaum, Switzerland
[5]Alfred-Wegener-Institut, Helmholtz-Zentrum für Polar- und Meeresforschung, Am Alten Hafen 26, 27568 Bremerhaven, Germany
[6]Laboratory of Applied Microbiology (LMA), Department for Environmental Constructions and Design (DACD), University of Applied Sciences and Arts of Southern Switzerland (SUPSI), via Mirasole 22a, 6500 Bellinzona, Switzerland
[7]Microbiology Unit, Department of Botany and Plant Biology, University of Geneva, 1211 Geneva, Switzerland
[8]Physics and Materials Science Research Unit, University of Luxembourg, 162 A, Avenue de la Faencerie, L-1511 Luxembourg City, Luxembourg
[9]Institute for Environmental Engineering, Department of Civil, Environmental and Geomatic Engineering, ETH Zurich, 8093 Zurich, Switzerland.

*corresponding author: Jasmine.Berg@mnhn.fr

ORIGINALITY-SIGNIFICANCE STATEMENT

This study reveals that dark aerobic sulfide oxidation within an anoxic layer dominated by anoxygenic phototrophic bacteria in the stratified water column of Lake Cadagno is to a large extent carried out by the anoxygenic phototrophic bacterium *Chromatium okenii*. Our findings imply that aerobic metabolisms may be more prevalent in anoxic zones than previously thought. We also present an environmental metagenome-assembled genome of *Chr. okenii* which is the first genome sequence for the genus *Chromatium* and reveals new interesting physiological features of this environmentally relevant organism, including its capacity for aerobic respiration.

SUMMARY

Anoxygenic phototrophic sulfide oxidation by green and purple sulfur bacteria (PSB) plays a key role in sulfide removal from anoxic shallow sediments and stratified waters. Although some PSB can also oxidize sulfide with nitrate and oxygen, little is known about the prevalence of this chemolithotrophic lifestyle in the environment. In this study, we investigated the role of these phototrophs in light-independent sulfide removal in the chemocline of Lake Cadagno. Our temporally resolved, high-resolution chemical profiles indicated that dark sulfide oxidation was coupled to high oxygen consumption rates of ~9 μM $O_2 \cdot h^{-1}$. Single-cell analyses of lake water incubated with $^{13}CO_2$ in the dark revealed that *Chr. okenii* was to a large extent responsible for aerobic sulfide oxidation and it accounted for up to 40 % of total dark carbon fixation. The genome of *Chr. okenii* reconstructed from the Lake Cadagno metagenome confirms its capacity for microaerophilic growth and provides further insights into its metabolic capabilities. Moreover, our genomic and single-cell data indicated that other PSB grow microaerobically in these apparently anoxic waters. Altogether, our observations suggest that aerobic respiration

may not only play an underappreciated role in anoxic environments, but also that organisms

typically considered strict anaerobes may be involved.

INTRODUCTION

Anoxygenic phototrophic bacteria oxidizing sulfide and fixing CO_2 with sunlight play an

important role in the carbon and sulfur cycles of sulfidic, shallow sediments and stratified water

columns. Phototrophic sulfur bacteria, for example, are believed to be responsible for 20-85% of

the total daily carbon fixation in anoxic lakes (summarized in Cohen *et al.,* 1977). This primary

production is so important that it can control the bulk C-isotope fractionation in the water

column, generating isotopic signatures that are transported and preserved in sediments (Posth

et al., 2017). Biomass from anoxygenic phototrophs both feeds grazing zooplankton in overlying

oxic waters (Sorokin 1966) and drives sulfate reduction in anoxic waters below (Pfennig 1975).

The phototrophic sulfur bacteria also remove toxic sulfide from the water column, enabling

aerobic life at the surface while recycling sulfur compounds for sulfate reducers. While their role

in sulfide detoxification has primarily been studied in stratified lakes, there are a few examples

of marine environments such as the Black Sea (Jørgensen *et al.,* 1991; Overmann *et al.,* 1992;

Manske *et al.,* 2005; Marschall *et al.,* 2010) and the Chesapeake Bay (Findlay *et al.,* 2015) where

phototrophic sulfur bacteria also significantly impact sulfur cycling.

Anoxygenic phototrophs generally inhabit illuminated, anoxic, reducing environments which has

been attributed to the toxicity of oxygen to these bacteria, and to the competition with abiotic

reactions involving oxygen for their electron donors. Abiotic sulfide oxidation with oxygen is

kinetically hampered and thus extremely slow (on the order of days) in trace-metal poor waters (e.g.

Millero 1986; Millero et al 1987; Luther et al 2011). However, the presence of metals such as Fe, and

less importantly Mn, has been shown to catalyze the reaction between sulfide and oxygen (Vazquez

et al 1989). Nonetheless, sulfide oxidation by microorganisms possessing enzymes which have

evolved to overcome these kinetic constraints likely remains the most important sulfide removal

process in most environments (Luther et al. 2011).

In fact, some anoxygenic phototrophs have evolved the capacity for chemotrophic growth

under microoxic conditions. Whereas the green sulfur bacteria (GSB) of the *Chlorobiaceae*

family are considered strict anaerobes, members of the *Proteobacteria* collectively known as

the purple sulfur bacteria (PSB) can be facultatively microaerobic (e.g. Kampf and Pfennig, 1980;

de Witt and Van Gemerden, 1990). Both the GSB and PSB are well adapted to fluctuating

environmental conditions, synthesizing and accumulating storage compounds during periods of

substrate/nutrient excess. The anoxygenic phototrophs are known to store zero-valent sulfur

(S^0), polyphosphate, glycogen, and in the case of the PSB alone, poly-3-hydroxyalkanoates (PHA)

(Mas and Van Gemerden, 1995). The macromolecular structure and metabolism of these

compounds have been intensely studied in laboratory pure cultures in order to understand

conditions leading to their accumulation and breakdown. It has been suggested that glycogen

may play a role in energy generation under dark conditions based on observations that cultured

Chromatium sp. utilize glycogen to reduce stored sulfur, yielding sulfide and PHA (Van

Gemerden, 1968).

Here we investigated the role of anoxygenic phototrophic bacteria in dark sulfur cycling

processes in Lake Cadagno, a permanently stratified lake with high sulfate concentrations of up

to 1-2 mM in the monimolimnion. Microbial reduction of sulfate produces large amounts of

sulfide in the anoxic bottom waters (up to ~300 µM) and sediments (>500 µM) which support

dense populations of GSB and PSB in the photic zone. These bacteria heavily influence the

chemistry of the lake, forming chemocline of 1-2 meters in thickness where sulfide and oxygen

remain mostly non-detectable. The PSB *Chromatium okenii* is by far the most active of these

bacteria, having been shown to play a disproportionately large role in inorganic carbon and

ammonium assimilation despite their low abundances (<1% of total cell numbers) in the

chemocline (Musat *et al.,* 2008; Posth *et al.,* 2017). In addition to their important contribution

to light-driven sulfide oxidation, previous studies have shown that the anoxygenic phototrophic

bacteria of Lake Cadagno remain active in the dark (Musat *et al.,* 2008; Halm *et al.,* 2009; Storelli

et al., 2013). However, their mechanism of energy generation in the absence of light is not yet

clear. We therefore combined high-resolution biogeochemical profiling with metagenomic

analyses to gain an overview of possible light-independent metabolic processes impacting the

sulfur biogeochemistry of Lake Cadagno. In addition to providing insights into the metabolism of

anoxygenic phototrophic bacteria *in situ*, we present a model to explain the mechanism of dark

sulfide oxidation in the chemocline of this meromictic lake.

RESULTS & DISCUSSION

Biogeochemistry of Lake Cadagno

Lake Cadagno is characterized by an oxic mixolimnion and a sulfidic monimolimnion spatially

separated from each other by a chemocline (defined by bold contour lines in Fig. 1A) free of

detectable oxygen (detection limit 50-100 nM) and containing very little sulfide (0-5 µM). In

August 2015, oxygen disappeared just above the chemocline close to 12 m depth. The daytime

increase in oxygen concentrations between 11-12 m depth denotes net photosynthesis and the

nighttime decrease denotes net respiration (Fig. 1A). The permanent absence of oxygen in the

chemocline indicated that oxygen was consumed both in the day and the night.

Steep gradients of sulfide diffusing into the chemocline varied independently of light-dark

periods and the total sulfide concentration in the chemocline did not exceed 5 µM at any time

point. Because the lake is meromictic, these stratified conditions were also present during other

sampling years (see Fig. S2 for 2013 and 2014 profiles). In 2015, the 0.5-1 m wide chemocline

was located around 11-12 m depth, with the exact location varying over the day most likely due

to the action of internal waves (Egli *et al.,* 1998). In previous years, the chemocline was up to 2

m wide (Fig. S2) and remained completely sulfide-free in the dark. Conservative properties such

as temperature and conductivity were constant throughout the chemocline in all years sampled

(Fig. S1&2) indicating mixing of this zone (Sommer *et al.,* 2017). Flat conductivity profiles

revealed stronger mixing of the chemocline in 2013 and 2014 (Fig. S2) than in 2015 (Fig. S1)

when the region of constant conductivity was reduced or absent.

Chr. okenii was the most significant microorganism in the chemocline both in terms of biomass,

accounting for ~60-80% of total microbial biovolume (Sommer *et al*. 2017), and activity,

accounting for 30-40% of inorganic carbon fixation (Fig. S3). The cell abundances of *Chr. okenii*

in the Lake Cadagno chemocline were enumerated by flow cytometry during 2 daily cycles (Fig.

1B). Higher densities of *Chr. okenii* were found in 2014 ($10^6 \cdot ml^{-1}$) than in 2015 ($10^5 \cdot ml^{-1}$). *Chr.*

okenii is highly motile, swimming at speeds of ~27 µm·s^{-1} and has been hypothesized to drive

the convection and mixing of the chemocline (Wüest, 1994; Sommer *et al.,* 2017). *Chromatium*

are known to migrate between gradients of sulfide, light, and oxygen by photo- and chemotaxis

(Pfennig *et al.,* 1968). We observed that *Chr. okenii* were positioned between oxygen and sulfide

gradients, regardless of changes in depth or light availability (Fig. 1A,B). The hourly variations in

cell numbers were likely the result of their heterogeneous distribution across the chemocline as

Chr. okenii can swim in a horizontal direction or be displaced by compression of the water

masses by internal waves (Egli et al., 1998). Other anoxygenic phototrophs that have been

consistently detected in the chemocline include the PSB *Lamprocystis, Thiocystis* and

Thiodictyon and several GSB of the genus *Chlorobium* (Tonolla *et al.,* 1999, 2004, 2005).

Together these bacteria constituted the majority of the total phototrophic cells ($10^6 \cdot ml^{-1}$) in

2015, but they are considerably smaller in size (and thus proportion of total biovolume) than

Chr. okenii.

The oxidation of sulfide by these anoxygenic phototrophs proceeds via the formation of S^0 as an

obligate intermediate (Mas and Van Gemerden, 1995). This S^0 was measured as particulate

sulfur on 0.7 μm filters and may comprise S^0 stored intracellularly by PSB and S^0 adhering

extracellularly to GSB. The highest concentrations of S^0 (up to 45 μM; 1C) were measured in the

middle of the chemocline. It is likely that this S^0 was present in the form of both elemental S and

polysulfides formed by the reaction of free sulfide with intra- and extracellular S^0, as has

previously been suggested in other euxinic lakes (Overmann, 1997). Our analytical method for

total S^0 did not distinguish between different forms of S^0 such as cyclooctasulfur and

polysulfides. However, we could confirm the presence of polysulfides inside live *Chr. okenii* cells

in environmental samples using Raman spectroscopy. The Raman spectrum of a sulfur inclusion

from *Chr. okenii* exhibited two weak peaks at 152 and 218 and a prominent peak at 462 cm^{-1}

(Fig. S4) which is characteristic of linear polysulfide species (Janz *et al.,* 1976). The Raman peak

at ~2900 cm^{-1} corresponds to the CH_2 and CH_3 stretching vibrations (Socrates, 2004), and its co-

occurrence with polysulfide peaks support the theory that the sulfur chains in these purple

sulfur bacteria are terminated by organic end groups as reported previously (Prange *et al.,*

1999).

Over two diurnal cycles, the S^0 inventory (Fig. S5A), or the total amount of particulate S^0 in the

chemocline, was much lower than expected from the sulfide gradients and corresponding

sulfide fluxes (discussed below), suggesting that stored S^0 served only as a transient

intermediate and was rapidly oxidized to sulfate. No day-night trends in S^0 accumulation were

apparent in the chemocline. Nevertheless, the increase in the S^0 inventory at several time points

during the night was indicative of dark sulfide oxidation.

In culture, *Chromatium* spp. are known to store carbon compounds like glycogen and

polyhydroxyalkanoates (PHAs) which have been proposed to be involved in dark sulfur

metabolism (Mas and van Gemerden, 1995). We therefore quantified glycogen and PHA

abundance in biomass samples from one day/night profile of the chemocline (Fig. 2A&B). We

could not detect any PHA, but the presence of glycogen during the day and night coincided with

Chr. okenii cell numbers (Fig. 2A&B). This is consistent with previous reports of glycogen storage

and an absence of PHA in natural populations of *Chr. okenii* (Del Don *et al.,* 1994). While the

highest potential cellular glycogen content (2.38 $\cdot$ 10^{-6} µg/cell) was found at the top of the

chemocline during the day, we observed little change in the cellular glycogen content between

day and night (Fig. S6). Average potential cellular glycogen decreased from 5.50 $\cdot$ 10^{-7} µg/cell

during the day to 5.33 $\cdot$ 10^{-7} µg/cell during the night, which represents a 3% reduction in cellular

glycogen reserves. *Chr. okenii* in Lake Cadagno were previously reported to decrease their

glycogen reserves by 50 % in the dark (Del Don *et al.,* 1994), but this may have been a result of

undersampling as our time- and depth-resolved biogeochemical profiles revealed light-dark

independent variations in *Chr. okenii* cell numbers and glycogen concentrations. While it has

been demonstrated that *Chromatium* sp. in pure cultures obtain energy from the reduction of S^0

with glycogen in the dark (Van Gemerden, 1968), we could not confirm this observation for *Chr.*

okenii in situ. From our data, we conclude that storage compounds did not play a significant role

in the dark respiratory metabolism of *Chr. okenii* in the Lake Cadagno chemocline.

Sulfate was measured as the end product of sulfide oxidation, but due to the high (1-2 mM)

background sulfate concentrations, the comparably small concentration changes resulting from

sulfide oxidation processes were non-detectable. To identify regions of sulfate production in

and around the chemocline, we therefore determined deviations from the sulfate-conductivity

mixing line drawn for each profile (see Fig. S7 for details). Strong mixing of the chemocline is

expected to produce a linear relationship between sulfate and conductivity, and large

digressions from this best-fit line indicated that sulfate was produced faster than the rate of

mixing. The expected sulfate concentration could be extrapolated based on measured

conductivity, and then subtracted from the measured sulfate concentration to give excess

sulfate:

measured $[SO_4^{2-}]$ − expected $[SO_4^{2-}]$ = excess $[SO_4^{2-}]$

This excess sulfate was attributed to biological sulfate production, as abiotic sulfide oxidation

has been shown to produce mainly S^0 and thiosulfate both in laboratory solutions (Chen and

Morris 1972; Gourmelon et al 1977) and in similarly stratified natural environments (Zopfi et al

2001; Ma et al 2006). Biogenic sulfate profiles from 2015 plotted over two diurnal cycles

exhibited a peak at the top of the chemocline in the region of oxygen depletion (Fig. 1D).

Interestingly, sulfate production was observed during the night and the overlap of excess sulfate

and oxygen in these profiles indicated that sulfide may be oxidized aerobically. Daytime sulfate

production in 2014 was related to photosynthetically active radiation (PAR) intensity (Fig. S2),

suggesting that sulfide and S^0 could either have been oxidized aerobically within the chemocline

using *in situ*-produced oxygen (Milucka *et al.,* 2015) or phototrophically. The comparatively

broad biogenic sulfate peak in the 2014 night profile likely reflects the broader vertical

distribution of the *Chr. okenii* population (Fig. S2).

The sulfate excess in the chemocline is not expected to be affected by sulfate reduction as no

sulfate reduction was detected within the chemocline in 2014 or 2015. The sulfate reduction

rates measured in the sulfidic zone 1 m below the chemocline were about 235 nM·d^{-1} and 375

nM·d^{-1} in 2014 and 2015, respectively.

To quantify biological sulfide consumption over time, we calculated the total sulfide flux into

the chemocline (Fig. S5B). Assuming that phototrophic sulfide oxidation ceases in the dark,

upwards-diffusing sulfide should accumulate in the chemocline at night. The expected sulfide

accumulation was calculated based on fluxes into the layer over a 10-h night period and

compared to the actual sulfide concentration observed in the layer. From an average sulfide flux

F = 0.15 µm·cm^{-2}h^{-1} (Fig. S5B), into a well-mixed layer of thickness H = 1 m over t = 10 hours, the

resulting sulfide concentration $C = F*t/H$ should be about 15 µM in the chemocline. However,

the average sulfide measured in the layer did not exceed 3 µM (Fig. 1A), or five times less than

expected, indicating that sulfide is consumed.

We therefore partitioned the total sulfide flux into two fractions: the flux of biologically

consumed sulfide and the flux of residual sulfide in the chemocline. First, the amount of residual

sulfide was calculated at each sampling time point by integrating sulfide concentrations within

the mixed layer (Fig. S5C). The rate of sulfide accumulation was then calculated for each 4-h

sampling interval and subtracted from the total sulfide flux to give the biologically consumed

sulfide flux. The flux of sulfide consumed in the dark was in the same range as in the day (0.03

to 0.22 $\mu mol \cdot cm^{-2}h^{-1}$) and the residual sulfide flux was very small in comparison (Fig. 2C). The

observed variations did not correlate with day-night cycles and the changes of sulfide gradients

could have been induced by internal waves, as mentioned above. Together, this indicates that

sulfide oxidation continued in the dark and seemed to be related to the total sulfide flux (Fig. S5

B) rather than the presence of sunlight. For comparison, the upwards flux of sulfide in previous

years was slightly lower, or 0.011-0.024 $\mu mol \cdot cm^{-2}h^{-1}$ in 2013 and 0.032-0.072 $\mu mol \cdot cm^{-2}h^{-1}$ in

2014.

It was not possible to calculate S^0 fluxes in Lake Cadagno because S^0 is actively transported by

the motile purple sulfur bacteria during chemo- and phototaxis (Pfennig *et al.*, 1968)

independent of diffusive processes. The total (upwards and downwards) biogenic sulfate flux

(Fig. S5 D) in this region was roughly equivalent to the sulfide flux and followed a similar trend.

Overall, our high-resolution profiles revealed that sulfide in Lake Cadagno was consumed during

the day and night, but only light-dependent sulfide oxidation has thus far been recognized as a

major sulfide-removing process in the lake. In the absence of light, it is also possible that

alternative electron acceptors such as NO_x^-, $Fe^{(III)}$, $Mn^{(IV)}$, or O_2 play a role in sulfide oxidation.

Nitrate and nitrite concentrations in the Lake Cadagno chemocline are negligible (Halm *et al.*,

2009; Milucka *et al.,* 2015). High fluxes of reduced, dissolved metals (0.027 µmol Fe·cm^{-2}·d^{-1} and

Mn 0.049 µmol Mn·cm^{-2}·d^{-1}) suggest that Fe- and Mn-oxides are rapidly reduced by

microorganisms or abiotically by sulfide in the chemocline (Berg *et al.,* 2016), but re-oxidation of

Fe and Mn would ultimately depend on oxygen in the dark. We therefore considered oxygen as

the principal direct (or indirect) oxidant responsible for observed dark sulfide oxidation.

The oxygen flux into the chemocline varied slightly between 0.022-0.071 µmol·cm^{-2}h^{-1} over the

period of 48 h (Fig. 2D). Oxygen fluxes measured in 2013 and 2014 were in the same range, or

0.013-0.048 µmol·cm^{-2}h^{-1} and 0.037-0.073 µmol·cm^{-2}h^{-1}, respectively.

Sulfide oxidation can occur in two steps:

(1) $O_2 + 2HS^- + 2H^+ \rightarrow 2S^0 + 2H_2O$

(2) $2S^0 + 3O_2 + H_2O \rightarrow 4H^+ + 2SO_4^{2-}$

Assuming that all sulfide is eventually oxidized to sulfate, we related oxygen fluxes to sulfide

consumption using the 2:1 stoichiometry for complete aerobic sulfide oxidation:

(3) $2O_2 + HS^- \rightarrow 1H^+ + SO_4^{2-}$

Calculated oxygen fluxes in 2013 and 2014 were sufficient to account for all the sulfide oxidized

in the dark. In 2015, aerobic sulfide respiration could account for up to 10-50% of sulfide

oxidized during the day and 5-45% of sulfide oxidized during the night (Fig. 2D). During the day,

the remainder of sulfide oxidation could be attributed to anoxygenic photosynthesis and/or

aerobic sulfide oxidation fueled by *in situ* oxygen production by photosynthetic algae. At several

time points in the dark, however, we could not explain the disappearance of roughly 60-90% of

upwards-diffusing sulfide. We hypothesize that the missing oxygen is supplied laterally from the

turbulent transport initiated by internal wave breaking at the lake boundaries. The convection

within the chemocline may be key to the transport of oxygen and sulfide to aerobic sulfide-

oxidizing bacteria in the chemocline. A weakening of the mixing regime was observed in August

2015 (Sommer *et al.,* 2017) which may have signified a slowed transport of electron acceptors,

thus contributing to the accumulation of sulfide in the chemocline.

To determine potential O_2 consumption rates in the Lake Cadagno chemocline, we performed

incubation experiments with chemocline water by sequentially injecting dissolved O_2 and H_2S

and monitoring O_2 concentrations online (Fig. S8). Background rates of O_2 consumption prior to

H_2S addition were 0.55 - 0.57 $\mu M \cdot h^{-1}$ (Fig. S8 B). Within 5 minutes of H_2S addition, the O_2

consumption rate increased 4- to 5-fold and remained high for 15-30 minutes. The response of

the H_2S addition was highly reproducible, resulting in mean rates of 8.7 ± 1.9 $\mu M\ O_2 \cdot h^{-1}$ for five

consecutive additions (Fig. S8 B). These potential aerobic respiration rates were well within the

range of the volumetric oxygen consumption rates (4.4-14.2 $\mu M\ O_2 \cdot h^{-1}$) determined from *in situ*

fluxes over a chemocline of 0.5 m thickness.

For each injection of 1.6 $\mu M\ H_2S$, O_2 concentrations fell by 1.5 to 2.5 μM which is equivalent to a

stoichiometry of O_2:H_2S between 1:1 and 2:1. This suggests a combination of sulfide oxidation to

sulfur and to sulfate (equations 1 and 3). Based on the experimental oxygen consumption rate,

we calculated the most conservative sulfide oxidation rate (using a 2:1 ratio) to be 4.35 μM

$H_2S \cdot h^{-1}$. These rates are much higher than would be expected from abiotic sulfide oxidation

alone (Millero 1986; et al. Millero 1987) implying that microorganisms play a key role in aerobic

sulfide oxidation in Lake Cadagno.

Dark activity of Chr. okenii in Lake Cadagno

While the sulfide-oxidizing *Chr. okenii* is one of the most active microorganisms in the Lake

Cadagno chemocline in terms of C-fixation in the light (Musat et al. 2008; Fig. S3), its metabolic

activity and contribution to sulfide removal in the dark has not yet been investigated. The

capacity for microaerophilic respiration is known for a few PSB, and it is therefore possible that

Chr. okenii grows chemoautotrophically on oxygen and sulfide in the absence of light. To test

this, we incubated Lake Cadagno chemocline water with added sulfide and $^{13}CO_2$ in the dark

under anoxic, low microoxic, and medium microoxic conditions and compared the C-fixation

activity of single *Chr. okenii* cells to that under light conditions. In order to mimic the diffusive

flux of O_2 in the lake, air was injected into the headspace of the microoxic incubations to

achieve expected concentrations of 5 µM and 15 µM dissolved O_2 in the low and medium

incubations, respectively. Although bottles were shaken at the start of the incubation to

equilibrate the water and headspace and then stirred constantly to enhance diffusion of O_2 into

the water, dissolved O_2 concentrations dropped rapidly within the first hour of incubation (Fig.

S9) at the same rate (8 µM $O_2 \cdot h^{-1}$) observed in our experimental rate measurements above. For

the remainder of the incubation, dissolved O_2 remained below 15 % of the targeted

concentrations, most likely because O_2 consumption outpaced diffusion resulting in O_2

concentrations close to those observed *in situ*.

The incorporation of $^{13}CO_2$ into *Chr. okenii* cells under dark, anoxic conditions was only slightly

above background (Fig. 3A). The mean growth rate of 0.009 ± 0.007 d^{-1} could be attributed to

trace oxygen contamination or to fermentation of carbon storage compounds as reported by

Van Gemerden (1968). A few small rod-shaped cells were also enriched in ^{13}C and were likely

sulfur-disproportionating bacteria such as *Desulfocapsa* and *Desulfobulbus* which have

previously been identified in Lake Cadagno (Tonolla et al. 2000; Peduzzi et al 2003). In the

microoxic incubations, dark C-fixation activity of *Chr. okenii* increased proportionally with the

amount of O_2 added (Fig 3B&C) confirming that these bacteria obtained energy from aerobic

respiration. The mean growth rate at an expected concentration of 15 µM O_2 (0.082 ± 0.021 d^-1)

was about three times higher than at an expected concentration of 5 µM O_2 (0.025 ± 0.008 d^{-1}).

Based on these growth rates and total cell abundances, we could estimate that *Chr. okenii*

accounted for 31-42 % of the dark C-assimilation measured by bulk analyses (Fig. S3).

Clusters of cells reminiscent of the PSB *Thiodictyon* as well as a few other bacterial filaments

and rod-shaped cells that were most likely aerobic, sulfide-oxidizing bacteria were also highly

enriched in ^{13}C (Fig 3B&C). This suggests that the capacity for aerobic respiration is

taxonomically widespread in the anoxic chemocline.

Surprisingly, *Chr. okenii* growth rates in the light (0.100 ± 0.033) were only slightly higher than

under microoxic conditions (Fig 3E) and their activity represented about 25 % of the total C-

assimilation (Fig. S3). Since this incubation was performed over a full light-dark cycle, it is also

possible that activity was restricted to the 13-h light period, which would yield a higher hourly

growth rate. While it is difficult to disentangle phototrophic from chemoautotrophic carbon

fixation in this case, the presence of abundant associations of *Chr. okenii* cells with eukaryotic

algae (Fig. 3D) indicates that these PSB can also thrive off local light-driven O_2 production. We

could show that algae were photosynthetically active via the incorporation of $^{13}CO_2$ into the

eukaryotic cells (Fig. S10). Diffusion of O_2 to attached *Chr. okenii* cells may have stimulated

chemoautotrophic growth as they were also highly enriched in ^{13}C.

Chr. okenii use energy from aerobic respiration not only for anabolic processes but for motility

as well. We could confirm that *Chr. okenii* are highly motile both in the day and the night by

performing dark field video microscopy (see Movie S1 in Supplementary Materials) of

environmental samples obtained during the night and monitored in a dark room to avoid light-

induced artefacts. Although the average night time swimming speed of *Chr. okenii* (9.9 μm s^{-1};

see Fig. S11) was a third of the day time swimming speed (27 μm s^{-1}; Sommer et al. 2017), it is

clear that *Chr. okenii* remains motile even under dark conditions. This motility may in fact be key

to bridging spatially separated gradients of oxygen and sulfide on which they grow.

Metagenomic insights into the* Chr. okenii *population in Lake Cadagno

To assess the genomic potential for light-independent, aerobic sulfide oxidation by *Chr. okenii* in

Lake Cadagno, we sequenced two metagenomes, one from the chemocline and one from a

phototrophic, sulfide-oxidizing enrichment culture obtained from the lake (Table S1). From a

combined metagenomics assembly, we reconstructed a high quality (90% complete, <1%

contaminated) metagenome-assembled genome (MAG) of a PSB highly abundant in the sulfur-

oxidizing enrichment culture (Fig. S12). The recovered MAG had a low average nucleotide

identity ANI (<70%) to any sequenced *Chromatiaceae* genomes (data not shown). However, it

encoded an rRNA operon, including a complete 16S rRNA gene with 99% sequence identity to

the 16S rRNA gene of *Chr. okenii* (Imhoff *et al.,* 1998; Tonolla *et al.,* 1999), and thus likely

represents a strain of *Chr. okenii* which is the type strain of the genus *Chromatium*. At this time,

Chr. okenii has not been successfully isolated in pure culture, nor is there any published genome

available for this organism.

The key metabolic process of *Chr. okenii* in Lake Cadagno is photoautotrophic sulfur oxidation.

In accordance, the *Chr. okenii* MAG contained genes encoding for a sulfide : quinone reductase

(*sqr*) and the full genomic inventory coding for a reverse-acting dissimilatory sulfite reductase

(rDSR) pathway (Fig. 4). The operon structure of the rDSR encoding genes (*dsrABEFHCMKLJOPN*)

was identical to the operon structure in the well described PSB model organism *Allochromatium*

vinosum (Dahl *et al.,* 2005), but no *dsrR* and *dsrS* gene were found. No genes encoding for sulfur

oxidation via the SOX pathway, or homologues of sulfur globule proteins (*sgpABC*) typically

found in PSB were detected in the draft genome. In line with its phototrophic metabolism, the

Chr. okenii MAG showed the genomic potential for photosynthesis, with the genes encoding for

a light harvesting complex 1 (*pufAB*) and a PSB-type photosynthetic reaction center (*pufLMC*)

encoded in a single operon. Furthermore, the full genomic repertoire for a NADP-Me type C4

photosynthetic carbon assimilation cycle, and all genes (with exception of *cbbS* encoding for the

small subunit of the ribulose-1,5-bisphosphate carboxylase/oxygenase) necessary for CO_2

assimilation via the Calvin-Benson-Bassham (CBB) Cycle were present (Fig. 4).

Many *Chromatiaceae* can grow chemoautotrophically, respiring oxygen under microoxic

conditions (Kämpf and Pfennig, 1980). Cytochrome (Cyt) *c*-containing oxidases (e.g. Cyt *aa3*, Cyt

cbb3) were not found in the *Chr. okenii* MAG. However, a Cyt *bd* type ubiqinol oxidase, known

to function as sulfide-resistant O_2-accepting oxidase in other *Gammaproteobacteria* (Forte *et*

al., 2016), was identified (Fig. 4). Further, a plethora of genes related to heme *b* (*gltX*,

hemALBCD, and *hemH*) and siroheme (*cysG*) synthesis, degradation (a heme oxygenase) and

export (ABC-type heme exporter, *ccmABCD*), as well as hemerythrin-like metal binding proteins

were encoded. Hemerythrin has been implicated in binding of oxygen for delivery to oxygen-

requiring enzymes, for detoxification, or for oxygen sensing in motile, microaerobic prokaryotes

(French *et al.,* 2007). The presence of these oxygen-dependent enzymes, as well as a key

oxidative stress defense enzyme superoxide dismutase (SOD), support the idea that *Chr. okenii*

may be facultatively microaerobic. A complete set of genes for flagellar biosynthesis

(*fliDEGHJKLMNOPQRW*, *flgABCDEFGHIK*, *flhAB*) and flagellar motor proteins (*motAB*) confer

motility to this bacterium.

Several other genes revealed interesting metabolic capacities of *Chr. okenii.* A cytosolic

bidirectional [NiFe] type 3d hydrogenase and a nitrogenase were encoded in the MAG (Fig. 4),

implicating the potential for involvement of *Chr. okenii* in nitrogen fixation and hydrogen

oxidation which has previously been overlooked. Additionally, the *Chr. okenii* MAG encoded a

glycogen synthase and a glycogen debranching enzyme, as well as the full genomic repertoire

necessary for polyhydroxyalkanoate (PHA) biosynthesis. This is consistent with the detection of

glycogen in our biogeochemical profiles of the chemocline. Finally, it is possible that novel

terminal oxidases are among the hypothetical genes that could not be assigned any known

function.

CONCLUSIONS

It is intriguing that oxygen should play a major role in sulfide oxidation in the ostensibly anoxic

chemocline of Lake Cadagno, especially by purple sulfur bacteria generally thought to lead an

anaerobic lifestyle. To explain the coupling of oxygen and sulfide consumption in the oxygen-

and sulfide-free chemocline of Lake Cadagno, we sketched a diagram of the transport processes

likely driving biological activity there (Fig. 5). As described in Sommer *et al.,* (2017), active

convection of the chemocline can be driven by the formation of sinking bacterial plumes.

Combined with turbulence induced by the breaking of internal waves at the sides of the lake

basin, these convective currents may entrain sulfide and oxygen at the boundaries of the

chemocline and fuel populations of sulfide-oxidizing *Chr. okenii* there.

Sulfur-oxidizing bacteria have previously been reported to bridge distances between pools of

electron donors and acceptors by intracellularly storing and transporting S^0 and NO_3^- between

redox zones (Fossing *et al.,* 1995; Jørgensen and Gallardo, 1999) and even by transferring

electrons along nanowires (Pfeffer *et al.,* 2012), but aerobic sulfide oxidation in Lake Cadagno

represents a new mechanism of electron acceptor/donor coupling across large distances. After

entrainment into the chemocline, dissolved oxygen and sulfide are consumed so rapidly that

they remain below detection limits. The physical and biological processes described here may

therefore provide clues to sulfide oxidation in other anoxic environments such as the Black Sea

where the mechanism of sulfide removal is not completely understood. Clearly, the biochemical

limits to oxygen utilization are far below current definitions of anoxia and demonstrate that

aerobic respiration is possible in so-called "anoxic" lacustrine (Milucka *et al.,* 2015) and marine

(Garcia-Robledo *et al.,* 2017) waters.

Although purple sulfur bacteria are generally considered anaerobes, we propose that *Chr. okenii*

in the Lake Cadagno chemocline aerobically respire upwards-diffusing sulfide in addition to

performing phototrophic sulfide oxidation as previously reported (e.g. Tonolla et al. 1999;

Musat et al 2008; Storelli et al. 2013). Our direct evidence for oxygen-driven C-fixation and

continued motility of *Chr. okenii* in darkness is also supported by the presence of a potential

high-affinity oxidase in the metagenome-assembled genome. In contrast to observations from

laboratory cultures, *Chr. okenii* may have a very different metabolism in the environment where

high fluxes of substrates rather than absolute substrate concentrations fuel microbial activity.

Our nanoSIMS measurements revealed that in addition to *Chr. okenii*, a number of phototrophic

and non-phototrophic cells are also highly active under aerobic, sulfide-oxidizing conditions.

Based on the diversity of genes encoding for high-O_2 affinity terminal oxidases found in our Lake

Cadagno metagenome, it appears that some other purple sulfur bacteria as well as a handful of

Gamma- and *Epsilonproteobacteria* might aerobically oxidize sulfide (Fig. S13). Interestingly,

two species of strictly anaerobic green sulfur bacteria from Lake Cadagno also possessed genes

encoding for terminal oxidases, though further study will be required to elucidate the role of

these different bacterial populations in chemotrophic sulfide oxidation.

EXPERIMENTAL PROCEDURES

Sampling

The meromictic Lake Cadagno is situated in the Piora Valley in the Swiss Alps at an altitude of

1921 m. Data presented here were collected during field campaigns in September 2013, August

2014, June 2015 August 2015, and August 2018. In 2013 and 2014 *in situ* measurements were

performed with a profiling ion analyzer (PIA; see Kirf *et al.,* 2014 for description) lowered from a

platform anchored at the deepest part of the lake (20.7 m). Conductivity, turbidity, depth

(pressure), temperature and pH were measured with a multi-parameter probe (XRX 620, RBR).

Dissolved oxygen was recorded online with a type PSt1 normal (detection limit 125 nM) micro-

optode and a type TOS7 trace (reliable detection limit 50-100 nM) micro-optode (PreSens). The

oxygen sensors were calibrated by parallel Winkler titrations. Water samples for chemical

analyses and cell counts were collected with a rosette syringe sampler equipped with twelve 60-

ml syringes triggered online at selected depths. Due to a technical failure of the PIA, the 6 AM

profile in August 2014 and all subsequent profiles were measured with an SBE 19 plus V2 CTD

probe (Sea-Bird Electronics) equipped with sensors for pressure, temperature and conductivity,

and with additional sensors for turbidity (WET Labs Eco), oxygen (SBE 43), pH (18-I) and two

fluorescence wavelengths (WET Labs ECO-AFL). The detection limit of the SBE 43 oxygen probe

was about 1 µmol/l. In parallel with *in situ* measurements, water for chemical analyses was

pumped to the surface through Neoprene tubing attached to the CTD and filled into 60-ml

syringes (rinsed 2 X with *in situ* water) on board. Two parallel metal plates of diameter ~15 cm

attached to the submersed end of the tubing served to channel water horizontally, resulting in

more discrete vertical profiling.

Water samples in syringes were aliquoted on board immediately after collection. Samples for

sulfate analyses were filtered (0.22 µm pore size) directly into sterile Eppendorf vials. Sulfide

samples were fixed with Zn-acetate to a final concentration of 0.1 % (w/v). Biomass was

concentrated onto GF/F filters (Whatman) and stored at -20°C for analyses of intracellularly

stored elemental sulfur and organic carbon compounds. Filtrate (0.22 µm pore size) was also

collected and frozen at -20°C for metabolome analysis of dissolved compounds. Water for

dissolved inorganic carbon (DIC) analysis was filled into 12-ml Exetainers (Labco Limited) and

capped before injecting 100 µl of saturated $HgCl_2$ solution to terminate microbial activity.

Biomass samples for determination of background ^{13}C in the particulate organic carbon (POC)

were concentrated onto pre-combusted (6 h at 600 °C) GF/F filters. Samples for DNA analysis

were collected from the chemocline in August 2014 by concentrating microbial cells on 0.22 µm

GTTP-type polycarbonate filters (Millipore) on site and freezing at -20°C until further processing.

Additional water for cultivation and motility experiments was pumped directly from the

chemocline into 1-L Duran bottles or 250-ml serum bottles and sealed with butyl rubber

stoppers without a headspace to maintain anoxic conditions.

Chemical Analyses

Sulfide was measured using the colorimetric method of Cline (1969). Sulfate was measured on a

761 Compact ion chromatograph (Metrohm) equipped with a Metrosep A SUPP 5 column.

Intracellular sulfur on filters was extracted by sonication in methanol for 15 min in an ice bath.

Samples were analyzed on an Acquity H-Class UPLC system (Waters Corporation) with an

Acquity UPLC BEH C18 column coupled to a photodiode array (PDA) detector using UPLC-grade

methanol as eluent. Data was acquired and processed using the Empower III software.

Intracellular glycogen was analyzed following the procedures of the assay kit (MAK016, Sigma

Aldrich). Briefly, cells were extracted by scraping them from GF/F filters and homogenizing in

200 µL extraction buffer and centrifuged two times to clear the supernatant. The supernatant

was analyzed fluorometrically after incubation with enzyme mix and fluorescent peroxidase

substrate. Intracellular PHA was analyzed using the protocol from Braunegg *et al*., (1978).

Hydrolyzation of the polymer and conversion to a methyl-ester of the monomeric

hydroxyalkanoate fraction was done in acidified alcohol solution (6% H_2SO_4 in methanol) and

chloroform under heating (100°C, 2 h). After addition of water and phase separation the organic

phase was analyzed with GC-MS (Agilent 7890B GC connected to Agilent 5977A MSD) to detect

the methylhydroxyalkanoates using the following settings: Agilent 30 m DB-5-MS column,

splitless injection of 1 µl, temperature program was 50°C for 1 min than heating 10°C/min until

120°C followed by 45°C/min until 320°C and hold for 5 minutes. Benzoic acid was used as

internal standard in each sample and quantification was done with pure polyhydroxybutyrate

standard (Sigma Aldrich).

Sulfate reduction rates were measured by adding the radiotracer $^{35}SO_4^{2-}$ (5 MBq) to anoxic lake

water in 50-ml glass syringes and incubated in the dark. A solution of unlabeled Na_2S was added

to a final concentration of ~50 $\mu mol \cdot l^{-1}$ as a background sulfide pool in case of sulfide re-

oxidation. At each sampling point, 10 ml of sample was dispensed into 5 ml of 20% (w/v) Zn-

acetate. Reduced sulfur species (e.g. sulfur and sulfide as ZnS) were separated out via the

chromium distillation method described in Kallmeyer *et al.,* (2004) and the radioactivity per

sample was determined via scintillation counting (Packard 2500 TR).

Confocal Raman spectroscopy

In a glove box under 90:10 N_2-CO_2 atmosphere, a drop of fresh sample from the chemocline was

mounted between two glass coverslips and sealed with electrical tape to prevent contact with

air. A polysulfide solution containing 5.06 g $Na_2S \cdot 9H_2O$ and 5.8 g elemental sulfur per 100 ml

H_2O, with a final pH of 9.5 and sulfide concentration of 210 mM, was used as reference.

Measurements were conducted with an NTEGRA Spectra confocal spectrometer (NT-MDT)

coupled to an inverted Olympus IX71 microscope. The excitation light from a 532-nm solid-state

laser was focused on the sample through an Olympus 100X (numerical aperture [NA], 1.3) oil

immersion objective. Raman scattered light was collected by an electron-multiplying charge-

coupled device (EMCCD) camera (Andor Technology) cooled to -70°C. Spectra were recorded

between 0 and 4,500 cm^{-1} with a spectral resolution of 0.2 cm^{-1} and analyzed with the NT-MDT

software Nova_Px 3.1.0.0.

Flux Calculations

Turbulent fluxes (J) of sulfide, sulfur, sulfate, and oxygen at the chemocline were calculated

assuming steady state by applying Fick's first law: $J=-D\partial C/\partial x$. For sulfide, sulfate, and oxygen we

used the turbulent diffusion coefficient (D) of 1.6×10^{-6} m^2 s^{-1} from (Wüest, 1994)

corresponding to turbulence at the Lake Cadagno chemocline boundaries. For sulfur gradients

within the well-mixed chemocline the coefficient $D = 1.5 \times 10^{-5}$ m^2 s^{-1} from Wüest (1994) was

used. The change in concentration (∂C) was computed for each species over the depths with the

steepest gradients. Oxygen and sulfide fluxes were determined for the regions immediately

above and below the chemocline, defined as the zone of constant conductivity.**Potential oxygen**

consumption rate measurements

Potential O_2 consumption rate measurements were conducted as described in Holtappels et al.

(2014). Incubations were performed in 250 ml Winkler bottles using trace oxygen sensor spots

(Ø5mm, TROXSP5, Pyroscience) glued to the inner glass wall and read from the outside using a

4-channel fiber-optic oxygen meter (FireStingO2, Pyroscience) recording O_2 concentration every

10 seconds. The glass stopper of the Winkler bottle was pierced with a glass tube (11 cm long,

inner diameter 1.5 mm), to allow for pressure compensation due to temperature-induced

volume changes and to facilitate the addition of small amounts of reagents. Oxygen transport

through this opening was limited by diffusion through a narrow passage over a long distance.

Tests with degassed sterile water showed negligible O_2 concentration changes of 3-25 nM·d^{-1}

(figures available on request). Water inside the incubation bottles was gently mixed by small

glass-coated magnets driven by an adjacent magnetic stirrer. The incubation bottles were

placed in a temperature-controlled water bath set to 4.8°C (+/- 0.1°C). A two-point calibration of

the trace optode spots was performed by using degassed water to which a known amount of

O_2-saturated water was added through the glass tube.

Anoxic chemocline water was pumped onboard a sampling platform and distributed directly

into incubation vessels via Neoprene tubing, avoiding atmospheric contamination as described

above. Bottles were placed immediately into the water bath and incubated in the dark for 18

hours. Initial O_2 values were between 6 and 8 µM. Over a period of 12 hours, H_2S was added 5

times by injecting up to 300 µl of a 4 mM Na_2S stock solution to a final concentration of 1.6 µM.

Between 4-6 ml O_2-saturated lake water from the same depth was periodically added to

maintain O_2 concentrations between 5 and 11 µM. O_2 consumption rates were calculated from

linear regressions over 10-30 min time intervals.

Stable isotope incubations

250-ml serum bottles containing Lake Cadagno chemocline water collected in August 2018 were

bubbled for 10 min with N_2 to remove any contaminant oxygen. Four different incubation

conditions were set up to test microbial carbon fixation under the following conditions: no O_2 +

dark, no O_2 + light, medium O_2 + dark, and high O_2 + dark. To each of the dark incubations, 50 ml

of water was removed and replaced with N_2 gas. Solutions of Na_2S and ^{13}C-DIC were added to

final concentrations of 25 µM and 1 mM (36-39 atom %), respectively, except for the light

incubation which received 200 µM (8 atom %) ^{13}C-DIC. Volumes of 0.918 ml and 2.8 ml air were

added to the dark, aerobic incubations to achieve headspace O_2 concentrations of 129 and 388

µM, respectively, resulting in expected dissolved O_2 concentrations of 5 and 15 µM,

respectively, at the start of the incubations (Garcia and Gordon, 1992). All bottles were gently

shaken for several minutes to equilibrate the water and headspace and then placed in a 10°C

water bath in the dark for 18 h, except for the light condition which was incubated separately

under natural light, 0.1 − 2 % incident daylight, for a full light-dark cycle. Equilibration with the

headspace was enhanced throughout the incubation by stirring with glass-coated magnetic stir-

bars at minimum speed.

In order to verify the consumption of oxygen, two bottles prepared without ^{13}C-DIC and injected

with 0.918 ml and 2.8 ml of air were incubated in parallel with needle optodes (FirestingO2,

Pyroscience) immersed in the liquid phase. Dissolved O_2 concentrations were recorded every 3

min and are shown in Fig. S9.

At the end of the experiment, bottles were sacrificed for analysis of bulk and single-cell ^{13}C

uptake. Water for ^{13}C-DIC and ^{13}C-POC was sampled as described above. For nanoSIMS, water

samples were fixed with formaldehyde (2 % v/v) overnight at 4°C before filtering onto 0.2 μm

Au/Pd-sputtered polycarbonate filters (Millipore) and stored at -20°C until processing.

Mass spectrometry isotope uptake analysis

The labeling % of ^{13}C-DIC was determined by diluting Hg-killed water 1:10 in MilliQ and

transferring 3 ml into a 12 ml Exetainer with an N_2 headspace. Samples were then acidified with

the addition of ~100 μl concentrated H_3PO_4 so the outgassed $^{13}CO_2$ could be analyzed by cavity

ring-down spectroscopy (G2201-i Picarro).

The incorporation of ^{13}C from $^{13}CO_2$ into biomass was measured by combustion of the

particulate organic carbon fraction on GF/F filters. Filters were decalcified by incubation with

37% fuming HCl overnight, dried in an oven at 60°C, and then packed into tin capsules for

combustion analysis. The C isotopic composition was determined on an automated elemental

analyzer (Thermo Flash EA, 1112 Series) coupled to a Delta Plus XP IRMS (Thermo Finnigan).

Nanometer-scale secondary ion mass-spectrometry

Circular pieces (5-mm diameter) were punched out of the Au/Pd-sputtered filter and regions of

interest were marked with a laser micro-dissection microscope (LMD) (DM 6000 B, Leica

Microsystems) before analysis with the nanoSIMS 50L (Cameca) at the Max Planck Institute for

Marine Microbiology in Bremen, Germany. For each field of view, the sample was pre-sputtered

with a primary Cs^+ beam of 300 pA, and then measured by rastering a primary Cs^+ beam with a

diameter of < 100 nm and a beam current between 1.0 and 1.5 pA over the area. Secondary ion

images of $12C^-$, $13C^-$, $12C14N^-$, $31P^-$ and $32S^-$ were recorded simultaneously using 5 detectors

together with the secondary electron (SE) image ranging from 10 x 10 μm to 50 x 50 μm in size

and corresponding to an image size of 256 x 256 and 512 x 512 pixels, respectively, using a dwell

time of 1 ms per pixel. Up to 60 planes were recorded for each imaged area.

Images were processed using the Matlab-based Look@NanoSIMS software (Polerecky *et al.,*

2012). Regions of interest (ROIs) were drawn around cells based on the $12C14N^-$ signal overlain

on SE images and *Chr. okenii* cells were identified based on morphology, size, and pigmentation

in parallel light-microscope images. Cellular ^{13}C atm. % was calculated from $^{13}C/^{12}C$ values of

individual ROIs. The background (cell-free polycarbonate filter) ^{13}C content was also evaluated

in every field of view for comparison. Growth rates were calculated assuming a linear growth

rate where one cell division results in a maximum of ½ the labeling percentage:

$g = (atm\ \%\ ^{13}C\ excess_{cells}/\ ½\ atm\ \%\ ^{13}C\ excess_{tot\ dissolved})\ /\ t_{hours}$

Cellular C-uptakes rates were calculated based on the relationship between biovolume and C-

content from Verity *et al.*, (1992), assuming an average biovolume of spherocylinders cells of 65

μm^3 based on direct measurements of 70 *Chr. okenii* cells.

. **Motility analysis**

Water samples containing *Chr. okenii* cells were collected under anoxic conditions from the

chemocline during the night, protected from artificial light with aluminum foil, and analyzed

immediately on site. Motile cells were transferred via a degassed glass syringe to a sealed

rectangular millimetric chamber (dimensions 20 mm × 10 mm × 2 mm) prepared using glass

slides separated by a 2-mm thick spacer, which provided an anoxic environment during motility

characterization. Experiments were conducted in a dark room, and imaging was performed

using the dark field microscopy mode at 25 fps, with the lowest intensity illumination. No

transient response was observed right at the start of the imaging, and the swimming velocity

remained steady throughout the duration of the measurements. This is in contrast to swimming

behavior at higher light intensities where the swimming cells exhibited a positive phototactic

response (Sommer *et al.*, 2017). We could therefore rule out a light-induced effect on motility at

the minimum illumination level used for our measurements. Videos of swimming cells were

acquired and subsequently analyzed using the ImageJ Particle Tracker routine to obtain the

coordinates of the cells (geometric centers) at each time interval. These were used to calculate

the swimming speeds and extract the trajectories of individual cells.

DNA extraction, sequencing, and analysis

Two metagenomes were sequenced: one from the chemocline and one from a phototrophic,

sulfide-oxidizing enrichment from the lake. The enrichment culture was obtained by preparing

agar-stabilized sulfide-gradient tubes as described in Schwedt et al. (2012) using sterile-filtered

Lake Cadagno water amended with vitamins and trace elements as described for cultivation of

purple sulfur bacteria (Eichler and Pfennig, 1988). After inoculation with water from the

chemocline, cultures were incubated for approximately 3 months in indirect light at $15^{\circ}C$ until

development of a deep purple color indicative of growth of PSB.

DNA was extracted from polycarbonate filters with the Ultra Clean MoBio PowerSoil DNA kit

(MoBio Laboratories, Carlsbad, USA) according to the manufacturer's protocol with the

following modification: the bead beating step was reduced to 30 sec followed by incubation on

ice for 30 sec, repeated 4x. The DNA was gel-purified using SYBR Green I Nucleic Acid Gel Stain

(Invitrogen) and the QIAquick Gel Extraction Kit (Qiagen) according to the accompanying

protocols. DNA concentration was determined fluorometrically at 260 nm, using the Qubit 2.0

Fluorometer and the Qubit dsDNA HS Assay KIT (Invitrogen) and sent to the Max Planck-

Genome Centre (Cologne, Germany) for sequencing. The metagenome was sequenced (100 bp

paired end reads) by Illumina HiSeq (Illumina Inc., USA) sequencing following a TruSeq library

preparation. Metagenomic reads were adapter- and quality-trimmed (phred score 15, bbduk

function of the BBMap package, https://sourceforge.net/projects/bbmap/) and paired-end

reads were *de novo* assembled with the uneven depth assembler IDBA-UD (Peng *et al.,* 2012).

The metagenome assembly was binned based on tetranucleotide frequencies, differential

coverage, taxonomic classification, and conserved single-copy gene profiles with the Metawatt

binning software (version 3.5.2; Strous *et al.,* 2012). The completeness and contamination of the

binned MAGs was evaluated with CheckM (Parks et al 2014). The bulk metagenome and the

MAG identified as *Chr. okenii* were automatically annotated in IMG (Markowitz et al 2011), and

the *Chr. okenii* MAG was manually screened for the presence of genes of interest to this study.

Assembled data is available in IMG, under the IMG genome IDs 3300010965 (bulk assembly) and

2700988602 (*Chr. okenii* MAG).

Cytochrome *c* and quinol oxidase subunit I and the cytochrome *bd* oxidase subunit I proteins

were extracted from the metagenome via Pfam motif search (PF00115 and PF01654,

respectively), and manually inspected to exclude NO reductase subunit I (NorB) proteins and

proteins shorter than 100 aa prior to downstream analysis. A total of 130 out of 178 identified

cytochrome *c* and quinol oxidase subunit I proteins and 76 out of 82 recovered cytochrome *bd*

oxidase subunit I proteins passed the manual curation step. For phylogenetic tree

reconstruction, the manually vetted protein sequences and a selection of reference sequences

recovered from NCBI were aligned using MAFFT (Katoh *et al.,* 2002) and IQ-TREE was used to

construct the phylogenetic tree with automated best model prediction and a total of 1000

bootstrap replicates (Trifinopoulos *et al.,* 2016; Hoang *et al.,* 2017). Thereafter, a selection of

sequences closely related to known sulfur oxidizing microorganisms alongside the respective

reference sequences were used for reconstruction of the phylogenetic trees presented in Fig.

S11. The output files were uploaded to iTOL for visualization (Letunic *et al.,*

2016).**ACKGNOWLEDGEMENTS**

We are grateful to the 2014 and 2015 Cadagno Field Expedition Teams from EAWAG and MPI

Bremen for assistance in the field, and to the Alpine Biology Center Foundation (Switzerland) for

use of its research facilities. We would especially like to thank Abiel Kidane and Sten Littman for

help with nanoSIMS measurements as well as Nadine Rujanski, Gabriele Klockgether, and Gaute

Lavik for mass spectrometry analyses. Funding was provided by the International Max Planck

Research School of Marine Microbiology, the Max Planck Society, and the Deutsche

Forschungsgemeinschaft (through the MARUM Center for Marine Environmental Sciences). A.S.

was supported by the Human Frontier Science Program (Cross Disciplinary Fellowship,

LT000993/2014-C).

REFERENCES

Berg, J.S., Schwedt, A., Kreutzmann, A.-C., Kuypers, M.M., and Milucka, J. (2014) Polysulfides as
Intermediates in the Oxidation of Sulfide to Sulfate by Beggiatoa spp. *Applied and Environmental*
Microbiology **80**: 629-636.

Braunegg, G., Sonnleitner, B. Y., and Lafferty, R. M. (1978). A rapid gas chromatographic method for the
determination of poly-β-hydroxybutyric acid in microbial biomass. *Applied Microbiology and*
Biotechnology **6:** 29-37.

Chen, K. Y., & Morris, J. C. (1972). Kinetics of oxidation of aqueous sulfide by oxygen. *Environmental*
Science & Technology **6:** 529-537.

Cline, J.D. (1969) Spectrophotometric determination of hydrogen sulfide in natural waters. *Limnology*
and Oceanography **14**: 454-458.

Dahl, C., Engels, S., Pott-Sperling, A. S., Schulte, A., Sander, J., Lübbe, Y., ... and Brune, D. C. (2005). Novel
genes of the dsr gene cluster and evidence for close interaction of Dsr proteins during sulfur oxidation in
the phototrophic sulfur bacterium *Allochromatium vinosum*. *Journal of Bacteriology* **187**: 1392-1404.

Dahl, T.W., Anbar, A.D., Gordon, G.W., Rosing, M.T., Frei, R., and Canfield, D.E. (2010) The behavior of
molybdenum and its isotopes across the chemocline and in the sediments of sulfidic Lake Cadagno,
Switzerland. *Geochimica et Cosmochimica Acta* **74**: 144-163.

De Witt, R., and Van Gemerden, H. (1990) Growth of the phototrophic purple sulfur bacterium
Thiocapsaroseopersicina under oxic/anoxic regimens in the light. *FEMS Microbiology Ecology* **6**: 69-76.

Del Don, Chr., Hanselmann, K.W., Peduzzi, R., and Bachofen, R. (1994) Biomass composition and
methods for the determination of metabolic reserve polymers in phototrophic sulfur bacteria. *Aquatic*
Sciences **56**: 1-15.

Egli, K., Wiggli, M., Klug, J., and Bachofen, R. (1998) Spatial and temporal dynamics of the cell density in a
plume of phototrophic microorganisms in their natural environment. *Doc Ist Ital Idrobiol* **63**: 121-126.

Eichler, B., & Pfennig, N. (1988). A new purple sulfur bacterium from stratified freshwater lakes,
Amoebobacter purpureus sp. nov. *Archives of Microbiology* **149:** 395-400.

Findlay, A.J., Bennett, A.J., Hanson, T.E., and Luther, G.W. (2015) Light-dependent sulfide oxidation in the
anoxic zone of the Chesapeake Bay can be explained by small populations of phototrophic bacteria.
Applied and Environmental Microbiology **81**: 7560-7569.

Forte, E., Borisov, V. B., Falabella, M., Colaço, H. G., Tinajero-Trejo, M., Poole, R. K., ... and Giuffrè, A.
(2016). The terminal oxidase cytochrome bd promotes sulfide-resistant bacterial respiration and growth.
Scientific Reports **6:** 23788.

Fossing, H., Gallardoi, V., Jorgensen, B., Hiittel, M., Nielsenl, L., Schulz, H. *et al.,* (1995) Concentration and
transport of nitrate by the mat-forming sulphur bacterium Thioploca. *Nature* **374**: 20.

French, C. E., Bell, J. M., and Ward, F. B. (2007). Diversity and distribution of hemerythrin-like proteins in
prokaryotes. *FEMS Microbiology Letters* **279**: 131-145.

Garcia, H. E., & Gordon, L. I. (1992). Oxygen solubility in seawater: Better fitting equations. *Limnology*
and Oceanography **37** 1307-1312.

Garcia-Robledo, E., Padilla, C. C., Aldunate, M., Stewart, F. J., Ulloa, O., Paulmier, A., ... and Revsbech, N.
P. (2017). Cryptic oxygen cycling in anoxic marine zones. *Proceedings of the National Academy of*
Sciences 201619844.

Gourmelon, C., Boulegue, J., & Michard, G. (1977). Oxydation partielle de l'hydrogène sulfuré en phase
aqueuse. *CR Acad Sci Paris* **284**: 269-272.

Halm, H., Musat, N., Lam, P., Langlois, R., Musat, F., Peduzzi, S. *et al.,* (2009) Co-occurrence of
denitrification and nitrogen fixation in a meromictic lake, Lake Cadagno (Switzerland). *Environmental*
Microbiology **11**: 1945-1958.

Hoang, D. T., Chernomor, O., von Haeseler, A., Minh, B. Q., & Vinh, L. S. (2017). UFBoot2: improving the
ultrafast bootstrap approximation. Molecular Biology and Evolution **35**: 518-522.

Holtappels, M., Tiano, L., Kalvelage, T., Lavik, G., Revsbech, N. P., & Kuypers, M. M. (2014). Aquatic
respiration rate measurements at low oxygen concentrations. *PLoS One* **9**: e89369.

Imhoff, J. F., Süling, J., and Petri, R. (1998). Phylogenetic relationships among the Chromatiaceae, their
taxonomic reclassification and description of the new genera *Allochromatium*, *Halochromatium*,
Isochromatium, *Marichromatium*, *Thiococcus*, *Thiohalocapsa* and *Thermochromatium*. *International*
Journal of Systematic and Evolutionary Microbiology **48**: 1129-1143.

Janz, G., Downey Jr, J., Roduner, E., Wasilczyk, G., Coutts, J., and Eluard, A. (1976) Raman studies of
sulfur-containing anions in inorganic polysulfides. Sodium polysulfides. *Inorganic Chemistry* **15**: 1759-
1763.

Jørgensen, B.B., Fossing, H., Wirsen, Chr.O., and Jannasch, H.W. (1991) Sulfide oxidation in the anoxic
Black Sea chemocline. *Deep Sea Research Part A, Oceanographic Research Papers* **38**: S1083-S1103.

Jørgensen, B.B., and Gallardo, V.A. (1999) Thioploca spp.: filamentous sulfur bacteria with nitrate
vacuoles. *FEMS Microbiology Ecology* **28**: 301-313.

Kallmeyer, J., Ferdelman, T.G., Weber, A., Fossing, H., and Jørgensen, B.B. (2004) A cold chromium
distillation procedure for radiolabeled sulfide applied to sulfate reduction measurements. *Limnology and*
Oceanography Methods **2**: 171-180.

Kampf, Chr., and Pfennig, N. (1980) Capacity of Chromatiaceae for chemotrophic growth. Specific
respiration rates of Thiocystis violacea and Chromatium vinosum. *Archives of Microbiology* **127**: 125-135.

Katoh, K., Misawa, K., Kuma, K. I., & Miyata, T. (2002). MAFFT: a novel method for rapid multiple
sequence alignment based on fast Fourier transform. *Nucleic Acids Research* **30:** 3059-3066.

Kirf, M.K., Dinkel, Chr., Schubert, Chr.J., and Wehrli, B. (2014) Submicromolar oxygen profiles at the oxic–
anoxic boundary of temperate lakes. *Aquatic Geochemistry* **20**: 39-57.

Letunic, I., & Bork, P. (2016). Interactive tree of life (iTOL) v3: an online tool for the display and
annotation of phylogenetic and other trees. Nucleic Acids Research **44**: W242-W245.

Luther, G. W., Findlay, A. J., MacDonald, D. J., Owings, S. M., Hanson, T. E., Beinart, R. A., & Girguis, P. R.
(2011). Thermodynamics and kinetics of sulfide oxidation by oxygen: a look at inorganically controlled
reactions and biologically mediated processes in the environment. *Frontiers in Microbiology* **2**: 62.

Ma, S., Noble, A., Butcher, D., Trouwborst, R. E., and Luther III, G. W. (2006). Removal of H2S via an iron
catalytic cycle and iron sulfide precipitation in the water column of dead end tributaries. *Estuarine,*
Coastal and Shelf Science **70**: 461-472.

Manske, A. K., Glaeser, J., Kuypers, M. M., and Overmann, J. (2005). Physiology and phylogeny of green
sulfur bacteria forming a monospecific phototrophic assemblage at a depth of 100 meters in the Black
Sea. *Applied and Environmental Microbiology* **71**: 8049-8060.

Markowitz, V.M., Chen, I.M.A., Palaniappan, K., Chu, K., Szeto, E., Grechkin, Y., Ratner, A., Jacob, B.,
Huang, J., Williams, P. and Huntemann, M., (2011). IMG: the integrated microbial genomes database and
comparative analysis system. *Nucleic Acids Research*, *40*(D1), pp.D115-D122.

Marschall, E., Jogler, M., Henßge, U., and Overmann, J. (2010). Large-scale distribution and activity
patterns of an extremely low-light-adapted population of green sulfur bacteria in the Black Sea.
Environmental Microbiology **12**: 1348-1362.

Mas, J., and Van Gemerden, H. (1995) Storage products in purple and green sulfur bacteria. In
Anoxygenic Photosynthetic Bacteria: Springer, pp. 973-990.

Millero, F. J. (1986). The thermodynamics and kinetics of the hydrogen sulfide system in natural waters.
Marine Chemistry, **18**: 121-147.

Millero, F. J., Hubinger, S., Fernandez, M., & Garnett, S. (1987). Oxidation of H_2S in seawater as a function
of temperature, pH, and ionic strength. *Environmental Science & Technology* **21**: 439-443.

Milucka, J., Kirf, M., Lu, L., Krupke, A., Lam, P., Littmann, S. *et al.,* (2015) Methane oxidation coupled to
oxygenic photosynthesis in anoxic waters. *The ISME Journal*.

Musat, N., Halm, H., Winterholler, B., Hoppe, P., Peduzzi, S., Hillion, F. *et al.,* (2008) A single-cell view on
the ecophysiology of anaerobic phototrophic bacteria. *Proceedings of the National Academy of Sciences*
105: 17861-17866.

Overmann, J., Cypionka, H., and Pfennig, N. (1992). An extremely low-light adapted phototrophic sulfur
bacterium from the Black Sea. *Limnology and Oceanography* **37**: 150-155.

Overmann, J. (1997) Mahoney Lake: a case study of the ecological significance of phototrophic sulfur
bacteria. In *Advances in Microbial Ecology*: Springer, pp. 251-288.

Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2014. Assessing the quality of microbial
genomes recovered from isolates, single cells, and metagenomes. *Genome Research* **25**: 1043-1055

Peduzzi, S., Tonolla, M., & Hahn, D. (2003). Vertical distribution of sulfate-reducing bacteria in the
chemocline of Lake Cadagno, Switzerland, over an annual cycle. *Aquatic Microbial Ecology* **30**: 295-302.

Peng, Y., Leung, H.Chr., Yiu, S.-M., and Chin, F.Y. (2012) IDBA-UD: a de novo assembler for single-cell and
metagenomic sequencing data with highly uneven depth. *Bioinformatics* **28**: 1420-1428.

Pfeffer, Chr., Larsen, S., Song, J., Dong, M., Besenbacher, F., Meyer, R.L. *et al.,* (2012) Filamentous
bacteria transport electrons over centimetre distances. *Nature* **491**: 218-221.

Pfennig, N. (1975). The phototrophic bacteria and their role in the sulfur cycle. *Plant and Soil*, *43*(1), 1-16.

Pfennig, N., Höfling, K.-H., and Kusmierz, H. (1968) *Chromatium okenii (Thiorhodaceae)-Biokonvektion,*
aero-und phototaktisches Verhalten: IWF.

Polerecky, L., Adam, B., Milucka, J., Musat, N., Vagner, T., and Kuypers, M.M. (2012) Look@ NanoSIMS–a
tool for the analysis of nanoSIMS data in environmental microbiology. *Environmental Microbiology* **14**:
1009-1023.

Posth, N. R., Bristow, L. A., Cox, R. P., Habicht, K. S., Danza, F., Tonolla, M., ... and Canfield, D. E. (2017).
Carbon isotope fractionation by anoxygenic phototrophic bacteria in euxinic Lake Cadagno. *Geobiology*
15: 798-816.

Prange, A., Arzberger, I., Engemann, Chr., Modrow, H., Schumann, O., Trüper, H.G. *et al.,* (1999). *In situ*
analysis of sulfur in the sulfur globules of phototrophic sulfur bacteria by X-ray absorption near edge
spectroscopy. *Biochimica et Biophysica Acta (BBA)-General Subjects* **1428**: 446-454.

Schwedt, A., Kreutzmann, A. C., Polerecky, L., and Schulz-Vogt, H. N. (2012). Sulfur respiration in a
marine chemolithoautotrophic Beggiatoa strain. *Frontiers in Microbiology*, *2*, 276.

Socrates, G. (2004). *Infrared and Raman Characteristic Group frequencies: Tables and Charts*: John Wiley
and Sons.

Sommer, T., Danza, F., Berg, J., Sengupta, A., Constantinescu, G., Tokyay, T., Bürgmann, H., Dressler, Y.,
Sepúlveda Steiner, O., Schubert, C.J. and Tonolla, M. (2017). Bacteria-induced mixing in natural waters.
Geophysical Research Letters 44: 9424-9432.

Storelli, N., Peduzzi, S., Saad, M.M., Frigaard, N.-U., Perret, X., and Tonolla, M. (2013) CO_2 assimilation in
the chemocline of Lake Cadagno is dominated by a few types of phototrophic purple sulfur bacteria.
FEMS Microbiology Ecology **84**: 421-432.

Strous, M., Kraft, B., Bisdorf, R., and Tegetmeyer, H. (2012) The binning of metagenomic contigs for
microbial physiology of mixed cultures. *Frontiers in Microbiology* **3**: 410.

Tonolla, M., Demarta, A., Peduzzi, R., and Hahn, D. (1999) In situ analysis of phototrophic sulfur bacteria
in the chemocline of meromictic Lake Cadagno (Switzerland). *Applied and Environmental Microbiology*
65: 1325-1330.

Tonolla, M., Peduzzi, S., Demarta, A., Peduzzi, R., and Dittmar, H. A. H. N. (2004). Phototropic sulfur and
sulfate-reducing bacteria in the chemocline of meromictic Lake Cadagno, Switzerland. *Journal of*
Limnology **63:** 161-170.

Tonolla, M., Peduzzi, R., and Hahn, D. (2005) Long-term population dynamics of phototrophic sulfur
bacteria in the chemocline of Lake Cadagno, Switzerland. *Applied and Environmental Microbiology* **71**:
3544-3550.

Trifinopoulos, J., Nguyen, L. T., von Haeseler, A., & Minh, B. Q. (2016). W-IQ-TREE: a fast online
phylogenetic tool for maximum likelihood analysis. Nucleic Acids Research **44**: W232-W235.

Vaituzis, Z., and Doetsch, R. (1969) Motility tracks: technique for quantitative study of bacterial
movement. *Applied Microbiology* **17**: 584-588.

Van Gemerden, H. (1968) On the ATP generation by Chromatium in darkness. *Archiv für Mikrobiologie*
64: 118-124.

Vazquez, G., Zhang, J. Z., & Millero, F. J. (1989). Effect of metals on the rate of the oxidation of H2S in
seawater. *Geophysical Research Letters* **16**: 1363-1366.

Verity, P. G., Robertson, C. Y., Tronzo, C. R., Andrews, M. G., Nelson, J. R., & Sieracki, M. E. (1992).
Relationships between cell volume and the carbon and nitrogen content of marine photosynthetic
nanoplankton. *Limnology and Oceanography* **37**: 1434-1446.

Wüest, A. (1994) Interactions in lakes: Biology as source of dominant physical forces. *Limnologica Jena*
24: 93-104.

FIGURES

Fig 1| (A) Combined oxygen (top) and sulfide (bottom) profiles of the Lake Cadagno water
column revealing the persistence of an oxygen- and sulfide- free zone over a period of 48 hours,
with contour lines indicating sulfide concentrations. The bold contour lines delimiting the region
with > 5 µM sulfide were used to define the chemocline in parallel profiles of *Chr. okenii* cell
counts (B), particulate S^0 (C), and sulfate (D). Black dots represent sampling points for all
parameters except O_2 which was measured with a microsensor mounted on a CTD probe.
Shaded boxes represent dark periods between sunset at ~20:50 and sunrise at ~6:10. Time plots
were interpolated from original profiles measured in August 2015 and are provided in Fig S1.

Figure 2| Representative day (A) and night (B) profiles through the chemocline illustrating
glycogen and S⁰ concentrations in relation to *Chr. okenii* cell numbers, oxygen, and sulfide
gradients in August 2015. No oxygen data is available for the day profile. Profiles were
measured at 4-h intervals over 2 day-night cycles and used to calculate chemical fluxes. (C) The
consumed sulfide flux (solid line) was calculated by subtracting the residual sulfide flux (dashed
line) from the total sulfide flux into the mixed layer. (D) The downwards oxygen flux into the
chemocline (solid line) was used to estimate the maximum % of sulfide aerobically respired
(dotted line), assuming the complete oxidation of sulfide to sulfate. Shaded regions represent
dark periods.

Figure 3| $^{13}CO_2$ uptake by individual *Chr. okenii* cells incubated in the dark with no O_2, 5 μM (expected) dissolved O_2, and 15 μM (expected) O_2, and in the light with no added O_2. Secondary ion images (A-C) show ratios of $^{13}C/^{12}C$. Scale bars are 3 μm and bacterial cells are outlined in white. Epifluorescence microscopy image (D) of DAPI-stained *Chr. okenii* associated with a consortium of autofluorescent algal cells enclosed within an organic sheath (dotted orange line, defined based on $^{12}C^{14}N$ image in Fig S8). White triangles denote cells identified as: *Chr* = *Chr. okenii*, *PSB* = other purple sulfur bacteria, and *np* = non-phototrophic. (E) Box-and-whisker plots showing the range and the median growth rate of *Chr. okenii* under each incubation condition calculated from ^{13}C-uptake per cell.

Figure 4| *Chr. okenii* cell illustration, showing the metabolic potential inferred from the metagenome-assembled genome with a particular focus on the genetic machinery implicated in photosynthesis, sulfur oxidation, aerobic metabolism, motility, glycogen and PHA storage, nitrogen fixation and transmembrane transport. The respiratory chain enzyme complexes are labeled with Roman numerals.

Figure 5: Schematic of phototrophic and aerobic sulfide oxidation processes in the Lake Cadagno
chemocline. Convection in the chemocline may be driven by a combination of turbulence and
sinking bacterial plumes, represented by the large number of descending *Chr. okenii* cells on the
left. As a result, oxygen and sulfide are entrained into the chemocline and immediately
consumed by purple sulfur bacteria, keeping concentrations of these compounds below
detection limits. *Chr. okenii* cells, depicted with internal sulfur globules (yellow dots), are pulled
in the direction of their flagellar bundle.