Systematic review of 50 years of coral disease research visualized through the scope of network theory

Luis M. Montilla[1,*] , Alfredo Ascanio-Moreno[1] , Alejandra Verde[1] , Aldo Croquer[1] ,

1 Laboratory of Experimental Ecology, Department of Environmental Studies, Simon Bolivar University. Caracas, Venezuela.

* luismmontilla@usb.ve

Abstract

Coral disease research encompasses five decades of undeniable progress. Since the first descriptions of anomalous signs, we have come to understand multiple processes and environmental drivers that interact with coral pathologies. In order to gain a better insight into the knowledge we already have, we explored how key topics in coral disease research have been related to each other using network analysis. We reviewed 719 papers and conference proceedings published from 1965 to 2017. From each study, four elements determined our network nodes: 1) studied disease(s); 2) host genus; 3) marine ecoregion(s) associated with the study site; and 4) research objectives. Basic properties of this network confirmed that there is a set of specific topics comprising the majority of research. The top five diseases, genera, and ecoregions studied accounted for over 48% of the research effort in all cases. The community structure analysis identified 15 clusters of topics with different degrees of overlap among them. These clusters represent the typical sets of elements that appear together for a given study. Our results show that while some coral diseases have been studied considering multiple aspects, the overall trend is for most diseases to be understood under a limited range of approaches, e.g. bacterial assemblages have been considerably studied in Yellow and Black band diseases while immune response has been better examined for the aspergillosis*Gorgonia* system. Thus, our challenge in the near future is to identify and resolve potential gaps in order to achieve a more comprehensive progress on coral disease research.

Keywords: coral disease, network analysis, research trends, community structure.

1 Introduction

Coral diseases have been an important factor responsible for the decline of coral reefs in the last decades (Rogers and J. Miller 2013). Although pathogens and diseases are part of the natural dynamics of ecosystems, including coral reefs, the interaction with other stressful environmental factors aggravates their negative effects (Ban et al. 2014), enhancing important losses of live coral cover (Lewis et al. 2017; Precht et al. 2016; Randall and Woesik 2015).This loss of coral cover has been particularly important in the Caribbean, which has been frequently called a "coral disease hot spot" because of the number of diseases and the range of affected species in comparison with the low coral cover in the region (Bruckner 2002; Green and Bruckner 2000).

The evolution of this body of knowledge has been compiled and updated frequently in multiple narrative reviews (Goreau et al. 1998; Bourne et al. 2009; Antonius 1981; Green and Bruckner 2000; Richardson 1998; Rosenberg and Ben-haim 2002; Santavy and Peters 1997; Weil et al. 2006; Woodley et al. 2015). Quantitative approaches (e.g. meta-analyses, systematic reviews) that complement these publications have been less frequent. One example is the result provided by Work and Meteyer 2014, who examined almost 500 coral disease papers, and classified the used methods in terms of six

broad categories. From their analysis they evidenced the scarcity of microscopical sign descriptions using histological techniques. In another example, Ward and Lafferty 2004 analyzed the frequency of coral disease papers as a potential proxy of a coral diseases incidence. Finally, Ban et al. 2014 reviewed the experimental research about at least two stressors simultaneously using network theory. These types of syntheses, although scarce, provide important benefits over narrative reviews such as offering a wide and more objective perspective of the research landscape, integrating the trends in subfields of a discipline, and potentially identifying research gaps and opportunities for new questions to be explored.(Lortie 2014).

Here we present a systematic review aimed at identifying groups of coral disease research topics that have been explored more frequently than others. To do this, we performed a network analysis approach. Network analysis is frequently used in systematic reviews (Borrett et al. 2014; Ohniwa et al. 2010) and allows researchers to address multiple issues in coral disease research, such as the evaluation of phage-bacteria interactions (Soffer et al. 2014) synergistic effects of environmental stressors (Ban et al. 2014), the analysis of microbial positive and negative interactions in healthy and diseased conditions (Sweet and Bulling 2017; Meyer et al. 2016), and the analysis of gene expression and regulation (Wright et al. 2015). We hypothesized that if the research topics typically addressed in coral research lack uniformity among several key aspects of epizootiology, then a network representing the co-apparition of these topics in coral disease research papers would exhibit a community structure, where the communities (also called clusters) of nodes would represent the different themes that have characterized most of coral disease research in the last 50 years.

Methods

Data acquisition

We reviewed a total of 719 publications spanning a period from 1969 to 2017. We searched for these in the search engines and databases *Google scholar*, *Meta*, and *Peerus*, using combinations of keywords with search modifiers, including "coral disease", "syndrome", "yellow", "black", "purple", "spot", "band", "pox", "dark", "plague", "growth", "trematode", "anomalies", "ciliate", "soft coral", and "aspergillosis". This included peer-reviewed papers and conference proceedings, since the latter would also provide information about research questions explored in a given time and a given location. We excluded thesis, preprints , and book chapters. Additionally, we compared our database with the list of papers analyzed by Work and Meteyer 2014 looking for potential omissions. The papers were included if the research question directly addressed some biological or ecological aspect of coral diseases or coral pathogens (Fig. 1). The comprehensive list of included papers is provided as supplementary material. For every paper, we assessed four fundamental questions, each one corresponding to one node category:

- **What was the studied disease?** Each disease was listed as an individual node and the cases where there was no specific disease of interest (e.g. general surveys), we assigned the node "multiple diseases". Additionally, we classified "white syndrome" as all the descriptions of pathologies involving tissue discoloration and loss from the Pacific *sensu* (Bourne et al. 2015). We applied similar criteria and classified "Pink syndrome" as the references to Pink spots, Pink line syndrome, Pink-Blue syndrome, and Pink-Blue spot syndrome, as these diseases have not been clearly distinguished from one another. We excluded papers specifically concerning thermally-induced bleaching but included the node "bleaching effects" for papers investigating the effects of bleaching over some disease topic or vice versa.

- **Where did the samples come from?** Or **where was the study conducted?** We reported the corresponding marine ecoregion *sensu* (Spalding et al. 2007). When no explicit sampling site was stated, we contacted the respective authors to verify the location.

- **What was the genus of the affected host?** If the study implicated multiple specific genera, each one was listed as an individual node. In the cases of baselines and similar studies

Figure 1. PRISMA flow diagram for coral disease papers included in the network analysis.

which lacked specific taxa of interest, we assigned the node "multiple genera".

- **What were the objectives of the study?** We specifically looked into the questions or specific objectives addressed in the introduction section of the papers and assigned a set of keywords. Each keyword represented one node in the network. The complete list of keywords is included as a supplementary file at `https://github.com/github.com/luismmontilla/CoDiRNet`.

Two of the authors (see author contributions) performed the assignation of nodes to each study, and in the case of disagreements, all the authors were consulted until a consensus was reached.

Network construction and community structure analysis

Using the extracted topics as vertices and their co-occurrence in a given paper as the edges, we built an undirected weighted network. The weight w_{ij} was the frequency of co-occurrence of the keywords i and j in the same article (figure 2). The network was constructed and analyzed using the R package *igraph* (Csardi and Nepusz 2006; R Core team 2016). Considering the size and complexity of the resulting graph, we used the network reduction algorithm proposed by Serrano et al. 2009, implemented in the package *disparityfilter* for R (Bessi 2015) to extract the backbone of our network. This resulted in a smaller graph that retains the multiscale properties of the original network.

Next, we used the link communities approach (Ahn et al. 2010) to obtain groups of nodes forming closely connected groups (hereafter, referred as communities), using the *linkcomm* package (Kalinka and Tomancak 2011). With this method overlapping communities may appear, which allows for several nodes to be part of multiple communities (the scripts for the used functions with their modifications are available at `https://github.com/luismmontilla/CoDiRNet`. We explored the similarity among the obtained communities, representing them as a new network of communities where each community was a node, and the edges had the value of the Jaccard coefficient for the number of shared nodes (hence, communities without shared topics would be disconnected).

In order to test the statistical robustness of the obtained communities, we measured the communities assortativity (r_{com}) using a modification of the method proposed by Shizuka and Farine (2016). In this case, this metric measures a proportion of derived replicates (permuting the network edges) that result in the same community structure as the network derived from the original data. The coefficient is a value ranging from -1 to 1, where values closer to 0 indicate that the community structure in the network of interest differed from the communities obtained in random permutations; values close to 1 indicate that the permuted networks have similar node communities as the original network; and values close to -1 indicate that the communities of the permutation-derived replicates are formed by different nodes in contrast with the original network. We used a

Figure 2. Network construction. **A** represents a hypothetical paper focusing on the prevalence of Yellow Band Disease in a country of the Southern Caribbean. Each paper produces a fully connected graph. In **B**, a second hypothetical paper generates its own graph, which has a link between *Orbicella* and 'Prevalence/Incidence' in common with **A**. **C** represents the resulting network, with the link between *Orbicella* and 'Prevalence/Incidence' representing the co-apparition in two papers and the remaining links representing one paper.

Table 1. Top five topics addressed in the coral disease research literature

Category	Topic	Frequency (%)
	Black band disease	17.1
	White syndromes	14.2
	White plague	6.4
Disease	Yellow band disease	5.6
	Growth anomalies	5.4
	Total	**48.7**
	Acropora	16.3
	Orbicella	12.0
Genus	*Porites*	9.5
	Montipora	7.6
	Gorgonia	7.4
	Total	**52.8**
	Floridian	15.9
	Eastern Caribbean	9.3
	Greater Antilles	9.2
Ecoregion	Central and Southern GBR	8.6
	Hawaii	7.6
	Total	**50.6**
	Prevalence/incidence	23.5
	Spatial patterns	13.1
	Sign descriptions	11.8
Objective	Transmission	10.0
	Temperature	9.9
	Total	**68.3**

modification (hereafter r_{oc}) that allows the use of this coefficient for overlapping communities (code available at `https://github.com/alspeed09/OverlapCommAssortativity`).

Results

The content of all the 719 coral disease papers yielded a network comprising 302 vertices and 4184 edges. The strength distribution (i.e. the distribution of the sum of all connections to the nodes) revealed that the coral disease research network, like other natural networks, presented a highly skewed distribution, where most research topics share few connections and a comparatively small group of topics are heavily represented in scientific publications. This pattern persisted for the specific distribution of each type of nodes, suggesting that over the past 50 years, coral disease research has focused on a reduced number of questions in each category (figure 3).

Considering this, we explored the identity of the five most represented topics in each case. For all the categories, five topics accounted for over 40% of the total frequency of apparition in all the examined papers (table 1). We show that the evaluations of prevalence and/or incidence have had a prominent place in research, together with questions about the spatial patterns, transmission, and descriptions of the signs (mostly visual but also histological descriptions) of diseases. The role of temperature stress had also received priority in comparison with other environmental factors.

After applying the disparity filter, the initial network was reduced to 93 vertices and 233 edges. The set of excluded research questions were included as supplementary material at `https://github.com/luismmontilla/CoDiRNet`. These remaining topics either encompass a set of highly-specific questions being scarcely explored (e.g. diseases on mesophotic corals, the effects of ocean acidification, or the role of commensal bacteria) or new questions altogether.

From this reduced network, we obtained a total of 15 node communities (hereafter, referred

Figure 3. Strength distributions for A) the global network, B) Disease nodes, C) Ecoregion nodes, D) Genus nodes, and E) Objective nodes.

as C1 to C15). We obtained a r_{oc} value of 0.23, indicating that the network effectively possess a community structure that departs from randomness. These groups ranged from small communities with low overlap, to rich and highly interconnected communities, e.g. C5 and C14 shared almost 50% of their members (figure 4).

The smaller communities were composed of six or less nodes. C3 represented four topics related to questions about temporal patterns; C6 represented four genera typically affected by Dark Spot Disease; C8 was comprised of studies dealing with stress temperature and zooxanthellae damage in the Great Barrier Reef and nearby locations. C9 showed that an important group of studies related to White Band Disease have been focused in determining the advance rate, transmission patterns, and usually a description of the signs. C10 included the effects of environmental factors on diseased and bleached corals (figure 5).

The medium-sized communities offered more clear pictures of several trends prevailing in the coral disease research. C1 was an *Acropora*-centered community. This included topics related to the descriptions of signs, bacterial assemblages, transmission mechanisms and pattern of several affections like White Syndromes, White Pox, and Brown Band Disease, in the Great Barrier Reef. C7 included studies around Caribbean Yellow band Disease, mostly about changes in the microbial community, and also temporal and spatial patterns of the disease. C11 was a community constituted by studies about *Vibrio* as a coral pathogen, especially infecting the genera *Oculina*, *Montipora*, and *Pocillopora*. C13 was a community that incorporated two research trends. The most important node in this community was White Syndrome, associated to topics like pathogen characterization, transmission experiments, and changes in bacterial assemblages of the hosts. These hosts were mainly *Montipora* and *Porites*, which were the connection to the second group: research about growth anomalies and trematode infections in Hawaii. C15 was a community centered on gorgonian affections and immune processes (figure 6.)

All the large communities (nodes > 5) had two nodes in common: Black Band Disease and Southern Caribbean. C2 was the most specific one, including objectives related to Black Band Disease only. C4 also included Yellow Band Disease and White Plague, in this case encompassing studies dealing with transmission experiments, effects of temperature and changes in the associated microbiota. C5 was a community covering reviews, meta-analysis, baseline studies, and outbreaks reports. C12 was very similar to C4, with the distinction of including studies about White Pox, pathogen characterization, particularly *Serratia marscescens*, pathogen metabolites, and the Bahamian ecoregion. C14 consisted of several topics shared with C5, with the particular difference of including studies on Acropora and White Syndromes (figure 7).

Discussion

The network approximation used in this study for analyzing the relationships between coral disease research topics provided a quantitative perspective on the intensity of the research and the extension of the field so far. The communities obtained summarized the dominant trends (considering the bulk of the historical research), with each community corresponding to the themes of interest that have accumulated most of the published papers up to 2017. It is important to notice that we expected some minimal community structure to arise in our network, for some diseases affect only a specific set of coral genera and in certain regions. However, our hypotheses was that the patterns of addressed questions would also contribute to the apparition of multiple communities in our network.

From this data we see some clear patterns. For example, Black Band Disease occupies a preponderant place in coral disease research. There are several reasons contributing to this. Black Band Disease affection has been studied since the emergence of the field (Antonius 1976; Garrett and Ducklow 1975) and it has consistently accumulated a body of knowledge that makes it the most studied coral disease so far. Its effects comprise a substantial number of coral hosts across different ecoregions and produce extensive mortality during disease epizootics (Diraviya Raj et al. 2016; Aeby et al. 2015; Yang et al. 2014; Sato et al. 2009; Hobbs et al. 2015), especially when combined with seasonal or anomalous temperature increases and determined light conditions (Chen et al. 2017;

Figure 4. Similarity network of 15 communities. Node size represents the number of topics in each community, and edge width represents the similarity between two given communities.

Figure 5. Membership matrix of communities C3, C6, C8, C9, and C10. The circles represent the value of the overall node centrality.

Figure 6. Membership matrix of communities C1, C7, C11, C13, and C15. The circles represent the value of the overall node centrality.

Figure 7. Membership matrix of communities C2, C4, C5, C12, and C14. The circles represent the value of the overall node centrality.

Lewis et al. 2017; Bhedi et al. 2017; A. W. Miller and Richardson 2015; Kuehl et al. 2011; Sato et al. 2011; Boyett et al. 2007). Additionally, there have been important advances in the field derived from studies about this disease. Studies about the pathobiome of Black Band Disease pioneered the concept of pathogenic consortia as etiological agents of coral diseases (Carlton and Richardson 1995), and the accumulated findings are allowing the proposal of etiological models (Sato et al. 2016). In contrast, other diseases have been important enough to appear in their own communities, but they have been associated with less diverse research themes. For example, Caribbean Yellow Band Disease has been studied mostly around the changes in its microbial assemblages, and soft coral Aspergillosis studies usually deal with immune processes.

Our findings also highlight the historical importance of environmental stressors for a broad number of diseases. In our network, temperature was consistently associated with several communities, and the role of sedimentation, turbidity, and nutrient loads (including input through sewage waters) were the main environmental factors associated with baseline studies. This was consistent with a previous review where the authors found that the temperature was the most frequently studied environmental variable in their sample, along with sedimentation, both variables having high influence over coral diseases (Ban et al. 2014).

This is where this contribution complements previous narrative reviews (*sensu* Lortie 2014; Petticrew 2001; Morais et al. 2018) about coral diseases; the previously mentioned low-to-mid topic diversity communities could be used as a reference of unexplored questions, locations, or affected genera in different coral diseases that could be tackled in future research. A theme that we can use as an example is the research about coral immunity. Although there have been a steady progress in this sub-field, there is still so much room for exploration, especially considering the complex symbiotic interactions that constitute the coral holobiont (Palmer 2018). This can also be considered in the context of diverse and increasing stressful environmental conditions that coral are expected to withstand. The role of high temperatures has been the main environmental objective addressed, however there are other conditions like oxygen depletion, ocean acidification, and pollution, whose interactive effects with temperature and diseases remain far less explored. Additionally, there are potential insights that can be gained through the examination of the topics excluded from the network backbone, such as emerging topics that could be relevant in the near future but are masked in our results because of the weight of the historical trends.

In summary, we obtained a generalized representation of the most explored topics in coral disease research. However, these predominant themes and questions are yet to be generalized to the range of potential coral hosts and their diseases. One additional contribution of this work is the availability of a growing database that can be used for further and more diverse exploration of the trends in this field, including the detection of emerging topics and temporal trends, which will help to gain better insights of the development of coral disease research.

Author contributions

- **Conceptualization:** LMM.

- **Literature search:** LMM, AV.

- **Data curation:** LMM, AV.

- **Formal analysis:** LMM, AA.

- **Investigation:** LMM, AA, AV.

- **Methodology:** LMM, AA, AV, AC.

- **Visualization:** LMM, AV.

- **Software:** LMM, AA, AV.

- **Writing:** LMM, AA, AV, AC.

Acknowledgments

We wish to thank Laurie Richardson, Longin Kaczmarsky, and Mike Sweet for the information provided. Also Stuart Sandin, Mark Vermeij, Eugenia Sanchez, Bill Gowacki, and Kourtney Barber for their comments on an early manuscript.

References

Aeby, Greta Smith et al. (2015). "First Record of Black Band Disease in the Hawaiian Archipelago: Response, Outbreak Status, Virulence, and a Method of Treatment". *Plos One* 10.3, e0120853. DOI: 10.1371/journal.pone.0120853.

Ahn, Yong-Yeol, James P. Bagrow, and Sune Lehmann (2010). "Link communities reveal multiscale complexity in networks". *Nature* 466.7307, pp. 761–764. DOI: 10.1038/nature09182.

Antonius, Arnfried (1976). "New observations on coral destruction in reefs". *10th Meeting of the Association of Island Marine Laboratories of the Caribbean*. Mayagüez, Puerto Rico.

— (1981). "Coral reef pathology: a review". *Proceedings of the 4th International Coral Reef Symposium*. Manila, pp. 3–6.

Ban, Stephen S., Nicholas A J Graham, and Sean R. Connolly (2014). "Evidence for multiple stressor interactions and effects on coral reefs". *Global Change Biology* 20.3, pp. 681–697. DOI: 10.1111/gcb.12453.

Bessi, Alessandro (2015). *disparityfilter: Disparity Filter Algorithm of Weighted Network*.

Bhedi, Chinmayee D et al. (2017). "Elevated temperature enhances short to medium chain acyl homoserine lactone production by Black Band Disease associated vibrios". *FEMS Microbiology Ecology* January, pp. 1–12. DOI: 10.1093/femsec/fix005.

Borrett, Stuart R., James Moody, and Achim Edelmann (2014). "The rise of Network Ecology: Maps of the topic diversity and scientific collaboration". *Ecological Modelling* 293, pp. 111–127. DOI: 10.1016/j.ecolmodel.2014.02.019. arXiv: 1311.1785.

Bourne, David G. et al. (2009). "Microbial disease and the coral holobiont". *Trends in Microbiology* 17.12, pp. 554–562. DOI: 10.1016/j.tim.2009.09.004.

Bourne, David G. et al. (2015). "Towards a better understanding of white syndromes and their causes on Indo-Pacific coral reefs". *Coral Reefs* 34.1, pp. 233–242. DOI: 10.1007/s00338-014-1239-x.

Boyett, Holly V, David G. Bourne, and Bette L. Willis (2007). "Elevated temperature and light enhance progression and spread of black band disease on staghorn corals of the Great Barrier Reef". *Marine Biology* 151.5, pp. 1711–1720. DOI: 10.1007/s00227-006-0603-y.

Bruckner, Andrew W (2002). *Priorities for effective management of coral diseases*. US Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service.

Carlton, Richard G and Laurie L. Richardson (1995). "Oxygen and sulfide dynamics in a horizontally migrating cyanobacterial mat: Black band disease of corals". *FEMS Microbiology Ecology* 18.2, pp. 155–162. DOI: 10.1016/0168-6496(95)00052-C.

Chen, Carla C.M. et al. (2017). "Modelling environmental drivers of black band disease outbreaks in populations of foliose corals in the genus Montipora". *PeerJ* 5.1, e3438. DOI: 10.7717/peerj.3438.

Csardi, Gabor and Tamas Nepusz (2006). "The igraph software package for complex network research". *InterJournal Complex Systems* 1695, p. 1695.

Diraviya Raj, K. et al. (2016). "Mass mortality of Montipora digitata (Scleractinia) in Vaan Island, Gulf of Mannar, southeast India". *Current Science* 110.8, pp. 1407–1408.

Garrett, Peter and Hugh Ducklow (1975). "Coral diseases in Bermuda". *Nature* 253.5490, pp. 349–350. DOI: 10.1038/253349a0.

Goreau, Thomas J et al. (1998). "Rapid spread of diseases in Caribbean coral reefs". *Revista de biología tropical* 46.5, pp. 157–171.

Green, Edmund P and Andrew W Bruckner (2000). "The significance of coral disease epizootiology for coral reef conservation". *Biological Conservation* 96.3, pp. 347–361. DOI: 10.1016/S0006-3207(00)00073-2.

Hobbs, Jean-Paul A et al. (2015). "Selective Impact of Disease on Coral Communities: Outbreak of White Syndrome Causes Significant Total Mortality of Acropora Plate Corals". *Plos One* 10.7, e0132528. DOI: 10.1371/journal.pone.0132528.

Kalinka, Alex T and P. Tomancak (2011). "linkcomm: an R package for the generation, visualization, and analysis of link communities in networks of arbitrary size and type". *Bioinformatics* 27.14, pp. 2011–2012. DOI: 10.1093/bioinformatics/btr311.

Kuehl, Kristin et al. (2011). "The roles of temperature and light in black band disease (BBD) progression on corals of the genus Diploria in Bermuda". *Journal of Invertebrate Pathology* 106.3, pp. 366–370. DOI: 10.1016/j.jip.2010.12.012.

Lewis, Cynthia L. et al. (2017). "Temporal dynamics of black band disease affecting pillar coral (Dendrogyra cylindrus) following two consecutive hyperthermal events on the Florida Reef Tract". *Coral Reefs*. DOI: 10.1007/s00338-017-1545-1.

Lortie, Christopher J. (2014). "Formalized synthesis opportunities for ecology: Systematic reviews and meta-analyses". *Oikos* 123.8, pp. 897–902. DOI: 10.1111/j.1600-0706.2013.00970.x. arXiv: peerj.preprints.270v1 [10.7287].

Meyer, Julie L et al. (2016). "Microbiome shifts and the inhibition of quorum sensing by Black Band Disease cyanobacteria". *The ISME Journal* 10.5, pp. 1204–1216. DOI: 10.1038/ismej.2015.184.

Miller, Aaron W and Laurie L. Richardson (2015). "Emerging coral diseases: a temperature-driven process?" *Marine Ecology* 36.3, pp. 278–291. DOI: 10.1111/maec.12142.

Morais, Juliano, Aline P.M. Medeiros, and Bráulio A. Santos (2018). "Research gaps of coral ecology in a changing world". *Marine Environmental Research* 140.June, pp. 243–250. DOI: 10.1016/j.marenvres.2018.06.021.

Ohniwa, Ryosuke L., Aiko Hibino, and Kunio Takeyasu (2010). "Trends in research foci in life science fields over the last 30 years monitored by emerging topics". *Scientometrics* 85.1, pp. 111–127. DOI: 10.1007/s11192-010-0252-2.

Palmer, Caroline V (2018). "Immunity and the coral crisis". *Communications Biology* 1.1, p. 91.

Petticrew, M. (2001). "Systematic reviews from astronomy to zoology: myths and misconceptions". *BMJ* 322.7278, pp. 98–101. DOI: 10.1136/bmj.322.7278.98.

Precht, William F. et al. (2016). "Unprecedented Disease-Related Coral Mortality in Southeastern Florida". *Scientific Reports* 6, p. 31374. DOI: 10.1038/srep31374.

R Core team (2016). *R: A language environment for statistical computing*. Vienna, Austria.

Randall, Carly J and Robert van Woesik (2015). "Contemporary white-band disease in Caribbean corals driven by climate change". *Nature Climate Change* 5.4, pp. 375–379. DOI: 10.1038/nclimate2530.

Richardson, Laurie L. (1998). "Coral diseases: what is really known?" *Trends in Ecology & Evolution* 13.11, pp. 438–443. DOI: 10.1016/S0169-5347(98)01460-8.

Rogers, Caroline S and Jeff Miller (2013). "Coral Diseases Cause Reef Decline". *Science* 340.June, pp. 1522–4. DOI: 10.1126/science.340.6140.1522-a.

Rosenberg, Eugene and Yael Ben-haim (2002). "Microbial disease of corals and global warming". *Environmental Microbiology* 4.6, pp. 318–326.

Santavy, Deborah L and Esther C. Peters (1997). "Microbial pests: coral disease in the western Atlantic". *Proceedings of the 8th international coral reef symposium*. Panama, pp. 607–6012.

Sato, Yui, David G. Bourne, and Bette L. Willis (2009). "Dynamics of seasonal outbreaks of black band disease in an assemblage of Montipora species at Pelorus Island (Great Barrier Reef, Australia)". *Proceedings of the Royal Society B: Biological Sciences* 276.1668, pp. 2795–2803. DOI: 10.1098/rspb.2009.0481.

— (2011). "Effects of temperature and light on the progression of black band disease on the reef coral, Montipora hispida". *Coral Reefs* 30.3, pp. 753–761. DOI: 10.1007/s00338-011-0751-5.

Sato, Yui et al. (2016). "Integrated approach to understanding the onset and pathogenesis of black band disease in corals". *Environmental Microbiology* 18.3, pp. 752–765. DOI: 10.1111/1462-2920.13122.

Serrano, M Angeles, Marian Boguna, and Alessandro Vespignani (2009). "Extracting the multiscale backbone of complex weighted networks." *Proceedings of the National Academy of Sciences of the United States of America* 106.16, pp. 6483–6488. DOI: 10.1073/pnas.0808904106.

Shizuka, Daizaburo and Damien R. Farine (2016). "Measuring the robustness of network community structure using assortativity". *Animal Behaviour* 112, pp. 237–246. DOI: 10.1016/j.anbehav.2015.12.007.

Soffer, Nitzan, Jesse R Zaneveld, and Rebecca Vega-Thurber (2014). "Phage-bacteria network analysis and its implication for the understanding of coral disease". *Environmental Microbiology*. DOI: 10.1111/1462-2920.12553.

Spalding, Mark D. et al. (2007). "Marine Ecoregions of the World: A Bioregionalization of Coastal and Shelf Areas". *BioScience* 57.7, p. 573. DOI: 10.1641/B570707.

Sweet, M.J. and Mark T. Bulling (2017). "On the Importance of the Microbiome and Pathobiome in Coral Health and Disease". *Frontiers in Marine Science* 4.January, p. 9. DOI: 10.3389/fmars.2017.00009.

Ward, Jessica R. and Kevin D. Lafferty (2004). "The elusive baseline of marine disease: Are diseases in ocean ecosystems increasing?" *PLoS Biology* 2.4, pp. 542–547. DOI: 10.1371/journal.pbio.0020120.

Weil, Ernesto, Garriet W. Smith, and Diego Luis Gil-Agudelo (2006). "Status and progress in coral reef disease research." *Diseases of aquatic organisms* 69.1, pp. 1–7. DOI: 10.3354/dao069001.

Woodley, Cheryl M et al., eds. (2015). *Diseases of Coral*. Hoboken, NJ: John Wiley & Sons, Inc. DOI: 10.1002/9781118828502.

Work, Thierry M. and Carol Meteyer (2014). "To Understand Coral Disease, Look at Coral Cells". *EcoHealth*. DOI: 10.1007/s10393-014-0931-1.

Wright, Rachel M et al. (2015). "Gene expression associated with white syndromes in a reef building coral, Acropora hyacinthus". *BMC Genomics* 16.1, p. 371. DOI: 10.1186/s12864-015-1540-2.

Yang, HongQiang et al. (2014). "Black band disease as a possible factor of the coral decline at the northern reef-flat of Yongxing Island, South China Sea". *Science China Earth Sciences* 57.4, pp. 569–578. DOI: 10.1007/s11430-013-4734-y.