High-definition transcranial direct current stimulation over the dorsal premotor cortex modulates motor network excitability

Stephanie Lefebvre[1], Kay Jann[2], Allie Schmiesing[1], Kaori Ito[1], Mayank Jog[2], Yuchuan Qiao[3], Ryan Cabeen[3], Yonggang Shi[3], Nicolas Schweighofer[4], Danny JJ Wang[2], *Sook-Lei Liew[1]

1   *Neural Plasticity and Neurorehabilitation Laboratory, University of Southern California, Los Angeles, CA United States,*

2   *Laboratory of FMRI Technology, USC Stevens Neuroimaging and Informatics Institute, Keck School of Medicine of USC, University of Southern California, Los Angeles, CA,*

3   *Laboratory of Neuro Imaging (LONI), USC Stevens Neuroimaging and Informatics Institute, Keck School of Medicine of USC, University of Southern California, Los Angeles, CA, USA.*

4   *Computational Neuro-Rehabilitation Laboratory, University of Southern California, Los Angeles, CA,*

*Corresponding author:

Sook-Lei Liew

sliew@chan.usc.edu

Abstract: 185/200 words

Keywords: brain stimulation, tDCS, MRI, primary motor cortex, cortical excitability

Running title: HD-tDCS to modulate motor network excitability

Abstract

The effectiveness of transcranial direct current stimulation (tDCS) over the primary motor cortex

(M1) to modulate motor network excitability is highly variable. The dorsal premotor cortex (PMd)

may be a promising alternative target, as it provides high-level motor control using multiple

pathways, including one independent of M1 and one with direct, influencing connections to M1.

This double-blind, placebo-controlled study aimed to differentially excite M1 and PMd using high-

definition tDCS (HD-tDCS) with concurrent functional magnetic resonance imaging. Both M1 and

PMd HD-tDCS demonstrated high inter-individual variability. However, PMd HD-tDCS led to a

slightly greater increase in responders and greater changes in local complexity than M1 HD-tDCS

or sham. PMd responsiveness was also positively correlated with the strength of the corticospinal

tract originating from PMd. An MRI analysis of individual motor hotspot locations revealed that

actual motor hotspots were often located between M1 and PMd, rather than directly over

anatomical M1. This heterogeneity in the stimulation site, despite using a canonical definition of

the motor hotspot, may contribute to the variability of tDCS results. Altogether, these findings

provide new considerations for improving tDCS effects on motor excitability.

Transcranial direct current stimulation (tDCS) is a noninvasive tool that can modulate cortical

excitability in human brain regions (Nitsche and Paulus 2000). The effects of tDCS on upper limb

motor rehabilitation have been widely studied, and more than 20 studies have shown promising

effects of tDCS applied over the primary motor cortex (M1) on post-stroke motor recovery

(Lefebvre and Liew 2017; Shaheiwola et al. 2018). However, a major issue that prevents tDCS

from being widely adopted in clinical practice is the high inter-individual variability shown in

behavioral or cortical physiological changes following M1 stimulation (Ammann et al. 2017;

Lopez-Alonso et al. 2018). The reasons for this inter-individual variability have not been fully

elucidated (for a review, see Lefebvre & Liew, 2017).

A potential solution to reduce the inter-individual variability is to use alternative stimulation

sites and/or more precise tDCS montages. That is, it is possible that stimulating another entry point

into the motor network may lead to more reliable changes following tDCS. To date, the majority

of tDCS studies for motor rehabilitation have focused on targeting M1 to alter motor network

excitability based on two primary reasons. First, the site for M1 stimulation can be easily

personalized to each individual using transcranial magnetic stimulation (TMS) to identify the

individual's motor hotspot, which is assumed to represent M1. Second, since M1 is the primary

region associated with descending motor pathways to the spinal cord (i.e., 50% of the corticospinal

tract (CST) neurons originate in M1 (Dum and Strick 1991)), it makes sense to target this region

to influence motor behavior. However, given the variable responses in stimulating M1, other

stimulation sites that have functional connections with M1 should also be considered.

The dorsal premotor cortex (PMd) is a promising alternative neural target for modulating motor

excitability. The PMd is the second major origin of the CST with 30% of CST projections (Barbas

and Pandya 1987; Dum and Strick 1991). As a higher motor area (Kantak et al. 2012), the PMd

contributes to motor control of the upper limb using a downstream pathway that is independent of

M1 (Dum and Strick 1991; Etkin et al. 2011; Fujiyama et al. 2013), leading to some high-level

control of motor commands (Thura and Cisek 2014). Moreover, neural inputs to the hand and

forelimb representation in M1 mostly originate from PMd (Dum and Strick 2005; Burman et al.

2014), creating a cortico-cortical circuitry that has influences on motor network excitability

(Stinear et al. 2009; Wymbs and Grafton 2013; Neva et al. 2014; Bestmann and Krakauer 2015).

Studies using monkeys have shown that if there are cortico-cortical connections between M1 and

premotor cortex, the onset of the movement-related activity is always earlier in the premotor cortex.

This suggests that these connections are initiated from the premotor cortex (Umilta et al. 2007;

Thura and Cisek 2014). As such, PMd may modulate motor output in two ways: first, through its

dedicated CST projections to the spinal cord; second, through its direct controlling connection to

M1. Thus, due to its strong involvement in motor learning (Hardwick et al. 2013), motor selection,

movement planning, and preparation (Etkin et al. 2011), and its controlling role over M1

excitability, PMd provides a promising alternative target for modulating motor network

excitability. Stimulating PMd could also make the motor network more susceptible to cortical

excitability changes. Only a few studies have examined the use of tDCS to modulate PMd, but most

of them suggest that targeting PMd with tDCS could enhance motor performance and motor

learning (for a review, see Buch et al., 2017). However, to our knowledge, there are no studies

directly comparing the effects of M1 and PMd-tDCS on motor network excitability.

This lack of direct comparison between M1 and PMd as targets for tDCS is likely due to a

major drawback of conventional tDCS montages, which is their poor spatial resolution. The

distribution of the electrical current using a standard tDCS montage is relatively diffuse, leading to

changes far beyond the stimulated regions (Nitsche et al. 2007). Therefore, trying to target M1 with

conventional tDCS likely involves stimulation of at least a part of PMd (Kuo et al. 2013). The

main reason for diffuse current densities in conventional tDCS montages is the large inter-electrode

distance, which is limited by the size of the conventional electrodes (e.g., 5x5 or 5x7 cm). In

contrast, recently introduced high-definition tDCS (HD-tDCS) devices provide the ability to

increase stimulation focality (Caparelli-Daquer et al. 2012) by delivering current through a 4x1

ring montage using smaller electrodes, making it easier to differentially target M1 and PMd.

Although there is still likely a little overlap in the stimulation, the ring configuration helps to focus

the peak stimulation over the targeted area.

We assessed changes in motor network excitability following M1 versus PMD tDCS using two

primary methods: (1) changes in cortical physiology using TMS and (2) changes in motor network

connectivity and complexity using resting-state fMRI (rs-fMRI). More specifically, TMS-induced

motor evoked potentials (MEP) (Nitsche and Paulus 2000) were used to evaluate cortical

excitability locally at the motor hotspot (M1), while rs-fMRI (Lefebvre et al. 2017) was used to

measure brain activity pre- and post-stimulation more broadly across the motor network. We used

rs-fMRI to estimate functional connectivity (FC) of the motor network as well as to calculate

multiscale entropy (MSE), which reflects the complexity of regional BOLD signals (Liang et al.

2014; Wang et al. 2018). FC and MSE are complementary measures of network integrity, and MSE

has been shown to reflect the dynamics of neural networks across multiple time scales or temporal

frequencies (Heisz et al. 2012). Specifically, MSE changes during rs-fMRI have been reported

following tDCS (Liang et al. 2014; Kang et al. 2018). Since the complexity of neural activity

fluctuations is associated with the probability of neural firing (Wang et al. 2018), this makes MSE

a particularly relevant measure for assessing excitability changes following anodal tDCS, which is

thought to increase the probability of neural firing (Giordano et al. 2017).

The present study harnesses recent methodological innovations to examine concurrent HD-

tDCS and fMRI. This technique provides the unique opportunity to not only explore whether

stimulation of different areas in the motor network (M1 vs PMd) may have different effects on

motor network cortical excitability but also to see if there were associated local or network changes

underlying these effects, allowing us to more carefully examine mechanisms of plasticity. An

additional benefit of this concurrent method is the ability to localize the stimulation electrodes on

each individual's brain to assess whether the stimulated region matched the targeted region. Finally,

using both functional and structural brain imaging, we were able to examine whether there were

any neuroimaging predictors of response to HD-tDCS in any of the groups (M1, PMd, sham).

In this study, we hypothesized that stimulating the left PMd with HD-tDCS would modulate

motor network excitability more reliably than stimulating left M1 in healthy individuals, due to the

additional motor pathways emerging from PMd. We also examined whether there were concurrent

changes in motor network connectivity and complexity related to tDCS responsiveness in each

group using both standard functional connectivity and newer MSE approaches. We further explored

whether there were any functional or structural neuroimaging predictors of HD-tDCS

responsiveness in each group. Finally, we examined each individual's stimulated location in

relation to the targeted stimulation locations to see whether inter-individual variability in the

stimulated location might relate to HD-tDCS responsiveness.

MATERIAL AND METHODS

Population

The protocol was approved by the University of Southern California Institutional Review

Board. All subjects gave written informed consent in accordance with the 1964 Declaration of

Helsinki. The inclusion criteria were the following: (1) being a healthy volunteer aged 18 – 50

years, (2) being right-handed. The exclusion criteria were the following: (1) having a pacemaker

or other piece of metal in the body, (2) pregnancy, (3) previous history of stroke or other brain

damage, (4) previous history or family history of epilepsy, (5) previous history of head injury, (6)

previous history of skull damage, (7) previous history of migraines, (8) currently taking any

medications that may affect cortical excitability (e.g. benzodiazepines), (9) inability to understand

the instructions. Forty-six healthy participants took part in this double-blind, placebo-controlled

study (neither the participants nor the primary experimenter knew the nature of the stimulation

(e.g., real or sham HD-tDCS)). At inclusion, a first experimenter randomly assigned each

participant into one of the following 3 groups: HD-tDCS applied over left M1, HD-tDCS applied

over left PMd or sham HD-tDCS (in which case, electrodes were randomized over M1 or PMd).

The second experimenter, who performed the data acquisition and analysis, was only aware of the

location of the electrodes but remained blinded to whether participants received real or sham tDCS.

Study design

The study involved a single three-hour visit which included the following 3 sessions (see Figure

1): (1) a pre-MRI TMS session, including the localization of M1 and PMd as well as the recording

of neurophysiological measurements, (2) a concurrent HD-tDCS/MRI session, including the

placement of the HD-tDCS cap, acquisition of structural MRI sequences, a pre-HD-tDCS rs-fMRI

scan, the double-blinded HD-tDCS stimulation with concurrent acquisition of rs-fMRI scan, as well

as a post-HD-tDCS rs-fMRI scan, and (3) a post-MRI TMS session to measure changes in

neurophysiological measurements following HD-tDCS. To our knowledge, this is the first known

randomized, group study using a concurrent HD-tDCS and fMRI paradigm in humans. The safety

of these concurrent designs was established in a pilot study from Gbadeyan and collaborators in

2016 using both phantoms and a single participant (Gbadeyan et al. 2016). They also examined the

tolerability of their specific double ring HD-tDCS montage in a behavioral session (without fMRI)

in 30 participants.

Figure 1: Study design. Each participant will take place in 3 sessions as follows: a first TMS

session, then an MRI/HD-tDCS session, and finally a second TMS session. *HD-tDCS: high-*

definition transcranial direct current stimulation, MEP: Motor Evoked Potential, RMT: Resting

Motor Threshold, rs-fMRI: resting-state functional magnetic resonance imaging

<u>Pre-MRI TMS session</u>

Briefly, we used single- and paired-pulse transcranial magnetic stimulation (TMS) (Magstim

200² device with BiStim module; Magstim Inc., UK) with a figure-of-eight coil, along with the

Brainsight neuronavigation system (Rogue Resolutions Ltd, UK) and surface electrodes to record

muscle activity (electromyography, or EMG; Delsys Trigno wireless sensors, Delsys Inc, MA). To

localize the motor hotspot, we recorded EMG at the right first dorsal interosseous (FDI). We then

recorded pre-HD-tDCS cortical excitability, including each individual's resting motor threshold

(RMT) and single-pulse motor-evoked potentials (MEPs) with a test stimulus at the intensity

needed to generate an MEP of 0.75mV peak-to-peak amplitude. Notably, this test stimulus is in

accordance with previous studies showing a range of test stimuli between 0.2mV and 4mV (Wagle-

Shukla et al. 2009; Schambra et al. 2015b). We also measured short-interval intracortical inhibition

(SICI) using a paired-pulse TMS paradigm (see Brain Stimulation Intervention section below for

details on the TMS paradigms). MEPs were recording using Signal 6.05 software (Cambridge

Electronic Design Ltd., UK).

MRI session

Immediately following the first TMS session, participants were fitted with the HD-tDCS cap

and began an MRI session including the acquisition of an anatomical image (T1w MPRAGE), a

T2-weighted anatomical MRI, a diffusion-weighted MRI, a 7-minute baseline rs-fMRI (pre-HD-

tDCS rs-fMRI), a second 7-minute rs-fMRI during which participants received sham or HD-tDCS

applied over M1 or PMd, depending on their group, and a third 7-minute rs-fMRI to assess the after

effects of HD-tDCS (see MRI section below for more details). An additional experimenter, who

was unblinded to the stimulation condition (real or sham), turned the tDCS stimulation device on

and off. In this way, the primary experimenter and participant remained blinded to the stimulation

condition.

Post-MRI TMS session

Immediately following the MRI session, the participants took part in a second TMS session to

quantify the HD-tDCS-induced changes on cortical excitability. The cortical excitability

measurements of this second TMS session started less than 10 minutes after the end of the third

resting state fMRI scan (i.e., time for the participant to leave the MRI, walk to the adjacent TMS

experimental room and have the HD-tDCS equipment removed and the TMS coil repositioned on

their M1 hotspot).

Brain stimulation intervention

<u>TMS paradigm</u>

We defined the motor hotspot as the spot at which the maximal MEP peak-to-peak amplitude

was recorded in the FDI muscle using a single TMS pulse at $50\% \pm 5\%$ of the maximum stimulator

output. We defined the RMT as the intensity of the stimulator that induced an MEP of at least 50

µV peak-to-peak amplitude in 5 out of 10 trials. We defined the test stimulus for the current study

as the intensity of the stimulator that induced an MEP of at least 0.75mV ($TS_{0.75mV}$) peak-to-peak

amplitude in 5 out of 10 trials.

After defining these thresholds, we used them in an MEP-SICI paradigm which consisted of 40

trials in a randomized order with a $7s \pm 10\%$ inter-trial duration. This paradigm included 20 single-

pulse trials using the $TS_{0.75mV}$ (the peak-to-peak amplitude of these trials will be referred to as MEP

amplitude), and 20 paired-pulse trials using a conditioning stimulus (CT) at 80% of the RMT 2.5ms

before the $TS_{0.75mV}$ (the peak-to-peak amplitude of these trials will be referred to as MEP amplitude

in SICI trials). We anticipated that the SICI trials should result in a decrease of MEP amplitude in

SICI trials compared to MEP trials. Therefore, to analyze HD-tDCS effects on intracortical

inhibition, we compared the ratio of MEP amplitude divided by MEP amplitude in SICI trials

between pre and post-HD-tDCS. (For SICI results, see *Supplementary Data 1*).

For 6 participants (2 in each group), we were unable to reach the intensity needed to elicit an

MEP of 0.75mV, or even 0.5mV. In these cases, we used the RMT as the test stimulus to acquire

MEP amplitude. For one additional participant in the M1 group, we were only able to localize the

motor hotspot but did not acquire any other neurophysiological measurements due to the

participant's reported discomfort. This participant was included only in the resting-state analyses.

<u>HD-tDCS experiment</u>

We used the Soterix MRI-compatible HD-tDCS system (4x1 ring configuration) to deliver

tDCS (1mA) for 7 minutes during an fMRI session (see Figure 1). We used 5 small gel-filled

electrodes (base diameter: 2.4 cm, gel-skin contact area of ~4.5 cm^2) positioned in adjacent

electrode locations within the cap, and we placed the center electrode on the cap over the location

of the neural target (as defined below). Each of the 4 cathodal ring electrodes delivered a -0.25mA

current leading to a 1mA stimulation over the center anodal electrode.

<u>Localization of the HD-tDCS targets</u>

For the M1 group, we positioned the center electrode over the motor hotspot, and for the PMd

group, we positioned the center electrode 2.5cm anterior to the motor hotspot, which is the standard

localization for PMd (Rizzo et al. 2004). The use of this MRI-compatible device also gave us the

unique opportunity to obtain an anatomical MRI with the exact position of the electrodes on the

participant's own head. We therefore also report the localization of individual stimulation sites

across participants in the Results section.

MRI session acquisition and preprocessing

MRI acquisition

MRI data were acquired on a 3T Prisma MRI scanner (Siemens, Germany) with a 32-channel

head coil using protocols from the Human Connectome Project (HCP)

(*http://protocols.humanconnectome.org/CCF/;https://www.humanconnectome.org/storage/app/m*

edia/documentation/lifespan-pilot/LSCMRR_3T_printout_2014.08.15.pdf). The MRI sequences

acquired included the following: T1-weighted MPRAGE scan (208 sagittal slices, 0.8mm thick,

TR = 2400ms, TE = 2.22ms, flip angle = 8°, voxel size 0.8 x 0.8 x 0.8mm); T2-weighted turbo

spin-echo scan (208 sagittal slices, 0.8mm thick, TR = 3200ms, TE = 523ms, flip angle = 8°, voxel

size 0.8 x 0.8 x 0.8mm); diffusion MRI scan (92 slices, 1.5 mm thick, TR = 3230ms, TE = 89.20ms,

multi-band factor = 4, flip angle = 78°, voxel size 1.5 x 1.5 x1.5mm, with a gradient protocol with

7 scans at b=0 s/mm2, 47 at b=1500 s/mm2, 46 at b=3000 s/mm2, and a complete repetition with

reversed phase encoding in the A-P direction); and three rs-fMRI sequences (7 minutes and 6

seconds covering 520 volumes of 72 slices per scan; TR = 800ms, TE = 37ms, flip angle = 52°,

voxel size 2 x 2 x 2mm).

Resting-state fMRI processing

We analyzed the rs-fMRI data using SPM12, the CONN toolbox (FC toolbox (Whitfield-

Gabrieli and Nieto-Castanon 2012)*; http://www.nitrc.org/projects/conn*) and the Complexity

Toolbox (http://loft-lab.org/index-5.html). The preprocessing steps included slice-timing

correction, motion realignment, noise correction using white matter, CSF and motion parameters

as regressors, and band-pass filtering (0.01-0.1 Hz). We also performed co-registration between

the functional scans and 3D-T1w MPRAGE scans of each participant. Finally, we normalized the

functional scans to MNI space and smoothed them using a Gaussian filter of 6mm.

To measure functional changes in the motor network, we compared the FC between regions of

interest (ROI-to-ROI analyses) and analyzed the neural complexity by computing the MSE of the

BOLD time series in each ROI (defined below) before and after HD-tDCS.

ROI-to-ROI analyses

We defined 4 ROIs as the left M1, right M1, left PMd and right PMd. We used a meta-analysis

from Hardwick and collaborators (Hardwick et al. 2013) to define the localization of the ROIs in

the left hemisphere and then flipped them to the right hemisphere. The exact coordinates of the

center of each ROI, which was defined as a sphere with a radius of 10mm, were: left M1 (x = -38,

y = -24, z = 56), right M1 (x = 38, y = -24, z = 56), left PMd (x = -26, y = 2, z = 60), right PMd (x

= 26, y = 2, z = 60).

In addition, as an exploratory analysis, we also defined an additional ROI to represent the whole

motor network, which was defined from a spatial map from an independent component analysis

(ICA) on the rs-fMRI data before HD-tDCS was administered (Figure 2). Although we primarily

use the main 4 Hardwick ROIs (left and right M1 and PMd) for most analyses, when analyzing the

difference between each ROI and the whole network, we use this ROI as the definition of the whole

network. Additional methods on the ICA analysis are presented in *Supplementary Methods 1*.

We extracted the cross-correlations of these 4 ROIs (left/right M1, left/right PMd) and ordered

them for each group and each condition based on the correlation intensity. To compare the

organization of the motor network following each stimulation session, we compared these orders

using Spearman's rank correlations coefficient.

We also used RM-ANOVAs (one RM-ANOVA for each ROI as a seed region) in the CONN

toolbox to compare combinations of pairs between the 4 ROIs (the 6 FC pairs were: left M1-right

M1, left PMd-right PMd, left M1-left PMd, left M1-right PMd, left PMd-right M1, right M1-right

PMd) using "Time" (pre-HD-tDCS, post-HD-tDCS) and "Group" (M1, PMd and sham) as factors.

Figure 2: Localization of the regions of interest. Panel A: Localization of the 4 ROIs based

on the meta-analysis from Hardwick and collaborators[Hardwick et al. 2013]. Each ROI was defined as a

sphere with a radius of 10mm around the reported peak coordinates for each region. These were:

left M1 (x = -38, y = -24, z = 56), right M1 (x = 38, y = -24, z = 56), left PMd (x = -26, y = 2, z =

60), right PMd (x = 26, y = 2, z = 60). Panel B: The whole motor network component as generated

from the pre-HD-tDCS spatial map from the ICA analysis. *L: left, M1: primary motor area, PMd:*

dorsal premotor cortex, R: right, ROI: region of interest

Multiscale-entropy

MSE was used to evaluate changes in neural complexity following HD-tDCS in each of the 4

ROIs (left M1, right M1, left PMd, right PMd). The MSE metric quantifies the regularity or

complexity of biological signals across a range of coarse-grained timescales or temporal

frequencies. Complex systems with 1/f power spectra exhibit constant entropy over various

timescales (due to their fractal properties), whereas random noise shows a marked decrease in

entropy at longer time scales (as random fluctuations are smoothed out) (Wang et al. 2018). We

used a pattern matching threshold (r) of 0.5 and a pattern length (m) of 2 (McDonough and Nashiro

2014b; Wang et al. 2018; Yang et al. 2018). In total, we obtained 20 coarse-sampled scales. Based

on the band-pass filter applied, we kept only 9 scales for the analyses (scales 12 to 20).

Structural Correlates of MRI Responsiveness

In addition to measuring HD-tDCS-induced changes in rs-fMRI (connectivity and complexity),

we also performed additional analyses to check for any potential anatomical factors that could

explain or predict HD-tDCS-induced effects. Therefore, we also measured the strength of the

corticospinal tract (using diffusion MR tractography) and the cortical thickness of the ROIs to see

if any of these measures influence HD-tDCS effects.

Diffusion MRI

For the diffusion MRI preprocessing, we followed the HCP processing pipeline

(*https://github.com/Washington-University/HCPpipelines/tree/master/DiffusionPreprocessing*).

Then, we modeled each voxel with a multi-compartment fiber orientation distribution (FOD)

approach (Shi and Toga 2017). Finally, we obtained the diffusion tensor fractional anisotropy (FA)

values using FSL and completed the deterministic FOD fiber tracking (Cabeen et al. 2016) using

the Quantitative Imaging Toolkit (QIT) (Cabeen RP et al. 2018). For each participant, we created

three tracts: a global corticospinal tract (CST) in the left hemisphere, a CST section that originated

from the left M1 and a CST section that originated from the left PMd (see Figure 3 for an example

in one participant). For each participant, we first drew the global left CST from an ROI covering

the left posterior limb of the internal capsule. Following this, we created two different sub-tracts

by defining the overlap between the individual's global CST with M1 and PMd template tracts

based on 100 subjects, as defined in Archer et al., 2018 (Archer et al. 2018). This resulted in a total

CST tract, an M1 tract, and a PMd tract for each individual. Finally, we extracted the FA value of

each tract for each individual to perform a correlation between FA and the physiological outcomes.

Figure 3: Example of the DTI tracts in one participant. Left panel: After positioning the PLIC

ROI, we extracted the CST of the participant. Middle panel: By overlapping the CST obtained and

the M1 or PMd template tract, we obtained the M1 and PMd tract for each participant. Right panel:

Superposition of the M1 and PMd tracts for one participant. Note that the M1 tract is localized at

the posterior part of the CST whereas the PMd tract is more anterior. The yellow division line is

shown to visualize the separation between the two tracts. *PLIC: posterior limb of the internal*

capsule, CST: corticospinal tract, M1: primary motor area, PMd: dorsal premotor cortex

Cortical thickness analysis

We used FreeSurfer software version 6 (*http://surfer.nmr.mgh.harvard.edu*) to automatically

create a 3D model of the cortical surface for cortical thickness measurements on the T1-3D scans

of all our participants (Fischl and Dale 2000). We manually inspected the individual surfaces for

potential segmentation errors between gray and white matter, and we manually corrected

segmentations as needed. We computed cortical thickness measurements on each of the 4 ROIs

(left M1, right M1, left PMd and right PMd) for each participant.

Statistical analyses

We used R (*https://www.r-project.org/about.html*) to perform all the statistical analyses, except

for the analyses performed using the CONN toolbox (rs-fMRI FC). We used the CONN toolbox

for ROI-to-ROI analysis, and the results are presented at a qFalse discovery rate = 0.05 (corrected

for multiple comparisons).

We performed a repeated measures RM-ANOVAs to compare changes in MEP amplitude

across groups, and separate RM-ANOVAS to compare changes in MSE for each of the 4 motor

ROIs. We used factors of "Time" (pre-HD-tDCS, post-HD-tDCS) and "Group" (M1, PMd and

sham) in all RM-ANOVAs. To evaluate HD-tDCS-induced changes within each group, we also

performed a priori analyses of pre/post-HD-tDCS differences in MEP amplitude for each of the 3

groups using two-way paired t-tests. We report the test statistics and p-values, along with the effect

size, using partial eta squared (η_P^2) for the ANOVA main effects, and Cohen's d for the t-tests. Any

post-hoc tests are corrected for multiple comparisons using the Tukey HSD. In addition, given the

large inter-individual variability in the data, we also reported the Bayes Factor (BF) for the group

level changes over time in MEP amplitude and MSE analyses using the R package "BayesFactor".

The BF is used to report a ratio of the strength of the experimental effect compared to the null

hypothesis and has recently been used with highly variable experimental data, including in brain

stimulation studies (Apsvalka et al. 2018; Liew et al. 2018). In our data, we are using the anovaBF

to determine which model explained the best the variation observed in the results. We tested 3

models: impact of "Time", impact of "Group" and impact of the interaction Group*Time. For each

anovaBF, the model with the highest BF value will be the best combination of models among these.

The strength of the impact of this model on the data is determined by the size of the BF value.

Briefly, a BF of 0-3 should be considered as weak evidence, a BF of 3-10 as substantial evidence,

a BF of 10-30 as strong evidence and a BF >30 as very strong evidence (Schonbrodt et al. 2017).

To determine which parameters might influence the HD-tDCS effects, we estimated Pearson's

correlation coefficients relating pre-HD-tDCS measurements (rs-fMRI FC, MSE, FA, cortical

thickness) with the MEP changes in each of the M1 and PMd groups. We reported correlation p-

values that are both uncorrected (p_{uncorr}) and corrected for multiple comparisons (p_{corr}) using the

Bonferroni correction for the 4 comparisons (rs-fMRI FC, MSE, FA, cortical thickness). Finally,

we performed two-way independent samples t-tests to compare each of these 4 measurements

between responders and non-responders. For the t-tests, we also reported both p_{uncorr} and p_{corr}. We

defined the "responders" subgroup as participants in the M1 or PMd group who showed an increase

in MEP amplitude following HD-tDCS [(averaged MEP amplitude over the 20 trials post-tDCS) /

(averaged MEP amplitude over the 20 trials pre-tDCS) > 1)]. The "non-responders" subgroup

refers to participants in the M1 or PMd group who demonstrated a decrease or no change in MEP

amplitude following HD-tDCS [(averaged MEP amplitude over the 20 trials post-tDCS) / (averaged

MEP amplitude over the 20 trials pre-tDCS) ≤ 1)]. Although there is a fine boundary between the

two, we note that it has been consistently used to define responders/non-responders in the literature

(Guerra et al. 2017; Lopez-Alonso et al. 2018). Due to the large variability observed in the TMS

outcomes, however, we also explored different methods to classify the responders/non-responders.

This additional analysis is summarized in *Supplementary Data 2*.

RESULTS

Neurophysiological measurements

<u>Baseline thresholds</u>

We first examined whether there were any differences within the sham group for individuals

with electrodes placed either over M1 or PMd. Using Student t-tests, we did not find any differences

between these two sham groups across any measures, including in RMT (t(14) = 0.68, p = 0.50),

$TS_{0.75mV}$ (t(12) = 0.57, p = 0.58), MEP amplitude pre-tDCS (t(14) = 0.05, p = 0.96), or MEP

amplitude post-tDCS (t(14) = 0.29, p = 0.78). Given this, we merged the data in a single sham

group for all of the following analyses (Table 1).

We then checked to see whether the initial cortical excitability between the 3 groups was similar

by comparing the pre-HD-tDCS RMT and $TS_{0.75mV}$ across groups (Table 1). Using two separate

one-way ANOVAs with "Group" (M1, PMd, sham) as a factor, we found that there was no

significant difference between the 3 groups in the stimulator intensity used to reach either the RMT

(n = 45, Group effect: F(2) = 0.50, $p = 0.61$) or the $TS_{0.75mV}$ threshold (n = 39, Group effect: F(2)

= 0.49, p = 0.62).

Subject	Group	RMT	TS0.75mV	MEP pre-tDCS (mV)	MEP post-tDCS (mV)
s01	M1	48	55	1.14	2.37
s02	M1	--	--	--	--
s03	M1	47	65	1.72	0.87
s04	M1	47	55	1.33	1.68
s05	M1	50	57	2.22	1.78
s06	M1	50	69	1.17	0.96
s07	M1	41	47	0.86	0.46
s08	M1	43	70	0.94	2.25
s09	M1	46	54	0.79	0.24
s10	M1	59	--	0.07	0.07
s11*	*M1*	*38*	*48*	*1.06*	*6.8*
s12	M1	47	--	0.18	0.15
s13	M1	58	75	1.64	1.92
s14	M1	46	66	0.91	1.63
s15	M1	37	64	0.95	1.39
Mean		*46.93*	*60.42*	*1.07 (*1.07)*	*1.21(*1.61)*
SD		*6.32*	*8.99*	*0.58 (*0.56)*	*0.80(*1.68)*
s16	PMd	38	57	1.18	3.26
s17	PMd	45	80	0.48	0.71
s18	PMd	50	63	0.38	0.49
s19	PMd	42	51	1.48	0.92

s20	PMd	44	53	0.98	2.56
s21	PMd	48	72	0.38	0.99
s22	PMd	38	44	1.17	2.32
s23	PMd	63	73	0.3	0.41
s24	PMd	54	--	0.11	0.43
s25	PMd	35	52	1.02	1.52
s26	PMd	51	--	0.21	0.08
s27	PMd	41	80	0.39	2.13
s28	PMd	38	39	0.52	0.25
s29	PMd	37	41	0.7	0.61
s30	PMd	39	56	0.49	1.38
Mean		*44.20*	*58.54*	*0.68*	*1.20*
SD		*7.74*	*14.06*	*0.41*	*0.96*
s31	sham-M1	60	78	1.39	0.68
s32	sham-M1	50	--	0.26	0.65
s33	sham-M1	40	50	2.14	0.62
s34	sham-M1	33	44	0.99	0.82
s35	sham-M1	48	60	0.61	0.23
s36	sham-M1	35	40	2.41	1.33
s37	sham-M1	49	53	0.92	3.37
s38	sham-M1	43	52	0.58	1.2
s39	sham-PMd	48	72	0.85	0.85
s40	sham-PMd	48	54	0.59	0.49

s41	sham-PMd	56	72	1.33	0.83
s42	sham-PMd	65	--	0.14	0.03
s43	sham-PMd	48	55	1.24	0.87
s44	sham-PMd	44	65	3.32	0.16
s45	sham-PMd	38	50	0.56	3.63
s46	sham-PMd	36	36	1.1	1.87
Mean sham		*46.31*	*55.79*	*1.15*	*1.05*
SD sham		*8.94*	*12.39*	*0.84*	*0.83*

Table 1: Initial cortical excitability measurements. SD: standard deviation; RMT: resting motor threshold, *outlier, -- indicates the measurement could not be elicited. For the M1 group, MEP mean and SD are presented excluding the outlier and with the outlier (*).

Changes in MEP amplitude

Inter-individual variability in HD-tDCS response

We then assessed inter-individual variability in response to HD-tDCS between the 3 groups, focusing on the number of individuals showing changes in MEP following HD-tDCS in each group. As noted in the Methods, a 'responder' was characterized as anyone with an increase in MEP amplitude compared to baseline, as per previous research (Guerra et al. 2017; Lopez-Alonso et al. 2018).

We found considerable inter-individual variability in MEP changes following HD-tDCS in the 3 groups (Figure 4). In the M1 group, 7 out of 14 (50%) participants showed an increase in MEP amplitude, while in the PMd group, 11 out of 15 participants (73.3%) showed an increase in MEP amplitude. Importantly, in the sham group, 5 out of 16 participants (31.25%) showed an increase

in MEP amplitude following sham HD-tDCS, suggesting that some individuals will show some

level of change in cortical excitability without stimulation. The number of participants with

increased MEP amplitude post-HD-tDCS showed a trending difference across the 3 groups (χ^2 (2)

= 5.497, *p = 0*.06). Additional, more stringent definitions of 'responders' were also explored (see

Supplementary Data 2), with generally similar results.

Due to the large inter-individual variability, especially in the M1 group, we ran an outlier

detection analysis, using the Quantile Range Outliers method to identify any outliers beyond 3

interquartile ranges (calculated on MEP amplitude across the entire cohort). In doing so, we

identified and removed one participant with an outlier response in the M1 group from further

excitability analyses (an increase in MEP amplitude following HD-tDCS of more than 3

interquartile ranges, or 4.2mV; see Figure 4).

Figure 4: Individual changes in MEP amplitude over time. The dot pairs reflect MEP amplitude

pre and post-tDCS for each participant in each of the 3 groups. Participants who showed a decrease

in MEP between pre and post-tDCS are displayed in purple. *In the M1 group, one participant

presented a very strong effect post-tDCS. After running an outlier detection, this participant was

removed from the rest of the analyses (indicated by the black arrow). *M1: primary motor area,*

MEP: motor evoked potential, mV: millivolts, PMd: dorsal premotor cortex, tDCS: transcranial

direct current stimulation.

Group level changes in MEP amplitude

We next evaluated the magnitude of HD-tDCS-induced changes in cortical excitability across

the 3 groups by comparing the amplitude of MEP amplitude before and after HD-tDCS (Table 1,

Figure 5).

A two-way RM-ANOVA was performed on MEP amplitude, with "Time" (pre-HD-tDCS,

post-HD-tDCS) and "Group" (M1, PMd and sham) as factors. There were no significant interaction

or group effects (RM-ANOVA: n = 44, Interaction Group*Time $F_{(2,41)}$ = 1.88, p = 0.17, η_p^2 =

0.31, BF = 0.5 ± 1.29 %; Group: $F_{(2,41)}$ = 0.52, p = 0.59, η_p^2 = 0.04, BF = 0.22 ± 0.54%; Time:

$F_{(1,41)}$ = 1.83, p = 0.18, η_p^2 = 0.0030, BF = 0.48 ± 1.13 %). The BF analysis confirmed that the

model exploring the effect of "Interaction Group*Time" on MEP amplitude is the one explaining

the best the results; however, these results are still very weak. We then examined a priori

comparisons of pre versus post HD-tDCS for each group. This demonstrated a significant increase

in MEP amplitude only following PMd HD-tDCS (*PMd post > pre-HD-tDCS t(41) = 2.56, p =*

0.03, Cohen's d = 0.73; M1 post > pre-HD-tDCS t(41) = 0.55, p = 0.58, Cohen's d = 0.20; sham

post > pre-HD-tDCS t(41) = 0.44, p = 0. 66, Cohen's d = 0.12).

Figure 5: Changes in MEP amplitude for each group across time. Box-and-whisker plots

of MEP amplitude per group over time (pre/post-tDCS). The center line represents the median

value, the lower bound of the box represents the 25th percentile, the upper bound of the box the

75th percentile, and the whiskers represent 3 times the interquartile range. Any outliers are

indicated by dots. Note that the only statistically significant difference is observed in the PMd

group reflecting an increase of MEP amplitude post-tDCS compared to pre-tDCS. *M1: primary*

motor area, MEP: motor evoked potential, mV: millivolts; PMd: dorsal premotor cortex, tDCS:

transcranial direct current stimulation.

Finally, we examined whether there was a difference in the magnitude of MEP changes within

the individuals who responded to M1 HD-tDCS versus PMd HD-tDCS (i.e., the responders). That

is, even though more individuals showed an increased MEP following HD-tDCS in the PMd group

than the M1 group, we wanted to evaluate whether the amplitude of the MEP changes was different

between these two groups. Using just responders from M1 (n=6) and PMd (n=11) groups, a two-

way RM-ANOVA with "Time" (pre-tDCS, post-tDCS) and "Group" (M1, PMd) as factors showed

an increase in cortical excitability following HD-tDCS, but no group effect or interaction (RM-

ANOVA: n = 17, Interaction Group*Time $F(1,27) = 0.13$, $p = 0.72$, $\eta_p^2=0.002$, BF = 270.8 ± 7.38

% ; Group: $F(1,15) = 2.74$, $p = 0.12$, $\eta_p^2 =0.12$, BF = 0.87 ± 1.76 %; Time: $F(1,15) = 26.72$, $p =$

0.0001, $\eta_p^2=0.30$, BF = 499.7 ± 1.2 %). The BF analysis confirmed that the model exploring the

effect of "Time" on MEP amplitude is the one that best explains the results. Then, we used post-

hoc tests to compare MEP amplitude pre versus post HD-tDCS, again just for the responders in

each group: *PMd post > pre-HD-tDCS* t(15) = 4.37, *p = 0.0005, Cohen's d=1.12; M1 post > M1*

pre-HD-tDCS t(15) = 2.72, *p = 0.014, Cohen's d=2.14.* This demonstrates that when individuals

responded to HD-tDCS, there were no significant differences in the magnitude of MEP amplitude

change between M1 and PMd response. Also, the increase in MEP amplitude for responders was

significantly greater than at baseline.

Impact of baseline excitability on tDCS-induced MEP improvements

As observed in Figure 5, tDCS effects on MEP amplitude reported in the PMd group could

be due to the fact that the lowest pre-tDCS MEP amplitudes were observed in some participants in

this group. This occurred despite the fact that the stimulator intensity was set to induce an MEP of

at least 0.75mV for all participants. However, due to intra-individual variability in MEPs, the

average MEP amplitude was sometimes higher or lower than the targeted amplitude. Thus, we

performed an additional analysis to explore the potential impact of individuals' initial excitability

levels on MEP amplitude changes after tDCS. The correlation between pre-MEP amplitude and

change in MEP after either M1 or PMd tDCS suggests a non-significant trend (n = 28, r = -0.35, *p*

*= 0.*06). This could be interpreted to suggest that the less excitable the motor cortex is at baseline,

the greater the change in MEP amplitude may be after tDCS. Preliminarily, this weak effect seems

to be limited to the actual stimulation groups (M1, PMd). In looking at the six individuals from the

sham group with a low baseline MEP amplitude (<0.75mV; Table 1), half show an increase in the

post-MEP measurement, while the other half show a decrease. This preliminarily suggests that a

low initial MEP may not necessarily be linked to a higher increase in the second MEP, in the

absence of an intervention.

Resting-state fMRI

<u>Resting-state functional connectivity</u>

We then examined whether there were overall changes in the functional organization of the

motor network by comparing cross-correlation rankings for each of the ROI-to ROI pairs before

versus after HD-tDCS. However, the global organization of the motor network did not change after

HD-tDCS in any of the groups, as demonstrated by a strong rank correlation within each group

between pre and post-HD-tDCS measurements (M1 group: n = 15, r = 0.88, *p = 0.*03, PMd group:

n = 15, r = 0.88, *p = 0.*03, sham group: n = 16, r = 0.94, *p = 0.*02). In addition, we compared

changes in rank correlations between groups using a Fisher's Z test(Diedenhofen and Musch 2015),

which also showed no significant differences in correlation ranks between groups (M1 > PMd: z =

0.0, *p =*1; M1 > sham: z = 0.78, *p = 0.*43, PMd > sham: z = 0.78, *p = 0.*43). This suggests that

there was no global reorganization of the relative ROI-to-ROI connectivity strengths in the motor

network following either M1 or PMd HD-tDCS.

We then examined changes in connectivity for each ROI-to-ROI pair using an ANOVA with

Time and Group as factors (n = 46, thresholded at qFDR = 0.05; CONN toolbox). Here we also

found no significant changes post-HD-tDCS in any of the 3 groups. Using left PMd as the seed

ROI, there were no significant changes in FC post-HD-tDCS in the M1 group $(F(3,41) = 1.47, p$

$= 0.67)$, the PMd group $(F(3,41) = 0.18, p = 0.91)$ or the sham group $(F(3,41) = 0.34, p = 0.94)$.

Using right PMd as the seed ROI, there were no significant changes in FC post-HD-tDCS in the

M1 group $(F(3,41) = 1.16, p = 0.67)$, the PMd group $(F(3,41) = 0.18, p = 0.91)$ or the sham group

$(F(3,41) = 0.21, p = 0.94)$. Using left M1 as the seed ROI, there were no significant changes in FC

post-HD-tDCS in the M1 group $(F(3,41) = 0.55, p = 0.73)$, the PMd group $(F(3,41) = 0.23, p =$

$0.91)$ or the sham group $(F(3,41) = 0.33, p = 0.94)$. Using right M1 as the seed ROI, there were no

significant changes in FC post-HD-tDCS in the M1 group $(F(3,41) = 0.43, p = 0.73)$, the PMd

group $(F(3,41) = 0.39, p = 0.91)$ or the sham group $(F(3,41) = 0.13, p = 0.94)$

Additional exploratory analyses focusing on ROI-to-ROI comparisons and ICA network

comparisons at a less stringent threshold (cluster-level corrected p<0.01) are reported in

Supplementary Data 3 & 4, Supplementary Figures 1 & 2, Supplementary Table 1, and

Supplementary Discussion.

<u>Multi-scale entropy of resting-state fMRI</u>

We then examined HD-tDCS-associated changes in MSE to examine potential changes of

motor network excitability (Figure 6).

Figure 6: Changes in MSE across groups. Box-and-whisker plots of the MSE per group over

time (pre/post-tDCS) in the left PMd ROI (in the left panel) and in the right PMd ROI (in the right

panel). The center line represents the median value, the lower bound of the box represents the 25th

percentile, the upper bound of the box the 75th percentile, and the whiskers represent 3 times the

interquartile range. Any outliers are indicated by dots. *L: left, M1: primary motor area, MSE:*

multiscale entropy, PMd: dorsal premotor cortex, R: right, tDCS: transcranial direct current

stimulation.

RM-ANOVAs with "Time" (pre-HD-tDCS, post-HD-tDCS) and "Group" (M1, PMd, sham) as

factors were performed for each of the 4 regions of interest (left/right M1, left/right PMd) and

demonstrated a significant interaction between group and time and significant main effect of group,

with a significant increase in MSE only in the left PMd following PMd HD-tDCS (Left PMd RM-

ANOVA: n = 46, Interaction Group*Time $F(2,43) = 9.06$, $p = 0.0005$, $\eta_p^2 = 0.14$, BF = 498 ± 6.4

%; Group: $F(2,43) = 7.5$ $p = 0.002$, η_p^2 0.19, BF = 11.19 ± 0.61 %; Time: $F(1,43) = 2.33$, $p = 0.13$,

$\eta_p^2 = 0.02$, BF = 0.59 ± 1.91 %. The BF analysis confirmed that the model exploring the effect of

the "Interaction Group*Time" on the left PMd MSE changes is the one that best explains the results.

Then, we used post-hoc tests to compare left PMd MSE pre versus post HD-tDCS for each group:

PMd post > pre-HD-tDCS t(43) =4.09, *p = 0.0002*, Cohen's d=1.31*, M1 post > pre-HD-tDCS t(43)*

=-1.94, p = 0.06, Cohen's d=-0.61, sham post > pre-HD-tDCS t(43) = 0.50, p = 0.61, Cohen's d=

0.17). There was also a trend towards a decrease in MSE in the left PMd following M1 stimulation,

but this was not significant.

There was also a significant interaction and main effect of group, with a decrease in MSE only

in the PMd group in the right PMd after HD-tDCS (Right PMd RM-ANOVA: n = 46, Interaction

Group*Time F(2,43) =3.89, *p = 0.03*, η_p^2= 0.09, BF= 0.41 ± 2.85 %; Group: F(2,43) = 0.02, *p =*

0.9, $\eta_p^2 = 0.0003$, BF= 0.14 ± 1.34 %; Time: F(1,43) =1.14, *p = 0.29*, η_p^2= 0.009, BF= 0.45 ± 1.3

%. The BF analysis suggested that the model exploring the effect of "Time" on the right PMd MSE

changes best explained the results, although the results are weak. Then, we used post-hoc tests to

compare right PMd MSE pre versus post HD-tDCS for each group: *PMd post > pre-HD-tDCS* t(43)

=-2.43, *p = 0.02*, Cohen's d =-0.78*, M1 post > pre-HD-tDCS t(43) =-0.71, p = 0.48, Cohen's d=-*

0.28, sham post > pre-HD-tDCS t(43) =1.48, p = 0.14, Cohen's d= 0.60. There were no significant

findings in either left or right M1, and there were no significant correlations between changes in

MSE with changes in MEP amplitude in either the M1 or PMd groups.

Correlation between MEP changes and baseline functional/structural brain measures

We next calculated correlations between the changes in MEP amplitude after HD-tDCS in M1

and PMd groups with a number of baseline functional connectivity and neuroanatomical measures

to determine if any initial functional or structural measurements were predictive of changes in

cortical excitability following HD-tDCS. Specifically, we assessed correlations between MEP

changes and the following factors: (1) baseline rs-fMRI FC between M1/PMd and the whole motor

network ROI, (2) the initial MSE complexity levels in M1/PMd, (3) the strength of the CST

originating from M1 or PMd (as measured by fractional anisotropy), and (4) the cortical thickness

in the M1/PMd. As a reminder, the whole motor network ROI was defined using an ICA spatial

map representing the motor network.

In the M1 group, MEP change did not correlate with either initial rs-fMRI FC between left M1

and the whole motor network (n = 13, r = 0.48, P_{uncorr} = 0.09, P_{corr} = 0.36), initial MSE in left M1

(n = 13, r = 0.18, P_{uncorr} = 0.56, P_{corr} > 0.9), FA in the left M1 tract (n = 13, r = 0.29, P_{uncorr} = 0.38,

P_{corr} > 0.9), or cortical thickness in left M1 (n = 13, r =-0.25, P_{uncorr} = 0.39, P_{corr} > 0.9).

In the PMd group, MEP change showed a positive correlation with the FA in the left PMd tract

(n = 15, r = 0.62, t(13) = 2.89, P_{uncorr} = 0.012, $P_{corr=0.048}$), but it did not correlate with the initial

resting-state functional connectivity between the left PMd and the whole motor network (n = 15, r

= -0.02, P_{uncorr} = 0.9, P_{corr} > 0.9), initial MSE in left PMd (n = 15, r = 0.19, P_{uncorr} = 0.49, P_{corr}

> 0.9), or cortical thickness in left PMd (n = 15, r =-0.13, P_{uncorr} = 0.65, P_{corr} > 0.9).

We also evaluated whether any of the above factors could explain the distinction between

responders and non-responders, and thus compared them between responders and non-responders

in the M1 and PMd groups (for full analyses, see *Supplementary Data 5*). Here, we also only found

that FA in the left PMd tract was higher in responders (0.55 ± 0.01) compared to non-responders

(0.52 ± 0.003; t(12.73) = 6.09, n = 15, P_{uncorr} < 0.001, P_{corr} < 0.001).

Comparison of the stimulation site localization

Finally, as we collected both TMS and MRI with tDCS stimulation sites localized on them, we performed a post-hoc analysis to examine if the individual locations of M1 and PMd, defined using standard TMS and tDCS brain stimulation protocols (see *Methods*), matched the functional/anatomical definitions for these regions on their MRIs. We used the same ROIs from the Hardwick meta-analysis (Hardwick et al. 2013) to define the standard functional M1 and PMd locations (see Figure 2). The standard functional M1 location matched the hand knob (Yousry et al. 1997), which is commonly thought to represent the anatomical location of the motor hand area. For PMd, due to the large extent of the anatomical PMd, the standard functional PMd was within its anatomical extent. However, some studies have found activation associated with motor performance and motor learning in a more anterior part of the anatomical PMd boundaries (Mayka et al. 2006) compared to Hardwick location (Hardwick et al. 2013). We quantified the distance of each individual's stimulated region from the standard M1/PMd ROIs by computing the minimum Hausdorff's distance (i.e., the minimum distance between the two ROI boundaries) (Fedorov et al. 2008).

In the M1 group, the average Hausdorff's distance between the individual stimulation site and the standard M1 was 3.79±4.64mm, and with the standard PMd, it was 3.46±4.89mm. In the PMd group, the average Hausdorff's distance between the individual stimulation site and the standard PMd was 4.93±4.80mm, and with the standard M1, it was 15.44±4.78mm.

In terms of the number of individuals with overlap with each ROI, in the M1 group, the M1 stimulation site overlapped with the standard M1 ROI in 6 participants and with the standard PMd ROI in 6 participants. In the remaining 3 participants, the stimulation site was localized between the M1 and PMd ROIs. In the PMd group, the PMd stimulation site overlapped with the standard

PMd ROI in 5 participants and between the M1 and PMd ROIs in 2 participants. In the other 8

participants, the stimulation site localization was localized a little anteriorly to the standard PMd

ROI, but still within the anatomical (atlas-based) extent of PMd. Given the wide variability in actual

stimulation site in the M1 group, in *Supplementary Data 6,* we performed an exploratory analysis

to compare the response of the participants who were initially assigned to the M1 group based on

the location of their actual stimulation (*Supplementary Figure 3)*. We re-analyzed the pre-post MEP

amplitude changes, adding the six individuals in the M1 group who actually received PMd

stimulation to the existing PMd group for a total of 20 individuals. We again found a significant

increase in MEP amplitude post-tDCS (t(14)=2.67, p=0.018), confirming previous results.

Finally, we computed Pearson correlations between the distance from each ROI and MEP

changes to see if distance from a specific ROI related to MEP response. This showed only non-

significant trends for each group. In the M1 group, participants with the greatest MEP changes also

tended to have the largest distance between the stimulation site and the PMd ROI (t(13) = 2.3, *n =*

15, r = 0.53, p = 0.06). In contrast, in the PMd group, participants with the greatest MEP changes

also tended to have the shortest distance between the stimulation site and the PMd ROI (t(13) =

1.8, *n = 15,* r =-0.48, p = 0.07). There were no significant correlations between these distances and

changes in MSE post-HD-tDCS in any group. To see a visualization of the overlap of each

individual stimulation site with the standard ROIs, see Figure 7. For more information and

correlation plots, see *Supplementary Figure 4*.

Figure 7: Localization of the stimulation site. This figure displays the localization of the

actual stimulation sites in all the participants and shows an overlap of these sites with the standard

left M1 or left PMd localization for both groups. The actual stimulation sites are presented in a heat

map, with individual subjects plotted in red and overlap between participants reflected in warmer

colors (with yellow representing the surface shared by most of the participants). The standard M1

and PMd ROIs are displayed in blue. The overlap between the actual and standard localization is

colored in purple. *M1: primary motor area, PMd: dorsal premotor cortex, L: left.*

Overall, this suggests that there is large inter-individual variability in M1 and PMd localization

using canonical methods (e.g., defining M1 as the TMS motor hotspot and PMd as 2.5 cm anterior

to the motor hotspot). In particular, many individuals in the M1 group had M1 stimulation sites that

were bordering or overlapping with PMd, or somewhere between M1 and PMd. This heterogeneity

in the stimulation site could contribute to the variability of responsiveness to HD-tDCS. We,

therefore, compared the location of the stimulation sites between responders and non-responders.

As shown in Figure 8, more than half of the responders, including participants in both the M1 and

PMd groups, received stimulation over the same area inside PMd anatomical boundaries. We found

that 10 responders had an overlapping stimulation site within PMd, regardless of group allocation

(see *Supplementary Figure 5*), while the location of the stimulation site in non-responders across

groups was more widespread and no stimulation sites were shared by more than 4 participants. This

suggests that even if participants responded to tDCS over M1 or somewhere between M1 and PMd,

the highest and most consistent number of responders was observed when stimulation was applied

over PMd. However, there is still large inter-individual variability in the stimulated location in

responders, and some individuals in the non-responder group also were stimulated within PMd,

suggesting that there is likely not a single target across all participants that would effectively

increase excitability.

Figure 8: Heat map of the stimulation site between responders and non-responders in M1 and PMd groups. This figure displays the localization of the actual stimulation sites in the participants allocated to the M1 group (n= 14) for the 7 responders (hot heat map) and the 7 non-responders (cool heat map), and to the PMd group (n=15) for the 11 responders and the 4 non-responders. The majority of the responders in both M1 and PMd groups received stimulation over the actual same spot within the PMd anatomical boundaries (see also *Supplementary Figure 5*).

DISCUSSION

In this study, we performed a systematic, double-blind investigation using concurrent HD-tDCS

and fMRI to examine changes in brain network connectivity and complexity before and after HD-

tDCS stimulation of either M1 or PMd. To our knowledge, this is the first known group study using

concurrent HD-tDCS and fMRI to compare M1 and PMd stimulation. In addition to concurrent

HD-tDCS and fMRI, we examined changes in cortical excitability using TMS before and after HD-

tDCS. We also investigated whether any baseline measures of cortical connectivity, complexity or

neuroanatomy correlated with response to HD-tDCS in either group. Finally, we explored the inter-

individual variability in the stimulation site using canonical localization techniques and reported

the distance between actual versus expected stimulation sites.

Overall, we observed that HD-tDCS applied over both left M1 or left PMd, produced highly

variable changes across individuals. Due to the large inter-individual variability in response to

tDCS, the magnitude of changes in cortical excitability was not significantly different between the

3 groups. However, despite this, we found that HD-tDCS over left PMd produced more slightly

reliable increases in cortical excitability than HD-tDCS over left M1 or sham (i.e., more

participants showed increased MEP amplitudes following HD-tDCS over PMd than over M1 or

sham), and we found that MEP amplitude increased significantly after HD-tDCS only in the PMd

group (not in M1 or sham groups). The slight increase of cortical excitability in the PMd group was

also accompanied by modulations in motor network complexity, observed only after PMd HD-

tDCS and not after M1 HD-tDCS or sham.

We also explored whether any functional or structural brain characteristics could influence

responsiveness to tDCS. We observed that the strength of the CST originating from left PMd

demonstrated a significant positive correlation with the magnitude of MEP changes following HD-

tDCS over PMd and was significantly different between responders versus non-responders in the

PMd group. In addition, motor cortex excitability slightly predicted the magnitude of the response

to M1 or PMd tDCS, such that a less excitable motor cortex at baseline correlated with a greater

change in MEP amplitude after tDCS.

Finally, we showed that there was significant variability in the location of the actual stimulation

sites compared to the expected stimulation sites based on canonical electrode placement – a factor

which may introduce great variability into studies of brain stimulation. Specifically, we showed

that many participants in the M1 group actually received stimulation over anatomical PMd or

between M1 and PMd. In addition, more than half of the participants who responded to tDCS,

across both M1 and PMd groups, received stimulation in the anatomical PMd boundaries, whereas

the participants who did not respond to tDCS received stimulation across the motor network with

no consistent location. Put another way, stimulation over PMd (regardless of the group assignment)

produced the greatest number of individuals with increased MEP amplitude following tDCS.

Altogether, these results suggest that PMd HD-tDCS may lead to more reliable changes in

cortical excitability and complexity than M1 HD-tDCS, but also that individual brain structural and

functional characteristics, such as corticospinal tract integrity, baseline motor cortex excitability

and the location of the functional motor hotspot, likely play a strong role in tDCS responsiveness.

PMd HD-tDCS may lead to small but more reliable changes in cortical excitability than M1

HD-tDCS

As previously observed, there was large variability in the MEP measurement even in the sham

group, suggesting that this measure is highly variable over time and across individuals. However,

in line with our initial hypothesis, we observed a trend of a slightly more reliable increase of MEP

amplitude following PMd-HD-tDCS, resulting in 75% responders (i.e., participants who

demonstrated an increase in MEP amplitude after HD-tDCS) compared to only 50% responders in

the M1 group (and 31% in the sham group). Moreover, the comparison between pre and post-tDCS

MEP amplitude demonstrated that only the PMd group showed a slightly significant increase post-

tDCS.

However, when examining just the participants who responded to the stimulation, both M1 and

PMd HD-tDCS led to significant increases in MEP amplitude after HD-tDCS, with no significant

differences in amplitude between the groups. This suggests that changing the entry point into the

motor network from M1 to PMd increases the reliability of HD-tDCS effects on cortical excitability

by increasing the number of participants who responded to HD-tDCS, but not the amplitude of

changes in the responders. Similar to what has been previously suggested (Lopez-Alonso et al.

2018), here we showed that M1 may not be the best target to consistently enhance cortical

excitability, even in healthy volunteers. Moreover, in line with recent studies from stroke

rehabilitation, targeting PMd may be an effective option to enhance the excitability of the motor

network (Cunningham et al. 2015). Notably, this study used a modest sample of 15-16 participants

per group and had relatively modest effect sizes; future studies with more participants are needed

to confirm this finding. However, we also observed that higher changes in MEP amplitude

following tDCS tend to be related to an initial lower excitability level of the motor cortex. Thus,

participants with a less excitable motor cortex may benefit the most from tDCS, suggesting that the

magnitude of tDCS effects may be related to an individual's basic cortical excitability at rest.

Interestingly, previously work has also shown that responders to anodal tDCS had lower baseline

MEPs than nonresponders (Lopez-Alonso et al. 2014). However, in the previous study, the baseline

MEP characteristic did not show a significant correlation with response using logistic regression.

Overall, this finding is preliminary given our limited sample and mixed previous results, and should

be confirmed with larger samples.

Network changes following HD-tDCS

Functional connectivity

Both M1 and PMd HD-tDCS led to no differences in motor network FC. Although previous

work has suggested that tDCS may induce changes in FC of the motor network (Sehm et al. 2013;

Amadi et al. 2014), our analyses showed no significant changes. Our exploratory analyses, at a

lower threshold, presented in *Supplementary Data 3 and 4, Supplementary Figures 1 and 2, and*

Supplementary Discussion, demonstrated some trends towards modifications in FC occurred after

HD-tDCS but these modifications are only significant at lower thresholds. It is possible that the

short duration of the HD-tDCS session (7 minutes) and low amplitude of current (1 mA) might not

have been sufficient to induce reliable and significantly different changes in the FC of the motor

network, which had been previously observed after 10 to 20 minutes of stimulation (Sehm et al.

2013; Amadi et al. 2014).

Multiscale Entropy

On the other hand, we did find changes in multiscale entropy in left PMd following left PMd

HD-tDCS. MSE, a method for measuring the complexity of finite length time series, quantifies the

regularity of a time series by calculating the complexity of a signal at different time scales (Costa

et al. 2005). This measure is used as a complexity measure for neuroimaging signals (McDonough

and Nashiro 2014a; Smith et al. 2014; Li et al. 2018) and reflects the dynamics of the neural

networks (Heisz et al. 2012). Previous work has shown that increasing brain complexity is

associated with brain adaptation (Liang et al. 2014; Wang et al. 2018). Importantly, MSE is thought

to measure fine-grained changes in brain function following tDCS (Liang et al. 2014; Lafontaine

et al. 2016; Kang et al. 2018). Despite a large variability in the baseline MSE measurements, both

the RM-ANOVA and the Bayesian Factor analysis suggested that MSE measurement in left PMd

presented changes following tDCS that were different across the 3 groups, while the changes in

right PMd were relatively weak. However, the post-hoc analyses confirmed that as hypothesized,

PMd HD-tDCS led to significant and more reliable changes in MSE than M1 HD-tDCS (i.e., a

strong increase of the MSE in the left PMd along with a weak decrease of the MSE in the right

PMd). On the other hand, no significant MSE changes were observed after M1 HD-tDCS. Given

that PMd HD-tDCS led to more reliable changes in cortical excitability, it is likely this was also

captured in the MSE changes found in both hemispheres (i..e., an enhancement of the excitability

in the stimulated hemisphere associated with a decrease of the excitability in the second hemisphere

(Shin and Pelled 2017; Mondini et al. 2018)). In line with previous studies (Wang et al. 2018),

MSE and FC seemed to be complementary metrics: MSE may be more sensitive and able to capture

changes in a node of the motor network whereas traditional rs-fMRI FC would capture changes

between nodes.

Fractional anisotropy of the corticospinal tract corresponds to HD-tDCS response in the PMd

group

Previous research has shown difficulty finding reliable biomarkers that predict responsiveness

to HD-tDCS. Examples of proposed but variable biomarkers include neurochemical or genetic

variants (Antal et al. 2010; Brunoni et al. 2013; Puri et al. 2015) and EEG-based measures of brain

activity (Nuzum et al. 2016). However, recent studies have suggested that functional connectivity

(Hordacre et al. 2017), as well as the integrity of the white matter fiber tracts originating from the

targeted area (Lin et al. 2017), could predict the response to HD-tDCS. In the present study, none

of the measured parameters tended to correlate with the change in MEP amplitude after M1 HD-

tDCS. On the other hand, in the PMd group, similar to what was observed previously (Lin et al.

2017), higher FA in the part of the CST that originated from left PMd was related to increased MEP

amplitude after PMd HD-tDCS. Moreover, stronger FA values in the left PMd-portion of the CST

were observed in the participants who responded to PMd HD-tDCS. This suggests that an

individual's structural integrity of the descending corticospinal tract, as measured by the fractional

anisotrpy of the corticospinal tract emerging from the stimulated region, may be a predictor of HD-

tDCS response. This should be further examined in a large sample.

Inter-individual variability is observed after tDCS

In the present study, we observed high inter-individual variability in HD-tDCS-induced

changes in cortical excitability. This variability is similar to what has been recently reported, with

up to 50% of participants showing no response to tDCS (Wiethoff et al. 2014; Lopez-Alonso et al.

2015; Ammann et al. 2017). Here, when we compared groups of M1 and PMd HD-tDCS, we found

only to a modest enhancement of the number of participants who responded to HD-tDCS. However,

when we examined individual stimulation locations, and reallocated individuals according to their

stimulation location, we found a consistent increase in MEP amplitude for the PMd group (for those

that actually received PMd HD-tDCS). We also consistently found the PMd to be the most-

stimulated location in responders. This suggests that stimulating the PMd to modulate the motor

network might help reduce inter-individual variability compared to stimulating M1, as long as it is

actually the PMd that is being stimulated.

That is, the position of the electrodes may also introduce considerable variability. In

accordance with the canonical methods for electrode placement, we localized M1 based on the

motor hotspot and PMd 2.5 centimeters anterior to the motor hotspot (Cunningham et al. 2015). As

displayed in Figure 7, despite using this established localization method, the position of the

stimulated area was highly variable across individuals when compared with their expected

anatomical location. Specifically, in the M1 group, in more than half of the participants, the actual

localization of the stimulation did not overlap with the standard M1 localization (Hardwick et al.

2013). In contrast, as the PMd was slightly wider and defined anteriorly to the motor hotspot, more

individuals fell within the PMd ROI definition or within its anatomical boundaries (slightly anterior

to the PMd ROI). Therefore, it seems that the one-size-fits-all method to localize the target area

may not be effective in everyone. It also suggests that the motor hotspot, which is thought to reflect

the brain area that controls the FDI muscle in our case, may not always be localized at M1 or the

hand knob, but may bleed into the adjacent premotor cortex, as previously observed (Ahdab et al.

2016). This is also supported by studies with primates showing that both M1 and the adjacent PMd

are involved in maps of motor control, without any clear functional boundaries between the regions

(Godschalk et al. 1995; Graziano et al. 2002a; Graziano et al. 2002b). This suggests that the motor

hotspot localization using TMS cannot be assumed to be M1 (hand knob) but may also be a location

within the premotor cortex. Moving forward, these data suggest that researchers should keep track

of where they are stimulating and report where they ultimately find the motor hotspot instead of

assuming it is always over M1. This would help to show the variability in hotspot locations across

individuals. Moreover, this also suggests that there is an individualized brain organization which,

similar to fingerprints, may be slightly different across individuals.

Also, since the PMd location was dependent on the location of the motor hotspot, this lead to

variation in PMd as well. In our study, the actual PMd stimulation location did not always overlap

with the functional PMd localization (as obtained from the Hardwick meta-analysis (Hardwick et

al. 2013)) but it still overlapped with the anatomical PMd boundaries. In future work, if it is possible

to first obtain an MRI, it may be advisable to use neuroanatomical constraints on the motor hotspot

localization using the individual's MRI and neuronavigation-guided TMS to find the spot within

the anatomical area that provides the strongest response. This might help to reduce inter-individual

variability in experimental parameters. In the practical constraint that an MRI cannot be obtained

prior to TMS, registering all participants to a template brain to guide and record motor hotspot

localization may be an alternative solution to at least note whether there is a common stimulation

site for responders.

Finally, our results suggest there is no single brain area that would ensure a positive

response to tDCS across all participants. Although more than half of the participants who did

respond to tDCS received stimulation over an overlapping PMd spot, for the other half, the

stimulation site was somewhere between M1 and PMd. Moreover, in the participants who did not

respond to tDCS, the stimulation site was all over the motor network, including in PMd, with no

strong overlap across participants. Again, this suggest that more than the absolute location,

differences in brain structure and function (Valizadeh et al. 2018) should be taken into account to

determine the responsiveness of each individual to tDCS. Future studies should determine which

stimulation parameters (neural target, nature of stimulation, duration, etc.,) will be the most

effective to induce changes in motor network excitability and function at the individual level

instead of following a one-size-fits-all method.

Limitations

The present study has several limitations. First, although the size of our groups (n=15-16 per

group) is relatively large for studies looking at cortical excitability following brain stimulation,

these groups are still relatively small, especially given the large variability within each group.

Replicating these results in a larger sample would be useful. A second limitation is that the current

sham group included half M1 and half PMd electrode positions, which showed no significant

differences with one another, and which we therefore combined into a single sham group. However,

to provide proper control for each experimental group, a matched full sham group should be used

in the future. Third, in the present study, we used a short 7-minute stimulation duration at 1 mA

current, which may not have been enough to induce a strong effect in cortical excitability or

network changes. However, this study demonstrated that stimulating a brain region other than M1

(in this case, PMd) even for a short period can lead to an enhancement of cortical excitability and

complexity, suggesting it was strong enough to elicit some changes. However, future work may

examine whole-brain network changes following longer durations or higher intensities of HD-

tDCS. Fourth, the difference we observed between PMd and M1 stimulation could have been

related to the structural and anatomical differences between M1 and PMd more than the targeted

area itself. Specifically, PMd stimulation is targeting a gyrus (Chouinard and Paus 2006), whereas

M1 stimulation is targeting a region within a sulcus (White et al. 1997; Chouinard and Paus 2006).

Future studies that more carefully target stimulation based on individual anatomy would help

address this issue. Fifth, we used a parallel study design, as opposed to a cross-over design. Each

participant received HD-tDCS over only one neural target (M1 or PMd). Consequently, we cannot

affirm that the participants who responded to PMd HD-tDCS would have not also have responded

to M1 HD-tDCS, or vice-versa. Future studies involving a larger sample size and a cross-over

design are needed to answer this specific question. Finally, in our study, the main outcome (MEP

amplitude over time as measured with TMS) suffers from both large intra- and inter-individual

variability. Previous reports (Schambra et al. 2015a) suggested that this variability was diminished

at the group-level, suggesting TMS measures may be best suited for detecting group-level changes.

Thus, due to the high variability in the TMS outcomes, the definition of "responders" as those with

any increase in MEP amplitude compared to the baseline measurement could be criticized as being

part of measurement noise. Accordingly, we tested several other thresholds for "responders" that

required more stringent thresholds (see *Supplementary Data 2*). Based on these additional methods,

we demonstrated that the responders/non-responders distinction was generally reliable across the

different methods, especially for the PMd group (which had consistently more responders in it than

the other groups). This confirmed the accuracy of the method we choose and also confirmed that

changes observed in the PMd group were more reliable than in the M1 group.

Conclusion

To our knowledge, this is the first large study employing HD-tDCS with fMRI in a double-

blind design with multiple stimulation sites. Using this design, we investigated the

neurophysiological effects of tDCS on the motor network immediately after stimulation. We

compared HD-tDCS over M1 versus PMd and showed that PMd may be a potential alternative

neural target to modulate motor network excitability and neural complexity with less variability

that what was observed following M1 HD-tDCS. These results also suggest that multiscale entropy

may be a sensitive measure of changes in neuronal excitability following noninvasive brain

stimulation within a single region and may be able to detect changes not detected by more standard

functional connectivity measures. We further showed that response to PMd HD-tDCS may be

predicted by the integrity of the cortical spinal tract originating from the targeted brain area. Finally,

we demonstrate that conventional methods for defining tDCS targets such as M1 and PMd may be

influenced by large inter-individual variability in the motor hotspot location. Future work should

carefully consider and account for the actual, versus assumed, stimulated location. Altogether, these

results open new considerations for improving the efficacy of tDCS on modulating the motor

network and suggest that individualized targeting based on each individual's brain structure and

function may result in more reliable excitability changes.

Data availability

The unthresholded statistical maps used to generate the motor network (i.e., ICA analyses) are stored on the NEUROVAULT repository (https://neurovault.org/collections/UTZRKARR/). The raw data supporting these analyses will be made available following reasonable requests to the corresponding author. These data are not publicly available due to IRB restrictions.

Fundings

This work was supported by our funding sources, including several National Institutes of Health grants that supported this work (NIH K01HD091283, R61MH110526 and R01MH111896) and the University of Southern California Provost's Postdoctoral Scholar Research Grant.

Acknowledgments

We would like to thank all the study participants, as well as Katherin Martin for her help during the MRI acquisition.

Author contributions

S-L L and S.L. conceived and designed the study with input from D.J.J.W., K.J., and N.S.,

S.L., M.J. K.I., A.S. acquired the data,

S.L. M.J., K.J., Y.Q., R.C., Y.S., and S.L.L. contributed to the data analyses and interpretation,

S.L. and S-L.L. wrote the paper with contributions from all the authors,

All the authors agreed with the submitted version or the paper.

Competing interests

The authors declare no competing interests.

Corresponding author:

Correspondence to Sook-Lei Liew

REFERENCES

Ahdab, R, Ayache SS, Brugieres P, Farhat WH and Lefaucheur JP. 2016. "The Hand Motor Hotspot is not Always Located in the Hand Knob: A Neuronavigated Transcranial Magnetic Stimulation Study." Brain Topogr **29**(4): 590-597.

Amadi, U, Ilie A, Johansen-Berg H and Stagg CJ. 2014. "Polarity-specific effects of motor transcranial direct current stimulation on fMRI resting state networks." Neuroimage **88**: 155-161.

Ammann, C, Lindquist MA and Celnik PA. 2017. "Response variability of different anodal transcranial direct current stimulation intensities across multiple sessions." Brain Stimul **10**(4): 757-763.

Antal, A, Chaieb L, Moliadze V, Monte-Silva K, Poreisz C, Thirugnanasambandam N, Nitsche MA, Shoukier M, Ludwig H and Paulus W. 2010. "Brain-derived neurotrophic factor (BDNF) gene polymorphisms shape cortical plasticity in humans." Brain Stimul **3**(4): 230-237.

Apsvalka, D, Ramsey R and Cross ES. 2018. "Anodal tDCS over Primary Motor Cortex Provides No Advantage to Learning Motor Sequences via Observation." Neural plasticity **2018**: 1237962.

Archer, DB, Vaillancourt DE and Coombes SA. 2018. "A Template and Probabilistic Atlas of the Human Sensorimotor Tracts using Diffusion MRI." Cereb Cortex **28**(5): 1685-1699.

Barbas, H and Pandya DN. 1987. "Architecture and frontal cortical connections of the premotor cortex (area 6) in the rhesus monkey." J Comp Neurol **256**(2): 211-228.

Bestmann, S and Krakauer JW. 2015. "The uses and interpretations of the motor-evoked potential for understanding behaviour." Exp Brain Res **233**(3): 679-689.

Brunoni, AR, Kemp AH, Shiozawa P, Cordeiro Q, Valiengo LC, Goulart AC, Coprerski B, Lotufo PA, Brunoni D, Perez AB, Fregni F and Bensenor IM. 2013. "Impact of 5-HTTLPR and BDNF polymorphisms on response to sertraline versus transcranial direct current stimulation: implications for the serotonergic system." Eur Neuropsychopharmacol **23**(11): 1530-1540.

Burman, KJ, Bakola S, Richardson KE, Reser DH and Rosa MG. 2014. "Patterns of cortical input to the primary motor area in the marmoset monkey." J Comp Neurol **522**(4): 811-843.

Cabeen RP, Laidlaw DH and AW T. 2018. "*Quantitative Imaging Toolkit: Software for Interactive 3D Visualization, Data Exploration, and Computational Analysis of Neuroimaging Datasets*." Proc International Society for Magnetic Resonance in Medicine (ISMRM) 2854.

Cabeen, RP, Bastin ME and Laidlaw DH. 2016. "Kernel regression estimation of fiber orientation mixtures in diffusion MRI." Neuroimage **127**: 158-172.

Caparelli-Daquer, EM, Zimmermann TJ, Mooshagian E, Parra LC, Rice JK, Datta A, Bikson M and Wassermann EM. 2012. "A pilot study on effects of 4x1 high-definition tDCS on motor cortex excitability." Conf Proc IEEE Eng Med Biol Soc **2012**: 735-738.

Chouinard, PA and Paus T. 2006. "The primary motor and premotor areas of the human cerebral cortex." The Neuroscientist : a review journal bringing neurobiology, neurology and psychiatry **12**(2): 143-152.

Costa, M, Goldberger AL and Peng CK. 2005. "Multiscale entropy analysis of biological signals." Phys Rev E Stat Nonlin Soft Matter Phys **71**(2 Pt 1): 021906.

Cunningham, DA, Varnerin N, Machado A, Bonnett C, Janini D, Roelle S, Potter-Baker K, Sankarasubramanian V, Wang X, Yue G and Plow EB. 2015. "Stimulation targeting higher motor areas in stroke rehabilitation: A proof-of-concept, randomized, double-blinded

placebo-controlled study of effectiveness and underlying mechanisms." Restor Neurol Neurosci **33**(6): 911-926.

Diedenhofen, B and Musch J. 2015. "cocor: a comprehensive solution for the statistical comparison of correlations." PLoS One **10**(3): e0121945.

Dum, RP and Strick PL. 1991. "The origin of corticospinal projections from the premotor areas in the frontal lobe." J Neurosci **11**(3): 667-689.

Dum, RP and Strick PL. 2005. "Frontal lobe inputs to the digit representations of the motor areas on the lateral surface of the hemisphere." J Neurosci **25**(6): 1375-1386.

Etkin, A, Egner T and Kalisch R. 2011. "Emotional processing in anterior cingulate and medial prefrontal cortex." Trends in cognitive sciences **15**(2): 85-93.

Fedorov, A, Billet E, Prastawa M, Gerig G, Radamanesh A, Warfield S, K.,, Kikinis R and Chrisochoides N. 2008. Evaluation of Brain MRI Alignment with the Robust Hausdorff Distance Measures. , Springer, Berlin, Heidelberg. **5358**.

Fischl, B and Dale AM. 2000. "Measuring the thickness of the human cerebral cortex from magnetic resonance images." Proc Natl Acad Sci U S A **97**(20): 11050-11055.

Fujiyama, H, Hinder MR and Summers JJ. 2013. "Functional role of left PMd and left M1 during preparation and execution of left hand movements in older adults." J Neurophysiol **110**(5): 1062-1069.

Gbadeyan, O, Steinhauser M, McMahon K and Meinzer M. 2016. "Safety, Tolerability, Blinding Efficacy and Behavioural Effects of a Novel MRI-Compatible, High-Definition tDCS Set-Up." Brain stimulation **9**(4): 545-552.

Giordano, J, Bikson M, Kappenman ES, Clark VP, Coslett HB, Hamblin MR, Hamilton R, Jankord R, Kozumbo WJ, McKinley RA, Nitsche MA, Reilly JP, Richardson J, Wurzman R and

Calabrese E. 2017. "Mechanisms and Effects of Transcranial Direct Current Stimulation." Dose-response : a publication of International Hormesis Society **15**(1): 1559325816685467.

Godschalk, M, Mitz AR, van Duin B and van der Burg H. 1995. "Somatotopy of monkey premotor cortex examined with microstimulation." Neurosci Res **23**(3): 269-279.

Graziano, MS, Taylor CS and Moore T. 2002a. "Complex movements evoked by microstimulation of precentral cortex." Neuron **34**(5): 841-851.

Graziano, MS, Taylor CS, Moore T and Cooke DF. 2002b. "The cortical control of movement revisited." Neuron **36**(3): 349-362.

Guerra, A, Lopez-Alonso V, Cheeran B and Suppa A. 2017. "Solutions for managing variability in non-invasive brain stimulation studies." Neurosci Lett.

Hardwick, RM, Rottschy C, Miall RC and Eickhoff SB. 2013. "A quantitative meta-analysis and review of motor learning in the human brain." Neuroimage **67**: 283-297.

Heisz, JJ, Shedden JM and McIntosh AR. 2012. "Relating brain signal variability to knowledge representation." Neuroimage **63**(3): 1384-1392.

Hordacre, B, Moezzi B, Goldsworthy MR, Rogasch NC, Graetz LJ and Ridding MC. 2017. "Resting state functional connectivity measures correlate with the response to anodal transcranial direct current stimulation." Eur J Neurosci **45**(6): 837-845.

Kang, J, Cai E, Han J, Tong Z, Li X, Sokhadze EM, Casanova MF and Ouyang G. 2018. "Transcranial Direct Current Stimulation (tDCS) Can Modulate EEG Complexity of Children With Autism Spectrum Disorder." Front Neurosci **12**: 201.

Kantak, SS, Stinear JW, Buch ER and Cohen LG. 2012. "Rewiring the brain: potential role of the

premotor cortex in motor control, learning, and recovery of function following brain

injury." Neurorehabilitation and neural repair **26**(3): 282-292.

Kuo, HI, Bikson M, Datta A, Minhas P, Paulus W, Kuo MF and Nitsche MA. 2013. "Comparing

cortical plasticity induced by conventional and high-definition 4 x 1 ring tDCS: a

neurophysiological study." Brain Stimul **6**(4): 644-648.

Lafontaine, MP, Lacourse K, Lina JM, McIntosh AR, Gosselin F, Theoret H and Lippe S. 2016.

"Brain signal complexity rises with repetition suppression in visual learning." Neuroscience

326: 1-9.

Lefebvre, S, Dricot L, Laloux P, Desfontaines P, Evrard F, Peeters A, Jamart J and Vandermeeren

Y. 2017. "Increased functional connectivity one week after motor learning and tDCS in

stroke patients." Neuroscience **340**: 424-435.

Lefebvre, S and Liew SL. 2017. "Anatomical Parameters of tDCS to Modulate the Motor System

after Stroke: A Review." Front Neurol **8**: 29.

Li, X, Zhu Z, Zhao W, Sun Y, Wen D, Xie Y, Liu X, Niu H and Han Y. 2018. "Decreased resting-

state brain signal complexity in patients with mild cognitive impairment and Alzheimer's

disease: a multiscale entropy analysis." Biomed Opt Express **9**(4): 1916-1929.

Liang, WK, Lo MT, Yang AC, Peng CK, Cheng SK, Tseng P and Juan CH. 2014. "Revealing the

brain's adaptability and the transcranial direct current stimulation facilitating effect in

inhibitory control by multiscale entropy." Neuroimage **90**: 218-234.

Liew, SL, Thompson T, Ramirez J, Butcher PA, Taylor JA and Celnik PA. 2018. "Variable Neural

Contributions to Explicit and Implicit Learning During Visuomotor Adaptation." Frontiers

in neuroscience **12**: 610.

Lin, RL, Douaud G, Filippini N, Okell TW, Stagg CJ and Tracey I. 2017. "Structural Connectivity

Variances Underlie Functional and Behavioral Changes During Pain Relief Induced by

Neuromodulation." Sci Rep **7**: 41603.

Lopez-Alonso, V, Cheeran B, Rio-Rodriguez D and Fernandez-Del-Olmo M. 2014. "Inter-

individual variability in response to non-invasive brain stimulation paradigms." Brain

stimulation **7**(3): 372-380.

Lopez-Alonso, V, Fernandez-Del-Olmo M, Costantini A, Gonzalez-Henriquez JJ and Cheeran B.

2015. "Intra-individual variability in the response to anodal transcranial direct current

stimulation." Clin Neurophysiol **126**(12): 2342-2347.

Lopez-Alonso, V, Liew SL, Fernandez Del Olmo M, Cheeran B, Sandrini M, Abe M and Cohen

LG. 2018. "A Preliminary Comparison of Motor Learning Across Different Non-invasive

Brain Stimulation Paradigms Shows No Consistent Modulations." Front Neurosci **12**: 253.

Mayka, MA, Corcos DM, Leurgans SE and Vaillancourt DE. 2006. "Three-dimensional locations

and boundaries of motor and premotor cortices as defined by functional brain imaging: a

meta-analysis." Neuroimage **31**(4): 1453-1474.

McDonough, IM and Nashiro K. 2014a. "Network complexity as a measure of information

processing across resting-state networks: evidence from the Human Connectome Project."

Front Hum Neurosci **8**: 409.

McDonough, IM and Nashiro K. 2014b. "Network complexity as a measure of information

processing across resting-state networks: evidence from the Human Connectome Project."

Frontiers in human neuroscience **8**: 409-409.

Mondini, V, Mangia AL and Cappello A. 2018. "Single-session tDCS over the dominant

hemisphere affects contralateral spectral EEG power, but does not enhance neurofeedback-

guided event-related desynchronization of the non-dominant hemisphere's sensorimotor rhythm." PLoS One **13**(3): e0193004.

Neva, JL, Vesia M, Singh AM and Staines WR. 2014. "Modulation of left primary motor cortex excitability after bimanual training and intermittent theta burst stimulation to left dorsal premotor cortex." Behav Brain Res **261**: 289-296.

Nitsche, MA, Doemkes S, Karakose T, Antal A, Liebetanz D, Lang N, Tergau F and Paulus W. 2007. "Shaping the effects of transcranial direct current stimulation of the human motor cortex." J Neurophysiol **97**(4): 3109-3117.

Nitsche, MA and Paulus W. 2000. "Excitability changes induced in the human motor cortex by weak transcranial direct current stimulation." J Physiol **527 Pt 3**: 633-639.

Nuzum, ND, Hendy AM, Russell AP and Teo WP. 2016. "Measures to Predict The Individual Variability of Corticospinal Responses Following Transcranial Direct Current Stimulation." Front Hum Neurosci **10**: 487.

Puri, R, Hinder MR and Summers JJ. 2015. "The BDNF Val66Met polymorphism modulates motor cortex plasticity, induced by prolonged anodal transcranial direct current stimulation, in older adults." Brain Stimulation: Basic, Translational, and Clinical Research in Neuromodulation **8**(2): 336.

Rizzo, V, Siebner HR, Modugno N, Pesenti A, Munchau A, Gerschlager W, Webb RM and Rothwell JC. 2004. "Shaping the excitability of human motor cortex with premotor rTMS." J Physiol **554**(Pt 2): 483-495.

Schambra, HM, Ogden RT, Martinez-Hernandez IE, Lin X, Chang YB, Rahman A, Edwards DJ and Krakauer JW. 2015a. "The reliability of repeated TMS measures in older adults and in patients with subacute and chronic stroke." Frontiers in cellular neuroscience **9**: 335.

Schambra, HM, Ogden RT, Martinez-Hernandez IE, Lin X, Chang YB, Rahman A, Edwards DJ and Krakauer JW. 2015b. "The reliability of repeated TMS measures in older adults and in patients with subacute and chronic stroke." Front Cell Neurosci **9**: 335.

Schonbrodt, FD, Wagenmakers EJ, Zehetleitner M and Perugini M. 2017. "Sequential hypothesis testing with Bayes factors: Efficiently testing mean differences." Psychological methods **22**(2): 322-339.

Sehm, B, Kipping J, Schafer A, Villringer A and Ragert P. 2013. "A Comparison between Uni- and Bilateral tDCS Effects on Functional Connectivity of the Human Motor Cortex." Front Hum Neurosci **7**: 183.

Shaheiwola, N, Zhang B, Jia J and Zhang D. 2018. "Using tDCS as an Add-On Treatment Prior to FES Therapy in Improving Upper Limb Function in Severe Chronic Stroke Patients: A Randomized Controlled Study." Front Hum Neurosci **12**: 233.

Shi, Y and Toga AW. 2017. "Connectome imaging for mapping human brain pathways." Mol Psychiatry **22**(9): 1230-1240.

Shin, SS and Pelled G. 2017. "Novel Neuromodulation Techniques to Assess Interhemispheric Communication in Neural Injury and Neurodegenerative Diseases." Front Neural Circuits **11**: 15.

Smith, RX, Yan L and Wang DJ. 2014. "Multiple time scale complexity analysis of resting state FMRI." Brain Imaging and Behavior **8**(2): 284-291.

Stinear, CM, Barber PA, Coxon JP, Verryt TS, Acharya PP and Byblow WD. 2009. "Repetitive stimulation of premotor cortex affects primary motor cortex excitability and movement preparation." Brain Stimul **2**(3): 152-162.

Thura, D and Cisek P. 2014. "Deliberation and commitment in the premotor and primary motor cortex during dynamic decision making." Neuron **81**(6): 1401-1416.

Umilta, MA, Brochier T, Spinks RL and Lemon RN. 2007. "Simultaneous recording of macaque premotor and primary motor cortex neuronal populations reveals different functional contributions to visuomotor grasp." J Neurophysiol **98**(1): 488-501.

Valizadeh, SA, Liem F, Merillat S, Hanggi J and Jancke L. 2018. "Identification of individual subjects on the basis of their brain anatomical features." Sci Rep **8**(1): 5611.

Wagle-Shukla, A, Ni Z, Gunraj CA, Bahl N and Chen R. 2009. "Effects of short interval intracortical inhibition and intracortical facilitation on short interval intracortical facilitation in human primary motor cortex." J Physiol **587**(Pt 23): 5665-5678.

Wang, DJJ, Jann K, Fan C, Qiao Y, Zang YF, Lu H and Yang Y. 2018. "Neurophysiological Basis of Multi-Scale Entropy of Brain Complexity and Its Relationship With Functional Connectivity." Front Neurosci **12**: 352.

White, LE, Andrews TJ, Hulette C, Richards A, Groelle M, Paydarfar J and Purves D. 1997. "Structure of the human sensorimotor system. I: Morphology and cytoarchitecture of the central sulcus." Cerebral cortex **7**(1): 18-30.

Whitfield-Gabrieli, S and Nieto-Castanon A. 2012. "Conn: a functional connectivity toolbox for correlated and anticorrelated brain networks." Brain Connect **2**(3): 125-141.

Wiethoff, S, Hamada M and Rothwell JC. 2014. "Variability in response to transcranial direct current stimulation of the motor cortex." Brain Stimul **7**(3): 468-475.

Wymbs, NF and Grafton ST. 2013. "Contributions from the left PMd and the SMA during sequence retrieval as determined by depth of training." Exp Brain Res **224**(1): 49-58.

1   Yang, AC, Tsai SJ, Lin CP and Peng CK. 2018. "A Strategy to Reduce Bias of Entropy Estimates

2   in Resting-State fMRI Signals." Frontiers in neuroscience **12**: 398.

3   Yousry, TA, Schmid UD, Alkadhi H, Schmidt D, Peraud A, Buettner A and Winkler P. 1997.

4   "Localization of the motor hand area to a knob on the precentral gyrus. A new landmark."

5   Brain **120 (Pt 1)**: 141-157.