Cell-to-cell variation in defective virus expression and effects on host responses during

influenza virus infection

Chang Wang[1], Christian V. Forst[2], Tsui-wen Chou[1], Adam Geber[1], Minghui Wang[2], Wissam

Hamou[2], Melissa Smith[2], Robert Sebra[2], Bin Zhang[2], Bin Zhou[1,#], and Elodie Ghedin[1,3*]

[1]Center for Genomics and Systems Biology, Department of Biology, New York University

[2]Mount Sinai Center for Transformative Disease Modeling, Department of Genetics and Genomic

Sciences, Icahn Institute of Genomics and Multiscale Biology, Icahn School of Medicine at Mount

Sinai

[3]College of Global Public Health, New York University, New York, New York

[#]Current address: Bin Zhou, nmb7@cdc.gov, Influenza Division, National Center for Immunization

and Respiratory Diseases, Centers for Disease Control and Prevention, Atlanta, Georgia

* Corresponding author

E-mail: elodie.ghedin@nyu.edu

ABSTRACT

Virus and host factors contribute to cell-to-cell variation in viral infections and determine the

outcome of the overall infection. However, the extent of the variability at the single cell level and

how it impacts virus-host interactions at a systems level are not well understood. To characterize

the dynamics of viral transcription and host responses, we used single-cell RNA sequencing to

quantify at multiple time points the host and viral transcriptomes of human A549 cells and primary

bronchial epithelial cells infected with influenza A virus. We observed substantial variability of viral

transcription between cells, including the accumulation of defective viral genomes (DVGs) that

impact viral replication. We show a correlation between DVGs and viral-induced variation of the

host transcriptional program and an association between differential induction of innate immune

response genes and attenuated viral transcription in subpopulations of cells. These observations

at the single cell level improve our understanding of the complex virus-host interplay during

influenza infection.

IMPORTANCE

Defective influenza virus particles generated during viral replication carry incomplete viral

genomes and can interfere with the replication of competent viruses. These defective genomes

are thought to modulate disease severity and pathogenicity of the influenza infection. Different

defective viral genomes also introduce another source of variation across a heterogeneous cell

population. Evaluating the impact of defective virus genomes on host cell responses cannot be

fully resolved at the population level, requiring single cell transcriptional profiling. Here we

characterized virus and host transcriptomes in individual influenza-infected cells, including that of

defective viruses that arise during influenza A virus infection. We established an association

between defective virus transcription and host responses and validated interfering and

immunostimulatory functions of identified dominant defective virus genome species *in vitro*. This

study demonstrates the intricate effects of defective viral genomes on host transcriptional

responses and highlights the importance of capturing host-virus interactions at the single-cell level.

INTRODUCTION

The productivity of viral replication at the cell population level is determined by cell-to-cell variation

in viral infection [1]. The genetically diverse nature of RNA viruses and the heterogeneity of host

cell states contribute to this inter-cell variability, which can impact therapeutic applications [2, 3].

Although previous studies of cell-to-cell variation during viral infection have mainly centered on

non-segmented viruses such as poliovirus [2, 4], vesicular stomatitis virus (VSV) [5, 6], or dengue

(DENV) and zika (ZIKV) viruses [7], heterogeneity across cells may be further complicated for

viruses with a segmented genome, such as influenza A virus (IAV). IAV-infected cells display

substantial cell-to-cell variation as it pertains to relative abundance of different viral genome

segments [1, 8], transcripts [9], and their encoded proteins [10], which can result in non-productive

infection in a large fraction of cells.

Defective interfering (DI) particles readily generated during successive high multiplicity of infection

(MOI) in cell culture passages [11-14] and observed in natural infections [15, 16] are likely by-

products of the IAV replication process and have a significant effect on productive infection. DI

viruses have incomplete viral genomes with large internal deletions and they possess the ability

to interfere with the replication of infectious viruses. Because DIs diminish the productivity of

infectious progenies and could impact disease outcome, they are of great interest for therapeutic

and prophylactic purposes (reviewed in [17]). A specific influenza DI (i.e., DI244) was shown to

effectively provide prophylactic and therapeutic protection against IAV infection in mice [18]. One

of the ways influenza DIs are thought to modulate viral infections is through interaction with a

cytosolic pathogen-recognition receptor (PRR), RIG-I, essential for interferon (IFN) induction [19].

Besides enhanced IFN induction during infection with DI-rich influenza virus populations observed

in vitro and *in vivo*, it has been assumed that DI virus could also compete with standard virus for

cellular resources (reviewed in [17, 20]).

While diverse DI viruses can arise during IAV infection [21], the emergence and accumulation of

distinct DIs, as well as other defective virus genomes (DVGs), has not been characterized at a

single cell resolution, although the diversity of DIs present could be contributing to the observed

cell-to-cell variation in host transcription.

We probed viral and host transcriptomes simultaneously in the same cells using single-cell RNA-

seq to monitor host-virus interactions in cultured cells over the course of the infection. This data

established a temporal association between the level of viral transcription and the alteration of

the host transcriptome, and characterized the diversity and accumulation of DVG transcripts.

RESULTS

Cell-to-cell variation in virus gene expression.

To determine how both the viral and host cell transcriptional programs relate to each other over

the course of an influenza infection, we infected two cell types—the adenocarcinomic human

alveolar basal epithelial A549 cell line and human primary bronchial epithelial cells HBEpC—at

high multiplicity of Infection (MOI; 5) with A/Puerto Rico/8/34(H1N1) (PR8) and performed single

cell and bulk RNA-seq expression analyses. A high MOI infection ensures that virtually all the

cells can be rapidly infected, promotes the accumulation of DVGs, and consequently enables the

characterization of both host response and DVG diversity. We first determined the percentage of

reads that uniquely aligned to viral genes from the total number of mapped reads to obtain the

relative abundance of virus transcripts within cells at each time point. Similar to what has been

observed at early stages of infection during a low MOI infection of IAV [9], the relative abundance

of virus transcripts was heterogeneous across cells from both cell types, with 0 to 70% of the total

reads in each cell being derived from virus transcripts and the relative abundance of these

transcripts increasing over time (**Supplementary Fig. 1a**). The same trend was also seen when

analyzing segment-specific virus transcripts within individual cells over the course of the infection

(**Supplementary Fig. 1b**).

Heterogeneity of defective virus segment expression across the cell population.

Since DVGs are known to accumulate in cell culture and can serve as templates for transcription,

we characterized their abundance and diversity by examining gap-spanning reads in the

sequencing data. To quantify the relative abundance of DVGs identified, we calculated the

frequency of each unique DVG transcript type (determined by 5' / 3' gap coordinates) across

single cells and the ratio of these to non-gap-spanning transcripts derived from the corresponding

viral segments (i.e., DVG/FL ratio). Although DVG transcripts could be detected in all viral

segments, those originating from the polymerase segments (i.e., PB2, PB1, and PA) occurred

with the highest abundance and frequency in both cell types over the course of the infection

(**Supplementary Fig. 2**), consistent with observations made in other studies [15, 22]. We thus

focused our analyses on the detection of DVG transcripts derived from those segments. We

collected reads with large internal deletions spanning $\geq$1000 nucleotides (nt) (**Fig. 1a**) and

identified the junction coordinates of these gap-spanning reads. We observed a diverse pool of

DVG transcripts, including some shared with the viral segments (vRNA) from the PR8 stock

(**Supplementary Fig. 3-5**). The sizes for the majority of these DVG transcripts are estimated to

be between 300nt and 1000nt. To evaluate the accumulation of DVG transcripts over the course

of the infection, we compared changes in the ratio of DVG transcripts to full-length (FL) transcripts

derived from a given polymerase segment (DVG/FL) across time points. The DVG/FL ratios

increased significantly between the early (6hpi) and late (24hpi) stages of the infection ($p < 2.2 \times$

10^{-16}) and displayed a high level of heterogeneity across single cells from both cell types

(**Supplementary Fig. 6**).

Interestingly, the junction sites aggregated in specific genomic regions, as they systematically

occurred in a high percentage of cells. We identified two predominant types of DVG transcripts

corresponding to PB2 and PB1 defective segments (**Fig. 1b-c** and **Supplementary Fig. 7-8**).

These same DVG transcripts were present in the stock, increased in prevalence over the course

of the infection, and were conserved in different cell types and at different MOIs (**Supplementary**

Fig. 9-10), indicating likely carryover from the stock virus rather than *de novo* formation in each

cell type. For PA, one particular DVG type that was present in the stock was found in A549 and

MDCK cells at different MOIs, but not in HBEpC cells. Its prevalence was also much lower than

for PB2 and PB1 DVGs (**Supplementary Fig. 11-12**). The dominant DVG PB2 and PB1

transcripts derived from corresponding defective segments in both the stock and the infected cells

showed stable relative abundance across individual cells over the course of the infection,

suggesting the persistence of the defective viral segments.

To determine if any of the inferred defective polymerase segments could lead to the generation

of defective interfering particles (DIs), we tested their interference potential experimentally at the

bulk level. We generated clonal PB8-DI162 and PR8-DI222 viruses carrying the deletion sites in

the PB2 segment identified as those in two dominant PB2-DVG species, and then co-infected

each of the DI viruses with PR8 wild-type (WT) virus in A549 cells. Both PR8-DI162 and PR8-

DI222 viruses inhibited the productivity of WT virus as well as PR8-DI244, known to be an

effective DI [18] (**Fig. 1d**). To confirm that inhibition is specific to the DIs and not due to

competition between co-infecting strains, we also tested a full-length PR8 virus with a frame-

shifting point mutation in the PB2 gene. Co-infection with this mutant virus did not affect the

productivity of WT virus (**Fig. 1d**). The inhibitory effect of PR8-DI222 virus could still be detected

at a DI/WT ratio of 1:1 (**Fig. 1d**). This confirmed the interfering ability of the two types of defective

PB2 segments identified in the single cell data. The interference test was not intended to mimic a

single-cell scenario given the difficulties in recreating single-cell infection conditions at a bulk level.

Cell-to-cell variation in concurrent virus-host transcriptional changes over the course of

the infection

Viral infection can trigger massive changes in the host transcriptional program, including the

activation of the interferon (IFN) response. Since the induction of IFNs is also subject to cell-to-

cell variation [23], we first evaluated the expression frequency of type I (i.e., IFN beta) and III (i.e.,

IFN lambda) IFNs to determine the extent of the variation during infection. While these were

significantly differentially expressed over the course of the infection at the population level, as

measured by bulk RNA-seq (**Fig. 2a**), the expression of IFNs was only detected in less than 3%

of cells until the late stage of infection, although this proportion increased over time (**Fig. 2b** and

Fig. 2c). The same expression profile was observed for a number of IFN-stimulated genes (ISGs),

such as *RSAD2*, *CXCL10*, *GBP4*, *GBP5*, *IDO1*, and *CH25H* in A549 cells, and *IFI44L*, *CMPK2*,

IFIT1, *BST2*, *OASL*, and *XAF1* in HBEpC (**Fig. 2**). In silico pooling of the single cell data to mimic

the population level measurement resembles the bulk RNA-seq data, thus excluding the

possibility of substantial technical limitation of single-cell RNA-seq (**Fig. 2a**).

To evaluate cell-to-cell variation in the host response manifested by the alteration in the host

transcriptome following IAV infection, we performed un-supervised cell clustering solely using the

host transcriptional data. We then annotated single cells in each subpopulation with information

about the relative abundance of viral transcripts and the DVG/FL ratio. We observed a correlative

pattern between the distribution of the relative abundance of virus transcripts and the host

transcriptional landscape across cells within a population, as cells with a high abundance of virus

transcripts formed a separate cluster by 12hpi in both cell types and this population structure,

marked by a drastic difference in the level of viral transcripts between clusters, was retained at

24hpi (**Fig. 3a** and **Fig. 3b**). However, cell-cycle status, which contributes to the heterogeneity of

the cell population by 12hpi and the formation of clusters in distinct cell-cycle stages

(**Supplementary Fig. 13a** and **Supplementary Fig. 13b**) with a similar abundance of virus

transcripts, does not seem to have a significant impact on population structure in A549 cells at

24hpi. Consistent with previous reports on G0/G1 cell-cycle arrest [24, 25], at 24hpi we observed

a drastic increase in the proportion of cells in the G0/G1 phase as compared to earlier time-points

(**Supplementary Fig. 13c**). Interestingly, this same pattern was not observed in HBEpC cells at

the same time-point; in these cells the host and viral transcriptome landscape at 24hpi is similar

to that in A549 cells at 12hpi, suggesting a temporal difference in host response and viral

replication between cell types.

To characterize host genes driving the changes associated with viral transcription, we assessed

the over-representation of Gene Ontology (GO) terms associated with differentially expressed

genes in each subpopulation of A549 and HBEpC cells compared to the rest of the cells at 24hpi.

In two subpopulations of A549 cells with a high relative abundance of virus transcripts (clusters 0

and 1 in **Fig. 4a**), there is an enrichment of genes involved in viral transcription, RNA processing,

translation, SRP-dependent co-translational protein targeting to the membrane, and mitochondrial

electron transport (**Fig. 4a**). These GO terms are also associated with genes highly expressed in

HBEpC cells with the highest abundance of virus transcripts at 24hpi (cluster 3 in **Fig. 4b**; p < 2.2

x 10^{-16} compared to the other clusters). In contrast, genes involved in antiviral responses, such

as the type I IFN signaling pathway and the negative regulation of viral genome replication, are

highly expressed in the other two subpopulations of A549 cells with a similar or severely reduced

relative abundance of virus transcripts (clusters 3 and 2, respectively in **Fig. 4a**). However, some

antiviral genes are differentially induced in cells with different levels of viral transcription. For

example, type I and III IFNs, as well as a subset of ISGs, are highly expressed in cluster 3, and

another subset of ISGs are highly expressed in cluster 2. In HBEpC cells, the antiviral responses,

such as the type I IFN signaling pathway, are primarily observed in a cluster that has a low

abundance of virus transcripts (cluster 2 in **Fig. 4b**) and is mostly comprised of cells in the G0/G1

phase (**Supplementary Fig. 13b** at 24hpi), as seen for A549 cells at 12hpi (**Supplementary Fig.**

13a and **Supplementary Fig. 14**).

To further understand how viral and host transcriptional activities are coordinated in clusters of

cells that are subjected to different levels of viral stimuli, we performed gene co-expression

network analyses using multiscale embedded gene co-expression network analysis (MEGENA)

[26] and subsequent correlation analyses to directly characterize virus-host interactions. This was

done for each cluster individually for A549 and HBEpC cells collected at 24hpi to identify modules

of co-expressed genes representing coherent functional pathways. The relevance of a module to

viral infection was evaluated by correlating the first principal component of a module with the

relative abundance of virus transcripts. Consistent with the processes identified in the GO term

enrichment approach using differentially expressed genes identified within a given cluster of cells

(**Fig. 4**), those related to the host cell cycle and viral life cycle, including transcription, translation,

RNA processing, metabolic process, and protein localization to membrane, were enriched in the

modules that were correlated with viral transcription and identified in the clusters of A549 cells

with a high abundance of virus transcripts (**Supplementary Fig. 15**). Conversely, the host innate

immune response (e.g., response to type I IFN) was associated with viral transcription-correlated

modules in cells with the lowest abundance of virus transcripts (cluster 2 of A549 cells at 24hpi in

Fig. 5a). For example, module M100 negatively correlated with relative abundance of virus

transcripts (ρ = -0.262, p = 2.9 x 10^{-5}) in cluster 2 of A549 cells has genes—including *ISG20,*

RNF213, IFI35, STAT1, *RBCK1*, *IFI27*, *OAS2*, *OAS3*, *XAF1*, *STAT2*, *IFIT5*—that are involved in

the innate antiviral response and that were significantly highly expressed, as compared to their

expression in other cells of the same population (**Fig. 5b**). In HBEpC cells, viral transcription-

correlated modules, such as M227, enriched for type I IFN signaling genes (e.g., *ISG15*, *MX1,*

MX2, IFI35, IFI27, IFIT1, IFITM3, OAS1, OASL, ISG20.) and positively correlated with relative

abundance of virus transcripts (ρ = 0.103, p = 0.021; **Fig. 5c** and **5d**), were detected in one cluster

of cells with a low abundance of virus transcripts (cluster 2 of HBEpC cells at 24hpi in **Fig. 5c**).

Modules correlated with viral transcription in the other two clusters of HBEpC cells with a low

abundance of virus transcripts (clusters 0 and 1 in **Supplementary Fig. 16**) were enriched for

genes associated with the host cell cycle, consistent with observations in A549 cells at 12hpi

(**Supplementary Fig. 14**), while the viral transcription correlated modules in HBEpC cells with a

high abundance of virus transcripts (cluster 3 in **Supplementary Fig. 16**) were enriched for genes

involved in the ER-associated ubiquitin dependent protein catabolic process, small molecule

metabolic process, vesicle mediated transport, etc. Overall, by identifying modules of genes that

have a direct correlation with virus transcript abundance, the gene co-expression network analysis

further demonstrates a complex association between virus transcript abundance and host

responses, consistent with the pattern revealed by GO term analysis of highly expressed genes

in individual clusters, and thus provides additional evidence of complex virus-host interactions.

Differential expression of host genes linked to defective viral genomes

Since DVGs are known to play an important role in IFN induction and inhibition of viral replication

(reviewed in [20]), we determined whether the accumulation of DVGs was associated with

attenuated viral transcription and induction of innate immune antiviral response genes. We

observed a subtle but consistent inverse trend between the relative abundance of virus transcripts

and the PB2 and PB1 DVG/FL ratios at 12hpi in A549 cells and at 24hpi in HBEpC cells (**Fig. 6a-**

d). The cluster of cells with the highest abundance of virus transcripts (cluster 1 of A549 cells at

12hpi in **Fig. 6a** and cluster 3 of HBEpC cells at 24hpi in **Fig. 6b**) has the lowest level of PB2 and

PB1 DVG/FL ratios (**Fig. 6c-d**; $p < 2.2 \times 10^{-16}$ comparing PB2 and PB1 DVG/FL ratios in cluster

1 of A549 cells or cluster 3 of HBEpC cells against the other clusters). Similarly, in A549 cells at

24hpi, we see the lowest abundance of virus transcripts in cells (cluster 2 in **Fig. 6e**) that have a

higher level of PB2 DVG/FL ratios (cluster 2 in **Fig. 6f**; $p < 2.2 \times 10^{-16}$ compared to clusters 0 and

1) and the highest level of PB1 DVG/FL ratios ($p = 1.943 \times 10^{-9}$ comparing cluster 2 against the

other clusters). Moreover, innate immune response genes are highly expressed in cells with either

an elevated DVG/FL ratio for both DVG PB2 and PB1 transcripts and a low abundance of virus

transcripts (cluster 2 of A549 cells at 24 hpi in **Fig. 6e-f**), or an elevated DVG/FL ratio for PB2

transcripts and a relatively high abundance of virus transcripts (cluster 3 in **Fig. 6e-f**; $p = 4.036 \times$

10^{-11} compared to clusters 0 and 1 in **Fig. 6f**). It suggests that both DVG and FL transcripts could

play a role in stimulating the innate immune response and that the enrichment of DVG transcripts

in the total virus transcript pool is associated with strong immunostimulation, independent of viral

transcription levels. Notably, the abundance of two dominant and ubiquitous PB2-DVG transcripts

in the total virus transcript pool varies between cells and across clusters. The DVG PB2 transcripts

carrying the same deletion sites as for PR8-DI222 were highly abundant in the cluster of cells with

the lowest abundance of virus transcripts (cluster 2 of A549 cells at 24 hpi in **Fig. 6g**; p < 2.2 x

10^{-16} compared to the other clusters), while the other DVG PB2 transcripts corresponding to PR8-

DI162 were highly abundant in both cluster 2 and cluster 3 (**Fig. 6g**; p = 2.067 x 10^{-10} and p =

0.001241, respectively, compared to clusters 0 and 1), suggesting a potential difference in the

induction of innate immune response genes by different DVG species. We did not detect a

difference in the relative abundance of defective PA transcripts among cells with different levels

of viral transcription (**Supplementary Fig. 17**).

Given the enrichment of PB2 DVGs and the over-expression of ISGs in cluster 2 of A549 cells at

24hpi compared to the rest of the cells in the same population, we experimentally validated the

immunostimulatory impact of PB2 DVGs on the host cells. We infected A549 cells with each of

our 3 PR8-DIs (PR8-DI162, -DI222, or -DI244) or with WT virus, and quantified the expression

level of the influenza virus M gene and of ISGs that were shown to be significantly highly

expressed in cluster 2 of A549 cells, including *IFI35, IFI27*, and *MX1*, the type I (IFN beta) and III

(IFN lambda) IFNs that were shown to be significantly highly expressed in cluster 3 of A549 cells

(**Supplementary Fig. 18a and 18b**). Due to their limited ability to replicate without a helper virus,

DI viruses cultured on their own exhibit a significantly lower level of viral transcription compared

to WT virus (**Supplementary Fig. 18c**). Despite the lower replication of DIs, there was a higher

induction of ISGs in DI-infected cells compared to cells infected with WT virus, while there was a

comparable or marginally higher induction of IFN beta (IFN-ß) and IFN lambda (IFN-λ) in the WT-

infected cells (**Supplementary Fig. 18c**). Together, these data demonstrate that at the late stage

of infection PB2 DVGs have a strong immunostimulatory effect on the host cells manifested by a

significant induction of ISGs even at very low levels of viral transcription and independent of type

I and III IFN expression levels.

DISCUSSION

The goal of this study was to quantitatively and qualitatively characterize viral and host factors that contribute to cell-to-cell transcriptional variation observed during IAV infection. We identified DVGs as potentially important factors in the temporal variation of the host cell transcriptome. The substantial cell-to-cell variation in viral replication and host response that we detected highlight the potential of single-cell virology to provide novel insight into complex virus-host interactions.

The inherent stochasticity of molecular processes, especially during the initial stages of infection, can contribute to the cell-to-cell variation in viral replication [1]. Evidence from studies examining low MOI infections with different virus strains and cell lines suggests that this intrinsic stochasticity can impact viral replication and result in the loss of viral genome segments, transcripts, or proteins [1, 8-10]. Similarly, at different time points of the high MOI infection, we observed cell-to-cell variation in viral transcription, including that in segment-specific transcription. Although the mechanisms leading to this observation are unclear based on our data, it may be explained by the following possibilities as discussed in [10], including (i) the absence of one or more viral genome segments in infecting virions, (ii) a deficiency in intracellular trafficking of incoming viral ribonucleoprotein complexes (vRNPs), (iii) the random degradation of vRNAs prior to transcription, or (iv) mutations resulting in decreased stability of viral mRNAs.

Another source of variability in viral replication stems from the emergence and accumulation of various vRNA species synthesized by the error-prone viral polymerase. Indeed, viral genetic diversity, including amino acid mutations in the NS1 and PB1 proteins, the absence of the NS gene, and an internal deletion in the PB1 gene, contributes to the cell-to-cell variation in the innate immune response as shown in a recent report on a low MOI infection [27]. Although the 3' sequencing strategy we used in our study limits the identification of mutations in the full-length segments, our experimental setup provides a unique opportunity to evaluate the diversity of DVGs

by characterizing the DVG transcripts, since a high MOI infection promotes the accumulation of

DVGs. Late stages of infection also emphasize the impact of accumulated DVGs on the host

response. We identified a diverse pool of DVG transcripts derived from the three polymerase

segments, including several dominant species. Notably, the fact that all the dominant DVG PB2

and PB1 transcripts identified in the infected cells were also present in the virus stock, and that

the dominant DVG PB1 transcripts were detected in an increasing proportion of cells over the

course of infection, suggests potential cellular transmission of DI viruses carrying these DVGs.

However, we cannot exclude the possibility that the regions where the hot spots were located are

most prone to polymerase skipping and that DVG enrichment over the course of the infection

resulted in their abundance above a detection threshold in an increasing number of cells.

The accumulation of DVGs can have consequences on viral replication by directly competing with

the full-length genomes and modulating host innate immune responses. The immune-modulatory

effect of DVGs has been attributed to efficient activation of the IFN induction cascade and antiviral

immunity, a mechanism well-known for the non-segmented negative-sense RNA viruses, such as

the murine parainfluenza virus Sendai (SeV) (reviewed in [20]). Consistent with previous reports,

we detected enrichment of highly expressed genes involved in the IFN response in one

subpopulation of cells with significantly attenuated viral transcription. In this cluster of cells we

also observed that DVG transcripts accumulated to significantly higher levels compared to FL

transcripts. The co-expression network analysis provides an opportunity to identify modules of co-

expressed host genes and directly correlate them with the level of viral transcription revealing

groups of innate response genes that are potentially differentially expressed under the effect of

viral transcription. The observation was further validated in our DI- or WT- infection assay, in

which we detected a significantly higher expression of some ISGs in DI-infected cells independent

of the level of viral transcription and that of IFNs at the late stage of the infection. The IFN mRNA

dose-independent ISG expression in DI-infected cells suggests a potential novel mechanism of

DVG immunostimulation that is independent of IFN induction. Expression of a subset of ISGs can

be induced in the absence of IFN signaling, as shown in other studies [28, 29]. It is likely to be at

least partially attributed to similar and overlapping consensus DNA-binding motifs for the

interferon regulatory factor (IRF) family, most notably IRF3 and IRF7 that are key regulators of

type I IFN production, and for IFN-stimulated gene factor 3 (ISGF3) that promotes the expression

of many ISGs in response to IFN signaling [30-33]. For example, a IRF3 ChIP-seq experiment

with adenovirus-infected human primary bronchial/tracheal epithelial cells demonstrated IRF3

binding to the promoter region of induced ISGs, including MX1 and ISG15, which have the IFN-

stimulated response element (ISRE) bound by IRFs and ISGF3 in the promoter regions [34].

Further investigation of ISG induction in the absence of IFN signaling during DI infection will

elucidate this potential mechanism of DI immunostimulation. In addition, given the fact that some

ISGs could be highly expressed in cells with a high level of viral transcription or a high abundance

of DVGs suggests that other viral factors, besides the accumulation of DVGs, may also play an

important role in triggering the innate immune response. Nevertheless, unlike SeV DVGs that

form "copy-back" structures due to complementary termini generated by "copying back" the

authentic 5' terminus at the 3' terminus [35, 36], IAV DVGs share identical termini with the full-

length genomes. In the copy-back SeV DVGs, a stretch of dsRNA adjacent to a 5'-triphosphate

serves as an effective RIG-I ligand [20]; however, the mechanism of IAV DVGs underlying a more

effective immunostimulation compared to the full-length genomes remains elusive [20]. A

proposed hypothesis is that the unencapsidated replication products of small DVGs can

potentially activate RIG-I if they can reach the cytoplasm [20], as observed *in vitro* and *in vivo*

where short influenza RNA templates can be replicated in the absence of nucleoprotein (NP) [37].

In addition, the interfering and immune-modulatory abilities of DVGs could also be attributed to

DVG-encoded proteins that directly interact with mitochondrial antiviral-signaling (MAVS) and act

independently of RIG-I in IFN induction, as previously reported [38].

Our approach enabled us to characterize and quantify the abundance of diverse DVG species

that accumulated in individual cells. It also allowed us to associate the enrichment of certain DVG

species with the attenuation of viral transcription and the induction of innate immune responses

at single-cell resolution. For example, the accumulation of PB2-DVGs, especially including one

type of dominant PB2-DVG species (PR8-DI222), in one subpopulation of cells was associated

with significantly attenuated viral transcription and a strong host innate immune response

manifested by high expression of some ISGs, although we cannot rule out the effect of other non-

dominant defective PB2-DVGs, DVGs derived from other polymerase segments, or certain

mutations in stimulating host responses. Diverse DVG species can play different roles in shaping

the virus-host interaction. However, further investigation is necessary to elucidate the dynamics

and impact of DVG species distribution during pathogenesis, enhancing the understanding of the

relationship between DVG dynamics and infection outcome. Our characterization of DVGs and

corresponding host responses reveals complex virus-host interactions and an underappreciated

single-cell level of immune-modulation by DVGs. The complex gene expression pattern

underlying the host innate immune response demonstrates the intricate effects of various viral

factors, including—but probably not limited to—the accumulation of DVGs, on shaping infection

outcome at the single-cell level.

MATERIALS AND METHODS

Cells and virus

Human lung adenocarcinoma epithelial A549 cells (ATCC, Virginia, USA) were maintained in Kaighn's modified Ham's F-12 medium (F-12K) supplemented with 10% fetal bovine serum (FBS). Primary human bronchial epithelial cells (HBEpC) (PromoCell, Heidelberg, Germany) were maintained in PromoCell airway epithelial cell growth media with SupplementMix (PromoCell). Madin-Darby canine kidney (MDCK) cells were maintained in minimum essential medium (MEM) supplemented with 5% FBS. Human embryonic kidney epithelial 293T cells (ATCC, Virginia, USA) were maintained in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% FBS. PB2 protein-expressing modified MDCK (AX4/PB2) [39] cells were maintained in MEM supplemented with 5% Newborn Calf Serum (NBCS), 2 µg/ml puromycin and 1 µg/ml blasticidin. 293T and AX4/PB2 cells were co-cultured with DMEM supplemented with 10% FBS. Viral strain A/Puerto Rico/8/34 (H1N1) was plaque purified and propagated in MDCK cells. Viral titers were determined by plaque assay in MDCK cells and sequences confirmed by Illumina MiSeq sequencing.

Infection assays

Subconfluent monolayers of A549 cells in T-25 flasks were washed with A549 infection media (F-12K supplemented with 0.15% bovine serum albumin (BSA) fraction V and 1% antibiotic-antimycotic) and infected with influenza virus at a multiplicity of infection (MOI) of 5 in 0.5 mL pre-cooled A549 infection media. The flasks were incubated at 4°C for 30 minutes with agitation every 5-10 minutes followed by addition of 4.5 mL pre-warmed A549 infection media and transferred to a 37°C, 5% CO_2 incubator for further incubation. HBEpC cells were infected similarly except the HBEpC infection media was PromoCell airway epithelial cell growth media. The inoculum was back-titrated to confirm the desired MOI was used. The virus-infected cells were collected at 6,

12, and 24 hours post infection (hpi) and the mock-infected cells were collected at 12 hpi. There

was no substantial cell death observed at any time point. Cells were extensively washed before

re-suspension in PBS containing 0.04% BSA. A proportion of cells from one of the two duplicate

flasks per time point were subject to 10X Genomics Chromium single-cell library preparation. The

remaining cells in the two flasks were used for conventional bulk RNA-seq library preparation.

Notably, due to a failure in the single-cell library preparation for the PR8-infected HBEpC cells

collected at 12hpi, we replaced this sample by repeating the same infection assay with HBEpC

cells.

Sample preparation, library construction, and sequencing

10X Genomics single cell library preparation with Chromium 3' v2 chemistry was performed

following the manufacturer's protocol. A range of 3,900-7,500 cells were used as input into each

single cell preparation, with a median of ~3,353 cells (range 2,075-5,254) obtained following

sequencing, as described below. Sequencing was performed on the HiSeq 2500 in HighOutput

mode (v2) with one library per lane following the manufacturer's recommended sequencing

configuration (i.e., paired-end read 1: 27 bp, read 2: 99 bp, and i7 index: 8 bp). An average of

59,200 reads per cell were obtained. For bulk mRNA sequencing, total RNA from two replicate

flasks per time point was extracted using RNeasy Mini Kit (Qiagen, Hilden, Germany) with on-

column DNA digestion using RNase-free DNase (Qiagen). The conventional bulk RNA-seq library

preparation was performed using NEBNext Ultra II RNA library prep kit for Illumina following

poly(A) mRNA enrichment. Libraries were multiplexed and sequenced on an Illumina NextSeq

500 in HighOutput 2x75 bp mode (v3). Viral genomic RNA (vRNA) of the virus stocks used in the

infection assay was amplified using multi-segment reverse-transcription PCR (M-RTPCR) [40].

The M-RTPCR product was subject to Illumina library preparation with Nextera DNA library prep

kit and sequenced on the HiSeq 2500 in Rapid 2x150 bp mode (v2).

Upstream computational analyses of single-cell RNA-seq data

The Cell Ranger (v2.1.0) single cell software suite (10X Genomics) was applied to the single-cell RNA-seq data to perform the alignment to the concatenated human (hg19) and influenza A virus (A/Puerto Rico/8/34) reference with STAR (v2.5.3a) [41] and gene counting with the processed UMIs for each cell. The UMI counts for the viral reads with the CIGAR string containing a gap (i.e., "MNM") and different UMI sequences comparing to the ungapped reads were added into the expression matrix later, since they were not considered during gene counting. Given the fact that we performed 3'-end sequencing, transcripts derived from alternative splicing of M and NS segments, encoding M1/M2 and NS1/NEP respectively, could not be distinguished in our analyses, as those transcribed from the same genes share the identical 3'-end.

To identify and exclude low quality cells in the single cell data from the downstream analyses, the following metrics were calculated for each sample and were employed to filter out cells that failed to meet these criteria: 1) remove cells with fewer than 2,000 detected genes; 2) remove cells with an alignment rate less than the mean minus 3 standard deviations; 3) remove cells with a number of reads after log10 transformation not within 3 standard deviations below or above the mean; 4) remove cells with the number of UMI after log10 transformation not within 3 standard deviations below or above the mean; 5) remove cells with the percentage of reads aligned to mitochondrial genes not within 3 standard deviations below or above the mean; 6) remove mock-infected cells with 1 viral read detected. Although we filtered cells based on the size of the total mRNA pool (i.e., total reads) as one of the criteria described above, we also checked the distribution of the size of the host mRNA pool (i.e., host reads) for individual cells. As reported in recent virus-infected single cell studies [9, 42], while virus-induced host shutoff of gene expression [43] is likely to be mainly mediated by host mRNA degradation, it remains unclear what effect viral transcription has on the size of the total mRNA pools. In our study, we observed that virtually all the cells with substantially smaller or larger host mRNA pools–where the number of host reads after log10

transformation was not within 3 standard deviations below or above the mean–were already being

excluded when applying the 6 criteria described above, thus we decided to filter cells based on

these. Approximately 120-250 cells were eliminated from each sample, with a median of 164 cells

per sample. We further removed genes that were detected in fewer than three cells. After initial

cell and gene quality control, the majority of samples had approximately 3,000-3,500 cells left for

downstream analysis. The expression levels of host genes were normalized based on the size of

the host mRNA pool.

Cell clustering with Seurat

An unsupervised cell clustering to identify subtle changes in the population structure after infection

was performed on each time point data separately following the procedures of the Seurat package

(v2.1.0) [44] using the normalized, scaled, and centered host gene expression matrix, without

considering the relative abundance of virus transcripts within individual cells. Briefly, the highly

variable genes with average expression < 4 and dispersion > 1 were used as input for the PCA.

The statistically significant and biological meaningful PCs determined by the built-in jackstraw and

elbow analyses and manually exploration were retained for visualization by t-distributed stochastic

neighbor (t-SNE) and subsequent clustering by a shared nearest neighbor (SNN) graph-based

approach. The legitimacy of the initially identified clusters was validated using the

"ValidateClusters" function in Seurat, which built a support vector machine (SVM) classifier with

significant PCs and then applied the accuracy cutoffs of 0.9 and the minimal connectivity threshold

of 0.001. To identify markers that are differentially expressed among clusters, the "FindMarkers"

function in Seurat was used with the different test options, including "bimod" [45], "poisson",

"negbinom", and "MAST" (v1.4.1) [46]. Only the genes that showed at least 0.25-fold difference

on the log-scale between two groups and were expressed in at least 25% of the cells in either

group were tested for differential expression. Significantly differentially expressed genes for each

cluster (i.e., the markers) were identified by applying the adjusted p-value cut-off of 0.05. Markers

identified by all four methods were retained. GO term over-representation analysis of the up-

regulated markers was performed with the online service DAVID [47, 48] by applying the

Benjamini p-value cut-off of 0.05. To determine the cell cycle stage associated with individual

cells, cell-cycle scoring and assignment were performed with Seurat using the "CellCycleScoring"

function based on the expression of canonical markers [49].

Computational analyses of bulk RNA-seq and virus stock sequencing data

The raw bulk RNA-seq and virus stock sequencing data was first trimmed with trimmomatic (v0.36)

[50] to remove the adaptors and trim off low quality bases. Reads with a minimal length of 36

bases in the trimmed bulk RNA-seq dataset were aligned to the concatenated human (hg19) and

influenza A/Puerto Rico/8/34 (H1N1) reference with STAR (v2.5.3a) [41] with the default

parameters, and counted with featureCounts [51] in the Subread package (v1.5.1) [52], while

reads in the virus stock sequencing dataset were aligned to the influenza A/Puerto Rico/8/34

(H1N1) reference with STAR (v2.5.3a) [41]. Differential expression analysis was performed with

DESeq2 (v1.18.1) [53] and edgeR (v3.20.5) [54, 55] using the bulk RNA-seq and the merged

single-cell RNA-seq data. Host genes with the adjusted p-value < 0.05 were identified as

significantly differentially expressed at each time point.

Deletion junction identification, filtering, and quantification

Reads that aligned to both ends of a viral segment, with each aligned portion comprised of at

least 10 nucleotides in length, were collected. Following UMI de-duplication, reads with junction

coordinates within a 10-nucleotide window were grouped together. We excluded from

downstream analyses reads with junction coordinates that occurred fewer than 10 times in each

sample. To compare quantitatively gap-spanning reads for each segment across cells and

samples, the number of gap-spanning reads was normalized to the total number of non-gap-

spanning reads aligned to a 100nt region centering the coverage peak at the 3' end. For the

polymerase segments, gap-spanning reads were also filtered by the size of deletions (i.e. ≥ 1000

nucleotides). The 1000 nucleotides threshold for the polymerase segments was chosen based on

the distribution of the size of the deletions (**Supplementary Fig. 19**), as the majority of the

deletions are larger than 1000 nucleotides. As cells with low infection (especially those harvested

at the early stage of infection or inoculated at a low MOI) typically have poor coverage for the viral

segments of interest, the DVG/FL ratio in those cells calculated as described above is typically

inflated and may even fail to be calculated because all the viral reads corresponding to a given

segment are gap-spanning or there are no viral reads. To mitigate this effect, we overwrote those

values to 0 in the datasets, including the dataset collected from HBEpC cells at 6hpi.

Generation of PR8-derived DI162, DI222, DI244, and PB2del virus

To generate the PR8 PB2-DI162, -DI222, and -DI244 reverse genetics plasmids, gBlocks Gene

Fragments (Integrated DNA Technologies, California, USA) were ordered using the

corresponding defective PB2 genomic sequences identified in this study (PR8-DI162 and PR8-

DI222) or previously reported (DI244) [18]. To generate the PR8 PB2del plasmid, a point mutation

at nucleotide position 265 was introduced into the PR8 PB2 genomic sequence using the

mutagenesis primer PR8-PB2-stopdel-F (5'-TTCATTTACTCCATTAAGTTTGTCCTTGCTC-3').

The reverse genetics plasmids were cloned into the pBZ61A18 reverse genetics vector as

previously described [56]. To rescue the PR8 DI viruses, each of the sequence-confirmed PB2-

DI or PB2del plasmids was co-transfected into 293T-AX4/PB2 co-cultured cells with the 7 reverse

genetics plasmids for PR8 PB1, PA, HA, NP, NA, M, and NS and a PB2 protein-expression

plasmid using Lipofectamine™ 3000 Transfection Reagent (Invitrogen, California, USA). The

supernatant was collected on day 2 post-infection and passaged twice in the AX4/PB2 cell line,

which expresses the PB2 protein in *trans* [39]. Viral titers were determined by TCID50 assay in

AX4/PB2 cells.

Interference test for PR8-DI162 and PR8-DI222 viruses and validation of their immunostimulatory effects

To test the interfering ability of PR8-DI162 and PR8-DI222 viruses that carry the deletions identified in two dominant PB2-DVG species from the single cell dataset, their inhibitory effects on wild type PR8 virus replication was quantified in cultured cells at different DI/WT ratios. Subconfluent monolayers of A549 cells in 12-well plates were washed with infection media (MEM supplemented with 0.15% BSA fraction V, 1% antibiotic-antimycotic and 1 µg/mL TPCK-treated trypsin) and each well was co-infected with one type of defective virus at a MOI of 5 and the PR8 virus either at a MOI of 0.005 for a DI/WT ratio of 1000:1 or at a MOI of 5 for a DI/WT ratio of 1:1 in 1 mL infection media. For a DI/WT ratio of 1000:1 co-infection, plates were transferred to a 37°C, 5% CO_2 incubator and supernatants were collected at 2 hours post infection, and 1, 2, and 3 days post infection. For a DI/WT ratio of 1:1 co-infection, infection was similar to that used for single-cell RNA-seq, including incubating at 4°C for 30 minutes with agitation every 5-10 minutes after infection followed by adding pre-warmed A549 infection media and transferred to a 37°C, 5% CO_2 incubator for further incubation. Supernatants were collected at 2, 6, 12, 24 hours post infection and titrated as described above. Wild type PR8 yield from collected supernatants was determined by TCID50 assay using MDCK cells, which do not support replication of DI viruses due to the lack of functional PB2 proteins.

To determine if the DI viruses could stimulate a higher expression of some ISGs identified in the network analysis, subconfluent monolayers of A549 cells in 12-well plates were infected with each DI virus (PR8-DI162, -DI222, or -DI244) or WT virus at a MOI of 10 in triplicate. Total RNA from cells collected at 24hpi was extracted using RNeasy Mini Kit (Qiagen, Hilden, Germany). 100ng RNA was subsequently used as template for reverse transcription using SuperScript IV system (Invitrogen, California, USA) with Oligo d(T)$_{20}$ primer. qPCR was performed using PowerUp SYBR

Green (Applied Biosystems, Massachusetts, USA) in triplicate with the following primers: IFI35:

Hs.PT.58.38490206 (IDT, Iowa, USA); IFI27: IFI27fw (5'-GCCTCTGGCTCTGCCGTAGTT-3')

and IFI27rev (5'-ATGGAGGACGAGGCGATTCC-3') [57]; MX1: MX1fw (5'-

CTTTCCAGTCCAGCTCGGCA-3') and MX1rev (5'-AGCTGCTGGCCGTACGTCTG-3') [57];

IFNB: IFNBfw (5'-TCTGGCACAACAGGTAGTAGGC-3') and IFNBrev (5'-

GAGAAGCACAACAGGAGAGCAA-3') [58]; IFNL: IFNLfw (5'-GCCCCCAAAAAGGAGTCCG-3')

and IFNLrev (5'-AGGTTCCCATCGGCCACATA-3') [58]; β-actin (ACTB): ACTBfw (5'-

GAACGGTGAAGGTGACAG-3') and ACTBrev (5'-TTTAGGATGGCAAGGGACT-3') [59]; FLU:

mCDC-FluA-F-PR8 (5'-GACCAATCCTGTCACCTCTGA-3') and mCDC-FluA-R (5'-

AGGGCATTTTGGACAAAGCGTCTA-3'). For all the targets, qPCR parameters according to

manufacturer's recommendation were: 95°C for 10 min and then 45 cycles of 95°C for 15 s, 57°C

for 15 s, and 60°C for 60 s. Fold change of target gene expression was calculated using the

$2^{-\Delta\Delta C_T}$ method normalized to ACTB.

Gene co-expression network analysis

Multiscale Embedded Gene Co-Expression Network Analysis (MEGENA) [26] was performed to

identify host modules of highly co-expressed genes in influenza infection. The MEGENA workflow

comprises 4 major steps: 1) Fast Planar Filtered Network construction (FPFNC), 2) Multiscale

Clustering Analysis (MCA), 3) Multiscale Hub Analysis (MHA), 4) and Cluster-Trait Association

Analysis (CTA). A cutoff of 0.05 after perturbation-based FDR calculation was used. The

differentially expressed genes are identified between virus-infected cells of a particular cluster

and the rest of the cells in the same population determined by a t-test. Only the genes that showed

at least 0.25-fold difference on the log-scale between two groups, and were expressed in at least

25% of the cells in either group, were tested for differential expression. Significantly differentially

expressed genes were identified by applying the adjusted p-value cut-off of 0.05.

Identification of enriched GO terms, key regulators in the host module, relative abundance

of virus transcripts associated with host modules

To functionally annotate gene signatures and gene modules identified in this study, enrichment

analysis was performed of the established gene ontology (GO) categories. The hub genes in each

subnetwork were identified using the adopted Fisher's inverse Chi-square approach in MEGENA;

Bonferroni-corrected p-values smaller than 0.05 were set as the threshold to identify significant

hubs.

The relative abundance of virus transcripts and the DVGs associated with the host modules were

identified using Spearman correlation between the first principal component of the gene

expression in the corresponding module and the relative abundance of viral transcripts.

Significantly associated traits were identified using the Benjamini-Hochberg FDR-corrected p-

value 0.05 as the cutoff.

STATISTICAL ANALYSIS

The statistical significance of the changes in the relative abundance of viral and DVG transcripts

between 3 or more groups of cells was first determined by the one- or two- tailed Kruskal-Wallis

rank sum test, followed by the one-tailed Wilcoxon rank sum test to calculate the pairwise

comparisons. The statistical significance of expression fold changes in the qPCR validation assay

was determined using the two-tailed Student's *t*-test. A p-value of ≤ 0.05 was considered

statistically significant.

CODE AVAILABILITY

The code used to generate all the results is available on Github

(https://github.com/GhedinLab/Single-Cell-IAV-infection-in-monolayer).

DATA AVAILABILITY

Sequencing data that support the findings of this study have been deposited in the Gene Expression Omnibus (GEO) repository with the accession codes GSE118773 (currently private record).

ACKNOWLEDGEMENTS

We thank members of the Ghedin laboratory for feedback and discussion. We thank Tara Rock, Nicholas Rouillard, Olivia Micci-Smith, and Mohammed Khalfan of the Genomics Core Facility at the Center for Genomics and Systems Biology, New York University for sequencing. We thank Dr. Yoshihiro Kawaoka for providing the AX4/PB2 cells and Dr. Peter Palese and Dr. Adolfo Garcia-Sastre for providing the PR8 RGs plasmids. We thank Dr. Meike Dittmann for her feedback on IFNs and ISGs. This work was supported by NIAID/NIH U01 AI111598. This work was also supported in part through the NYU IT High Performance Computing resources, services, and staff expertise.

AUTHOR CONTRIBUTIONS

All authors read and approved the manuscript. E.G. conceived and designed the experiments, supervised research, and wrote the manuscript. B. Zhou conceived and designed the experiments, supervised research, performed the infection assays, and wrote the manuscript. C.W. performed the infection assays and bulk RNA-seq library preparation, analyzed the data, and wrote the manuscript. C.V.F. performed the gene co-expression network reconstruction and module-based functional enrichment analysis and wrote the manuscript. T.C. performed the defective virus generation, the interference and validation assays, and wrote the manuscript. A.G. performed the sequence library preparation and contributed to the analyses. M.W. and B. Zhang contributed to

637   data analyses. W.H., M.S., and R.S. performed the single-cell library preparation and contributed

638   to the analyses.

REFERENCES

1. Heldt FS, Kupke SY, Dorl S, Reichl U, Frensing T. Single-cell analysis and stochastic modelling unveil large cell-to-cell variability in influenza A virus infection. Nature Communications. 2015;6:8938.

2. Guo F, Li S, Caglar MU, Mao Z, Liu W, Woodman A, et al. Single-cell virology: On-chip investigation of viral infection dynamics. Cell Reports. 2017;21(6):1692-704.

3. Raj A, van Oudenaarden A. Nature, nurture, or chance: Stochastic gene expression and its consequences. Cell. 2008;135(2):216-26.

4. Lwoff A, Dulbecco R, Vogt M, Lwoff M. Kinetics of the release of poliomyelitis virus from single cells. Virology. 1955;1(1):128-39.

5. Zhu Y, Yongky A, Yin J. Growth of an RNA virus in single cells reveals a broad fitness distribution. Virology. 2009;385(1):39-46.

6. Timm A, Yin J. Kinetics of virus production from single cells. Virology. 2012;424(1):11-7.

7. Zanini F, Pu S-Y, Bekerman E, Einav S, Quake SR. Single-cell transcriptional dynamics of flavivirus infection. eLife. 2018;7:e32942.

8. Dou D, Hernandez-Neuta I, Wang H, Ostbye H, Qian X, Thiele S, et al. Analysis of IAV replication and co-infection dynamics by a versatile RNA viral genome labeling method. Cell Reports. 2017;20(1):251-63.

9. Russell AB, Trapnell C, Bloom JD. Extreme heterogeneity of influenza virus infection in single cells. eLife. 2018;7:e32303.

10. Brooke CB, Ince WL, Wrammert J, Ahmed R, Wilson PC, Bennink JR, et al. Most influenza a virions fail to express at least one essential viral protein. Journal of Virology. 2013;87(6):3155-62.

11. Henle W, Henle G. Interference of inactive virus with the propagation of virus of influenza A. Science. 1943;98(2534):87-9.

12. von Magnus P. Incomplete Forms of Influenza Virus. Advances in Virus Research Volume 2. Advances in Virus Research1954. p. 59-79.

13. Davis AR, Nayak DP. Sequence relationships among defective interfering influenza viral RNAs. Proceedings of the National Academy of Sciences of the United States of America. 1979;76(7):3092-6.

14. Nayak DP, Chambers TM, Akkina RK. Defective-Interfering (DI) RNAs of Influenza Viruses: Origin, Structure, Expression, and Interference. Current Topics in Microbiology and Immunology. 114. 1st ed. Berlin Heidelberg: Springer-Verlag; 1985. p. 103-51.

15. Saira K, Lin X, DePasse JV, Halpin R, Twaddle A, Stockwell T, et al. Sequence analysis of in vivo defective interfering-like RNA of influenza A H1N1 pandemic virus. Journal of Virology. 2013;87(14):8064-74.

16. Vasilijevic J, Zamarreno N, Oliveros JC, Rodriguez-Frandsen A, Gomez G, Rodriguez G, et al. Reduced accumulation of defective viral genomes contributes to severe outcome in influenza virus infected patients. PLos Pathogens. 2017;13(10):e1006650.

17. Dimmock NJ, Easton AJ. Defective interfering influenza virus RNAs: time to reevaluate their clinical potential as broad-spectrum antivirals? Journal of Virology. 2014;88(10):5217-27.

18. Dimmock NJ, Rainsford EW, Scott PD, Marriott AC. Influenza virus protecting RNA: an effective prophylactic and therapeutic antiviral. Journal of Virology. 2008;82(17):8570-8.

19. Baum A, Sachidanandam R, García-Sastre A. Preference of RIG-I for short viral RNA molecules in infected cells revealed by next-generation sequencing. PNAS. 2010;107(37):16303-8.

20. Killip MJ, Fodor E, Randall RE. Influenza virus activation of the interferon system. Virus Research. 2015;209:11-22.

21. Janda MJ, Davis AR, Nayak DP, De BK. Diversity and generation of defective interfering influenza virus particles. Virology. 1979;95(1):48-58.

22.   Alnaji FG, Holmes JR, Rendon G, Vera JC, Fields C, Martin BE, et al. Sequencing framework for the sensitive detection and precise mapping of defective interfering particle-associated deletions across influenza A and B viruses. Journal of Virology. 2019.

23.   Chen S, Short JA, Young DF, Killip MJ, Schneider M, Goodbourn S, et al. Heterocellular induction of interferon by negative-sense RNA viruses. Virology. 2010;407(2):247-55.

24.   He Y, Xu K, Keiner B, Zhou J, Czudai V, Li T, et al. Influenza A virus replication induces cell cycle arrest in G0/G1 phase. Journal of Virology. 2010;84(24):12832-40.

25.   Jiang W, Wang Q, Chen S, Gao s, Song L, Liu P, et al. Influenza A virus NS1 induces G0/G1 cell cycle arrest by inhibiting the expression and activity of RhoA protein. Journal of Virology. 2013;87(6):3039-52.

26.   Song W-m, Zhang B. Multiscale embedded gene co-expression network analysis. PLOS Computational Biology. 2015;11(11):e1004574.

27.   Russell AB, Elshina E, Kowalsky JR, te Velthuis AJW, Bloom JD. Single-cell virus sequencing of influenza infections that trigger innate immunity. Journal of Virology. 2019;93:e00500-19.

28.   Schmid S, Mordstein M, Kochs G, García-Sastre A, tenOever BR. Transcription factor redundancy ensures induction of the antiviral state. Journal of Biological Chemistry. 2010;285(53):42013-22.

29.   Wang N, DOng Q, Li J, Jangra RK, Fan M, Brasier AR, et al. Viral induction of the zinc finger antiviral protein is IRF3-dependent but NF-κB-independent. Journal of Biological Chemistry. 2010;285(9):6080-90.

30.   Ourthiague DR, Birnbaum H, Ortenlof N, Vargas JD, Wollman R, Hoffmann A. Limited specificity of IRF3 and ISGF3 in the transcriptional innate-immune response to double-stranded RNA. Journal of Leukocyte Biology. 2015;98(1):119-28.

31.   Grandvaux N, Servant MJ, tenOever B, Sen GC, Balachandran S, Barber GN, et al. Transcriptional profiling of interferon regulatory factor 3 target genes: direct involvement in the regulation of interferon-stimulated genes. Journal of Virology. 2002;76(11):5532-9.

32.   Ning S, Pagano JS, Barber GN. IRF7: activation, regulation, modification and function. Genes & Immunity. 2011;12(6):399-414.

33.   Yanai H, Negishi H, Taniguchi T. The IRF family of transcription factors. Oncoimmunology. 2012;1(8):1376-86.

34.   Zemke NR, Berk AJ. The adenovirus E1A C terminus suppresses a delayed antiviral response and modulates Ras signaling. Cell Host Microbe. 2017;22(6):789-800 e5.

35.   Re GG, Gupta KC, Kingsbury DW. Genomic and copy-back 3' termini in Sendai virus defective interfering RNA species. Journal of Virology. 1983;45(2):659-64.

36.   Strahle L, Garcin D, Kolakofsky D. Sendai virus defective-interfering genomes and the activation of interferon-beta. Virology. 2006;351(1):101-11.

37.   Turrell L, Lyall JW, Tiley LS, Fodor E, Vreede FT. The role and assembly mechanism of nucleoprotein in influenza A virus ribonucleoprotein complexes. Nature Communications. 2013;4:1591.

38.   Boergeling Y, Rozhdestvensky TS, Schmolke M, Resa-Infante P, Robeck T, Randau G, et al. Evidence for a novel mechanism of influenza virus-induced type I interferon expression by a defective RNA-encoded protein. PLoS Pathogens. 2015;11(5):e1004924.

39.   Ozawa M, Victor ST, Taft AS, Yamada S, Li C, Hatta M, et al. Replication-incompetent influenza A viruses that stably express a foreign gene. Journal of General Virology. 2011;92:2879-88.

40.   Zhou B, Donnelly ME, Scholes DT, George KS, Hatta M, Kawaoka Y, et al. Single-reaction genomic amplification accelerates sequencing and vaccine production for classical and swine origin human influenza A viruses. Journal of Virology. 2009;83(19):10309-13.

41.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-seq aligner. Bioinformatics. 2013;29(1):15-21.

42. Steuerman Y, Cohen M, Peshes-Yaloz N, Valadarsky L, Cohn O, David E, et al.
Dissection of influenza infection in vivo by single-cell RNA sequencing. Cell Systems.
2018;6(6):679-91 e4.
43. Bercovich-Kinori A, Tai J, Gelbart IA, Shitrit A, Ben-Moshe S, Drori Y, et al. A systematic
view on influenza induced host shutoff. eLife. 2016;5:e18311.
44. Satija R, Farrell JA, Gennert D, Schier AF, Regev A. Spatial reconstruction of single-cell
gene expression data. Nature Biotechnology. 2015;33(5):495-502.
45. McDavid A, Finak G, Chattopadyay PK, Dominguez M, Lamoreaux L, Ma SS, et al. Data
exploration, quality control and testing in single-cell qPCR-based gene expression experiments.
Bioinformatics. 2013;29(4):461-7.
46. Finak G, McDavid A, Yajima M, Deng J, Gersuk V, Shalek AK, et al. MAST: a flexible
statistical framework for assessing transcriptional changes and characterizing heterogeneity in
single-cell RNA sequencing data. Genome Biology. 2015;16:278.
47. Huang DW, Sherman BT, Lempicki RA. Systematic and integrative analysis of large
gene lists using DAVID Bioinformatics Resources. Nature Protocol. 2009;4(1):44-57.
48. Huang DW, Sherman BT, Lempicki RA. Bioinformatics enrichment tools: paths toward
the comprehensive functional analysis of large gene lists. Nucleic Acids Research.
2009;37(1):1-13.
49. Kowalczyk MS, Tirosh I, Heckl D, Rao TN, Dixit A, Haas BJ, et al. Single-cell RNA-seq
reveals changes in cell cycle and differentiation programs upon aging of hematopoietic stem
cells. Genome Research. 2015;25(12):1860-72.
50. Bolger A, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence
data. Bioinformatics. 2014;30(15):2114-20.
51. Liao Y, Smyth G, Shi W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. Bioinformatics. 2014;30(7):923-30.
52. Liao Y, Smyth G, Shi W. The Subread aligner: fast, accurate and scalable read mapping
by seed-and-vote. Nucleic Acids Research. 2013;41(10):e108.
53. Love M, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biology. 2014;15:550.
54. Robinson M, McCarthy D, Smyth G. edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. Bioinformatics. 2010;26(1):139-40.
55. McCarthy D, Chen Y, Smyth G. Differential expression analysis of multifactor RNA-seq
experiments with respect to biological variation. Nucleic Acids Research. 2012;40(10):4288-97.
56. Zhou B, Wentworth DE. Influenza A Virus Molecular Virology Techniques. In: Kawaoka
Y, Neumann G, editors. Influenza Virus Methods in Molecular Biology (Methods and Protocols).
865: Humana Press; 2012.
57. Cheon H, Holvey-Bates EG, Schoggins JW, Forster S, Hertzog P, Imanaka N, et al.
IFNβ-dependent increases in STAT1, STAT2, and IRF9 mediate resistance to viruses and DNA
damage. EMBO Journal. 2013;32(30):2751-63.
58. Zhou B, Li Y, Belser JA, Pearce MB, Schmolke M, Subba AX, et al. NS-based live
attenuated H1N1 pandemic vaccines protect mice and ferrets. Vaccine. 2010;28(50):8015-25.
59. Mayanagi T, Morita T, Hayashi K, Fukumoto K, Sobue K. Glucocorticoid receptor-
mediated expression of caldesmon regulates cell migration via the reorganization of the actin
cytoskeleton. Journal of Biological Chemistry. 2008;283(45):31183-96.
786
787

    FIGURES

Fig 1.

Figure 1. Characterization of DVG transcripts and validation of interfering capacity. (a)

Selection criteria for reads considered to be representatives of DVGs. Given the length of the

reads, we required that each portion of the gap-spanning reads were at least 10 bp in length.

Considering that DVGs typically lose approximately 80% of their original sequences and

considering the distribution in the size of the deletions detected in this study, we also required

that the size of the gap should be at least 1 kb. **(b-c)** Schematic diagram showing two types of

defective viral transcripts derived from **(b)** PB2 and **(c)** PB1 segments, respectively. The caret "^"

denotes a deletion event and the numbers before and following it indicate the coordinates of the

junction sites. **(d)** Interfering capacity of generated defective interfering particles (DIs) carrying

identified deletion sites in the PB2 segment during co-infection with the WT-PR8 virus. A549 cells

were co-infected with WT virus and one type of DI virus at a ratio of 1:1000 or 1:1. Viral titers were

measured by the TCID50 assay on MDCK cells. Four types of defective viruses were tested,

including two DI viruses identified in this study, a previously reported DI244 virus [18] that has

been considered as a potential candidate for antiviral therapy [17], and a PB2 polymerase

defective PR8 virus mutant (i.e., PB2del) that carries 1 nucleotide deletion in the coding sequence

of the PB2 gene. For each type of DI virus, the information about the junction sites was denoted

in parentheses. The error bar representing the standard deviation of the mean was obtained from

three biological replicates. Each panel shows an independent interference assay.

Fig 2.

Figure 2. Comparison of the induction of interferons (IFNs) and interferon-stimulated genes (ISGs) at bulk and single-cell levels. (a) Log2 fold changes of the top 10 differentially

expressed genes with the greatest fold change by 24hpi, including type I and III IFNs and some

ISGs, compared to mock-infected cells detected in A549 and HBEpC cells over the course of the

infection at the bulk level and their log2 fold changes in merged single-cell data that mimics the

bulk level measurements. The results from edgeR are shown. **(b)** The expression of these genes

at the single-cell level over the course of the infection in two cell types. Expression was calculated

as log-transformed UMI counts normalized by the host library size. The violins below the dots are

colored by samples. **(c)** Proportion of cells in which these genes are detected over the course of

the infection in two cell types. The colors are consistent with those in panel (b).

Fig 3.

821 **Figure 3. Comparison of cell clustering patterns and the distribution of viral transcript relative abundance.** PR8-infected **(a)**

822 A549 and **(b)** HBEpC cells harvested at each time point were subjected to un-supervised clustering based on the host transcriptome

823   and visualized on a t-SNE plot. Each dot on a t-SNE plot represents a cell. Cell clustering patterns, in which cells were colored by their

824   cluster identity, are shown in the left panels, and the distribution of the relative abundance of viral transcripts at different time points in

825   individual cells is shown in the right panels.

Fig 4.

Figure 4. Enriched biological process gene ontology (GO) terms in A549 and HBEpC cells at 24hpi. (a-b) Biological process GO terms associated with over-expressed genes in each cluster of (**a**) A549 and (**b**) HBEpC cells. The top 10 over-represented GO terms in each cluster are shown in the upper panel and another over-represented GO term related to type I IFN signaling pathway in clusters 2 of HBEpC cells are shown in the lower panel. Each GO term is denoted by a dot. The color intensity of each dot indicates the fold enrichment of the corresponding GO term and the size corresponds to the ratio of queried genes in the gene set associated with a

832   given GO term. The GO terms enriched in each cluster are color-coded by clusters. The asterisks in the upper panel denote the GO

833   terms enriched in two clusters and their colors indicate the identity of the other cluster besides the one denoted by the color of the GO

834   term. The bar chart above the bubble chart of the GO terms shows the median relative abundance of the viral transcripts across cells

835   in each cluster. The colors of the bar are consistent with that of the GO terms.

Fig 5.

Figure 5. Functional MEGENA modules in clusters of A549 and HBEpC cells at 24hpi. Left panels show enriched biological

process GO terms in the modules correlated with the relative abundance of virus transcripts in **(a)** cluster 2 of A549 cells and **(c)** cluster

2 of HBEpC cells at 24hpi. Each GO term is denoted by a dot. The color intensity of each dot indicates the fold enrichment of the

corresponding GO term and the size corresponds to the -log10 transformed corrected p-value for a given GO term. Red and blue color-

bars above the dot plot denote positive or negative correlation of viral transcription with corresponding modules, respectively. Right

panels show the MEGENA network of modules enriched with the innate immune response, including **(b)** module M100 in cluster 2 of

A549 cells and **(d)** module M227 in cluster 2 of HBEpC cells. Red and blue nodes represent significantly up- or down-regulated genes

compared to the other cells in the population, while light -green nodes denote genes that are not significantly differentially expressed.

Diamond nodes indicate key regulators. The size of the nodes indicates node strength after Multiscale Hub Analysis within the

MEGENA pipeline. Modules that have a significant correlation with the relative abundance of virus transcripts and enriched GO

biological process terms are shown in the GO term plots (**a** and **c**). Module member information for all viral transcription correlated-

modules   can   be   found   on   Github   (https://github.com/GhedinLab/Single-Cell-IAV-infection-in-

monolayer/tree/master/SupplementaryTables).

Fig 6.

Figure 6. Distribution of the relative abundance of viral transcripts and DVG/FL ratio for

DVG PB2 and PB1 transcripts in each cluster of cells. (a, b, and e) Box-plot of the relative

abundance of viral transcripts in each cluster of **(a)** A549 cells at 12hpi, **(e)** 24hpi, and **(b)** HBEpC

cells at 24hpi. **(c, d, and f)** The DVG/FL ratio for DVG PB2 and PB1 transcripts calculated as the

ratio of gap-spanning reads to non-gap-spanning reads aligned to a 3' coverage peak region

derived from the corresponding viral segments in each cluster. **(g)** The DVG/FL ratio for DVG PB2

transcripts carrying deletion sites similar to those in PR8-DI222 and PR8-DI162 viruses in each

cluster of A549 cells at 24hpi. All box plots show the first and third quantiles as the lower and

upper hinges, the median in the center, and 1.5 * inter-quartile range (IQR) from the first and third

quantiles as the whiskers. The significance levels of pairwise comparisons determined by one-

tailed Wilcoxon rank sum test were denoted by the asterisks: * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$,

**** $p \leq 0.0001$, and "ns" as not significant.

SUPPORTING INFORMATION

Supplementary Fig 1.

(a)

(b)

Supplementary Figure 1. Distribution of the relative abundance of viral transcripts within individual cells over the course of the infection. **(a)** The overall relative abundance of viral transcripts was calculated as the percentage of viral reads in the pool of all the reads for each cell. The dots are colored by the time points. All pairwise comparisons were performed with one-tailed Wilcoxon rank sum test, for which the null hypothesis was that cells harvested at an earlier time point have a lower median relative abundance of viral transcripts than those harvested at a later time point. The significance levels were denoted by the asterisks: * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$, and "ns" as not significant. The data for HBEpC cells at 12hpi was collected from a repeated infection assay, due to the initial failure of single-cell library preparation

875   for the corresponding sample. **(b)** Distribution of the relative abundance of viral transcripts derived

876   from each segment within individual cells over the course of the infection. All box plots show the

877   first and third quantiles as the lower and upper hinges, the median in the center, and 1.5 * inter-

878   quartile range (IQR) from the first and third quantiles as the whiskers.

Supplementary Fig 2.

Supplementary Figure 2. Distribution of DVG/FL ratios for each viral segment in two cell types over the course of the infection. DVG transcripts represented by gap-spanning reads were grouped by their junction coordinates and filtered by their frequency in each sample (i.e., >= 10 times)

Supplementary Fig 3.

Supplementary Figure 3. Distribution of junction sites in the PB2 segment and the medians of their DVG/FL ratio in cells where a given defective viral transcript was detected. **(a)** The defective PB2 segments detected in the PR8 virus stock used for infection. **(b-c)**

886   The DVG PB2 transcripts detected in **(b)** A549 and **(c)** HBEpC cells over the course of the infection. Transcripts carrying the same

887   junction sites as seen in the virus stock were denoted by asterisks.

Supplementary Fig 4.

Supplementary Figure 4. Distribution of junction sites in the PB1 segment and the medians of their DVG/FL ratio in cells where a given defective viral transcript was detected.

Supplementary Fig 5.

893　**Supplementary Figure 5.** Distribution of junction sites in the PA segment and the medians of their DVG/FL ratio in cells where a given

894　defective viral transcript was detected.

Supplementary Fig 6.

(a)

Supplementary Figure 6. Distribution of the DVG/FL ratio in A549 and HBEPC cells over the course of the infection. All pairwise comparisons were performed with two-tailed Wilcoxon rank sum test, for which the null hypothesis was that cells harvested at two different time points have different median relative abundance of viral transcripts. The significance levels were denoted by

900   the asterisks: * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$, and "ns" as not significant. All

901   box plots show the first and third quantiles as the lower and upper hinges, the median in the

902   center, and 1.5 * inter-quartile range (IQR) from the first and third quantiles as the whiskers.

Supplementary Fig 7.

Supplementary Figure 7. Percentage of A549 and HBEpC cells in which a given DVG PB2 transcript was detected versus the median of the DVG/FL ratio for that transcript in those cells over the course of the infection. Each type of DVG PB2 transcript was represented

906   by a dot or triangle. A dot denoted the DVG transcripts also seen in the virus stock and a filled triangle denoted the *de novo* generated

907   transcripts. Two dominant DVG PB2 species were highlighted in slate blue and orange.

910 **Supplementary Figure 8.** Percentage of A549 and HBEpC cells in which a given DVG PB1 transcript was detected versus the median

911 of the DVG/FL ratio for that transcript in those cells over the course of the infection. Two dominant DVG PB1 species were highlighted

912 in yellow and pink.

Supplementary Fig 9.

Supplementary Figure 9. Percentage of MDCK and A549 cells in which a given DVG PB2 transcript was detected versus the median

of the DVG/FL ratio for that transcript in those cells. MDCK and A549 cells were infected with the same PR8 virus stock at high (5) or

916   low (0.2) MOI and harvested at 6hpi followed by 10X Genomics 3' single-cell library preparation and sequencing. Two dominant DVG

917   PB2 species were highlighted in slate blue and orange.

Supplementary Fig 10.

920　**Supplementary Figure 10.** Percentage of MDCK and A549 cells in which a given DVG PB1 transcript was detected versus the median

921　of the DVG/FL ratio for that transcript in those cells. MDCK and A549 cells were infected with the same PR8 virus stock at high (5) or

922   low (0.2) MOI and harvested at 6hpi followed by 10X Genomics 3' single-cell library preparation and sequencing. Two dominant DVG

923   PB1 species were highlighted in yellow and pink.

Supplementary Fig 11.

Supplementary Figure 11. Percentage of A549 and HBEpC cells in which a given DVG PA transcript was detected versus the median

of the DVG/FL ratio for that transcript in those cells over the course of the infection. The dominant DVG PA species in each cell type

were highlighted.

Supplementary Figure 12. Percentage of MDCK and A549 cells in which a given DVG PA transcript was detected versus the median

of the DVG/FL ratio for that transcript in those cells. MDCK and A549 cells were infected with the same PR8 virus stock at high (5) or

low (0.2) MOI and harvested at 6hpi followed by 10X Genomics 3' single-cell library preparation and sequencing. The DVG PA

transcripts were not detected in MDCK cells infected at low MOI. The dominant DVG PA species was highlighted in chartreuse.

Supplementary Fig 13.

Supplementary Figure 13. Visualization of un-supervised cell clustering on a t-SNE plot for

mock-infected **(a)** A549 and **(b)** HBEpC cells and the distribution of cell-cycle within the population

937   at each time point in two cell types. (**a-b**) Distribution of the cell-cycle stage for each cell

938   represented by a dot on the t-SNE plot. Dots were colored by either the cluster identities for the

939   mock-infected cells, or the cell-cycle stages for mock- and PR8-infected cells. The cell-cycle stage

940   was assigned to each cell based on the expression level of a list of cell-cycle markers. (**c**)

941   Distribution of the percentage of cells in each cell-cycle stage in two cell types.

Supplementary Fig. 14

Supplementary Figure 14. Enriched biological process gene ontology (GO) terms in A549 at

12hpi. The top 10 over-represented GO terms in each cluster are denoted by dots. The color

intensity of each dot indicates the fold enrichment of the corresponding GO term and the size

corresponds to the ratio of queried genes in the gene set associated with a given GO term. The

GO terms enriched in each cluster are color-coded by clusters. The bar chart above the bubble

chart of the GO terms shows the median relative abundance of the viral transcripts across cells

in each cluster. The colors of the bar are consistent with that of the GO terms.

Supplementary Fig 15.

Supplementary Figure 15. Functional MEGENA modules in clusters 0 and 1 of A549 at 24hpi.

Each panel shows enriched biological process GO terms in the modules correlated with the

relative abundance of virus transcripts in corresponding clusters. Each GO term is denoted by a

dot. The color intensity of each dot indicates the fold enrichment of the corresponding GO term

and the size corresponds to the -log10 transformed corrected p-value for a given GO term. Red

and blue color-bars above the dot plot denote positive or negative correlation of viral transcription

with corresponding modules, respectively. Modules that have a significant correlation with the

relative abundance of virus transcripts and enriched GO biological process terms are shown in

the GO term plots. Module member information for all viral transcription correlated-modules can

be   found   on   Github   (https://github.com/GhedinLab/Single-Cell-IAV-infection-in-

monolayer/tree/master/SupplementaryTables).

Supplementary Fig 16.

965 **Supplementary Figure 16.** Functional MEGENA modules in clusters 0, 1, and 3 of HBEpC at

966 24hpi. Each panel shows enriched biological process GO terms in the modules correlated with

967 the relative abundance of virus transcripts in corresponding clusters. Each GO term is denoted

968 by a dot. The color intensity of each dot indicates the fold enrichment of the corresponding GO

969 term and the size corresponds to the -log10 transformed corrected p-value for a given GO term.

970 Red and blue color-bars above the dot plot denote positive or negative correlation of viral

971 transcription with corresponding modules, respectively. Modules that have a significant

972 correlation with the relative abundance of virus transcripts and enriched GO biological process

973 terms are shown in the GO term plots. Module member information for all of viral transcription

974 correlated-modules can be found on Github (https://github.com/GhedinLab/Single-Cell-IAV-

975 infection-in-monolayer/tree/master/SupplementaryTables).

Supplementary Fig 17.

Supplementary Figure 17. Distribution of the DVG/FL ratio for the DVG PA transcripts in each

cluster of A549 cells at 12hpi and 24hpi and HBEpC cells at 24hpi. All box plots show the first and

third quantiles as the lower and upper hinges, the median in the center, and 1.5 * inter-quartile

981   range (IQR) from the first and third quantiles as the whiskers. The significance levels of pairwise

982   comparisons determined by one-tailed Wilcoxon rank sum test were denoted by the asterisks: *

983     $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$, and "ns" as not significant.

Supplementary Fig 18.

Supplementary Figure 18. Expression of ISGs and IFNs in DI-enriched A549 cells at 24hpi. **(a-**

b) The expression level of **(a)** significantly over-expressed genes in cluster 2, which are

associated with GO terms related to the type I IFN signaling pathway and have the fold change

on the log-scale >= 1, and **(b)** type I and III IFNs, in each cluster of A549 cells at 24hpi. The dot

represents the normalized expression level of a gene in each cell. The violin shade is colored by

the cluster identity. **(c)** Validation of over-expression of selected ISGs, IFNs, and the M gene of

IAV (i.e., FLU) by qPCR. Subconfluent A549 cells were either mock-infected or infected with one

type of PR8-DI virus (PR8-DI162, -DI222, or -DI244) or PR8-WT virus at a MOI of 10. Total RNA

was extracted from cells collected at 24hpi. The levels of ISGs, IFNs, and the IAV M gene mRNA

were determined by qPCR and normalized to the level of β-actin (ACTB). The error bar

representing the standard deviation of the mean was obtained from three biological replicates.

Statistical tests were done using two-tailed Student's t-test in R to compare the differences in fold

change between PR8-WT and PR8-DI infection. The significance levels were denoted by the

asterisks: * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$, and "ns" as not significant.

Supplementary Fig 19.

Supplementary Figure 19. Distribution of the size of the deletions in the gap-spanning reads aligned to the polymerase segments in

two cell types over the course of the infection. The frequency of each size detected in a sample was represented by the height of the

bar shown on the y-axis.