The Spatio-Temporal Control of Zygotic Genome Activation

George E. Gentsch,[1,*] Nick D. L. Owens,[2] and James C. Smith[1,3,*]

[1]Developmental Biology Laboratory, Francis Crick Institute, London, NW1 1AT, UK
[2]Department of Stem Cell and Developmental Biology, Pasteur Institute, 75015 Paris, France
[3]Lead Contact
*Correspondence: george.gentsch@crick.ac.uk (G.E.G.) and jim.smith@crick.ac.uk (J.C.S.)

SUMMARY

One of the earliest and most significant events in embryonic development is zygotic genome activation (ZGA). In several species, bulk transcription begins at the mid-blastula transition (MBT) when, after a certain number of cleavages, the embryo attains a particular nuclear-to-cytoplasmic (N/C) ratio, maternal repressors become sufficiently diluted, and the cell cycle slows down. Here we resolve the frog ZGA in time and space by profiling RNA polymerase II (RNAPII) engagement and its transcriptional readout. We detect a gradual increase in both the quantity and the length of RNAPII elongation before the MBT, revealing that >1,000 zygotic genes disregard the N/C timer for their activation, and that the sizes of newly transcribed genes are not necessarily constrained by cell cycle duration. We also find that Wnt, Nodal and BMP signaling together generate most of the spatio-temporal dynamics of regional ZGA, directing the formation of orthogonal body axes and proportionate germ layers.

INTRODUCTION

The genomes of multicellular organisms are transcriptionally silent at the time of fertilisation, and the events of early development, including zygotic (also known as embryonic) genome activation (ZGA), are directed by maternal gene products (De Iaco et al., 2019; Eckersley-Maslin et al., 2019; Gentsch et al., 2018b; Lee et al., 2013; Leichsenring et al., 2013; Liang et al., 2008). The number of cell cycles after which ZGA becomes essential for development (at which embryos arrest if transcription is inhibited) is highly reproducible within each species. In the zebrafish, the frog *Xenopus* and the fruit fly *Drosophila*, this occurs after 10, 12 and 13 cell cycles, respectively, at the so-called mid-blastula transition (MBT) (Blythe and Wieschaus, 2015; Gentsch et al., 2018b; Kane and Kimmel, 1993; Newport and Kirschner, 1982a). Early development in these species occurs with no gain in cytoplasmic volume, and studies in *Xenopus* suggest that ZGA is triggered at a particular nuclear-to-cytoplasmic (N/C) ratio, when the increasing amount of nuclear DNA titrates out maternally deposited repressors (Newport

and Kirschner, 1982b). Slower-developing mammalian embryos show major waves of RNA polymerase II (RNAPII)-mediated transcription as early as the 2-cell stage in mice (Bolton et al., 1984; Hamatani et al., 2004) and 4- to 8-cell stage in humans (Braude et al., 1988; Vanessa et al., 2011). This occurs days before the formation of the blastocyst, which, like the blastula, contains the pluripotent cells that form the embryo proper.

In *Xenopus*, ZGA is associated with changes in cell behaviour after the MBT. First, rapid and nearly synchronous cell cleavages give way to longer and asynchronous cell divisions (Anderson et al., 2017; Newport and Kirschner, 1982a). Second, embryonic cells acquire the ability to respond to inductive signaling (Gentsch et al., 2018b), causing them to become motile, to establish dorso-ventral patterning, and to contribute to one or two of the three germ layers (endoderm, mesoderm and ectoderm). These germ layers emerge first during gastrulation and are the primordia of all organs. Third, embryos show accelerated degradation of maternal RNA, and fourth, cells gain apoptotic (Stack and Newport, 1997) and immunogenic (Gentsch et al., 2018a) capabilities.

While large-scale ZGA occurs at the MBT, some genes escape the repressive environment of the early embryo, and nascent transcripts can be detected in *Xenopus* during rapid cleavage stages. For example, primary microRNA transcripts of the polycistronic MIR-427 gene (Lund et al., 2009) are detectable in *X. tropicalis* after just three cell divisions (Owens et al., 2016). MIR-427, like its zebrafish equivalent MIR-430, is activated at early stages by the synergistic and pioneering activities of maternal members of the SoxB1 and Pou5F (Oct4) transcription factor (TF) families (Gentsch et al., 2018b; Heyn et al., 2014; Lee et al., 2013). These core pluripotency TFs, represented by Sox3 and Pou5f3 in *Xenopus*, are characterized by ubiquitous and high translation frequencies in pre-MBT embryos (Gentsch et al., 2018b; Lee et al., 2013). Zygotic transcription of the Nodal-encoding genes *nodal3/5/6*, and of the homeobox genes *siamois1/2*, is initiated by nuclear β-catenin as early as the 32-cell stage (Owens et al., 2016; Skirkanich et al., 2011; Yang et al., 2002).

While miR-427 (and miR-430 in zebrafish) contributes to the clearance of maternal RNA (Giraldez et al., 2006; Lund et al., 2009), *nodal* and *siamois* gene initiate the formation of the germ layers and body axes (Agius et al., 2000; Lemaire et al., 1995). All these genes, and other early-activated genes in *Drosophila*, *Xenopus* and zebrafish, have coding sequences of <1 kb and either no introns or just a few (Heyn et al., 2014). It has been suggested that the early rapid cell cycles cause DNA replication machinery to interfere with the transcription of larger genes (Shermoen and O'Farrell, 1991), a suggestion supported, to date, by the profiling of nascent transcripts (Heyn et al., 2014). We note, however, that the detection and temporal resolution of de novo transcription can be particularly challenging for genes that have both maternal and zygotic transcripts.

Here we use the continuous occupancy of RNAPII along gene bodies as a method to record ZGA. In contrast to transcript profiling techniques, this method (1) directly determines the

activity of every gene; (2) is independent of metabolic labeling (Heyn et al., 2014) and of any

gene feature such as introns (Lee et al., 2013), single nucleotide polymorphisms (Harvey et al.,

2013; Lott et al., 2011) or transcript half-lives; and (3) circumvents difficulties in detecting

nascent transcripts in a large pool of maternal transcripts. By combining these data with the

profiling of the transcriptome along the primary body axes (Blitz et al., 2017), we resolve ZGA

in time and space for wild-type and various loss-of-function embryos. We provide evidence that

runs counter to our original understanding of the cell cycle or of the N/C ratio in constraining

gene expression before MBT. And finally, we show how signaling initiates and coordinates

spatio-temporal ZGA in the *Xenopus* embryo.

RESULTS

RNAPII Profiling Reveals Exponential ZGA before MBT

In an effort to resolve the progression of ZGA, we profiled chromatin for RNAPII engagement

on hand-sorted *X. tropicalis* embryos over six developmental stages from the 32-cell to the late

gastrula stage (Figure 1A,B). RNAPII was localized on the genome by chromatin

immunoprecipitation followed by deep sequencing (ChIP-Seq). We complemented RNAPII

profiling with high time-resolution transcriptomics (Owens et al., 2016) counting both exonic

and intronic RNA at 30-min intervals from fertilization to the late gastrula stage (Figures 1A,B

and S1A,B). For both maternal and zygotic genes, the detection threshold was set to ≥3

transcripts per million (TPM) averaged over any 1-h window during this developmental period

to avoid genes with general low-level expression. This restricted the analysis to 13,042 genes

(Figure 1B). These genes were considered active when we detected simultaneously RNAPII

enrichment along their full length (see Transparent Methods) as well as the presence of the

corresponding transcripts. In doing so, we used a low threshold of ≥0.1 TPM so as not to miss

the onset of gene transcription. RNAPII-guided ZGA profiling was verified in part by active post-

translational histone marks (Hontelez et al., 2015) and by differential expression methods

aiming at detecting nascent transcripts. Thus, zygotic transcript depletion (by blocking RNAPII

elongation with α-amanitin) (Gentsch et al., 2018b) or enrichment (by selecting 4-thiouridine

[4sU] tagged transcripts at the MBT and the mid-gastrula stage) showed substantial overlaps

and positive correlations with RNAPII-covered genes (Figures 1A,B and S1C,D and Tables S1

and S2).

This analysis revealed an exponential ZGA before the MBT with 27, 144 and 1,044 active genes

after 5 (32-cell, ~2.5 hpf), 7 (128-cell, ~3 hpf) and 10 (1,024-cell, ~4 hpf) cell cycles,

respectively. Gene activation reached its peak at the MBT (~4.5 hpf), with 1,854 newly-

activated genes, before dropping to 724 genes at the early-to-mid gastrula stage (~7.5 hpf) and

increasing again to 1,214 genes towards the end of gastrulation (~10 hpf) (Figures 1B,C and

S1E and Table S2). The dramatic increase in transcriptional activity that occurs in the 1.5 hours

between the 128-cell stage and the MBT can be illustrated by Hilbert curves (Figure 1C), which

provide a genome-wide overview of RNAPII enrichment by folding chromosomes into two-
dimensional space-filling and position-preserving plots (Gu et al., 2016). While most zygotic
genes remain active beyond the mid-gastrula stage, 197 (including *siamois2* [*sia2*], *nodal5* and
znf470) of the 4,836 zygotic genes (~4%) are deactivated within ~6 h of development (Figures
1C and S1F,G). Slightly less than one third of the activated genes were differentially expressed
along either or both of the animal-vegetal and dorso-ventral axes (Figures 1D and S1F).

The temporal order of enriched biological processes supported by ZGA matched the regulatory
flow of gene expression, starting with nucleosome assembly, nucleic acid synthesis, mRNA
metabolism and production, post-translational modification and degradation of proteins (Figure
1E). The earliest transcriptional engagement, beginning at the 32 to 128-cell stages, was
detected in gene clusters of tens to hundreds of kilobases (Figures 1A,C and S1H-J). These
clusters featured close relatives of the same genes, some of which are critical to Nodal signaling
(Nodal ligands), the formation of the Spemann organiser (Siamois homeobox transcription
factors), nucleosome assembly (histones), mRNA decay (MIR-427), and ongoing gene
regulation (zinc finger [ZF] transcription factors with on average 10 Cys2-His2 [C2H2] domains;
Figure S1J). These earliest activated genes were shorter and encoded smaller proteins than
those within the maternal pool or those that are activated post-MBT (Figures 1F and S1K). The
non-coding features that contributed most to the differences in length were the 3' UTRs and
introns (Figure S1K).

We noted that the shorter zygotic genes observed before the MBT did not correlate strictly with
the time constraints imposed by short cell cycles. We detected increasing and wider spread of
de novo recruitment of RNAPII before the MBT, when cleavages continue to occur at rapid and
near-constant pace (Figures 1F and S1K). During this period, the median length of activated
genes (and their coding sequences) increases from ~0.9 kb (~0.4 kb) to ~5.9 kb (~0.9 kb).
However, it was not until after the MBT that the overall architecture of zygotic and maternal
genes became indistinguishable (Figures 1F and S1K). Temporal comparison of RNAPII
engagement and total RNA profiling suggested that the zygotic contribution to the transcriptome
(as calculated by the number of zygotic genes divided by the number of genes with ≥0.1 TPM
maternal transcripts averaged over the first hour post-fertilisation when the entire zygotic
genome is still transcriptionally inactive) rose within seven cell cycles from ~0.2% at the 32-cell
stage to ~22% at the MBT (Figure 1G). Further maternal degradation and more moderate
transcriptional engagement extended the zygotic contribution to about one third of the
transcriptome by the late gastrula stage. Maternal transcripts (≥0.1 TPM, see above) were
detected for ~67% of newly activated genes (18 out of 27 genes) at the 32-cell stage, ~85%
(99/117) at the 128-cell stage, ~87% (780/900) at the 1,024-cell stage, ~95% (1,754/1,854) at
the MBT, ~89% (644/724) at the mid-gastrula stage and ~90% (1,094/1,214) at the late gastrula
stage (Table S2). Altogether ~91% (4,389/4,836) of newly activated genes have ≥0.1 TPM
maternal contribution.

Wnt, Nodal and BMP Signals Are Key Drivers of Regional ZGA

We next sought to investigate the single and combined effects of different inductive signals on the spatio-temporal dynamics of ZGA. The early vertebrate embryo employs canonical Wnt, Nodal and BMP signals and their key transcriptional effectors β-catenin, Smad2 and Smad1, respectively, to establish the primary body axes and the three germ layers (reviewed by Arnold and Robertson, 2009 and Kimelman, 2006). In *Xenopus*, β-catenin first translocates to the nuclei of dorsal blastomeres at the 32-cell stage (Larabell et al., 1997; Schneider et al., 1996) (Figure 2A). After the MBT, zygotic Wnt8a causes more nuclear β-catenin to accumulate around the forming blastopore lip (Christian and Moon, 1993; Schohl and Fagotto, 2002). The nuclear translocation of Smad1 and Smad2 is triggered around the MBT by various BMP and Nodal ligands. Nuclear Smad1 is primarily detected on the ventral side and the blastocoel roof of the embryo while nuclear Smad2 is detected within the vegetal hemisphere (VH) and the marginal zone (MZ) (Faure et al., 2000; Schohl and Fagotto, 2002) (Figure 2A).

In an effort to inhibit canonical Wnt signaling, we injected into the *X. tropicalis* zygote a previously validated antisense morpholino oligonucleotide (MO) which interferes with β-catenin protein synthesis by annealing to the translation start codon (Heasman et al., 2000). Nodal and BMP signals were selectively blocked by incubating dejellied embryos from the 8-cell stage in the cell-permeable inhibitors SB431542 (Ho et al., 2006; Inman et al., 2002) and LDN193189 (Cuny et al., 2008; Young et al., 2017), respectively. The morphological phenotypes of these single loss-of-function (LOF) treatments were consistent with previous observations and ranged from impaired axial elongation causing the loss of tail structures (BMP, Reversade et al., 2005) to severe gastrulation defects (Wnt, Heasman et al., 2000, and Nodal, Ho et al., 2006) as shown in Figure 2B. Briefly, Nodal LOF impaired blastopore lip formation and bulk tissue movements of gastrulation (bullet points in Figure 2B). However, it did not preclude subsequent elongation of the antero-posterior axis. By contrast, Wnt LOF embryos underwent gastrulation (albeit delayed and more circumferentially rather than in a dorsal to ventral wave), but failed to form an antero-posterior axis, with both head and tail being absent. With respect to the joint effects of Wnt, Nodal and BMP signaling, most dual and triple LOFs combined their individual morphological defects such that, for example, Wnt/Nodal LOF resulted in the complete loss of gastrulation and axial elongation. By contrast, Wnt/BMP LOF produced defects such as non-fusing neural folds (arrowheads in Figure 2B), structures that were either absent in Wnt LOF embryos or normal in BMP LOF embryos.

Changes to ZGA caused by the single or combined LOF of Wnt, Nodal and/or BMP were then determined at the late blastula stage on a transcriptome-wide scale using deep RNA sequencing (RNA-Seq). Analysis was limited to the 3,315 zygotic genes for which spatio-temporal expression data is available (Blitz et al., 2017; Owens et al., 2016) and where reduced expression (≥50% loss of exonic and/or intronic transcript counts, FDR ≤10%) could be detected in α-amanitin-injected embryos (Figure 2C and Table S3) (Gentsch et al., 2018b). α-Amanitin-mediated inhibition of RNAPII elongation impedes the morphogenetic tissue

movements of gastrulation and ultimately leads to early embryonic death (Gentsch et al., 2018b). Spatial gene expression patterns were inferred from experiments comparing the transcriptomes of embryos dissected along their animal-vegetal and dorso-ventral axes (Blitz et al., 2017); we did not include the left-right axis because there were no significant differences in gene expression across this axis at the gastrula stage (Blitz et al., 2017). The signal-mediated transcriptional effects (1.5-fold change from control RNA level) on zygotic genes, 86% (2840/3315) of which have ≥0.1 TPM maternal contribution, ranged from ~1.5% (~1.3% down and ~0.2% up) to ~26% (~19% down and 7% up) for single BMP LOF and triple Wnt/Nodal/BMP LOFs, respectively (Figure 2C). As expected, the transcript levels of genes whose expression is solely zygotic were more strongly affected than those of zygotic genes with maternally contributed transcripts (Figures 2C and 4A). The extent of ZGA mis-regulation largely reflected the severity of the resulting morphological phenotypes at the late gastrula and the mid-tailbud stage (Figure 2B,C).

In comparison, the LOFs of critical maternal TFs like Pou5f3/Sox3 or VegT (Gentsch et al., 2018b) caused the mis-regulation of 61% (~24% down and ~37% up) and 13% (~6% down and ~7% up) of zygotic genes, respectively. The LOFs of four zygotic T-box TFs (zVegT, Eomes, Tbxt and Tbxt.2), all of which require Nodal signaling for their expression, caused slight mis-regulation in 19% (~9% down and ~10% up) of the zygotic genes as detected over three consecutive developmental time points during gastrulation (Table S3). Among the ZGA-enriched biological functions (Figure 1D), Wnt, Nodal and BMP signals, like maternal Pou5f3/Sox3 and VegT, strongly affected zygotic genes associated with cell migration, gastrulation, dorso-ventral and antero-posterior body axis formation and regionalization (Figure 2D). Impaired tissue movements during gastrulation, as observed in various LOFs (Figure 2B and Movie S1), was prefigured by a strong enrichment for cell migration-associated genes. The genes suppressed or unaffected by the selected signals and maternal TFs were enriched for the ZGA-critical biological processes of mRNA metabolism and translation. For instance, the transient activation of the entire zinc finger cluster (Figure S2A) was not affected by any tested LOF. Because family members are frequently cross-regulated, and the MBT-staged chromatin contains many Krüppel-like zinc finger 'footprints' (Gentsch et al., 2018b), it is conceivable that the unaffected, tissue-nonspecific part of ZGA is regulated by maternal zinc finger TFs. This vertebrate gene regulatory branch may be more ancient than that of Pou5F/SoxB1 as zinc finger TFs like Zelda are also key to ZGA of the invertebrate *Drosophila* (Liang et al., 2008).

Next, signal-dependent ZGA was resolved in time and space based on (i) the profiling of RNAPII-engaged chromatin from the 32-cell stage to the MBT and (ii) known gene expression patterns along the animal-vegetal and dorso-ventral axes (Blitz et al., 2017) (Figures 2E and S2B-F). In line with the nuclear translocation of their signal mediators (Figure 2A), Wnt, Nodal and BMP proved to be required for gene activation in different spatio-temporal domains of the early embryo: β-catenin was needed for ~87% and ~46% of genes preferentially expressed on the dorsal side and in the VH/MZ, respectively. Some of its target genes like *nodal3.1* and *sia2*

were already active by the 32-cell stage (Figures 1C and 2E). Upon Wnt LOF, the early
transcriptional down-regulation was followed by the mis-regulation of opposing cell fate
specifiers: the upregulation of ventral genes (e.g., *id2*, *szl*) and the downregulation of dorsal
genes (e.g., *chrd*, *otx2*). These observations suggest that β-catenin protects dorsal cells from
ventralization (Figures 2E and S2C).

Along similar lines, Nodal LOF embryos predominantly displayed a down-regulation of dorsal
(~63%) and VH/MZ-specific (~73%) genes, although there was no effect of Nodal LOF on the
earliest-activated genes at the 32-cell stage or on opposing cell fate regulators (Figure S2A,B).
Among the first genes to be sensitive to Nodal LOF was the MZ-specific FGF ligand *fgf20*,
activated by the 128-cell stage (Figures 1C and S2A,B). By contrast, BMP LOF caused a
decrease in ventrally-expressed gene expression (~45%) from the 1,024-cell stage onwards
(Figure S2A,C).

As a comparison, the ubiquitous expression of maternal Pou5f3/Sox3 was required for
transcription in all spatio-temporal domains, including, for example, the uniform expression of
miR-427 (Figure S2A,D). The requirement for Pou5f3/Sox3 was more marked, however, for
region-specific genes, in particular those expressed within animal- (~55%) and ventral-specific
(~67%) domains (Figure S2A,D). The maternal TF VegT promoted vegetal identity by activating
~40% of genes transcriptionally enriched within its own expression domain, the vegetal
hemisphere, while suppressing genes that are expressed in the animal hemisphere like *foxi4.2*.
The requirement for VegT was similar in ventral- and dorsal-specific genes (~31% and ~30%
respectively).

Wnt/BMP Synergy Enables Uniform ZGA Across the Dorso-Ventral Axis
The relationships between inductive signals with respect to spatial aspects of ZGA was
explored by comparing zygotic transcriptomes in single and double LOF experiments.
Interestingly, while simultaneous loss of both Nodal and BMP function, or both Nodal and Wnt,
caused additive effects on gene expression compared to the single LOFs (Figures 3E and S3A-
N), simultaneous loss of both BMP and Wnt signaling showed a more synergistic effect (Figure
3A,F). These observations are consistent with the morphological consequences of single and
double LOFs (Figure 2B). Single LOF experiments revealed very little overlap between Wnt
and BMP gene targets (Figure 3B), a result consistent with their domains of activity which are,
initially, at opposite ends of the dorso-ventral axis. However, dual Wnt/BMP inhibition increased
the number of downregulated genes by 292, a rise of ~118% and ~664% with respect to
individual Wnt- and BMP-dependent genes respectively (Figure 3B). Interestingly, this synergy
affected 166 Nodal-dependent genes, most of which had uniform expression levels across the
dorso-ventral axis (Figure 3C,D,F-H). Thus, spatially-restricted Wnt, BMP and Nodal signals
act together to establish dorso-ventral expression uniformity of genes such as *tbxt* and *eomes*
(Figure 3I).
261

Overall, the loss of canonical Wnt, Nodal and/or BMP signaling caused the misregulation of
~39% (~22.1% down, ~2.1% down/up and ~14.4% up) of genes activated at ZGA (Figure 4A).
These signals were required for most regional ZGA on the dorsal side (~89%) and in the VH
and MZ (~82%). Notably, their input affected virtually all genes (~98%, 56/57) with enriched
expression in the dorso-vegetal/MZ quadrant (Figure 4B,C). Thus, Wnt, Nodal and BMP
substantially contribute to regional ZGA in most anatomical domains of the early gastrula
embryo with the exception of animally enriched transcription (~19%). Animal- and ventral-
specific gene expression relies strongly on both activation by ubiquitous maternal TFs (e.g.
Pou5f3/Sox3) and on repression by signals (Figure 4B,D) and other maternal TFs (e.g. VegT)
on the opposite side (Figure S2F).

DISCUSSION

Our study provides two major insights into the mechanisms by which ZGA is initiated in time
and space in *Xenopus tropicalis*. The first concerns the temporal aspects, where we find that
RNAPII can be detected across gene bodies well before the MBT, during the period when rapid
synchronous cell divisions divide the zygote into 4,096 blastomeres (Figure 4E). The average
length of genes covered by RNAPII grows during this time, from ~1 kb at the 32-cell stage to
~6 kb at the 1,024-cell stage; these figures compare with an average size of ~16 kb for
maternally-expressed genes and for genes expressed after the MBT. Recent long-read
sequencing of the zebrafish transcriptome at pre-MBT stages identified transcripts as long as
8 kb spanning multiple pri-miR-430 elements (Nudelman et al., 2018). Furthermore, RNAPII
elongation in pre-MBT *Drosophila* embryos occurred at rates of 2.4 to 3.0 kb/min (Chen et al.,
2013; Fukaya et al., 2017).

We do not know why RNAPII, despite its high abundance and its ability to promote rapid
elongation, is restricted at early stages from transcribing more genes and longer genes. It may,
perhaps, be a consequence of the gradual nature of the chromatin remodeling that occurs
during these stages, from the accessibility of *cis*-regulatory elements (Gentsch et al., 2018b;
Liu et al., 2018; Lu et al., 2016; Wu et al., 2016) to the spatial organisation of an initially
unstructured or highly variable chromatin landscape (Du et al., 2017; Flyamer et al., 2017; Hug
et al., 2017; Kaaij et al., 2018; Ke et al., 2017). Whatever the reason, the increase of elongated
RNAPII engagement between the 128-cell stage and the MBT indicates that a significant
component of ZGA disregards the N/C ratio which was originally thought to underlie the onset
of transcription at the MBT (Newport and Kirschner, 1982b). Similar conclusions have been
drawn from profiling the zygotic transcriptome of haploid *Drosophila* (Lu et al., 2009) and cell
cycle-arrested zebrafish (Chan et al., 2018). Thus, it is becoming clear from work in flies, fish
and frogs that ZGA starts before the MBT and accelerates thereafter (Ali-Murthy et al., 2013;
Collart et al., 2014; Mathavan et al., 2005; Owens et al., 2017; Tan et al., 2013), reaching a
peak at the MBT (reviewed by Jukam et al., 2017 and Langley et al., 2014).

These observations notwithstanding, it remains possible that cell cycles do contribute to the
temporal progression of ZGA and the exponential increase in the number of activated genes
before the MBT. In particular, cell cycles may accelerate chromatin remodeling by displacing
suppressors in mitotic chromatin and providing unique access to TFs (Halley-Stott et al., 2014)
and structural proteins of high-order chromatin (Ke et al., 2017). For example, maternal core
histones have been shown to prevent premature ZGA by competing with specific TFs (Joseph
et al., 2017).

In addition to the small sizes of the earliest activated genes, we observed that most of these
genes, which have no or few introns, code for groups of related factors like histones or zinc
finger TFs, and that they appear as clusters spanning up to several hundred kilobases. This is
in line with previous findings of the earliest active multicopy and intron-poor genes like miR-427
and *nodal5/6* in *Xenopus* embryos (Collart et al., 2014; Lund et al., 2009; Owens et al., 2016;
Skirkanich et al., 2011; Takahashi et al., 2006; Yanai et al., 2011; Yang et al. 2002) and miR-
430 in zebrafish and Medaka fish (Giraldez et al., 2005; Heyn et al., 2014; Tani et al., 2010).
The number and spatial proximity of clustered genes enhances transcriptional output by
allowing the sharing of multiple cis-regulatory elements (arranged as super-enhancers) (Whyte
et al., 2013) and by fortifying transcriptional condensates of TFs, coactivators and RNAPII
(Boija et al., 2018; Cho et al., 2018; Chong et al., 2018; Sabari et al., 2018; Shrinivas et al.,
2018). Overall, based on enriched gene functions, we discovered that ZGA exerts a temporal
control of gene expression from nucleosome remodeling before the MBT to protein degradation
after the MBT.

Our second insight concerns spatial ZGA, and the observation that we can assign a large
proportion of spatio-temporal ZGA to key signaling pathways (reviewed by Arnold and
Robertson, 2009 and Kimelman, 2006). Canonical Wnt, Nodal and BMP signaling govern
regional ZGA in line with the nuclear translocation of their signal mediators (Faure et al., 2000;
Larabell et al., 1997; Schohl and Fagotto, 2002). Thus, Nodal signaling predominantly affects
transcription within the vegetal hemisphere and marginal zone, while Wnt and BMP initiate
transcription in dorsal and ventral regions, respectively. The timing of regional ZGA is defined
by the sequential translocation of signal mediators such that nuclear β-catenin directs regional
ZGA at the 32-cell stage, followed by nuclear Smad2 at the 128-cell stage and Smad1 at the
1,024-cell stage. While Smad2-mediated signal transduction depends on the zygotic
transcription of its six Nodal ligands (Faure et al., 2000; Gentsch et al., 2018b; Jones et al.,
1995; Yang et al., 2002), canonical Wnt and BMP signaling are initiated by the maternally
inherited ligands Wnt11 and BMP2/4/7, respectively (Faure et al., 2000; Heasman, 2006; Tao
et al., 2005).

We also show that the synergy of opposing signals of the Wnt and BMP pathway affects many
Nodal-dependent genes with uniform expression along the dorso-ventral axis such as *eomes*
and *Brachyury* (*tbxt*). It is not yet clear whether Wnt/BMP synergy arises from joint chromatin

engagement or from mutual or post-translational interactions. For instance, Wnt8a signal can
enhance BMP transcriptional readouts by inhibiting the phosphorylation of GSK3, which
normally targets Smad1 for degradation (Fuentealba et al., 2007). However, the analysis of
Brachyury gene regulation in zebrafish suggests that Wnt and BMP can be integrated at a
single cis-regulatory DNA element and together with a separate Nodal-responsive DNA
element they can establish uniform dorso-ventral expression (Harvey et al., 2010). This is
further corroborated by our analysis of genome-wide chromatin engagement (Gentsch et al.,
2018b): the canonical DNA recognition motif for the Wnt-associated basic helix-span-helix
(bHSH) TF AP-2 was more enriched at Smad1 than Smad2 binding sites (Figure S3O).

We therefore propose that Wnt, BMP and Nodal signal mediators are critical to regional ZGA
and that they balance initially opposing cell fate commitments. However, we have previously
shown that signal integration also relies on maternal pioneer TFs like Pou5f3 and Sox3 to make
signal-responsive *cis*-regulatory elements accessible for signal mediator binding. For example,
Nodal-induced transcription of the *Brachyury* gene depends on the pioneering roles of maternal
Pou5f3 and Sox3, and less on their transcriptional activities (Gentsch et al., 2018b).

Overall, we demonstrate that the temporal and spatial dynamics of regional ZGA are regulated
by the sequential and spatially restricted translocation of Wnt, Nodal and BMP signal mediators.
These events establish the formation of the primary body axes and germ layers of the embryo.
Temporal RNAPII profiling indicates that >1,000 genes of increasing length are activated before
MBT and that this substantial portion of ZGA is independent of both the classic N/C ratio and
of cell cycle lengthening.

LIMITATIONS OF THE STUDY

We detected a dramatic increase in genome-wide recruitment of RNAPII over the cleavage
stages during which the genome begins to be transcribed. We used within-sample
normalization to scale developmental stage-specific RNAPII profiles. However, because of the
large differences in total RNAPII enrichment between samples, chromatin spike-ins are now
considered a more accurate method to normalize ChIP-Seq profiles across consecutive
developmental stages (Chen et al., 2016). We combined separate whole-embryo
determinations of RNAPII engagement and transcript levels to reveal the temporal dynamics of
ZGA. This approach could be improved by profiling RNAPII-associated RNA to directly couple
RNAPII elongation with transcript accumulation (e.g., Churchman and Weissman, 2011). In
addition, the spatial resolution of ZGA, which is based on transcriptomics of dissected
embryonic parts in our study, could be enhanced by various deep single-cell profiling and super-
resolution imaging technologies. We show that most regional ZGA depends on Wnt, Nodal and
BMP signals, but an important question remains: How are these signals integrated at the
chromatin level to sustain RNAPII-mediated transcript elongation? In part, this could be

investigated by targeted genome editing to increase our understanding of signal-responsive

gene regulatory DNA.

ACKNOWLEDGEMENTS

We thank Abdul Sesay, Leena Bhaw, Harsha Jani, Deborah Jackson and Meena Anissi for

deep sequencing; Mareike Thompson for critical reading of the manuscript; and the Smith lab

for discussions and advice. G.E.G and J.C.S. were supported by the Medical Research Council

(program number U117597140) and are now supported by the Francis Crick Institute, which

receives its core funding from Cancer Research UK (FC001-157), the UK Medical Research

Council (FC001-157), and the Wellcome Trust (FC001-157).

AUTHOR CONTRIBUTIONS

Conceptualization, G.E.G.; Methodology, G.E.G.; Computational Code, G.E.G.; Formal

Analysis, G.E.G. and N.D.L.O.; Investigation, G.E.G.; Writing – Original Draft, G.E.G. and

J.C.S.; Writing – Review & Editing, G.E.G and J.C.S.; Funding Acquisition, J.C.S.

DECLARATION OF INTERESTS

The authors declare no competing interests.

REFERENCES

Agius, E., Oelgeschlager, M., Wessely, O., Kemp, C., and De Robertis, E.M. (2000).
Endodermal Nodal-related signals and mesoderm induction in Xenopus. Development *127*,
1173–1183.

Ali-Murthy, Z., Lott, S.E., Eisen, M.B., and Kornberg, T.B. (2013). An essential role for zygotic
expression in the pre-cellular Drosophila embryo. PLoS Genet. *9*, e1003428.

Anderson, G.A., Gelens, L., Baker, J.C., and Ferrell, J.E. (2017). Desynchronizing Embryonic
Cell Division Waves Reveals the Robustness of Xenopus laevis Development. Cell Rep. *21*,
37–46.

Arnold, S.J., and Robertson, E.J. (2009). Making a commitment: cell lineage allocation and
axis patterning in the early mouse embryo. Nat. Rev. Mol. Cell Biol. *10*, 91–103.

Boija, A., Klein, I.A., Sabari, B.R., Dall'Agnese, A., Coffey, E.L., Zamudio, A.V., Li, C.H.,
Shrinivas, K., Manteiga, J.C., Hannett, N.M., et al. (2018). Transcription Factors Activate
Genes through the Phase-Separation Capacity of Their Activation Domains. Cell *175*, 1842–
1855.e16.

Blitz, I.L., Paraiso, K.D., Patrushev, I., Chiu, W.T.Y., Cho, K.W.Y., and Gilchrist, M.J. (2017).
A catalog of Xenopus tropicalis transcription factors and their regional expression in the early
gastrula stage embryo. Dev. Biol. *426*, 409–417.

Blythe, S.A., and Wieschaus, E.F. (2015). Zygotic genome activation triggers the DNA
replication checkpoint at the midblastula transition. Cell *160*, 1169–1181.

Bolton, V.N., Oades, P.J., and Johnson, M.H. (1984). The relationship between cleavage,
DNA replication, and gene expression in the mouse 2-cell embryo. J. Embryol. Exp. Morph.
79, 139–163.

Braude, P., Bolton, V., and Moore, S. (1988). Human gene expression first occurs between
the four- and eight-cell stages of preimplantation development. Nature *332*, 459–461.

Chan, S.H., Tang, Y., Miao, L., Darwich-Codore, H., Vejnar, C.E., Beaudoin, J.-D., Musaev,
D., Fernandez, J.P., Moreno-Mateos, M.A., and Giraldez, A.J. (2018). Brd4 and P300
regulate zygotic genome activation through histone acetylation. bioRxiv 369231. DOI:
https://doi.org/10.1101/369231

Chen, K., Hu, Z., Xia, Z., Zhao, D., Li, W., and Tyler, J.K. (2015). The Overlooked Fact:
Fundamental Need for Spike-In Control for Virtually All Genome-Wide Analyses. Mol. Cell.
Biol. *36*, 662–667.

Chen, K., Johnston, J., Shao, W., Meier, S., Staber, C., and Zeitlinger, J. (2013). A global
change in RNA polymerase II pausing during the Drosophila midblastula transition. Elife *2*,
e00861.

Chiu, W.T., Le, R.C., Blitz, I.L., Fish, M.B., Li, Y., Biesinger, J., Xie, X., and Cho, K.W.Y.
(2014). Genome-wide view of TGFβ/Foxh1 regulation of the early mesendoderm program.
Development *141*, 1–114.

Cho, W.-K., Spille, J.-H., Hecht, M., Lee, C., Li, C., Grube, V., and Cisse, I.I. (2018). Mediator
and RNA polymerase II clusters associate in transcription-dependent condensates. Science
361, 412–415.

Chong, S., Dugast-Darzacq, C., Liu, Z., Dong, P., Dailey, G.M., Cattoglio, C., Heckert, A.,
Banala, S., Lavis, L., Darzacq, X., et al. (2018). Imaging dynamic and selective low-
complexity domain interactions that control gene transcription. Science *361*, eaar2555.

Christian, J.L., and Moon, R.T. (1993). Interactions between Xwnt-8 and Spemann organizer
signaling pathways generate dorsoventral pattern in the embryonic mesoderm of Xenopus.
Genes Dev. *7*, 13–28.

Collart, C., Owens, N.D.L., Bhaw-Rosun, L., Cooper, B., De Domenico, E., Patrushev, I.,
Sesay, A.K., Smith, J.N., Smith, J.C., and Gilchrist, M.J. (2014). High-resolution analysis of
gene activity during the Xenopus mid-blastula transition. Development *141*, 1927–1939.

Cuny, G.D., Yu, P.B., Laha, J.K., Xing, X., Liu, J.-F., Lai, C.S., Deng, D.Y., Sachidanandan,
C., Bloch, K.D., and Peterson, R.T. (2008). Structure-activity relationship study of bone
morphogenetic protein (BMP) signaling inhibitors. Bioorg. Med. Chem. Lett. *18*, 4388–4392.

De Iaco, A., Coudray, A., Duc, J., and Trono, D. (2019). DPPA2 and DPPA4 are necessary to
establish a 2C-like state in mouse embryonic stem cells. EMBO Reports e47382.

Du, Z., Zheng, H., Huang, B., Ma, R., Wu, J., Zhang, X., He, J., Xiang, Y., Wang, Q., Li, Y., et
al. (2017). Allelic reprogramming of 3D chromatin architecture during early mammalian
development. Nature *547*, 232–235.

Eckersley-Maslin, M., Alda-Catalinas, C., Blotenburg, M., Kreibich, E., Krueger, C., and Reik,
W. (2019). Dppa2 and Dppa4 directly regulate the Dux-driven zygotic transcriptional program.
Genes Dev. *33*, 194–208.

Faure, S., Lee, M.A., Keller, T., Dijke, ten, P., and Whitman, M. (2000). Endogenous patterns
of TGFbeta superfamily signaling during early Xenopus development. Development *127*,

2917–2931.

Flyamer, I.M., Gassler, J., Imakaev, M., Brandão, H.B., Ulianov, S.V., Abdennur, N., Razin,
S.V., Mirny, L.A., and Tachibana-Konwalski, K. (2017). Single-nucleus Hi-C reveals unique
chromatin reorganization at oocyte-to-zygote transition. Nature *544*, 110–114.

Fuentealba, L.C., Eivers, E., Ikeda, A., Hurtado, C., Kuroda, H., Pera, E.M., and De Robertis,
E.M. (2007). Integrating patterning signals: Wnt/GSK3 regulates the duration of the
BMP/Smad1 signal. Cell *131*, 980–993.

Fukaya, T., Lim, B., and Levine, M. (2017). Rapid Rates of Pol II Elongation in the Drosophila
Embryo. Curr. Biol. *27*, 1387–1391.

Gentsch, G.E., Spruce, T., Monteiro, R.S., Owens, N.D.L., Martin, S.R., and Smith, J.C.
(2018a). Innate Immune Response and Off-Target Mis-splicing Are Common Morpholino-
Induced Side Effects in Xenopus. Dev. Cell *44*, 597–610.e10.

Gentsch, G.E., Spruce, T., Owens, N.D.L., and Smith, J.C. (2018b). The role of maternal
pioneer factors in predefining first zygotic responses to inductive signals. bioRxiv 306803.
DOI: https://doi.org/10.1101/306803

Giraldez, A.J., Cinalli, R.M., Glasner, M.E., Enright, A.J., Thomson, J.M., Baskerville, S.,
Hammond, S.M., Bartel, D.P., and Schier, A.F. (2005). MicroRNAs regulate brain
morphogenesis in zebrafish. Science *308*, 833–838.

Giraldez, A.J., Mishima, Y., Rihel, J., Grocock, R.J., Van Dongen, S., Inoue, K., Enright, A.J.,
and Schier, A.F. (2006). Zebrafish MiR-430 promotes deadenylation and clearance of
maternal mRNAs. Science *312*, 75–79.

Gu, Z., Eils, R., and Schlesner, M. (2016). HilbertCurve: an R/Bioconductor package for high-
resolution visualization of genomic data. Bioinformatics *32*, 2372–2374.

Halley-Stott, R.P., Jullien, J., Pasque, V., and Gurdon, J. (2014). Mitosis gives a brief window
of opportunity for a change in gene transcription. PLoS Biol. *12*, e1001914.

Hamatani, T., Carter, M.G., Sharov, A.A., and Ko, M.S.H. (2004). Dynamics of global gene
expression changes during mouse preimplantation development. Dev. Cell *6*, 117–131.

Harvey, S.A., Sealy, I., Kettleborough, R., Fenyes, F., White, R., Stemple, D., and Smith, J.C.
(2013). Identification of the zebrafish maternal and paternal transcriptomes. Development
140, 2703–2710.

Harvey, S.A., Tümpel, S., Dubrulle, J., Schier, A.F., and Smith, J.C. (2010). no tail integrates
two modes of mesoderm induction. Development *137*, 1127–1135.

Heasman, J. (2006). Patterning the early Xenopus embryo. Development *133*, 1205–1217.

Heasman, J., Kofron, M., and Wylie, C. (2000). βCatenin Signaling Activity Dissected in the
Early Xenopus Embryo: A Novel Antisense Approach. Dev. Biol. *222*, 124–134.

Heyn, P., Kircher, M., Dahl, A., Kelso, J., Tomancak, P., Kalinka, A.T., and Neugebauer, K.M.
(2014). The earliest transcribed zygotic genes are short, newly evolved, and different across
species. Cell Rep. *6*, 285–292.

Ho, D.M., Chan, J., Bayliss, P., and Whitman, M. (2006). Inhibitor-resistant type I receptors
reveal specific requirements for TGF-beta signaling in vivo. Dev. Biol. *295*, 730–742.

Hontelez, S., van Kruijsbergen, I., Georgiou, G., van Heeringen, S.J., Bogdanović, O., Lister,
R., and Veenstra, G.J.C. (2015). Embryonic transcription is controlled by maternally defined

chromatin state. Nat. Commun. *6*, 10148.

Hug, C.B., Grimaldi, A.G., Kruse, K., and Vaquerizas, J.M. (2017). Chromatin Architecture
Emerges during Zygotic Genome Activation Independent of Transcription. Cell *169*, 216–
228.e219.

Inman, G.J., Nicolás, F.J., Callahan, J.F., Harling, J.D., Gaster, L.M., Reith, A.D., Laping,
N.J., and Hill, C.S. (2002). SB-431542 is a potent and specific inhibitor of transforming growth
factor-beta superfamily type I activin receptor-like kinase (ALK) receptors ALK4, ALK5, and
ALK7. Mol. Pharmacol. *62*, 65–74.

Jones, C.M., Kuehn, M.R., Hogan, B.L., Smith, J.C., and Wright, C.V. (1995). Nodal-related
signals induce axial mesoderm and dorsalize mesoderm during gastrulation. Development
121, 3651–3662.

Joseph, S.R., Pálfy, M., Hilbert, L., Kumar, M., Karschau, J., Zaburdaev, V., Shevchenko, A.,
and Vastenhouw, N.L. (2017). Competition between histone and transcription factor binding
regulates the onset of transcription in zebrafish embryos. Elife *6*, e23326.

Jukam, D., Shariati, S.A.M., and Skotheim, J.M. (2017). Zygotic Genome Activation in
Vertebrates. Dev. Cell *42*, 316–332.

Kaaij, L.J.T., van der Weide, R.H., Ketting, R.F., and de Wit, E. (2018). Systemic Loss and
Gain of Chromatin Architecture throughout Zebrafish Development. Cell Rep. *24*, 1–10.e4.

Kane, D.A., and Kimmel, C.B. (1993). The zebrafish midblastula transition. Development *119*,
447–456.

Ke, Y., Xu, Y., Chen, X., Feng, S., Liu, Z., Sun, Y., Yao, X., Li, F., Zhu, W., Gao, L., et al.
(2017). 3D Chromatin Structures of Mature Gametes and Structural Reprogramming during
Mammalian Embryogenesis. Cell *170*, 367–381.e20.

Kimelman, D. (2006). Mesoderm induction: from caps to chips. Nat. Rev. Genet. *7*, 360–372.

Langley, A.R., Smith, J.C., Stemple, D.L., and Harvey, S.A. (2014). New insights into the
maternal to zygotic transition. Development *141*, 3834–3841.

Larabell, C.A., Torres, M., Rowning, B.A., Yost, C., Miller, J.R., Wu, M., Kimelman, D., and
Moon, R.T. (1997). Establishment of the dorso-ventral axis in Xenopus embryos is presaged
by early asymmetries in beta-catenin that are modulated by the Wnt signaling pathway. J. Cell
Biol. *136*, 1123–1136.

Lemaire, P., Garrett, N., and Gurdon, J.B. (1995). Expression cloning of Siamois, a Xenopus
homeobox gene expressed in dorsal-vegetal cells of blastulae and able to induce a complete
secondary axis. Cell *81*, 85–94.

Lee, M.T., Bonneau, A.R., Takacs, C.M., Bazzini, A.A., Divito, K.R., Fleming, E.S., and
Giraldez, A.J. (2013). Nanog, Pou5f1 and SoxB1 activate zygotic gene expression during the
maternal-to-zygotic transition. Nature *503*, 360–364.

Leichsenring, M., Maes, J., Mössner, R., Driever, W., and Onichtchouk, D. (2013). Pou5f1
transcription factor controls zygotic gene activation in vertebrates. Science *341*, 1005–1009.

Liang, H.-L., Nien, C.-Y., Liu, H.-Y., Metzstein, M.M., Kirov, N., and Rushlow, C. (2008). The
zinc-finger protein Zelda is a key activator of the early zygotic genome in Drosophila. Nature
456, 400–403.

Liu, G., Wang, W., Hu, S., Wang, X., and Zhang, Y. (2018). Inherited DNA methylation primes
the establishment of accessible chromatin during genome activation. Genome Res. *28*, 998–
1007.

Lott, S.E., Villalta, J.E., Schroth, G.P., Luo, S., Tonkin, L.A., and Eisen, M.B. (2011).
Noncanonical compensation of zygotic X transcription in early Drosophila melanogaster
development revealed through single-embryo RNA-seq. PLoS Biol. *9*, e1000590.

Lu, F., Liu, Y., Inoue, A., Suzuki, T., Zhao, K., and Zhang, Y. (2016). Establishing Chromatin
Regulatory Landscape during Mouse Preimplantation Development. Cell *165*, 1375–1388.

Lu, X., Li, J.M., Elemento, O., Tavazoie, S., and Wieschaus, E.F. (2009). Coupling of zygotic
transcription to mitotic control at the Drosophila mid-blastula transition. Development *136*,
2101–2110.

Lund, E., Liu, M., Hartley, R.S., Sheets, M.D., and Dahlberg, J.E. (2009). Deadenylation of
maternal mRNAs mediated by miR-427 in Xenopus laevis embryos. RNA *15*, 2351–2363.

Mathavan, S., Lee, S.G.P., Mak, A., Miller, L.D., Murthy, K.R.K., Govindarajan, K.R., Tong,
Y., Wu, Y.L., Lam, S.H., Yang, H., et al. (2005). Transcriptome analysis of zebrafish
embryogenesis using microarrays. PLoS Genet. *1*, 260–276.

Newport, J., and Kirschner, M.W. (1982a). A major developmental transition in early xenopus
embryos: I. characterization and timing of cellular changes at the midblastula stage. Cell *30*,
675–686.

Newport, J., and Kirschner, M.W. (1982b). A major developmental transition in early xenopus
embryos: II. control of the onset of transcription. Cell *30*, 687–696.

Nudelman, G., Frasca, A., Kent, B., Sadler, K.C., Sealfon, S.C., Walsh, M.J., and Zaslavsky,
E. (2018). High resolution annotation of zebrafish transcriptome using long-read sequencing.
Genome Res. *28*, 1415-1425.

Owens, N.D.L., Blitz, I.L., Lane, M.A., Patrushev, I., Overton, J.D., Gilchrist, M.J., Cho,
K.W.Y., and Khokha, M.K. (2016). Measuring Absolute RNA Copy Numbers at High Temporal
Resolution Reveals Transcriptome Kinetics in Development. Cell Rep. *14*, 632–647.

Reversade, B., Kuroda, H., Lee, H., Mays, A., and De Robertis, E.M. (2005). Depletion of
Bmp2, Bmp4, Bmp7 and Spemann organizer signals induces massive brain formation in
Xenopus embryos. Development *132*, 3381–3392.

Sabari, B.R., Dall'Agnese, A., Boija, A., Klein, I.A., Coffey, E.L., Shrinivas, K., Abraham, B.J.,
Hannett, N.M., Zamudio, A.V., Manteiga, J.C., et al. (2018). Coactivator condensation at
super-enhancers links phase separation and gene control. Science *361*, eaar3958.

Schneider, S., Steinbeisser, H., Warga, R., and Hausen, P. (1996). Beta-catenin translocation
into nuclei demarcates the dorsalizing centers in frog and fish embryos. Mech. Dev. *57*, 191–
198.

Schohl, A., and Fagotto, F. (2002). Beta-catenin, MAPK and Smad signaling during early
Xenopus development. Development *129*, 37–52.

Shermoen, A.W., and O'Farrell, P.H. (1991). Progression of the cell cycle through mitosis
leads to abortion of nascent transcripts. Cell *67*, 303–310.

Shrinivas, K., Sabari, B.R., Coffey, E.L., Klein, I.A., Boija, A., Zamudio, A.V., Schuijers, J.,
Hannett, N.M., Sharp, P.A., Young, R.A., et al. (2018). Enhancer features that drive formation
of transcriptional condensates. bioRxiv 495606. DOI: https://doi.org/10.1101/495606

Skirkanich, J., Luxardi, G., Yang, J., Kodjabachian, L., and Klein, P.S. (2011). An essential
role for transcription before the MBT in Xenopus laevis. Dev. Biol. *357*, 478–491.

Stack, J.H., and Newport, J.W. (1997). Developmentally regulated activation of apoptosis
early in Xenopus gastrulation results in cyclin A degradation during interphase of the cell

cycle. Development *124*, 3185–3195.

Takahashi, S., Onuma, Y., Yokota, C., Westmoreland, J.J., Asashima, M., and Wright, C.V.E.
(2006). Nodal-related gene Xnr5 is amplified in the Xenopus genome. Genesis *44*, 309–321.

Tan, M.H., Au, K.F., Yablonovitch, A.L., Wills, A.E., Chuang, J., Baker, J.C., Wong, W.H., and
Li, J.B. (2013). RNA sequencing reveals a diverse and dynamic repertoire of the Xenopus
tropicalis transcriptome over development. Genome Res. *23*, 201–216.

Tani, S., Kusakabe, R., Naruse, K., Sakamoto, H., and Inoue, K. (2010). Genomic
organization and embryonic expression of miR-430 in medaka (Oryzias latipes): insights into
the post-transcriptional gene regulation in early development. Gene *449*, 41–49.

Tao, Q., Yokota, C., Puck, H., Kofron, M., Birsoy, B., Yan, D., Asashima, M., Wylie, C.C., Lin,
X., and Heasman, J. (2005). Maternal wnt11 activates the canonical wnt signaling pathway
required for axis formation in Xenopus embryos. Cell *120*, 857–871.

Vassena, R., Boué, S., González-Roca, E., Aran, B., Auer, H., Veiga, A., and Izpisúa
Belmonte, J.C. (2011). Waves of early transcriptional activation and pluripotency program
initiation during human preimplantation development. Development *138*, 3699–3709.

Whyte, W.A., Orlando, D.A., Hnisz, D., Abraham, B.J., Lin, C.Y., Kagey, M.H., Rahl, P.B.,
Lee, T.I., and Young, R.A. (2013). Master transcription factors and mediator establish super-
enhancers at key cell identity genes. Cell *153*, 307–319.

Wu, J., Huang, B., Chen, H., Yin, Q., Liu, Y., Xiang, Y., Zhang, B., Liu, B., Wang, Q., Xia, W.,
et al. (2016). The landscape of accessible chromatin in mammalian preimplantation embryos.
Nature *534*, 652–657.

Yanai, I., Peshkin, L., Jorgensen, P., and Kirschner, M.W. (2011). Mapping gene expression
in two Xenopus species: evolutionary constraints and developmental flexibility. Dev. Cell *20*,
483–496.

Yang, J., Tan, C., Darken, R.S., Wilson, P.A., and Klein, P.S. (2002). Beta-catenin/Tcf-
regulated transcription prior to the midblastula transition. Development *129*, 5743–5752.

Young, J.J., Kjolby, R.A.S., Wu, G., Wong, D., Hsu, S.-W., and Harland, R.M. (2017). Noggin
is required for first pharyngeal arch differentiation in the frog Xenopus tropicalis. Dev. Biol.
426, 245–254.

MAIN FIGURES

Figure 1. Dynamics and Architecture of ZGA in *X. tropicalis*

(A) (i) Genome-wide profiling of RNAPII and total RNA (Owens et al., 2016) to determine temporal ZGA dynamics. (ii) Complementary approach to detect transcriptionally active genes by α-amanitin-induced loss and 4sU-enrichment of nascent (zygotic) transcripts.

(B) Progression of ZGA from the 32-cell to the late gastrula stage based on (i) whole gene body (full-length) occupancy of RNAPII (i.e., RNAPII was enriched across entire gene bodies; see Transparent Methods). Co-aligned: (ii) High time-resolution of total RNA, separated by intron- and exon-derived signals, from fertilization to the late gastrula stage, and (iii) enrichments of 4sU-tagged RNA at the MBT and the mid-gastrula stage. Numbers below RNAPII heat map represent counts of active (blue) and activated (red) genes at the indicated developmental stages. The horizontal dotted line separates RNAPII-engaged (RNAPII^{+}) from non-engaged (RNAPII^{-}) genes as detected until the late gastrula stage. The vertical dotted lines in the total RNA plots indicate the developmental time points of the MBT (white) and the late gastrula stage (black), respectively.

(C) 2D space-filling (Hilbert) curves showing RNAPII recruitment to chromosomes (Chr) at the 128-cell stage and the MBT. A few zygotic genes are highlighted as being active (ON) or not (OFF) based on their engagement with RNAPII.

(D) Alluvial diagram of spatio-temporal ZGA. Tissue-specificity inferred from regional transcript enrichments along the animal-vegetal or the dorso-ventral axes or both (Blitz et al., 2017).

(E) ZGA-associated enrichment of biological processes.

(F) Length of maternal and/or zygotic genes. *, p <1.9e-7 (Wilcoxon rank-sum test against maternal and post-MBT activated genes); r_{effect} 0.06 ('MBT' vs 'Egg') - 0.48 ('128' vs 'mG').

(G) Maternal/zygotic contribution to the transcriptome deduced from full-length RNAPII occupancy and maternally inherited RNA.

Abbreviations: 32, 32-cell; 128, 128-cell; 1K, 1,024-cell; MBT, mid-blastula transition; mG, mid-gastrula; lG, late gastrula; 4sU, 4-thiouridine; Mdn, median; TPM, transcripts per million.

See also Figure S1 and Tables S1 and S2.

Figure 2. Spatio-temporal ZGA Regulated by Canonical Wnt, Nodal and BMP Signals

(A) Spatio-temporal ZGA and nuclear localization of signal mediators β-catenin (canonical Wnt), Smad2 (Nodal) and Smad1 (BMP) (Faure et al., 2000; Larabell et al., 1997; Schohl and Fagotto, 2002) from the 32-cell to the late gastrula stage.

(B) Morphological phenotypes of single and combined signal LOFs at the late gastrula and the mid-tailbud stage. Left ('control') pictures are taken from Gentsch et al. (2018b). Bullet points, failed blastopore formation. Arrowheads, excessive neural fold formation. Scale bar, 0.5 mm.

(C) Heat map (i) and bar graph summary (ii) of ZGA mis-regulated in various LOF embryos: α-amanitin, positive control; control MO, negative control. Abbreviations: B, BMP; N, Nodal; W, canonical Wnt; mPS, maternal Pou5f3/Sox3; mV, maternal VegT; 4x zT, four zygotic T-box TFs (zygotic VegT, Eomes, Tbxt and Tbxt.2).

(D) Biological processes enriched with mis-regulated and control (not down-regulated or

unaffected by the loss of maternal TFs or signaling) sets of zygotic genes under indicated LOFs.

(E) Summary (i) and temporal resolution (ii) of Wnt LOF effects on regional ZGA. Percentages

only refer to the down-regulated genes among all zygotic genes with similar expression ratios

along the animal-vegetal or the dorso-ventral axes.

See also Figure S2, Tables S1 and S3, and Movie S1.

Figure 3. Uniform Gene Expression Across Dorso-Ventral Axis Achieved by Wnt/BMP

Synergy.

(A,C,E,F,G,J) Scatter plots of relative (% to control) transcript levels between indicated LOFs

with each dot (gene) color-coded according to a third attribute: (A,B) relative (% to control)

transcript levels, (E,F) synergy factors between single inductive signals, and (G,J) regional

expression ratios between opposite ends of the indicated axis.

(B,D) Venn diagram of down-regulated genes by indicated LOFs.

(H,K) Box and beeswarm plots of regional expression (as measured along the indicated axes)

depending on increased Wnt-BMP synergy.

(I) WMISH of *tbxt* and *eomes* transcripts under various LOFs. Control and Nodal LOF pictures

are from Gentsch et al. (2018b).

See also Figure S3 and Table S3.

Figure 4. Canonical Wnt, Nodal and BMP Signals Induce the Majority of Regional ZGA.

(A) Total percentage of active (zygotic only and maternal-zygotic) genes mis-regulated by Wnt, Nodal and/or BMP LOF.

(B) Graphical explanations of figure panels (C) and (D).

(C,D) Summary (i) and temporal resolution (ii) of minimal (C) and maximal (D) transcript levels of active genes (separated by regional expression along the primary body axes) detected among Wnt, Nodal and/or BMP LOFs. Percentages only refer to the down-regulated (C) or up-regulated (D) genes among all zygotic genes with the same range of expression ratios along the animal-vegetal or dorso-ventral axes.

(E) Exponential activation of gradually longer genes before the MBT (bulk ZGA) when cell divisions occur at rapid and nearly constant intervals (~20 min at 28ºC). Sequential induction

682 of the canonical Wnt, Nodal and BMP pathway is critical to high percentages of regional ZGA
683 (as measured along the two primary body axes within the indicated halves and quadrants of an
684 early gastrula embryo): e.g. ~89% or ~98% of gene expression enriched in the dorsal half or
685 vegetal-dorsal quadrant, respectively.

Figure S1. Dynamics and Architecture of ZGA in *X. tropicalis*, Related to Figure 1.

(A) Progression of ZGA from the 32-cell to the late gastrula stage based on (i) whole gene body (full-length) occupancy of RNAPII (i.e., RNAPII was enriched across entire gene bodies; see Transparent Methods). Co-aligned: Active histone marks H3K4me1 (ii) and H3K36me3 (iii) (Hontelez et al., 2015) and intronic signal from the high time-resolution profile of poly(A) RNA (iv) (Owens et al., 2016). The horizontal dotted line separates RNAPII-engaged (RNAPII$^+$) from non-engaged (RNAPII$^-$) genes as detected until the late gastrula stage. The vertical dotted lines in the poly(A) RNA plot indicate the developmental time points of the MBT (white) and the late gastrula stage (black), respectively.

(B) Transcript feature levels (mean +/- SD) during the maternal-to-zygotic transition. Asterisk, polyadenylation immediately after fertilization (Collart et al., 2014) transiently increased the intronic signal obtained from the poly(A) RNA samples.

(C) Venn diagram of zygotic genes detected by full-length RNAPII occupancy or reduced exonic or intronic transcript counts upon blocking RNAPII-mediated transcription with α-amanitin.

(D) Pairwise Spearman's correlations (R_s) of enrichment values resulting from RNAPII profiling and 4sU tagging to detect zygotic genes at indicated developmental stages.

(E) Plots of the number of active and newly activated genes (i) or newly activated genes versus the developmental time (i), the number of completed cell divisions (ii) or formed blastomeres (iii).

(F) Alluvial diagram of spatio-temporal ZGA including maternally inherited RNA transcripts of genes not activated by the late gastrula stage. Tissue-specificity inferred from regional transcript enrichments along the animal-vegetal or the dorso-ventral or both axes (Blitz et al., 2017).

(G) Numbers of genes with full-length RNAPII occupancy at indicated developmental stages. Examples of ubiquitously (black) and tissue-specifically (orange) expressed genes are listed to the right.

(H) RNAPII dynamics at the Cys2-His2 [C2H2] zinc finger (ZF) cluster from the 32-cell to the early neurula stage.

(I) Expression dynamics of C2H2 ZF genes normalized to maximal transcript levels recorded between fertilization and 23.5 hpf (Owens et al., 2016).

(J) Phylogenetic tree of the C2H2 ZF genes shown in (H,I). Maternal VegT (mVegT), outgroup TF of this phylogenetic tree.

(K) Box and beeswarm plots showing various metrics of the zygotic/maternal genes (i) and mRNA (ii) during ZGA. Asterisks, significant Wilcoxon rank-sum tests against maternal and post-MBT activated genes and corresponding effect sizes (r_{effect}): # introns, p <2.1e-13, r_{effect} 0.07-0.43; Ø intron (kb), p <7.9e-7, r_{effect} 0.07-0.42; CDS (kb), p <3e-7, r_{effect} 0.05-0.27; mRNA (kb), p <7e-11, r_{effect} 0.06-0.32; 5' UTR (kb), p <0.015, r_{effect} 0.03-0.15; and 3' UTR (kb), p <1.4e-6, r_{effect} 0.04-0.21.

(L) Meta-profiles (mean +/- SD) of RNAPII (separated by developmental stage) and input (negative control) densities at zygotic genes.

728   Abbreviations: 32, 32-cell; 128, 128-cell; 1K, 1,024-cell; MBT, mid-blastula transition; mG, mid-
729   gastrula; lG, late gastrula; eN, early neurula; 4sU, 4-thiouridine; Mdn, median; TPM, transcripts
730   per million.

Figure S2. Effect of Canonical Wnt, Nodal and BMP Signals on ZGA, Related to Figure 2.

(A) Poly(A) RNA profiles of the C2H2 ZF cluster (Figure S1H) for indicated control and LOFs at

the MBT.

(B) Graphical explanations of figure panels (C-F).

(C-F) Summary (i) and temporal resolution (ii) of gene mis-regulations upon the LOF of Nodal

(C) or BMP (D) signaling or maternal Pou5f3/Sox3 (mPou5f3/Sox3) (E) or VegT (mVegT) (F).

Percentages only refer to the down-regulated genes (by ≥1/3 compared to control expression

level) among all zygotic genes with the same range of expression ratios along the animal-

vegetal or the dorso-ventral axes.

741
Figure S3. Relationship between Canonical Wnt, Nodal and BMP to Control Regional

ZGA, Related to Figure 3.

(A-C,E,H-J,L) Scatter plots of relative (% to control) transcript levels between indicated LOFs

with each dot (gene) color-coded according to a third attribute: (A,H,I) relative (% to control)

transcript levels, (B) synergy factors between single inductive signals, and (C,E,J,L) regional

expression ratios between opposite ends of the indicated axis.

(G,N) Venn diagram of down-regulated genes by indicated LOFs.

(D,F,K,M) Box and beeswarm plots of regional expression (as measured along the indicated

axes) depending on increased Wnt-Nodal (D,F) or Nodal-BMP (K,M) synergy.

(O) Coverage and enrichment of Smad and β-catenin-associated DNA motifs (SMAD and

bHSH motifs) at endogenous binding sites of β-catenin, Smad1, Smad2 (Gentsch et al., 2018b)

at indicated developmental stages (color-coded).

SUPPLEMENTAL TABLES

Table S1. Summary of Deep sequencing and Read Alignments, Related to Figure 1 and
2.

The meta-data of ChIP-Seq, 4sU-Seq and RNA-Seq runs includes the developmental stage,
condition, read type, and total and genome-aligned read numbers. *, all rows represent
manually collected biological sample: a, b and c mark biological replicates. †, library was
sequenced twice. Reads were pooled from both sequencing runs. ¶, reported numbers are
reads that are non-redundantly aligned to the genome assembly 7.1 with a mapping quality
(MAPQ) of 10 (DNA) and 255 (RNA). NF, staging according to Nieuwkoop and Faber (1994).
PE/SExx, paired-end or single-end sequencing plus read length.

Sample*	Library	Stage	NF	Condition	PE/SExx	Total reads	Aligned reads¶
1†	RNAPII ChIP	32-cell	6	None	SE50	45,073,468	25,451,597
2†	RNAPII ChIP	128-cell	7	None	SE50/100	65,757,131	41,352,403
3†	RNAPII ChIP	1,024-cell	8	None	SE50	60,445,294	37,661,064
4	RNAPII ChIP	MBT	8.5	None	SE50	29,196,397	18,402,411
5	RNAPII ChIP	mid-gastrula	11	None	SE50	29,268,648	20,085,357
6†	RNAPII ChIP	late gastrula	12+	None	SE50	95,959,570	67,591,256
7	RNAPII ChIP	early neurula	13+	None	SE50	47,039,656	32,889,013
8	4sU-tagged total RNA	Mid-blastula	8+	4sU (4-thiouridine)	PE50	22,202,216	2,738,397
9	4sU-tagged total RNA	Early gastrula	10+	4sU (4-thiouridine)	PE50	20,388,030	5,212,782
10	Total RNA	Mid-blastula	8+	4sU (4-thiouridine), input	PE50	15,087,016	11,451,545
11	Total RNA	Early gastrula	10+	4sU (4-thiouridine), input	PE50	14,902,023	9,771,572
12a	Poly(A)+ RNA	Late blastula	9+	Wnt LOF (β-catenin MO)	PE76	39,439,343	35,540,116
12b	Poly(A)+ RNA	Late blastula	9+	Wnt LOF (β-catenin MO)	PE76	53,523,555	48,364,696
12c	Poly(A)+ RNA	Late blastula	9+	Wnt LOF (β-catenin MO)	PE76	50,568,196	45,534,243
13a	Poly(A)+ RNA	Late blastula	9+	Wnt/Nodal LOF (β-catenin MO, SB431542)	PE76	43,980,513	39,732,439
13b	Poly(A)+ RNA	Late blastula	9+	Wnt/Nodal LOF (β-catenin MO, SB431542)	PE76	44,582,821	40,373,045
13c	Poly(A)+ RNA	Late blastula	9+	Wnt/Nodal LOF (β-catenin MO, SB431542)	PE76	37,243,519	33,648,196
14a	Poly(A)+ RNA	Late blastula	9+	Wnt/BMP LOF (β-catenin MO, LDN193189)	PE76	48,440,801	44,080,313
14b	Poly(A)+ RNA	Late blastula	9+	Wnt/BMP LOF (β-catenin MO, LDN193189)	PE76	40,989,331	37,188,245
14c	Poly(A)+ RNA	Late blastula	9+	Wnt/BMP LOF (β-catenin MO, LDN193189)	PE76	42,494,161	38,527,873
15a	Poly(A)+ RNA	Late blastula	9+	Wnt/Nodal/BMP LOF (β-catenin MO, SB431542, LDN193189)	PE76	52,761,134	47,815,313
15b	Poly(A)+ RNA	Late blastula	9+	Wnt/Nodal/BMP LOF (β-catenin MO, SB431542, LDN193189)	PE76	55,128,794	50,165,817
15c	Poly(A)+ RNA	Late blastula	9+	Wnt/Nodal/BMP LOF (β-catenin MO, SB431542, LDN193189)	PE76	57,414,981	52,101,210
16a	Poly(A)+ RNA	Late blastula	9+	Control (uninjected, DMSO)	PE76	42,248,469	37,989,215
16b	Poly(A)+ RNA	Late blastula	9+	Control (uninjected, DMSO)	PE76	52,066,825	47,105,847
16c	Poly(A)+ RNA	Late blastula	9+	Control (uninjected, DMSO)	PE76	42,875,722	38,893,706

17a	Poly(A)+ RNA	Late blastula	9+	Nodal LOF (SB431542)	PE76	47,946,077	43,367,783
17b	Poly(A)+ RNA	Late blastula	9+	Nodal LOF (SB431542)	PE76	43,201,587	39,270,622
17c	Poly(A)+ RNA	Late blastula	9+	Nodal LOF (SB431542)	PE76	47,497,608	43,105,868
18a	Poly(A)+ RNA	Late blastula	9+	Bmp LOF (LDN193189)	PE76	49,171,585	44,430,698
18b	Poly(A)+ RNA	Late blastula	9+	Bmp LOF (LDN193189)	PE76	35,247,462	31,937,946
18c	Poly(A)+ RNA	Late blastula	9+	Bmp LOF (LDN193189)	PE76	40,645,238	36,899,368
19a	Poly(A)+ RNA	Late blastula	9+	Nodal/BMP LOF (SB431542, LDN193189)	PE76	49,865,847	45,127,683
19b	Poly(A)+ RNA	Late blastula	9+	Nodal/BMP LOF (SB431542, LDN193189)	PE76	48,875,533	44,167,262
19c	Poly(A)+ RNA	Late blastula	9+	Nodal/BMP LOF (SB431542, LDN193189)	PE76	49,397,754	44,710,470
20a	Poly(A)+ RNA	Early gastrula	10+	4x zT (4 zygotic T-box TFs) LOF (zVegT, Eomes, Tbxt, Tbxt.2 MO)	PE76	70,597,621	63,789,737
20b	Poly(A)+ RNA	Early gastrula	10+	4x zT (4 zygotic T-box TFs) LOF (zVegT, Eomes, Tbxt, Tbxt.2 MO)	PE76	62,453,340	56,799,027
20c	Poly(A)+ RNA	Early gastrula	10+	4x zT (4 zygotic T-box TFs) LOF (zVegT, Eomes, Tbxt, Tbxt.2 MO)	PE76	66,378,689	60,011,884
20d	Poly(A)+ RNA	Early gastrula	10+	4x zT (4 zygotic T-box TFs) LOF (zVegT, Eomes, Tbxt, Tbxt.2 MO)	PE76	53,693,870	48,621,416
21a	Poly(A)+ RNA	Mid-gastrula	11+	4x zT (4 zygotic T-box TFs) LOF (zVegT, Eomes, Tbxt, Tbxt.2 MO)	PE76	60,071,724	53,786,168
21b	Poly(A)+ RNA	Mid-gastrula	11+	4x zT (4 zygotic T-box TFs) LOF (zVegT, Eomes, Tbxt, Tbxt.2 MO)	PE76	49,856,218	44,780,174
21c	Poly(A)+ RNA	Mid-gastrula	11+	4x zT (4 zygotic T-box TFs) LOF (zVegT, Eomes, Tbxt, Tbxt.2 MO)	PE76	57,420,101	51,441,091
21d	Poly(A)+ RNA	Mid-gastrula	11+	4x zT (4 zygotic T-box TFs) LOF (zVegT, Eomes, Tbxt, Tbxt.2 MO)	PE76	57,421,959	51,207,446
22a	Poly(A)+ RNA	Late gastrula	12+	4x zT (4 zygotic T-box TFs) LOF (zVegT, Eomes, Tbxt, Tbxt.2 MO)	PE76	59,905,689	53,050,487
22b	Poly(A)+ RNA	Late gastrula	12+	4x zT (4 zygotic T-box TFs) LOF (zVegT, Eomes, Tbxt, Tbxt.2 MO)	PE76	76,564,052	67,801,662
22c	Poly(A)+ RNA	Late gastrula	12+	4x zT (4 zygotic T-box TFs) LOF (zVegT, Eomes, Tbxt, Tbxt.2 MO)	PE76	54,924,184	48,697,757
22d	Poly(A)+ RNA	Late gastrula	12+	4x zT (4 zygotic T-box TFs) LOF (zVegT, Eomes, Tbxt, Tbxt.2 MO)	PE76	56,546,178	49,962,419
23a	Poly(A)+ RNA	Early gastrula	10+	Control (standard control MO)	PE76	57,881,388	52,311,548
23b	Poly(A)+ RNA	Early gastrula	10+	Control (standard control MO)	PE76	49,193,836	44,649,709
23c	Poly(A)+ RNA	Early gastrula	10+	Control (standard control MO)	PE76	78,605,062	70,838,519
23d	Poly(A)+ RNA	Early gastrula	10+	Control (standard control MO)	PE76	67,143,066	60,397,172
24a	Poly(A)+ RNA	Mid-gastrula	11+	Control (standard control MO)	PE76	50,011,338	44,604,698
24b	Poly(A)+ RNA	Mid-gastrula	11+	Control (standard control MO)	PE76	54,847,378	48,981,425
24c	Poly(A)+ RNA	Mid-gastrula	11+	Control (standard control MO)	PE76	56,633,663	50,510,544
24d	Poly(A)+ RNA	Mid-gastrula	11+	Control (standard control MO)	PE76	69,504,876	61,831,845
25a	Poly(A)+ RNA	Late gastrula	12+	Control (standard control MO)	PE76	53,566,350	47,558,871
25b	Poly(A)+ RNA	Late gastrula	12+	Control (standard control MO)	PE76	36,043,121	31,918,744
25c	Poly(A)+ RNA	Late gastrula	12+	Control (standard control MO)	PE76	48,184,154	42,775,598
25d	Poly(A)+ RNA	Late gastrula	12+	Control (standard control MO)	PE76	39,860,708	35,323,477
Total						2,948,234,870	2,524,336,789

Table S2. Temporal Progression of ZGA, Related to Figure 1.

Activated genes are listed according to the earliest developmental stage of full-length RNAPII occupancy. The lists also include the overall enrichment of RNAPII across the gene body, the maternal contribution (TPM) (Owens et al., 2016) and the enrichment of de novo synthesized transcripts determined by 4sU tagging at the MBT and the early gastrula stage.

Table S3. Differential ZGA Analysis, Related to Figure 2, 3 and 4.

Differential expression analysis of genes showing ≥50% (FDR ≤10%) reduced transcript levels
in α-amanitin-injected embryos. Normalized transcript levels (inferred from exon or intron
counts) are scaled (percentage, %) to the expression level in control embryos. The list also
includes FDRs, expression ratios across the animal-vegetal or dorso-ventral axis (Blitz et al.,
2017), the earliest developmental stage of full-length RNAPII occupancy, and average
expression levels between 0 and 1 hpf (maternal) and 5 and 9 hpf (from the MBT to the mid-
gastrula stage).

Movie S1. Quadruple LOF of Zygotic T-box TFs, Related to Figure 2.
Simultaneous filming of the vegetal (top row) and animal (bottom row) hemisphere of 4x zT
LOF (labelled as T-box KD in the movie) (left) and control (right) embryos from early gastrula
to mid-tailbud stage.

TRANSPARENT METHODS

CONTACT FOR REAGENTS AND RESOURCE SHARING

Further information and requests for resources and reagents should be directed to and will be
fulfilled by the Lead Contact, James C. Smith (jim.smith@crick.ac.uk).

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Xenopus tropicalis Manipulation
Standard procedures were used for ovulation, fertilization, and manipulation and incubation of
embryos (Khokha et al., 2002; Sive et al., 2000). Briefly, frogs were obtained from Nasco
(Wisconsin, USA). Ovulation was induced by injecting serum gonadotropin (Intervet) and
chorionic gonadotropin (Intervet) into the dorsal lymph sac of mature female frogs. Eggs were
fertilized *in vitro* with sperm solution consisting of 90% Leibovitz's L-15 medium (Thermo Fisher
Scientific, Cat#11415064) and 10% fetal bovine serum (Thermo Fisher Scientific,
Cat#10500056). After 10 min, fertilized eggs were de-jellied with 2.2% (w/v) L-cysteine (Merck,
Cat#168149) equilibrated to pH 8.0. Embryos were cultured in 5% Marc's Modified Ringer's
solution (MMR) (5 mM NaCl, 0.1 mM KCl, 0.1 mM $CaCl_2$, 0.05 mM $MgSO_4$ and 0.25 mM HEPES
pH7.5) at 21ºC-28ºC. Embryos were staged according to Nieuwkoop and Faber (1994). All
Xenopus work fully complied with the UK Animals (Scientific Procedures) Act 1986 as
implemented by the Francis Crick Institute.

Chromatin immunoprecipitation (ChIP)
ChIP was carried out as detailed previously (Gentsch and Smith, 2017). Briefly, de-jellied *X.*
tropicalis embryos were fixed at room temperature with 1% formaldehyde (Merck, Cat#F8775)
in 1% MMR for 25 min. The fixation time was extended to 45 min for pre-gastrula stages. The
following number of embryos were used for ChIP-Seq: 1,400 at the 32-cell stage, 1,000 at the

128-cell stage, 700 at the 1,024-cell stage, 450 at the MBT and 350 for the post-MBT stages.

Fixation was terminated by rinsing embryos three times with ice-cold 1% MMR. Fixed embryos

were homogenized in CEWB1 (150 mM NaCl, 1 mM EDTA, 1% (v/v) Igepal CA-630 [Merck,

Cat#I3021], 0.25% (w/v) sodium deoxycholate [Merck, Cat#SRE0046], 0.1% (w/v) sodium

dodecyl sulfate [Merck, Cat#71729] and 10 mM Tris-HCl pH 8.0) supplemented with 0.5 mM

DL-Dithiothreitol (Fluorochem, Cat#M02712) and protease inhibitors (Roche,

Cat#11873580001). The homogenate was left on ice for 5 min and then centrifuged at 1,000 g

(4ºC) for 5 min. Homogenization and centrifugation was repeated once before resuspending

the pellet in 1-3 ml CEWB1. Chromatin was solubilized and fragmented by microtip-mediated

ultra-sonication (Misonix 3000 sonicator with a tapered 1/16-inch microtip). The solution of

fragmented chromatin was cleared by centrifuging at 16,000 g (4ºC) for 5 min. About 1% of the

cleared chromatin extract was set aside for the input sample (negative control). The remaining

chromatin was incubated overnight at 4ºC on a vertical rotor (10 rpm) with 20 µl of the mouse

monoclonal anti-RNAPII (8WG16) (Covance, Cat#MMS-126R; RRID: AB_10013665) antibody.

After adding 100 µl of washed protein G magnetic beads (Thermo Fisher Scientific,

Cat#10003D) the solution was incubated for another 4 h at 4ºC on a vertical rotor (10 rpm). The

beads were washed eight times in CEWB1 and once in TEN (10 mM Tris-HCl pH 8.0, 150 mM

NaCl and 1 mM EDTA) at 4ºC. ChIP was eluted off the beads twice with 100 µl SDS elution

buffer (50 mM Tris-HCl pH 8.0, 1 mM EDTA and 1% (w/v) sodium dodecyl sulfate) at 65ºC.

ChIP eluates were pooled before reversing DNA-protein cross-links. Input (filled up to 200 µl

with SDS elution buffer) and ChIP samples were supplemented with 10 µl 5 M NaCl and

incubated at 65ºC for 6-16 h. Samples were treated with proteinase K (Thermo Fisher Scientific,

Cat#AM2548) and RNase A (Thermo Fisher Scientific, Cat#12091021) to remove any proteins

and RNA from the co-immunoprecipitated DNA fragments. The DNA was purified with

phenol:chloroform:isoamyl alcohol (25:24:1, pH 7.9) (Thermo Fisher Scientific, Cat#AM9730)

using 2.0-ml Phase Lock Gel Heavy microcentrifuge tubes (VWR, Cat#733-2478) for phase

separation and precipitated with 1/70 volume of 5 M NaCl, 2 volumes of absolute ethanol and

15 µg GlycoBlue (Thermo Fisher Scientific, Cat#AM9516). After centrifugation, the DNA pellet

was air-dried and dissolved in 11 µl elution buffer (10 mM Tris-HCl pH 8.5). The DNA

concentration was determined on a Qubit fluorometer using high-sensitivity reagents for

detecting double-stranded DNA (10 pg/µl to 100 ng/µl) (Thermo Fisher Scientific, Cat#Q33231).

ChIP-Seq Library Preparation

Using the KAPA Hyper Prep Kit (Roche, Cat#KK8504), 2.5-5 ng ChIP DNA or 5 ng input DNA were

converted into indexed paired-end libraries as previously described (Gentsch and Smith, 2017).

Briefly, DNA fragments were end-repaired and A-tailed for 30 min at 20ºC followed by 30 min at

65ºC before cooling to 4ºC. 7.5 pmol TruSeq (single index) Y-adapters (IDT) were ligated to the

DNA fragment ends for 20 min at 20ºC. The DNA ligation product was extracted with 0.8x SPRI

(solid phase reversible immobilisation) beads (Beckman Coulter, Cat#A63882) and amplified in

five PCR cycles (15 sec at 98ºC, 30 sec at 60ºC and 30 sec at 72ºC) using the KAPA high-fidelity

polymerase master mix (Roche, Cat#KK2602) and 25 pmol Illumina P5 (forward) and P7 (reverse)

primers (IDT). After cleaning up the PCR reaction with 1x SPRI beads, the DNA library was size-

separated by electrophoresis using E-gel EX agarose gels (Thermo Fisher Scientific,

Cat#G401002). A gel slice containing DNA ranging from 250 to 450 bp in size was dissolved

shaking in 350 µl QG buffer (Qiagen) using a thermomixer (1,000 rpm) at room temperature. The

DNA was purified with MinElute columns (Qiagen, Cat#28604) and eluted off these columns twice

using 11 µl elution buffer (10 mM Tris-HCl pH 8.5). The library was re-amplified using another 6-8

PCR cycles yielding 100-200 ng DNA without adapter dimer contamination. The DNA library was

cleaned up with 1x SPRI beads.

Illumina Sequencing

All sequencing libraries were quality controlled: The DNA yield and fragment size distribution were

determined by fluorometry and chip-based capillary electrophoresis, respectively. ChIP-Seq and

RNA-Seq libraries were sequenced on the Illumina HiSeq 2500 and 4000, respectively, by the

Advanced Sequencing Facility of the Francis Crick Institute. Sequencing samples and read

alignment results are summarized in Table S1.

Post-Sequencing Analysis of ChIP-Seq

Single reads of maximal 50 bases were processed using trim_galore v0.4.2 (Babraham Institute,

UK) to trim off low-quality bases (default Phred score of 20, i.e. error probability was 0.01) and

adapter contamination from the 3' end. Processed reads were aligned to the *X. tropicalis* genome

assembly v7.1 and v9.1 (for Hilbert curves) running Bowtie2 v2.2.9 (Langmead and Salzberg,

2012) with default settings (Table S1). Alignments were converted to the HOMER's tag density

format (Heinz et al., 2010) with redundant reads being removed (makeTagDirectory -single -tbp 1

-unique -mapq 10 -fragLength 175 -totalReads all). Only uniquely aligned reads (i.e. MAPQ ≥10)

were processed. We pooled all input alignments from various developmental stages (Gentsch et

al., 2018b). This created a comprehensive mappability profile that covered ~400 million unique

base pair positions. For Hilbert curves, tag densities were generated across the genome v9.1 using

sliding (200-bp increments) 400-bp window. Background signals (<0.3 reads per 1 million mapped

reads) were removed. Blacklisted (Gentsch et al., 2018b) regions (except for MIR-427) were

excluded using intersectBed (-v -f 0.5) from BEDtools v2.25.0 (Quinlan and Hall, 2010).

Detecting Zygotic and Maternal Genes Using RNAPII Profiling and High Time-Resolution

Transcriptomics

Normalized RNAPII and input tag densities were calculated across the gene body in 10 bins of

equal size. Gene annotations v7.1 were altered based on a few known zygotic isoforms and some

corrections obtained from assembling total and poly(A) RNA (Owens et al., 2016) from stage 6 to

stage 12.5 *de novo* (Pertea et al., 2016). A few genes had previously been annotated as gene

clusters due to assembly uncertainties. We reduced the annotation of polycistronic MIR-427 to the

minus arm (scaffold_3b:3516900-3523400) and only monitored *nodal3.5* and *nodal5.3* within their

respective gene clusters. Gene bodies with <40% mappability were removed. Here, the threshold
of mappability per bin was set at 10% of the input read density averaged across all gene bodies in
use. Subsequently, enrichment values were only obtained for all mappable bins by dividing read
densities of RNAPII and input. Further, we restricted the analysis to genes for which ≥3 transcripts
per million (TPM) could be detected on average over three consecutive time points (i.e. over the
developmental time of 1 h) of a high-resolution profile of total RNA (Owens et al., 2016) from
fertilization to after gastrulation (stage 13). Genes were considered active when RNAPII
enrichments along their full length (see thresholds below) and corresponding transcripts (≥0.1
TPM) were simultaneously detected. Transcript levels were calculated over three consecutive time
points +/- 1 h from the developmental stage of RNAPII profiling. RNAPII enrichment covered ≥80%
of the mappable gene body and reached at least one of the following thresholds: (1) 2.6-fold, (2)
1.8-fold and 1.4-fold at the next or previous stage, (3) 1.4-fold and 1.8-fold at the next or previous
stage, or (4) 1.4-fold over three consecutive stages. The heatmap (Figure 1B and S1A) was sorted
by the developmental stage (1st) and the overall fold (2nd) of RNAPII enrichment. Zygotic and
maternal contributions to transcriptome (Figure 1G) were based on RNAPII enrichment (see
above) and mean transcript levels (≥0.1 TPM) detected between 0 and 1 hpf, respectively.
900
Peak Calling and Motif Enrichment Analysis
Peak calling and motif enrichment analysis were carried out as previously reported (Gentsch et
al., 2018b). Briefly, HOMER v4.8.3 (Heinz et al., 2010) was used to identify the binding sites of
Smad1 (Gentsch et al., 2018b), Smad2 (Chiu et al., 2014; Gentsch et al., 2018b; Yoon et al., 2011)
and β-catenin (Gentsch et al., 2018b; Nakamura et al., 2016) by virtue of ChIP-enriched read
alignments (hereafter called peaks): findpeaks -style factor -minDist 175 -fragLength 175 -
inputFragLength 175 -fdr 0.001 -gsize 1.435e9 -F 3 -L 1 -C 0.97. This means that both ChIP and
input alignments were extended 3' to 175 bp for the detection of significant (FDR ≤0.1%) peaks
being separated by ≥175 bp. The effective size of the *X. tropicalis* genome assembly v7.1 was set
to 1.435 billion bp, an estimate obtained from the mappability profile (Gentsch et al., 2018b). These
peaks showed equal or higher tag density than the surrounding 10 kb, ≥3-fold more tags than the
input and ≥0.97 unique tag positions relative to the expected number of tags. To further eliminate
any false positive peaks, we removed any peaks with <0.5 CPM and those falling into blacklisted
regions showing equivocal mappability due to genome assembly errors, gaps or simple/tandem
repeats. Regions of equivocal mappability were identified by a two-fold lower (poor) or three-fold
higher (excessive) read coverage than the average detected in 400-bp windows sliding at 200-bp
intervals through normalized ChIP input and DNase-digested naked genomic DNA (Gentsch et al.,
2018b). All identified regions ≤800 bp apart were subsequently merged. Gap coordinates were
obtained from the Francis Crick mirror site of the UCSC genome browser
(http://genomes.crick.ac.uk). Simple repeats were masked with RepeatMasker v4.0.6 (Smit et al.)
using the crossmatch search engine v1.090518 (Phil Green) and the following settings:
RepeatMasker -species "xenopus silurana tropicalis" -s -xsmall. Tandem repeats were masked
with Jim Kent's trfBig wrapper script of the Tandem Repeat Finder v4.09 (Benson, 1999) using the

following settings: weight for match, 2; weight for mismatch, 7; delta, 7; matching probability, 80;

indel probability, 10; minimal alignment score, 50; maximum period size, 2,000; and longest

tandem repeat array (-l), 2 [million bp]. The enrichment and occurrence of predetermined DNA

binding motifs was calculated using 100 bp centred across the top 2,000 peaks per chromatin

feature and developmental stage: findMotifsGenome.pl -size 100 -mknown -nomotif.

Injections and Treatments of Embryos

Microinjections were performed using calibrated needles and embryos equilibrated in 4% (w/v)

Ficoll PM-400 (Merck, Cat#F4375) in 5% MMR. Microinjection needles were generated from

borosilicate glass capillaries (Harvard Apparatus, GC120-15) using the micropipette puller

Sutter p97. Maximally three nanolitres were injected into the animal hemisphere of de-jellied

zygotes using the microinjector Narishige IM-300. Embryos were transferred to fresh 5% MMR

(without Ficoll PM-400) once they reached about the mid-blastula stage.

For profiling the nascent transcriptome, embryos were injected with 75 ng 4-thiouridine-5'-

triphosphate (4sU) (TriLink BioTechnologies, Cat#N-1025), which is incorporated into newly

synthesized transcripts.

Loss-of-functions (LOFs) were generated by treating embryos with small molecule inhibitors

and/or injecting them with morpholinos (MOs) or α-amanitin. MOs were designed and produced

by Gene Tools (Oregon, USA) to block splicing (MO_{splice}) or translation (MO_{transl}): maternal

Pou5f3/Sox3 (mPou5f3/Sox3) LOF, 5 ng *Pou5f3.2* MO_{transl} (Chiu et al., 2014; Gentsch et al.,

2018b; GCTGTTGGCTGTACATAGTGTC), 5 ng *Pou5f3.3* MO_{transl} (Chiu et al., 2014; Gentsch

et al., 2018b; TACATTGGGTGCAGGGACCCTCTCA) and 5 ng *Sox3* MO_{transl} (Gentsch et al.,

2018b; GTCTGTGTCCAACATGCTATACATC); maternal VegT (mVegT) LOF, 10 ng m*VegT*

MO_{transl} (Gentsch et al., 2018b; Rana et al., 2006; TGTGTTCCTGACAGCAGTTTCTCAT);

canonical Wnt LOF, 5 ng *β-catenin* MO_{transl} (Heasman et al., 2000;

TTTCAACAGTTTCCAAAGAACCAGG); LOF of four zygotic T-box TFs (4x zT LOF), 2.5 ng *tbxt*

(Xbra, t) MO_{splice} (Gentsch et al., 2013; TGGAGAGACCCTGATCTTACCTTCC), 2.5 ng *tbxt*

(Xbra, t) MO_{transl} (Gentsch et al., 2013; GGCTTCCAAGCGCACACACTGGG), 2.5 ng *tbxt.2*

(Xbra3, t2) MO_{splice} (Gentsch et al., 2013; GAAAGGTCCATATTCTCTTACCTTC), 2.5 ng *tbxt.2*

(*Xbra3, t2*) MO_{transl} (Gentsch et al., 2013; AGCTGTGCCTGTGCTCATTGTATTG), 5 ng z*VegT*

MO_{transl} (Fukuda et al., 2010; Gentsch et al., 2013; CATCCGGCAGAGAGTGCATGTTCCT) and

5 ng *eomes* MO_{splice} (Fukuda et al., 2010; Gentsch et al., 2013;

GAACATCCTCCTGCAAAGCAAAGAC); control MO, 5-20 ng standard control MO

(CCTCTTACCTCAGTTACAATTTATA) according to the dose used for the β-catenin, mVegT

and 4x zT LOF experiment; and 30 pg α-amanitin (BioChemica, Cat#A14850001). To block

Nodal (Nodal LOF) and BMP (BMP LOF) signaling, embryos were treated with 100 µM

SB431542 (Tocris, Cat#1614) and/or 10 µM LDN193189 (Selleckchem, Cat#S2618) from the

8-cell stage onwards. Control embryos were treated accordingly with DMSO, in which these

antagonists were dissolved. Transcriptional effects of combinatorial signal LOF were

determined at late blastula stage (stage 9$^+$), while those of all other maternal LOFs were

determined over three consecutive time points: the MBT (stage 8+), the late blastula (stage 9$^+$)

and the early gastrula (stage 10$^+$) stage. The 4x zT LOF was transcriptionally profiled at early,

mid and late gastrula stage (stage 10$^+$, 11$^+$ and 12$^+$). The 4x zT LOF comparison has four

biological replicates (n=4). All other comparisons entail three biological replicates (n=3).

Extraction of Total RNA

Embryos were homogenized in 800 µl TRIzol reagent (Thermo Fisher Scientific,

Cat#15596018) by vortexing. The homogenate was either snap-frozen in liquid nitrogen and

stored at -80ºC or processed immediately. For phase separation, the homogenate together with

0.2x volume of chloroform was transferred to pre-spun 2.0-ml Phase Lock Gel Heavy

microcentrifuge tubes (VWR), shaken vigorously for 15 sec, left on the bench for 2 min and

spun at ~16,000 g (4ºC) for 5 min. The upper phase was mixed well with one volume of 95-

100% ethanol and spun through the columns of the RNA Clean & Concentrator 25 Kit (Zymo

Research, Cat#R1017) at ~12,000 g for 30 sec. Next, the manufacturer's instructions were

followed for the recovery of total RNA (>17 nt) with minor modifications. First, the flow-through

of the first spin was re-applied to the column. Second, the RNA was treated in-column with 3 U

Turbo DNase (Thermo Fisher Scientific, Cat#AM2238). Third, the RNA was eluted twice with

25 µl molecular-grade water. The concentration was determined on the NanoDrop 1000

spectrophotometer or by fluorometry before depleting ribosomal RNA from total RNA (Profiling

the Nascent Transcriptome).

Tagging the Nascent Transcriptome

Thirty 4sU-injected embryos were collected at the MBT and the early-to-mid gastrula stage.

Total RNA was extracted as outlined above. The 4sU-tagging was performed according to Gay

et al. (2014) with few minor modifications. The RNA Clean & Concentrator 5 Kit (Zymo

Research, Cat#R1013) was used to purify RNA. Briefly, the Ribo-Zero Gold rRNA Removal Kit

(Illumina, Cat#MRZG126) was used according to the manufacturer's instructions to deplete

ribosomal RNA from ~10 µg total RNA. The RNA was purified and fragmented for 4 min at 95ºC

using the NEBNext Magnesium RNA Fragmentation Module (NEB, Cat#E6150). The RNA was

purified again before conjugating HDPD-Biotin (Thermo Fisher Scientific, Cat#21341) to 4sU

via disulfide bonds for 3 h in the dark. Purified RNA was mixed with Streptavidin beads (Thermo

Fisher Scientific, Cat#65305) to pull down biotin-tagged RNA. The RNA was eluted off the

beads by treating them twice with 100 µl pre-heated (80ºC) 100 mM β-mercaptoethanol (Merck,

Cat#M6250), which breaks the disulfide bond between Biotin and 4sU. Subsequently, the RNA

was converted into a deep sequencing library by following the manual instructions (Rev. C,

8/2014) of the ScriptSeq v2 RNA-Seq Library Preparation Kit (Illumina, Cat#SSV21106) starting

with 4.1.A. (Anneal the cDNA Synthesis Primer) and 4.1.B. (Synthesize cDNA), RNA and

ending with part 3.C (Synthesize 3'-Tagged DNA) to 3.G. (Assess Library Quantity and Quality).

cDNA was purified using 1.8x SPRI beads. Input and 4sU-enriched cDNA were PCR-amplified

with 11 and 15 cycles, respectively. The 4sU RNA-Seq library was purified with 1x SPRI beads.

Post-Sequencing Analysis of 4sU Tagging

Paired-end reads were aligned to the *X. tropicalis* transcriptome assembly v7.1 running Bowtie2 (Langmead and Salzberg, 2012) with the following constraints: -k 200 (maximal allowed number of alignments per fragment) -X 800 (maximum fragment length in bp) --rdg 6,5 (penalty for read gaps of length N, 6+N*5) --rfg 6,5 (penalty for reference gaps of length N, 6+N*5) --score-main L,-.6,-.4 (minimal alignment score as a linear function of the read length x, f(x) = -0.6 - 0.4*x) --no-discordant (no paired-end read alignments breaching maximum fragment length X) --no-mixed (only concordant alignment of paired-end reads). Only read pairs that uniquely align to one gene were counted. Raw read counts were normalized with DESeq2 v1.22.1 (Love et al., 2014) and then scaled to the input.

Poly(A) RNA-Seq Profiling

10-15 embryos were collected per stage and condition. Total RNA was extracted as outlined above. Libraries were made from ~1 μg total RNA by following the low-sample protocol of the TruSeq RNA Library Prep Kit v2 (Illumina, Cat#RS-122-2001) with a few modifications. First, 1 μl cDNA purified after second strand synthesis was quantified on a Qubit fluorometer using high-sensitivity reagents for detecting double-stranded DNA (10 pg/μl to 100 ng/μl). By this stage, the yield was ~10 ng. Second, the number of PCR cycles was reduced to eight to avoid products of over-amplification such as chimera fragments.

Poly(A) RNA-Seq Read Alignment

Paired-end reads were aligned to the *X. tropicalis* genome assembly v7.1 using STAR v2.5.3a (Dobin et al., 2013) with default settings. The alignment was guided by a revised version of the gene models v7.2 (Collart et al., 2014) to improve mapping accuracy across splice junctions. The alignments were sorted by read name using the sort function of Samtools v1.3.1 (Li et al., 2009). Exon and intron counts (-t 'exon;intron') were extracted from unstranded (-s 0) alignment files using VERSE v0.1.5 (Zhu et al., 2016) in featureCounts (default) mode (-z 0). Intron coordinates were adjusted to exclude any overlap with exon annotation. For visualization, genomic BAM files of biological replicates were merged using Samtools and converted to the bigWig format. These genome tracks were normalized to the wigsum of 1 billion excluding any reads with mapping quality <10 using the python script bam2wig.py from RSeQC v2.6.4 (Wang et al., 2012).

Differential Gene Expression Analysis

Differential expression analysis was performed with both raw exon and intron counts excluding those belonging to ribosomal and mitochondrial RNA using the Bioconductor/R package DESeq2 v1.22.1 (Love et al., 2014). In an effort to find genes with consistent fold changes over time, p-values were generated according to a likelihood ratio test reflecting the probability of rejecting the reduced (~ developmental stage) over the full (~ developmental stage + condition) model. Resulting p-values were adjusted to obtain false discovery rates (FDR) according to the Benjamini-Hochburg procedure with thresholds on Cook's distances and independent filtering being switched off. Equally, combinatorial LOF profiling and regional expression datasets (Blitz

et al., 2017) without time series were subjected to likelihood ratio tests with reduced (~ 1) and
full (~ condition) models for statistical analysis. Fold changes of intronic and exonic transcript
levels were calculated for each developmental stage and condition using the mean of DESeq2-
normalized read counts from biological replicates. Both intronic and exonic datasets were
filtered for ≥10 DESeq2-normalized read counts that were detected at least at one
developmental stage in all uninjected or DMSO-treated samples. Gene-specific fold changes
were removed at developmental stages that yielded <10 normalized read counts in
corresponding control samples. Next, the means of intronic and exonic fold changes were
calculated across developmental stages. The whole dataset was confined to 3,318 genes for
which at least 50% reductions (FDR ≤10%) in exonic (default) or intronic counts could be
detected in α-amanitin-injected embryos. Regional expression was based on exonic read
counts by default unless the intronic fold changes were significantly (FDR ≤10%) larger than
the exonic fold changes (Table S3). For the hierarchical clustering of relative gene expression
(Figure 2C), increased transcript levels were masked and only data points from signal LOFs,
mPou5f3/Sox3 LOF and 4x zT LOF embryos were used. Euclidean distance-derived clusters
were linked according to Ward's criterion and sorted using the optimal leaf ordering (OLO)
algorithm. The synergy factor (SF) between signals x and y (Figures 3 and S3) were calculated
as follows: $SF_{xy} = \Delta_{xy} / (\Delta_x + \Delta_y)$. Δ is the relative loss of gene expression caused by signal
depletion. For these calculations, any gene upregulations were neutralised (i.e. set to 1).

Analysis of Enriched Gene Ontology (GO) Terms
Over-represented GO terms were found by applying hypergeometric tests of the
Bioconductor/R package GOstats v2.42.0 (Falcon and Gentleman, 2007) on gene lists. The
process was also supported by the Bioconductor/R packages GSEABase v1.44.0 (Morgan et
al., 2017) and GO.db v3.4.1 (Carlson et al., 2007). The gene universe was associated with GO
terms by means of BLAST2GO (Conesa et al., 2005) as previously outlined (Collart et al., 2014;
Gentsch et al., 2015).

Generation of Hybridization Probes
Plasmids *X. laevis eomes* pCRII-TOPO (Gentsch et al., 2013) and *X. laevis tbxt (Xbra, t)* pSP73
(Smith et al., 1991) were linearized by restriction digestion (*Bam*HI and *Bgl*II, respectively) and
purified using the QIAquick PCR Purification Kit (Qiagen, Cat#28104). The hybridization probes
were transcribed from ~1 µg linearized plasmid using 1x digoxigenin-11-UTP (Roche,
Cat#11277065910), 40 U RiboLock RNase inhibitor (Thermo Fisher Scientific, Cat#EO0381), 1x
transcription buffer (Roche) and T7 RNA polymerase (Roche, Cat#10881767001) at 37ºC for 2 h.
The probe was treated with 2 U Turbo DNase (Thermo Fisher Scientific) to remove the DNA
template and purified by LiCl precipitation. RNA was diluted to 10 ng/µl (10x stock) with
hybridization buffer. The hybridization buffer (stored at -20ºC) consists of 50% formamide (Fisher
Scientific, Cat#10052370), 5x saline sodium citrate (SSC), 1x Denhardt's solution (Thermo Fisher
Scientific, Cat#750018), 10 mM EDTA, 1 mg/ml torula RNA (Merck, Cat#R6625), 100 µg/ml

heparin (Merck, Cat#H4784), 0.1% (v/v) Tween-20 (Merck, Cat#P9416) and 0.1% (w/v) CHAPS

(Merck, Cat#C3023).

1082

Whole-Mount In Situ Hybridization (WMISH)

WMISH was conducted using digoxigenin-labeled RNA probes (Monsoro-Burq, 2007; Sive et

al., 2000). Briefly, *X. tropicalis* embryos were fixed in MEMFA (100 mM MOPS pH 7.4, 2 mM

EDTA, 1 mM $MgSO_4$ and 3.7% formaldehyde) at room temperature for 1 h. The embryos were

then washed once in 1x PBS and two to three times in ethanol. Fixed and dehydrated embryos

were kept at -20ºC for ≥24 h to ensure proper dehydration before starting hybridization.

Dehydrated embryos were washed once more in ethanol before rehydrating them in two steps

to PBT (1x PBS and 0.1% (v/v) Tween-20). Embryos were treated with 5 µg/ml proteinase K

(Thermo Fisher Scientific) in PBT for 6-8 min, washed briefly in PBT, fixed again in MEMFA for

20 min and washed three times in PBT. Embryos were transferred into baskets, which were

kept in an 8x8 microcentrifuge tube holder sitting inside a 10x10 slot plastic box filled with PBT.

Baskets were built by replacing the round bottom of 2-ml microcentrifuge tubes with a Sefar

Nitex mesh. This container system was used to readily process several batches of embryos at

once. These baskets were maximally loaded with 40 to 50 *X. tropicalis* embryos. The

microcentrifuge tube holder was used to transfer all baskets at once and to submerge embryos

into subsequent buffers of the WMISH protocol. Next, the embryos were incubated in 500 µl

hybridization buffer (see recipe above) for 2 h in a hybridization oven set to 60ºC. After this pre-

hybridization step, the embryos were transferred into 500 µl digoxigenin-labeled probe (1 ng/µl)

preheated to 60ºC and further incubated overnight at 60ºC. The pre-hybridization buffer was

kept at 60ºC. The next day embryos were transferred back into the pre-hybridization buffer and

incubated at 60ºC for 10 min. Subsequently, they were washed three times in 2x SSC/0.1%

Tween-20 at 60ºC for 10 min, twice in 0.2x SSC/0.1% Tween-20 at 60ºC for 20 min and twice

in 1x maleic acid buffer (MAB) at room temperature for 5 min. Next, the embryos were treated

with blocking solution (2% Blocking Reagent [Merck, Cat#11096176001] in 1x MAB) at room

temperature for 30 min, and incubated in antibody solution (10% lamb serum [Thermo Fisher

Scientific, Cat#16070096], 2% Blocking Reagent [Merck], 1x MAB and 1:2,000 Fab fragments

from polyclonal anti-digoxigenin antibodies conjugated to alkaline phosphatase [Roche,

Cat#11093274910; RRID:AB_514497]) at room temperature for 4 h. The embryos were then

washed four times in 1x MAB for 10 min before leaving them in 1x MAB overnight at 4ºC.

1112

On the final day of the WMISH protocol, the embryos were washed another three times in 1x

MAB for 20 min and equilibrated to working conditions of alkaline phosphatase (AP) for a total

of 10 min by submerging embryos twice into AP buffer (50 mM $MgCl_2$, 100 mM NaCl, 100 mM

Tris-HCl pH 9.5 and 1% (v/v) Tween-20). At this stage, the embryos were transferred to 5-ml

glass vials for monitoring the progression of the AP-catalyzed colorimetric reaction. Any

residual AP buffer was discarded before adding 700 µl staining solution (AP buffer, 340 µg/ml

nitro-blue tetrazolium chloride [Roche, Cat#11383213001] and 175 µg/ml 5-bromo-4-chloro-3'-

indolyphosphate [Roche, Cat#11383221001]). The colorimetric reaction was developed at room temperature in the dark. Once the staining was clear and intense enough, the color reaction was stopped by two washes in 1x MAB. To stabilize and preserve morphological features, the embryos were fixed with Bouin's fixative without picric acid (9% formaldehyde and 5% glacial acetic acid [Fisher Scientific, Cat#10171460]) at room temperature for 30 min. Next, the embryos were washed twice in 70% ethanol/PBT to remove the fixative and residual chromogens. After rehydration to PBT in two steps, the embryos were treated with weak Curis solution (1% (v/v) hydrogen peroxide [Merck, Cat#1072090500], 0.5x SSC and 5% formamide) at 4°C in the dark overnight. Finally, the embryos were washed twice in PBS before imaging them in PBS on a thick agarose dish by light microscopy.

Processing of External Datasets

High-time (30-min intervals) resolution of total and poly(A) RNA-Seq (GSE65785) was processed as reported in the original publication (Owens et al., 2016). In addition, intron read counts were corrected by spike RNA-derived normalization factors. For visualization normalized exon and intron counts were scaled to the maximal count detected across the time course and fitted using cubic smoothing splines from 0 to 23.5 hpf: smooth.spline(1:48, x, spar=0.6). Other RNA-Seq (GSE81458) and ChIP-Seq (GSE67974, GSE30146, GSE53654 and GSE72657) were processed as described in detail above except for H3K4me3 and H3K36me3 whose enriched regions were detected as follows: findPeaks -style histone - fragLength 175 -inputFragLength 175 -fdr 0.001 -gsize 1.435e9 -F 2 -C 1 -region -size 350 - minDist 500. Thus, we detected significant regions of histone modifications (-style histone) of at least the lengths of two DNA fragments (-size 350) and being separated by at least 500 bp from each other.

Generation of Plots and Heatmaps

Genomic snapshots were generated with the IGV genome browser v2.4-rc6 (Robinson et al., 2011). All plots and heatmaps were generated using R v3.5.1 (http://cran.r-project.org/). The following add-on R and Bioconductor packages were used for sorting and graphical visualization of data: alluvial v0.1-2 (Michał Bojanowsk), beeswarm v0.2.3 (Aron Eklund), circlize v0.4.5 (Gu et al., 2014), complexHeatmap v1.20.0 (Gu et al., 2016a), dplyr v0.7.8, ggplot2 v3.1.0 (Wickham, 2016), gplots v3.0.1 (Gregory Warnes and colleagues), GenomicFeatures v1.34.1 (Lawrence et al., 2013), GenomicRanges v1.38.0 (Lawrence et al., 2013), HilbertCurve v1.12.0 (Gu et al., 2016b), limma v3.38.2 (Ritchie et al., 2015), rtracklayer 1.42.1 (Lawrence et al., 2009) and seriation v1.2-3 (Hahsler et al., 2008).

QUANTIFICATION AND STATISTICAL ANALYSIS

No statistical method was used for determining sample size; rather, we followed the literature to select the appropriate sample size. The experiments were not randomized. Due to the nature of experiments, the authors were not blinded to group allocation during data collection and

analysis. Only viable embryos were included in the analysis. Frequencies of shown morphological phenotypes and WMISH patterns are included in every image. The significance of over-represented GO terms was based on hypergeometric tests. Significances of non-normally distributed data points (gene features) across ZGA were calculated using paired Wilcoxon rank-sum tests (alternative hypothesis 'less'). The effect size (r_{effect}) was estimated from the standard normal deviate of the Wilcoxon p-value (p) as previously described (Rosenthal, 1991), $r_{effect}=Z/sqrt(N)$, where $Z=qnorm(1-p/2)$ is the standardized Z-score and N is the number of observations.

For RNA-Seq, biological triplicates were used to account for transcriptional variability between clutches. Each LOF experiment has its own control embryos collected in parallel from the same mothers: exp. #1 (α-amanitin), uninjected embryos; exp. #2 (BMP or Nodal LOF), DMSO-treated embryos; exp. #3 (Wnt LOF), uninjected embryos; exp. #4 (mPou5f3/Sox3 LOF), uninjected embryos; exp. #5 (mVegT LOF), uninjected embryos; exp. #6 (single and combinatorial LOFs of Wnt, Nodal and BMP), DMSO-treated embryos; exp. #7 (combinatorial LOF of 4 zygotic T-box TFs), control MO-injected embryos. The gene expressions of control MO-injected embryos of exp. #2 and #5 were normalized to their corresponding uninjected embryos. The mean of these normalizations and conservative FDR estimations (i.e., higher FDR of the two likelihood ratio tests) were used for the comparison with LOF conditions. RNA-Seq libraries from each experiment were generated simultaneously to mitigate any batch effects. The FDR was controlled for multiple comparisons according to the Benjamini-Hochberg procedure. The exact computational implementation of differential expression analysis is outlined on GitHub (see below).

DATA AND SOFTWARE AVAILABILITY

Sequencing reads (FASTQ files) and raw RNA-Seq read counts reported in this paper are available in the GEO database (www.ncbi.nlm.nih.gov/geo) under the accession numbers GSE113186 and GSE122551. All analyses were performed in R v3.5.1 (Bioconductor v3.8), Perl v5.18.2 (https://www.perl.org) and Python v2.7.12 (http://www.python.org) as detailed above. The R code, genome annotation, intermediate datasets and graphs are available on GitHub at https://github.com/gegentsch/SpatioTemporalControlZGA. Original datasets are also available on Mendeley Data at http://dx.doi.org/10.17632/jn466b4n8v.1.

SUPPLEMENTAL REFERENCES

Benson, G. (1999). Tandem repeats finder: a program to analyze DNA sequences. Nucleic Acids Res. *27*, 573–580.

Blitz, I.L., Paraiso, K.D., Patrushev, I., Chiu, W.T.Y., Cho, K.W.Y., and Gilchrist, M.J. (2017). A catalog of Xenopus tropicalis transcription factors and their regional expression in the early gastrula stage embryo. Dev. Biol. *426*, 409–417.

Carlson, M., Falcon, S., Pages, H., and Li, N. (2007). Bioconductor - GO.db.

Chiu, W.T., Le, R.C., Blitz, I.L., Fish, M.B., Li, Y., Biesinger, J., Xie, X., and Cho, K.W.Y. (2014). Genome-wide view of TGFβ/Foxh1 regulation of the early mesendoderm program. Development *141*, 1–114.

Collart, C., Owens, N.D.L., Bhaw-Rosun, L., Cooper, B., De Domenico, E., Patrushev, I., Sesay, A.K., Smith, J.N., Smith, J.C., and Gilchrist, M.J. (2014). High-resolution analysis of gene activity during the Xenopus mid-blastula transition. Development *141*, 1927–1939.

Conesa, A., Götz, S., García-Gómez, J.M., Terol, J., Talón, M., and Robles, M. (2005). Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. Bioinformatics *21*, 3674–3676.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*, 15–21.

Falcon, S., and Gentleman, R. (2007). Using GOstats to test gene lists for GO term association. Bioinformatics *23*, 257–258.

Fukuda, M., Takahashi, S., Haramoto, Y., Onuma, Y., Kim, Y.-J., Yeo, C.-Y., Ishiura, S., and Asashima, M. (2010). Zygotic VegT is required for Xenopus paraxial mesoderm formation and is regulated by Nodal signaling and Eomesodermin. Int. J. Dev. Biol. *54*, 81–92.

Gay, L., Karfilis, K.V., Miller, M.R., Doe, C.Q., and Stankunas, K. (2014). Applying thiouracil tagging to mouse transcriptome analysis. Nat. Protoc. *9*, 410–420.

Gentsch, G.E., and Smith, J.C. (2017). Efficient Preparation of High-Complexity ChIP-Seq Profiles from Early Xenopus Embryos. Methods Mol. Biol. *1507*, 23–42.

Gentsch, G.E., Owens, N.D.L., Martin, S.R., Piccinelli, P., Faial, T., Trotter, M.W.B., Gilchrist, M.J., and Smith, J.C. (2013). In vivo T-box transcription factor profiling reveals joint regulation of embryonic neuromesodermal bipotency. Cell Rep. *4*, 1185–1196.

Gentsch, G.E., Patrushev, I., and Smith, J.C. (2015). Genome-wide snapshot of chromatin regulators and states in Xenopus embryos by ChIP-Seq. J. Vis. Exp. *96*, e52535.

Gentsch, G.E., Spruce, T., Monteiro, R.S., Owens, N.D.L., Martin, S.R., and Smith, J.C. (2018a). Innate Immune Response and Off-Target Mis-splicing Are Common Morpholino-Induced Side Effects in Xenopus. Dev. Cell *44*, 597–610.e10.

Gentsch, G.E., Spruce, T., Owens, N.D.L., and Smith, J.C. (2018b). The role of maternal pioneer factors in predefining first zygotic responses to inductive signals. bioRxiv 306803. DOI: https://doi.org/10.1101/306803

Gu, Z. Eils, R., and Schlesner, M. (2016a). Complex heatmaps reveal patterns and correlations in multidimensional genomic data. Bioinformatics *32*, 2847-2849.

Gu, Z., Eils, R., and Schlesner, M. (2016b). HilbertCurve: an R/Bioconductor package for high-resolution visualization of genomic data. Bioinformatics *32*, 2372–2374.

Gu, Z., Gu, L., Eils, R., Schlesner, M., and Brors, B. (2014). circlize Implements and enhances circular visualization in R. Bioinformatics *30*, 2811–2812.

Hahsler, M., Hornik, K., and Buchta, C. (2008). Getting Things in Order: An Introduction to the R Package seriation. J. Stat. Softw. *25*, 1–34.

Heasman, J., Kofron, M., and Wylie, C. (2000). βCatenin Signaling Activity Dissected in the Early Xenopus Embryo: A Novel Antisense Approach. Dev. Biol. *222*, 124–134.

Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre, C.,
Singh, H., and Glass, C.K. (2010). Simple combinations of lineage-determining transcription
factors prime cis-regulatory elements required for macrophage and B cell identities. Mol. Cell
38, 576–589.

Hontelez, S., van Kruijsbergen, I., Georgiou, G., van Heeringen, S.J., Bogdanović, O., Lister,
R., and Veenstra, G.J.C. (2015). Embryonic transcription is controlled by maternally defined
chromatin state. Nat. Commun. *6*, 10148.

Khokha, M., Chung, C., Bustamante, E., Gaw, L., Trott, K., Yeh, J., Lim, N., Lin, J., Taverner,
N., Amaya, E., et al. (2002). Techniques and probes for the study of Xenopus tropicalis
development. Dev. Dyn. *225*, 499–510.

Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat.
Methods *9*, 357–359.

Lawrence, M., Gentleman, R., and Carey, V. (2009). rtracklayer: an R package for interfacing
with genome browsers. Bioinformatics *25*, 1841-1842.

Lawrence, M., Huber, W., Pagès, H., Aboyoun, P., Carlson, M., Gentleman, R., Morgan, M.,
and Carey, V. (2013). Software for Computing and Annotating Genomic Ranges. PLoS
Comput. Biol. *9*, e1003118.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,
Durbin, R., 1000 Genome Project Data Processing Subgroup (2009). The Sequence
Alignment/Map format and SAMtools. Bioinformatics *25*, 2078–2079.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol. *15*, 550.

Monsoro-Burq, A.H. (2007). A Rapid Protocol for Whole-Mount In Situ Hybridization on
Xenopus Embryos. Cold Spring Harb. Protoc. *2007*, pdb.prot4809.

Morgan, M., Falcon, S., and Gentleman, R. (2017). *GSEABase: Gene set enrichment data
structures and methods*.

Nakamura, Y., de Paiva Alves, E., Veenstra, G.J.C., and Hoppler, S. (2016). Tissue- and
stage-specific Wnt target gene expression is controlled subsequent to β-catenin recruitment
to cis-regulatory modules. Development *143*, 1914–1925.

Nieuwkoop, P., and Faber, J. (1994). Normal table of Xenopus laevis (Daudin): a systematical
and chronological survey of the development from the fertilized egg till the end of
metamorphosis. Garland.

Owens, N.D.L., Blitz, I.L., Lane, M.A., Patrushev, I., Overton, J.D., Gilchrist, M.J., Cho,
K.W.Y., and Khokha, M.K. (2016). Measuring Absolute RNA Copy Numbers at High Temporal
Resolution Reveals Transcriptome Kinetics in Development. Cell Rep. *14*, 632–647.

Pertea, M., Kim, D., Pertea, G.M., Leek, J.T., and Salzberg, S.L. (2016). Transcript-level
expression analysis of RNA-seq experiments with HISAT, StringTie and Ballgown. Nat.
Protoc. *11*, 1650–1667.

Quinlan, A.R., and Hall, I.M. (2010). BEDTools: a flexible suite of utilities for comparing
genomic features. Bioinformatics *26*, 841–842.

Rana, A., Collart, C., Gilchrist, M., and Smith, J. (2006). Defining synphenotype groups in
Xenopus tropicalis by use of antisense morpholino oligonucleotides. PLoS Genet. *2*, e193.

Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W., and Smyth, G.K. (2015). limma
powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic

Acids Res. *43*, e47.

Robinson, J.T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E.S., Getz, G., and
Mesirov, J.P. (2011). Integrative genomics viewer. Nat. Biotechnol. *29*, 24–26.

Rosenthal, R. (1991). Meta-Analytic Procedures for Social Research. SAGE Publishing.

Sive, H., Grainger, R., and Harland, R. (2000). Early development of Xenopus laevis: A
laboratory manual. Cold Spring Harbor Laboratory Press.

Smit, A., Hubley, R., and Green, P. RepeatMasker Open-4.0. 2013-2015.

Smith, J., Price, B., Green, J., Weigel, D., and Herrmann, B. (1991). Expression of a Xenopus
homolog of Brachyury (T) is an immediate-early response to mesoderm induction. Cell *67*,
79–87.

Wang, L., Wang, S., and Li, W. (2012). RSeQC: quality control of RNA-seq experiments.
Bioinformatics *28*, 2184–2185.

Wickham, H. (2016). ggplot2: Elegant Graphics for Data Analysis. Springer.

Yoon, S.-J., Wills, A.E., Chuong, E., Gupta, R., and Baker, J.C. (2011). HEB and E2A
function as SMAD/FOXH1 cofactors. Genes Dev. *25*, 1654–1661.

Zhu, Q., Fisher, S.A., Shallcross, J., and Kim, J. (2016). VERSE: a versatile and efficient RNA-
Seq read counting tool. bioRxiv 053306. DOI: https://doi.org/10.1101/053306