In vivo microscopy reveals the impact of *Pseudomonas aeruginosa*

social interactions on host colonization

Chiara Rezzoagli[1,2*], Elisa T. Granato[3], Rolf Kümmerli[1,2*]

[1]Department of Plant and Microbial Biology, University of Zurich, Zurich,

Switzerland

[2]Department of Quantitative Biomedicine, University of Zurich, Zurich,

Switzerland

[3]Department of Zoology, University of Oxford, Oxford, United Kingdom

Running title: Bacterial social dynamics and host colonization

* Corresponding authors:

Chiara Rezzoagli or Rolf Kümmerli, Department of Quantitative Biomedicine,

University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland.

Email: chiara.rezzoagli@uzh.ch (CR), rolf.kuemmerli@uzh.ch (RK).

Phone: +41 44 635 48 01.

Conflict of interest: The authors declare no conflict of interest.

Abstract

Pathogenic bacteria engage in social interactions to colonize hosts, which include quorum-sensing-mediated communication and the secretion of virulence factors that can be shared as "public goods" between individuals. While in-vitro studies demonstrated that cooperative individuals can be displaced by "cheating" mutants freeriding on social acts, we know little about social interactions in infections. Here, we developed a live imaging system to track virulence factor expression and social strain interactions in the human pathogen *Pseudomonas aeruginosa* colonizing the gut of *Caenorhabditis elegans*. We found that shareable siderophores and quorum-sensing systems are expressed during infections, affect host gut colonization, and benefit non-producers. However, non-producers were unable to cheat and outcompete producers. Our results indicate that the limited success of cheats is due to the spatial segregation of strains within the gut, the down-regulation of virulence factors over the course of the infection, and the fact that each virulence factor examined is important but not essential for host colonization. Our results shed new light on bacterial social interactions in infections and reveal potential limits of therapeutic approaches that aim to capitalize on social dynamics between strains for infection control.

Introduction

During infections, pathogenic bacteria secrete a wide range of extracellular virulence factors to colonize and grow inside the host (Rahme *et al.*, 1995; Wu *et al.*, 2008). Secreted molecules include siderophores for iron scavenging, signaling molecules for quorum sensing (QS), toxins to attack host cells, and matrix compounds for biofilm formation (Diggle *et al.*, 2007; Flemming *et al.*, 2016; Henkel *et al.*, 2010; Granato *et al.*, 2016). In-vitro studies have shown that extracellular virulence factors can be shared as "public goods" between cells, and thereby benefit individuals other than the producing cell (Köhler *et al.*, 2009; Raymond *et al.*, 2012; Harrison, 2013). There has been enormous interest in understanding how this form of bacterial cooperation can be evolutionarily stable since secreted public goods can be exploited by non-cooperative mutants called "cheats", which do not engage in cooperation yet still benefit from the molecules produced by others (West *et al.*, 2007; Ghoul *et al.*, 2014; Özkaya *et al.*, 2018).

There is increasing evidence that social interactions and cooperator-cheat dynamics might also matter within hosts (Buckling and Brockhurst, 2008; Harrison, 2013; Leggett *et al.*, 2014). For instance, in controlled infection experiments engineered non-producers, deficient for the production of specific virulence factors, could outcompete producers and thereby reduce virulence (Harrison *et al.*, 2006; Rumbaugh *et al.*, 2009, 2012; Pollitt *et al.*, 2014), but in other cases the success of non-producers was compromised (Zhou *et al.*, 2014; Harrison *et al.*, 2017). Other studies have followed chronic human infections within patients over time and reported that virulence-factor-negative

mutants emerge and spread, with the mutational patterns suggesting

cooperator-cheat dynamics (Köhler *et al.*, 2009; Andersen *et al.*, 2015, 2018).

These findings spurred ideas of how social interactions within hosts could be

manipulated for therapeutic purposes (Brown *et al.*, 2009; Allen *et al.*, 2014;

Leggett *et al.*, 2014). Suggested approaches include: inducing cooperator-

cheat dynamics to steer infections towards lower virulence (Köhler *et al.*, 2009;

Granato *et al.*, 2018); introducing less virulent strains with medically beneficial

alleles into established infections (Brown *et al.*, 2009); and targeting secreted

virulence factors to control infections and constrain the evolution of resistance

(André and Godelle, 2005; Clatworthy *et al.*, 2007; Rasko and Sperandio,

2010; Pepper, 2012; Allen *et al.*, 2014; Rezzoagli *et al.*, 2018).

However, all these approaches explicitly rely on the assumption that the social

traits of interest are: (i) expressed inside hosts; (ii) important for host

colonization; (iii) exploitable; and (iv) induce cooperator-cheat dynamics as

observed in vitro (Harrison, 2013) – assumptions that have not yet been tested

in real time inside living hosts. Here, we explicitly test the importance of

bacterial social interactions within hosts by using in-vivo fluorescence

microscopy to monitor bacterial virulence factor production, host colonization

and strain interactions, using the opportunistic pathogen *Pseudomonas*

aeruginosa and its nematode model host *Caenorhabditis elegans* (Tan and

Ausubel, 2000; Ewbank, 2002; Papaioannou *et al.*, 2013)*.

C. elegans naturally preys on bacteria (Félix and Braendle, 2010). While most

bacteria are killed during ingestion, a small fraction of cells survives (Portal-

Celhay *et al.*, 2012), which can, in the case of pathogenic bacteria, establish

an infection in the gut (Tan *et al.*, 1999). *P. aeruginosa* deploys an arsenal of

virulence factors that facilitate successful host colonization (Nadal Jimenez *et*

al., 2012). For example, the two siderophores pyoverdine and pyochelin

scavenge host-bound iron during acute infections to enable pathogen growth

(Meyer *et al.*, 1996; Takase *et al.*, 2000; Cornelis and Dingemans, 2013;

Granato *et al.*, 2016). *P. aeruginosa* further secretes the protease elastase,

the toxin pyocyanin, and rhamnolipid biosurfactants to attack host tissue

(Smith and Iglewski, 2003; Alibaud *et al.*, 2008; Köhler *et al.*, 2009).

Production of these latter virulence factors only occurs at high cell densities

and is controlled by the Las and the Rhl QS-systems (Lee and Zhang, 2015).

Because both QS-regulons and siderophores were shown to be involved in *C.*

elegans killing (Tan *et al.*, 1999; Zaborin *et al.*, 2009; Kirienko *et al.*, 2013;

Cezairliyan *et al.*, 2013; Zhu *et al.*, 2015), we used them as focal traits for our

study.

To tackle our questions, we first conducted experiments with fluorescently

tagged *P. aeruginosa* bacteria (PAO1) to follow infection dynamics, from the

first uptake through feeding up to a progressed state of gut infection. We then

constructed promoter-reporter fusions for genes involved in the synthesis of

the two siderophores (pyoverdine and pyochelin) and the two QS-regulators

(LasR and RhlR) to track in vivo virulence factor gene expression during host

colonization. Subsequently, we used mutant strains deficient for virulence

factors to determine whether they show compromised colonization abilities.

Finally, we followed mixed infections of wildtype and mutants over time to

determine the extent of strain co-localization in the gut, and to test whether

secreted virulence factors are indeed exploitable by non-producers in the host.

Material and methods

Strain and bacterial growth conditions

Bacterial strains, primers and plasmids used in this study are listed in the

Supplementary Tables S1-S3. Details on strain construction are given in the

Supplementary Methods. For all experiments, overnight cultures were grown

in 8 ml Lysogeny broth (LB) in 50 ml tubes, incubated at 37°C, 220 rpm for 18

hours, washed with 0.8% NaCl solution and adjusted them to OD_{600} = 1.

Nematode Growth Media (NGM) plates [0.25% Peptone, 50 mM NaCl, 25mM

$[PO_4^-]$, 5 µg/ml Cholesterol, 1mM $CaCl_2$, 1mM $MgSO_4$ supplemented with

1.5% agar, 6 cm diameter] were seeded with 50 µl of bacterial culture and

incubated at 25°C for 24 hours. All *P. aeruginosa* strains used in this study

showed equal growth on NGM exposure plates (Supplementary Figure S1.

Peptone was purchased from BD Biosciences, Switzerland, all other

chemicals from Sigma Aldrich, Switzerland).

Nematode culture

We used the temperature-sensitive, reproductively sterile *C. elegans* strain

JK509 (glp-1(q231) III). This strain reproduces at 16°C, but does not develop

gonads and is therefore sterile at 25°C. Worms were maintained fertile at 16°C

on High Growth Media (HGM) agar plates (2% Peptone, 50 mM NaCl, 25mM

$[PO_4^-]$, 20 µg/ml Cholesterol, 1mM $CaCl_2$, 1mM $MgSO_4$) seeded with the

standard food source *E. coli* strain OP50 (Stiernagle, 2006). For age

synchronization, plates were washed with sterile distilled water and adult

worms were killed with hypochlorite-sodium hydroxide solution to isolate eggs

(Stiernagle, 2006). Eggs were placed in M9 buffer (20 mM KH_2PO_4, 40 mM

Na_2HPO_4, 80 mM NaCl, 1 mM $MgSO_4$) and incubated at 16°C for 16-18 hours

to hatch. Then, L1 larvae were transferred to OP50-seeded HGM plates and

incubated at 25°C for 28 hours to reach L4 developmental stage. Worms and

OP50 bacteria were provided by the Caenorhabditis Genetic Center (CGC),

which is supported by the National Institutes of Health - Office of Research

Infrastructure Programs (P40 OD010440).

C. elegans **infection protocol**

Synchronized L4 worms were washed from HGM plates with M9 buffer + 50

µg/ml kanamycin (M9-Kan), and washed three times with M9-Kan for worm

surface-disinfection. Viable worms were further separated from any debris by

sucrose flotation (Portman, 2006) and rinsed three time in M9 buffer to remove

sucrose. The worm handling protocol for the main experiments is depicted in

Figure 1A. Specifically, approximately 200 worms were moved to seeded

NGM plates and incubated for 24 hours at 25°C. After this period of exposure

to pathogens, infected worms were extensively washed with M9 buffer + 50

µg/ml chloramphenicol (M9-Cm) followed by M9 buffer, subsequently

transferred to individual wells of a 6-well plate filled with sterile M9 buffer + 5

µg/ml cholesterol (M9+Ch Buffer) where they were kept up to 48 hours post

exposure (hpe) and imaged after 0, 6 and 30 hpe.

Nematode survival assay

Our goal was to observe infections inside living hosts. To verify that worms

stayed alive during the experiment (up to 48 hpe), we tracked the survival of

infected population (50-90 worms) in M9-Ch buffer. Worms were observed for

motility at 0, 24 and 48 hpe, by prodding them with a platinum wire. Worms

were considered dead when they no longer responded to touches. Each

bacterial strain was tested in three replicates and three independent

experiments were carried out. We used *E. coli* OP50 as a negative control for

killing. During this observation period, worms experienced only negligible

killing by the colonizing bacteria, and we found no significant difference in

killing between the non-pathogenic *E. coli* food strain and the *P. aeruginosa*

strains (Supplementary Figure S2).

Microscopy setup and imaging

For microscopy, we picked individual worms from the M9+Ch buffer and

paralyzed them with 25 mM sodium azide before transferring them to an 18-

well μ-slide (Ibidi, Germany). All experiments were carried out at the Center for

Microscope and Image Analysis of the University Zürich (ZMB). For the

colonization experiment, images were acquired on a Leica LX inverted

widefield light microscope system with Leica-TX2 filter cube for mCherry

(emission: 560 ± 40 nm, excitation: 645 ± 75 nm, DM = 595) and Leica-DFC-

350-FX, cooled fluorescence monochrome camera (resolution: 1392 × 1040

pixels) for image recording (16-bit color depth). For gene expression

experiments, microscopy was performed on the InCell Analyzer 2500HS (GE

Healthcare) automated imaging system, using polychroic beam splitter

BGRFR_2 (for mCherry, excitation: 575 ± 25 nm, emission: 607.5 ± 19 nm)

and PCO – sCMOS camera (resolution: 2048 x 2048 pixels, 16-bit).

Image processing and analysis

To extract fluorescence measurements from individual worms, images were

segmented into objects and background, using an automated image

segmentation workflow with *ilastik* software (Sommer *et al.*, 2011). Segmented

images were then imported in *Fiji* (Schindelin *et al.*, 2012) to determine the

fluorescence intensity (as "Raw Integrated Density", i.e. the sum of pixels

values in the selection) and area of each worm. Images obtained from the

InCell microscope entailed 64 frames (8x8 grid) with 10% overlap. These

frames were stitched together using a macro-automated version of the

Stitching plugin in *Fiji* (Preibisch *et al.*, 2009) prior to segmentation and

analysis. To correct for background and host-tissue autofluorescence, we

imaged, at each time point, worms infected with non-fluorescent strains (i.e.

OP50 or PAO1), and used the mean intensity of these control infections to

subtract background fluorescence values from worms infected with fluorescent

strains.

Competition assay in the host

For in-vivo competitions between PAO1-mCherry and PAO1 *ΔpvdDΔpchEF* or

PAO1 *ΔlasR*, overnight monocultures were washed twice with 0.8% NaCl

solution, adjusted to $OD_{600} = 1$ and mixed at a 1:1 ratio. To control for fitness

effects of the mCherry marker, we also competed PAO1-mCherry against the

untagged PAO1. NGM plates were then seeded with 50 µl of mixed culture

and incubated at 25°C for 24 hours. Worms were exposed to the mix for 24

hours and then recovered as previously described. After 6 and 48 hours post-

exposure, individual worms were picked, immobilized with sodium azide and

washed for 5 minutes with M9 + 0.003% NaOCl. Worms were washed twice

with M9 buffer. We then transferred each individual worm to a 1.5 ml screw-

cap micro tube (Sarstedt, Switzerland) containing sterilized glass beads (1 mm

diameter, Sigma Aldrich). Worms were disrupted using a bead-beater

(TissueLyser II, QIAGEN, Germany), shaking at 30 Hz for 1.5 min before

flipping the tubes and shaking for an additional 1.5 min to ensure even

disruption (adapted from Vega and Gore, 2017). Tubes were then centrifuged

at 2000 x g for 2 min, the content was re-suspended in 200 µl of 0.8% NaCl

and plated on two LB 1.2 % agar plates for each sample. Plates were

incubated overnight at 37°C and left at room temperature for another 24 h to

allow the fluorescent marker to fully mature. We then distinguished between

fluorescent and non-fluorescent colonies using a custom built fluorescence

imaging device (*Infinity 3* camera, Lumenera, Canada). We then calculated the

relative fitness of PAO1-mCherry as $\ln(v)=\ln\{[a_{48}\times(1-a_6)]/[a_6\times(1-a_{48})]\}$, where

a_6 and a_{48} are the frequency of PAO1-mCherry at 6 and 48 hours after

recovery, respectively (Ross-Gillespie *et al.*, 2007). Values of $\ln(v)<0$ or

$\ln(v)>0$ indicate whether the frequency of PAO1-mCherry increased ($\ln(v)< 0$)

or decreased ($\ln(v)>0$) relative to its competitor.

Co-localization analysis

To determine the degree of co-localization of two different bacterial strains in

the host, we transferred nematodes to NGM plates seeded with a 1:1 mix of

PAO1-gfp with either PAO1-mCherry, PAO1 *ΔpvdDΔpchEF*–mCherr*y,* or

PAO1 *ΔlasR-mCherry*. After a 24 hours grazing period, we picked single

worms and imaged both mCherry- and GFP channels, using the InCell

Analyzer 2500HS microscope. We used *Fiji* to straighten each worm with the

Straighten plugin (Kocsis *et al.*, 1991), and extracted fluorescence intensity

values in the GFP and mCherry channels for each pixel from tail ($X = 0$) to

head ($X = 1$) of the worm. To ensure that we only measure areas where

bacteria were present, we restricted our analysis to the region of the worm gut,

where bacterial colonization takes place. We then calculated Spearman

correlation coefficients between the fluorescent signals, as a proxy for strain

co-localization using RStudio v. 3.3.0 (R Development Core Team, 2013).

Statistical analysis

All statistical analyses were performed with RStudio. We used Pearson

correlations to test for associations between PAO1-mCherry fluorescence

intensities and (a) recovered bacteria from the gut; and (b) total bacterial load

in mixed infections. We used analysis of variance (ANOVA) to compare

fluorescence values between observation times, strains and for comparisons

to non-fluorescent controls. P-values were corrected for multiple comparisons

using the post-hoc Tukey HSD test. To compare promoter expression data

between PAO1 WT and mutant strains, and to compare relative fitness values

between competitors in the competition assay, we used Welch's two-sample t-

test. To measure co-localization, we calculated the Spearman correlation

coefficient between the intensity of mCherry and GFP signals across the worm

gut, and used ANOVA to test for differences between treatments.

Results

PAO1 colonization dynamics in the *C. elegans* gut

For all infection experiments, we followed the protocol depicted in Figure 1A-C. We first exposed worms to *P. aeruginosa* for 24 hours on NGM plates. Subsequently, worms were removed, washed, and treated with antibiotics to kill external bacteria. We then imaged infected worms under the microscope at different time points and quantified bacterial density and gene expression using fluorescent mCherry markers. We first confirmed that mCherry fluorescence is a suitable proxy for the number of live bacteria in *C. elegans*, by comparing fluorescence intensities in whole worms (Figure 1B) to the number of live bacteria recovered from the worms' gut. Fluorescence intensity values positively correlated with the bacterial load inside the nematodes, both immediately after recovering the worms from the exposure plates and at 6 hours post exposure (hpe; Supplementary Figure S3, Pearson correlation coefficient at 0 hpe: $r = 0.49$, $t_{28} = 3.02$, $p = 0.0053$; at 6 hpe: $r = 0.713$, $t_{23} = 4.88$, $p < 0.0001$). As our goal was to image infections in living hosts, we further confirmed that worms stayed alive during the observation period (Supplementary Figure S2).

When following host colonization by PAO1-mCherry over time, we observed that immediately after removal from the exposure plate, worms carried large amounts of bacteria in their gut (Figure 1D). Subsequently, bacterial load significantly declined when worms were kept in buffer for 6 hours (ANOVA: t_{391} = -8.55, $p < 0.001$) and then remained constant for the next 24 hours (t_{391} =

0.61, p = 0.529). This pattern suggests that a large number of bacteria are

taken up during the feeding phase, followed by the shedding of a high

proportion of cells, leaving behind a fraction of live bacteria that establishes an

infection and colonizes the worm gut.

PAO1 expresses siderophore biosynthesis genes and QS regulators in

the host

We then examined whether genes involved in the synthesis of pyoverdine

(*pvdA*) and pyochelin (*pchEF*), and the genes encoding the QS-regulators

lasR and *rhlR*, are expressed inside hosts. Worms were exposed to four

different PAO1 strains, each carrying a specific promoter-mCherry fusion.

Imaging after the initial uptake phase (0 hpe) revealed that, with the exception

of *pchEF*, all genes were significantly expressed in the host (Figure 2;

ANOVA, comparisons to the non-fluorescent control, for *pvdA*: t_{754} = 4.23, p

<0.001; for *pchEF*: t_{754} = 0.74, p = 0.461; for *lasR*: t_{754} = 2.96, p = 0.003*;* for

rhlR: t_{754} = 10.37, p <0.001*).* Although fluorescence intensity declined over

time (linear model, $F_{1,1795}$ = 48.98, p < 0.001), we observed that apart from

pchEF, all genes were still significantly expressed at 30 hpe (Figure 2;

ANOVA, for *pvdA*: t_{754} = 4.87, p <0.001*;* for *pchEF*: t_{754} = 0.684, p = 0.461; for

lasR: t_{754} = 3.01, p = 0.003; for *rhlR*: t_{754} = 16.68, p <0.001). These results

suggest that pyoverdine and QS-regulated traits seem important for both the

initial uptake and the subsequent colonisation of the host.

Regulatory links between social traits operate inside the host

We know that regulatory links exist between the virulence traits studied here.

While pyoverdine synthesis suppresses pyochelin production under stringent

iron limitation (Dumas *et al.*, 2013), the Las-QS system positively activates the

Rhl-QS system (Lee and Zhang, 2015). To test whether these links operate

inside the host, we measured gene expression of each trait in the negative

background of the co-regulated trait (Figure 3). For *pvdA*, we observed

significant gene expression levels in both the wildtype PAO1 and the

pyochelin-deficient PAO1 Δ*pchEF* strain (Figure 3A), albeit the overall

expression was slightly reduced in PAO1 Δ*pchEF* (t-test, t_{253} = 8.67, p <

0.001). For *pchEF*, expression patterns confirm the suppressive nature of

pyoverdine: the pyochelin synthesis gene was not expressed in the wildtype

but significantly upregulated in the pyoverdine-deficient PAO1 Δ*pvdD* strain

(Figure 3B; t_{296} = -19.68, p < 0.001). For *lasR*, we found that gene expression

was not significantly different in wildtype PAO1 compared to the Rhl-negative

mutant PAO1 Δ*rhlR*, confirming that the Las-QS system is at the top of the

hierarchy and not influenced by the Rhl-system (Figure 3C; t_{211} = -1.50, p =

0.136). Conversely, the expression of *rhlR* was strongly dependent on a

functional Las-system, and therefore only expressed in PAO1, but repressed

in the Las-negative mutant PAO1 Δ*lasR* (Figure 3D; t_{156} = 19.04, p < 0.001).

These results show that (i) iron-limitation is strong in *C. elegans* as PAO1

primarily invests in the more potent siderophore pyoverdine; (ii) pyochelin can

have compensatory effects when pyoverdine is lacking; and (iii) the loss of the

Las-system leads to the concomitant collapse of the Rhl-system.

Virulence-factor-negative mutants show trait-specific deficiencies in host colonization

To examine whether the ability to produce shared virulence factors is important for initial bacterial uptake and host colonization, we exposed *C. elegans* to five isogenic mutants of PAO1-mCherry, either impaired in the production of pyoverdine (Δ*pvdD*), pyochelin (Δ*pchEF*), both siderophores (Δ*pvdD*Δ*pchEF*), the QS receptor LasR (Δ*lasR*), or the QS receptor RhlR (Δ*rhlR*). After the feeding phase, the bacterial load of the wildtype and all three siderophores mutants were equally abundant inside hosts, whereas bacterial load was significantly reduced for the two QS-mutants compared to the wildtype (Figure 4A; ANOVA, significant variation among strains $F_{5,736} = 10.50$, $p < 0.001$; post-hoc Tukey test for multiple comparisons: $p > 0.05$ for all siderophore mutants, $p = 0.021$ for PAO1 Δ*lasR*, $p < 0.001$ for PAO1 Δ*rhlR*).

As previously described for PAO1 colonization (Figure 1D), we observed that the bacterial load of all strains declined at 6 hpe (Supplementary Figure S4) and 30 hpe (Figure 4B) following worm removal from the exposure plates. This decline was significantly more pronounced for the double-siderophore knockout PAO1 Δ*pvdD*Δ*pchEF* than for the wildtype (Figure 4B; ANOVA, post-hoc Tukey test $p < 0.001$). In contrast, mutants deficient in pyochelin (PAO1 Δ*pchEF*) and RhlR (PAO1 Δ*rhlR*) production showed a significantly higher ability to remain in the host than the wildtype (Figure 4B; ANOVA, post-hoc Tukey test $p <0.001$ for both strains). Taken together, our findings suggest that the two siderophores can complement each other, and that only the

siderophore double mutant and the LasR-deficient strain have an overall

disadvantage in colonizing worms.

Mixed communities are formed inside hosts, but exploitation of social

traits is constrained

Given our findings on colonization deficiencies, we reasoned that the

siderophore-double mutant (PAO1 $\Delta pvdD\Delta pchEF$) and the Las-deficient

mutant (PAO1 $\Delta lasR$) could act as cheats and benefit from the exploitation of

virulence factors produced by the wildtype in mixed infections. To test this

hypothesis, we first competed the PAO1-mcherry strain against the untagged

wildtype in the host, and found that the mCherry tag had a small negative

effect on PAO1 fitness (Figure 5A; one sample t-test, t_{24} = -4.12, p < 0.001).

We then competed PAO1-mcherry against the two putative cheats and found

that neither of them could gain a significant fitness advantage over the

wildtype, but also did not lose out (Figure 5A; ANOVA, $F_{2,70}$ = 0.517, p =

0.598). These results indicate that virulence-factor-negative mutants, initially

compromised in host colonization, can indeed benefit from the presence of the

wildtype producer, but not to an extent that would allow them to increase in

frequency and displace producers.

Since the wildtype alone was able to maintain higher bacterial loads in the

worms compared to the two mutants (Figure 4B), we hypothesized that

frequencies of the wildtype in mixed infections should positively correlate with

the total bacterial load in the gut. Consistent with this prediction, we found that

at 6 hpe, the frequency of PAO1-mCherry positively correlated with bacterial

load in mixed infections with the two non-producers, but not in the control

mixed infections with the untagged wildtype (Pearson correlation coefficient, r

= 0.54, t_{17} = 2.67, p = 0.016 (PAO1 *ΔlasR),* r = 0.40, t_{17} = 1.77, p = 0.031

(PAO1 *ΔpvdDΔpchEF*); r = 0.12, t_{17} = 0.47, p = 0.639 (PAO1)). These

correlations disappeared at the later colonization stage (48 hpe; Pearson

correlation coefficient r < 0 for all strains).

Spatial distribution and strain co-localization varies substantially across

host individuals

In-vitro studies have shown that spatial proximity of cells is crucial for efficient

compound sharing (Van Gestel *et al.*, 2014; Weigert and Kümmerli, 2017). We

thus assessed the spatial proximity of co-infecting strains in vivo by exposing

worms to a mix of two differentially labeled (either mCherry or GFP) strains

(Figure 6). We found that all worms were colonized by both strains, but their

co-localization varied greatly across individuals. Specifically, correlation

between mCherry and GFP fluorescence from tail to head ranged from almost

perfect co-localization in some worms (Figure 6A; r = 1) to near complete

spatial segregation in others (Figure 6B; r < 0). Similar co-localization patterns

emerged for all three strain combinations tested, highlighting that the type of

competitor did not influence the degree of strain co-localization in the host gut

(Figure 6C; ANOVA, $F_{2,102}$ = 2.17, p =0.119). Important to note is that we

calculated co-localization based on a 2-dimensional projection of a 3-

dimensional organ (i.e. the gut), which might overestimate the level of co-

localization along the z-axis.

Discussion

We developed a live imaging system that allows us to track host colonization by pathogenic bacteria (*P. aeruginosa*) and their expression of virulence factors inside hosts (*C. elegans*). We used this system to focus on the role of secreted virulence factors, which can be shared as public goods between bacterial cells, and examined competitive dynamics between virulence factor producing and non-producing strains in the host. We found that siderophores (pyoverdine and pyochelin) and the Las and Rhl QS-systems: (i) are expressed inside the host, (ii) affect the ability to colonize and reside within the nematodes; (iii) allow non-producers to benefit from virulence factors secreted by producers in mixed infections; but (iv) do not allow non-producers to cheat and outcompete producers. Our results have implications for both the understanding of bacterial social interactions within hosts, and therapeutic approaches that aim to manipulate social dynamics between strains for infection control.

Numerous in-vitro studies have shown that bacterial cooperation can be exploited by cheating mutants that no longer express the social trait, but benefit from the cooperative acts performed by others (Griffin *et al.*, 2004; Diggle *et al.*, 2007; Ross-Gillespie *et al.*, 2007; Sandoz *et al.*, 2007; Kümmerli *et al.*, 2009a; Popat *et al.*, 2012; Ghoul *et al.*, 2014; O'Brien *et al.*, 2017; Özkaya *et al.*, 2018). These findings contrast with our observations that the spread of non-producers was constrained within infections. There are multiple ways to explain this constraint. First, increased spatial structure is known to limit bacterial dispersal and the diffusion of secreted metabolites (Kümmerli *et*

al., 2009a). As a consequence, metabolite sharing becomes more local, which restricts non-producers to access the public good. In infection systems such as ours, where bacteria attach to host tissue, spatial structure is likely high (as shown in Fig. 6), and public good sharing might be reduced. Our results match those by Zhou *et al* (2014), who showed that QS-mutants of *Bacillus thuringiensis* infecting caterpillars could not exploit metabolites from producers, presumably because they were spatially separated in the host. Moreover, Vega and Gore (2017) showed that the likelihood of two competiting strains to end up in the same host is drastically reduced when bacteria occur at low frequency in the environment, leading to a complete spatial segretion of bacterial strains.

Second, negative frequency-dependent selection could explain why the spread of virulence factor negative mutants is constrained (Ross-Gillespie *et al.*, 2007). This scenario predicts that cheaters only experience a selective advantage when rare, but not when common. The reasoning is that non-producers can only efficiently exploit public goods when surrounded by many producers. Our competition experiments indeed provide indirect evidence for negative frequency-dependent selection in the nematode gut (Figure 5B). Specifically, we observed that bacterial load was reduced when producers occurred at low frequency early during infection (6 hpe), but not later on (48 hpe). This pattern indicates that rare producers have a selective advantage, increase in relative frequency and restore bacterial load.

Third, the relatively low bacterial density in the gut could further compromise

the ability of non-producers to cheat (Figure 1D, 5B). Low cell density restricts

the sharing and exploitation of secreted compounds (Ross-Gillespie *et al.*,

2009; Scholz and Greenberg, 2015). Mechanisms responsible for the low

bacterial density in the gut (Figure 1D, 5B) could include the peristaltic activity

of the gut, expelling a part of the pathogen population and the host immune

system, killing a fraction of the bacteria (Pukkila-Worley and Ausubel, 2012).

Fourth, our analysis reveals that the expression of pyoverdine and QS-

systems decline over time (Figure 2). This could mean that the costs and

benefits of shared virulence factors are reduced at later stages of the infection,

or that bacteria switch from producing to recycling already secreted public

goods (Imperi *et al.*, 2009; Kümmerli and Brown, 2010). The spread of non-

producers might be hampered in this case because the exploitability of a trait

depends on its expression level (Kümmerli *et al.*, 2009b; Jiricny *et al.*, 2010).

Finally, we show that the regulatory linkage between traits is an important

factor to consider when predicting the putative advantage of non-producers

(Lindsay *et al.*, 2016; dos Santos *et al.*, 2018). For instance, *P. aeruginosa*

pyoverdine-negative-mutants upregulated pyochelin production to compensate

for the lack of their primary siderophore (Figure 3). Thus, if pyoverdine-

negative mutants evolve *de novo*, their spread as cheats could be hampered

because they invest in pyochelin as an alternative siderophore (Ross-Gillespie

et al., 2015). For QS, meanwhile, we observed that the absence of a functional

Las-system resulted in the concomitant collapse of the Rhl-system. Although

lasR mutants could be potent cheats, as they are deficient for multiple social traits, their spread might be hampered because QS-systems also regulate non-social traits, important for individual fitness (Dandekar *et al.*, 2012).

When relating our insights to previous studies, it turns out that earlier work produced mixed results with regard to the question whether siderophore- and QS-deficient mutants can spread within infections. While Harrison *et al.* (2006, 2017; pyoverdine, *P. aeruginosa* in *Galleria mellonella* and ex-vivo infection models) and Zhou *et al.* (2014; QS, *B. thuringiensis* in *Plutella xylostella*) showed that the spread of non-producers is constrained, Rumbaugh *et al.* (2009, 2012; QS; *P. aeruginosa* in mice) and Pollitt *et al.* (2014; QS, *Staphylococcus aureus* in *G. mellonella*) demonstrated cases where non-producers spread to high frequencies in host populations. While the reported results were based on strain frequency counts before and after competition, we here show that information on social trait expression, temporal infection dynamics and physical interactions among strains within hosts are essential to understand whether social traits are important and exploitable in a given system. We thus posit that more such detailed approaches are required to understand the importance of bacterial social interactions across host systems and infection contexts and explain differences between them.

A deeper understanding of bacterial social interactions inside hosts is particularly relevant for novel therapeutic approaches that seek to take advantage of cooperator-cheat dynamics inside hosts to control infections. For instance, it was proposed that strains deficient for virulence factors could be

introduced into established infections (Brown *et al.*, 2009). These strains are

expected to spread because of cheating, thereby reducing the overall

virulence factor availability in the population and the damage to the host. Our

results reveal that virulence-factor-negative strains, although gaining a benefit

from producer strains, are unable to spread in populations. Another

therapeutic approach involves the specific targeting of secreted virulence

factors (André and Godelle, 2005; Allen *et al.*, 2014). This approach is thought

to reduce damage to the host and to compromise resistance evolution

(Pepper, 2012). Resistant mutants, resuming virulence factor production, are

not expected to spread because they would act as cooperators, sharing the

benefit of secreted goods with susceptible strains (Mellbye and Schuster,

2011; Gerdt and Blackwell, 2014; Ross-Gillespie *et al.*, 2014). Our results yet

indicate that such resistant mutants could get local benefits and thus increase

to a certain frequency in the population (Rezzoagli *et al.*, 2018). These

confrontations show that the identification of key parameters driving social

interactions across hosts and infection types is of utmost importance to predict

the success of 'cheat therapies' and anti-virulence strategies targeting

secreted public goods.

Acknowledgments

We thank the Center of Microscopy and Image Analysis (University of Zürich)

for support with image acquisition and advice on image analysis.

Funding

This project has received funding from the Swiss National Science Foundation

(grant no. PP00P3_165835 to RK and no. P2ZHP3_174751 to EG), and the

European Research Council under the grant agreement no. 681295 (to RK).

Competing Interests

537 The authors have no competing interests to declare.

Supplementary Information

540 Supplementary information is available at the journal's website.

References:

Alibaud L, Köhler T, Coudray A, Prigent-Combaret C, Bergeret E, Perrin J, *et*

al. (2008). Pseudomonas aeruginosa virulence genes identified in a

Dictyostelium host model. *Cell Microbiol* **10**: 729–740.

Allen RC, Popat R, Diggle SP, Brown SP. (2014). Targeting virulence: can we

make evolution-proof drugs? *Nat Rev Microbiol* **12**: 300–308.

Andersen SB, Ghoul M, Marvig RL, Lee Z-B, Molin S, Johansen HK, *et al.*

(2018). Privatisation rescues function following loss of cooperation. *bioRxiv*. e-

pub ahead of print, doi: 10.1101/326165.

Andersen SB, Marvig RL, Molin S, Krogh Johansen H, Griffin AS. (2015).

Long-term social dynamics drive loss of function in pathogenic bacteria. *Proc*

Natl Acad Sci **112**: 10756–10761.

André JB, Godelle B. (2005). Multicellular organization in bacteria as a target

for drug therapy. *Ecol Lett* **8**: 800–810.

Brown SP, West S a, Diggle SP, Griffin AS. (2009). Social evolution in micro-

organisms and a Trojan horse approach to medical intervention strategies.

Philos Trans R Soc Lond B Biol Sci **364**: 3157–3168.

Buckling A, Brockhurst MA. (2008). Kin selection and the evolution of

virulence. *Heredity (Edinb)* **100**: 484–488.

Cezairliyan B, Vinayavekhin N, Grenfell-Lee D, Yuen GJ, Saghatelian A,
Ausubel FM. (2013). Identification of Pseudomonas aeruginosa phenazines
that kill Caenorhabditis elegans. *PLoS Pathog* **9**: e1003101.

Clatworthy AE, Pierson E, Hung DT. (2007). Targeting virulence: a new
paradigm for antimicrobial therapy. *Nat Chem Biol* **3**: 541–548.

Cornelis P, Dingemans J. (2013). *Pseudomonas aeruginosa* adapts its iron
uptake strategies in function of the type of infections. *Front Cell Infect*
Microbiol **3**: 1–7.

Dandekar AA, Chugani S, Greenberg PE. (2012). Bacterial Quorum Sensing
and Metabolic Incentives to Cooperate. *Science (80-)* **338**: 264–266.

Diggle SP, Griffin AS, Campbell GS, West S a. (2007). Cooperation and
conflict in quorum-sensing bacterial populations. *Nature* **450**: 411–414.

Dumas Z, Ross-Gillespie A, Kümmerli R. (2013). Switching between
apparently redundant iron-uptake mechanisms benefits bacteria in changeable
environments. *Proc Biol Sci* **280**: 20131055.

Ewbank JJ. (2002). Tackling both sides of the host – pathogen equation with
Caenorhabditis elegans. **4**: 247–256.

Félix M-A, Braendle C. (2010). The natural history of Caenorhabditis elegans.
Curr Biol **20**: R965–R969.

Flemming HC, Wingender J, Szewzyk U, Steinberg P, Rice SA, Kjelleberg S.
(2016). Biofilms: An emergent form of bacterial life. *Nat Rev Microbiol* **14**:
563–575.

Gerdt JP, Blackwell HE. (2014). Competition Studies Confirm Two Major
Barriers That Can Preclude the Spread of Resistance to Quorum-Sensing
Inhibitors in Bacteria. *ACS Chem Biol* **9**: 2291–2299.

Van Gestel J, Weissing FJ, Kuipers OP, Kovács ÁT. (2014). Density of
founder cells affects spatial pattern formation and cooperation in Bacillus
subtilis biofilms. *ISME J* **8**: 2069–2079.

Ghoul M, Griffin AS, West SA. (2014). Toward an evolutionary definition of

cheating. *Evolution (N Y)* **68**: 318–331.

Granato ET, Harrison F, Kümmerli R, Ross-Gillespie A. (2016). Do Bacterial

"Virulence Factors" Always Increase Virulence? A Meta-Analysis of

Pyoverdine Production in Pseudomonas aeruginosa As a Test Case. *Front*

Microbiol **7**: 1952.

Granato ET, Ziegenhain C, Marvig RL, Kümmerli R. (2018). Low spatial

structure and selection against secreted virulence factors attenuates

pathogenicity in Pseudomonas aeruginosa. *ISME J*. e-pub ahead of print, doi:

10.1038/s41396-018-0231-9.

Griffin AS, West SA, Buckling A. (2004). Cooperation and competition in

pathogenic bacteria. *Nature* **430**: 1024–1027.

Harrison F. (2013). Bacterial cooperation in the wild and in the clinic: Are

pathogen social behaviours relevant outside the laboratory? *BioEssays* **35**:

108–112.

Harrison F, Browning LE, Vos M, Buckling A. (2006). Cooperation and

virulence in acute Pseudomonas aeruginosa infections. *BMC Biol*. e-pub

ahead of print, doi: 10.1186/1741-7007-4-21.

Harrison F, McNally A, Da Silva AC, Heeb S, Diggle SP. (2017). Optimised

chronic infection models demonstrate that siderophore 'cheating' in

Pseudomonas aeruginosa is context specific. *ISME J* **11**: 2492–2509.

Henkel JS, Baldwin MR, Barbieri JT. (2010). Toxins from bacteria. *EXS* **100**:

1–29.

Imperi F, Tiburzi F, Visca P. (2009). Molecular basis of pyoverdine

siderophore recycling in Pseudomonas aeruginosa. *Proc Natl Acad Sci* **106**:

20440–20445.

Jiricny N, Diggle SP, West SA, Evans BA, Ballantyne G, Ross-Gillespie A, *et*

al. (2010). Fitness correlates with the extent of cheating in a bacterium. *J Evol*

Biol **23**: 738–747.

Kirienko N V., Kirienko DR, Larkins-Ford J, W??hlby C, Ruvkun G, Ausubel

FM. (2013). Pseudomonas aeruginosa disrupts Caenorhabditis elegans iron

homeostasis, causing a hypoxic response and death. *Cell Host Microbe* **13**:

406–416.

Kocsis E, Trus BL, Steer CJ, Bisher ME, Steven AC. (1991). Image averaging

of flexible fibrous macromolecules: the clathrin triskelion has an elastic

proximal segment. *J Struct Biol* **107**: 6–14.

Köhler T, Buckling A, van Delden C. (2009). Cooperation and virulence of

clinical Pseudomonas aeruginosa populations. *Proc Natl Acad Sci* **106**: 6339–

6344.

Kümmerli R, Brown SP. (2010). Molecular and regulatory properties of a public

good shape the evolution of cooperation. *Proc Natl Acad Sci U S A* **107**:

18921–6.

Kümmerli R, Griffin AS, West S a, Buckling A, Harrison F. (2009a). Viscous

medium promotes cooperation in the pathogenic bacterium *Pseudomonas*

aeruginosa. *Proc Biol Sci* **276**: 3531–3538.

Kümmerli R, Jiricny N, Clarke LS, West SA, Griffin AS. (2009b). Phenotypic

plasticity of a cooperative behaviour in bacteria. *J Evol Biol* **22**: 589–598.

Lee J, Zhang L. (2015). The hierarchy quorum sensing network in

Pseudomonas aeruginosa. *Protein Cell* **6**: 26–41.

Leggett HC, Brown SP, Reece SE. (2014). War and peace□: social

interactions in infections. *Philos Trans R Soc B* **369**: 20130365.

Lindsay RJ, Kershaw MJ, Pawlowska BJ, Talbot NJ, Gudelj I. (2016).

Harbouring public good mutants within a pathogen population can increase

both fitness and virulence. *Elife* **5**: 1–25.

Mellbye B, Schuster M. (2011). The sociomicrobiology of antivirulence drug

resistance: a proof of concept. *MBio* **2**: e00131-11.

Meyer JM, Neely A, Stintzi A, Georges C, Holder IA. (1996). Pyoverdin is

essential for virulence of Pseudomonas aeruginosa. *Infect Immun* **64**: 518–

523.

Nadal Jimenez P, Koch G, Thompson J a., Xavier KB, Cool RH, Quax WJ.

(2012). The Multiple Signaling Systems Regulating Virulence in Pseudomonas

aeruginosa. *Microbiol Mol Biol Rev* **76**: 46–65.

O'Brien S, Luján AM, Paterson S, Cant MA, Buckling A. (2017). Adaptation to
public goods cheats in Pseudomonas aeruginosa. *Proc R Soc B Biol Sci* **284**:
20171089.

Özkaya Ö, Balbontín R, Gordo I, Xavier KB. (2018). Cheating on Cheaters
Stabilizes Cooperation in Pseudomonas aeruginosa. *Curr Biol* **26**: 2070–2080.

Papaioannou E, Utari P, Quax W. (2013). Choosing an Appropriate Infection
Model to Study Quorum Sensing Inhibition in Pseudomonas Infections. *Int J*
Mol Sci **14**: 19309–19340.

Pepper JW. (2012). Drugs that target pathogen public goods are robust
against evolved drug resistance. *Evol Appl* **5**: 757–761.

Pollitt EJG, West SA, Crusz SA, Burton-Chellew MN, Diggle SP. (2014).
Cooperation, quorum sensing, and evolution of virulence in Staphylococcus
aureus. *Infect Immun* **82**: 1045–1051.

Popat R, Crusz S a., Messina M, Williams P, West S a., Diggle SP. (2012).
Quorum-sensing and cheating in bacterial biofilms. *Proc R Soc B Biol Sci* **279**:
4765–4771.

Portal-Celhay C, Bradley ER, Blaser MJ. (2012). Control of intestinal bacterial
proliferation in regulation of lifespan in Caenorhabditis elegans. *BMC Microbiol*
12: 49.

Portman DS. (2006). Profiling C. elegans gene expression with DNA
microarrays. In: Community TC elegan. R (ed). *WormBook: The Online*
Review of C. elegans Biology. WormBook.

Preibisch S, Saalfeld S, Tomancak P. (2009). Globally optimal stitching of tiled
3D microscopic image acquisitions. *Bioinformatics* **25**: 1463–1465.

Pukkila-Worley R, Ausubel FM. (2012). Immune defense mechanisms in the
Caenorhabditis elegans intestinal epithelium. *Curr Opin Immunol* **24**: 3–9.

R Development Core Team. (2013). R: A language and environment for
statistical computing. In: *R Foundation for Statistical Computing.* Vienna,
Austria.

Rahme LG, Stevens EJ, Wolfort SF, Shao J, Tompkins RG, Ausubel FM.

(1995). Common virulence factors for bacterial pathogenicity in plants and

animals. *Science (80-)* **268**: 1899–902.

Rasko DA, Sperandio V. (2010). Anti-virulence strategies to combat bacteria-

mediated disease. *Nat Rev Drug Discov* **9**: 117–128.

Raymond B, West SA, Griffin AS, Bonsall MB. (2012). The dynamics of

cooperative bacterial virulence in the field. *Science (80-)* **337**: 85–88.

Rezzoagli C, Wilson D, Weigert M, Wyder S, Kümmerli R. (2018). Probing the

evolutionary robustness of two repurposed drugs targeting iron uptake in

Pseudomonas aeruginosa. *Evol Med Public Heal* **1**: 246–259.

Ross-Gillespie A, Dumas Z, Kümmerli R. (2015). Evolutionary dynamics of

interlinked public goods traits: An experimental study of siderophore

production in Pseudomonas aeruginosa. *J Evol Biol* **28**: 29–39.

Ross-Gillespie A, Gardner A, Buckling A, West SA, Griffin AS. (2009). Density

dependence and cooperation: theory and a test with bacteria. *Evolution* **63**:

2315–2325.

Ross-Gillespie A, Gardner A, West SA, Griffin AS. (2007). Frequency

dependence and cooperation: theory and a test with bacteria. *Am Nat* **170**:

331–342.

Ross-Gillespie A, Weigert M, Brown SP, Kümmerli R. (2014). Gallium-

mediated siderophore quenching as an evolutionarily robust antibacterial

treatment. *Evol Med Public Heal* **2014**: 18–29.

Rumbaugh KP, Diggle SP, Watters CM, Ross-Gillespie A, Griffin AS, West

SA. (2009). Quorum Sensing and the Social Evolution of Bacterial Virulence.

Curr Biol **19**: 341–345.

Rumbaugh KP, Trivedi U, Watters C, Burton-Chellew MN, Diggle SP, West

SA. (2012). Kin selection, quorum sensing and virulence in pathogenic

bacteria. *Proc R Soc B Biol Sci* **279**: 3584–3588.

Sandoz KM, Mitzimberg SM, Schuster M. (2007). Social cheating in

Pseudomonas aeruginosa quorum sensing. *Proc Natl Acad Sci U S A* **104**:

15876–81.

dos Santos M, Ghoul M, West SA. (2018). Pleiotropy, cooperation and the

social evolution of genetic architecture. *PLoS Biol* **16**: e2006671.

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, *et*

al. (2012). Fiji: an open-source platform for biological-image analysis. *Nat*

Methods **9**: 676–82.

Scholz RL, Greenberg EP. (2015). Sociality in Escherichia coli: Enterochelin Is

a Private Good at Low Cell Density and Can Be Shared at High Cell Density. *J*

Bacteriol **197**: 2122–2128.

Smith RS, Iglewski BH. (2003). *P. aeruginosa* quorum-sensing systems and

virulence. *Curr Opin Microbiol* **6**: 56–60.

Sommer C, Strähle C, Köthe U, Hamprecht FA. (2011). ilastik: Interactive

Learning and Segmentation Toolkit. In: *Eighth IEEE International Symposium*

on Biomedical Imaging (ISBI). Proceedings. pp 230–233.

Stiernagle T. (2006). Maintenance of C. elegans. In: The C.elegans Research

Community (ed). *WormBook: The Online Review of C. elegans Biology.*

Takase H, Nitanai H, Hoshino K, Otani T. (2000). Impact of Siderophore

Production on Pseudomonas aeruginosa Infections in Immunosuppressed

Mice. *Infect Immun* **68**: 1834–1839.

Tan MW, Ausubel FM. (2000). Caenorhabditis elegans: A model genetic host

to study Pseudomonas aeruginosa pathogenesis. *Curr Opin Microbiol* **3**: 29–

34.

Tan MW, Mahajan-Miklos S, Ausubel FM. (1999). Killing of Caenorhabditis

elegans by Pseudomonas aeruginosa used to model mammalian bacterial

pathogenesis. *Proc Natl Acad Sci U S A* **96**: 715–20.

Vega NM, Gore J. (2017). Stochastic assembly produces heterogeneous

communities in the Caenorhabditis elegans intestine. *PLOS Biol* **15**:

e2000633.

Weigert M, Kümmerli R. (2017). The physical boundaries of public goods

cooperation between surface-attached bacterial cells. *Proc R Soc B Biol Sci*

284: 20170631.

West SA, Diggle SP, Buckling A, Gardner A, Griffin AS. (2007). The Social

Lives of Microbes. *Annu Rev Ecol Evol Syst* **38**: 53–77.

Wu HJ, Wang AHJ, Jennings MP. (2008). Discovery of virulence factors of

pathogenic bacteria. *Curr Opin Chem Biol* **12**: 93–101.

Zaborin A, Romanowski K, Gerdes S, Holbrook C, Lepine F, Long J, *et al.*

(2009). Red death in Caenorhabditis elegans caused by Pseudomonas

aeruginosa PAO1. *Proc Natl Acad Sci* **106**: 6327–32.

Zhou L, Slamti L, Nielsen-LeRoux C, Lereclus D, Raymond B. (2014). The

Social Biology of Quorum Sensing in a Naturalistic Host Pathogen System.

Curr Biol **24**: 2417–2422.

Zhu J, Cai X, Harris TL, Gooyit M, Wood M, Lardy M, *et al.* (2015). Disarming

pseudomonas aeruginosa virulence factor lasb by leveraging a caenorhabditis

elegans infection model. *Chem Biol* **22**: 483–491.

 Figure captions

Figure 1. Quantifying *P. aeruginosa* infections in the *C. elegans* gut. (A)

Experimental procedure: we used fluorescently tagged *P. aeruginosa* strains

to examine bacterial colonization of the *C. elegans* gut. Per experiment, we

exposed approximately 200 *C. elegans* nematodes to a lawn of mCherry-

tagged PAO1 strains for 24 hours. Subsequently, nematodes were removed

from the bacterial plate, surface washed and collected in sterile buffer for

monitoring. After 0, 6, or 30 hours post exposure (hpe), approximately 30

nematodes were immobilized and transferred to microscopy slides for imaging.

(B) Brightfield and fluorescence channel merged image depicting mCherry-

fluorescent bacteria inside the host gut. (C) Bacterial load inside the nematode

was quantified as the sum of fluorescence intensity across pixels in the region

of interest "ROI" (yellow outline) and standardised by total worm area. (D)

Colonization dynamics of the wildtype strain PAO1-mCherry: immediately after

removal from the exposure plate (0 hpe), worms showed high bacterial loads

inside their guts. Bacterial load first declined when the worms where kept in

buffer for 6 hours, but then remained constant for the next 24 hours. Grey

shaded area indicates background fluorescence (mean +/- standard deviation)

of worms exposed to the non-fluorescent, non-pathogenic *E.coli* OP50. N =

number of worms from four independent experiments. *** = $p < 0.001$, n.s. =

not statistically significant.

Figure 2. *P. aeruginosa* expresses genes for pyoverdine synthesis and

quorum sensing regulators in the host gut. To quantify the expression of

virulence factor genes inside hosts, worms were exposed to four PAO1

strains, each containing a promoter::mCherry fusion for either *pvdA*

*(*pyoverdine synthesis*), pchEF* (pyochelin synthesis)*, lasR* or *rhlR* (quorum

sensing regulators). With the exception of *pchEF*, all genes were significantly

expressed in the host, both at 0 and 30 hpe. Expression levels were

standardised for bacterial load. Grey shaded areas depict background

fluorescence (mean +/- standard deviation) of worms exposed to the non-

fluorescent, non-pathogenic *E.coli* OP50. N = number of worms from four

independent experiments. * = $p < 0.05$; ** = $p < 0.01$, *** = $p < 0.001$, n.s. = not

statistically significant.

Figure 3. *P. aeruginosa* can switch between siderophores, while quorum

sensing regulators act hierarchically. Because virulence traits are linked at

the regulatory level, we measured gene expression of each trait in the

negative background of the co-regulated trait. (A) The expression of the

pyoverdine synthetic gene *pvdA* is significantly expressed in the wildtype and

the pyochelin-negative background, but slightly reduced in the latter. (B) The

pyochelin synthetic gene *pchEF* is significantly expressed in the pyoverdine-

negative background, but silent in the wildtype. (C) The expression of the QS-

regulator gene *lasR* is unchanged in the Rhl-negative background compared

to the wildtype. (D) The expression of the QS-regulator gene *rhlR* is reduced in

the Las-negative background. Expression levels were standardised for

bacterial load. Grey shaded areas depict background fluorescence (mean +/-

standard deviation) of worms exposed to the non-fluorescent, non-pathogenic

E.coli OP50. N = number of worms form four independent experiments. * = $p <$

0.05; ** = $p < 0.01$, *** = $p < 0.001$, n.s. = not statistically significant.

Figure 4. Virulence factor production affects bacterial uptake and host colonization ability (A) Bacterial load inside *C. elegans* guts measured immediately after the recovery of worms from the exposure plates (0 hours post exposure; hpe). Comparisons across isogenic PAO1 mutant strains, each deficient for the production of one or two virulence factors, reveal that the two quorum-sensing mutants PAO1 Δ*lasR* and PAO1 Δ*rhlR* reached lower bacterial densities than the wildtype. (B) Comparison of the relative colonization success of strains (ratio of bacterial loads at 0 hpe versus 30 hpe) revealed that the siderophore-negative strain PAO1 Δ*pvdD*Δ*pchEF* showed significantly reduced ability to remain in the host compared to the wildtype. In contrast, the colonisation success of PAO1 Δ*pchEF* and PAO1 Δ*rhlR* was increased relative to the wildtype. Grey shaded areas depict background fluorescence (mean +/- standard deviation) of worms exposed to the non-fluorescent, non-pathogenic *E.coli* OP50. N = number of worms form four independent experiments. * = $p < 0.05$; ** = $p < 0.01$, *** = $p < 0.001$.

Figure 5. Mixed infections reveal social strain dynamics but no successful cheating (A) Relative fitness of the wildtype PAO1-mCherry after 42 hours of competition inside the *C. elegans* gut against an untagged PAO1 control strain (white circles); the siderophore-negative strain PAO1 Δ*pvdD*Δ*pchEF* (green diamonds); and the Las-negative strain PAO1 Δ*lasR* (blue squares). The control competition revealed a mild but significant negative effect of the mCherry tag on wildtype fitness. When accounting for these mCherry costs, we found that the putative cheat strains PAO1 Δ*lasR*

and PAO1 Δ*pvdD*Δ*pchEF* performed equally well compared to the wildtype,

but could not outcompete it. This suggests that virulence factor deficient

strains benefit from the presence of non-producers but cannot successfully

cheat on them. (B) At 6 hpe, wildtype frequency in mixed infections correlated

positively with total bacterial load inside hosts in competition with PAO1 Δ*lasR*

(blue line) and PAO1 Δ*pvdD*Δ*pchEF* (green line) but not in the control

competition (grey line). (C) These correlations disappeared at 48 hpe. Each

data point represents an individual worm. Data stems from three independent

experiments, with 8 replicates each.

Figure 6. Spatial structure of mixed infections in the nematode gut. (A, B)

Illustrative examples of *C. elegans* individuals infected with a mixture of GFP-

and mCherry-labelled strains. Each worm was computationally straightened

and fluorescence intensity values were extracted for each pixel from tail (X=0)

to head (X=1). We then calculated the Spearman correlation coefficient

between the intensity values in the two fluorescence channels across pixels,

as our estimate of strain colocalization. Examples show worms with high (A)

and low (B) degrees of colocalization. (C) Patterns of colocalization levels

varied substantially between individuals, but did not differ across strain

combinations (p = 0.119: wildtype PAO1-mCherry versus: (i) wildtype PAO1-

gfp (white circles), (ii) PAO1 Δ*pvdD*Δ*pchEF*-mCherry (green diamonds) or (iii)

PAO1 Δ*lasR*-mCherry (blue squares). Each data point represents an individual

worm. Data stems from 3 independent experiments, with 12 replicates each.

Supplementary Figures captions

Supplementary Figure S1. Growth of *P. aeruginosa* WT and mutant

strains for social traits on NGM plates. To assess the growth ability of

PAO1 and the mutant strains on NGM plates, after 24 hours of incubation at

25°C, we collected the bacterial lawn in sterile NaCl solution and the OD600

was measured as proxy for cell growth. We found no statistically significant

difference between strains (linear model $F_{5,60}$ = 2.27, p = 0.0588). Data is

shown as mean across four independent experiments, with three replicates

(i.e. plates) per strain for each experiment. Error bars denote the standard

error of the mean.

Supplementary Figure S2. Survival assay of worms infected with various

***P. aeruginosa* strains.** After washing worms off the exposure plates, we

estimated host survival over 48 hours. For this purpose, we observed 50 to 90

worms and checked for viability every 24 hours by prodding them with a

platinum wire three times. Worms were considered dead if they no longer

moved upon stimulation with the wire . We found no significant difference in

the survival rate between any of the strains compared to the food strain *E.coli*

OP50-I (linear model, $F_{6,210}$ = 0.60, p = 0.7296). However, we found a small

but significant difference in the survival of worms colonized by the three

siderophore-mutants, which was higher compared to the survival of worms

exposed to PAO1 (Tukey's range test, p = 0.0213 for PAO1 Δ*pchEF*, p =

0.0054 for PAO1 Δ*pvdD* and p = 0.0062 for PAO1 Δ*pvdD*Δ*pchEF*). Data

points depict average survival across three independent experiments. In each

experiment, we had three replicates for each strain. Gray areas represent the

standard error of the mean.

Supplementary Figure S3. Fluorescence intensity significantly correlates

with live bacteria inside the host gut. To assess the relationship between

fluorescence signal and bacterial load inside *C. elegans*, colonized nematodes

were observed using fluorescence microscopy and disrupted to extract live

bacteria from the gut. Fluorescence intensity significantly correlated with the

number of bacteria present in the host gut. (A) The correlation was moderate

when the worms were observed immediately after exposure (0 hours post

exposure; hpe) (Pearson correlation coefficient r = 0.496; test for association

between paired samples t_{28} = 3.02, p = 0.0053). (B) At 6 hpe, fluorescence

intensity correlated more strongly with bacterial load in the host gut (Pearson

correlation coefficient r = 0.713; test for association between paired samples

t_{23} = 4.88, p < 0.0001). In total, 65 worms were observed in two independent

experiments. Fluorescence intensity values were blank corrected, using

worms infected with the untagged strain PAO1 as non-fluorescent controls.

Supplementary Figure S4. Bacterial load declines at 6 hours post

exposure (hpe). We kept infected nematodes in sterile buffer and determined

the fluorescence intensity in the host gut, reflecting the number of bacteria that

managed to colonize the worms at 6 hpe. When scaled to the bacterial load at

0 hpe, we found that all strains showed a significant reduction in bacterial load.

Moreover, the siderophore-negative strain PAO1 Δ*pvdD*Δ*pchEF* showed

significantly reduced ability to remain in the host compared to PAO1 (ANOVA

with post-hoc Tukey test, t_{888} = -2.25, p = 0.0025). For each individual strain,

relative fluorescence is expressed as fluorescence intensity at 6 hpe scaled for

the intensity at 0 hpe.

Supplementary Figure S5. Interaction between social traits inside the

host at 30 hours post exposure. We compared the expression of promoter

fusions inserted either in the wildtype PAO1 or in mutant strains, which lack

either the second siderophore (light grey boxplot) or the second QS-regulator

(white bars), at 30 hpe. Although values are generally lower, we observed the

same trend as in Figure 3. Values are corrected for cell density. N = number of

worms tested. Error bars represent standard errors of the mean. Grey shaded

areas indicate the non-fluorescent background (mean +/- standard deviation).

*** = p < 0.001 and n.s. = not statistically significant, based on Welch's 2-

sample t-test between PAO1 and the respective mutant strain.

A Nematodes washed with buffer + antibiotic

0 hpe →
6 hpe →
30 hpe →

Nematodes immobilized (25 mM NaN₃)

Exposure
Host exposed to mCherry-tagged *P. aeruginosa* strains for 24 h

Observation and measurements
Removal from exposure plate, collection in sterile buffer and microscopy measurement taken at 0, 6 and 30 hours post exposure (hpe).

B

C

D

Initial uptake   Host colonization

***

n.s.

Bacterial load (Log(fluorescent units/area))

N=118   N=154   N=131

Observation time (hours post exposure)