In vivo* microscopy reveals the impact of *Pseudomonas aeruginosa

social interactions on host colonization

Chiara Rezzoagli[1,2*], Elisa T. Granato[3], Rolf Kümmerli[1,2*]

[1]Department of Plant and Microbial Biology, University of Zurich, Zurich,

Switzerland

[2]Department of Quantitative Biomedicine, University of Zurich, Zurich,

Switzerland

[3]Department of Zoology, University of Oxford, Oxford, United Kingdom

Running title: Bacterial social dynamics and host colonization

* Corresponding authors:

Chiara Rezzoagli or Rolf Kümmerli, Department of Quantitative Biomedicine,

University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland.

Email: chiara.rezzoagli@uzh.ch (CR), rolf.kuemmerli@uzh.ch (RK).

Phone: +41 44 635 48 01.

Conflict of interest: The authors declare no conflict of interest.

Abstract

Pathogenic bacteria engage in social interactions to colonize hosts, which include quorum-sensing-mediated communication and the secretion of virulence factors that can be shared as "public goods" between individuals. While *in-vitro* studies demonstrated that cooperative individuals can be displaced by "cheating" mutants freeriding on social acts, we know less about social interactions in infections. Here, we developed a live imaging system to track virulence factor expression and social strain interactions in the human pathogen *Pseudomonas aeruginosa* colonizing the gut of *Caenorhabditis elegans*. We found that shareable siderophores and quorum-sensing systems are expressed during infections, affect host gut colonization, and benefit non-producers. However, non-producers were unable to cheat and outcompete producers. Our results indicate that the limited success of cheats is due to a combination of the down-regulation of virulence factors over the course of the infection, the fact that each virulence factor examined contributed to but was not essential for host colonization, and the potential for negative-frequency dependent selection. Our findings shed new light on bacterial social interactions in infections and reveal potential limits of therapeutic approaches that aim to capitalize on social dynamics between strains for infection control.

Introduction

During infections, pathogenic bacteria secrete a wide range of extracellular virulence factors to colonize and grow inside the host [1, 2]. Secreted molecules include siderophores for iron scavenging, signaling molecules for quorum sensing (QS), toxins to attack host cells, and matrix compounds for biofilm formation [3–6]. *In-vitro* studies have shown that extracellular virulence factors can be shared as "public goods" between cells, and thereby benefit individuals other than the producing cell [7–9]. There has been enormous interest in understanding how this form of bacterial cooperation can be evolutionarily stable since secreted public goods can be exploited by non-cooperative mutants called "cheats", which do not engage in cooperation yet still benefit from the molecules produced by others [10–12].

There is increasing evidence that social interactions and cooperator-cheat dynamics might also matter within hosts [9, 13, 14]. For instance, in controlled infection experiments engineered non-producers, deficient for the production of specific virulence factors, could outcompete producers and thereby reduce virulence [15–19], but in other cases the success of non-producers was compromised [20, 21]. Other studies have followed chronic human infections within patients over time and reported that virulence-factor-negative mutants emerge and spread, with the mutational patterns suggesting cooperator-cheat dynamics [7, 22, 23]. These findings spurred ideas of how social interactions within hosts could be manipulated for therapeutic purposes [14, 24, 25]. Suggested approaches include: inducing cooperator-cheat dynamics to steer infections towards lower virulence [7, 26]; introducing less virulent strains with

medically beneficial alleles into established infections [24]; and targeting

secreted virulence factors to control infections and constrain the evolution of

resistance [25, 27–31].

However, all these approaches explicitly rely on the assumption that the social

traits of interest are: (i) expressed inside hosts; (ii) important for host

colonization; (iii) exploitable; and (iv) induce cooperator-cheat dynamics as

observed *in vitro* [9] – assumptions that have not yet been tested in real time

inside living hosts. Here, we explicitly test the importance of bacterial social

interactions within hosts by using *in-vivo* fluorescence microscopy to monitor

bacterial virulence factor production, host colonization and strain interactions,

using the opportunistic pathogen *Pseudomonas aeruginosa* and its nematode

model host *Caenorhabditis elegans* [32–34].

C. elegans naturally preys on bacteria [35]. While most bacteria are killed

during ingestion, a small fraction of cells survives [36], which can, in the case

of pathogenic bacteria, establish an infection in the gut [37]. *P. aeruginosa*

deploys an arsenal of virulence factors that facilitate successful host

colonization [38]. For example, the two siderophores pyoverdine and pyochelin

scavenge host-bound iron during acute infections to enable pathogen growth

[6, 39–41]. *P. aeruginosa* further secretes the protease elastase, the toxin

pyocyanin, and rhamnolipid biosurfactants to attack host tissue [7, 42, 43].

Production of these latter virulence factors only occurs at high cell densities

and is controlled by the Las and the Rhl QS-systems [44]. Because both QS-

regulated virulence factors and siderophores were shown to be involved in *C.*

elegans killing [37, 45–48], we used them as focal traits for our study.

To tackle our questions, we first conducted experiments with fluorescently

tagged *P. aeruginosa* bacteria (PAO1) to follow infection dynamics, from the

first uptake through feeding up to a progressed state of gut infection. We then

constructed promoter-reporter fusions for genes involved in the synthesis of

the two siderophores (pyoverdine and pyochelin) and the two QS-regulators

(LasR and RhlR) to track *in vivo* virulence factor gene expression during host

colonization. Subsequently, we used mutant strains deficient for virulence

factors to determine whether they show compromised colonization abilities.

Finally, we followed mixed infections of wildtype and mutants over time to

determine the extent of strain co-localization in the gut, and to test whether

secreted virulence factors are indeed exploitable by non-producers in the host.

Material and methods

Strain and bacterial growth conditions

Bacterial strains, primers and plasmids used in this study are listed in the

Supplementary Tables S1-S3. Details on strain construction are given in the

Supplementary Methods. For all experiments, overnight cultures were grown

in 8 ml Lysogeny broth (LB) in 50 ml tubes, incubated at 37°C, 220 rpm for 18

hours, washed with 0.8% NaCl solution and adjusted them to OD_{600} = 1.

Nematode Growth Media (NGM) plates [0.25% Peptone, 50 mM NaCl, 25mM

[PO_4^-], 5 µg/ml Cholesterol, 1mM $CaCl_2$, 1mM $MgSO_4$ supplemented with

1.5% agar, 6 cm diameter] were seeded with 50 µl of bacterial culture and

incubated at 25°C for 24 hours. All *P. aeruginosa* strains used in this study

showed equal growth on NGM exposure plates (Supplementary Figure S1).

Peptone was purchased from BD Biosciences, Switzerland, all other

chemicals from Sigma Aldrich, Switzerland.

Nematode culture

We used the temperature-sensitive, reproductively sterile *C. elegans* strain

JK509 (*glp-1*(*q231*) III). This strain reproduces at 16°C, but does not develop

gonads and is therefore sterile at 25°C. Worms were maintained fertile at 16°C

on High Growth Media (HGM) agar plates (2% Peptone, 50 mM NaCl, 25mM

$[PO_4^-]$, 20 µg/ml Cholesterol, 1mM $CaCl_2$, 1mM $MgSO_4$) seeded with the

standard food source *E. coli* strain OP50 [49]. For age synchronization, plates

were washed with sterile distilled water and adult worms were killed with

hypochlorite-sodium hydroxide solution to isolate eggs [49]. Eggs were placed

in M9 buffer (20 mM KH_2PO_4, 40 mM Na_2HPO_4, 80 mM NaCl, 1 mM $MgSO_4$)

and incubated at 16°C for 16-18 hours to hatch. Then, L1 larvae were

transferred to OP50-seeded HGM plates and incubated at 25°C for 28 hours

to reach L4 developmental stage. Worms and OP50 bacteria were provided by

the Caenorhabditis Genetic Center (CGC), which is supported by the National

Institutes of Health - Office of Research Infrastructure Programs (P40

OD010440).

C. elegans **infection protocol**

Synchronized L4 worms were washed from HGM plates with M9 buffer + 50

µg/ml kanamycin (M9-Kan), and washed three times with M9-Kan for worm

surface-disinfection. Viable worms were further separated from any debris by

sucrose flotation [50] and rinsed three time in M9 buffer to remove sucrose.

The worm handling protocol for the main experiments is depicted in Figure 1A.

Specifically, approximately 200 worms were moved to seeded NGM plates

and incubated for 24 hours at 25°C. After this period of exposure to

pathogens, infected worms were extensively washed with M9 buffer + 50

µg/ml chloramphenicol (M9-Cm) followed by M9 buffer, subsequently

transferred to individual wells of a 6-well plate filled with sterile M9 buffer + 5

µg/ml cholesterol (M9+Ch Buffer) where they were kept up to 48 hours post

exposure (hpe) and imaged after 0, 6 and 30 hpe.

148   Nematode survival assay

Our goal was to observe infections inside living hosts. To verify that worms

stayed alive during the experiment (up to 48 hpe), we tracked the survival of

infected population (50-90 worms) in M9-Ch buffer. Worms were observed for

motility at 0, 24 and 48 hpe, by prodding them with a platinum wire. Worms

were considered dead when they no longer responded to touches. Each

bacterial strain was tested in three replicates and three independent

experiments were carried out. We used *E. coli* OP50 as a negative control for

killing. During this observation period, worms experienced only negligible

killing by the colonizing bacteria, and we found no significant difference in

killing between the non-pathogenic *E. coli* food strain and the *P. aeruginosa*

strains (Supplementary Figure S2).

161   Microscopy setup and imaging

For microscopy, we picked individual worms from the M9+Ch buffer and

paralyzed them with 25 mM sodium azide before transferring them to an 18-

well μ-slide (Ibidi, Germany). All experiments were carried out at the Center for

Microscope and Image Analysis of the University Zürich (ZMB). For the

colonization experiment, images were acquired on a Leica LX inverted

widefield light microscope system with Leica-TX2 filter cube for mCherry

(emission: 560 ± 40 nm, excitation: 645 ± 75 nm, DM = 595) and Leica-DFC-

350-FX, cooled fluorescence monochrome camera (resolution: 1392 × 1040

pixels) for image recording (16-bit color depth). For gene expression

experiments, microscopy was performed on the InCell Analyzer 2500HS (GE

Healthcare) automated imaging system, using polychroic beam splitter

BGRFR_2 (for mCherry, excitation: 575 ± 25 nm, emission: 607.5 ± 19 nm)

and PCO – sCMOS camera (resolution: 2048 x 2048 pixels, 16-bit).

Image processing and analysis

To extract fluorescence measurements from individual worms, images were

segmented into objects and background, using an automated image

segmentation workflow with *ilastik* software [51]. Segmented images were

then imported in *Fiji* [52] to determine the fluorescence intensity (as "Raw

Integrated Density", i.e. the sum of pixels values in the selection) and area of

each worm. Images obtained from the InCell microscope entailed 64 frames

(8x8 grid) with 10% overlap. These frames were stitched together using a

macro-automated version of the Stitching plugin in *Fiji* [53] prior to

segmentation and analysis. To correct for background and host-tissue

autofluorescence, we imaged, at each time point, worms infected with non-

fluorescent strains (i.e. OP50 or PAO1), and used the mean intensity of these

control infections to subtract background fluorescence values from worms

infected with fluorescent strains.

Competition assay in the host

For *in-vivo* competitions between PAO1-*mCherry* and PAO1*ΔpvdDΔpchEF* or

PAO1*ΔlasR*, overnight monocultures were washed twice with 0.8% NaCl

solution, adjusted to $OD_{600} = 1$ and mixed at a 1:1 ratio. To control for fitness

effects of the mCherry marker, we also competed PAO1-*mCherry* against the

untagged PAO1. NGM plates were then seeded with 50 µl of mixed culture

and incubated at 25°C for 24 hours. Worms were exposed to the mix for 24

hours and then recovered as previously described. After 6 and 48 hours post-

exposure, individual worms were picked, immobilized with sodium azide and

washed for 5 minutes with M9 + 0.003% NaOCl. Worms were washed twice

with M9 buffer. We then transferred each individual worm to a 1.5 ml screw-

cap micro tube (Sarstedt, Switzerland) containing sterilized glass beads (1 mm

diameter, Sigma Aldrich). Worms were disrupted using a bead-beater

(TissueLyser II, QIAGEN, Germany), shaking at 30 Hz for 1.5 min before

flipping the tubes and shaking for an additional 1.5 min to ensure even

disruption (adapted from [54]). Tubes were then centrifuged at 2000 x g for 2

min, the content was re-suspended in 200 µl of 0.8% NaCl and plated on two

LB 1.2 % agar plates for each sample. Plates were incubated overnight at

37°C and left at room temperature for another 24 h to allow the fluorescent

marker to fully mature. We then distinguished between fluorescent and non-

fluorescent colonies using a custom built fluorescence imaging device (*Infinity*

3 camera, Lumenera, Canada). We then calculated the relative fitness of

PAO1-*mCherry* as $\ln(v)=\ln\{[a_{48}\times(1-a_6)]/[a_6\times(1-a_{48})]\}$, where a_6 and a_{48} are the

frequency of PAO1-*mCherry* at 6 and 48 hours after recovery, respectively

[55]. Values of $\ln(v)<0$ or $\ln(v)>0$ indicate whether the frequency of PAO1-

mCherry increased ($\ln(v)<0$) or decreased ($\ln(v)>0$) relative to its competitor.

Co-localization analysis

To determine the degree of co-localization of two different bacterial strains in

the host, we transferred nematodes to NGM plates seeded with a 1:1 mix of

PAO1-*gfp* with either PAO1-*mCherry*, PAO1*ΔpvdDΔpchEF–mCherry,* or

PAO1*ΔlasR-mCherry*. After a 24 hours grazing period, we picked single

worms and imaged both mCherry- and GFP channels, using the InCell

Analyzer 2500HS microscope. We used *Fiji* to straighten each worm with the

Straighten plugin [56], and extracted fluorescence intensity values in the GFP

and mCherry channels for each pixel from tail ($X = 0$) to head ($X = 1$) of the

worm. To ensure that we only measure areas where bacteria were present, we

restricted our analysis to the region of the worm gut, where bacterial

colonization takes place. We then calculated Spearman correlation coefficients

between the fluorescent signals, as a proxy for strain co-localization using

RStudio v. 3.3.0 [57].

Statistical analysis

All statistical analyses were performed with RStudio. We used Pearson

correlations to test for associations between PAO1-*mCherry* fluorescence

intensities and (a) recovered bacteria from the gut; and (b) total bacterial load

in mixed infections. We used analysis of variance (ANOVA) to compare

fluorescence values between observation times, strains and for comparisons

to non-fluorescent controls. *P*-values were corrected for multiple comparisons

using the post-hoc Tukey HSD test. To compare promoter expression data

between PAO1 WT and mutant strains, and to compare relative fitness values

between competitors in the competition assay, we used Welch's two-sample t-

test. To measure co-localization, we calculated the Spearman correlation

coefficient ρ between the intensity of mCherry and GFP signals across the

worm gut, and used ANOVA to test for differences between treatments.

Results

PAO1 colonization dynamics in the *C. elegans* gut

For all infection experiments, we followed the protocol depicted in Figure 1A-

C. We first exposed worms to *P. aeruginosa* for 24 hours on NGM plates.

Subsequently, worms were removed, washed, and treated with antibiotics to

kill external bacteria. We then imaged infected worms under the microscope at

different time points and quantified bacterial density and gene expression

using fluorescent mCherry markers. We first confirmed that mCherry

fluorescence is a suitable proxy for the number of live bacteria in *C. elegans*,

by comparing fluorescence intensities in whole worms (Figure 1B) to the

number of live bacteria recovered from the worms' gut. Fluorescence intensity

values positively correlated with the bacterial load inside the nematodes, both

immediately after recovering the worms from the exposure plates and at 6

hours post exposure (hpe; Supplementary Figure S3, Pearson correlation

coefficient at 0 hpe: $r = 0.49$, $t_{28} = 3.02$, $p = 0.0053$; at 6 hpe: $r = 0.713$, $t_{23} =$

4.88, p < 0.0001). As our goal was to image infections in living hosts, we

further confirmed that worms stayed alive during the observation period

(Supplementary Figure S2).

When following host colonization by PAO1-*mCherry* over time, we observed

that immediately after removal from the exposure plate, worms carried large

amounts of bacteria in their gut (Figure 1D). Subsequently, bacterial load

significantly declined when worms were kept in buffer for 6 hours (ANOVA: t_{391}

= -8.55, p < 0.001) and then remained constant for the next 24 hours (t_{391} =

0.61, p = 0.529). This pattern suggests that a large number of bacteria are

taken up during the feeding phase, followed by the shedding of a high

proportion of cells, leaving behind a fraction of live bacteria that establishes an

infection and colonizes the worm gut.

PAO1 expresses siderophore biosynthesis genes and QS regulators in

the host

We then examined whether genes involved in the synthesis of pyoverdine

(*pvdA*) and pyochelin (*pchEF*), and the genes encoding the QS-regulators

lasR and *rhlR*, are expressed inside hosts. Worms were exposed to four

different PAO1 strains, each carrying a specific promoter-*mCherry* fusion.

Imaging after the initial uptake phase (0 hpe) revealed that, with the exception

of *pchEF*, all genes were significantly expressed in the host (Figure 2;

ANOVA, comparisons to the non-fluorescent control, for *pvdA*: t_{754} = 4.23, p <

0.001; for *pchEF*: t_{754} = 0.74, p = 0.461; for *lasR*: t_{754} = 2.96, p = 0.003*; for*

rhlR: t_{754} = 10.37, p <0.001*).* Although fluorescence intensity declined over

time (linear model, $F_{1,1795}$ = 48.98, p < 0.001), we observed that apart from

pchEF, all genes were still significantly expressed during the subsequent

colonisation of the host at 30 hpe. (Figure 2; ANOVA, for *pvdA*: t_{754} = 4.87, p <

0.001*;* for *pchEF*: t_{754} = 0.684, p = 0.461; for *lasR*: t_{754} = 3.01, p = 0.003; for

rhlR: t_{754} = 16.68, p < 0.001).

Regulatory links between social traits operate inside the host

We know that regulatory links exist between the virulence traits studied here.

While pyoverdine synthesis suppresses pyochelin production under stringent

iron limitation [58], the Las-QS system positively activates the Rhl-QS system

[44]. To test whether these links operate inside the host, we measured gene

expression of each trait in the negative background of the co-regulated trait

(Figure 3). For *pvdA*, we observed significant gene expression levels in both

the wildtype PAO1 and the pyochelin-deficient PAO1Δ*pchEF* strain (Figure

3A), albeit the overall expression was slightly reduced in PAO1Δ*pchEF* (t-test,

t_{253} = 8.67, p < 0.001). For *pchEF*, expression patterns confirm the

suppressive nature of pyoverdine: the pyochelin synthesis gene was not

expressed in the wildtype but significantly upregulated in the pyoverdine-

deficient PAO1Δ*pvdD* strain (Figure 3B; t_{296} = -19.68, p < 0.001). For *lasR*, we

found that gene expression was not significantly different in wildtype PAO1

compared to the Rhl-negative mutant PAO1Δ*rhlR*, confirming that the Las-QS

system is at the top of the hierarchy and not influenced by the Rhl-system

(Figure 3C; t_{211} = -1.50, p = 0.136). Conversely, the expression of *rhlR* was

strongly dependent on a functional Las-system, and therefore only expressed

in PAO1, but repressed in the Las-negative mutant PAO1Δ*lasR* (Figure 3D;

$t_{156} = 19.04$, p < 0.001). These results show that (i) iron-limitation is strong in

C. elegans as PAO1 primarily invests in the more potent siderophore

pyoverdine; (ii) pyochelin can have compensatory effects when pyoverdine is

lacking; and (iii) the loss of the Las-system leads to the concomitant collapse

of the Rhl-system.

Virulence-factor-negative mutants show trait-specific deficiencies in host

colonization

To examine whether the ability to produce shared virulence factors is

important for initial bacterial uptake and host colonization, we exposed

C. elegans to five isogenic mutants of PAO1-*mCherry*, either impaired in the

production of pyoverdine (Δ*pvdD*), pyochelin (Δ*pchEF*), both siderophores

(Δ*pvdD*Δ*pchEF*), the QS receptor LasR (Δ*lasR*), or the QS receptor RhlR

(Δ*rhlR*). After the feeding phase, the bacterial load of the wildtype and all three

siderophores mutants were equally abundant inside hosts, whereas bacterial

load was significantly reduced for the two QS-mutants compared to the

wildtype (Figure 4A; ANOVA, significant variation among strains $F_{5,736} = 10.50$,

p < 0.001; post-hoc Tukey test for multiple comparisons: p > 0.05 for all

siderophore mutants, p = 0.021 for PAO1Δ*lasR*, p < 0.001 for PAO1Δ*rhlR*).

As previously described for PAO1 colonization (Figure 1D), we observed that

the bacterial load of all strains declined at 6 hpe (Supplementary Figure S4)

and 30 hpe (Figure 4B) following worm removal from the exposure plates. This

decline was significantly more pronounced for the double-siderophore

knockout PAO1Δ*pvdD*Δ*pchEF* than for the wildtype (Figure 4B; ANOVA, post-

hoc Tukey test p < 0.001). In contrast, mutants deficient in pyochelin

(PAO1ΔpchEF) and RhlR (PAO1ΔrhlR) production showed a significantly

higher ability to remain in the host than the wildtype (Figure 4B; ANOVA, post-

hoc Tukey test p <0.001 for both strains). Taken together, our findings suggest

that the two siderophores can complement each other, and that only the

siderophore double mutant and the LasR-deficient strain have an overall

disadvantage in colonizing worms.

Mixed communities are formed inside hosts, but exploitation of social

traits is constrained

Given our findings on colonization deficiencies, we reasoned that the

siderophore-double mutant (PAO1ΔpvdDΔpchEF) and the Las-deficient

mutant (PAO1ΔlasR) could act as cheats and benefit from the exploitation of

virulence factors produced by the wildtype in mixed infections. To test this

hypothesis, we first competed the PAO1-mcherry strain against the untagged

wildtype in the host, and found that the mCherry tag had a small negative

effect on PAO1 fitness (Figure 5A; one sample t-test, t_{24} = -4.12, p < 0.001).

We then competed PAO1-mCherry against the two putative cheats and found

that neither of them could gain a significant fitness advantage over the

wildtype, but also did not lose out (Figure 5A; ANOVA, $F_{2,70}$ = 0.517, p =

0.598). These results indicate that virulence-factor-negative mutants, initially

compromised in host colonization, can indeed benefit from the presence of the

wildtype producer, but not to an extent that would allow them to increase in

frequency and displace producers.

Since our mono-infection experiments showed that the wildtype can maintain

higher bacterial loads in the worms compared to the two mutants (Figure 4B),

we hypothesized that worms, which have initially taken up higher frequencies

of the wildtype relative to the mutant should carry increased bacterial loads in

the gut. We found this prediction to hold true at 6 hpe in mixed infections with

the two non-producers, but not in the control mixed infections with the

untagged wildtype (Figure 5B, 6hpe; Pearson correlation coefficient, for mixed

infection with PAO1Δ*lasR*: r = 0.54, t_{17} = 2.67, p = 0.016; with

PAO1Δ*pvdD*Δ*pchEF*: r = 0.40, t_{17} = 1.77, p = 0.031; with control PAO1: r =

0.12, t_{17} = 0.47, p = 0.639). These correlations disappeared at the later

colonization stage (Figure 5B, 48 hpe; Pearson correlation coefficient r < 0, p

> 0.05 for all strains). The loss of these correlations indicates that rare

producers experienced a selective advantage during competition and

increased in relative frequency, while common producers might have lost and

decreased in frequency.

Strain co-localization is generally high within the host, but varies

substantially across individuals

In-vitro studies have shown that spatial proximity of cells is crucial for efficient

compound sharing [59, 60]. We thus assessed the co-localization of strains in

mixed infections inside the gut (Figure 6). We found that all worms were

colonized by both strains, and that the level of co-localization ρ (from tail to

head) was generally high, although it varied substantially across individuals

(Figure 6A-B). Similar co-localization patterns emerged for all three strain

combinations tested, highlighting that the type of competitor did not influence

the degree of strain co-localization in the host gut (Figure 6C; ANOVA, $F_{2,102} =$

2.17, p = 0.119). While our measure of co-localization has some limitations as

it does not reveal physical proximity at the single cell level, and is based on a

2D projection of a 3D organ, it clearly suggests that competing cells are close

to one another, and that social interactions could occur between them.

Discussion

We developed a live imaging system that allows us to track host colonization

by pathogenic bacteria (*P. aeruginosa*) and their expression of virulence

factors inside hosts (*C. elegans*). We used this system to focus on the role of

secreted virulence factors, which can be shared as public goods between

bacterial cells, and examined competitive dynamics between virulence factor

producing and non-producing strains in the host. We found that siderophores

(pyoverdine and pyochelin) and the Las and Rhl QS-systems (i) are expressed

inside the host; (ii) affect the ability to colonize and reside within the

nematodes; (iii) allow non-producers to benefit from virulence factors secreted

by producers in mixed infections; but (iv) do not allow non-producers to cheat

and outcompete producers. Our results have implications for both the

understanding of bacterial social interactions within hosts, and therapeutic

approaches that aim to manipulate social dynamics between strains for

infection control.

Numerous *in-vitro* studies have shown that bacterial cooperation can be

exploited by cheating mutants that no longer express the social trait, but

benefit from the cooperative acts performed by others [3, 11, 12, 55, 61–65].

These findings contrast with our observations that the spread of non-
producers was constrained within infections. There are multiple ways to
explain this constraint. First, increased spatial structure can limit the diffusion
of secreted metabolites and lead to the physical separation of cooperators and
cheats [63]. Both effects result in metabolites being shared more locally
among cooperative individuals. The physical separation of strains seemed to
explain the results of Zhou *et al* [20], where QS-mutants of *Bacillus*
thuringiensis infecting caterpillars could not exploit metabolites from
producers. Conversely, physical separation seemed low in our study system
(Figure 6), and therefore unlikely explains why cheats could not spread. While
we solely focussed on proximity patterns inside hosts, it is important to note
that processes at the meta-population level such as bottlenecking [54, 66], can
reduce the probability of different strains ending up in the same host, and thus
further compromise cheat success.

Second, negative frequency-dependent selection could explain why the
spread of virulence factor negative mutants is constrained [55]. This scenario
predicts that cheats only experience a selective advantage when rare, but not
when common. The reasoning is that non-producers can only efficiently exploit
public goods when surrounded by many producers. Our competition
experiments indeed provide indirect evidence for negative frequency-
dependent selection in the nematode gut (Figure 5B). Specifically, we
observed that bacterial load was reduced when producers occurred at low
frequency early during infection (6 hpe), a result confirming that non-producers
are worse host colonizers than producers. These correlations disappeared

during the competition period (48 hpe), indicating that rare producers might

have experienced a selective advantage and increased in relative frequency,

while common producers lost and decreased in frequency.

Third, the relatively low bacterial density in the gut could further compromise

the ability of non-producers to cheat (Figure 1D, 5B). Low cell density restricts

the sharing and exploitation of secreted compounds [67, 68]. Mechanisms

responsible for the low bacterial density in the gut (Figure 1D, 5B) could

include the peristaltic activity of the gut, expelling a part of the pathogen

population and the host immune system, killing a fraction of the bacteria [69].

Fourth, our analysis reveals that, although siderophores and QS-systems play

a role in host colonization, they are not essential (Figure 4). Moreover, the

expression of pyoverdine and QS-systems declined over time (Figure 2).

These two observations indicate that the benefits of cheating might be fairly

low, and that the costs of virulence factor production are reduced at later

stages of the infection. Thus, bacteria might switch from production to

recycling of already secreted public goods [70, 71], an effect that can hamper

the spread of cheats.

Finally, we show that the regulatory linkage between traits is an important

factor to consider when predicting the putative advantage of non-producers

[72,   73].   For   instance,   *P. aeruginosa*   pyoverdine-negative-mutants

upregulated pyochelin production to compensate for the lack of their primary

siderophore (Figure 3). Thus, if pyoverdine-negative mutants evolve *de novo*,

their spread as cheats could be hampered because they invest in pyochelin as

an alternative siderophore [74]. For QS, meanwhile, we observed that the

absence of a functional Las-system resulted in the concomitant collapse of the

Rhl-system. Although *lasR* mutants could be potent cheats, as they are

deficient for multiple social traits, their spread might be hampered because

QS-systems also regulate non-social traits, important for individual fitness [75].

When relating our insights to previous studies, it turns out that earlier work

produced mixed results with regard to the question whether siderophore- and

QS-deficient mutants can spread within infections. While Harrison *et al.* ([15,

21]; pyoverdine, *P. aeruginosa* in *Galleria mellonella* and *ex-vivo* infection

models) and Zhou *et al.* ([20]; QS, *B. thuringiensis* in *Plutella xylostella*)

showed that the spread of non-producers is constrained, Rumbaugh *et al.*

([16, 17]; QS; *P. aeruginosa* in mice), Pollitt *et al.* ([19]; QS, *Staphylococcus*

aureus in *G. mellonella*) and Diard *et al.* ([18], T3SS-driven inflammation,

Salmonella typhimorium in mice) demonstrated cases where non-producers

spread to high frequencies in host populations. While the reported results were

based on strain frequency counts before and after competition, we here show

that information on social trait expression, temporal infection dynamics and

physical interactions among strains within hosts are essential to understand

whether social traits are important and exploitable in a given system. We thus

posit that more such detailed approaches are required to understand the

importance of bacterial social interactions across host systems and infection

contexts and explain differences between them.

A deeper understanding of bacterial social interactions inside hosts is

particularly relevant for novel therapeutic approaches that seek to take

advantage of cooperator-cheat dynamics inside hosts to control infections. For

instance, it was proposed that strains deficient for virulence factors could be

introduced into established infections [24]. These strains are expected to

spread because of cheating, thereby reducing the overall virulence factor

availability in the population and the damage to the host. Our results reveal

that virulence-factor-negative strains, although eventually gaining a benefit

from producer strains, are unable to spread in populations. Another

therapeutic approach involves the specific targeting of secreted virulence

factors [25, 27]. This approach is thought to reduce damage to the host and to

compromise resistance evolution [30]. Resistant mutants, resuming virulence

factor production, are not expected to spread because they would act as

cooperators, sharing the benefit of secreted goods with susceptible strains

[76–78]. Our results yet indicate that such resistant mutants could get local

benefits and thus increase to a certain frequency in the population [31]. These

confrontations show that the identification of key parameters driving social

interactions across hosts and infection types is of utmost importance to predict

the success of 'cheat therapies' and anti-virulence strategies targeting

secreted public goods.

Acknowledgments

We thank two anonymous reviewers for constructive comments and the

Center of Microscopy and Image Analysis (University of Zürich) for support

with image acquisition and advice on image analysis.

Funding

This project has received funding from the Swiss National Science Foundation

(grant no. PP00P3_165835 and 31003A_182499 to RK and no.

P2ZHP3_174751 to EG), and the European Research Council under the grant

agreement no. 681295 (to RK).

Competing Interests

The authors have no competing interests to declare.

Supplementary Information

Supplementary information is available at the journal's website.

References:

1.   Rahme LG, Stevens EJ, Wolfort SF, Shao J, Tompkins RG, Ausubel
FM. Common virulence factors for bacterial pathogenicity in plants and
animals. *Science* 1995; **268**: 1899–902.

2.   Wu HJ, Wang AHJ, Jennings MP. Discovery of virulence factors of
pathogenic bacteria. *Curr Opin Chem Biol* 2008; **12**: 93–101.

3.   Diggle SP, Griffin AS, Campbell GS, West SA. Cooperation and conflict
in quorum-sensing bacterial populations. *Nature* 2007; **450**: 411–414.

4.   Flemming HC, Wingender J, Szewzyk U, Steinberg P, Rice SA,
Kjelleberg S. Biofilms: An emergent form of bacterial life. *Nat Rev*
Microbiol 2016; **14**: 563–575.

5.   Henkel JS, Baldwin MR, Barbieri JT. Toxins from bacteria. *EXS* 2010;
100: 1–29.

6.   Granato ET, Harrison F, Kümmerli R, Ross-Gillespie A. Do Bacterial

"Virulence Factors" Always Increase Virulence? A Meta-Analysis of

Pyoverdine Production in *Pseudomonas aeruginosa* As a Test Case.

Front Microbiol 2016; **7**: 1952.

7.   Köhler T, Buckling A, van Delden C. Cooperation and virulence of

clinical *Pseudomonas aeruginosa* populations. *Proc Natl Acad Sci USA*

2009; **106**: 6339–6344.

8.   Raymond B, West SA, Griffin AS, Bonsall MB. The dynamics of

cooperative bacterial virulence in the field. *Science* 2012; **337**: 85–88.

9.   Harrison F. Bacterial cooperation in the wild and in the clinic: Are

pathogen social behaviours relevant outside the laboratory? *BioEssays*

2013; **35**: 108–112.

10.   West SA, Diggle SP, Buckling A, Gardner A, Griffin AS. The Social Lives

of Microbes. *Annu Rev Ecol Evol Syst* 2007; **38**: 53–77.

11.   Ghoul M, Griffin AS, West SA. Toward an evolutionary definition of

cheating. *Evolution* 2014; **68**: 318–331.

12.   Özkaya Ö, Balbontín R, Gordo I, Xavier KB. Cheating on Cheaters

Stabilizes Cooperation in *Pseudomonas aeruginosa*. *Curr Biol* 2018; **26**:

2070–2080.

13.   Buckling A, Brockhurst MA. Kin selection and the evolution of virulence.

Heredity 2008; **100**: 484–488.

14.   Leggett HC, Brown SP, Reece SE. War and peace: social interactions in

infections. *Philos Trans R Soc Lond B Biol Sci* 2014; **369**: 20130365.

15.   Harrison F, Browning LE, Vos M, Buckling A. Cooperation and virulence

in acute *Pseudomonas aeruginosa* infections. *BMC Biol* 2006; **4**: 21.

16.   Rumbaugh KP, Diggle SP, Watters CM, Ross-Gillespie A, Griffin AS,

West SA. Quorum Sensing and the Social Evolution of Bacterial

Virulence. *Curr Biol* 2009; **19**: 341–345.

17.   Rumbaugh KP, Trivedi U, Watters C, Burton-Chellew MN, Diggle SP,

West SA. Kin selection, quorum sensing and virulence in pathogenic

bacteria. *Proc R Soc B Biol Sci* 2012; **279**: 3584–3588.

18.   Diard M, Garcia V, Maier L, Remus-Emsermann MNP, Regoes RR,
570   Ackermann M, et al. Stabilization of cooperative virulence by the
571   expression of an avirulent phenotype. *Nature* 2013; **494**: 353–356.

572   19.   Pollitt EJG, West SA, Crusz SA, Burton-Chellew MN, Diggle SP.
573   Cooperation, quorum sensing, and evolution of virulence in
574   *Staphylococcus aureus*. *Infect Immun* 2014; **82**: 1045–1051.

575   20.   Zhou L, Slamti L, Nielsen-LeRoux C, Lereclus D, Raymond B. The
576   Social Biology of Quorum Sensing in a Naturalistic Host Pathogen
577   System. *Curr Biol* 2014; **24**: 2417–2422.

578   21.   Harrison F, McNally A, Da Silva AC, Heeb S, Diggle SP. Optimised
579   chronic infection models demonstrate that siderophore 'cheating' in
580   *Pseudomonas aeruginosa* is context specific. *ISME J* 2017; **11**: 2492–
581   2509.

582   22.   Andersen SB, Marvig RL, Molin S, Krogh Johansen H, Griffin AS. Long-
583   term social dynamics drive loss of function in pathogenic bacteria. *Proc*
584   *Natl Acad Sci USA* 2015; **112**: 10756–10761.

585   23.   Andersen SB, Ghoul M, Marvig RL, Lee Z Bin, Molin S, Johansen HK, et
586   al. Privatisation rescues function following loss of cooperation. *Elife*
587   2018; **7**: e38594.

588   24.   Brown SP, West S a, Diggle SP, Griffin AS. Social evolution in micro-
589   organisms and a Trojan horse approach to medical intervention
590   strategies. *Philos Trans R Soc Lond B Biol Sci* 2009; **364**: 3157–3168.

591   25.   Allen RC, Popat R, Diggle SP, Brown SP. Targeting virulence: can we
592   make evolution-proof drugs? *Nat Rev Microbiol* 2014; **12**: 300–308.

593   26.   Granato ET, Ziegenhain C, Marvig RL, Kümmerli R. Low spatial
594   structure and selection against secreted virulence factors attenuates
595   pathogenicity in *Pseudomonas aeruginosa*. *ISME J* 2018; **12**: 2907–
596   2918.

597   27.   André JB, Godelle B. Multicellular organization in bacteria as a target for
598   drug therapy. *Ecol Lett* 2005; **8**: 800–810.

28. Clatworthy AE, Pierson E, Hung DT. Targeting virulence: a new
paradigm for antimicrobial therapy. *Nat Chem Biol* 2007; **3**: 541–548.

29. Rasko DA, Sperandio V. Anti-virulence strategies to combat bacteria-
mediated disease. *Nat Rev Drug Discov* 2010; **9**: 117–128.

30. Pepper JW. Drugs that target pathogen public goods are robust against
evolved drug resistance. *Evol Appl* 2012; **5**: 757–761.

31. Rezzoagli C, Wilson D, Weigert M, Wyder S, Kümmerli R. Probing the
evolutionary robustness of two repurposed drugs targeting iron uptake in
Pseudomonas aeruginosa. *Evol Med Public Heal* 2018; **1**: 246–259.

32. Tan MW, Ausubel FM. *Caenorhabditis elegans*: A model genetic host to
study *Pseudomonas aeruginosa* pathogenesis. *Curr Opin Microbiol*
2000; **3**: 29–34.

33. Ewbank JJ. Tackling both sides of the host – pathogen equation with
Caenorhabditis elegans. *Microbes Infect* 2002; **4**: 247–256.

34. Papaioannou E, Utari P, Quax W. Choosing an Appropriate Infection
Model to Study Quorum Sensing Inhibition in *Pseudomonas Infections*.
Int J Mol Sci 2013; **14**: 19309–19340.

35. Félix M-A, Braendle C. The natural history of *Caenorhabditis elegans*.
Curr Biol 2010; **20**: R965–R969.

36. Portal-Celhay C, Bradley ER, Blaser MJ. Control of intestinal bacterial
proliferation in regulation of lifespan in *Caenorhabditis elegans*. *BMC*
Microbiol 2012; **12**: 49.

37. Tan MW, Mahajan-Miklos S, Ausubel FM. Killing of *Caenorhabditis*
elegans by *Pseudomonas aeruginosa* used to model mammalian
bacterial pathogenesis. *Proc Natl Acad Sci USA* 1999; **96**: 715–20.

38. Jimenez PN, Koch G, Thompson JA, Xavier KB, Cool RH, Quax WJ.
The multiple signaling systems regulating virulence in *Pseudomonas*
aeruginosa. *Microbiol Mol Biol Rev* 2012; **76**: 46–65.

39. Meyer JM, Neely A, Stintzi A, Georges C, Holder IA. Pyoverdin is
essential for virulence of *Pseudomonas aeruginosa*. *Infect Immun* 1996;

64: 518–523.

40.   Takase H, Nitanai H, Hoshino K, Otani T. Impact of Siderophore
631   Production on *Pseudomonas aeruginosa* Infections in
632   Immunosuppressed Mice. *Infect Immun* 2000; **68**: 1834–1839.

633   41.   Cornelis P, Dingemans J. *Pseudomonas aeruginosa* adapts its iron
634   uptake strategies in function of the type of infections. *Front Cell Infect*
635   *Microbiol* 2013; **3**: 1–7.

636   42.   Smith RS, Iglewski BH. *P. aeruginosa* quorum-sensing systems and
637   virulence. *Curr Opin Microbiol* 2003; **6**: 56–60.

638   43.   Alibaud L, Köhler T, Coudray A, Prigent-Combaret C, Bergeret E, Perrin
639   J, et al. *Pseudomonas aeruginosa* virulence genes identified in a
640   *Dictyostelium* host model. *Cell Microbiol* 2008; **10**: 729–740.

641   44.   Lee J, Zhang L. The hierarchy quorum sensing network in
642   Pseudomonas aeruginosa. *Protein Cell* 2015; **6**: 26–41.

643   45.   Zaborin A, Romanowski K, Gerdes S, Holbrook C, Lepine F, Long J, et
644   al. Red death in *Caenorhabditis elegans* caused by *Pseudomonas*
645   *aeruginosa* PAO1. *Proc Natl Acad Sci USA* 2009; **106**: 6327–32.

646   46.   Kirienko N V., Kirienko DR, Larkins-Ford J, W??hlby C, Ruvkun G,
647   Ausubel FM. Pseudomonas aeruginosa disrupts Caenorhabditis elegans
648   iron homeostasis, causing a hypoxic response and death. *Cell Host*
649   *Microbe* 2013; **13**: 406–416.

650   47.   Cezairliyan B, Vinayavekhin N, Grenfell-Lee D, Yuen GJ, Saghatelian A,
651   Ausubel FM. Identification of *Pseudomonas aeruginosa* phenazines that
652   kill *Caenorhabditis elegans*. *PLOS Pathog* 2013; **9**: e1003101.

653   48.   Zhu J, Cai X, Harris TL, Gooyit M, Wood M, Lardy M, et al. Disarming
654   *Pseudomonas aeruginosa* Virulence Factor LasB by Leveraging a
655   *Caenorhabditis elegans* Infection Model. *Chem Biol* 2015; **22**: 483–491.

656   49.   Stiernagle T. Maintenance of C. elegans. In: The *C.elegans* Research
657   Community (ed). *WormBook: The Online Review of C. elegans Biology.*
658   2006.

659   50.   Portman DS. Profiling C. elegans gene expression with DNA
660   microarrays. In: The C.elegans Research Community (ed). *WormBook:*
661   *The Online Review of C. elegans Biology.* 2006.

662   51.   Sommer C, Strähle C, Köthe U, Hamprecht FA. ilastik: Interactive
663   Learning and Segmentation Toolkit. *Eighth IEEE Int. Symp. Biomed.*
664   *Imaging (ISBI). Proc.* 2011. pp 230–233.

665   52.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch
666   T, et al. Fiji: an open-source platform for biological-image analysis. *Nat*
667   *Methods* 2012; **9**: 676–82.

668   53.   Preibisch S, Saalfeld S, Tomancak P. Globally optimal stitching of tiled
669   3D microscopic image acquisitions. *Bioinformatics* 2009; **25**: 1463–
670   1465.

671   54.   Vega NM, Gore J. Stochastic assembly produces heterogeneous
672   communities in the *Caenorhabditis elegans* intestine. *PLOS Biol* 2017;
673   **15**: e2000633.

674   55.   Ross-Gillespie A, Gardner A, West SA, Griffin AS. Frequency
675   dependence and cooperation: theory and a test with bacteria. *Am Nat*
676   2007; **170**: 331–342.

677   56.   Kocsis E, Trus BL, Steer CJ, Bisher ME, Steven AC. Image averaging of
678   flexible fibrous macromolecules: the clathrin triskelion has an elastic
679   proximal segment. *J Struct Biol* 1991; **107**: 6–14.

680   57.   R Development Core Team. R: A language and environment for
681   statistical computing. *R Foundation for Statistical Computing.* 2013.
682   Vienna, Austria.

683   58.   Dumas Z, Ross-Gillespie A, Kümmerli R. Switching between apparently
684   redundant iron-uptake mechanisms benefits bacteria in changeable
685   environments. *Proc Biol Sci* 2013; **280**: 20131055.

686   59.   Van Gestel J, Weissing FJ, Kuipers OP, Kovács ÁT. Density of founder
687   cells affects spatial pattern formation and cooperation in *Bacillus subtilis*
688   biofilms. *ISME J* 2014; **8**: 2069–2079.

60. Weigert M, Kümmerli R. The physical boundaries of public goods
cooperation between surface-attached bacterial cells. *Proc R Soc B Biol*
Sci 2017; **284**: 20170631.

61. Griffin AS, West SA, Buckling A. Cooperation and competition in
pathogenic bacteria. *Nature* 2004; **430**: 1024–1027.

62. Sandoz KM, Mitzimberg SM, Schuster M. Social cheating in
Pseudomonas aeruginosa quorum sensing. *Proc Natl Acad Sci USA*
2007; **104**: 15876–81.

63. Kümmerli R, Griffin AS, West S a, Buckling A, Harrison F. Viscous
medium promotes cooperation in the pathogenic bacterium
Pseudomonas aeruginosa. Proc Biol Sci 2009; **276**: 3531–3538.

64. Popat R, Crusz SA, Messina M, Williams P, West SA, Diggle SP.
Quorum-sensing and cheating in bacterial biofilms. *Proc R Soc B Biol*
Sci 2012; **279**: 4765–4771.

65. O'Brien S, Luján AM, Paterson S, Cant MA, Buckling A. Adaptation to
public goods cheats in *Pseudomonas aeruginosa. Proc R Soc B Biol Sci*
2017; **284**: 20171089.

66. van Leeuwen E, O'Neill S, Matthews A, Raymond B. Making pathogens
sociable: The emergence of high relatedness through limited host
invasibility. *ISME J* 2015; **9**: 2315–2323.

67. Ross-Gillespie A, Gardner A, Buckling A, West SA, Griffin AS. Density
dependence and cooperation: theory and a test with bacteria. *Evolution*
2009; **63**: 2315–2325.

68. Scholz RL, Greenberg EP. Sociality in *Escherichia coli*: Enterochelin Is a
Private Good at Low Cell Density and Can Be Shared at High Cell
Density. *J Bacteriol* 2015; **197**: 2122–2128.

69. Pukkila-Worley R, Ausubel FM. Immune defense mechanisms in the
Caenorhabditis elegans intestinal epithelium. *Curr Opin Immunol* 2012;
24: 3–9.

70. Imperi F, Tiburzi F, Visca P. Molecular basis of pyoverdine siderophore

recycling in *Pseudomonas aeruginosa. Proc Natl Acad Sci USA* 2009;
106: 20440–20445.

71. Kümmerli R, Brown SP. Molecular and regulatory properties of a public
good shape the evolution of cooperation. *Proc Natl Acad Sci USA* 2010;
107: 18921–6.

72. Lindsay RJ, Kershaw MJ, Pawlowska BJ, Talbot NJ, Gudelj I.
Harbouring public good mutants within a pathogen population can
increase both fitness and virulence. *Elife* 2016; **5**: 1–25.

73. dos Santos M, Ghoul M, West SA. Pleiotropy, cooperation and the social
evolution of genetic architecture. *PLOS Biol* 2018; **16**: e2006671.

74. Ross-Gillespie A, Dumas Z, Kümmerli R. Evolutionary dynamics of
interlinked public goods traits: An experimental study of siderophore
production in *Pseudomonas aeruginosa. J Evol Biol* 2015; **28**: 29–39.

75. Dandekar AA, Chugani S, Greenberg PE. Bacterial Quorum Sensing
and Metabolic Incentives to Cooperate. *Science* 2012; **338**: 264–266.

76. Mellbye B, Schuster M. The sociomicrobiology of antivirulence drug
resistance: a proof of concept. *MBio* 2011; **2**: e00131-11.

77. Gerdt JP, Blackwell HE. Competition Studies Confirm Two Major
Barriers That Can Preclude the Spread of Resistance to Quorum-
Sensing Inhibitors in Bacteria. *ACS Chem Biol* 2014; **9**: 2291–2299.

78. Ross-Gillespie A, Weigert M, Brown SP, Kümmerli R. Gallium-mediated
siderophore quenching as an evolutionarily robust antibacterial
treatment. *Evol Med Public Heal* 2014; **2014**: 18–29.

Figure captions

Figure 1. Quantifying *P. aeruginosa* infections in the *C. elegans* gut. (A) Experimental procedure: we used fluorescently tagged *P. aeruginosa* strains to examine bacterial colonization of the *C. elegans* gut. Per experiment, we exposed approximately 200 *C. elegans* nematodes to a lawn of mCherry-tagged PAO1 strains for 24 hours. Subsequently, nematodes were removed from the bacterial plate, surface washed and collected in sterile buffer for monitoring. After 0, 6, or 30 hours post exposure (hpe), approximately 30 nematodes were immobilized and transferred to microscopy slides for imaging. (B) Brightfield and fluorescence channel merged image depicting mCherry-fluorescent bacteria inside the host gut. (C) Bacterial load inside the nematode was quantified as the sum of fluorescence intensity across pixels in the region of interest "ROI" (yellow outline) and standardised by total worm area. (D) Colonization dynamics of the wildtype strain PAO1-*mCherry*: immediately after removal from the exposure plate (0 hpe), worms showed high bacterial loads inside their guts. Bacterial load first declined when the worms where kept in buffer for 6 hours, but then remained constant for the next 24 hours. Grey shaded area indicates background fluorescence (mean +/- standard deviation) of worms exposed to the non-fluorescent, non-pathogenic *E.coli* OP50. N = number of worms from four independent experiments. *** = $p < 0.001$, n.s. = not statistically significant.

Figure 2. *P. aeruginosa* expresses genes for pyoverdine synthesis and quorum sensing regulators in the host gut. To quantify the expression of virulence factor genes inside hosts, worms were exposed to four PAO1

strains, each containing a promoter::*mCherry* fusion for either *pvdA*

(pyoverdine synthesis*), pchEF* (pyochelin synthesis)*, lasR* or *rhlR* (quorum

sensing regulators). With the exception of *pchEF*, all genes were significantly

expressed in the host, both at 0 and 30 hpe. Expression levels were

standardised for bacterial load. Grey shaded areas depict background

fluorescence (mean +/- standard deviation) of worms exposed to the non-

fluorescent, non-pathogenic *E.coli* OP50. N = number of worms from four

independent experiments. * = p < 0.05; ** = p < 0.01, *** = p < 0.001, n.s. = not

statistically significant.

Figure 3. *P. aeruginosa* can switch between siderophores, while quorum

sensing regulators act hierarchically. Because virulence traits are linked at

the regulatory level, we measured gene expression of each trait in the

negative background of the co-regulated trait. (A) The expression of the

pyoverdine synthetic gene *pvdA* is significantly expressed in the wildtype and

the pyochelin-negative background, but slightly reduced in the latter. (B) The

pyochelin synthetic gene *pchEF* is significantly expressed in the pyoverdine-

negative background, but silent in the wildtype. (C) The expression of the QS-

regulator gene *lasR* is unchanged in the Rhl-negative background compared

to the wildtype. (D) The expression of the QS-regulator gene *rhlR* is reduced in

the Las-negative background. Expression levels were standardised for

bacterial load. Grey shaded areas depict background fluorescence (mean +/-

standard deviation) of worms exposed to the non-fluorescent, non-pathogenic

E.coli OP50. N = number of worms form four independent experiments. * = p <

0.05; ** = p < 0.01, *** = p < 0.001, n.s. = not statistically significant.

Figure 4. Virulence factor production affects bacterial uptake and host

colonization ability. (A) Bacterial load inside *C. elegans* guts measured

immediately after the recovery of worms from the exposure plates (0 hours

post exposure; hpe). Comparisons across isogenic PAO1 mutant strains, each

deficient for the production of one or two virulence factors, reveal that the two

quorum-sensing mutants PAO1Δ*lasR* and PAO1Δ*rhlR* reached lower bacterial

densities than the wildtype. (B) Comparison of the relative colonization

success of strains (ratio of bacterial loads at 0 hpe versus 30 hpe) revealed

that the siderophore-negative strain PAO1Δ*pvdD*Δ*pchEF* showed significantly

reduced ability to remain in the host compared to the wildtype. In contrast, the

colonisation success of PAO1Δ*pchEF* and PAO1Δ*rhlR* was increased relative

to the wildtype. Grey shaded areas depict background fluorescence (mean +/-

standard deviation) of worms exposed to the non-fluorescent, non-pathogenic

E.coli OP50. N = number of worms form four independent experiments. * = p <

0.05; ** = p < 0.01, *** = p < 0.001.

Figure 5. Mixed infections reveal social strain dynamics but no

successful cheating. (A) Relative fitness of the wildtype PAO1-*mCherry* after

42 hours of competition inside the *C. elegans* gut against an untagged PAO1

control strain; the siderophore-negative strain PAO1Δ*pvdD*Δ*pchEF*; and the

Las-negative strain PAO1Δ*lasR*. The control competition revealed a mild but

significant negative effect of the mCherry tag on wildtype fitness. When

accounting for these mCherry costs, we found that the putative cheat strains

PAO1 Δ*lasR* and PAO1 Δ*pvdD*Δ*pchEF* performed equally well compared to

the wildtype, but could not outcompete it. This suggests that virulence factor

deficient strains benefit from the presence of non-producers but cannot

successfully cheat on them. (B) At 6 hpe, wildtype frequency in mixed

infections correlated positively with total bacterial load inside hosts in

competition with PAO1Δ*lasR* (squares and dashed lines) and

PAO1Δ*pvdD*Δ*pchEF* (diamonds and dotted lines) but not in the control

competition (circles and solid lines). These correlations disappeared at 48 hpe.

Each data point represents an individual worm. Data shown in A+B stem from

the same three independent experiments.

Figure 6. Spatial structure of mixed infections in the nematode gut. (A, B)

Illustrative examples of *C. elegans* individuals infected with a mixture of GFP-

and mCherry-labelled strains. Each worm was computationally straightened

and fluorescence intensity values were extracted for each pixel from tail (X=0)

to head (X=1). We then calculated the Spearman correlation coefficient ρ

between the intensity values in the two fluorescence channels across pixels,

as our estimate of strain colocalization. Examples show worms with high (A)

and low (B) degrees of colocalization. (C) Patterns of colocalization levels

varied substantially between individuals, but did not differ across strain

combinations (p = 0.119: wildtype PAO1-*mCherry* versus: (i) wildtype PAO1-

gfp (circles), (ii) PAO1Δ*pvdD*Δ*pchEF*-*mCherry* (diamonds) or (iii) PAO1Δ*lasR*-

mCherry (squares). Each data point represents an individual worm. Data

stems from 3 independent experiments, with 12 replicates each.

A Exposure — Host exposed to mCherry-tagged *P. aeruginosa* strains for 24 h

Nematodes washed with buffer + antibiotic

0 hpe
6 hpe
30 hpe

Nematodes immobilized (25 mM NaN₃)

Observation and measurements — Removal from exposure plate, collection in sterile buffer and microscopy measurement taken at 0, 6 and 30 hours post exposure (hpe).

B

C

D

A — Initial uptake of bacteria

B — Colonization ability

A

Fitness of PAO1−*mCherry* (ln(fitness))

n.s.

n.s.

N=25   N=24   N=26

PAO1   PAO1Δ*pvdD*Δ*pchEF*   PAO1Δ*lasR*

Competitor

B

Bacterial load in host gut (CFU)

Early colonization (6 hpe)   Late colonization (48 hpe)

r = 0.40   r = − 0.03   PAO1Δ*pvdD*Δ*pchEF*

r = 0.54   r = − 0.09   PAO1Δ*lasR*

r = 0.12   r = − 0.09   PAO1

Frequency of PAO1−*mCherry* in competition with

A – High co–localization, ρ = 0.903

B – Low co–localization, ρ = −0.04

C