1

Genetics of adaptation of the ascomycetous fungus

Podospora anserina to submerged cultivation

Olga A. Kudryavtseva[1,¶], Ksenia R. Safina[2,3,¶*], Olga A. Vakhrusheva[2,3], Maria D. Logacheva[2,4],

Aleksey A. Penin[3,4,5,6], Tatiana V. Neretina[4], Viktoria N. Moskalenko[7], Elena S. Glagoleva[8],

Georgii A. Bazykin[2,3], Alexey S. Kondrashov[9]

[1] Department of Mycology and Phycology, Faculty of Biology, Lomonosov Moscow State

University, Moscow, Russia

[2] Department of Life Sciences, Skolkovo Institute of Science and Technology, Skolkovo, Russia

[3] Sector of Molecular Evolution, Institute for Information Transmission Problems of the Russian

Academy of Sciences, Moscow, Russia

[4] Belozersky Institute of Physico-Chemical Biology, Lomonosov Moscow State University,

Moscow, Russia

[5] Department of Genetics, Faculty of Biology, Lomonosov Moscow State University, Moscow,

Russia

[6] Extreme Biology Laboratory, Institute of Fundamental Medicine and Biology, Kazan Federal

University, Kazan, Russia

[7] Zoological Museum, Lomonosov Moscow State University, Moscow, Russia

[8] Department of Plant Physiology, Faculty of Biology, Lomonosov Moscow State University,

Moscow, Russia

[9] Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor,

Michigan, USA

* Corresponding author

E-mail: k.r.safina@gmail.com

¶ These authors contributed equally to this work.

Abstract

Podospora anserina is a model ascomycetous fungus which shows pronounced phenotypic senescence when grown on solid medium but possesses unlimited lifespan under submerged cultivation. In order to study the genetic aspects of adaptation of *P. anserina* to submerged cultivation, we initiated a long-term evolution experiment. In the course of the first four years of the experiment, 125 single-nucleotide substitutions and 23 short indels were fixed in eight independently evolving populations. Six proteins that affect fungal growth and development evolved in more than one population; in particular, the G-protein alpha subunit FadA evolved in seven out of eight experimental populations. Parallel evolution at the level of genes and pathways, an excess of nonsense and missense substitutions, and an elevated conservation of proteins and their sites where the changes occurred suggest that many of the observed allele replacements were adaptive and driven by positive selection.

Author summary

Living beings adapt to novel conditions that are far from their original environments in different ways. Studying mechanisms of adaptation is crucial for our understanding of evolution. The object of our interest is a multicellular fungus *Podospora anserina*. This fungus is known for its pronounced senescence and a definite lifespan, but it demonstrates an unlimited lifespan and no signs of senescence when grown under submerged conditions. Soon after transition to submerged cultivation, the rate of growth of *P. anserina* increases and its pigmentation changes. We wanted to find out whether there are any genetic changes that contribute to adaptation of *P. anserina* to these novel conditions and initiated a long-term evolutionary experiment on eight

independent populations. Over the first four years of the experiment, 148 mutations were fixed in

these populations. Many of these mutations lead to inactivation of the part of the developmental

pathway in *P. anserina*, probably reallocating resources to vegetative proliferation in liquid

medium. Our observations imply that strong positive selection drives changes in at least some of

the affected protein-coding genes.

Introduction

Over the last several decades, experimental evolution became an essential part of

evolutionary biology, providing an opportunity to observe and study changes that occur in

controlled populations in the course of many generations. In particular, the *E. coli* long-term

evolution experiment led by Richard Lenski since 1988 yielded a number of important results,

including that on morphological evolution of cells (1,2), long-term fitness gain under a constant

environment (1,3), and ecological specialization (4,5). The existence of twelve replicas made it

possible to study changes in genotypes and phenotypes that occurred in parallel in independently

evolving bacterial populations (6–8).

Being easily cultured and fast-growing, *E. coli* serves as a convenient model organism for

experimental evolution. Still, the range of questions that can be studied on it and on other

prokaryotes is limited by their relative simplicity. Eukaryotes such as protozoans or fungi

possess more complex gene regulation, morphology and physiology and have also been used in

the studies of experimental evolution (9–12).

Podospora anserina is a model ascomycete fungus that possesses a definite lifespan with

pronounced phenotypic senescence when grown on solid medium (13). In contrast, under

conditions of submerged cultivation, *P. anserina* displays unlimited lifespan with no signs of

senescence (14,15). Isolates taken from such cultures revert to senescence, indicating that the

unlimited lifespan is not associated with genetic changes, instead being a product of epigenetic

changes or environmental influences. Still, such isolates demonstrate increased lifespan when

plated on solid medium and become sterile, only rarely forming mostly nonfunctional

microconidia and protoperithecia (14,15). After some period of submerged cultivation,

P. anserina cultures acquire higher rates of biomass accumulation and change their

pigmentation. The mechanisms that alter *P. anserina* senescence program and affect its

phenotype under such conditions are unknown.

In order to study the genetic aspects of adaptation that takes place during the submerged

cultivation of *P. anserina*, we started in 2012 a long-term evolutionary experiment. We used two

wild strains of *P. anserina* to establish eight independent experimental populations. Here, we

analyse the genetic changes that took place in the first four years of this experiment.

Results

Allele replacements

We used two founder genotypes, denoted A and B, to establish eight replicate

experimental populations (five from A and three from B, referred to as A populations and B

populations). Populations were cultivated via serial passages in the standard synthetic medium

M2 as described in Materials and Methods. After the short period of initial adaptation to the

conditions of the submerged cultivation, all the populations displayed identical phenotypic

changes, including changes in colony morphology and pigmentation (15). In order to identify

genetic changes acquired by the populations during the course of the experiment, we sequenced

both the founder genotypes and each of the populations at four time points (Fig 1).

The summary of the observed allele replacements is provided in Table 1 and their full list

is presented in Table S1. Our set of allele replacements comprises 85 (73% transitions) and 40

(53% transitions) nucleotide substitutions and 13 and 10 indels in A and B populations,

respectively. Among the 148 identified allele replacements, 52 result in an amino acid

substitution, 11 are single-nucleotide substitutions leading to a premature inframe stop codon,

and 13 cause a frameshift, comprising 35.1%, 7.4% and 8.8% of all allele replacements,

respectively.

Table 1. Summary of allele replacements that occurred in the experimental populations.

Type of a replacement	Experimental populations A	Experimental populations B
Total	98	50
By mutation type (single nucleotide mutations):		
Transition	62	21
Transversion	23	19
By mutation effect:		
Missense nucleotide substitutions	31	21
Synonymous nucleotide substitutions	6	7
Nonsense nucleotide substitutions	8	3
Nucleotide substitutions within noncoding regions	26	5
Nucleotide substitutions within genome regions that cannot be aligned with the reference genome	9	4
Nucleotide substitutions within introns	3	0

Nucleotide substitutions in incomplete proteins	2	0
Short insertions/deletions (including frameshifts)	13 (9)	10 (4)

The dynamics of accumulation of allele replacements is summarized in Fig 2 and Table

S2. There are no obvious deviations from their uniform rate in the course of the experiments

(linear regression's time variable p-value is 0.0613). Furthermore, there were no mutators: the

numbers of the mutations in each population were all drawn from the same barrel and there were

no difference between A and B populations (two-way ANOVA test, see Table S2).

Distribution of allele replacements within the genome

We then focused on the analysis of single nucleotide substitutions and first computed dN/

dS ratios for populations A and B. dN/dS ratio equals 1.77 and 1.11 for populations A and B,

respectively (Table S3). While dN/dS > 1 is commonly thought to indicate positive selection, our

estimates may be unreliable because of the scarcity of the data. We next investigated how allele

replacements are distributed in the *P. anserina* genome. We produced null-distributions for the

number of missense and nonsense substitutions, using permutation procedure (see Materials and

Methods). Null-distributions were obtained separately for A and B populations, assuming that

mutations strike the genome and become fixed at random, with the uniform per site rates. There

are significantly more nonsense substitutions than expected by chance in both A and B

populations, and significantly more missense substitutions in B populations (Fig 3).

Moreover, despite the fact that there were very few mutations overall, a substantial

proportion of protein-altering allele replacements affected the same gene and protein in more

than one population. Totally, six proteins were subject to such parallel evolution (Table 2). The

probability of observing by chance the same or a higher number of genes carrying just two

single-nucleotide protein-altering substitutions by chance is very low (Fig 4), and four out of six

proteins that underwent parallel evolution accepted more than two allele replacements. Clearly,

positive natural selection was responsible for this phenomenon.

Table 2. Proteins affected independently in more than one population.

Protein identifier and its function	The number of independent allele replacements
Pa_1_10140, Putative transcriptional regulatory protein pro-1	2 nonsense substitutions
Pa_1_23950, Guanine nucleotide-binding protein alpha subunit encoded by the mod-D gene	1 nonsense substitutions, 2 missense mutations
Pa_3_400, putative protein with unknown function	6 frameshifts
Pa_3_6550, Putative protein homologous to VeA of *Aspergillus nidulans*	2 missense substitutions
Pa_6_3340, Putative developmental regulator flbA	2 nonsense substitutions, 1 missense substitutions
Pa_7_7970, Putative guanine nucleotide-binding protein alpha-1 subunit	7 missense substitutions

This parallelism cannot be an artefact of ancestral polymorphism segregating within our

lineages, since we conservatively excluded a number of positions with identical parallel

mutations in multiple populations founded from the same ancestral lineage. For the same reason,

we could lose some actual functional parallelism, making our estimates for the frequency of

parallel substitutions conservative. We did observe two identical nucleotide mutations between

two lineages formed from two different ancestral lineages (Table S1); those are unlikely to be

due to ancestral polymorphism.

Proteins carrying missense and nonsense substitutions are more conservative

To study the rate of long-term evolution of proteins that evolved in our experiment, we

searched for pairs of orthologs pairs encoded by *P. anserina* and *Neurospora crassa* genomes.

Out of 10230 proteins in the reference *P. anserina* genome, 4713 have an ortholog in *N. crassa*

genome, such that their alignments contain no more than 20% of gaps. Both nonsense and, to a

lesser extent, missense allele replacements preferentially occurred in those *P. anserina* proteins

that have reliable orthologs in *N. crassa* (Table 3).

Moreover, missense allele replacements within such proteins preferentially occurred at

amino acid sites conserved between the two species. While the overall proportion of such sites

for 4713 alignments of reliable orthologs is 62%, 87% (27 out of 31) of replaced amino acids

occupied conservative sites (p = 0.003).

Table 3. Statistics of orthologization of *P. anserina* и *N. crassa* genomes.

	Total # of proteins	# of proteins carrying	# of proteins carrying missense

	nonsense substitutions	substitutions
# of proteins in *P. anserina* genome	10230	9	44
# of such proteins that have a reliable ortholog in *N. crassa*	4713	9 (0.000940)	27 (0.048609)

Values specified in parentheses correspond to p-values of Fisher's exact test

Possible phenotypic effects of parallel allele replacements

Proteins that underwent allele replacements in more than one experimental population

(Table 2) are of particular interest. Indeed, their evolution must have been adaptive (Fig 4) and,

therefore, must have led to changes in the phenotype. Moreover, they probably affected the

protein structures radically. Indeed, the differences in physicochemical properties (computed as

Miyata's distance) for pairs of amino acids were higher for missense substitutions that occurred

during our experiment than for those that occurred in the course of divergence between

P. anserina and *N. crassa* (mean values 1.979811 vs. 1.472333, two-tail t-test p-value 0.002).

Let us discuss what these changes could be.

The most striking example of parallel evolution is protein Pa_7_7970 which experienced

nucleotide substitutions in seven independent populations out of eight. Pa_7_7970 is a putative

alpha-1 subunit of a guanine nucleotide-binding protein (Table 2). There are three classes of G

protein alpha subunits in ascomycetes fungi (16), each affecting fungal growth and development

differently. Pa_7_7970 belongs to class 1 and is 93% identical to its well-studied orthologous

protein FadA of *Aspergillus nidulans*. FadA stimulates vegetative growth (17) and probably

suppresses sexual reproduction (18), which agrees well with intensive vegetative growth during

the experiment suggesting that constitutive activation of Pa_7_7970 may be adaptive under the

conditions of submerged cultivation.

Only four sites in Pa_7_7970 evolved in the course of the experiment: arg-178 was

substituted in three experimental populations, gly-183 was substituted twice, and ser-206 and

lys-210 were each substituted once. All these sites have already been studied. Arg-178 is a

catalytic residue crucial for GTP hydrolysis and alpha subunit inactivation (19) and its mutations

result in constitutive activation of alpha subunit (20). Gly-183 takes part in the interaction

between alpha subunit and regulator of G protein signalling (RGS) that activates GTP hydrolysis,

and Gly->Ser substitution (observed in one case) hinders this process drastically (21). Ser-206

and Lys-210 are also involved in the interaction with RGS (22,23) although no specific effects of

their mutations have yet been described.

Remarkably, we also observed one missense and two nonsense substitutions in the

ortholog of RGS (Pa_6_3340) which belongs to the same pathway as FadA and likely

contributes to *P. anserina* adaptation in similar fashion.

Another protein that is worth mentioning is a transcription factor pro-1 (Pa_1_10140)

which acquired two loss-of-function allele replacements. pro-1 interacts with promoters of many

developmental genes including *esdC* gene that stimulates sexual development and is suppressed

by FadA (22,23), and deletion of pro-1 results in sexual sterility. Since *P. anserina* proliferates

only vegetatively during the experiment, defects in sexual development pathway may be

beneficial.

We observed two missense substitutions in VeA homolog (Pa_3_6550). Although we

cannot suggest the effect of two missense substitutions on protein function, VeA in *A. nidulans*

positively regulates expression of the previously mentioned *esdC* gene (18) and might affect *P.*

anserina metabolism in some way.

We also observed one protein with a surprising amount of parallel evolution via

frameshifts. Pa_3_400 protein experienced frameshift in six independent populations. Its

homologs in *N. crassa* (24) and *A. nidulans* are involved in asexual reproduction (25). As

P. anserina lacks asexual reproduction, we cannot suggest the role of Pa_3_400 inactivation in

P. anserina adaptation, although in *P. anserina* it can be involved in other developmental

pathways.

As for Pa_1_23950, it belongs to class 3 alpha subunits of fungal G proteins. Mutations

in this protein cause defects in vegetative proliferation in *P. anserina* (26) making it difficult to

interpret the observed substitutions in this protein in the context of adaptation.

All proteins discussed above are involved in growth and development processes in

various fungi. Taken with the excess of coding substitutions and the number of protein changed

in parallel, it favors the possible adaptiveness of some of the observed allele replacements

toward the conditions of submerged cultivation. Yet mutant construction is needed to claim it

with certainty.

Discussion

In populations of *P. anserina*, transition to prolonged submerged cultivation inevitably

results in the same set of morphophysiological changes (14). These changes always occur early

in the course of an experiment (15) and therefore are most likely caused by modifications of

epigenetic patterns that alter gene expression. Yet at a longer timescale, one can also expect

P. anserina to adapt to the conditions that are far from its native environment through fixations

of newly acquired mutations.

We analysed small-scale changes that occurred in the *P. anserina* genome in the course

of 268 experimental passages. We observed and confirmed the appearance of 125 single-

nucleotide substitutions and 23 short indels. It is hard to tell which of these mutations are

advantageous for submerged cultivation. Experimental populations of *P. anserina* cannot

reproduce sexually under conditions of the experiment, and *P. anserina* genus itself lacks the

ability to form asexual reproductive structures, macroconidia. Thus, the propagation of our

experimental populations occurred exclusively through vegetative proliferation, so that they must

possess clonal population structure. Thus, hitch-hiking of neutral and even slightly deleterious

mutations that arose in high-fitness genotypes destined to fixation was likely common (27).

Nevertheless, two observations clearly indicate that positive selection drove many of the

observed allele replacements which, therefore, contributed to adaptation to the experimental

environment.

First, evolution of our eight experimental populations often occurred in parallel, with the

same gene and protein being affected by a mutation up to seven times (Table 2). Parallel

evolution of phenotypes is common in nature, although it does not necessarily involve parallel

genetic changes (Steiner et al. 2009). In contrast, in evolutionary experiments parallel genetic

changes have being reported several times (28–31). Parallelisms at nucleotide level are rare, and

we observed only two such events in Pa_7_7970 gene, which might be mutational hotspots but

considering the importance of the mutated protein sites clearly indicate their adaptive role.

Some of the six proteins that underwent parallel changes in our experiment are involved

in the same pathways, and all of them seem to be associated with growth and development

processes in fungi. A prominent role of nonsense and frameshift alleles, as well as the nature of

the fixed missense alleles, indicate that this parallel evolution mostly, if not exclusively, led to

inactivation of the affected proteins. At least five missense substitutions in Pa_7_7970, as well as

nonsense substitutions in Pa_6_3340, alter developmental pathways probably stimulating

vegetative growth and inhibiting sexual reproduction at a very early stage. A similar pattern was

observed in experimental yeast populations, where parallel evolution affected genes of the

mating pathway and small GTPases of the Ras protein family (29). Loss-of-function mutations of

large effects can provide a selective advantage inactivating parts of regulatory/metabolic network

that became useless under the current environmental conditions (32). For example, in yeasts

mutations resulting in sexual sterility provide advantage in growth rate (33).

Still, the extent of parallel evolution in our experiment is remarkable. Fungi tend to not

differentiate in liquid medium (34), so that sexual reproduction was already suppressed before

experimental evolution ever begun. Thus, here the situation is very different from parallel loss of

eyes in many cave populations of *Astyanax mexicana*, where the cost of maintaining even a

useless capacity to see was estimated as up to 15% of resting metabolism (35), which makes its

elimination obviously beneficial. In contrast, we observed that positive selection strongly

favored mutations that damage a top-level metabolic pathway which already lacks a salient

phenotypic expression. Perhaps, before loss-of-function mutations in regulatory proteins occur,

some cryptic activity of this pathway remains, and fixation of such mutations allows cells to

reallocate their resources to vegetative proliferation. Thus, the role of positive selection in

degenerative evolution may be even more prominent than thought previously.

The second reason to believe that many of the observed allele replacements were

adaptive is overrepresentation among them of nonsense and missense mutations which must have

disproportionally strong impact on function (Fig 3). Moreover, among missense mutations, those

that involve radical amino acid replacements and those that occurred at conservative sites and in

more conservative proteins are also more common than expected. Because such mutations have a

reduced chance of being effectively neutral, this pattern implies that positive selection drove

most of their fixations.

Surprisingly, there is still no clarity regarding the role of functionally radical mutations in

the adaptive evolution in nature. The radical-conservative ratio of missense substitutions has

been studied in the context of both positive and negative selection, and while there are some

hints that it correlates with the strength of negative selection (36–38), its relation to positive

selection remains ambiguous. Radical-conservative ratio has been proposed as a measure to

detect positive selection almost three decades ago (39). However, this measure is vulnerable to

various factors not related to selection (40) and studies of different species did not find enough

evidences for the excess of radical compared to conservative substitutions at positively selected

sites (36,41,42).

A prominent role of mutations at conservative sites and proteins in adaptation to the

experimental environment was reported for *E. coli* (43). An obvious possibility is that these, as

well as ours, data reflect how adaptation occurs in nature. Indeed, substitutions at conservative

sites that occur in the course of natural evolution are enriched with those driven by positive

selection (44). Alternatively, a disproportional contribution of radical mutations to experimental

evolution may also be due to simplified laboratory environments, under which adaptation mostly

involves shedding functions that are no longer needed. Clearly, this issue deserves to be studied

further.

At the moment, there is no obvious trend in mutation accumulation rate. Experimental

populations acquired two times less mutations between time points 3 and 4 than they did in the

previous time interval. However, the reason for that is yet unclear, and further observations

might shed some light on this tendency. The experiment lasts for six years already, and to our

knowledge, there are not much long-lasting evolution experiments conducted on multicellular

organisms. We are planning to continue the experiment and analyse changes in experimental

populations of *P. anserina*.

Material and Methods

Evolutionary experiment

Two homokaryotic vegetatively incompatible wild strains *s* and *S* (denoted here as A and B) were kindly provided by Annie Sainsard-Chanet and Carole H. Sellem (Département Biologie Cellulaire et Intégrative, Centre de Génétique Moléculaire, CNRS, Gif-sur-Yvette Cedex, France) and gave rise to five and three independent experimental populations, respectively (Fig 1). Experimental populations were maintained via serial passages and were grown in the dark on the standard synthetic medium M2 (15). Submerged cultivation was carried out in 750ml Erlenmeyer flasks with 100 ml of the medium at $27 \pm 1°C$ on a 200rpm rotor shaker. Cultures were transferred to fresh medium every 4-7 days. During the experiment, mycelium samples were plated on agarized (20 g/l) M2 medium in Petri dishes. Initial founder strains are stored on ararized medium.

We denoted 64, 130, 200, and 268 passages as the first, the second, the third, and the fourth time points of the experiment, respectively. Whole-genome sequencing was performed for both the founder populations, for populations B at the points 1, 2, and 4, and for populations A only at the point 4. The allele replacements that were observed from whole genome sequencing data were subsequently verified by Sanger sequencing (see below). Several samples of passage 64 were lost due to contamination; for that reason, we substituted passage 64 with the closest available passage 75 and refer to passage 75 as the first time point in the Results section. The scheme of the experiment is shown in Fig 1.

Whole-genome sequencing

DNA was extracted using CTAB-based method (45). For library preparation for the first

series of samples (founder genotype B and experimental populations B at the first time point), we

used Truseq DNA sample preparation kit (Illumina), for the second series of samples

(experimental populations B at the second time point) we used NEBNext Ultra DNA kit (New

England Biolabs) according to manufacturer instructions. For the third series (founder genotype

A and all eight experimental populations at the fourth time point), we constructed PCRfree

libraries using Accel-NGS 2S PCR-Free DNA Library Kit (Swift Biosciences). The first and

second series were sequenced on HiSeq2000 (Illumina) using Truseq v.3 reagents. The third

series was sequenced using Nextseq500 and v.2 reagents.

De novo genome assembly

Pair-end reads from HiSeq (100bp) and NextSeq (150bp) were trimmed using

Trimmomatic (46) with options (ILLUMINACLIP:adapters:2:30:10 LEADING:7 TRAILING:7

SLIDINGWINDOW:5:10 MINLEN:50). Duplicate reads were excluded using fastq-mcf (with -

D 50 option).

We performed *de novo* genome assembly of both founder genotypes with SPAdes (47)

(with -only-assembler option). The initial assemblies were then filtered using the published

reference genome (48), v. 6.31, last accessed ~ Dec 2014). We used blastn (49) to find the best

hits for each contig in published genome and kept only those contigs that had a local alignment

of at least 5kb with published genome with identity of at least 95%. Finally, we performed

correction of assemblies using ICORN2 (50). Statistics on assemblies is provided in Table S4.

Variant calling

For each of the sequenced samples, we mapped its pair-end reads onto the corresponding

founder genome using bwa (51) and removed reads that had non-unique alignments. Filtered

mappings were used for variant calling. We used samtools (52) to call SNPs and InDels. The

resulting calls were filtered by coverage (>=10) and variant frequency (>=80%) to produce high-

quality variant set for each of the experimental populations. We then excluded non-unique

variants that were supported by at least two reads in more than one experimental population due

to extremely low probability of mutations affecting the same base pair in several independent

populations. We further masked low complexity regions of founder genomes by repeatmasker

(53) and excluded indels that fell into the masked sites and adjacent regions ten base pairs around

them. In populations A4 and B2, we observed a couple of mutations in adjacent genomic sites;

we excluded these variants as probable dinucleotide mutations.

Sanger sequencing

Sanger sequencing was used to check the presence of the called variants (138 SNPs and

24 indels) at each of the four time points. Several samples of passage 64 were lost due to

Penicillium contamination, so we used the closest available passage 75 as the first time point

passage. As each of the mutations observed at earlier time points in B populations were always

present in later populations, we considered a mutation as verified if it was confirmed by Sanger

sequencing in any of four time points (assuming that the mutation that appeared once became

fixed at later time points). For B populations, we believed in NGS data for passages 64 and 130

if the mutation was identified by Sanger sequencing at any time point. For A populations, we had

to check each of the four time points.

Primers for Sanger sequencing were designed using primer3 (54,55) and tntblast (56)

software and synthesized in Evrogen. PCR was performed using Encyclo kit (Evrogen).

Amplification products were sequenced in both directions on ABI Prism 3500 sequencer

(Applied Biosystems). Chromatograms were validated manually in CodonCode Aligner (57).

The detailed results of Sanger sequencing are included in Table S1. The number of

confirmed mutations is provided in Table S2. We were unable to confirm neither the presence

nor the absence of 15 mutations and excluded them from any further analyses. The majority of

mutations that we failed to validate comprise mutations in noncoding/not annotated regions.

Also, there were three mutations in A populations for which we couldn't track the time of their

appearance.

Variant annotation

We mapped reference CDS sequences (48) (v. 6.31, last accessed ~ Dec 2014) onto the

reference genome and obtained coding regions coordinates using Splign (58). Out of 10,635

reference protein-coding genes, we were able to identify 10230 genes in the reference genome

that coded complete proteins with proper start and stop codons. We then performed pairwise

genome alignment of A and B founder genotype assemblies with published genome using Multiz

(59) and identified coding regions coordinates in the assembled genomes using Splign annotation

results. 9732 and 9732 (numbers are indeed the same but protein sets differ) complete genes

were identified in A and B founder genotypes, respectively, with a tiny amount of incomplete

genes (627 and 530 genes for A and B populations, respectively). Based on the produced

annotations, the identified genomic variants were classified as exonic, intronic or intergenic.

Exonic variants in complete genes (i.e., genes with all of their exons completely identified in the

founder genome) were further classified as missense, nonsense or silent mutations in case of

SNPs and frameshift or amino acid insertion/deletion mutations in case of indels.

Permutations

In order to estimate the probability to obtain the observed number of protein-altering

variants by chance, we generated null distributions for the number of missense and nonsense

mutations in A and B founder genomes using permutations. A permutation procedure was

performed as follows. 85 and 40 observed SNP mutations were thrown at random onto A and B

genomes, respectively, and annotated as described above. To control for possible confounding

effects of mutation context, we added three-nucleotide sequence contexts to permutations. The

procedure was repeated 10,000 times and produced null distributions for the expected number of

nonsense and missense mutations in each of the founder genomes (Fig 3). To obtain the expected

number of proteins evolving in parallel, we combined permutation results from A and B

populations and counted the number of proteins that acquired at least two single-nucleotide

protein-changing mutations in eight populations during the permutation procedure.

Protein conservation analysis

We used OrthoMCL (60) to identify orthologous groups between protein sets from

P. anserina and *N. crassa* (61) reference genomes. Orthologous protein pairs were then aligned

using MAFFT (62) (with -localpair option), and for each protein carrying ortholog in *N. crassa*

genome, we calculated its conservation as a fraction of conserved positions in the alignment. We

also measured the conservation of mutated positions as a fraction of conserved positions among

the whole set of mutated positions.

References

1.   Lenski RE, Travisano M. Dynamics of adaptation and diversification: a 10,000-generation

experiment with bacterial populations. Proc Natl Acad Sci U S A. 1994 Jul 19;91(15):6808–14.

2.   Philippe N, Pelosi L, Lenski RE, Schneider D. Evolution of penicillin-binding protein 2

concentration and cell shape during a long-term experiment with Escherichia coli. J Bacteriol. 2009

Feb;191(3):909–21.

3.   Lenski RE, Wiser MJ, Ribeck N, Blount ZD, Nahum JR, Morris JJ, et al. Sustained fitness gains and

variability in fitness trajectories in the long-term evolution experiment with Escherichia coli

[Internet]. 2015. Available from: http://dx.doi.org/10.1101/027391

4.   Blount ZD, Barrick JE, Davidson CJ, Lenski RE. Genomic analysis of a key innovation in an

experimental Escherichia coli population. Nature. 2012;489(7417):513–8.

5.   Leiby N, Marx CJ. Metabolic Erosion Primarily Through Mutation Accumulation, and Not

Tradeoffs, Drives Limited Evolution of Substrate Specificity in Escherichia coli. PLoS Biol.

2014;12(2):e1001789.

6.   Lenski RE, Travisano M. Dynamics of adaptation and diversification: a 10,000-generation

experiment with bacterial populations. Proc Natl Acad Sci U S A. 1994 Jul 19;91(15):6808–14.

7.   Woods R, Schneider D, Winkworth CL, Riley MA, Lenski RE. Tests of parallel molecular evolution

in a long-term experiment with Escherichia coli. Proceedings of the National Academy of Sciences.

2006;103(24):9107–12.

8.   Cooper TF, Remold SK, Lenski RE, Schneider D. Expression profiles reveal parallel evolution of

epistatic interactions involving the CRP regulon in Escherichia coli. PLoS Genet. 2008 Feb;4(2):e35.

9.   Gray JC, Cutter AD. Mainstreaming Caenorhabditis elegans in experimental evolution. Proc Biol

Sci. 2014 Mar 7;281(1778):20133055.

10. McDonald MJ, Rice DP, Desai MM. Sex speeds adaptation by altering the dynamics of molecular

evolution. Nature. 2016 Mar 10;531(7593):233–6.

11. Rundle HD, Chenoweth SF, Doughty P, Blows MW. Divergent selection and the evolution of signal

traits and mating preferences. PLoS Biol. 2005 Nov;3(11):e368.

12. Dettman JR, Anderson JB, Kohn LM. Divergent adaptation promotes reproductive isolation among

experimental populations of the filamentous fungus Neurospora. BMC Evol Biol. 2008 Jan 31;8:35.

13. Lorin S, Dufour E, Sainsard-Chanet A. Mitochondrial metabolism and aging in the filamentous

fungus Podospora anserina. Biochimica et Biophysica Acta (BBA) - Bioenergetics. 2006;1757(5-

6):604–10.

14. Turker MS, Cummings DJ. Podospora anserina does not senesce when serially passaged in liquid

culture. J Bacteriol. 1987 Feb;169(2):454–60.

15. Kudryavtseva OA, Mazheika IS, Solovchenko AE, Kamzolkina OV. Genetic instability of the short-

living ascomycetous fungus Podospora anserina induced by prolonged submerged cultivation.

Microbiology. 2011;80(6):784–96.

16. Bölker M. Sex and crime: heterotrimeric G proteins in fungal mating and pathogenesis. Fungal

Genet Biol. 1998 Dec;25(3):143–56.

17. Krijgsheld P, Bleichrodt R, van Veluw GJ, Wang F, Müller WH, Dijksterhuis J, et al. Development

in Aspergillus. Stud Mycol. 2013 Mar 15;74(1):1–29.

18. Han K-H, Kim JH, Moon H, Kim S, Lee S-S, Han D-M, et al. The Aspergillus nidulans esdC (early

sexual development) gene is necessary for sexual development and is controlled by veA and a

heterotrimeric G protein. Fungal Genet Biol. 2008;45(3):310–8.

19. Coleman DE, Berghuis AM, Lee E, Linder ME, Gilman AG, Sprang SR. Structures of active

conformations of Gi alpha 1 and the mechanism of GTP hydrolysis. Science. 1994 Sep

2;265(5177):1405–12.

20. Landis CA, Masters SB, Spada A, Pace AM, Bourne HR, Vallar L. GTPase inhibiting mutations

activate the α chain of Gs and stimulate adenylyl cyclase in human pituitary tumours. Nature.

1989;340(6236):692–6.

21. Lan KL, Sarvazyan NA, Taussig R, Mackenzie RG, DiBello PR, Dohlman HG, et al. A point

mutation in Galphao and Galphai1 blocks interaction with regulator of G protein signaling proteins. J

Biol Chem. 1998 May 22;273(21):12794–7.

22. Khafizov K, Lattanzi G, Carloni P. G protein inactive and active forms investigated by simulation

methods. Proteins. 2009 Jun;75(4):919–30.

23. Clark MA, Lambert NA. Endogenous regulator of G-protein signaling proteins regulate the kinetics

of Galphaq/11-mediated modulation of ion channels in central nervous system neurons. Mol

Pharmacol. 2006 Apr;69(4):1280–7.

24. Springer ML, Yanofsky C. A morphological and genetic analysis of conidiophore development in

Neurospora crassa. Genes Dev. 1989 Apr;3(4):559–71.

25. Chung D-W, Greenwald C, Upadhyay S, Ding S, Wilkinson HH, Ebbole DJ, et al. acon-3, the

Neurospora crassa ortholog of the developmental modifier, medA, complements the conidiation

defect of the Aspergillus nidulans mutant. Fungal Genet Biol. 2011 Apr;48(4):370–6.

26. Loubradou G, Bégueret J, Turcq B. MOD-D, a Galpha subunit of the fungus Podospora anserina, is

involved in both regulation of development and vegetative incompatibility. Genetics. 1999

Jun;152(2):519–28.

27. Barton NH. Genetic hitchhiking. Philos Trans R Soc Lond B Biol Sci. 2000 Nov

29;355(1403):1553–62.

28. Woods R, Schneider D, Winkworth CL, Riley MA, Lenski RE. Tests of parallel molecular evolution

in a long-term experiment with Escherichia coli. Proceedings of the National Academy of Sciences.

2006;103(24):9107–12.

29. Lang GI, Rice DP, Hickman MJ, Sodergren E, Weinstock GM, Botstein D, et al. Pervasive genetic

hitchhiking and clonal interference in forty evolving yeast populations. Nature. 2013 Aug

29;500(7464):571–4.

30. Wichman HA, Badgett MR, Scott LA, Boulianne CM, Bull JJ. Different trajectories of parallel

evolution during viral adaptation. Science. 1999 Jul 16;285(5426):422–4.

31. Lang GI, Desai MM. The spectrum of adaptive mutations in experimental evolution. Genomics.

2014 Dec;104(6 Pt A):412–6.

32. Hottes AK, Freddolino PL, Khare A, Donnell ZN, Liu JC, Tavazoie S. Bacterial Adaptation through

Loss of Function. PLoS Genet. 2013;9(7):e1003617.

33. Lang GI, Murray AW, Botstein D. The cost of gene expression underlies a fitness trade-off in yeast.

Proc Natl Acad Sci U S A. 2009 Apr 7;106(14):5755–60.

34.   Osiewacz HD. Molecular Biology of Fungal Development. CRC Press; 2002. 608 p.

35.   Moran D, Softley R, Warrant EJ. The energetic cost of vision and the evolution of eyeless Mexican

cavefish. Sci Adv. 2015 Sep;1(8):e1500363.

36.   Smith NGC. Are radical and conservative substitution rates useful statistics in molecular evolution?

J Mol Evol. 2003 Oct;57(4):467–78.

37.   Wernegreen JJ. Reduced selective constraint in endosymbionts: elevation in radical amino acid

replacements occurs genome-wide. PLoS One. 2011 Dec 14;6(12):e28905.

38.   Popadin K, Polishchuk LV, Mamirova L, Knorre D, Gunbin K. Accumulation of slightly deleterious

mutations in mitochondrial protein-coding genes of large versus small mammals. Proc Natl Acad Sci

U S A. 2007 Aug 14;104(33):13390–5.

39.   Hughes AL, Ota T, Nei M. Positive Darwinian selection promotes charge profile diversity in the

antigen-binding cleft of class I major-histocompatibility-complex molecules. Mol Biol Evol. 1990

Nov;7(6):515–24.

40.   Dagan T, Talmor Y, Graur D. Ratios of Radical to Conservative Amino Acid Replacement are

Affected by Mutational and Compositional Factors and May Not Be Indicative of Positive

Darwinian Selection. Mol Biol Evol. 2002;19(7):1022–5.

41.   Hanada K, Gojobori T, Li W-H. Radical amino acid change versus positive selection in the evolution

of viral envelope proteins. Gene. 2006 Dec 30;385:83–8.

42.   Suzuki Y. Inferring natural selection operating on conservative and radical substitution at single

amino acid sites. Genes Genet Syst. 2007 Aug;82(4):341–60.

43.   Maddamsetti R, Hatcher PJ, Green AG, Williams BL, Marks DS, Lenski RE. Core Genes Evolve

Rapidly in the Long-term Evolution Experiment with Escherichia coli. Genome Biol Evol [Internet].

2017 Apr 4; Available from: http://dx.doi.org/10.1093/gbe/evx064

44.   Bazykin GA, Kondrashov AS. Major role of positive selection in the evolution of conservative

segments of Drosophila proteins. Proc Biol Sci. 2012 Sep 7;279(1742):3409–17.

45.   Doyle JJ, Doyle JL. A rapid DNA isolation procedure for small quantities of fresh leaf tissue.

Phytochemical Bulletin. 1987;19:11–5.

46. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.

Bioinformatics. 2014 Aug 1;30(15):2114–20.

47. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a new

genome assembly algorithm and its applications to single-cell sequencing. J Comput Biol. 2012

May;19(5):455–77.

48. Podospora anserina Genome Project [Internet]. [cited 2018 Jun 16]. Available from:

http://podospora.i2bc.paris-saclay.fr/download.php

49. Altschul S. Basic Local Alignment Search Tool. J Mol Biol. 1990;215(3):403–10.

50. Otto TD, Sanders M, Berriman M, Newbold C. Iterative Correction of Reference Nucleotides

(iCORN) using second generation sequencing technology. Bioinformatics. 2010 Jul 15;26(14):1704–

7.

51. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics. 2009 Jul 15;25(14):1754–60.

52. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Alignment/Map

format and SAMtools. Bioinformatics. 2009 Aug 15;25(16):2078–9.

53. RepeatMasker Home Page [Internet]. [cited 2018 Jun 16]. Available from:

http://www.repeatmasker.org

54. Untergasser A, Cutcutache I, Koressaar T, Ye J, Faircloth BC, Remm M, et al. Primer3—new

capabilities and interfaces. Nucleic Acids Res. 2012;40(15):e115–e115.

55. Koressaar T, Remm M. Enhancements and modifications of primer design program Primer3.

Bioinformatics. 2007 May 15;23(10):1289–91.

56. Gans JD, Wolinsky M. Improved assay-dependent searching of nucleic acid sequence databases.

Nucleic Acids Res. 2008 Jul;36(12):e74.

57. Website [Internet]. [cited 2018 Jun 16]. Available from: www.codoncode.com

58. Kapustin Y, Souvorov A, Tatusova T, Lipman D. Splign: algorithms for computing spliced

alignments with identification of paralogs. Biol Direct. 2008;3(1):20.

59. Blanchette M, Kent WJ, Riemer C, Elnitski L, Smit AFA, Roskin KM, et al. Aligning multiple

genomic sequences with the threaded blockset aligner. Genome Res. 2004 Apr;14(4):708–15.

60. Li L. OrthoMCL: Identification of Ortholog Groups for Eukaryotic Genomes. Genome Res.

2003;13(9):2178–89.

61. Krumlau R, Marzluf GA. Characterization of the sequence complexity and organization of the

Neurospora crassa genome. Biochemistry. 1979;18(17):3705–13.

62. Katoh K. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier

transform. Nucleic Acids Res. 2002;30(14):3059–66.

Figures

Fig 1. Scheme of the experiment.

Eight experimental populations were established from two founder lineages A and B. After 75,

130, 200, and 268 passages, we determined genotypes of the experimental populations (dashed

lines; see Methods).

Fig 2. Accumulation of allele replacements during the experiment.

Each dot reflects a number of replacements observed in a population at a particular time point.

Populations are shown as thin lines. Thick blue line corresponds to the linear regression line with

confidence interval depicted as shaded area.

Fig 3. The observed number of missense and nonsense substitutions is greater than

expected by chance.

Distributions of the expected number of nonsense (A) and missense (B) substitutions in A (left

panel) and B (right panel) experimental populations in 10,000 permutations. Red line

corresponds to the observed number of nonsense or missense substitutions on (A) and (B),

respectively. *p* (p-value) is the area to the right of the red line.

Fig 4. The excess of the number of proteins with parallel changes.

Distribution of the number of genes with two protein-altering single-nucleotide substitutions in

10,000 random permutations. Red line shows the observed number of genes with two or more

such substitutions occurred.

Supplementary Information

Table S1. The summary of the observed allele replacements. Sheets A1-A5 and B1-B3

contain the results of the Sanger sequencing validation procedure. The two last sheets comprise

annotation of the allele replacements in A and B populations.

Table S2. Accumulation of allele replacements. Each entry provides the number of newly

observed replacements, relatively to the previous time point.

Table S3. dN/dS calculation.

Table S4. Statistics on founder genotypes assemblies.

Figure 1

Figure 2

Figure 3

Figure 4