Classification: Biological Sciences; Ecology

The conflict between adaptation and dispersal for maintaining biodiversity in changing

environments

Patrick L. Thompson[1*] and Emanuel A. Fronhofer[2]

1: ORCID ID: 0000-0002-5278-9045

patrick.thompson@zoology.ubc.ca

Department of Zoology, Biodiversity Research Centre, University of British Columbia,

#4200-6270 University Blvd., Vancouver, Canada, V6T 1Z4

2: ORCID ID: 0000-0002-2219-784X

emanuel.fronhofer@umontpellier.fr

ISEM, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France.

*corresponding author

Department of Zoology

University of British Columbia

#4200-6270 University Blvd.

Vancouver, B.C.

Canada V6T 1Z4

Phone: +1-604-822-2131

Fax: +1-604-822-2416

Abstract

Dispersal and adaptation both allow species to persist in changing environments. Yet, we have limited understanding of how these processes interact to affect species persistence, especially in diverse communities where biotic interactions greatly complicate responses to environmental change. Here we use a stochastic metacommunity model to demonstrate how dispersal and adaptation to environmental change independently and interactively contribute to biodiversity maintenance. Dispersal provides spatial insurance, whereby species persist on the landscape by shifting their distributions to track favourable conditions. In contrast, adaptation allows species to persist by allowing for evolutionary rescue. But, when species both adapt and disperse, dispersal and adaptation do not combine positively to affect biodiversity maintenance, even if they do increase the persistence of individual species. This occurs because faster adapting species evolve to hold onto their initial ranges (i.e. monopolization effects), thus impeding slower adapting species from shifting their ranges and thereby causing extinctions. Importantly, these differences in adaptation speed emerge as the result of competition, which alters population sizes and colonization success. By demonstrating how dispersal and adaptation each independently and interactively contribute to the maintenance of biodiversity, we provide a framework that links the theories of spatial insurance, evolutionary rescue, and monopolization. This highlights the expectation that the maintenance of biodiversity in changing environments depends jointly on rates of dispersal and adaptation, and, critically, the interaction between these processes.

Significance Statement

Species can persist when the environment changes by shifting their ranges through dispersal or by adapting to the new conditions that they experience. Thus, we might expect that dispersal and adaptation in combination would increase persistence. Using a simulation model, we show that this may not be the case. Instead, species competition causes dispersal and adaptation to have conflicting contributions to biodiversity maintenance. Dispersal and adaptation each independently increase biodiversity maintenance. But when species both disperse and evolve, faster adapting species persist in their current ranges, preventing others from shifting their ranges to track environmental change. These findings highlight the need to consider ecological and evolutionary processes together, or we risk underestimating how global change will impact biodiversity.

Introduction

When environmental conditions change, what determines whether biodiversity will be maintained? This question remains a great and pressing challenge facing ecologists and evolutionary biologists (1) because of the unprecedented magnitude of anthropogenic environmental change (2). From an ecological perspective, metacommunity theory has established that biodiversity can be maintained in changing conditions by dispersal which allows species to track favourable conditions by shifting their ranges (i.e. spatial insurance; 3). From an evolutionary perspective, genetic changes can allow populations to persist in stressful environmental conditions, which would otherwise cause extinction (i.e. evolutionary rescue; 4-6). But of course, species exist as members of ecological communities, and so their interactions with one another also impact their ability to persist (7-9). Most of our current knowledge on how

species respond to environmental change is based on studies that have only considered a subset

of dispersal, adaptation, and biotic interactions (1). Thus, there is pressing need for theory that

integrates the processes of dispersal and evolutionary change in the context of diverse

communities, if we wish to understand how biodiversity can be maintained (10).

Theory that has integrated eco-evolutionary feedbacks in competitive communities suggests that

the contributions of dispersal and adaptation to the maintenance of biodiversity may be reduced

when these processes interact. When dispersal rates are fast compared to rates of local

adaptation, dispersal can inhibit adaptation by offering an alternative strategy for dealing with

environmental change (11). In contrast, when local adaptation is fast compared to dispersal,

evolutionary mediated priority effects can emerge. Also known as monopolization effects, these

occur when early-arriving species adapt to the local conditions fast enough to prevent pre-

adapted but later-arriving species from colonizing (12, 13). Such monopolization effects have

been demonstrated to be strong drivers of community dynamics in simulated metacommunities

experiencing environmental fluctuations or disturbances (14-17). Evidence of monopolization

effects has also been found in a number of empirical studies (18-21). Urban et al. (10) applied

this idea to changing climates, hypothesizing that monopolization effects could prevent species

from tracking their climate niches. But they did not formally model this hypothesis so it is

unclear how strong this effect should be and how it should depend on rates of evolution and

dispersal.

Our best expectations for how dispersal and evolution should interact to affect the persistence of

multispecies communities under climate change comes from a model by Norberg et al. (22).

These authors found that evolution minimized extinction risks and that this was greatest when

dispersal rates were low. But, in contrast with predictions from the spatial insurance hypothesis

(3, 23), they found that dispersal did not reduce extinctions because it allowed competitively

superior species to expand their ranges to the detriment of other species. Thus, reconciliation of

these conflicting predictions is a priority for advancing our understanding of how dispersal and

evolution contribute to the maintenance of biodiversity in changing conditions.

A further gap in our knowledge is on how stochasticity alters the ability of dispersal and

evolution to maintain biodiversity in changing conditions. The predictions of Norberg et al. (22)

and models of spatial insurance effects have all been deterministic (3, 23). Yet, stochasticity is

key to the emergence of monopolization effects in disturbed or fluctuating environments (14-17).

Thus, although monopolization effects are hypothesized to prevent species from tracking

environmental change through dispersal (10), we do not have good expectations of how this will

depend on the rates of dispersal or levels of genetic variation. Nor do we understand the

consequences of monopolization effects for the maintenance of entire assemblages of interacting

species as the environment changes.

Here we use a stochastic individual-based metacommunity model to test the hypotheses that 1)

when species disperse, but do not evolve their environmental optimum, biodiversity is preserved

through spatial insurance, 2) when species evolve their environmental optimum, but do not

disperse, biodiversity is preserved through evolutionary rescue, and 3) when species both evolve

their environmental optimum and disperse, biodiversity is preserved through a combination of

spatial insurance and evolutionary rescue, but monopolization effects lead to the loss of

biodiversity. Thus, we expect that fast adapting species will monopolize local habitats and

impede species sorting, and thus threaten the persistence of slower adapting species.

Furthermore, we expect that dispersal will only provide spatial insurance in regions of the

metacommunity that contain analogue environments—that is, post change conditions that fall

within the initial range of environmental conditions present in the metacommunity (24). In

contrast, evolutionary rescue should be possible under both analogue and non-analogue

conditions (22). Monopolization effects should occur whenever evolution and dispersal combine

(13).

Results and Discussion

Overall, we find that dispersal and adaptation both independently allow species to persist during

environmental change (Fig. 1a, b), consistent with the spatial insurance and evolutionary rescue

hypotheses and recent findings (3, 4, 23, 25). However, in combination, dispersal and adaptive

evolution can facilitate monopolization effects, which results in fewer species persisting than

when either process operates on its own (Fig. 1c). Dispersal, in the absence of evolution,

increases the proportion of species that persist (Fig. 2a), by providing spatial insurance (3),

whereby species track their environmental optimum through shifting their ranges. Spatial

insurance is only possible when local conditions after environmental change fall within the range

of initial conditions in the metacommunity (i.e., analogue environments). We see that the

positive effect of dispersal on species persistence increases with dispersal rate, only decreasing

slightly at the very highest dispersal rates, when source-sink dynamics become so strong as to

detrimentally impact source population sizes. Evolution, with no dispersal, also results in an

increased proportion of species that persist, by allowing for evolutionary rescue (4) as species

change their environmental optima through adaptation to new environmental conditions (Fig. 1b,

2).

When species both disperse and evolve their environmental optima, we find that these processes

generally conflict in their contributions to the maintenance of biodiversity in changing conditions

(Fig. 2). That is, we see that almost all combinations of dispersal and adaptive potential result in

a reduction in the proportion of species that persist, compared to the persistence that is possible

with one of the two processes acting in isolation. First focusing on analogue environments, high

adaptive potential ($\geq$1.08 x 10^{-4}) without dispersal is sufficient to allow the majority of species to

persist (Fig. 2a). When species also disperse, the number of species that persist at these levels of

adaptive potential is reduced (Fig. S2a, upper-left quadrant). Such high levels of adaptive

potential may correspond to a wide array of organisms, ranging from short lived organisms with

large population sizes such as plankton and other microbes, where evolutionary rescue has been

commonly witnessed (6), to vertebrates with high levels of standing genetic variation (e.g.

adaptation of cold tolerance in sticklebacks (26) and local adaptation across and elevational

gradient in parrotbeaks (27)).

In contrast, adaptation can either increase or decrease persistence with a given dispersal rate (Fig.

2a). In particular, we see that all but the very highest levels of adaptive potential reduce species

persistence when dispersal rates are intermediate (Fig. S2a, lower-left quadrant). For example,

when dispersal is 0.01, increasing adaptive potential from 0 to 1.08 x 10^{-4} reduces the number of

species that persist regionally from 0.74 to 0.59 (Fig. 2a). This is an intermediate level of

dispersal, which has been demonstrated to maximize species sorting and environmental tracking

in other metacommunity models (3, 28) and which is expected by theory to lead to classical

metapopulation dynamics (29). Likewise, intermediate dispersal rates reduce the effectiveness of

intermediate levels of adaptive potential in preserving species; with an adaptive potential of 1.08

x 10^{-4}, the proportion of species that persist decreases from 0.75 when dispersal is 0 to 0.59 when

dispersal is 0.01. Such intermediate levels of dispersal and adaptive potential likely correspond

to species that are already shifting their ranges or showing increased tolerance to warming

conditions, but where rates may not be high enough to ensure long term persistence (30-32). It is

only with high rates of dispersal ($\geq$0.1; e.g. as reported in metapopulations of the butterfly

Melitaea cinxia (33)) and intermediate adaptive potential (3.6 x 10^{-5} or 1.08 x10^{-4}) that

persistence is greater when evolution and dispersal combine, compared to what is possible with

either process in isolation (Fig. S2a, upper-right quadrant).

This antagonistic interaction between dispersal and adaptive potential occurs as the result of

monopolization effects (13), whereby species that are able to adapt faster can remain in place as

the environment changes, making it harder for slower adapting species to persist by shifting their

distributions. Differences in the speed of adaptation are an emergent property of our stochastic

model: the first species in which an advantageous mutation arises and spreads will increase in

number which increases the population scaled adaptive potential and reduces genetic drift,

leading to a positive feedback loop in terms of potential for adaptation. In contrast, species that

by chance don't receive adaptive mutations early on will suffer from maladaptation and decrease

in size, leading to a negative feedback loop (34, 35). As species are lost from the landscape,

dispersal allows the remaining species to expand their ranges (Fig. 3a) taking advantage of

reduced interspecific competition. Thus, we see the greatest range expansions when dispersal

rates are intermediate and low adaptive potential results in low species persistence.

A signature of the monopolization effect is interspecific variation in the degree to which species

expand their ranges (Fig. 1c, 3b). When dispersal or adaptation occur in isolation, we see

relatively little variation in the number of patches that the remaining species occupy. This

variation increases with increasing rates of dispersal (Fig. 3b), but intermediate adaptive

potential results in even greater variation in range size change at any given dispersal rate because

they facilitate monopolization. Monopolization occurs when faster evolving species expand their

distributions as conditions change (Fig. 1c), with adaptation allowing them to remain in place on

their trailing (e.g., warm) edge, and dispersal allowing them to expand their ranges on their

leading (e.g., cold) edge (Fig. 3c). This monopolization of the landscape by a few species causes

the other species to go extinct or to be restricted to only a few patches. Consequently, we see the

greatest difference in interspecific variation in range size compared to the no adaptation

scenarios when adaptive potential and dispersal rates are intermediate (Fig. 3b). These scenarios

correspond to the region of parameter space where dispersal and adaptation potential have the

greatest conflict in the contribution to species persistence (Fig. 2a).

An additional signature of the monopolization effect is that we see leading range boundaries

shifting faster than trailing range boundaries, with intermediate adaptive potential and dispersal

(Fig. 1c, 3c). Species hold onto their trailing edge (i.e., warm boundary) by adapting as the

environment changes, but the range expands on the leading edge (i.e., cold boundary) through

dispersal. This pattern corresponds with the observation that species ranges tend to shift faster at

leading edges compared to trailing edges in response to climate change (36-38) (but see 39).

In non-analogue conditions, that is, regions where the final environmental conditions exceed

those present in the initial metacommunity, adaptive potential is equally effective at preserving

species diversity, but dispersal is far less effective (Fig. 2b). Furthermore, dispersal reduces the

effectiveness of adaptive potential for preserving species richness (Fig. 2b) because it reduces

rates of adaptation through gene swamping (Fig. S3). The exception occurs when adaptive

potential is zero. In this case, persistence in non-analogue environments increases with dispersal,

but this is simply due to the fact that high dispersal facilitates source-sink dynamics so that

species are able to persist in non-analogue patches that are in close proximity to analogue

patches, where population growth rates are positive. This follows the conclusions of Norberg et

al. (22), that biodiversity maintenance will depend most on evolutionary processes in regions of

the planet where climate change creates non-analogue conditions (40), and that dispersal has the

potential to reduce evolutionary rescue in these regions. Maintaining biodiversity in regions that

have no current climate analogue will likely be a major challenge. However, as non-analogue

climatic regions are expected to cover a minority of the globe (40), we have elected to focus

mostly on analogue regions here.

Biotic interactions and persistence under change

We find that diversity losses result from asymmetric responses of the species to environmental

change that are driven by competition between species. Without interspecific interactions (Fig.

S4) all species are able to persist, except in the limiting cases when dispersal or adaptive

potential is so low that persistence is not possible. Competition results in asymmetric responses

to environmental change through two mechanisms. First, it causes unequal reductions in

equilibrium abundances. Species with larger populations are more likely to persist because they

tend to contain more genetic variation, making adaptation faster (4) and because they produce a

greater number of dispersing individuals, making range shifts more likely (23). Second,

competition alters the response of species to local environmental conditions and environmental

change (7, 9). Thus, competition results in interspecific differences in the ability of species to

colonize new habitats in order to track environmental change through species sorting. These

differences in colonization abiolity occur because the resident community acts as a filter,

allowing some species to colonize, while repelling others, even if colonizing species are equally

adapted to the local environment.

The interaction between dispersal and biotic interactions

Dispersal acts to maintain biodiversity as the environment changes by providing spatial

insurance, whereby species shift their distributions to ensure that they are locally adapted.

Without interspecific competition, relatively low rates of dispersal (dispersal ≥ 0.001) are

sufficient to allow almost all species to persist as the environment changes (Fig. S4). However,

with interspecific competition, higher rates of dispersal are required for maintaining biodiversity

(Fig. 2a). This occurs because high rates of dispersal generate source-sink dynamics, which

counteract the effects of biotic interactions in two ways. First, dispersal spreads populations out

across more patches (Fig. S5a), reducing the strength of intraspecific competition. Reduced

intraspecific competition increases regional population sizes (Fig. S4a; 41), making species less

prone to extinction. Second, dispersal allows species to maintain sink populations in marginal

conditions, where abiotic conditions are suboptimal, but persistence would be possible without

competition. Dispersal overcomes the resistance of the biotic community by providing a constant

flow of immigrants. Then, if the environment changes, these populations are already in place and

pre-adapted to the new conditions which makes them new potential source populations. We see

that the positive effect of dispersal on regional species richness increases with dispersal (Fig 2a),

only decreasing slightly at the very highest dispersal rates, when source-sink dynamics become

so strong as to detrimentally impact source population sizes (see also 23). This beneficial effect

of source-sink dynamics is somewhat unexpected, as we tend to consider sink populations to be a

drain on the overall metapopulation. However, our results suggest that source-sink dynamics

may play a key role in providing spatial insurance in changing environments.

The interaction between evolution and biotic interactions

In contrast, adaptation acts to maintain biodiversity by facilitating evolutionary rescue. Without

interspecific competition, moderate levels of adaptive potential are sufficient for maintaining all

species (Fig. S4). With competition, higher levels are required (Fig. 2a) because competition

reduces population sizes, increasing drift and resulting in fewer mutations (42). Of course, biotic

interactions could also increase the potential for evolutionary rescue if these interactions select

against maladapted individuals (42, 43). However, in our case this selective boost does not occur.

Rather, competition acts to slow down rates of evolutionary rescue by reducing population sizes.

Dispersal, evolution, and biotic interactions

When dispersal, adaptation, and competition are combined, monopolization effects are possible,

which reduce the ability of spatial insurance and evolutionary rescue to preserve regional

biodiversity. Monopolization effects occur when adaptive potential and dispersal rates are

intermediate (Fig. 2a) and are only possible because of interspecific competition. Competition

leads to interspecific differences in population sizes. Species with larger population sizes are

more likely to adapt to the changing conditions and they have a higher probability of colonizing

new habitats because they produce more dispersers. These species begin to occupy more space

on the landscape, to the detriment of their competitors, which can lead to extinctions and the loss

of diversity (Fig. 2a). Previous consideration of the monopolization hypothesis has mostly

focused on static or fluctuating environments, with studies highlighting the potential for early

colonizers of a habitat to become locally adapted and then repel later arriving colonists (12-14,

17, 44). However, our results support the hypothesis of Urban et al. (10) that monopolization

effects may impede species sorting under climate change.

General discussion

Our findings allow us to link the concepts of spatial insurance, evolutionary rescue, and

monopolization within a common theoretical framework (Fig. 4). These processes have each

been demonstrated to independently mediate the persistence of species in changing environments

(3, 4, 15), but how they relate and interact with each other has not been previously shown.

Critically, our study explores a range of dispersal rates and levels of adaptive potential that

covers the full range of species persistence outcomes: from full extinction to levels where further

increases in dispersal or adaptive potential would not increase persistence. This distinguishes our

study from that of Norberg et al. (22) who demonstrated clear benefits of evolutionary responses

to climate warming, but not spatial insurance. Rather, they found negative effects of dispersal on

species persistence, thus precluding any potential for monopolization to reduce the effectiveness

of spatial insurance. The lack of a positive effect of dispersal on species persistence is surprising,

and contrasts with our findings and those of previous studies that have demonstrated spatial

insurance effects (3, 23). However, it seems likely that the negative effect of dispersal in Norberg

et al. (22) is due to the fact that environmental change appears to have had a relatively low

impact on population performance relative to the effects of competition in their model. Without

competition, all species persist and are present in all patches in the landscape, before and after

environmental change. Thus, species do not need to shift their distributions in order to persist,

but they can be driven extinct by competition with another species that does track its

environmental optima through dispersal. Such competitive exclusion is likely in their model

because of the assumption of equal inter- and intraspecific competition, which precludes stable

local coexistence. Thus, the contrast between models highlights the fact that the outcome of

dispersal is likely to depend on the balance in strength between the direct effects of

environmental conditions and the density dependent effects of competition on the response of

species to environmental change. Dispersal should have little benefit for species persistence if

species distributions and the composition of species are predominantly structured by biotic

interactions. But we predict that the contribution of dispersal to the maintenance of biodiversity

should increase with the degree to which environmental change impacts population performance.

Although we find that high levels of either dispersal or adaptive potential are sufficient to

preserve almost all species in analogue environments (Fig. 2a), we believe that this is unlikely to

occur in many natural communities. Empirical estimates suggest that dispersal generally limits

species climate ranges over latitudinal gradients, although less so over elevation gradients (45).

Furthermore, estimates suggest that rates of climatic niche evolution in vertebrates and many

plants are far too slow to keep pace with climate change (31, 32). Instead, we expect that rates

will be limiting enough so that dispersal and evolution interactively determine whether species

are able to persist under environmental change. Limiting rates of adaptation are especially likely

for organisms with long generation times, small populations, and limited dispersal ability, or in

fragmented landscapes. Even in bacterial monocultures, where we expect rapid adaptation to

occur, evolutionary rescue is facilitated by dispersal (46). Likewise, current evidence for

evolutionary rescue in communities of interacting species suggests that both adaptation and

dispersal play key roles (25).

Our simulations are simplified representations of how ecological communities respond to

ongoing environmental change. However, our findings are qualitatively unchanged by increasing

the spatial complexity of our landscapes (Fig. S6) or increasing genetic complexity (Fig. S7).

Furthermore, our model also results in rates of phenotypic change (0 to 0.025 haldanes; Fig. S8)

that fall within the range estimated in natural populations (47) and within the range of

evolutionary rates which can be sustained indefinitely under directional environmental change

(34). The degree to which dispersal and adaptive potential have conflicting contributions to the

maintenance of biodiversity under environmental change depends on the specific parameters in

our model but is pervasive across all the combinations of parameter strengths in our sensitivity

analyses (Fig. S10, S11). Of course, additional complexities that are not captured by our model

may modify these effects. We suspect that interspecific variability in dispersal and adaptive

potential should lead to stronger monopolization effects (22). However, if high variability in

rates erodes equalizing coexistence mechanisms (48), pre-change diversity may be reduced, thus

precluding strong monopolization effects. Furthermore, we also know that dispersal rates and

distance are under selection in changing environments (49) as well as mutation rates (50) and

this has the potential to also increase the likelihood of interactions between dispersal and the

evolution. Specifically, dispersal is well known to evolve during range shifts (49, 51). However,

dispersal can also be highly plastic with respect to environmental gradients (52) and other

species (53). While these complexities are beyond the scope of the present study, future

theoretical and empirical work will have to account for the eco-evolutionary details of the

dispersal process. Despite these open questions, our findings clearly highlight the fact that

dispersal and evolution of environmental optima can interact to produce monopolization effects

in the community context, and that ignoring this interaction may lead us to overpredict species

persistence under future environments.

Conclusions

Climate change poses a major risk for biodiversity (54) and is already causing the reorganization

of ecosystems globally (55). Whether species will keep pace and persist in this changing climate

remains uncertain (56) but is expected to depend on both ecological and evolutionary processes

(1, 22). Our results clearly highlight dispersal and evolution can interact, creating the potential

for monopolization effects, which can result in the loss of biodiversity. Together, our findings

provide a more general understanding of the processes that act to maintain biodiversity in a

changing world (Fig. 4). This understanding highlights the need for more focus and study on the

interactions between ecological and evolutionary processes and how they jointly determine how

species and communities will respond to future environmental conditions.

Methods

We use an individual based model to simulate the temporal dynamics of 80 species in a

landscape comprised of 30 habitat patches (see Supplementary Methods for details). To avoid

edge effects, patches in the metacommunity are arranged in a ring, with each patch connected to

its two adjacent patches (for 2D – 5x30 landscapes see Supplemental Materials Fig. S6). We

assume that environmental values increase over the first half of the ring from 0 to $M/2 - 1$ in

integer steps and decrease following the same rule in the second half. These local conditions are

held constant for an initial period of 10,000 generations to allow the simulation to reach quasi-

equilibrium. We then gradually increase the environmental conditions in all patches by a total of

5 units over the next 5000 generations. Individuals reproduce sexually and generations are non-

overlapping. The reproductive output of each female depends on the match between its

phenotype and the local environmental conditions, as well as inter and intraspecific competition

with other individuals that are present in the same patch (Supplemental Methods - Equation 1).

Individuals are diploid and inherit one randomly chosen allele from each parent. The phenotype

is determined by the average of 2 diploid loci (for simulations with 20 loci see Supplemental

Materials Fig. S7). During inheritance, new allele values are drawn at a certain rate m from a

normal distribution with a mean value equal to that of the parent, and a standard deviation equal

to a mutation width of σ_{mut}. Thus, we define adaptive potential as $m\sigma_{mut}{}^2$, which is the per allele

variation generated in each generation due to mutations. Dispersal is between adjacent patches

with a probability d. We contrasted dispersal rates (0 to 0.5) and adaptive potential (0 to 3.6 x 10^{-}

^{4}) that span the range of outcomes from full extinction, to near full persistence in our model.

These levels of adaptive potential resulted in rates of phenotypic evolution between 0 and 0.025

haldanes, which falls within the range estimated in natural populations (47). We quantified the

proportion of species that persisted over the period of environmental change, distinguishing

between regions of the landscape where the final conditions fell within the range present on the

landscape (analogue environments) or not (non-analogue environments).

Acknowledgements

We thank Matthew Osmond, Luc De Meester, Sally Otto, Mary O'Connor and Nathaniel Sharp

for valuable discussions and feedback. PLT is supported by Killam and NSERC Postdoctoral

fellowships. This is publication ISEM-YYYY-XXX of the Institut des Sciences de l'Evolution -

Montpellier.

Citations

1. Urban MC, et al. (2016) Improving the forecast for biodiversity under climate change. *Science* 353(6304):aad8466‑aad8466.

2. IPCC (2013) *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* eds Stocker TF, et al. (Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA).

3. Loreau M, Mouquet N, Gonzalez A (2003) Biodiversity as spatial insurance in heterogeneous landscapes. *P Natl Acad Sci USA* 100(22):12765‑12770.

4. Gomulkiewicz R, Holt RD (1995) When Does Evolution by Natural-Selection Prevent Extinction. *Evolution* 49(1):201‑207.

5. Gonzalez A, Ronce O, Ferriere R, Hochberg ME (2012) Evolutionary rescue: an emerging focus at the intersection between ecology and evolution. *Phil Trans R Soc B* 368(1610):20120404‑20120404.

6. Bell G (2017) Evolutionary Rescue. *Annu Rev Ecol Evol Syst* 48(1):605‑627.

7. Ives AR, Cardinale BJ (2004) Food-web interactions govern the resistance of communities after non-random extinctions. *Nature* 429(6988):174‑177.

8. Davis AJ, Jenkinson LS, Lawton JH, Shorrocks B, Wood S (1998) Making mistakes when predicting shifts in species range in response to global warming. *Nature* 391(6669):783‑786.

9. Alexander JM, Diez JM, Levine JM (2015) Novel competitors shape species' responses to climate change. *Nature* 525(7570):515‑518.

10. Urban MC, De Meester L, Vellend M, Stoks R, Vanoverbeke J (2011) A crucial step toward realism: responses to climate change from an evolving metacommunity perspective. *Evolutionary Applications* 5(2):154‑167.

11. de Mazancourt C, Johnson E, Barrowman (2008) Biodiversity inhibits species' evolutionary responses to changing environments. *Ecology Letters* 11(4):380‑388.

12. De Meester L, Gomez A, Okamura B, Schwenk K (2002) The Monopolization Hypothesis and the dispersal-gene flow paradox in aquatic organisms. *Acta Oecologica-International Journal of Ecology* 23(3):121‑135.

13. De Meester L, Vanoverbeke J, Kilsdonk LJ, Urban MC (2016) Evolving Perspectives on Monopolization and Priority Effects. *Trends Ecol Evolut* 31(2):136‑146.

14. Loeuille N, Leibold MA (2008) Evolution in Metacommunities: On the Relative
Importance of Species Sorting and Monopolization in Structuring Communities. *Am Nat*
171(6):788‑799.

15. Urban MC, De Meester L (2009) Community monopolization: local adaptation enhances
priority effects in an evolving metacommunity. *Proc Biol Sci* 276(1676):4129‑4138.

16. Vanoverbeke J, Urban MC, De Meester L (2015) Community assembly is a race between
immigration and adaptation: eco-evolutionary interactions across spatial scales.
Ecography 39(9):858‑870.

17. Leibold MA, Urban MC, De Meester L, Klausmeier CA, Vanoverbeke J (2019) Regional
neutrality evolves through local adaptive niche evolution. *P Natl Acad Sci USA*
116(7):2612‑2617.

18. De Meester L, Louette G, Duvivier C, Van Damme C, Michels E (2007) Genetic
composition of resident populations influences establishment success of immigrant
species. *Oecologia* 153(2):431‑440.

19. Fukami T, Beaumont HJE, Zhang X-X, Rainey PB (2007) Immigration history controls
diversification in experimental adaptive radiation. *Nature* 446(7134):436‑439.

20. Crutsinger GM, Souza L, Sanders NJ (2008) Intraspecific diversity and dominant
genotypes resist plant invasions. *Ecology Letters* 11(1):16‑23.

21. Pantel JH, Duvivier C, Meester LD (2015) Rapid local adaptation mediates zooplankton
community assembly in experimental mesocosms. *Ecology Letters* 18(10):992‑1000.

22. Norberg J, Urban MC, Vellend M, Klausmeier CA, Loeuille N (2012) Eco-evolutionary
responses of biodiversity to climate change. *Nature Climate Change* 2(10):747‑751.

23. Thompson PL, Gonzalez A (2017) Dispersal governs the reorganization of ecological
networks under environmental change. *Nature Ecology and Evolution* 1(6):0162.

24. Williams JW, Jackson ST (2007) Novel climates, no-analog communities, and ecological
surprises. *Front Ecol Environ* 5(9):475‑482.

25. Low-Décarie E, et al. (2015) Community rescue in experimental metacommunities. *P Natl
Acad Sci USA* 112(46):14307‑14312.

26. Barrett RDH, et al. (2010) Rapid evolution of cold tolerance in stickleback. *Proc R Soc B*
278(1703):233‑238.

27. Lai Y-T, et al. (2019) Standing genetic variation as the predominant source for adaptation
of a songbird. *P Natl Acad Sci USA* 116(6):2152‑2157.

28. Thompson PL, Rayfield B, Gonzalez A (2017) Loss of habitat and connectivity erodes
species diversity, ecosystem functioning, and stability in metacommunity networks.
Ecography 40(1):98‑108.

29. Fronhofer EA, Altermatt F (2017) Classical metapopulation dynamics and eco-
evolutionary feedbacks in dendritic networks. *Ecography* 40(12):1455‑1466.

30. Diamond SE (2018) Contemporary climate-driven range shifts: Putting evolution back on
the table. *Funct Ecology* 32(7):1652‑1665.

31. Quintero I, Wiens JJ (2013) Rates of projected climate change dramatically exceed past
rates of climatic niche evolution among vertebrate species. *Ecology Letters* 16(8):1095‑
1103.

32. Franks SJ, Weber JJ, Aitken SN (2013) Evolutionary and plastic responses to climate
change in terrestrial plant populations. *Evolutionary Applications* 7(1):123‑139.

33. Hanski I (1999) *Metapopulation Ecology* (Oxford University Press, Oxford).

34. Bürger R, Lynch M (1995) Evolution and Extinction in a Changing Environment: a
Quantitative-Genetic Analysis. *Evolution* 49(1):151‑163.

35. Osmond MM, Klausmeier CA (2017) An evolutionary tipping point in a changing
environment. *Evolution* 71(12):2930‑2941.

36. Sunday JM, Bates AE, Dulvy NK (2012) Thermal tolerance and the global redistribution
of animals. *Nature Climate Change* 2(9):686‑690.

37. Ordonez A, Williams JW (2013) Climatic and biotic velocities for woody taxa
distributions over the last 16 000 years in eastern North America. *Ecology Letters*
16(6):773‑781.

38. Poloczanska ES, et al. (2013) Global imprint of climate change on marine life. *Nature*
Climate Change 3(10):919‑925.

39. Robinson LM, Hobday AJ, Possingham HP, Richardson AJ (2015) Trailing edges
projected to move faster than leading edges for large pelagic fish habitats under climate
change. *Deep-Sea Research Part II* 113(C):225‑234.

40. Mahony CR, Cannon AJ, Wang T, Aitken SN (2017) A closer look at novel climates: new
methods and insights at continental to landscape scales. *Global Change Biology*
23(9):3934‑3955.

41. Holt RD (1985) Population dynamics in two-patch environments: Some anomalous
consequences of an optimal habitat distribution. *Theoretical Population Biology*
28(2):181‑208.

42.   Osmond MM, de Mazancourt C (2012) How competition affects evolutionary rescue. *Phil*
Trans R Soc B 368(1610):20120085‑20120085.

43.   Osmond MM, Otto SP, Klausmeier CA (2017) When Predators Help Prey Adapt and
Persist in a Changing Environment. *Am Nat* 190(1):83‑98.

44.   Vanoverbeke J, Urban MC, De Meester L (2015) Community assembly is a race between
immigration and adaptation: eco-evolutionary interactions across spatial scales.
Ecography 39(9):858‑870.

45.   Hargreaves AL, Samis KE, Eckert CG (2014) Are species' range limits simply niche limits
writ large? A review of transplant experiments beyond the range. *Am Nat* 183:157‑173.

46.   Bell G, Gonzalez A (2011) Adaptation and Evolutionary Rescue in Metapopulations
Experiencing Environmental Deterioration. *Science* 332(6035):1327‑1330.

47.   Hendry A, Kinnison MT (1999) Perspective: The pace of modern life: Measuring rates of
contemporary microevolution. *Evolution* 53(6):1637‑1653.

48.   Chesson P (2000) General Theory of Competitive Coexistence in Spatially-Varying
Environments. *Theoretical Population Biology* 58:211‑237.

49.   Kubisch A, Holt RD, Poethke H-J, Fronhofer EA (2014) Where am I and why?
Synthesizing range biology and the eco-evolutionary dynamics of dispersal. *Oikos*
123(1):5‑22.

50.   Cobben MMP, Mitesser O, Kubisch A (2017) Evolving mutation rate advances the
invasion speed of a sexual species. *BMC Evol Biol* 17(1):150.

51.   Phillips BL, Brown GP, Webb JK, Shine R (2006) Invasion and the evolution of speed in
toads. *Nature* 439(7078):803‑803.

52.   Fronhofer EA, Nitsche N, Altermatt F (2017) Information use shapes the dynamics of
range expansions into environmental gradients. *Global Ecol Biogeogr* 26(4):400‑411.

53.   Fronhofer EA, et al. (2018) Bottom-up and top-down control of dispersal across major
organismal groups. *Nature Ecology and Evolution* 2(12):1859‑1863.

54.   Urban MC (2015) Accelerating extinction risk from climate change. *Science* 348:571‑
573.

55.   Walther GR (2010) Community and ecosystem responses to recent climate change. *Phil*
Trans R Soc B 365(1549):2019‑2024.

56.   Loarie SR, et al. (2009) The velocity of climate change. *Nature* 462(7276):1052‑1055.

Figure 1. Illustration of how dispersal and evolution of environmental optimum, in isolation

(a,b, respectively) and in combination (c), affect how species respond to environmental change.

Only one half of the landscape ring is shown (see Fig. S1 for full landscape). Species are

arranged on the x-axis by their pre-change latitude. Circles and diamonds are paired for each

species and indicate the latitude and size of each population prior to and after environmental

change, respectively. Species with circles but not diamonds failed to persist. The colour shows

the mean environmental optimum in each population. Panel a shows a scenario where dispersal

is intermediate (0.01) and the adaptive potential is zero; here, species respond to environmental

change by shifting to higher latitudes to maintain the match between their phenotype and their

local environmental conditions. Panel b shows a scenario where dispersal is 0 and adaptive

potential is intermediate (1.08×10^{-4}); here species respond to environmental change through

adaptation (change in colour), with no change in latitude. Panel c shows a scenario where

dispersal (0.01) and adaptive potential (1.08×10^{-4}) are both intermediate; here species respond

through a combination of shifting latitude and through adaptation. Results shown are from one

representative simulation run with standard parameter values (Table S1). To explore additional

combinations of dispersal and adaptive potential in a Shiny app visit -

https://shiney.zoology.ubc.ca/pthompson/Meta_eco_evo_shiny/.

Figure 2. The proportion of species that are maintained following environmental change

depending on dispersal and adaptive potential (colour). The proportion of species maintained

was calculated as the number of species that were present in the region (a - analogue or b - non-

analogue) after environmental change, divided by the number of species that were present

before. Therefore, species that were only present in analogue patches would not be included in

the non-analogue diversity. The lines show the median value across 50 replicate simulations with

standard parameter values (Table S1) and the bands show the interquartile range. This figure

shows patterns for regional scale diversity. Local scale patterns are shown in Fig. S9.

Figure 3. Change in the average number of habitat patches occupied (range size) by the species

that persist during environmental change (a), the interspecific variation in range size change,

excluding species that go extinct (b), and the leading vs. trailing edge asymmetry of the range

expansion (c), depending on dispersal and adaptive potential (colour). Positive (negative) values

of range change asymmetry indicate that the centroid of the range shifted towards warmer

(colder) conditions, relative to the mean environmental optima of the species. The lines show the

median value across 50 replicate simulations with standard parameter values (Table S1) and the

bands show the interquartile range. Analogue and non-analogue regions are included together in

these estimates. The lines for adaptive potential = 0 do not extend to the lowest dispersal rates

because all species went extinct during environmental change.

Figure 4. Conceptual illustration of how dispersal and the evolution of environmental optima

independently and interactively act to maintain biodiversity in changing environmental

conditions. Panel a shows the distribution and abundance of three species spanning a climate

gradient (e.g., warm to cold). Mean position shown by vertical dashed lines. Each species is

locally adapted to a different part of this gradient as indicated by the warmth of the color of the

curves. Panels b, c, and d show hypothetical scenarios after the environment has changed. In

panel b, with dispersal but no evolution of environmental optima, the species persist by shifting

upwards (spatial insurance). In panel c, with evolution of environmental optima but no dispersal,

the species persist by adapting to the changed conditions (change in color - evolutionary rescue),

but do not shift their ranges. In panel d, with both dispersal and evolution, the two outer species

evolve faster than the middle species, holding onto their initial trailing edge through adaptation

but expanding on their leading edge through dispersal (monopolization). By monopolizing the

landscape, they drive the middle species extinct (dashed curve). Panel e shows how dispersal and

the evolution of environmental optima each allow for species persistence via spatial insurance

and evolutionary rescue, respectively (based on Fig. 2a). But together they can lead to

monopolization effects, which can reduce biodiversity in changing environments.

Figure 1

589 **Figure 2**

592   **Figure 3**

    Figure 4