Pan-Cancer and Single-Cell modelling of genomic alterations through gene expression

Daniele Mercatelli[1], Forest Ray[2] & Federico M. Giorgi[1, *]

[1] Department of Pharmacy and Biotechnology, University of Bologna, Via Selmi 3, 40138, Bologna, Italy

[2] Columbia University Medical Center, New York, NY 10032, USA

* Corresponding Author

E-mail: federico.giorgi@unibo.it

Short Title

Pan-Cancer and Single-Cell modelling of genomic alterations through gene expression

Abstract

Cancer is a disease often characterized by the presence of multiple genomic alterations, which trigger altered transcriptional patterns and gene expression, which in turn sustain the processes of tumorigenesis, tumor progression and tumor maintenance. The links between genomic alterations and gene expression profiles can be utilized as the basis to build specific molecular tumorigenic relationships. In this study we perform pan-cancer predictions of the presence of single somatic mutations and copy number variations using machine learning approaches on gene expression profiles. We show that gene expression can be used to predict genomic alterations in every tumor type, where some alterations are more predictable than others. We propose gene aggregation as a tool to improve the accuracy of alteration prediction models from gene expression profiles. Ultimately, we show how this principle can be beneficial in intrinsically noisy datasets, such as those based on single cell sequencing.

Author Summary

In this article we show that transcript abundance can be used to predict the presence or absence of the majority of genomic alterations present in human cancer. We also show how these predictions can be improved by aggregating genes into small networks to counteract the effects of transcript measurement noise.

1 Introduction

Cancer is a molecular disease occurring when a cell or group of cells acquire uncontrolled proliferative behavior, conferred by a multitude of deregulations in specific pathways [1]. As is implied by such a broad definition, cancer is a highly heterogeneous disease, showing remarkably different molecular, histological, genetic and clinical properties, even when comparing tumors originating from the same tissue [2]. Many cancers are characterized by the presence of single nucleotide or short indel mutations and/or copy number alterations, which appear somatically at the early stages of oncogenesis and can drive tumor progression [3]. Cancers can be broadly divided in two classes: the M class, where point mutations are prevalent, and the C class, where copy number variations (CNVs) are more numerous and are often associated with TP53 mutations. Tumor class influences anatomic location. Most ovarian cancers, for example, belong to the C class, while most colorectal cancers belong to the M class, although many exceptions do exist [4].

The Cancer Genome Atlas (TCGA) project [5] has recently undergone a major effort to collect vast amounts of information on thousands of distinct tumor samples. The TCGA data collection, commonly referred to as the "Pan-cancer" dataset, provided the scientific community with an avalanche of data on DNA alterations, gene expression, methylation status and protein abundances among others, with the critical mass necessary to identify rarer driver tumorigenesis effects in many types of cancers [6–8]. By combining all 33 TCGA datasets, Bailey and colleagues [9] recently outlined a pan-cancer map of which mutations can be drivers for the progression of cancer.

The availability of thousands of samples measuring many different variables in cancer has allowed scientists to generate statistical models of relationships between different

molecular species. A pan-cancer correlation network between coding genes and long noncoding RNAs, for example, sheds light on the function of non-coding parts of the transcriptome [10]. More recently, mutations on transcription factors (TFs) have been linked to altered gene expressions and phosphoprotein levels in 12 TCGA tumor type datasets [11]. Network approaches have been applied to identify clusters of coexpressed genes, shared by multiple cancer types [12]. Several studies have sought to characterize the relationships between genomic status and expression levels in cancer, trying to identify commonalities across different cancer types [13,14]. In particular, Alvarez and colleagues [15] have postulated that the effect of genomic alterations in cancer can be more readily assessed by aggregating gene expression profiles into transcriptional networks, rather than by profiles taken separately.

While the association between genomic events and gene expression is proven in several scenarios, it remains to be seen if it can be assessed in scenarios where fully quantitative readouts are unavailable, such as low coverage samples. One of these scenarios is Single Cell Sequencing [16], often carried out in experiments where thousands of mutations are generated via a system of pooled CRISPR-Cas9 knockouts [17].

To our knowledge there is no study trying to identify relationships between all genomic alteration events (somatic mutations/indels and CNVs) and global gene expression across cancers. In this study, we use 24 TCGA tumor datasets to investigate whether gene expression can be used to predict the presence of specific genomic alterations in several cancer tissue contexts. To this end, we leverage the current availability of a vast family of machine learning algorithms [18]. We investigate whether some gene alterations can be better modelled than others, and whether using grouped gene expression profiles as

1 aggregated variables can effectively identify specific genomic alterations. Finally, we test

2 whether predicting mutations and CNVs can be carried out in an intrinsically noisy single cell

3 RNA-Seq (scRNA-Seq) transcriptomics datasets.

Results

Collection of Pan-Cancer Dataset

We downloaded the most recent version of the TCGA datasets available on Firehose

(v2016_01_28), encompassing mutational, CNV and gene expression data. Using T-

distributed stochastic neighbor embedding (TSNE [19]) clustering on gene expression data

(9642 samples), we observed how different tumor types cluster separately from each other

(Figure 1A). However, two tumour types segregate into two subgroups: breast cancer, which

subdivides into a major luminal cluster and a smaller (in terms of samples collected) basal

cluster [20]; and esophageal carcinoma, which roughly subdivides into adenocarcinomas

and squamous cell carcinomas [21].

We then aggregated the single nucleotide and short indel somatic mutation data from the

same samples for which we had collected gene expression. As is widely known, TP53 is the

most mutated gene in human cancer (Figure 1B), followed by PIK3CA, SYNE1 and KRAS. As

shown before [4] some tumor types are characterized by a high presence of somatic

mutations. In particular, lung squamous carcinoma, mesothelioma and esophageal cancer

carry at least one of these events in almost 100% of the samples in the TCGA dataset. In the

figure, we filtered out commonly known non-driver mutations [22], such as those happening

in long genes like TTN and OBSCN, but we kept them in all following analyses for the sake of

completion. A representation of all mutated genes, including blacklisted ones, is available in

Figure S1. Some tumors are characterized by the prevalence of a mutation in a specific gene,

such as the G-protein coding BRAF in thyroid carcinoma [23] or IDH1, translating into

isocitrate dehydrogenase, in low grade glioma [24].

Finally, we obtained readouts of CNV status for all TCGA samples. CNVs can have different

extensions in terms of nucleotides affected and can sometimes encompass entire

chromosomes [25] and the thousands of genes therein. In order to limit the number of

variables to a more meaningful subset, we assigned a CNV score to every gene, according

the copy number score of the genomic region most overlapping with the UCSC annotated

gene boundaries (genome version hg19). We then tested models for all genes affected by a

CNV in at least 10 samples (extending what previously done in [26]). In order to make CNV

variables comparable to the mutational ones, we defined a cut-off for presence or absence

by using the $\log_2$(CNV) threshold of 0.5, which roughly corresponds to at least one copy gain

for amplifications, and at least one copy loss for deletions (see Materials and Methods). We

then reported their abundance in the pan-cancer dataset, distinguishing between

amplifications (Figure 1C) and deletions (Figure 1D). As previously shown [4], virtually all

ovarian cancer samples are characterized by at least one CNV event. Among the most

amplified genes, we find the oncogenes SOX2 [27], EGFR [28] and MDM2 [29], and also a

non-coding gene, PVT1, the most amplified gene in breast cancer, with proven but as-of-yet

uncharacterized proto-oncogenic effects [30,31]. Amongst the most deleted genes (Fig.1D)

we observe well known tumor-suppressor genes, such as CDKN2A [32,33] and PTEN [34,35].

Modelling Cancer Alterations with gene expression

After collecting all the expression and genomic alteration data from TCGA, we set out to generate models able to predict the presence or absence of each event by virtue of gene expression data in the contexts of all collected tumor types.

We tested several modelling algorithms for classification using the aggregator platform for machine learning caret [18] in the bladder cancer mutational dataset [36]. We observed that all models provide better-than-random predictions for the majority of mutational events, in terms of area under the ROC curve (AUROC)(Figure 2) [37]. We chose the top-scoring algorithm in this test, the Gradient Boost Modelling algorithm (gbm), a robust tree-based boosting model [38], due to its robustness and speed of implementation.

We calculated gbm models for all tumour types of at least 100 samples with co-measured expression and CNV or mutations, which included 24 of the 33 TCGA tumor types. The models were predictive of genomic events observed in no less than 5% and no more than 95% of the patients in the dataset, and at least in 10 samples. Our results show that in all tumour types, a machine learning algorithm based on gene expression is consistently better than a random predictor (AUROC line at 0.5) at correctly classifying tumour samples for the presence or absence of specific genomic alteration events (Figure 3 and Supplementary Table S1). In particular, TP53 mutations are well modelled in many of these tumor types, being the most well predicted mutational event in both acute myeloid leukemia and low grade glioma. We could also model the presence of a copy loss of TP53 in sarcoma, which can be predicted with an accuracy of 70%. Ovarian and pancreatic cancer datasets presented exceptional cases, in that each contained such high TP53 mutation rates (next to

95% detected) [39,40] that our algorithms could not distinguish sufficient differences within

each dataset to train a model. Also KRAS-targeting events are well modelled, specifically in

colon, lung and stomach cancer, and cervical squamous carcinoma [41]. We noted a

tendency where models for more frequent CNV events yielded a greater predictive power

(Figure S2), a tendency not observed for somatic mutation models. We then tested if known

tumor-related genes, such as those curated by the Cancer Gene Census [42] are better

modelled than the rest of the genome. There is no difference in mutation and amplification

results, but for deletion events, oncogenes yield weaker models (Wilcoxon Test, p=0.0037,

figure S3) and tumor suppressor genes yield generally stronger models (p=0.00050). This is

in agreement with the central paradigm of cancer, where a tumor suppressor gene deletion

can be one of the driving events of tumorigenesis and tumor progression [43]. On the other

hand, deletion of tumor-promoting oncogenes is generally unfavourable for tumor

progression, and so, generally speaking it should be present only as a passenger event,

unlikely to determine global gene expression and tumor fate.

16   Modelling specific alterations with noise addition

In order to understand whether cancer-related genomic alterations can be modelled by

gene expression in scenarios with lower signal-to-noise ratio, we artificially perturbed the

TCGA gene expression dataset via the addition of Gaussian noise, and then proceeded to

build models to predict the presence of TP53 mutations in breast cancer, the largest dataset

in TCGA by number of samples.

As expected, the addition of uniform random gaussian noise to the gene expression matrix

has a detrimental effect on the amount of information left for modelling the presence of

TP53 somatic mutations (Figure 4A).

We then decided to test several permutations of noise addition on the same breast cancer

expression data, by each time aggregating genes into networks defined a priori in the same

context, using a Tukey Biweight Robust Average method [44] on Weighted Gene Correlation

Network Analysis (WGCNA) clusters [45] and the VIPER algorithm [15] on ARACNe-AP

networks [46]. It is important to note that WGCNA clusters are completely non-overlapping

and yield generally a lower number of aggregated variables than VIPER clusters, which are

groups of genes possibly shared by other transcription factor clusters and that collectively

yield the global expression of a transcription factor target set (dubbed as a proxy for "TF

activity" in the original VIPER manuscript [15]).

Our results show that gene expression, VIPER activity and WGCNA clusters yield very similar

models for predicting TP53 mutations in breast cancer (figure S4). The amount of

information contained in the input variables is therefore comparable. Adding noise to the

input expression matrix, however, and then aggregating the resulting noise-burdened genes

into VIPER or WGCNA clusters (see Materials and Methods), provides robustness to the

models (Figure 4B). Similar results with higher variances (possibly due to the smaller size of

the datasets) can be observed for EGFR amplifications in glioblastoma (Figure S5) and lung

squamous carcinoma (Figure S6), for PVT1 amplifications in ovarian cancer (Figure S7) and

for PTEN deletions in sarcoma (Figure S8). In all these examples, however, the performance

of the simple WGCNA/Tukey aggregation is closer (if not worse) to that of simple gene

expression.

An alternative way to reduce the information content from an NGS gene expression dataset is to reduce the number of read counts from each sample. This operation reflects either a low coverage bulk RNA-Seq experiment or an experiment arising from Single-Cell sequencing [47]. In particular, single-cell RNA-Seq (scRNA-Seq) is characterized by the dropout phenomenon [48] wherein genes expressed in the cells are sometimes not detected at all. In order to simulate such scenarios, we down-sampled each RNA-Seq gene count profile from the largest TCGA dataset (Breast Cancer) to a target aligned read number using a beta function, which allows for reduction coupled with random complete gene dropouts (Figure 5A). We then modelled again the presence of TP53 mutations using gene expression (Figure 5B). We found out that models based on standard unaggregated gene expression experience an accuracy drop at around 30M reads, while aggregating genes using VIPER (but not with WGCNA) allows for better-than-random accuracies even at 3M reads, confirming the benefits of gene aggregation in low coverage RNA-Seq, as previously found e.g. for sample clustering [49].

Mutation prediction in single-cell data

Based on the results from the pan-cancer analysis, where we predicted sample mutations based on pooled RNA-Seq gene expression patterns, we decided to extend the same approach on single-cell datasets. Recently, the CROP-Seq methodology has been introduced [17], allowing for the measurement of cell-specific transcriptome-wide gene expression and mutations induced by CRISPR-Cas9 [50], thanks to the concurrent sequencing of CRISPR-Cas9 guide RNAs. We therefore tested the capability of gbm models to predict mutations using gene expression variables in two independent single-cell datasets. The first dataset (dubbed "Datlinger") was extracted from the Jurkat cell line derived from human T lymphocytes [17]. The second one (dubbed "Shifrut") derive from primary unstimulated T

cells from a human donor [51]. We removed cell UMI (Unique Molecular Identifier) counts

and cell cycle as common confounding effects of single cell datasets [52] (Figure S10). We

generated a regulatory transcription network using ARACNe-AP on the RNA-Seq Cancer Cell

Line Encyclopedia dataset (CCLE [53]), which comprises 1021 distinct human cell lines. Using

the CCLE network, we aggregated gene expression from the single cell datasets using the

VIPER algorithm and implemented the resulting TF-centered VIPER activity profiles to build

prediction models for the Crop-Seq detected mutations. Parallelly, we built models using

un-aggregated VST-normalized gene expression data. Our results show that gbm models

based on VIPER activity variables globally achieve a significantly higher performance in both

the Datlinger (p=8.0x10^{-85}) and Shifrut datasets (p=2.2x10^{-117}) when compared with models

obtained from gene expression data (Figure 6). For specific mutations (TUBB gene, CDKN1B)

the VIPER aggregation based on CCLE ARACNe networks seem to be particularly beneficial to

increase the performance of mutation prediction models based on gene expression, while

for a few mutations, such as RUNX1, the CCLE-based networks significantly decrease the

model performance.

Discussion

In this paper, we tested a framework to investigate the complex relationships between

genetic events and transcriptional deregulation through machine learning approaches. We

demonstrated as a generalized proof-of-principle that genomic alterations can be modeled

by gene expression across several human cancers through several machine learning

algorithms, and specifically that a gradient boost modeling approach seems optimal for the

task. In the process, we generated a collection of models for each genomic alteration in

each cancer context, showing that the best predicted alterations are not necessarily

targeting known oncogenes or tumor suppressors. Interestingly, we show how the

aggregation of gene expression profiles in groups of coexpressed genes, via the

ARACNe/VIPER or WGCNA methods, makes the models more robust and more resistant to

perturbations such as gaussian noise or artificial downsampling. Finally, we have shown how

the same aggregation principle can have beneficial effects in predicting the presence of

mutations in intrinsically noisy scenarios, both with artificial noise introduction and read

reduction. At the same time, we have shown that expression-based mutation prediction can

be modeled out in single-cell sequencing contexts, which can be considered as real cases of

noisy datasets. The capability of predicting mutations based on single-cell RNA-Seq is

however reduced when compared to datasets derived from pooled cells sequencing, as

those provided by the TCGA dataset: the average performances of TCGA models (Figure 3)

generally rest on a range between 0.6 and 0.9 area under the ROC curve, while the

performance of CROP-Seq models fall on an average value of 0.55 (Figure 6).

The performance of gene aggregation methods has been tested before for sample clustering

in RNA-Seq read reduction scenarios [15,49], but never in this specific task nor in a pan-

cancer or a single-cell context. As a principle, the usage of robust averages of pre-defined

co-expressed genes can be applied in any context where reliability of gene expression data

is necessary, from differential expression to pathway enrichment analyses. The notion that

relationships between genomic alterations and gene expression profiles can be robustly

modelled across different cancer scenarios, as well as in single-cell and noisy contexts, can

have important repercussions in diagnostics and quantification studies of heterogeneous

cell populations, where theoretically a single quantitative expression experiment can be

used to predict the presence or absence of a mutation.

1 Materials and Methods

2 Data processing

We obtained raw expression counts, mutation and CNV raw data from TCGA using the

Firehose portal (gdac.broadinstitute.org). Raw counts were normalized using Variance

Stabilizing Transformation as described before [54]. Somatic mutations not changing the

aminoacid sequence of the protein product were discarded. We flagged genes blacklisted by

the MutSig project [22], such as TTN, ORs, MUCs as false positives, and removed them from

further analysis (except the most mutated in the pan-cancer dataset, shown in Figure S1).

CNV tracks were associated to the targeted gene using the GenomicRanges R package [55].

Gene-centered CNVs were then associated to the expression profile of the gene itself.

Genes affected by a CNV in more than 10 samples were used in the rest of the analysis.

Samples with more than 0.5% of the genes in the genome somatically amplified, deleted or

mutated were deemed "hypermodified" and the total number was shown in Figure 1

bottom bars.

Clustering analysis was carried out on the TCGA tumor samples using the expression profiles

of 1172 Transcription Factors defined by Gene Ontology terms "transcription factor activity,

sequence-specific DNA binding" (GO:0003700) and "nuclear location" (GO:0005634) [56].

The dataset expression profiles were visualized after TSNE transformation [19] with 1000

iterations using a 2D kernel density estimate for coloring different tumor types [57].

Oncogenes and Tumor Suppressor genes were obtained from the COSMIC Cancer Gene

Census in October 2018 [42].

1 Modeling

We used the R caret package [18] v 6.0-81 as the platform to run all our predictive models in

a standardized and reproducible way. Default parameters for model training were used.

Binary classifiers were built to predict the presence/absence of mutation, amplification and

deletion events. The CNV value provided by TCGA corresponds to log2(tumor coverage) −

genomic median coverage. The threshold for amplification/deletion presence was set to 0.5.

Data partitioning was performed once for each tumor type, with 75% of the samples used

for training and 25% for test purposes. Training was performed using 10-fold Cross

Validation. Recursive Feature Elimination was carried out by the default caret

implementation on the 10,000 highest variance gene expression tracks. The algorithms used

(and R packages implementing theme) were:

• Bayesian Generalized Linear Model (bayesglm)

• Tree Models from Genetic Algorithms (evtree)

• Gradient Boost Modeling (gbm)

• Generalized Linear Model (glm)

• k-Nearest Neighbors (kknn)

• Linear Discriminant Analysis (lda)

• Neural Networks (mxnet)

• Neural Networks with Feature Extraction (pcaNNet)

• Random Forest (rf)

• Linear Support Vector Machine (svmLinear)

• Radial Support Vector Machine (svmRadial)

In order to reduce information from the gene expression profiles, we adopted two

strategies. The first, shown e.g. in Figure 4B, adds random gaussian noise to the expression

tracks, with a variable standard deviation (indicated as "Gaussian Noise Level"). Each model

run after noise addition was run 100 times to allow for various data partitions. The second

strategy (Figure 5) reduced the number of reads mapped to each gene in order to obtain

expression samples with decreased total gene counts. In order to do so, we applied to each

gene in each sample a downsampling factor sample from a beta distribution:

$$\frac{1}{B(\alpha, \beta)} x^{\alpha-1}(1-x)^{\beta-1}$$

9   Where B is the Beta function, acting as a normalization constant, x is the raw gene

10   expression count in a particular sample, α is the first shape parameter and β the second

11   shape parameter. In order to reduce the total sample coverage to the desired level, β is set

12   to 0.1 and α is set to:

$$\alpha = \frac{\beta f/r}{1 - f/r}$$

Where f is the desired number of reads and r is the total number of reads in the sample. A

real case example of this beta distribution is shown in Figure S9.

Aggregation algorithms

We used ARACNe-AP [46] to generate TF-centered networks on each of the VST-normalized

TCGA expression datasets. TFs were selected via Gene Ontology as described before, with p-

value for each network edge set to 10^{-8}. ARACNe networks were then used to obtain an

aggregated value of TF activity for each sample using the VIPER algorithm [15] which reports

the collective gene expression level changes of each TF-centered network vs. the mean

expression of each gene in the dataset. Only TF networks with at least 10 genes (excluding

the TF) were included.

WGCNA clusters of genes were constructed using the wcgna package [45] with default

parameters and minimum network size set to 10 To obtain a robust median expression

value for each WGCNA cluster in each sample we used Tukey's Biweight function as

implemented by the R *affy* package [58].

Single Cell analysis

We generated TF regulatory networks using ARACNe-AP as described before on the CCLE

dataset available at https://portals.broadinstitute.org/ccle/data, raw counts version 2018-

09-29, normalized by Variance-Stabilizing Transformation [58].

We downloaded raw RNA-Seq counts and gRNA mutation data from single-cell CROP-Seq

datasets, specifically: 1) the Datlinger dataset available on Gene Expression Omnibus (GEO)

series GSE92872 [17] and 2) the Shifrut dataset was obtained from a healthy donor and is

available as raw counts and cell-sepcific gRNA from GEO sample GSM3375483 [51]. Both

single cell CROP-Seq datasets were normalized using the R package Seurat with default

parameters, as follows: a global-scaling normalization method ("LogNormalize") was applied

on raw gene counts for each cell, then the values were multiplied by a scale factor (10,000

by default) and the results were log-normalized. These values were then regressed by two

variables: UMI counts and cell cycle, using cell cycle markers from [52]. As an example of the

Seurat regression, the TSNE representation of the Datlinger dataset before and after

normalization clearly shows the removal of cell-cycle bias effects (Figure S10).

Gradient boost modelling (gbm) was applied to each CROP-Seq dataset by aggregating cells

carrying mutations on the same genes and using wild-type cells as control. Performance of

gbm models using VIPER and expression variables was compared using a two-tailed

Wilcoxon test on 100 repetitions of training/test set splits before cross-validation for model

testing [59].

Methods Availability

7 All code used to generate the analysis and the figures of this paper is available in the online

8 materials as Supplementary Code.

Figure Legends

Figure 1. The TCGA dataset used. A: TSNE clustering of TCGA samples based on gene

expression profiles. The 2D median of each tumor type is indicated using the TCGA tumor

code. Subset size is indicated in brackets next to tumor type names to the right. B: table of

most somatically mutated genes across TCGA tumor samples, in terms of number of

samples where the gene is somatically mutated with altered protein product sequence. C:

table of most amplified genes across TCGA tumor samples. D: table of most deleted genes

across TCGA tumor samples. The fraction of total TCGA samples carrying a gene-targeting

event is indicated to the right of panels B-D, and the fraction of samples where more than

0.5% of the genes is affected by the panel event type is indicated to the bottom of panels B-

D.

Figure 2. Performance of 11 machine learning algorithms in binary classification of

mutated/nonmutated samples using gene expression predictor variables in the Bladder

Cancer dataset. Each point corresponds to a specific mutation/model. Performance is

indicated as AUROC: Area Under the Receiver Operating Characteristic curve.

Figure 3. Performance of gbm models for each genomic alteration event in TCGA, predicted

as a function of each tumor gene expression. Boxplots indicate distribution median, upper

and lower quartile. Alterations targeting TP53 and KRAS are indicated.

Figure 4. Performance of a TP53 somatic mutation gbm model upon gaussian noise

addiction. A: ROC curves (and AUC) upon addition of increasing levels (in terms of SD of a

gaussian distribution with mean=0) of gaussian noise. B: AUROCs of the model with

increasing noise, calculated using gene expression (black line) or aggregated gene

expression using the WGCNA (green line) or VIPER (red line) algorithms. Pseudocounts of

0.1 are added in order to show zero counts as -1 in $\log_{10}$ scale.

Figure 5. Performance of a TP53 mutation gbm model upon downsampling of the TCGA

breast cancer RNA-Seq dataset. A: for a single TCGA sample (TCGA-A1-A0SB-01) with 43.8

gene mapping reads, the downsampling algorithm is applied for multiple target read

quantities. X-axis shows the count for each gene in the original sample, and Y-axis in the

downsampled output. B: AUROCs of the model with decreasing read numbers, calculated

using gene expression (black line) or aggregated gene expression using the WGCNA (green

line) or VIPER (red line) algorithms.

Figure 6. Performance as AUROC of gbm models to predict mutations in CROP-Seq datasets

using gene expression (red bars) and VIPER activity (blue bars) derived from CCLE expression

data in Datlinger (A) and Shifrut (B) datasets. The p-value of paired Wilcoxon tests between

all VIPER and Expression AUROCs in each dataset is reported, as well as the average of all

expression models (red solid line) and all VIPER activity models (blue dashed line). Error bars

report the standard deviation of 100 AUROCs generated from multiple partitioning of training/test sets.

Acknowledgments

We acknowledge the CINECA award (projects HP10CB1R7T and HP10CPQJBV) under the ISCRA initiative, for the support and availability of high-performance computing resources. We also thank Lupo Giorgi, Dr. Luca Pestarino and Jordan Pflugh Kraft for the fruitful discussions.

References

[1] Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. Cell 2011;144:646–74. doi:10.1016/j.cell.2011.02.013.

[2] Meacham CE, Morrison SJ. Tumour heterogeneity and cancer cell plasticity. Nature 2013;501:328–37. doi:10.1038/nature12624.

[3] Bozic I, Antal T, Ohtsuki H, Carter H, Kim D, Chen S, et al. Accumulation of driver and passenger mutations during tumor progression. Proc Natl Acad Sci U S A 2010;107:18545–50. doi:10.1073/pnas.1010978107.

[4] Ciriello G, Miller ML, Aksoy BA, Senbabaoglu Y, Schultz N, Sander C. Emerging landscape of oncogenic signatures across human cancers. Nat Genet 2013;45:1127–33. doi:10.1038/ng.2762.

[5] Chang K, Creighton CJ, Davis C, Donehower L, Drummond J, Wheeler D, et al. The Cancer Genome Atlas Pan-Cancer analysis project. Nature 2013.

[6] Brennan CW, Verhaak RGW, McKenna A, Campos B, Noushmehr H, Salama SR, et al. The Somatic Genomic Landscape of Glioblastoma. Cell 2013;155:462–77. doi:10.1016/j.cell.2013.09.034.

[7] Cancer Genome Atlas Network. Genomic Classification of Cutaneous Melanoma. Cell 2015;161:1681–96. doi:10.1016/j.cell.2015.05.044.

[8] Leiserson MDM, Vandin F, Wu H-T, Dobson JR, Eldridge JV, Thomas JL, et al. Pan-cancer network analysis identifies combinations of rare somatic mutations across pathways and protein complexes. Nat Genet 2015;47:106–14. doi:10.1038/ng.3168.

[9] Bailey MH, Tokheim C, Porta-Pardo E, Sengupta S, Bertrand D, Weerasinghe A, et al. Comprehensive Characterization of Cancer Driver Genes and Mutations. Cell 2018;174:1034–5. doi:10.1016/j.cell.2018.07.034.

[10] Liu Y, Zhao M. lnCaNet: pan-cancer co-expression network for human lncRNA and cancer genes. Bioinformatics 2016;32:1595–7. doi:10.1093/bioinformatics/btw017.

[11] Osmanbeyoglu HU, Toska E, Chan C, Baselga J, Leslie CS. Pancancer modelling predicts the context-specific impact of somatic mutations on transcriptional programs. Nat Commun 2017;8:14249. doi:10.1038/ncomms14249.

[12]   Kim H, Kim Y-M. Pan-cancer analysis of somatic mutations and transcriptomes reveals common functional gene clusters shared by multiple cancer types. Sci Rep 2018;8:6041. doi:10.1038/s41598-018-24379-y.

[13]   Ghazanfar S, Yang JYH. Characterizing mutation-expression network relationships in multiple cancers. Comput Biol Chem 2016;63:73–82. doi:10.1016/j.compbiolchem.2016.02.009.

[14]   Sharma A, Jiang C, De S. Dissecting the sources of gene expression variation in a pan-cancer analysis identifies novel regulatory mutations. Nucleic Acids Res 2018;46:4370–81. doi:10.1093/nar/gky271.

[15]   Alvarez MJ, Shen Y, Giorgi FM, Lachmann A, Ding BB, Ye BH, et al. Functional characterization of somatic mutations in cancer using network-based inference of protein activity. Nat Genet 2016;48:838–47. doi:10.1038/ng.3593.

[16]   Nawy T. Single-cell sequencing. Nat Methods 2013;11:18. doi:10.1038/nmeth.2771.

[17]   Datlinger P, Rendeiro AF, Schmidl C, Krausgruber T, Traxler P, Klughammer J, et al. Pooled CRISPR screening with single-cell transcriptome readout. Nat Methods 2017;14:297–301. doi:10.1038/nmeth.4177.

[18]   Kuhn M. Building Predictive Models in R Using the caret Package. J Stat Softw 2008;028.

[19]   Maaten L van der, Hinton G. Visualizing Data using t-SNE. J Mach Learn Res 2008;9:2579–605.

[20]   Perou CM, Sørlie T, Eisen MB, van de Rijn M, Jeffrey SS, Rees CA, et al. Molecular portraits of human breast tumours. Nature 2000;406:747–52. doi:10.1038/35021093.

[21]   Not All Esophageal Tumors Equal. Cancer Discov 2017;7:238. doi:10.1158/2159-8290.CD-NB2017-006.

[22]   Lawrence MS, Stojanov P, Polak P, Kryukov GV, Cibulskis K, Sivachenko A, et al. Mutational heterogeneity in cancer and the search for new cancer-associated genes. Nature 2013;499:214–8. doi:10.1038/nature12213.

[23]   Kimura ET, Nikiforova MN, Zhu Z, Knauf JA, Nikiforov YE, Fagin JA. High Prevalence of BRAF Mutations in Thyroid Cancer: Genetic Evidence for Constitutive Activation of the RET/PTC-RAS-BRAF Signaling Pathway in Papillary Thyroid Carcinoma. Cancer Res 2003;63:1454–7.

[24]   Yan H, Parsons DW, Jin G, McLendon R, Rasheed BA, Yuan W, et al. IDH1 and IDH2 Mutations in Gliomas. N Engl J Med 2009;360:765–73. doi:10.1056/NEJMoa0808710.

[25]   Shlien A, Malkin D. Copy number variations and cancer. Genome Med 2009;1:62. doi:10.1186/gm62.

[26]   Chen JC, Alvarez MJ, Talos F, Dhruv H, Rieckhof GE, Iyer A, et al. Identification of Causal Genetic Drivers of Human Disease through Systems-Level Analysis of Regulatory Networks. Cell 2014;159:402–14. doi:10.1016/j.cell.2014.09.021.

[27]   Bass AJ, Watanabe H, Mermel CH, Yu S, Perner S, Verhaak RG, et al. SOX2 is an amplified lineage-survival oncogene in lung and esophageal squamous cell carcinomas. Nat Genet 2009;41:1238–42. doi:10.1038/ng.465.

[28]   Bell DW, Lynch TJ, Haserlat SM, Harris PL, Okimoto RA, Brannigan BW, et al. Epidermal Growth Factor Receptor Mutations and Gene Amplification in Non–Small-Cell Lung Cancer: Molecular Analysis of the IDEAL/INTACT Gefitinib Trials. J Clin Oncol 2005;23:8081–92. doi:10.1200/JCO.2005.02.7078.

[29]   Momand J, Jung D, Wilczynski S, Niland J. The MDM2 gene amplification database. Nucleic Acids Res 1998;26:3453–9. doi:10.1093/nar/26.15.3453.

[30]   Colombo T, Farina L, Macino G, Paci P. PVT1: a rising star among oncogenic long noncoding RNAs. BioMed Res Int 2015;2015:304208. doi:10.1155/2015/304208.

[31]   Li X, Chen W, Wang H, Wei Q, Ding X, Li W. Amplification and the clinical significance of circulating cell-free DNA of PVT1 in breast cancer. Oncol Rep 2017;38:465–71. doi:10.3892/or.2017.5650.

[32]   Usvasalo A, Savola S, Räty R, Vettenranta K, Harila-Saari A, Koistinen P, et al. CDKN2A deletions in acute lymphoblastic leukemia of adolescents and young adults—An array CGH study. Leuk Res 2008;32:1228–35. doi:10.1016/j.leukres.2008.01.014.

[33]   Mistry M, Zhukova N, Merico D, Rakopoulos P, Krishnatry R, Shago M, et al. BRAF Mutation and CDKN2A Deletion Define a Clinically Distinct Subgroup of Childhood Secondary High-Grade Glioma. J Clin Oncol 2015;33:1015–22. doi:10.1200/JCO.2014.58.3922.

[34]   Zhao D, Lu X, Wang G, Lan Z, Liao W, Li J, et al. Synthetic essentiality of chromatin remodelling factor CHD1 in PTEN-deficient cancer. Nature 2017;542:484–8. doi:10.1038/nature21357.

[35]   Wang X, Cao X, Sun R, Tang C, Tzankov A, Zhang J, et al. Clinical Significance of PTEN Deletion, Mutation, and Loss of PTEN Expression in De Novo Diffuse Large B-Cell Lymphoma. Neoplasia 2018;20:574–93. doi:10.1016/j.neo.2018.03.002.

[36]   Robertson AG, Kim J, Al-Ahmadie H, Bellmunt J, Guo G, Cherniack AD, et al. Comprehensive Molecular Characterization of Muscle-Invasive Bladder Cancer. Cell 2017;171:540-556.e25. doi:10.1016/j.cell.2017.09.007.

[37]   Fawcett T. An introduction to ROC analysis. Pattern Recognit Lett 2006;27:861–74. doi:10.1016/j.patrec.2005.10.010.

[38]   Friedman JH. Greedy Function Approximation: A Gradient Boosting Machine. Ann Stat 2001;29:1189–232.

[39]   Cole AJ, Dwight T, Gill AJ, Dickson K-A, Zhu Y, Clarkson A, et al. Assessing mutant p53 in primary high-grade serous ovarian cancer using immunohistochemistry and massively parallel sequencing. Sci Rep 2016;6:26191. doi:10.1038/srep26191.

[40]   Cicenas J, Kvederaviciute K, Meskinyte I, Meskinyte-Kausiliene E, Skeberdyte A, Cicenas J. KRAS, TP53, CDKN2A, SMAD4, BRCA1, and BRCA2 Mutations in Pancreatic Cancer. Cancers 2017;9. doi:10.3390/cancers9050042.

[41]   Prior IA, Lewis PD, Mattos C. A comprehensive survey of Ras mutations in cancer. Cancer Res 2012;72:2457–67. doi:10.1158/0008-5472.CAN-11-2612.

[42]   Futreal PA, Coin L, Marshall M, Down T, Hubbard T, Wooster R, et al. A census of human cancer genes. Nat Rev Cancer 2004;4:177–83. doi:10.1038/nrc1299.

[43]   Sager R. Tumor suppressor genes: the puzzle and the promise. Science 1989;246:1406–12. doi:10.1126/science.2574499.

[44]   Irizarry RA, Wu Z, Jaffee HA. Comparison of Affymetrix GeneChip expression measures. Bioinformatics 2006;22:789–94. doi:10.1093/bioinformatics/btk046.

[45]   Langfelder P, Horvath S. WGCNA: an R package for weighted correlation network analysis. BMC Bioinformatics 2008;9:559. doi:10.1186/1471-2105-9-559.

[46]   Lachmann A, Giorgi FM, Lopez G, Califano A. ARACNe-AP: gene network reverse engineering through adaptive partitioning inference of mutual information. Bioinforma Oxf Engl 2016;32:2233–5. doi:10.1093/bioinformatics/btw216.

[47]   Pollen AA, Nowakowski TJ, Shuga J, Wang X, Leyrat AA, Lui JH, et al. Low-coverage single-cell mRNA sequencing reveals cellular heterogeneity and activated signaling

pathways in developing cerebral cortex. Nat Biotechnol 2014;32:1053–8. doi:10.1038/nbt.2967.

[48] Risso D, Perraudeau F, Gribkova S, Dudoit S, Vert J-P. A general and flexible method for signal extraction from single-cell RNA-seq data. Nat Commun 2018;9. doi:10.1038/s41467-017-02554-5.

[49] Bush EC, Ray F, Alvarez MJ, Realubit R, Li H, Karan C, et al. PLATE-Seq for genome-wide regulatory network analysis of high-throughput screens | Nature Communications. Nat Commun 2017;8. doi:doi.org/10.1038/s41467-017-00136-z.

[50] Ran FA, Hsu PD, Wright J, Agarwala V, Scott DA, Zhang F. Genome engineering using the CRISPR-Cas9 system. Nat Protoc 2013;8:2281–308. doi:10.1038/nprot.2013.143.

[51] Shifrut E, Carnevale J, Tobin V, Roth TL, Woo JM, Bui CT, et al. Genome-wide CRISPR Screens in Primary Human T Cells Reveal Key Regulators of Immune Function. Cell 2018;175:1958-1971.e15. doi:10.1016/j.cell.2018.10.024.

[52] Tirosh I, Izar B, Prakadan SM, Wadsworth MH, Treacy D, Trombetta JJ, et al. Dissecting the multicellular ecosystem of metastatic melanoma by single-cell RNA-seq. Science 2016;352:189–96. doi:10.1126/science.aad0501.

[53] Barretina J, Caponigro G, Stransky N, Venkatesan K, Margolin AA, Kim S, et al. The Cancer Cell Line Encyclopedia enables predictive modelling of anticancer drug sensitivity. Nature 2012;483:603–7. doi:10.1038/nature11003.

[54] Giorgi FM, Del Fabbro C, Licausi F. Comparative study of RNA-seq- and microarray-derived coexpression networks in Arabidopsis thaliana. Bioinforma Oxf Engl 2013;29:717–24. doi:10.1093/bioinformatics/btt053.

[55] Lawrence M, Huber W, Pagès H, Aboyoun P, Carlson M, Gentleman R, et al. Software for Computing and Annotating Genomic Ranges. PLOS Comput Biol 2013;9:e1003118. doi:10.1371/journal.pcbi.1003118.

[56] Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene Ontology: tool for the unification of biology. Nat Genet 2000;25:25–9. doi:10.1038/75556.

[57] Duong T. ks: Kernel Density Estimation and Kernel Discriminant Analysis for Multivariate Data in R. J Stat Softw 2007;021.

[58] Gautier L, Cope L, Bolstad BM, Irizarry RA. affy—analysis of Affymetrix GeneChip data at the probe level. Bioinformatics 2004;20:307–15. doi:10.1093/bioinformatics/btg405.

[59] Hanley JA, McNeil BJ. The meaning and use of the area under a receiver operating characteristic (ROC) curve. Radiology 1982;143:29–36. doi:10.1148/radiology.143.1.7063747.

[60] Satija R, Farrell JA, Gennert D, Schier AF, Regev A. Spatial reconstruction of single-cell gene expression data. Nat Biotechnol 2015;33:495–502. doi:10.1038/nbt.3192.

Supporting Information Legends

Figure S1. Table of most somatically mutated genes across TCGA tumor samples, in terms of number of samples where the gene is somatically mutated with altered protein product sequence. This table includes also MutSig-blacklisted genes (in grey) such as Titin (TTN), Obscurin (OBSCN) and Mucin genes.

Figure S2. Relationship between alteration models and alteration frequency in the Pan-cancer dataset, for mutations (left), amplifications (center) and deletions (right).

Figure S3. Performance of Pan-cancer alterations models globally (left) and for MutSig genes, COSMIC oncogenes and COSMIC tumor suppressors. The Y-Axis indicates rank-transformed AUROC values. Asterisks indicate a significant (<0.01) difference between a distribution and the global "Other Genes" distribution according to Two-tailed Wilcoxon tests.

Figure S4. ROC curves for gbm TP53 models in Breast Cancer, using original expression data, VIPER aggregation (TF "activity") and WGCNA aggregation (robust tukey biweight average of clusters).

Figure S5. AUROCs of EGFR amplication gbm prediction models in Glioblastoma with increasing noise, calculated using gene expression (black line) or aggregated gene expression using the WGCNA (green line) or VIPER (red line) algorithms.

Figure S6. AUROCs of EGFR amplication gbm prediction models in Lung Squamous Carcinoma (LUSC) with increasing noise, calculated using gene expression (black line) or aggregated gene expression using the WGCNA (green line) or VIPER (red line) algorithms.

Figure S7. AUROCs of PVT1 amplication gbm prediction models in Ovarian Cancer with

increasing noise, calculated using gene expression (black line) or aggregated gene

expression using the WGCNA (green line) or VIPER (red line) algorithms.

Figure S8. AUROCs of PTEN deletion gbm prediction models in Sarcoma with increasing

noise, calculated using gene expression (black line) or aggregated gene expression using the

WGCNA (green line) or VIPER (red line) algorithms.

Figure S9. Beta distribution used to down-sample the 43.8M reads breast cancer sample

TCGA-A1-A0SB-01 to 10M reads. The grey line shows the ratio between the target coverage

and the original coverage.

Figure S10. TSNE representation of the Datlinger CROP-Seq dataset before (A) and after (B)

removal of cell-cycle specific markers. Colors indicated the predicted cell cycle phase

according to the Seurat pipeline [60].

Supplementary Table S1. AUROCs for each event in the Pan-Cancer TCGA dataset (24 tumor

types with at least 100 samples with co-measured genomic and expression data. The Sheet

name indicates the tumor type and genomic alteration type (mut: somatic mutation, amp:

amplification, del: deletion).

Supplementary Code. R and bash code snippets used in this study.