Polysaccharide A-dependent opposing effects of mucosal and systemic exposures to human gut commensal *Bacteroides fragilis* in type 1 diabetes

M. Hanief. Sofi[1], Benjamin M. Johnson[1], Radhika R. Gudi[1], [1], Amy Jolly[1], Marie-Claude Gaudreau[1], and Chenthamarakshan Vasu[1,3]

[1]Microbiology and Immunology, College of Medicine, Medical University of South Carolina, Charleston, SC-29425

[3]Surgery, College of Medicine, Medical University of South Carolina, Charleston, SC-29425

Running title: *Bacteroides fragilis* in type 1 diabetes.

[5]Address Correspondence: Chenthamarakshan Vasu
Microbiology and Immunology
Medical University of South Carolina
173 Ashley Avenue, BSB208E
Charleston, SC-29425
Phone: 843-792-1032, Fax: 843-792-9588
Email: vasu@musc.edu

Abstract

Bacteroides fragilis (BF), a Gram-negative bacterium, is an integral component of the human colonic commensal microbiota. BF is also the most commonly isolated organism from clinical cases of intra-abdominal abscesses suggesting its potential to induce pro-inflammatory responses, upon accessing the systemic compartment. Hence, we examined the impact of intestinal and systemic exposure to BF on type 1 diabetes (T1D) incidence in non-obese diabetic (NOD) mice. The impact of intestinal exposure to BF under chemically-induced enhanced gut permeability condition, which permits microbial translocation to systemic compartment, in T1D was also examined. While oral administration of pre-diabetic mice with heat-killed (HK) BF caused enhanced immune regulation and significant suppression of autoimmunity resulting in delayed hyperglycemia, mice that received small amounts of HK BF by i.v. injection showed pro-inflammatory immune response and rapid disease progression. Importantly, polysaccharide-A deficient (ΔPSA) BF failed to produce these opposing effects upon oral and systemic deliveries. Further, BF induced modulation of disease progression was observed in WT, but not TLR2-deficient, NOD mice. Interestingly, oral administration of BF under chemically-induced enhanced gut permeability condition resulted in accelerated disease progression and rapid onset of hyperglycemia in NOD mice. Overall, these observations suggest that BF-like gut commensals can cause pro-inflammatory responses upon gaining access to systemic compartment and trigger/accelerate autoimmunity and contribute to T1D in at-risk subjects.

Introduction

While gut commensal microbes contribute profoundly to human health and immune system maturation[1-4], many recent reports also support the notion that gut commensals can contribute to the pathogenesis of autoimmune diseases like type 1 diabetes (T1D)[5-8]. It has been proposed that three mutually linked features, viz: 1) aberrant gut microbiota, 2) compromised intestinal epithelial barrier, and 3) altered immune responsiveness are linked to the contribution of commensal organisms to these diseases[9]. In addition to many environmental and genetic factors, gut pathobionts and pro-inflammatory immune responses can compromise gut integrity leading to the translocation of microbial components from the intestine to systemic compartment, and consequently systemic inflammation and autoimmune progression[10-14].

Bacteroides fragilis (BF)*,* a Gram-negative anaerobe and an integral component of the colonic commensal microbiota of many mammals[15], has been widely studied as a model organism to determine the influence of commensal bacteria in immune regulation[16-18]. Studies using different pre-clinical models of various inflammatory diseases have demonstrated many health benefits of this bacterium and its capsular polysaccharide[16,17,19,20]. Although *Bacteroides* members are highly abundant in the human intestine, BF represents a very small portion of the gut microbes[21]. Nevertheless, BF is the most commonly isolated bacterium from clinical cases of intra-abdominal abscesses[22,23], perhaps due to its ability to colonize the colonic crypt[16]. It has been shown that capsular polysaccharide of BF is necessary and sufficient for abscess formation[23-25]. While capsular polysaccharides are also found to be essential for its growth[26], these components, polysaccharide A (PSA) in particular, have been demonstrated to have the ability to promote immune regulation by directly acting on dendritic cells (DCs) and regulatory T cells (Tregs) in a TLR2 dependent manner[16,27-29]. In addition, it can modulate inflammation and suppress colitis and EAE in rodent models[20,30]. Overall, these paradoxical effects of BF in intestinal immune regulation and peritoneal abscess suggest that intestinal mucosa and systemic compartments may produce different responses upon exposure to this commensal. Importantly, many recent studies have associated higher prevalence of Bacteroidetes phylum members to T1D disease incidence / progression in humans and pre-clinical models[6,7,31-37]. However, the direct impact of BF on T1D disease has not been reported.

Here, using heat-killed (HK) BF, we show that intestinal and systemic exposures to this bacterium have opposing effects on autoimmune progression and T1D incidence in the non-obese diabetic (NOD) mouse model. While oral administration of pre-diabetic mice with heat-killed (HK) BF caused enhanced immune regulation and significant suppression of autoimmunity resulting in delayed hyperglycemia, mice that received small amounts of HK BF by i.v. injection showed pro-inflammatory immune response and rapid disease progression. Interestingly, only the wild-type (WT), but not PSA deficient (ΔPSA), BF produced opposing immune responses and disease outcomes upon oral and systemic administrations. Furthermore, these effects were observed only in WT, but not TLR2 deficient, NOD mice. Most importantly, oral administration of NOD mice with BF under enhanced gut permeability recapitulated the T1D accelerating effect of systemic administration with HK BF. These results suggest that systemic access by host beneficial gut symbiont such as BF, in the event of compromised gut barrier function, can act as a trigger and a catalyst for systemic inflammation and autoimmunity in T1D at-risk subjects.

Results

Exposure of gut mucosa to BF results in suppression of autoimmune progression and delayed hyperglycemia.

NOD mice show infiltration of immune cells into the pancreatic islets at as early as 4-weeks of age and it progresses to >90% of the insulin producing islets being completely destroyed by the time the mice turn diabetic. Hyperglycemia is detected in female NOD mice as early as 12 weeks of age and 80-100% of the mice turn overt hyperglycemic before the age of 30 weeks. Here, we used 10-week-old pre-diabetic NOD mice to determine the impact of oral administration of HK BF on autoimmune progression and T1D incidence. As shown in Fig. 1A, oral administration of NOD mice with HK BF (500 µg/day) for 15 days caused a significant delay in hyperglycemia and overall T1D incidence. While more than 80% of the mice remained diabetes free at 15 weeks post-treatment, more than 60% of the control mice developed diabetes within this period. To determine the impact of orally administered BF on islet autoimmunity, one set of euglycemic mice were euthanized 30 days post-treatment and the pancreatic tissues were examined for degree of insulitis. Delayed onset and low incidence of the disease in BF treated mice were

found to correlate with suppressed immune cell infiltration and a profoundly higher number of islets with no or modest immune cell infiltration (grades ≤2 insulitis) (Fig. 1B).

To determine if the autoreactivity of immune cells is modulated in mice that were orally administered with HK BF, PnLN cells were activated ex vivo using β-cell antigen peptide cocktail. PnLN cells from BF treated mice produced significantly higher amounts of IL10, and showed profoundly higher IL10 to IFNγ production ratio, compared to these cells from control mice (Fig. 1C). Oral BF treatment induced modulation of systemic immune function was also assessed by examining the splenic T cells for their expression of cytokines and Foxp3. As shown in Fig. 1D, IFNγ+ T cell frequencies were lower and Foxp3+ cell frequencies were higher in BF treated mice. Further, overall IL10+:IFNγ+ and Foxp3+:IFNγ+ cell ratios were significantly higher in these mice compared to controls. To further assess if oral administration of BF impacts the diabetogenic function of systemic immune cells, splenic T cells from BF treated mice were adoptively transferred to NOD-*Rag1* mice. Fig. 1E shows that NOD-*Rag1* mice that received T cells from BF-fed mice developed hyperglycemia at a significantly slower rate compared to mice that received T cells from control mice. Overall, these observations suggest that exposure of gut mucosa to BF results in enhanced immune regulation in the systemic compartment, including pancreatic microenvironment, resulting in suppression of autoimmunity, insulitis and T1D incidence in NOD mice.

Exposure of systemic compartment to BF results in enhanced pro-inflammatory response and accelerated T1D onset

Since BF is known to cause peritonitis, and intra-abdominal infections and abscesses upon gaining access to these compartments[22-25,38,39], we determined the effect of systemic exposure to HK BF on autoimmune progression and T1D incidence. Ten-week-old pre-diabetic NOD mice were treated i.v. with small amounts of BF (10 µg/day) every 3 days for 15 days and monitored for hyperglycemia. As observed in Fig. 2A, mice that received i.v. injection of HK BF showed, compared to controls, rapid onset of hyperglycemia and significantly high overall diabetic incidence. Consistent with this observation, mice that were treated i.v. showed profoundly higher degree of insulitis at 30 days post-treatment (Fig. 2B). Importantly, PnLN cells from mice that received systemic treatment showed significantly high IFNγ production, and overall low IL10 to

IFNγ production ratio, upon ex vivo activation using β-cell antigen peptide cocktail, compared to cells from control mice (Fig. 2C). Further, compared to control mice, significantly higher IFNγ+ and lower Foxp3+ T cell frequencies were detected in the spleen of mice that were exposed to BF systemically (Fig. 2D). In addition, *NOD-Rag1* mice that received splenic T cells from BF injected mice showed relatively faster onset of hyperglycemia, albeit not significant statistically, compared to those that received cells from control mice (Fig. 2E).

The systemic effects of i.v. injection with BF in NOD mice (Fig. 2) are opposite to that produced by oral treatment with this agent (Fig. 1). To determine if such contrasting effects are produced by other bacteria, pre-diabetic NOD mice were treated with HK *Listeria monocytogenes* (LM) by oral gavage or i.v. injection and examined for T1D incidence and insulitis. As observed in Fig. S1, orally and systemically administered HK LM produced disease incidence and insulitis trends in NOD mice similar to that of HK BF treated mice, however these observations were, unlike that from BF treated mice, not significant statistically by log-rank test and Fisher's exact test. Overall, above observations suggest that certain gut bacteria, including commensals, trigger opposing immune responses in the gut mucosa and systemic compartment and, under genetic susceptibility, can impact the autoimmune disease process differently.

Orally and systemically administered BF induces distinct immune responses in the gut mucosa and spleen.

Since NOD mice that were treated with HK BF orally and systemically showed opposing effects on insulitis and hyperglycemia onset, intestinal and splenic immune responses of mice that were exposed to BF for 3 days were determined. qPCR assay of distal ileum and distal colon from mice that received BF orally and spleen cells from mice that received BF systemically show that BF exposure triggers the expression of mRNA for a large number of cytokines, chemokines, and immune regulatory enzyme such as Aldh1A2 in both systemic and intestinal compartments. Notably, compared to controls, higher *Il10* mRNA expression was induced in BF exposed intestine as well as spleen. However, most intriguingly, *Il6* expression was profoundly higher in the spleen of mice that received i.v. injection, but significantly low in the intestine of oral treatment recipients, of BF compared to their control counterparts. Further, BF exposure associated *Il12* and *Il1b* mRNA expressions were significantly higher in the spleen of i.v. injected mice, but not in the intestine of orally fed mice.

To determine the impact of BF induced intestinal and systemic immune responses on T cell phenotype, mesenteric lymph nodes (MLN) from NOD-Foxp3-GFP mice that received oral administration and spleens from mice that received systemic administration of BF for 3 days were compared for Foxp3 (GFP) and intracellular cytokine positive T cell frequencies on day 7. As observed in Fig. 3B, while MLN of mice that received BF orally showed significantly higher frequencies of GFP+, IL10+ and IL17+ T cells compared to their control counterparts, spleens of mice that received systemic BF showed profoundly lower frequencies of GFP+ cells, and higher frequencies of IFNγ+ cells compared to controls. Further, mice that received BF systemically for 3 days also showed higher splenic IL17+ and IL10+ cell frequencies. In a similar experiment, mice that were subjected to prolonged (15 day) oral and systemic treatments as described for Figs. 1 and 2, were euthanized within 24 h and examined for T cell phenotype. Fig. S2 shows significantly higher Foxp3+ and IL10+, and lower IFNγ+ CD4+ cells, in the gut associated lymphoid tissues (GALT) of mice that received prolonged treatment. On the other hand, prolonged systemic treatment caused significantly reduced Foxp3+ and IL10+, and increased IFNγ+, CD4+ T cell frequencies in the spleen. Overall, these observations, along with the cytokine expression profile data of Fig. 3A, suggest an overall regulatory immune response by gut mucosa and pro-inflammatory immune response by systemic immune cells upon exposure to BF.

BF exposed intestinal and splenic APCs produce opposing effects on Tregs upon antigen presentation ex vivo and in vivo.

Since exposure to HK BF resulted in contrasting immune profiles in the intestinal mucosa and systemic compartment, impacts of exposure to BF on intestinal and splenic antigen presenting cells and antigen specific T cell activation were determined. NOD mice were treated HK BF for 3 consecutive days and euthanized on day 4 as described for Fig. 3A. CD11c+ dendritic cells (DC) were enriched from the small intestine of BF fed and spleen of BF injected mice along with respective control mice, and used in qPCR and antigen presentation assays. Fig. 4A shows DCs isolated from the intestine of mice that received BF orally expressed significantly lower *Il6* and *Il12*, and higher *Il10* and *Tnfa* mRNA compared to DCs from control mouse intestines. On the other hand, splenic DCs of mice that received systemic treatment with BF expressed higher *Il6*, *Il12*, *Il10* and *Tnfa* mRNA compared to their counterparts from control mice.

To examine the impact of ex vivo antigen presentation by APCs that were exposed to BF in vivo, BDC2.5 peptide pulsed DCs were cultured with T cells from NOD-BDC2.5-Foxp3-GFP mice and the Foxp3 (GFP)+ T cell frequencies were determined in these cultures. As shown in Fig. 4B, while intestinal DCs from mice that received BF orally caused a significant increase in GFP+ CD4 T cells compared to these cells from the control mice, GFP+ CD4 T cell frequencies were significantly lower in cultures where splenic DCs from BF injected mice were used. In addition, as shown in Fig. S3, activation of antigen specific T cells by intestinal DCs from BF fed mice resulted in relatively lower IFNγ and significantly high IL10 release, and an overall higher IL10:IFNγ ratio, in the cultures. On the other hand, activation of T cells by splenic DCs from BF injected mice resulted in higher IFNγ and lower IL10 release, and profoundly low IL10 to IFNγ ratio, in the cultures.

To determine the impact of antigen presentation by BF exposed APCs in vivo, NOD-BDC2.5-Foxp3-GFP mice were treated with BDC2.5 peptide alone or along with BF orally or systemically as detailed in Fig. 4B, and examined for GFP+ CD4 T cell frequencies in GALT and spleen respectively. Compared to the BDC2.5 peptide alone recipients, mice that received oral gavage of BDC2.5 peptide along with BF showed significantly higher frequencies of GFP+ CD4 cells in the GALT including MLN, PP and SiLP. However, mice that received systemic injection of BDC2.5 peptide along with BF had profoundly diminished frequencies of GFP+ CD4 T cells in the spleen than those that received BDC2.5 peptide alone. These observations, in association with the results on T1D incidences in BF treated mice (shown in Figs. 1 & 2), indicate that while BF- induced immune response in the gut mucosa promotes the generation of T cells with regulatory properties leading to eventual suppression of pancreatic autoimmunity, systemic immune response against BF promotes generation of pro-inflammatory T cells and suppression of Treg function contributing to acceleration of the disease process in T1D.

Oral or systemic administration of PSA deficient (ΔPSA)-BF failed to impact autoimmune progression and T1D incidence in NOD mice

Since PSA has been considered as the major symbiotic factor and host immune modulator of BF[16,19,40,41], we determined if isogenic ΔPSA mutant strain[25] of BF produces immune responses in the intestinal and systemic compartments similar to that by BF. Cohorts of mice were treated orally or systemically with HK BF and ΔPSA-BF for 3 days and cytokine expression profiles in

the intestine and spleen respectively were determined by qPCR as described for Fig. 3A. Fig. S4 shows that mice that received ΔPSA-BF orally for 3 days, compared to BF recipients, showed significantly higher expression of *Il6* and *Il12* mRNA, and lower expression of *Il10* mRNA in the intestine. On the other hand, spleen cells of mice that received ΔPSA-BF systemically showed diminished expression of mRNA for most cytokines, albeit not statistically significant except for *Il10*, and profoundly increased expression of *Il23* compared to spleen cells from BF recipient counterparts.

To determine the intestinal and systemic impacts of PSA deficiency in BF on T1D incidence, pre-diabetic female NOD mice were given HK BF or its ΔPSA mutant by oral gavage or by i.v. injection for 15 days as described for Figs. 1 and 2. Pancreatic tissues from cohorts of these mice euthanized 30 days post-treatment were examined for the degree of immune cell infiltration and islet function in terms of insulin production. Fig. 5A shows, as observed for Fig. 1B, significant suppression of insulitis as well as higher frequencies of insulin positive islets in mice that were orally administered with BF. Mice that received BF systemically showed, as observed for Fig. 2B, more severe insulitis as well as lower frequencies of insulin positive islets. Interestingly, mice that received ΔPSA orally and systemically showed islet function and immune cell infiltration comparable to that of control mice. Examination of PnLN cells showed that Foxp3+ T cell frequencies were significantly increased upon oral treatment and diminished upon systemic administration compared to controls only in BF, but not ΔPSA-BF, recipient mice (Fig. 5B). Further, only PnLN cells from BF, but not ΔPSA-BF, recipient mice showed significant differences in the β-cell antigen activation induced release of IL10 and IFNγ as compared to these cells from control mice (Fig. 5C). In concurrence with these observations on immune cell phenotypes, insulitis and β-cell function, timing of hyperglycemia and the disease incidence rate in NOD mice that received ΔPSA-BF orally and systemically were comparable to that of their control counterparts (Fig. 5D). Overall, these studies using HK ΔPSA-BF suggest that PSA is the primary factor responsible for the direct effect of BF induced opposing responses of gut mucosa and systemic immune cells, and the overall disease outcomes.

TLR2 deficient NOD mice, unlike WT mice, failed to show opposing T1D modulatory effects upon systemic and oral administration of BF.

Interaction of BF-PSA with TLR2 plays a key role in the symbiotic function of BF[16]. Therefore, the contribution of TLR2 in protecting NOD mice from T1D upon oral administration and accelerating the disease progression upon systemic administration of HK BF was determined using NOD-TLR2-KO mice. Ten week old TLR2 KO and WT NOD mice were given HK BF orally or i.v. as described for Figs. 1 and 2 and monitored for hyperglycemia. Fig. S5 shows that, as reported by others[5,8,42], NOD-TLR2 KO mice developed hyperglycemia at a relatively lower pace than that by WT mice. However, unlike WT NOD mice, neither systemic nor oral treatment with BF produced a modulatory effect on the onset of hyperglycemia or T1D incidence in TLR2 KO mice. Similarly, examination of pancreatic tissues of similarly treated mice 30 days post-treatment showed that control and BF treated NOD-TLR2KO mice had comparable insulitis. These results, in conjunction with the data of figure 6, suggest a direct mechanistic role for PSA-TLR2 interaction in the opposing T1D modulatory effects observed upon exposure of gut mucosa and systemic compartment to BF.

Oral administration of HK BF under enhanced gut permeability results in accelerated T1D.

Compromised gut epithelial barrier function and translocation of microbial components to the systemic compartment can occur under many circumstances including stress and medications. Recent reports have suggested that leaky-gut and microbial translocation may contribute to initiation and/or progression of different autoimmune conditions[12-14,43,44]. Since systemic administrations of HK BF caused rapid disease onset in NOD mice, we determined the impact of oral administration of this agent under chemical induced enhanced gut permeability conditions. Pre-diabetic NOD mice were given low-dose Dextran sulfate sodium (DSS) in drinking water to cause injury to epithelial barrier and to increase gut permeability, followed by oral administration of HK BF. DSS treated mice showed higher translocation of orally administered FITC-dextran as well as microbial DNA in to blood (Fig. S6) suggesting that they have higher gut permeability. While HK BF recipients, as observed in Fig.1, showed protection from T1D, DSS treatment alone did not impact the disease progression in NOD mice of our SPF facility (Fig. 6A). However, compared to mice that received DSS alone, mice that received both DSS and HK BF showed rapid progression of the disease and early onset of hyperglycemia. Further, pancreas of

mice that received both DSS and BF showed significantly higher number of islets with severe insultis compared to DSS recipient controls (Fig. 6B). Examination of PnLN cells revealed significantly low Foxp3+ T cell frequency (Fig. 6C) and profoundly higher β-cell antigen induced IFNγ production (Fig. 6D) in DSS and BF treated mice compared to DSS treated controls. Overall, these results suggest that while transient translocation of gut microbial components normally may not have a profound impact on autoimmune progression, systemic access to specific gut microbes such as BF, especially under gut permeability compromised conditions, can accelerate the disease process.

Discussion

Many years of studies have demonstrated unique immune-modulatory/-regulatory properties and symbiotic nature of a key human colonic bacterium belonging to Bacteroidetes phylum, BF[16-18,45,46]. However, BF is also the primary microbe detected in most clinical intra-abdominal infections/abscesses[22,23,38,39]. This suggested to us that, while BF-like commensals may promote immune regulation when restricted to the gut, their escape to the systemic compartment could produce strong pro-inflammatory response and may impact autoimmune progression. Here, we tested this notion, in a NOD mouse model of T1D, using HK BF and employing oral and i.v. administrations to mimic gut mucosa and systemic exposures to this organism. We also employed a chemical approach to establish enhanced gut permeability and test the translocation of orally administered BF components to the systemic compartment for assessing the potential impacts of systemic exposure to BF on autoimmunity. We show that, in contrast to exposure of gut mucosa to BF which produced protection from T1D, exposure of systemic compartment to BF resulted in pro-inflammatory response, rapid insulitis progression and early onset of hyperglycemia. Using a PSA deficient mutant of BF and NOD-TLR2-KO mice, we also show that these opposing effects of BF on T1D upon oral and systemic administrations are primarily bacterial PSA and host TLR2 dependent. Our observations suggest that higher prevalence of BF-like commensals in the gut of T1D prone subjects can allow the translocation of bacteria as well as their polysaccharide like factors to the systemic compartment, under various leaky-gut inducing conditions such as stress, medications, infection and inflammation[47-52], and perpetuate systemic inflammation and autoimmunity.

T1D is an autoimmune disorder resulting from destruction of pancreatic β-cells in genetically susceptible subjects and both genetic and environmental factors contribute to it. While it has now been established that the gut microbiota is an important environmental factor that modulates the disease outcomes[2,5,9,36,53-55], the impacts of specific microbial communities and factors on T1D disease initiation and progression are not fully understood. Studies in human subjects and mouse models that have shown the occurrence of dysbiosis very close to disease onset[32,56] and the impacts of alterations in gut microbial communities caused by dietary manipulations and antibiotic interventions on T1D incidence[36,57-62] have helped identify pro-diabetic changes in the microbiota. In general, higher abundance of Bacteroidetes phyla members and a reduction in firmicutes are associated with T1D[31-37,57,63-65]. However, protection from T1D has also been positively correlated with higher abundance of multiple species belonging to Bacteroidetes phylum, Provetella genus particularly[32,34]. The fact that BF, which is known to colonize the colonic crypt and promote gut immune regulation[16,26], is the most commonly isolated gut bacterium in clinical intra-abdominal infections and inflammation suggests its potential negative impact on autoimmune outcomes, especially upon reaching systemic compartment, in T1D at-risk subjects. In fact, higher gut permeability and microbial translocation have been detected in T1D-prone individuals long before the disease onset[13,66,67], suggesting that BF could be one of the major gut commensals which gains access to the systemic compartment in them. However, it is not known if and how the systemic immune response to this beneficial gut bacterium contributes to autoimmunity in T1D. In this regard, our observations from studies using HK BF in NOD mice, for the first time, show that exposure of the systemic compartment to this agent leads to a profound pro-inflammatory response, diminished Treg function and accelerated autoimmune progression. Our observations from studies using HK BF, in fact, substantiate some of the previously reported immune regulatory features of gut colonization by live BF[16,30]; however, suggests the adverse effects of this bacterium in T1D prone subjects upon reaching the systemic compartment.

The ability of gut and systemic immune cells to produce different types of immune responses has extensively been studied in the past and the gut mucosa is generally considered tolerant to commensals and dietary antigens[68,69]. In fact, oral tolerogenic vaccination strategies using self-antigens including insulin have been employed to induce antigen specific tolerance[70-72]. Hence,

although the impacts on T1D disease outcomes are intriguing, our observation that distinct immune responses are induced against HK BF by mucosal and systemic immune compartments is not surprising. However, it is important to note that not all gut microbial factors promote T1D upon exposing the systemic compartment to them. It has been shown that pathogenic bacteria and their components such as lipopolysaccharide (LPS) (TLR4 ligand) injection induce protection from T1D in NOD mice[73-75]. Further, modest and no impact of systemically administered HK LM and mutant BF respectively on T1D in NOD mice suggest that our observation on accelerated autoimmune process upon systemic administration is, perhaps, unique to BF-like microbes and specific molecular interactions. This notion has been validated further by our observation that enhanced gut permeability and translocation of components of normal gut microbiota of NOD mice in our SPF facility alone does not impact T1D.

Importantly, the opposing effects of systemic and oral administrations of HK BF in the mouse model of T1D appear to be primarily PSA driven, and its likely interaction with TLR2. BF PSA interacts with TLR2 on APCs as well as T cells and is known to promote immune regulation[16,27-29]. The ability of PSA to activate conventional and plasmacytoid DCs that further activate/induce Tregs has been described[28,76]. PSA is also known to be critical for gut colonization by BF and to suppress the colonization by pathogenic bacteria[11,30,77,78]. Interestingly, it has been shown that not only does TLR2 deficiency protect NOD mice from T1D, TLR2-gut microbiota interaction promotes autoimmunity progression in NOD mice[5,8,42]. Hence, our observation that opposing effects on T1D disease outcomes upon exposure of the gut mucosa and the systemic compartment to BF is detectable only in WT, but not TLR2-KO, NOD mice validates the previous reports on the involvement of TLR2 in T1D. Our observations also suggest a paradoxical role for TLR2 signaling in the gut mucosa and in the systemic compartment.

Unlike human gut Gram-negative bacteria such as *E. coli* that produce highly immunogenic LPS, which activates TLR4 signaling[77], LPS of BF is considered a weak immune activator and does not require TLR4[78]. Recently, it has been shown that variation in microbiome and LPS immunogenicity can impact T1D in humans[6]. Immune response to *E. coli* LPS, but not the LPS of *Bacteroides* Spp, *B. dorei* particularly, is associated with early immune education and protection from T1D in young children[6]. In agreement with these reports, not only that TLR4 KO NOD mice show rapid onset of hyperglycemia[5,79], but also TLR4 binding LPS preparations from

Gram-negative bacteria protects NOD mice from T1D[73,74]. Our observations that oral or systemic administrations of PSA deficient HK BF failed to impact insulitis or hyperglycemia in NOD mice indicate, in addition to the previously reported low immunogenicity and the passive role of its BF LPS, PSA plays an active role in modulating the T1D disease outcomes.

Overall, our observations from this study which exposed the gut mucosa and thesystemic compartment of a T1D-prone mouse model in an SPF facility to HK BF suggest that certain gut symbionts such as BF, while promoting immune regulation when restricted to its natural habitat in the distal colon, can contribute to initiation and/or progression of autoimmunity in at-risk subjects upon accessing the systemic compartment. This is evident from the fact that while mice that received HK BF orally under normal conditions were protected from T1D, administration of this agent under enhanced gut permeability accelerated the disease progression. A large body of circumstantial evidence indicates that gut barrier dysfunction and microbial translocation are associated with the pathogenesis of autoimmune diseases such as T1D and MS, and gut inflammatory conditions such as IBD[9,13,43,44,67,80-82]. In humans, leaky-gut can be caused by antibiotic and other medications, enteric bacterial and viral infections, altered microbiota and stress[41,58-61]. Previous studies have linked enteric viral infections to autoimmune diseases like T1D[83-87]. With further relevance to T1D, which is initiated mostly at juvenile age, reports show infants at breast-feeding age, those on formula diet particularly, have higher gut permeability; and animal studies have shown that suckling-weanling transition results in normal gut closure[88-91]. While NOD mice housed in an SPF facility may not encounter these gut permeability inducing events, one or more of the above described factors alone may be sufficient to cause transient microbial translocation across the gut barrier and produce inflammatory effects and instigate autoimmunity in at-risk subjects. Importantly, our observations that enhanced gut permeability and microbial translocation alone may not impact the disease outcomes in NOD mice also indicate that specific molecular interactions and immune responses may be critical to influence the disease initiation and progression. Of note, higher abundance of Bacteroidetes phyla members including *Bacteroides* Spp has been detected not only in the gut of rodent models of T1D, but also in T1D patients and at-risk children who have progressed towards developing disease symptoms[6,31-37,56,57,63-65], prompting the suggestion that *Bacteroides* members could exert pro-autoimmune effects under T1D susceptibility. However, while additional studies in SPF and germ-free NOD mice using live BF and other *Bacteroides* members are needed, our results

obtained using HK BF model and previous studies on BF suggest that these bacteria play a critical role in gut and systemic immune regulation under normal circumstances, but could promote autoimmunity under the circumstances such as abnormal epithelial barrier function and higher gut permeability.

Materials and Methods

Mice and Bacteria

Wild-type NOD/ShiLtJ (NOD), NOD-BDC2.5 TCR transgenic (NOD-BDC2.5), and NOD-*Rag 1* deficient (NOD-*Rag1*) mice were purchased from the Jackson laboratory (Maine, USA). Foxp3-GFP-knockin (Foxp3-GFP)[92] mice in the B6 background were kindly provided by Dr. Kuchroo (Harvard Medical School). Breeding colonies of these strains were established and maintained in the pathogen free facility of the Medical University of South Carolina (MUSC). NOD-WT mice from our breeding colony were used in this study. NOD-Foxp3-GFP mice were generated by backcrossing B6-Foxp3-GFP mice to the NOD background for 12 generations. NOD-BDC2.5-Foxp3-GFP mice were generated by crossing NOD-BDC2.5 mice with NOD-Foxp3-GFP mice. NOD-TLR2 knockout (NOD-TLR2 KO) mice[5] were provided by Dr. Chervonsky (University of Chicago). All animal studies were approved by the animal care and use committee of MUSC. To detect hyperglycemia in NOD-WT, NOD-TLR2 KO and NOD-Rag1 KO mice, glucose levels in blood collected from the tail vein were determined at timely intervals using the Ascensia Micro-fill blood glucose test strips and an Ascensia Contour blood glucose meter (Bayer, USA). Mice with glucose level of >250 mg/dl for two consecutive testing were considered diabetic.

BF (ATCC 25285; NCTC 9343) and isogenic ΔPSA-BF (kindly provided by Dr. Laurie Comstock, Harvard Medical School) were cultured from single colonies in complete brain-heart infusion (BHI) medium under anaerobic condition as described before[93] for up to 72 h, diluted this initial culture 50-fold using complete medium and cultured for an additional 16 h. Bacterial cells were pelleted and washed with PBS twice by centrifugation and incubated at 65°C for 30 min for heat inactivation. Heat-killed (HK) bacterial preparations were tested for viable bacteria, if any, by plating and culturing aliquots of these preparations on BHI agar pates for 72 h. *Listeria monocytogenes* (ATCC 19111) cultured in BHI medium under aerobic condition was also processed similarly. These HK bacterial preparations were subjected to an additional quick wash

in sterile distilled water to minimize salt content, and the pellet was air-dried and dry weight was determined, before suspending in PBS at 20 mg/ml concentration (w/v), as stock suspension.

Peptide antigens, cell lines, and antibodies

Immunodominant β-cell antigen peptides [viz., 1. Insulin B (9-23), 2. GAD65 (206-220), 3. GAD65 (524-543), 4. IA-2beta (755-777), 5. IGRP (123-145), and 6. BDC2.5 TCR reactive peptide (YVRPLWVRME; referred to as BDC-peptide) were custom synthesized (Genescript Inc) and used in this study. Peptides 1-5 were pooled at an equal molar ratio and used as β-cell-Ag peptide cocktail as described in our earlier studies[94]. PMA, ionomycin, Brefeldin A, monensin, magnetic bead based T cell and dendritic cell (total and sub-population) enrichment kits, ELISA and Magnetic bead based multiplex cytokine kits and unlabeled and fluorochrome labeled antibodies, FITC-Dextran, and other key reagents were purchased from Sigma-Aldrich, BD Biosciences, eBioscience, Invitrogen, Millipore, Miltenyi Biotec, StemCell Technologies, R&D Systems, Biolegend, etc. ELISA assays were read using Biorad iMark 96 well plate-reader. Luminex technology based multiplex assays were read using FlexMap3D or BioPlex 100 instruments. Real-time PCR assays were performed using transcript-specific or bacterial 16S rDNA specific primer sets and SYBR-green PCR mastermix and iCycler IQ (Bio-Rad). Flow cytometry data was acquired using FACS Calibur, FACS Verse or CyAn-ADP instruments and analyzed by using Summit or CytoBank applications.

Treatment of mice with HK BF and peptides

Female NOD-WT mice were administered orally (500 μg/mouse/day for 15 days) or injected intravenously (i.v.; 10 μg/mouse/day every 3rd day for 15 days) with HK BF (WT or ΔPSA) and examined for blood glucose levels every week for over 30 weeks post-treatment initiation to detect hyperglycemia. Control mice were left untreated or received PBS as carrier buffer. In some experiments, mice that were treated as described above for 15 days or for 3 consecutive days, were euthanized either 24 h or 15 days post-treatment. In some experiments, NOD-WT, NOD-Foxp3-GFP, or NOD-BDC2.5-Foxp3-GFP mice were fed or i.v. injected with bacteria alone or along with β-cell antigen peptide cocktail or BDC2.5 peptide. Spleen, pancreatic LN (PnLN), Mesenteric LN (MLN), Peyer's patch (PP), small intestinal lamina propria (SiLP) and/or large intestinal lamina propria (LiLP) cells from treated and control mice were examined for cytokine responses and/or T cell phenotypes. Pancreatic tissues were examined for insulitis

after staining by H&E method, or with anti-insulin antibody. In some experiments, serum samples were subjected to Luminex multiplex assays to detect cytokine or fluorometry to detect FITC-dextran levels.

Dendritic cells (DCs), T cells and in vitro and in vivo assays

CD11c+ DCs were enriched from spleen cells using magnetic separation kits and reagents from Miltenyi Biotec and/or StemCell Technologies. Small and large intestinal lamina propria (SiLP and LiLP) cells were prepared by collagenase digestion, density gradient centrifugation for enriching immune cells[95], and followed by CD11c+ DCs by magnetic separation. T cells were enriched from spleens of treated and untreated control mice using negative selection kits from Invitrogen, Miltenyi Biotec, and/or StemCell Technologies. In antigen presentation assays, DCs that were exposed to HK-BF in vivo were pulsed with BDC2.5 peptide, washed and incubated with CD4+ T cells from NOD-BDC2.5 or NOD-BDC2.5-Foxp3-GFP mice in 96 well plates. After 4 days of culture, cells were stained for CD4 and examined for Foxp3 or GFP expression, and/or intracellular cytokines (after 4 h stimulation using PMA and ionomycin) as described in our previous reports[96,97] by FACS.

Adoptive T cell transfer experiment

Total T cells isolated from spleens of control and HK BF treated mice were transferred into 8-week old NOD-*Rag1* deficient mice (i.v.; 1×10^6 cells/mouse) and tested for blood glucose levels every week to determine the diabetogenic properties of T cells.

Induction of increased gut permeability

Mice were given 0.5% Dextran sulfate sodium (DSS) (w/v) in drinking water for 5 days and switched to regular drinking water for rest of the monitoring period. Control and DSS treated mice were given HK-BF (500 µg/mouse/day) by oral gavage every day for up to 15 days starting day 0. Cohorts of these mice were euthanized on day 6, day 30, or monitored for up to 20 weeks post-treatment. To determine gut permeability, in some experiments, DSS treated and control mice were given FITC dextran (4KD; 10 mg/mouse) on day 6 by oral gavage, euthanized after 4 h, and plasma samples were examined for FITC-Dextran concentration by flurimetry against control samples spiked with known concentration of FITC-Dextran. In some assays, whole blood collected from different groups of euthanized mice by cardiac puncture using sterile syringe into

sodium citrate vials were subjected to qPCR assay using universal as well as BF specific 16S rDNA specific primer sets to determine the levels of bacterial DNA in the circulation.

Histochemical and immunofluorescence analysis of pancreatic tissues

Pancreata were fixed in 10% formaldehyde, 5-µm paraffin sections were made, and stained with hematoxylin and eosin (H&E). Stained sections were analyzed using a 0-4 grading system as described in our earlier studies[36,96,98]. Areas that appeared to have completely lost islets were not included in this grading approach. At least 100 islets were examined for every group. In some experiments, pancreatic sections were stained using anti-insulin antibody followed by Alexa fluor 488- or 568- linked secondary antibodies and DAPI, and scored for insulitis based on DAPI-positive cells in islet areas and insulin expression. Insulitis was scored as described for H&E stained sections and insulin positive and negative islets were counted. Areas that appeared to have completely lost islets were considered as grade 5 and included in this grading approach.

Statistical analysis

Mean, SD, and statistical significance (*p-value*) were calculated using GraphPad Prism, Microsoft Excel, and/or online statistical applications. Unpaired *t*-test was employed, unless specified, for values from *in vitro* and *ex vivo* assays. Log-rank analysis (http://bioinf.wehi.edu.au/software/russell/logrank/) was performed to compare T1D incidence (hyperglycemia) of the test group with that of respective control group. Fisher's exact test was used for comparing the total number of severely infiltrated islets (grades $\geq$3) relative to total number of islets with low or no infiltration (grades $\leq$2) in test vs. control groups. A *p* value of $\leq$0.05 was considered statistically significant.

Conflict of Interest statement: Authors do not have any conflict(s) of interest to disclose.

Author contribution: M.H.S. researched and analyzed data, R.G. researched and analyzed data and edited the manuscript, B.M.J researched data, A. J. assisted in experiments, M.C.G. assisted in experiments, and C.V. designed experiments, researched and analyzed data, and wrote/edited manuscript.

Footnote: This work was supported by unrestricted research funds from MUSC and National Institutes of Health (NIH) grant R21AI133798. Dr. Vasu is the guarantor of this work and, as such, has full access to all the data in the study and takes responsibility for the integrity of the data and accuracy of the data analysis. The authors are thankful to Cell and Molecular Imaging, Pathology, Proteomics, immune monitoring and discovery, and flow cytometry cores of MUSC for the histology service, microscopy, FACS and multiplex assay instrumentation support.

Abbreviations: **TID:** Type 1 Diabetes; **NOD:** non-obese diabetic; **TLR:** Toll-like receptor; **PSA:** polysaccharide A, **BF:** *Bacteroides fragilis*; **ΔPSA:** PSA deficient; **HK:** heat-killed; **MLN:** mesenteric lymph node; **PnLN:** pancreatic lymph node; **SiLP:** small intestinal lamina propria; **LiLP:** large intestinal lamina propria; **PP:** Peyer's patch; **DSS:** Dextran sulfate sodium

References

1.   Charbonneau, M.R. *et al.* A microbial perspective of human developmental biology. *Nature* **535**, 48-55 (2016).
2.   Chervonsky, A.V. Microbiota and autoimmunity. *Cold Spring Harb Perspect Biol* **5**, a007294 (2013).
3.   Arpaia, N. *et al.* Metabolites produced by commensal bacteria promote peripheral regulatory T-cell generation. *Nature* **504**, 451-5 (2013).
4.   Chiba, T. & Seno, H. Indigenous clostridium species regulate systemic immune responses by induction of colonic regulatory T cells. *Gastroenterology* **141**, 1114-6 (2011).
5.   Wen, L. *et al.* Innate immunity and intestinal microbiota in the development of Type 1 diabetes. *Nature* **455**, 1109-13 (2008).
6.   Vatanen, T. *et al.* Variation in Microbiome LPS Immunogenicity Contributes to Autoimmunity in Humans. *Cell* **165**, 842-53 (2016).
7.   Vatanen, T. *et al.* The human gut microbiome in early-onset type 1 diabetes from the TEDDY study. *Nature* **562**, 589-594 (2018).
8.   Burrows, M.P., Volchkov, P., Kobayashi, K.S. & Chervonsky, A.V. Microbiota regulates type 1 diabetes through Toll-like receptors. *Proc Natl Acad Sci U S A* **112**, 9973-7 (2015).
9.   Vaarala, O., Atkinson, M.A. & Neu, J. The "perfect storm" for type 1 diabetes: the complex interplay between intestinal microbiota, gut permeability, and mucosal immunity. *Diabetes* **57**, 2555-62 (2008).
10.   Vaarala, O. Gut microbiota and type 1 diabetes. *Rev Diabet Stud* **9**, 251-9 (2012).
11.   Maffeis, C. *et al.* Association between intestinal permeability and faecal microbiota composition in Italian children with beta cell autoimmunity at risk for type 1 diabetes. *Diabetes Metab Res Rev* **32**, 700-709 (2016).
12.   Manfredo Vieira, S. *et al.* Translocation of a gut pathobiont drives autoimmunity in mice and humans. *Science* **359**, 1156-1161 (2018).
13.   Bosi, E. *et al.* Increased intestinal permeability precedes clinical onset of type 1 diabetes. *Diabetologia* **49**, 2824-7 (2006).
14.   Costa, F.R. *et al.* Gut microbiota translocation to the pancreatic lymph nodes triggers NOD2 activation and contributes to T1D onset. *J Exp Med* **213**, 1223-39 (2016).
15.   Ley, R.E. *et al.* Evolution of mammals and their gut microbes. *Science* **320**, 1647-51 (2008).
16.   Round, J.L. *et al.* The Toll-like receptor 2 pathway establishes colonization by a commensal of the human microbiota. *Science* **332**, 974-7 (2011).
17.   Chiu, C.C. *et al.* Monocolonization of germ-free mice with Bacteroides fragilis protects against dextran sulfate sodium-induced acute colitis. *Biomed Res Int* **2014**, 675786 (2014).
18.   Mazmanian, S.K., Liu, C.H., Tzianabos, A.O. & Kasper, D.L. An immunomodulatory molecule of symbiotic bacteria directs maturation of the host immune system. *Cell* **122**, 107-18 (2005).
19.   Troy, E.B. & Kasper, D.L. Beneficial effects of Bacteroides fragilis polysaccharides on the immune system. *Front Biosci (Landmark Ed)* **15**, 25-34 (2010).

20.   Ochoa-Reparaz, J. *et al.* A polysaccharide from the human commensal Bacteroides fragilis protects against CNS demyelinating disease. *Mucosal Immunol* **3**, 487-95 (2010).
21.   Moore, W.E. & Holdeman, L.V. Human fecal flora: the normal flora of 20 Japanese-Hawaiians. *Appl Microbiol* **27**, 961-79 (1974).
22.   Nichols, R.L., Schumer, W., Nythus, L.M., Bartlett, J.G. & Gorbach, S.L. Anaerobic Infections. *Am Fam Physician* **14**, 100-10 (1976).
23.   Bartlett, J.G., Onderdonk, A.B., Louie, T., Kasper, D.L. & Gorbach, S.L. A review. Lessons from an animal model of intra-abdominal sepsis. *Arch Surg* **113**, 853-7 (1978).
24.   Onderdonk, A.B., Kasper, D.L., Cisneros, R.L. & Bartlett, J.G. The capsular polysaccharide of Bacteroides fragilis as a virulence factor: comparison of the pathogenic potential of encapsulated and unencapsulated strains. *J Infect Dis* **136**, 82-9 (1977).
25.   Coyne, M.J. *et al.* Polysaccharide biosynthesis locus required for virulence of Bacteroides fragilis. *Infect Immun* **69**, 4342-50 (2001).
26.   Liu, C.H., Lee, S.M., Vanlare, J.M., Kasper, D.L. & Mazmanian, S.K. Regulation of surface architecture by symbiotic bacteria mediates host colonization. *Proc Natl Acad Sci U S A* **105**, 3951-6 (2008).
27.   Kayama, H. & Takeda, K. Polysaccharide A of Bacteroides fragilis: actions on dendritic cells and T cells. *Mol Cell* **54**, 206-7 (2014).
28.   Dasgupta, S., Erturk-Hasdemir, D., Ochoa-Reparaz, J., Reinecker, H.C. & Kasper, D.L. Plasmacytoid dendritic cells mediate anti-inflammatory responses to a gut commensal molecule via both innate and adaptive mechanisms. *Cell Host Microbe* **15**, 413-23 (2014).
29.   Wang, Q. *et al.* A bacterial carbohydrate links innate and adaptive responses through Toll-like receptor 2. *J Exp Med* **203**, 2853-63 (2006).
30.   Mazmanian, S.K., Round, J.L. & Kasper, D.L. A microbial symbiosis factor prevents intestinal inflammatory disease. *Nature* **453**, 620-5 (2008).
31.   Cinek, O. *et al.* Imbalance of bacteriome profiles within the Finnish Diabetes Prediction and Prevention study: Parallel use of 16S profiling and virome sequencing in stool samples from children with islet autoimmunity and matched controls. *Pediatr Diabetes* (2016).
32.   Alkanani, A.K. *et al.* Alterations in Intestinal Microbiota Correlate With Susceptibility to Type 1 Diabetes. *Diabetes* **64**, 3510-20 (2015).
33.   Davis-Richardson, A.G. *et al.* Bacteroides dorei dominates gut microbiome prior to autoimmunity in Finnish children at high risk for type 1 diabetes. *Front Microbiol* **5**, 678 (2014).
34.   Mejia-Leon, M.E., Petrosino, J.F., Ajami, N.J., Dominguez-Bello, M.G. & de la Barca, A.M. Fecal microbiota imbalance in Mexican children with type 1 diabetes. *Sci Rep* **4**, 3814 (2014).
35.   de Goffau, M.C. *et al.* Fecal microbiota composition differs between children with beta-cell autoimmunity and those without. *Diabetes* **62**, 1238-44 (2013).
36.   Sofi, M.H. *et al.* pH of drinking water influences the composition of gut microbiome and type 1 diabetes incidence. *Diabetes* **63**, 632-44 (2014).
37.   de Goffau, M.C. *et al.* Aberrant gut microbiota composition at the onset of type 1 diabetes in young children. *Diabetologia* **57**, 1569-77 (2014).
38.   Horn, J., Bender, B.S. & Bartlett, J.G. Role of anaerobic bacteria in perimandibular space infections. *Ann Otol Rhinol Laryngol Suppl* **154**, 34-9 (1991).

39. Polk, B.F. & Kasper, D.L. Bacteroides fragilis subspecies in clinical isolates. *Ann Intern Med* **86**, 569-71 (1977).

40. Surana, N.K. & Kasper, D.L. The yin yang of bacterial polysaccharides: lessons learned from B. fragilis PSA. *Immunol Rev* **245**, 13-26 (2012).

41. Erturk-Hasdemir, D. & Kasper, D.L. Finding a needle in a haystack: Bacteroides fragilis polysaccharide A as the archetypical symbiosis factor. *Ann N Y Acad Sci* **1417**, 116-129 (2018).

42. Kim, H.S. *et al.* Toll-like receptor 2 senses beta-cell death and contributes to the initiation of autoimmune diabetes. *Immunity* **27**, 321-33 (2007).

43. Fasano, A. Leaky gut and autoimmune diseases. *Clin Rev Allergy Immunol* **42**, 71-8 (2012).

44. Meddings, J.B., Jarand, J., Urbanski, S.J., Hardin, J. & Gall, D.G. Increased gastrointestinal permeability is an early lesion in the spontaneously diabetic BB rat. *Am J Physiol* **276**, G951-7 (1999).

45. Telesford, K.M. *et al.* A commensal symbiotic factor derived from Bacteroides fragilis promotes human CD39(+)Foxp3(+) T cells and Treg function. *Gut Microbes* **6**, 234-42 (2015).

46. Mazmanian, S.K. Capsular polysaccharides of symbiotic bacteria modulate immune responses during experimental colitis. *J Pediatr Gastroenterol Nutr* **46 Suppl 1**, E11-2 (2008).

47. Kelly, J.R. *et al.* Breaking down the barriers: the gut microbiome, intestinal permeability and stress-related psychiatric disorders. *Front Cell Neurosci* **9**, 392 (2015).

48. Eutamene, H. *et al.* Synergy between Lactobacillus paracasei and its bacterial products to counteract stress-induced gut permeability and sensitivity increase in rats. *J Nutr* **137**, 1901-7 (2007).

49. Cani, P.D. *et al.* Changes in gut microbiota control metabolic endotoxemia-induced inflammation in high-fat diet-induced obesity and diabetes in mice. *Diabetes* **57**, 1470-81 (2008).

50. Deitch, E.A., Berg, R. & Specian, R. Endotoxin promotes the translocation of bacteria from the gut. *Arch Surg* **122**, 185-90 (1987).

51. Lee, A.S. *et al.* Gut barrier disruption by an enteric bacterial pathogen accelerates insulitis in NOD mice. *Diabetologia* **53**, 741-8 (2010).

52. Serrander, R., Magnusson, K.E., Kihlstrom, E. & Sundqvist, T. Acute yersinia infections in man increase intestinal permeability for low-molecular weight polyethylene glycols (PEG 400). *Scand J Infect Dis* **18**, 409-13 (1986).

53. Yurkovetskiy, L. *et al.* Gender bias in autoimmunity is influenced by microbiota. *Immunity* **39**, 400-12 (2013).

54. Dunne, J.L. *et al.* The intestinal microbiome in type 1 diabetes. *Clin Exp Immunol* **177**, 30-7 (2014).

55. Peng, J. *et al.* Long term effect of gut microbiota transfer on diabetes development. *J Autoimmun* **53**, 85-94 (2014).

56. Kostic, A.D. *et al.* The dynamics of the human infant gut microbiome in development and in progression toward type 1 diabetes. *Cell Host Microbe* **17**, 260-73 (2015).

57. Livanos, A.E. *et al.* Antibiotic-mediated gut microbiome perturbation accelerates development of type 1 diabetes in mice. *Nat Microbiol* **1**, 16140 (2016).

58. Candon, S. *et al.* Antibiotics in early life alter the gut microbiome and increase disease incidence in a spontaneous mouse model of autoimmune insulin-dependent diabetes. *PLoS One* **10**, e0125448 (2015).

59. Brown, K. *et al.* Prolonged antibiotic treatment induces a diabetogenic intestinal microbiome that accelerates diabetes in NOD mice. *ISME J* **10**, 321-32 (2016).

60. Hu, Y. *et al.* Maternal Antibiotic Treatment Protects Offspring from Diabetes Development in Nonobese Diabetic Mice by Generation of Tolerogenic APCs. *J Immunol* **195**, 4176-84 (2015).

61. Mejia-Leon, M.E., Lopez-Dominguez, L., Aguayo-Patron, S.V., Caire-Juvera, G. & Calderon de la Barca, A.M. Dietary Changes and Gut Dysbiosis in Children With Type 1 Diabetes. *J Am Coll Nutr* **37**, 501-507 (2018).

62. Henschel, A.M. *et al.* Modulation of the diet and gastrointestinal microbiota normalizes systemic inflammation and beta-cell chemokine expression associated with autoimmune diabetes susceptibility. *PLoS One* **13**, e0190351 (2018).

63. Endesfelder, D. *et al.* Towards a functional hypothesis relating anti-islet cell autoimmunity to the dietary impact on microbial communities and butyrate production. *Microbiome* **4**, 17 (2016).

64. Davis-Richardson, A.G. & Triplett, E.W. A model for the role of gut bacteria in the development of autoimmunity for type 1 diabetes. *Diabetologia* **58**, 1386-93 (2015).

65. Murri, M. *et al.* Gut microbiota in children with type 1 diabetes differs from that in healthy children: a case-control study. *BMC Med* **11**, 46 (2013).

66. Neu, J. *et al.* Changes in intestinal morphology and permeability in the biobreeding rat before the onset of type 1 diabetes. *J Pediatr Gastroenterol Nutr* **40**, 589-95 (2005).

67. Kuitunen, M., Saukkonen, T., Ilonen, J., Akerblom, H.K. & Savilahti, E. Intestinal permeability to mannitol and lactulose in children with type 1 diabetes with the HLA-DQB1*02 allele. *Autoimmunity* **35**, 365-8 (2002).

68. Pabst, O. & Mowat, A.M. Oral tolerance to food protein. *Mucosal Immunol* **5**, 232-9 (2012).

69. Castro-Sanchez, P. & Martin-Villa, J.M. Gut immune system and oral tolerance. *Br J Nutr* **109 Suppl 2**, S3-11 (2013).

70. Ramiya, V.K., Shang, X.Z., Wasserfall, C.H. & Maclaren, N.K. Effect of oral and intravenous insulin and glutamic acid decarboxylase in NOD mice. *Autoimmunity* **26**, 139-51 (1997).

71. Chaillous, L. *et al.* Oral insulin administration and residual beta-cell function in recent-onset type 1 diabetes: a multicentre randomised controlled trial. Diabete Insuline Orale group. *Lancet* **356**, 545-9 (2000).

72. Bonifacio, E. *et al.* Effects of high-dose oral insulin on immune responses in children at high risk for type 1 diabetes: the Pre-POINT randomized clinical trial. *JAMA* **313**, 1541-9 (2015).

73. Caramalho, I. *et al.* Regulatory T cells contribute to diabetes protection in lipopolysaccharide-treated non-obese diabetic mice. *Scand J Immunol* **74**, 585-95 (2011).

74. Aumeunier, A. *et al.* Systemic Toll-like receptor stimulation suppresses experimental allergic asthma and autoimmune diabetes in NOD mice. *PLoS One* **5**, e11484 (2010).

75. Wang, J. *et al.* Multiple mechanisms involved in diabetes protection by lipopolysaccharide in non-obese diabetic mice. *Toxicol Appl Pharmacol* **285**, 149-58 (2015).

76. Bloem, K. *et al.* Interaction of the Capsular Polysaccharide A from Bacteroides fragilis with DC-SIGN on Human Dendritic Cells is Necessary for Its Processing and Presentation to T Cells. *Front Immunol* **4**, 103 (2013).

77. Park, B.S. & Lee, J.O. Recognition of lipopolysaccharide pattern by TLR4 complexes. *Exp Mol Med* **45**, e66 (2013).

78. Tezuka, H. & Ohteki, T. Regulation of intestinal homeostasis by dendritic cells. *Immunol Rev* **234**, 247-58 (2010).

79. Gulden, E. *et al.* Toll-like receptor 4 deficiency accelerates the development of insulin-deficient diabetes in non-obese diabetic mice. *PLoS One* **8**, e75385 (2013).

80. Westerholm-Ormio, M., Vaarala, O., Pihkala, P., Ilonen, J. & Savilahti, E. Immunologic activity in the small intestinal mucosa of pediatric patients with type 1 diabetes. *Diabetes* **52**, 2287-95 (2003).

81. Vaarala, O. The gut as a regulator of early inflammation in type 1 diabetes. *Curr Opin Endocrinol Diabetes Obes* **18**, 241-7 (2011).

82. Visser, J., Rozing, J., Sapone, A., Lammers, K. & Fasano, A. Tight junctions, intestinal permeability, and autoimmunity: celiac disease and type 1 diabetes paradigms. *Ann N Y Acad Sci* **1165**, 195-205 (2009).

83. Rodriguez-Calvo, T., Sabouri, S., Anquetil, F. & von Herrath, M.G. The viral paradigm in type 1 diabetes: Who are the main suspects? *Autoimmun Rev* **15**, 964-9 (2016).

84. Kerr, J.R. The role of parvovirus B19 in the pathogenesis of autoimmunity and autoimmune disease. *J Clin Pathol* **69**, 279-91 (2016).

85. Drescher, K.M., von Herrath, M. & Tracy, S. Enteroviruses, hygiene and type 1 diabetes: toward a preventive vaccine. *Rev Med Virol* **25**, 19-32 (2015).

86. Chobot, A. *et al.* ATP4A autoimmunity and Helicobacter pylori infection in children with type 1 diabetes. *Clin Exp Immunol* **177**, 598-602 (2014).

87. Christen, U. & von Herrath, M.G. Induction, acceleration or prevention of autoimmunity by molecular mimicry. *Mol Immunol* **40**, 1113-20 (2004).

88. Le Huerou-Luron, I., Blat, S. & Boudry, G. Breast- v. formula-feeding: impacts on the digestive tract and immediate and long-term health effects. *Nutr Res Rev* **23**, 23-36 (2010).

89. O'Sullivan, A., Farver, M. & Smilowitz, J.T. The Influence of Early Infant-Feeding Practices on the Intestinal Microbiome and Body Composition in Infants. *Nutr Metab Insights* **8**, 1-9 (2015).

90. Palanch, A.C. & Alvares, E.P. Feeding manipulation elicits different proliferative responses in the gastrointestinal tract of suckling and weanling rats. *Braz J Med Biol Res* **31**, 565-72 (1998).

91. Bell, R.C., Hoedl, A. & Turchinsky, J. Fructose feeding in the suckling-weaning transition in rats: effects on hyperlipidemia in adulthood. *Arch Physiol Biochem* **111**, 17-22 (2003).

92. Korn, T. *et al.* Myelin-specific regulatory T cells accumulate in the CNS but fail to control autoimmune inflammation. *Nat Med* **13**, 423-31 (2007).

93. Comstock, L.E. *et al.* Analysis of a capsular polysaccharide biosynthesis locus of Bacteroides fragilis. *Infect Immun* **67**, 3525-32 (1999).

94. Perez, N. *et al.* Preferential costimulation by CD80 results in IL-10-dependent TGF-beta1(+) -adaptive regulatory T cell generation. *J Immunol* **180**, 6566-76 (2008).

95. Goodyear, A.W., Kumar, A., Dow, S. & Ryan, E.P. Optimization of murine small intestine leukocyte isolation for global immune phenotype analysis. *J Immunol Methods* **405**, 97-108 (2014).

96. Karumuthil-Melethil, S. *et al.* TLR2- and Dectin 1-associated innate immune response modulates T-cell response to pancreatic beta-cell antigen and prevents type 1 diabetes. *Diabetes* **64**, 1341-57 (2015).

97. Karumuthil-Melethil, S., Gudi, R., Johnson, B.M., Perez, N. & Vasu, C. Fungal beta-glucan, a Dectin-1 ligand, promotes protection from type 1 diabetes by inducing regulatory innate immune response. *J Immunol* **193**, 3308-21 (2014).

98. Karumuthil-Melethil, S., Perez, N., Li, R. & Vasu, C. Induction of innate immune response through TLR2 and dectin 1 prevents type 1 diabetes. *J Immunol* **181**, 8323-34 (2008).

Legends for Figures

FIGURE 1: Oral administration of BF results in suppression of autoimmunity and protection of NOD mice from T1D. Ten-week-old prediabetic female NOD mice were given HK BF for 15 consecutive days (500 µg/mouse/day; dry-weight equivalent) **A)** Cohorts of control (n=17) and BF treated (n=14) mice were examined for blood glucose levels every week and the mice with glucose levels of >250mg/dl for two consecutive weeks were considered diabetic. Percentage of mice that developed diabetic at weeks post-treatment initiation are shown. Statistical significance was assessed by log-rank test for up to 15 and 30 weeks of monitoring period. **B)** Cohorts of mice (n=5/group) were euthanized 30 days post-treatment, pancreatic tissue sections were subjected to H&E staining and examined for insulitis. Examples of islets with different insulitis grades (left panel) and the percentage islets with different insulitis grades (right panel) are shown. Grades: 0 = no evidence of infiltration, 1 = peri-islet infiltration (<5%), 2= 5-25% islet infiltration, 3 = 25–50% islet infiltration, and 4 = >50% islet infiltration. At least 100 islets from intermittent sections were examined for each group. Statistical significance was assessed by Fisher's exact test comparing relative numbers of islets with insulitis grades ≤2 and ≥3 between groups. **C)** PnLN cells of mice euthanized 15 days post-treatment (n=6 mice/group) were cultured in the presence of β-cell Ag peptide cocktail for 48 h and the spent media was examined for self-antigen induced cytokine release by Luminex multiplex assay. Cytokine concentrations and relative ratios were compared and the statistical significance values were calculated by unpaired *t*-test. **D)** Fresh spleen cells from mice euthanized at 15 days post-treatment (n=5 mice/group) were activated ex vivo with PMA and ionomycin for 4 h and stained for intracellular cytokine levels, or examined for Foxp3 levels without ex vivo activation. Percentage and relative ratios of cells that are positive for specific markers are shown and the statistical significance was assessed by unpaired *t*-test. **E)** Enriched splenic T cells from control and BF treated mice were adoptively transferred into 6 week old NOD-Rag1 KO mice (1x10^6 cells/mouse) and the mice were monitored (n=6 mice/group) for hyperglycemia as described above. Log-rank test was employed for determining statistical significance.

FIGURE 2: Systemic (i.v.) administration of BF results in aggravated autoimmune response and rapid T1D onset in NOD mice. Ten-week-old pre-diabetic female NOD mice

were given heat-killed BF every 3rd day during 15 day period (10 µg/mouse/injection; dry-weight equivalent) by i.v. injection and studied as detailed for Fig. 1. **A)** Cohorts of control (n=17) and BF treated (n=14) mice were monitored for hyperglycemia as described for Fig. 1. Statistical significance was calculated by log-rank test for up to 15 and 30 weeks of monitoring periods. **B)** Cohorts of mice (n= 5/group) were euthanized 30 days post-treatment and pancreatic tissue sections were examined for insulitis, and the statistical significance was calculated by Fisher's exact test. **C)** PnLN cells of mice (n=6/group) were examined for self-antigen induced cytokine levels by Luminex multiplex assay and the statistical significance values were calculated by unpaired *t*-test. **D)** Fresh spleen cells from BF treated mice (n=5/group) and controls, 15 days post-treatment were examined for intracellular cytokines and Foxp3 levels, and the statistical significance was calculated by unpaired *t*-test. **E)** Enriched splenic T cells from control and BF treated mice were adoptively transferred into 6 week old NOD-Rag1 KO mice (n=6/group) and monitored for hyperglycemia.

FIGURE 3: Orally and systemically administered BF induces distinct immune responses. Eight week old female NOD mice were given HK BF for 3 consecutive days by oral gavage (500 µg/mouse/day) or by i.v. injection (10 µg/mouse/injection). **A)** On day 4, one set of mice were euthanized and cDNA prepared from distal ileum and distal colon of control and BF fed (oral) mice and spleen cells from control and BF injected (systemic) mice were subjected to qPCR assay and the expression levels of cytokines and non-cytokine factors were compared. Expression levels relative to β-actin expression were plotted as heatmaps with Morpheus application. n= 3 mice/group and the assay was performed in triplicate for each mouse. **B)** A set of NOD-Foxp3-GFP mice from a similar experiment (treated with BF for 3 days) were euthanized on day 7. MLN cells of BF fed and spleen cells from BF injected and control mice were examined for GFP+CD4+ cells, or ex vivo stimulated with PMA and ionomycin for 4 h and examined for intracellular cytokines, by FACS. Representative FACS plots (left panels) and plots showing percentage and relative ratios of cells that are positive for specific markers (right panels) are shown. n= 5 mice/group and the assay was performed in duplicate for each mouse. The statistical significance was assessed by unpaired t-test for all panels. **Note:** Fig. S2 shows Foxp3+, IFNγ+ and IL10+ positive CD4+ T cell frequencies of PP, SiLP and LiLP of mice that received BF orally and spleen of mice that received i.v. injection of BF for an extended period.

FIGURE 4: BF exposed intestinal and splenic APCs produce opposing effects on Tregs upon antigen presentation ex vivo and in vivo. Eight week old female NOD mice were given HK BF for 3 consecutive days by oral gavage or by i.v. injection as described for Fig. 3. **A)** One set of mice were euthanized within 24 h of treatment and CD11c+ DCs were enriched from the ileum of control and BF fed (oral) mice and spleen cells of control and BF injected (systemic) mice and subjected to qPCR assay to detect the expression levels of cytokines. n= 3 mice/group and the assay was performed in duplicate for cells from each mouse. The statistical significance was assessed by unpaired t-test. **B)** DCs were enriched from cohorts of mice similar to that of panel A, pulsed with BDC2.5 peptide and cultured with T cells enriched from BDC2.5-Foxp3-GFP mice for 4 days and examined for GFP+CD4+ cells by FACS. n= 3 independent experiments using DCs enriched and pooled from 2 mice/group in each experiment. Each assay was performed in triplicate, and the result of a representative assay is shown. The statistical significance value was determined by paired t-test which compared control and BF groups within each assay. **Note:** Fig. S3 shows cytokine levels in the culture supernatants from these assays, tested by ELISA. DCs isolated from large intestine showed similar properties in terms of cytokine expression and T cell activation (not shown). **C)** BDC2.5-Foxp3-GFP mice that were treated for 3 days as described above received BDC2.5 peptide by oral gavage (20 µg/mouse) or by i.v. injection (2 µg/mouse) on days 2 and 3. These mice were euthanized on day 7, GFP+CD4+ T cell frequencies in MLN, PP and SiLP of orally treated and spleen cells from i.v. treated mice were determined by FACS. n= 5 mice/group and the assay was performed in duplicate for each mouse. The statistical significance was assessed by unpaired t-test.

FIGURE 5: Oral or systemic administration of ΔPSA-BF has no major impact on autoimmune progression and T1D incidence in NOD mice. Ten-week-old female NOD mice were given heat-killed BF or ΔPSA-BF by oral gavage or by i.v. injection and studied as described for Figs. 1&2. **A)** Cohorts of (4 mice/group) were euthanized 30 days post-treatment, pancreatic tissue sections were stained using anti-insulin Ab (red) and the nuclear stain DAPI (blue) and examined for insulitis and insulin positive islets. Insulitis was graded 0-5 based on islet structure and immune cell infiltration (based on DAPI) and insulin staining (+ or -). 0+ (no infiltration/insulin+), 1+ (<5% infiltration/insulin+), 2+ (5-25% infiltration/insulin+), 3+ (25-50% infiltration/insulin+), 4+ (50-100% infiltration/insulin+), 4- (50-100% infiltration/insulin-), and 5- (only islet remnants left/insulin-). At least 100 islets from multiple intermittent sections

were examined for each group. Statistical significance was calculated by Fisher's exact test comparing relative numbers of islets with insulitis grades ≤2+ and ≥3+ between control vs BF and BF vs ΔPSA-BF groups. **B)** PnLN cells of mice (n=5/group) euthanized 15 days post-treatment were examined for Foxp3+CD4+ T cell frequencies by FACS, and the statistical significance was calculated by unpaired *t*-test. **C)** PnLN cells (n=5 mice/group) were also cultured overnight in the presence of soluble anti-CD3 antibody for 24 h and the supernatants were tested for cytokine levels by Luminex multiplex assay. Cytokine levels and relative ratios are shown and the statistical significance was calculated by unpaired *t*-test. **D)** Cohorts of control (n=17) and BF treated (n=14) mice were monitored for hyperglycemia as described for Figs. 1 and 2. Statistical significance was calculated by log-rank test. **Note:** This set of experiments was carried out in parallel to the experiments of Figs. 1A and 2A and the BF group was not included in Fig 5D to avoid repeated presentation of the same data. Fig. S5 shows the cytokine mRNA expression profile of intestinal tissues from mice that were fed or injected with BF and ΔPSA-BF for 3 days.

FIGURE 6: Oral administration of BF under enhanced gut permeability results in accelerated autoimmune progression and early onset of hyperglycemia in NOD mice. Eight week old female NOD mice were maintained on regular drinking water or drinking water containing DSS (0.5%W/V) for 5 days and switched to regular water. A) Cohorts of these mice (n=10/group) were left untreated, or given daily oral gavage of HK BF (500 μg/day/mouse) for 15 days (during and post-DSS treatment), and monitored for hyperglycemia as described for Figs. 1&2. Statistical significance was calculated by log-rank test. **B)** Cohorts of mice (n= 4/group) were euthanized 30 days post-treatment, H&E stained pancreatic tissue sections were examined for insulitis as described for Fig. 1, and the statistical significance was calculated by Fisher's exact test. **C)** PnLN cells of mice (n=4/group) euthanized 30 days post-treatment initiation were examined for Foxp3+CD4+ T cell frequencies by FACS, and the statistical significance was assessed by unpaired *t*-test. **D)** PnLN cells (n=4 mice/group) were also cultured overnight in the presence of β-cell Ag peptide cocktail for 48 h and the supernatants were tested for cytokine levels by Luminex multiplex assay. The statistical significance was calculated by unpaired *t*-test. **Notes:** Fig. S6 shows the degree of gut permeability in low-dose DSS treated and control mice. High dose (2.5% W/V), but not low dose (0.5%), DSS treated NOD mice showed loss of body weight, albeit modest compared to B6 mice (not shown).

Fig. 1

Fig. 2

Fig. 3

A

Distal ileum

Control | BF

	1	2	3	1	2	3	
Il1b							
Il6							p=0.002
Il10							p=0.004
Il12p70							
Tnfa							
Il21							p=0.001
Il22							
Il23							p=0.005
Tgfb1							
Mcp1							
Gmcsf							
Rantes							
Mip1a							
Mip1b							
Cxcl12							
Cxcl1							
Aldh1a2							p=0.032

Distal colon

Control | BF

p=0.002
p=0.017

p=0.027
p=0.014

p=0.005

p=0.033

Spleen

Control | BF

p=0.013
p=0.002
P<0.001
p=0.001
p=0.013
p=0.021
p=0.008
p=0.017
p=0.002

p=0.034
p=0.003

Row Minimum — Row Maximum

B

Oral (MLN)

Control | BF

GFP 5.58% | 7.56% — % GFP+ cells *p*=0.004

IL10 1.15% | 3.01% — % IL10+ cells *p*=0.003

IFNγ 4.72% | 4.60% — % IFNγ+ cells

IL17 0.73% | 1.37% — % IL17+ cells *p*=0.005

CD4

i.v. (Spleen)

Control | BF

GFP 13.7% | 10.2% — % GFP+ cells *p*=0.003

IL10 1.24% | 1.75% — % IL10+ cells *p*=0.026

IFNγ 5.78% | 9.07% — % IFNγ+ cells *p*=0.002

IL17 1.67% | 2.43% — % IL17+ cells *p*=0.020

CD4

Control | BF

Fig. 4

Fig. 5

Fig. 6