Netrin/UNC-6 triggers actin assembly and non-muscle myosin activity to drive dendrite

retraction in the self-avoidance response.

Lakshmi Sundararajan,[1] Cody J. Smith,[1,2] Joseph D. Watson[3,4], Bryan A. Millis[1,5,6], Matthew J.

Tyska,[1] David M. Miller, III[1,4,7]

[1]Department of Cell and Developmental Biology, Vanderbilt University, Nashville, TN

[2] Current address: Department of Biological Sciences, Notre Dame University, South Bend, IN

[3]Current address: Rho, Chapel Hill, NC

[4]Neuroscience Graduate Program, Vanderbilt University, Nashville, TN

[5] Cell Imaging Shared Resource, Vanderbilt University, Nashville, TN

10 [6] Vanderbilt Biophotonics Center, Vanderbilt University, Nashville, TN

11 [7]Corresponding author

SUMMARY

Dendrite growth is constrained by the self-avoidance response but the downstream pathways that

balance these opposing mechanisms are unknown. We have proposed that the diffusible cue UNC-

6(Netrin) is captured by UNC-40 (DCC) for a short-range interaction with UNC-5 to trigger self-

avoidance in the *C. elegans* PVD neuron. Here we report that the actin-polymerizing proteins

UNC-34(Ena/VASP), WSP-1(WASP), UNC-73(Trio), MIG-10(Lamellipodin) and the Arp2/3

complex effect dendrite retraction in the self-avoidance response mediated by UNC-6(Netrin). The

paradoxical idea that actin polymerization results in shorter rather than longer dendrites is

explained by our finding that NMY-1 (non-muscle myosin II) is necessary for retraction and could

therefore mediate this effect in a contractile mechanism. Our results also show that dendrite length

is determined by the antagonistic effects on the actin cytoskeleton of separate sets of effectors for

retraction mediated by UNC-6(Netrin) versus outgrowth promoted by the DMA-1 receptor. Thus,

our findings suggest that the dendrite length depends on an intrinsic mechanism that balances

distinct modes of actin assembly for growth versus retraction.

INTRODUCTION

Dendritic arbors are defined by the balanced effects of outgrowth which expands the structure versus retraction which constrains the size of the receptive field. Microtubules and filamentous actin (F-actin) are prominent dendritic components and have been implicated as key drivers of dendritic growth and maintenance by the finding that treatments that perturb cytoskeletal dynamics may also disrupt dendritic structure [1-3]. For many neurons, dendritic growth is highly exuberant with multiple tiers of branches projecting outward from the cell soma. Ultimately, growth may be terminated by external cues. For example, neurons with similar functions are typically limited to separate domains by tiling mechanisms in which mutual contact induces dendrite retraction. The related phenomenon of self-avoidance is widely employed to prevent overlaps among sister dendrites arising from a single neuron [4]. Homotypic interactions between the membrane components Dscam and protocaderins can mediate the self-avoidance response [5-7]. Surprisingly, in some instances, soluble axon guidance cues and their canonical receptors are also required. For example, self-avoidance for the highly-branched Purkinje neuron depends on both protocadherins and also repulsive interactions between sister dendrites decorated with the Robo receptor and its diffusible ligand, Slit [8]. In another example, we have shown that UNC-6/Netrin and its cognate receptors, UNC-5 and UNC-40/DCC, mediate self-avoidance for PVD nociceptive neurons in *C. elegans* [9]. Although multiple cell-surface interactions are now known to trigger self-avoidance, little is known of the downstream effectors that drive dendrite retraction in this mechanism.

The PVD neurons, one on each side of the body, build complex dendritic arbors through a series of successive 1°, 2°, 3° and 4° orthogonal branching events [10][11,12]. Self-avoidance ensures that adjacent 3° branches do not overgrow one another[9,11] (Figure 1). Our previous results

suggested a novel mechanism of self-avoidance in which UNC-40/DCC captures UNC-6/Netrin

at the PVD cell surface and then triggers retraction by interacting with UNC-5 on the neighboring

3° branch. UNC-40/DCC may also effect self-avoidance by acting in a separate pathway that does

not involve UNC-6/Netrin[9]. Here we describe a cell biological model of dendrite retraction in

the PVD self-avoidance response that depends on actin polymerization.

The structure of the actin cytoskeleton is controlled by a wide array of effector proteins that

regulate specific modes of actin polymerization. For example, Ena/VASP enhances F-actin

elongation at the plus-end [13] and the Arp2/3 complex functions with WASP (Wiskott-Aldrich

syndrome protein) and the Wave Regulatory Complex (WRC) to promote F-actin branching[14].

Upstream regulators of WASP and the WRC include Rho family GTPases[15] and their

activators, the GEFs (Guanine nucleotide Exchange Factors) UNC-73/Trio[16,17] and TIAM (T-

cell Lymphoma Invasion and Metastasis Factor) [18,19]. Members of the Lamellipodin/Lpd

family recruit Ena/VASP to localize F-actin assembly at the leading edge of migrating cells[20].

The fact that all of these components (UNC-34/Ena/VASP, Arp2/3 complex, WSP-1/WASP,

WRC, UNC-73/Trio, TIAM/TIAM-1, MIG-10/lamellipodin) have been previously shown to

function in *C. elegans* to mediate axon guidance underscores the key role of the actin

cytoskeleton in growth cone steering [16,19,21-24].

Our results show that Ena/VASP functions downstream of UNC-5 to mediate PVD self-avoidance.

Genetic evidence detecting roles for the Arp2/3 complex and its upstream regulator, WASP/WSP-

1, indicates that the formation of branched actin networks may contribute to dendrite retraction.

A necessary role for actin polymerization is also indicated by the defective PVD self-avoidance

response of mutants with disabled UNC-73/Trio or MIG-10/lamellipodin. We show that PVD self-

avoidance also requires NMY-1/non-muscle myosin II which we propose effects dendrite

retraction in a contractile mechanism that drives the reorganization of the nascent actin

cytoskeleton. The necessary role for myosin could explain how actin polymerization can result in

shorter rather than longer dendrites in the self-avoidance response. Thus, we propose that UNC-

6/Netrin triggers self-avoidance by simultaneously stimulating actin assembly and non-muscle

myosin activity in 3° dendrites.

Because dendrite growth depends on actin polymerization, our finding that the actin cytoskeleton

is also necessary for dendrite self-avoidance points to different modes of actin polymerization for

growth vs retraction. Recent work has shown that a multicomponent complex involving the PVD

membrane proteins DMA-1 and HPO-30 drives dendritic growth by linking the PVD actin

cytoskeleton to adhesive cues on the adjacent epidermis[25]. Dendrite elongation, in this case,

depends on the WRC and TIAM-1, both of which are known to promote actin polymerization. In

contrast, our work has revealed that effectors of actin polymerization that are required for 3^0 branch

self-avoidance (e.g., UNC-34/Ena/VASP, MIG-10/lamellipodin, UNC-73/Trio, Arp2/3 complex,

WSP-1/WASP) are not necessary for 3^0 dendrite outgrowth. Additionally, we show that UNC-

6/Netrin signaling antagonizes the DMA-1-dependent mechanism of dendritic growth. These

findings are significant because they suggest that overall dendrite length is defined by the relative

strengths of opposing pathways that utilize separate sets of effectors to differentially regulate actin

polymerization for either growth or retraction.

RESULTS

UNC-6/Netrin mediates PVD sister dendrite self-avoidance.

To visualize PVD morphology, we utilized a previously characterized GFP marker driven by a PVD-specific promoter (*PVD*::GFP) (Figure 1A-B)[11]. Each PVD neuron adopts a stereotypical morphology characterized by a series of orthogonal junctions between adjacent branches: 1° dendrites extend laterally from the PVD cell soma; each 2° dendrite projects along the dorsal-ventral axis to generate a T-shaped junction comprised of two 3° dendrites each with either an anterior or posterior trajectory; terminal 4° dendrites occupy interstitial locations between the epidermis and body muscles. The net result of this branching pattern is a series of tree-like structures or menorahs distributed along the 1° dendrite. In contrast to the highly branched dendritic architecture, a single axon projects from the PVD cell soma to join the ventral nerve cord[10-12]. Despite the complexity of this network, dendrites rarely overlap (~2%) in the mature PVD neuron (Figure 1A). Self-avoidance depends on an active process in which sister PVD dendrites retract upon physical contact with one another (Figure 1C). In the wild type, 3° branches arising from adjacent menorahs initially grow outward toward one another but then retract after mutual contact. The net result is a characteristic gap between the tips of 3° dendrites in neighboring menorahs[11]. We have previously shown that this outcome depends in part on the diffusible cue UNC-6/Netrin and its canonical receptors, UNC-40/DCC and UNC-5[9]. Mutations that disable *unc-6*, for example, result in a significant increase in the fraction of overlapping 3° dendrites (arrowheads, Figure 1B, Figure 2D). Similar self-avoidance defects were observed for mutants of either *unc-40* or *unc-5*. Our results are consistent with a model in which UNC-40 captures UNC-6 for contact with UNC-5 at the tips of adjacent 3° dendrites (Figure 1D)[9]. Here we address the

question of how the activated UNC-5 receptor then modifies the actin cytoskeleton to drive

dendrite retraction.

Constitutively active UNC-5 drives dendrite retraction:

We have proposed that UNC-6 triggers dendrite retraction by activating UNC-5[9]. This model

predicts that a constitutively active form of UNC-5 should result in a 'hyper'-retraction phenotype.

To test this idea, we attached a myristoylation sequence to the cytoplasmic domain of UNC-5[26].

The resultant MYR::UNC-5 protein is effectively tethered to the cytoplasmic membrane thus

inducing activation of the UNC-5 intracellular domain. Expression of MYR::UNC-5 in PVD

results in substantially shorter 3° dendrites in comparison to wild type at the late L3 stage (Figure

2A-C). This phenotype suggests that constitutive activation of UNC-5 prevents normal outgrowth

of 3° dendrites. We tested this idea by scoring the dynamic status of 3° dendrites. To quantify

movement, we measured the distance from a fiduciary point to the tip of a 3° dendrite at intervals

over time (delta t = 240 sec) (Figure 2E) (Movies S1-S2). Examples of these measurements are

shown in Figure 2E. We observed that wild-type 3° dendrites typically display saltatory movement

with periodic bouts of extension and retraction that result in net elongation. In contrast, 3° dendrites

of PVD neurons expressing MYR::UNC-5 grow more slowly and rarely show periods of active

extension and retraction (Figure 2E and 2F). The higher overall motility of 3^0 dendrites in wild-

type vs MYR::UNC-5 is evident from a comparison of the standard deviation of the net movement

of 3° dendrites from a fiduciary point in each strain (Figure 2F). Because MYR::UNC-5 lacks the

UNC-5 extracellular ligand-binding domain, it should function independently of *unc-6*. We

confirmed this prediction by demonstrating that MYR::UNC-5 rescues the self-avoidance defects

observed in an *unc-6* mutant background (Figure 2G). Together, these results are consistent with

the hypothesis that UNC-6/Netrin triggers the self-avoidance response by activating the

downstream UNC-5 receptor to drive dendrite retraction.

UNC-34/Ena/VASP mediates self-avoidance in actin-containing PVD dendrites.

To explore potential modifications of the PVD cytoskeleton as effectors of the UNC-6/Netrin-

dependent self-avoidance response, we used live cell markers to visualize actin (PVD::ACT-

1::GFP) and microtubules (*pdes-2*::TBA-1::mCherry) in PVD[27]. We observed a robust ACT-

1::GFP signal throughout the PVD neuron including the axon and all dendrites (Figure 3A). In

contrast, the TBA-1::mCherry signal was strongest in the PVD axon and 1° dendrite. This result

confirms an earlier report suggesting that dendritic microtubules are most abundant in the 1°

branch (Figure 3A) [27]. We thus considered the possibility that the self-avoidance response could

depend on the regulation of actin dynamics. To test this idea, we examined the PVD self-avoidance

phenotype for a mutation that disables the actin binding protein, Ena/VASP. Ena/VASP promotes

the elongation of actin filaments and the *C. elegans* homolog UNC-34[28] is enriched in a gene

expression profile of PVD neurons[11]. A strong loss-of-function allele of *unc-34*/Ena/VASP [29]

alters overall PVD morphology (e.g., truncated or misplaced 1^O dendrite, fewer 2^O branches)

(Figure 3B-C, Figure S1) but does not produce the severe lateral branching defects arising from

mutations in other regulators of actin polymerization (e.g, *tiam-1*) that promote dendrite

outgrowth[25]. Notably, PVD neurons in *unc-34* mutant animals showed robust self-avoidance

defects (Figure 3B-C). PVD-specific expression of mCherry-tagged UNC-34 protein rescued this

unc-34 mutant phenotype suggesting that *unc-34* function is cell-autonomous for the self-

avoidance response (Figure 3C). *unc-34* mutant 1^O and 2^O branch defects were not rescued by PVD

expression of UNC-34, however, which is indicative of UNC-34 function in other nearby cell types

that influence PVD morphogenesis (Figure 3B, Figure S1). A downstream role for *unc-34* in the

UNC-6/Netrin pathway is consistent with our finding that the Unc-34 self-avoidance defect is

epistatic to the hyper-retraction phenotype of MYR::UNC-5 (Figure 3C).

For visualizing UNC-34/Ena/VASP in developing dendrites, we labeled UNC-34 with mCherry

and co-expressed it in PVD (PVD::mCherry::UNC-34) with LifeAct::GFP to detect actin-

containing structures. mCherry::UNC-34 is functional because it rescues the *unc-34* self-

avoidance defect (Fig 3C). iSIM super resolution imaging detected punctate mCherry::UNC-34

throughout the PVD dendritic architecture. Notably, mCherry::UNC-34 puncta are frequently

detected at the tips of 3^O dendrites (Figure 4A-C), a finding consistent with the observed distal

location of Ena/VASP in F-actin containing filopodial structures [30,31]. We used time-lapse

imaging to detect potential trafficking of mCherry::UNC-34 (Movie S6). Kymographs detected

anterograde as well as retrograde movements of mCherry::UNC-34 in 3° dendrites near the tip

(green arrowheads) whereas stationary puncta were characteristically located in the shaft region

(red arrowheads) (Figure 4C-D). These results are consistent with the idea that UNC-

34/Ena/VASP is delivered to 3° dendrites where it mediates actin assembly for branch retraction

in the self-avoidance response.

Mutations that disable actin-polymerizing proteins disrupt dendrite self-avoidance.

Genetic tests of additional regulators of actin polymerization also revealed PVD self-avoidance

phenotypes. Lamellipodin (Lpd) interacts with ENA/VASP at the tips of filopodia and

lamellopodia to extend actin filaments. Mutants of *mig-10*/Lpd display robust PVD self-avoidance

defects (Figure 3B, E). Similarly, self-avoidance is disrupted by a mutation that eliminates the Rac

GEF activity of UNC-73/Trio, a potent regulator of actin dynamics[16]. We note that both *mig-*

10/Lpd and *unc-73*/Trio mutants also show other defects in PVD morphology (e.g., displaced 1^O

dendrite, fewer 2^O branches) (Figure S1) but do not impair outgrowth of higher order dendrites as

was also observed for *unc-34* (**Fig. 3**). To test for potential roles for branched actin polymerization,

we examined a null allele of *wsp-1/WASP* (Wiskott-Aldrich Syndrome protein) and observed

defective PVD self-avoidance. If *wsp-1*/WASP functions in a common pathway with *unc-6*, then

the self-avoidance defect of the double mutant, *wsp-1; unc-6*, should be no more severe than that

of either *wsp-1* or *unc-6* single mutants. We confirmed this prediction by scoring PVD self-

avoidance in *wsp-1; unc-6* mutant animals (Figure 3D). Because WASP is known to activate the

Arp2/3 complex[15,32], we also performed RNAi knock down of the conserved Arp2/3

component ARX-5/p21 and detected significant PVD self-avoidance defects (Figure 3E). The

canonical roles of WASP and the Arp2/3 complex suggest that self-avoidance depends on the

formation of branched F-actin networks. Recent studies have shown that actin is required for PVD

dendrite outgrowth[25]. Here our genetic results point to the paradoxical idea that elongation and

branching of F-actin filaments are also necessary for dendrite retraction and that this mechanism

is activated by UNC-6/Netrin in the self-avoidance response.

The actin cytoskeleton is highly dynamic in growing and retracting PVD dendrites.

To test the idea that the actin cytoskeleton is differentially modulated for both outgrowth and

retraction we expressed LifeAct::GFP in PVD neurons to monitor actin dynamics during dendrite

morphogenesis. Although LifeAct::GFP binds both filamentous actin (F-actin) and monomeric G-

actin with high affinity, bright LifeAct::GFP-labeled foci typically correspond to F-actin-

containing structures[33]. We observed that the LifeAct::GFP signal was consistently brightest

nearest the tips of growing dendrites in comparison to internal dendritic regions (Fig. 5B, C) as

also reported in other studies [25,34]. Time-lapse imaging of PVD::LifeAct::GFP by spinning disk

confocal microscopy during the late L3/early L4 stage detected striking, dynamic fluctuations in

the LifeAct::GFP signal (Movie S3). We quantified this effect by comparing time-dependent

changes in LifeAct::GFP fluorescence versus that of a PVD::mCherry cytosolic marker. The

relatively constant intensity of PVD::mCherry signal in these experiments favors the idea that the

observed changes in LifeAct::GFP fluorescence could be due to actin dynamics in growing

dendrites (Figure 5A, D-E).

Because genetic knockdown of actin polymerizing proteins (UNC-34/Ena/VASP, MIG-10/Lpd,

UNC-73/Trio, WSP-1/WASP, ARX-5/P21) (**Figure 3**) disrupts self-avoidance and thus suggests

that F-actin is required, we monitored LifeAct::GFP in 3° dendrites during the period (L3-L4

transition) in which UNC-6/Netrin mediates dendrite retraction[9]. Time-lapse imaging detected

dynamic fluctuations in LifeAct::GFP fluorescence in these distal regions (Figure 6) (Movie S3).

For these measurements, the LifeAct::GFP signal near the tips of 3° dendrites was quantified at 1-

1.25 minute intervals over a 10 minute period spanning at least one contact event. A cytoplasmic

mCherry marker was used to detect contact events as instances in which the measured gap between

left and right 3° dendrites approached zero. In some cases, a burst of LifeAct::GFP signal was

detected in at least one of the adjacent dendrites during the contact period suggesting a role for

actin polymerization in retraction (Figure 6 A, B) (Movies S4 and S5). In other instances, however,

we did not observe an increase in LifeAct::GFP fluorescence with contact (Figure 6 C, D) and no

correlation was detected in quantitative comparisons (n = 8) between the LifeAct::GFP signal

before *versus* after contact (**Figure 6F**). An alternative approach for quantifying LifeAct::GFP in

3^O dendrites at shorter time intervals (35 s) [34] also did not detect a consistent correlation between

LifeAct::GFP fluorescence intensity and dendrite retraction (Figure S2). The simplest explanation

of these results is that actin polymerization drives both outgrowth and retraction which are tightly

coupled over time and thus difficult to resolve as separate events.

NMY-1/Non-muscle myosin II drives dendrite retraction in the self-avoidance mechanism.

Our results indicate that F-actin is the dominant cytoskeletal structure in growing PVD dendrites

and that F-actin assembly is also likely required for dendrite retraction. In considering a

mechanism to account for dendrite shortening, we hypothesized that a newly synthesized actin

cytoskeleton could mediate retraction by providing a substrate for non-muscle myosin II to drive

contraction. To test this idea, we examined loss-of-function alleles that disable the *C. elegans* non-

muscle myosins, NMY-1 and NMY-2. This experiment determined that mutations in *nmy-1* but

not *nmy-2* (data not shown) show PVD self-avoidance defects (Figure 7A, D). This effect is cell

autonomous for *nmy-1* because PVD-specific RNAi of *nmy-1* also results in overlapping 3°

dendrites and PVD-specific expression of GFP-tagged NMY-1 is sufficient to rescue the *nmy-1*

self-avoidance defect (Figure 7 B-D). We conclude that a specific non-muscle myosin, NMY-1,

promotes dendrite retraction in the self-avoidance mechanism and could act by driving the

translocation of F-actin bundles at the tips of 3° dendrites.

Phosphomimetic activation of MLC-4 shortens PVD 3° dendrites.

Non-muscle myosin II motor proteins are composed of myosin heavy chain (MHC) and light

chains, the essential light chain (ELC) and regulatory light chain (RLC) [35,36]. Myosin motor

activity is triggered by RLC phosphorylation that induces a conformational change to allow

assembly of the myosin complex into active bipolar filaments (Figure 8A) [37]. Having shown

that NMY-1/non-muscle myosin is required for self-avoidance, we next performed an additional

experiment to ask if constitutive non-muscle myosin activity is sufficient to induce dendrite

retraction. For this test, we expressed a phosphomimetic mutant of the MLC-4, the *C. elegans* RLC

homolog. MLC-4/myosin regulatory light chain contains serine/S and threonine/T phosphorylation

sites in a highly-conserved domain. Both S and T residues were mutated to Aspartate/D and the

resultant phosphomimetic construct MLC-4DD was fused to GFP for transgenic expression in

PVD (Figure 8B-C) [38] PVD neurons that expressed MLC-4DD showed significantly shorter 3°

dendrites in comparison to wild type as well as a concurrent increase in the width of gaps between

3° dendrites from adjacent menorahs (Figure 8D-G). These phenotypic traits are consistent with

the proposed role of non-muscle myosin in dendrite retraction and closely resemble that of PVD

neurons that express MYR::UNC-5 which functions downstream of UNC-6/Netrin to trigger the

self-avoidance response. Our additional finding that MLC-4DD rescues the self-avoidance defect

of *unc-6* mutants suggests that NMY-1/non-muscle myosin II also functions downstream of *unc-*

6 for dendrite retraction (Figure 8H).

Antagonistic pathways regulate dendrite outgrowth vs self-avoidance.

Our time lapse imaging studies detected dynamic actin polymerization in growing PVD dendrites

(Movie S3). This observation and the recent finding that PVD dendritic architecture is significantly

perturbed by mutations in either a specific actin structural gene, *act-4*, or that disable the F-actin

nucleating WAVE complex or TIAM-1/GEF, suggest that dendritic outgrowth depends on actin

assembly[25]. Actin assembly is also likely required for dendrite retraction given our results

showing that genetic knockdown of a separate set of actin-binding proteins (i.e., UNC-

34/Ena/VASP, MIG-10/Lpd, UNC-73/Trio, WSP-1/WASP, ARX-5/p21) (**Fig. 3**), blocks self-

avoidance but not dendrite outgrowth [34]. A shared role for actin polymerization in both

outgrowth and retraction argues that the net length of PVD dendrites must depend on an intrinsic

mechanism that regulates actin assembly to balance these competing effects.

Our results suggest that UNC-6/Netrin signaling promotes dendrite retraction in a mechanism

involving F-actin assembly. PVD dendritic outgrowth is driven by a separate pathway in which

DMA-1, a membrane protein expressed in PVD, interacts with a multicomponent receptor

anchored in the adjacent epidermis[39-43]. Dendritic growth depends on binding of DMA-1 and

the claudin-like protein, HPO-30, to specific regulators of actin assembly (i.e., WAVE complex,

TIAM-1/GEF) in the PVD cytoplasm[25,44]. These interactions may be modulated by the Furin

KPC-1 which antagonizes DMA-1 surface expression in PVD [45-47]. *kpc-1* activity is proposed

to temporally weaken adhesion with the epidermis to facilitate changes in trajectory at specific

stages in dendritic outgrowth [41,47]. For example, the tips of 3° branches typically execute a

right-angle turn after the self-avoidance response to produce a 4° dendrite[11]. This morphological

transition largely fails in *kpc-1* mutants in which 3° dendrites elongate abnormally and show severe

self-avoidance defects due to over-expression of DMA-1 [41,47]. Thus, we considered the idea

that self-avoidance is achieved by the combined effects of *kpc-1*-dependent downregulation of

DMA-1 and a separate mechanism that antagonizes outgrowth and requires UNC-6/Netrin. We

performed a series of genetic experiments to test this model.

The hypomorphic loss-of-function allele, *kpc-1(xr58),* results in over-expression of DMA-1 and a

consequent, robust self-avoidance defect with ~42% of sister 3° branches overlapping one another

(Figure 9A, D)[46,47]. This defect is not observed, however, in *kpc-1*/+ heterozygotes (Figure

9D). Similarly, self-avoidance in *unc-6*/+ heterozygous animals is indistinguishable from wild

type (Figure 9D). Thus, a 2-fold reduction in gene dosage for either *kpc-1* or *unc-6* does not perturb

self-avoidance. We reasoned, however, that if expression of UNC-6/Netrin and KPC-1 were

simultaneously reduced by half, then self-avoidance should be at least partially disabled since both

unc-6 and *kpc-1* normally function to antagonize dendrite outgrowth. To test this prediction, we

constructed double heterozygous mutants of *kpc-1* and *unc-6* (e.g., *kpc-1*/+; *unc-6*/+) and

quantified the self-avoidance phenotype. This experiment revealed that ~30% of adjacent sister 3°

dendrites fail to self-avoid in *kpc-1*/+; *unc-6*/+ animals which is significantly greater than that of

either *kpc-1*/+ or *unc-6*/+ single mutants (Figure 9B, D). This genetic "enhancer" effect is

consistent with the idea that both KPC-1 and UNC-6 function to limit overgrowth of 3° dendrites.

If this effect is due to elevated levels of DMA-1 arising from reduced KPC-1 activity in the *kpc-*

1/+; *unc-6*/+ background, then a concomitant diminution of DMA-1 activity could restore normal

self-avoidance. We tested this idea by constructing the triple heterozygote, *kpc-1* +/+ *dma-1; unc-*

6/+, and detected robust suppression of the self-avoidance defect observed in *kpc-1*/+; *unc-6*/+

animals (Figure 9C, D). Thus, these results suggest that DMA-1-mediated adhesion to the

epidermis antagonizes UNC-6/Netrin-driven dendrite retraction and that the relative strengths of

these opposing pathways must be finely tuned to achieve the self-avoidance effect (Figure 9E).

The key role of actin polymerization in both UNC-6/Netrin-mediated retraction and DMA-1-

dependent outgrowth argues that this balancing mechanism likely regulates both modes of actin

assembly.

DISCUSSION

Dendrites arising from the same neuron typically do not overlap. This phenomenon of dendrite

self-avoidance is universally observed and derives from an active mechanism in which contact

between sister dendrites triggers mutual retraction[4] As might be predicted for an event that

depends on physical proximity, cell-surface associated proteins have been shown to mediate

dendrite self-avoidance[34,48]. These include Dscam and protocadherins which are expressed in

multiple alternative forms to distinguish self from non-self among contacting dendrites [5-7]. Self-

avoidance for certain neurons can also depend on the interaction of diffusible cues with their

cognate receptors. For example, we have shown that the axon guidance signal, UNC-6/Netrin, and

its receptors, UNC-5 and UNC-40/DCC, function together to mediate self-avoidance for PVD

sensory neurons in *C. elegans*. We have proposed that UNC-40/DCC captures UNC-6/Netrin on

the membrane surface to facilitate repulsion through physical interaction with UNC-5 on PVD

dendrites[9] (Figure 1D). Little is known, however of the downstream effectors of dendrite

retraction in the self-avoidance response. Here we report that UNC-6/Netrin promotes actin-

polymerization to drive dendrite retraction. Our finding that non-muscle myosin is also involved

resolves the paradoxical observation that UNC-6/Netrin dependent actin polymerization shortens

rather than lengthens PVD 3° dendrites. We propose that non-muscle myosin engages the F-actin

cytoskeleton at the tips of PVD dendrites to drive retraction.

Regulators of actin polymerization are required for dendrite self-avoidance.

The proposed role for actin polymerization in the self-avoidance response derives from our finding

that multiple regulators of F-actin assembly are necessary for efficient dendrite retraction. For

example, a mutation that disables UNC-34, the *C. elegans* homolog of Ena/VASP, disrupts PVD

self-avoidance. Ena/VASP promotes actin polymerization by preventing capping proteins from

blocking the addition of actin monomers to the plus-end of elongating actin filaments. In axon

growth cones, Ena/VASP localizes to the tips of actin bundles in growing filopodia[13,31]. We

observed punctate Ena/VASP localization in PVD dendrites and active trafficking from the cell

soma (Figure 4). Our results showing that an *unc-34* mutant blocks the hyper-retraction phenotype

of a constitutively active UNC-5 receptor (Figure 3) argues that *unc-34* functions downstream to

mediate UNC-6/Netrin-dependent self-avoidance. Additional genetic evidence indicates that

WSP-1/WASP also acts in a common pathway with UNC-6/Netrin (Figure 3). The established role

for WASP of activating the Arp2/3 complex [49] is consistent with our finding that RNAi

knockdown of *arx-5/p21*, a conserved Arp2/3 component, partially disables the self-avoidance

mechanism (Figure 3). Dual roles for UNC-34 and WSP-1 in actin polymerization, as suggested

here, are consistent with previous work indicating that Ena/VASP can activate Arp2/3 through

direct interaction with WASP proteins[28,50-52]. Thus, taken together, our results suggest that

Ena/VASP and WSP-1 coordinate the creation of branched actin networks at the tips of retracting

PVD dendrites in the UNC-6/Netrin dependent self-avoidance response (Figure 9E).

A role for Ena/VASP in dendrite retraction, as have we have proposed here, parallels earlier

findings of a necessary function for Ena/VASP in growth cone repulsion. For example, axon

guidance defects arising from ectopic expression of UNC-5 in *C. elegans* touch neurons requires

UNC-34/Ena/VASP[53], a finding confirmed by our observation that the dendritic shortening

phenotype induced by MYR::UNC-5 is prevented by a loss-of-function mutation in *unc-34*.

Ena/VASP is also required for axon repulsion mediated by Slit and its receptor Robo[21,54,55].

Although Ena/VASP function can be readily integrated into models of axonal attraction, the well-

established role of Ena/VASP in F-actin assembly has been more difficult to rationalize in

mechanisms of axonal repulsion which are presumed to involve actin depolymerization[13,56]. A

partial explanation for this paradox is provided by a recent study showing that Slit, acting through

Robo, induces transient filopodial outgrowth in migrating axonal growth cones and that this effect

requires Ena/VASP[55]. The functional necessity for filopodial growth in this mechanism is

unclear, but the central role of Ena/VASP-induced actin assembly in the overall repulsive response

to Slit mirrors our finding that dendrite retraction in PVD neurons requires UNC-34/Ena/VASP

and F-actin assembly.

Our studies also detected necessary roles in self-avoidance for MIG-10/LPD which has been

previously shown to function with UNC-34/Ena/VASP to promote F-actin polymerization and

UNC-73/Trio, a Rho/Rac GEF that acts in UNC-6/Netrin pathways to regulate the actin

cytoskeleton[16,20,21,26]. Altogether, our genetic analysis identified five known effectors of actin

polymerization (UNC-34/Ena/VASP, WSP-1/WASP, Arp2/3 complex, MIG-10/Lpd, UNC-

73/Trio) that drive retraction of 3^O dendrites in the self-avoidance response but which are not

required for 3^O branch extension during outgrowth.

Non-muscle myosin II drives dendrite retraction in the self-avoidance response.

Our genetic analysis detected a cell-autonomous role for NMY-1/non-muscle myosin II in dendrite

self-avoidance (Figure 7). Because our results revealed F-actin at the tips of PVD dendrites (Figure

5, 6), we suggest that NMY-1 mediates retraction by interacting with F-actin to drive a contractile

mechanism. For example, NMY-1 could accelerate retrograde flow, a non-muscle myosin-

dependent effect that also drives filopodial retraction in growth cones and at the leading edge of

migrating cells[57,58]. Alternatively, NMY-1 could shorten PVD dendrites by inducing the

reorganization of the nascent actin cytoskeleton into a more compact structure[59].

Phosphorylation of myosin regulatory light chain (RLC) activates non-muscle myosin[37] and we

showed that MLC-4DD, a phosphomimetic, and, thus, constitutively active form of the *C. elegans*

homolog of RLC, results in hyper-retraction of PVD 3° dendrites (Figure 8). The excessive

retraction induced by chronic activation of MLC-4 suggests a model in which NMY-1 function is

selectively triggered by contact between sister dendrites and subsequent UNC-6/Netrin signaling.

A downstream function for non-muscle myosin II in the UNC-6/Netrin pathway is consistent with

our finding that the Unc-6 self-avoidance defect is rescued by MLC-4DD (Figure 8). Parallel roles

for these components in axon guidance are suggested by recent results showing that non-muscle

myosin II is required for Slit and Netrin-dependent midline avoidance in the developing vertebrate

spinal cord[60]. Similarly, non-muscle myosin II mediates Semaphorin-induced axonal repulsion.

The mechanism of Semaphorin action in this case also involves the assembly of actin bundles in

the axon shaft presumably to facilitate non-muscle myosin-driven retraction[61-63]. Thus, our

findings suggest that key drivers of axonal repulsion including nascent actin assembly and non-

muscle myosin may also mediate dendrite retraction in the self-avoidance response.

RLC phosphorylation activates non-muscle myosin II by releasing myosin monomers for

assembly into bipolar structures or myosin "stacks" that interact with actin filaments to induce

translocation[37,64]. Although our results are consistent with a necessary role for non-muscle

myosin in dendrite retraction, the tips of PVD 3° dendrites are likely too narrow (~50 nm) [65] to

accommodate myosin stacks (~300 nm) [64,66]. We suggest that the actin-branching function of

Arp2/3, which we have shown mediates PVD dendrite retraction, could induce transient

expansion of the dendritic tip to allow assembly of myosin stacks. In this model, actin-

polymerization involving Arp2/3 could effectively limit ectopic NMY-1 activation in the

dendritic shaft by restricting the assembly of myosin stacks to the tips of contacting sister

dendrites. Alternatively, recent evidence indicates that myosin can also induce contraction in its

monomeric or unpolymerized form[67] which should have ready access to the dendrite tip. This

idea is consistent would our observation that GFP-tagged NMY-1 is active (e.g., rescues the self-

avoidance defect of *nmy-1*) and did not display the punctate appearance characteristic of myosin

stacks (Fenix et al., 2016) but is diffusely localized throughout the PVD cytoplasm (Figure 7B).

Parallel-acting pathways drive dendrite self-avoidance.

Although self-avoidance is perturbed in UNC-6/Netrin pathway mutants, a significant fraction

(~70%) (Figure 2D) of PVD 3° dendrites show apparently normal self-avoidance behavior[9]. One

explanation for this observation is provided by the recent discovery of an independent pathway,

mediated by MIG/14/Wntless, that is necessary for self-avoidance in an additional fraction of PVD

3° dendrites[34]. Parallel acting pathways also mediate self-avoidance in mammalian Purkinje

neurons where slit-robo signaling acts in concert with protocadherins (Gibson et al., 2014;

Lefebvre et al., 2012). The likelihood of additional effectors of self-avoidance is also suggested

by the incompletely penetrant effects of Dscam mutants in Drosophila sensory neuron self-

avoidance (Matthews and Grueber, 2011). Thus, our results in *C. elegans* mirror findings in other

organisms which together suggest that multiple mechanisms have evolved to insure dendrite self-

avoidance.

453 **The actin cytoskeleton is highly dynamic in growing and retracting PVD dendrites.**

454 We used time-lapse imaging with LifeAct::GFP, a live-cell marker for actin {Riedl:2008gw}, to

455 monitor actin PVD dendrites. The LifeAct::GFP signal is typically brighter near the distal ends of

456 growing branches in comparison to interstitial regions (Figure 5) and is highly dynamic (Movie

457 S3). The distal localization and rapidly fluctuating LifeAct::GFP signal in PVD dendrites

458 resembles recently reported "actin blobs" that actively migrate in Drosophila sensory neurons with

459 complex dendrite arbors and that appear to presage points of branch initiation [68].

460

461 **Antagonistic pathways regulate dendrite outgrowth vs retraction.**

462 Dendritic architecture is defined by the combined effects of outgrowth which expands the arbor vs

463 retraction which limits the size of the receptive field. This interaction of positive and negative

464 effects is readily observed in the development of 3° PVD dendrites. Initially, 3° dendrites

465 emanating from adjacent menorahs grow out along the body axis until contacting one another and

466 then retract to avoid overlap[9,11]. Outgrowth in this case is promoted by a multicomponent

467 receptor-ligand complex that mediates adhesive interactions with the adjacent epidermis (Díaz-

468 Balzac et al., 2016; Dong et al., 2013; Salzberg et al., 2013; Zou et al., 2016) (Figure 9E). When

469 this pathway is dysregulated by over-expression of the DMA-1 receptor in PVD neurons, for

470 example, 3° dendrites continue to adhere to the epidermis and overgrow one another[47,69]. 3°

471 branch self-avoidance also fails in mutants that disable UNC-6/Netrin signaling which we have

472 shown promotes dendrite retraction[9]. Thus, these results suggest that the net length and

473 placement of each 3° branch depends on the balanced effects of locally acting signals that either

474 extend (e.g., DMA-1) or shorten (e.g., UNC-6/Netrin) the dendrite. We confirmed this idea in

475 genetic experiments that detected strong dose-sensitive interactions between these opposing

pathways. For example, a 50% reduction of UNC-6/Netrin expression in a heterozygous *unc-6/+*

mutant does not perturb self-avoidance. Similarly, a genetic mutant (*kpc-1/+*) that partially

elevates DMA-1 also shows normal 3^o branch outgrowth. The combination of both mutations in

the double heterozygote, *unc-6/+; kpc-1/+*, however, produces a strong self-avoidance defect. Our

genetic results identified multiple regulators of actin polymerization (UNC-34/Ena/VASP, WSP-

1/WASP, ARX-5/p21, MIG-10/Lamelipodin, UNC-73/Trio) that are required for self-avoidance.

We thus propose that UNC-6/Netrin promotes actin polymerization to drive dendrite retraction

and that non-muscle myosin II interacts with a nascent actin cytoskeleton to translocate each 3^o

dendrite away from its neighbor. F-actin is also abundant and highly dynamic in growing PVD

dendrites. Notably, a different set of F-actin promoting factors (WRC, TIAM/GEF) interacts with

the DMA-1 receptor complex to drive PVD dendritic growth (Figure 9E) [25]. Thus, the challenge

for future studies is to elucidate the cell biological machinery through which opposing pathways

manipulate the actin cytoskeleton to effect either dendrite growth or retraction.

MATERIALS AND METHODS

Strains and genetics

All the strains used were maintained at 20°C and cultured as previously described (Brenner, 1974). We used the N2 Bristol strain as wild-type.

The previously described strains used in this study include: NC1404 [*wdIs52* (*pF49H12.4*::GFP)] ,NW434 [*unc-6(ev400)X;wdIs52(pF49H12.4::GFP)*], TV17,200 [*kpc-1(xr58); wyIs592(pser2prom3::GFP)*], TV9656 [*dma-1(wy686); unc-119(ed3)III; wyEx3355(pser2prom3::GFP+odr-1::RFP)*], CX1248 [*KyEx3482 (pdes-2::TBA-1::mCherry+coel::RFP*)] [9], CLP928 [*twnEx382(Pser2.3::LifeAct::EGFP, Pser2.3::mCherry, Pgcy-8::gfp)*].

Additional strains generated for this study: NC2580 *unc-34(gm104)V*; *wdIs52*, NG324 [*wsp-1(gm324)*; *wdIs52*], +/*szT1 lon-2(e678) I;* HR1184 [*nmy-1(sb115) dpy-8(e130)/szT1 X*], NC2726 [*nmy-1(sb113);wdIs52*], *unc-73(rh40)*]; *wdIs52*.

NC3284 [*wdEx1011* (*pF49H12.4*::myrUNC-5::GFP)], was produced by microinjection of plasmids pLSR18 (pF49H12.4::myrUNC-5::GFP), *pmyo-2*::mCherry and pCJS04 (pF49H12.4::mCherry).

NC3048 [*wdIs98* (pF49H12.4::ACT-1::GFP)] was produced by microinjection of plasmids pLSR03 (pF49H12.4::ACT-1::GFP) and *pceh-22*::GFP and integrated as previously described (Miller and Niemeyer, 1995).

NC3085 [*wdEx967* (pF49H12.4::mCherry::UNC-34)] was obtained by microinjection of plasmids

pCJS78(pF49H12.4::mCherry::UNC-34) and *pmyo-2*::mCherry.

NC3090 [*wdEx972* (pF49H12.4::LifeAct::GFP)] was acquired by microinjecting plasmids

pLSR06 (pF49H12.4::LifeAct::GFP) and *pmyo-2*::mCherry.

NC3029 [(pF49H12.4::*nmy-1(+)* + pF49H12.4::*nmy-1(-)*)] was obtained by microinjection of

pLSR09 (pF49H12.4::*nmy-1(+)*), pLSR10 (pF49H12.4::*nmy-1(-)*), pmyo-2::mCherry and pCJS04

(pF49H12.4::mCherry)

NC3087 [*wdEx969* (pF49H12.4::GFP::Linker::NMY-1)] was produced by microinjecting

plasmids pLSR11 (pF49H12.4::GFP::Linker::NMY-1), pCJS04 (pF49H12.4::mCherry) and

pmyo-2::mCherry.

wdEx1009 (pF49H12.4::GFP::Linker::MLC-4DD) [NC3283] was obtained by microinjection of

plasmids pLSR17 (pF49H12.4::GFP::Linker::MLC-4), pCJS04 (pF49H12.4::mCherry) and

pmyo-2::mCherry.

Molecular Biology:

Building pLSR11 (*pF49H12.4::*GFP::Linker::NMY-1::*unc-10* 3'UTR): We used the

Clonetech in-fusion HD cloning kit (Cat # 638910) to build this plasmid. The *nmy-1* genomic

region was amplified from pACP01 (*pF49H12.4::*NMY-1::mCherry) with overlapping regions

corresponding to pCJS95 (*pF49H12.4::*GFP::CED-10::*unc-10* 3'UTR). We also PCR amplified

pCJS95 with primers overlapping the *nmy-1* genomic sequences to swap CED-10 with NMY-1.

We then used the NEB site-directed mutagenesis kit to insert a 24-nucleotide glycine rich linker

(see Table 1). The resultant pLSR11 plasmid was sequenced to confirm that the linker was inserted

between the GFP sequence and *nmy-1*. See table 1 for primer sequences.

Constructing pLSR17 (*pF49H12.4*::GFP::Linker::MLC-4DD): We used the Clonetech in-
fusion HD cloning kit (Cat # 638910) to construct this plasmid. The *mlc-4* genomic region was
amplified from N2 DNA using primers with overlapping adapter complementary to pLSR11
(*pF49H12.4*::GFP::Linker::NMY-1). pLSR11 was amplified with primers containing adapter
sequences complementary to *mlc-4*. The amplified fragments were combined for the infusion
cloning to replace *nmy-1* with *mlc-4*. We then used the NEB site directed mutagenesis kit to
change MLC-4 codons for Serine 18 and Threonine 17 to nucleotide sequences that encode
Aspartate acid. The resultant pLSR17 plasmid was confirmed by sequencing.

Confocal microscopy and image analysis: Animals were immobilized with 15 mM levamisole/
0.05% tricaine and mounted on 2% agarose pads in M9 buffer as previously described [11]. Z
stack images were collected using a Nikon TiE inverted fluorescent microscope equipped with an
A1R point scanning confocal and a Plan Apo 40X 1.3NA oil immersion objective or on a Leica
TCS SP5 confocal microscope with a 40X or 63X objective. Z stacks were collected at either 0.5
or 1μm depth from the ventral nerve cord to the top of PVD cell body. Maximum projection images
were collated with either LAS-AF (Leica) or with NIS elements software (Nikon). Dendrite length
was measured using the line tool in Fiji/image J. Co-localization images were assembled using
image J. Z stack images that were collected using the LAS-AF or NIS elements software was
imported into ImageJ/Fiji for further analysis.

TIRF microscopy and analysis. Animals were immobilized with 15mM levamisole/ 0.05%
tricane and mounted on 2% agarose pads in M9 buffer. The coverslip placed on top of the agarose

pad encasing the immobilized worms is 1mm in thickness. Time-lapse datasets were acquired

using a Nikon TiE inverted fluorescence microscope outfitted with a TIRF illuminator, Apo TIRF

100x, 1.49NA oil immersion objective, and NIS-Elements software (Nikon Instruments, Inc.). The

images for the time lapse video were captured in a single plane and the TIRF angle was adjusted

for oblique illumination of 3° PVD dendrites . Datasets were imported into FIJI/ImageJ for further

analysis, including intensity quantification. The rectangle tool was used to encompass left and right

pairs of adjacent 3° dendrites with separate ROI (Regions of Interest) for image frame and the

intensities measured using the analyze tool. Resultant fluorescent intensity values were generated

with prism graphical software.

iSIM super-resolution imaging and analysis. Animals were immobilized with 0.4mM

levamisole and mounted on 10% agarose pads in M9 buffer. Z stack images were captured using

a Hamamatsu sCMOS camera on a Leica iSIM and a 100x, 1.45 NA objective. Images were

deconvolved and processed using Metamorph and Fiji/ImageJ.

Spinning disk confocal microscopy: Animals were immobilized with 0.4mM levamisole and

mounted on 10% agarose pads in M9 buffer. Volumes over time were captured using an automated

TiE inverted fluorescence microscope platform with an X1 spinning disk head (Yokogawa) and

DU-897 EM-CCD (Andor Technology), piezo-electric Z stage (Mad City Labs) and Apo TIRF

100x 1.49 NA oil immersion objective (Nikon Instruments, Inc.). Analysis of fluorescence

intensity was accomplished using NIS-Elements and Fiji/Image J software.

Quantification of fluorescence intensity in growing vs. non-growing dendrites: L2-L3 larvae were immobilized with 0.4mM levamisole and mounted on 10% agarose pads in M9 buffer for imaging with the spinning disk confocal microscope. Z-stacks of PVD dendrites were imaged at either 1 min or 1.30 minute intervals with the 100X 1.49 NA oil immersion objective. PVD dendrites (2^o, 3^o and 4^o) undergoing outgrowth with a succession of rapid extensions and retractions during the length of the time lapse videos were analyzed. NIS elements was used for measurements of LifeAct::GFP from a 1μm ROI at the tips of growing dendrites. In the same videos, fluorescence intensity was measured for an identical 1μm ROI of PVD branches (1^o, 2^o and 3^o) that were not undergoing outgrowth. Fluorescence intensity measurements for the outgrowth and no-outgrowth datasets were normalized against the maximum overall LifeAct::GFP value (Figure 5).

Quantification of fluorescence intensity in 3^o dendrites undergoing self-avoidance: Late L3 larvae were immobilized as above for imaging in the spinning disk confocal microscope. Z-stacks of PVD dendrites were imaged at either 1 or 1.30 minute intervals with the a 100X objective. Contact is defined by the point at which the maximum ferret (i.e., gap) between adjacent PVD 3^o dendrites in the mCherry channel approached zero. Nikon NIS Elements was used to obtain LifeAct::GFP and mCherry fluorescence measurements from a 1μm ROI at the tips of the left and right 3^o dendrites undergoing contact. Fluorescence intensity measurements of LifeAct::GFP and mCherry for the opposing dendrites were normalized against the maximum value for each fluorophore and plotted vs time (Figure 6A-E).

For comparisons of LifeAct::GFP signals at the tips of 3^o dendrites before and during contact, fluorescence intensities were determined from a 5 μm ROI centered at the point of contact and

from a contiguous control (non-contact) 5 μm ROI positioned either to the left or right of the

contact region (Figure 6F). The change in fluorescence intensity (ΔF) for each contact and control

region was calculated as the difference between the signal at contact (F_t) and the fluorescence

value at two time points prior to contact (F_{t-2}) normalized to F_{t-2} or $\Delta F = (F_t - F_{t-2})/F_{t-2}$ [34]. For

all measurements, the cytoplasmic mCherry marker was monitored to measure the Max Feret (in

this case, maximum distance between two points) between the tips of adjacent 3° dendrites and the

point of contact determined when the Max Feret approaches zero. Statistical comparisons (N = 8)

were performed with a 2-way ANOVA with Bonferroni correction.

Statistics: Student's t test was used for comparisons between 2 groups and ANOVA for

comparisons between three or more groups with either Tukey posthoc or Bonferroni correction for

multiple comparisons. Kolmogorov-Smirnov test was performed for comparisons between

frequency distributions of 3° dendrite length and gaps between 3° dendrites (Figures 2C and 8F,

G) .

Table 1.

Primer name	Primer sequences
myrunc-5 Vec. For.	CGGCCGCGGATAACAAATTTCA
myrunc-5 Vec. Rev.	GGATCCCATCCCGGGGAT
myrunc-5 Frag. For.	CCCGGGATGGGATCCATGggatcttctaagtctgctagcAAAC
myrunc-5 Frag. Rev.	TGTTATCCGCGGCCGCTATTTGTATAGTTCATCCATGCCATGTG
nmy-1 linker F1	GGATGCCCCTCCgtacagctcgtccatg

nmy-1 linker R1	GCAGGGGGAGGTGATGGGTGACCTGCAGTAC
mlc-4DD F1	CATGTTCGATCAGGCTCAAATTCAAG
mlc-4DD R1	GCGAACACATTGTCATCGGCTCTTTG
mlc-4 Vec. For.	GGATAACAAATTTCTATGTTTTTGTTTGTTTTTTGTCTTAATTACCA
mlc-4 Vec. Rev.	ACCTCCCCCTGCGGATG
mlc-4 Frag. For.	TCCGCAGGGGGAGGTATGGCCTCCCGCAAAACC
mlc-4 Frag. Rev.	agaaatttgttatccTTAAGCCTCATCCTTGTCCTTGGTTCC

ACKNOWLEDGMENTS

We thank C. Bargmann, K. Shen, J. Plastino, Dr. T.J. Kubiseski, E. Lundquist, P. Mains and H. Buelow for strains and reagents. We thank Chun-Liang Pan and Chien-Po Liao for sharing their work, reagents and advice prior to publication, Dylan Burnette and members of his laboratory, Aidan Fenix and Nilay Taneja, for helpful suggestions and members of the Miller lab for comments on the manuscript. Spinning disk images were obtained in the Nikon Center of Excellence at Vanderbilt University. This work was supported by NIH Grants R01 NS079611 to D. M. M, F31 NS071801 to C.J.S and F31 NS49743 to J. D. W. Some of the strains used in this work were provided by the *C. elegans* Genetics Center, which is supported by the US National Institutes of Health (NIH) National Center for Research Resources. L.S., C.J.S and J.D.W. performed experiments with advice from D.M.M., M.J.T. provided imaging expertise and advice; B.A.M

REFERENCES

1.   Delandre C, Amikura R, Moore AW. Microtubule nucleation and organization in
dendrites. cc. 2016;15: 1685–1692. doi:10.1080/15384101.2016.1172158

2.   Jan YN, Jan LY. Branching out: mechanisms of dendritic arborization. Nat Rev
Neurosci. 2010;11: 316–328. doi:10.1038/nrn2836

3.   Konietzny A, Bär J, Mikhaylova M. Dendritic Actin Cytoskeleton: Structure,
Functions, and Regulations. Front Cell Neurosci. 2017;11: 147.
doi:10.3389/fncel.2017.00147

4.   Grueber WB, Sagasti A. Self-avoidance and tiling: Mechanisms of dendrite and
axon spacing. Cold Spring Harb Perspect Biol. 2010;2: a001750–a001750.
doi:10.1101/cshperspect.a001750

5.   Fuerst PG, Bruce F, Tian M, Wei W, Elstrott J, Feller MB, et al. DSCAM and
DSCAML1 function in self-avoidance in multiple cell types in the developing
mouse retina. Neuron. 2009;64: 484–497. doi:10.1016/j.neuron.2009.09.027

6.   Lefebvre JL, Kostadinov D, Chen WV, Maniatis T, Sanes JR. Protocadherins
mediate dendritic self-avoidance in the mammalian nervous system. Nature. Nature
Publishing Group; 2012;488: 517–521. doi:10.1038/nature11305

7.   Matthews BJ, Kim ME, Flanagan JJ, Hattori D, Clemens JC, Zipursky SL, et al.
Dendrite self-avoidance is controlled by Dscam. Cell. 2007;129: 593–604.
doi:10.1016/j.cell.2007.04.013

8.   Gibson DA, Tymanskyj S, Yuan RC, Leung HC, Lefebvre JL, Sanes JR, et al.
Dendrite Self-Avoidance Requires Cell-Autonomous Slit/Robo Signaling in
Cerebellar Purkinje Cells. Neuron. Elsevier Inc; 2014;81: 1040–1056.
doi:10.1016/j.neuron.2014.01.009

9.   Smith CJ, Watson JD, Vanhoven MK, Colón-Ramos DA, Miller DM III. Netrin
(UNC-6) mediates dendritic self-avoidance. Nat Neurosci. 2012;15: 731–737.
doi:10.1038/nn.3065

10. Albeg A, Smith CJ, Chatzigeorgiou M, Feitelson DG, Hall DH, Schafer WR, et al.
C. elegans multi-dendritic sensory neurons: morphology and function. Mol Cell
Neurosci. 2011;46: 308–317. doi:10.1016/j.mcn.2010.10.001

11. Smith CJ, Watson JD, Spencer WC, O'Brien T, Cha B, Albeg A, et al. Time-lapse
imaging and cell-specific expression profiling reveal dynamic branching and
molecular determinants of a multi-dendritic nociceptor in C. elegans. Developmental
Biology. 2010;345: 18–33. doi:10.1016/j.ydbio.2010.05.502

12. Oren-Suissa M, Hall DH, Treinin M, Shemer G, Podbilewicz B. The fusogen EFF-1
controls sculpting of mechanosensory dendrites. Science. 2010;328: 1285–1288.
doi:10.1126/science.1189095

13.   Drees F, Gertler FB. Ena/VASP: proteins at the tip of the nervous system. Current
Opinion in Neurobiology. 2008;18: 53–59. doi:10.1016/j.conb.2008.05.007

14.   Goley ED, Welch MD. The ARP2/3 complex: an actin nucleator comes of age. Nat
Rev Mol Cell Biol. 2006;7: 713–726. doi:10.1038/nrm2026

15.   Takenawa T, Suetsugu S. The WASP-WAVE protein network: connecting the
membrane to the cytoskeleton. Nat Rev Mol Cell Biol. 2007;8: 37–48.
doi:10.1038/nrm2069

16.   Steven R, Kubiseski TJ, Zheng H, Kulkarni S, Mancillas J, Ruiz Morales A, et al.
UNC-73 activates the Rac GTPase and is required for cell and growth cone
migrations in C. elegans. Cell. 1998;92: 785–795.

17.   Hu S, Pawson T, Steven RM. UNC-73/Trio RhoGEF-2 Activity Modulates
Caenorhabditis elegans Motility Through Changes in Neurotransmitter Signaling
Upstream of the GSA-1/G s Pathway. Genetics. 2011;189: 137–151.
doi:10.1534/genetics.111.131227

18.   Matsuo N, Hoshino M, Yoshizawa M, Nabeshima Y-I. Characterization of STEF, a
guanine nucleotide exchange factor for Rac1, required for neurite growth. J Biol
Chem. 2002;277: 2860–2868. doi:10.1074/jbc.M106186200

19.   Demarco RS, Struckhoff EC, Lundquist EA. The Rac GTP exchange factor TIAM-1
acts with CDC-42 and the guidance receptor UNC-40/DCC in neuronal protrusion
and axon guidance. PLoS Genetics. Public Library of Science; 2012;8: e1002665.
doi:10.1371/journal.pgen.1002665

20.   Krause M, Leslie JD, Stewart M, Lafuente EM, Valderrama F, Jagannathan R, et al.
Lamellipodin, an Ena/VASP ligand, is implicated in the regulation of lamellipodial
dynamics. Developmental Cell. 2004;7: 571–583. doi:10.1016/j.devcel.2004.07.024

21.   Chang C, Adler CE, Krause M, Clark SG, Gertler FB, Tessier-Lavigne M, et al.
MIG-10/lamellipodin and AGE-1/PI3K promote axon guidance and outgrowth in
response to slit and netrin. Curr Biol. 2006;16: 854–862.
doi:10.1016/j.cub.2006.03.083

22.   Demarco RS, Lundquist EA. RACK-1 Acts with Rac GTPase Signaling and UNC-
115/ abLIM in Caenorhabditiselegans Axon Pathfinding and Cell Migration. PLoS
Genetics. 2010;6: 1–17. doi:doi:10.1371/journal.pgen.100121

23.   Shakir MA, Jiang K, Struckhoff EC, Demarco RS, Patel FB, Soto MC, et al. The
Arp2/3 activators WAVE and WASP have distinct genetic interactions with Rac
GTPases in Caenorhabditis elegans axon guidance. Genetics. 2008;179: 1957–1971.
doi:10.1534/genetics.108.088963

24.   Levy-Strumpf N, Culotti JG. VAB-8, UNC-73 and MIG-2 regulate axon polarity
and cell migration functions of UNC-40 in C. elegans. Nature Publishing Group.
2007;10: 161–168. doi:10.1038/nn1835

25.   Zou W, Dong X, Broederdorf TR, Shen A, Kramer DA, Shi R, et al. A Dendritic
Guidance Receptor Complex Brings Together Distinct Actin Regulators to Drive
Efficient F-Actin Assembly and Branching. Developmental Cell. 2018;45: 362–
375.e3. doi:10.1016/j.devcel.2018.04.008

26.   Norris AD, Sundararajan L, Morgan DE, Roberts ZJ, Lundquist EA. The UNC-
6/Netrin receptors UNC-40/DCC and UNC-5 inhibit growth cone filopodial
protrusion via UNC-73/Trio, Rac-like GTPases and UNC-33/CRMP. Development.
2014;141: 4395–4405. doi:10.1242/dev.110437

27.   Maniar TA, Kaplan M, Wang GJ, Shen K, Wei L, Shaw JE, et al. UNC-33 (CRMP)
and ankyrin organize microtubules and localize kinesin to polarize axon-dendrite
sorting. Nat Neurosci. Nature Publishing Group; 2011;15: 48–56.
doi:10.1038/nn.2970

28.   Withee J, Galligan B, Hawkins N, Garriga G. Caenorhabditis elegans WASP and
Ena/VASP proteins play compensatory roles in morphogenesis and neuronal cell
migration. Genetics. 2004;167: 1165–1176. doi:10.1534/genetics.103.025676

29.   Fleming T, Chien S-C, Vanderzalm PJ, Dell M, Gavin MK, Forrester WC, et al. The
role of C. elegans Ena/VASP homolog UNC-34 in neuronal polarity and motility.
Developmental Biology. 2010;344: 94–106. doi:10.1016/j.ydbio.2010.04.025

30.   Applewhite DA, Barzik M, Kojima S-I, Svitkina TM, Gertler FB, Borisy GG.
Ena/VASP proteins have an anti-capping independent function in filopodia
formation. Molecular Biology of the Cell. 2007;18: 2579–2591.
doi:10.1091/mbc.E06-11-0990

31.   Gates J, Mahaffey JP, Rogers SL, Emerson M, Rogers EM, Sottile SL, et al.
Enabled plays key roles in embryonic epithelial morphogenesis in Drosophila.
Development. 2007;134: 2027–2039. doi:10.1242/dev.02849

32.   Zhang W, Wu Y, Du L, Tang DD, Gunst SJ. Activation of the Arp2/3 complex by
N-WASp is required for actin polymerization and contraction in smooth muscle. Am
J Physiol, Cell Physiol. 2005;288: C1145–60. doi:10.1152/ajpcell.00387.2004

33.   Riedl J, Crevenna AH, Kessenbrock K, Yu JH, Neukirchen D, Bista M, et al.
Lifeact: a versatile marker to visualize F-actin. Nat Meth. 2008;5: 605–607.
doi:10.1038/nmeth.1220

34.   Liao C-P, Li H, Lee H-H, Chien C-T, Pan C-L. Cell-Autonomous Regulation of
Dendrite Self-Avoidance by the Wnt Secretory Factor MIG-14/Wntless. Neuron.
2018;98: 320–334.e6. doi:10.1016/j.neuron.2018.03.031

35. Piekny AJ, Mains PE. Rho-binding kinase (LET-502) and myosin phosphatase (MEL-11) regulate cytokinesis in the early Caenorhabditis elegans embryo. J Cell Sci. 2002;115: 2271–2282.

36. Vicente-Manzanares M, Ma X, Adelstein RS, Horwitz AR. Non-muscle myosin II takes centre stage in cell adhesion and migration. Nat Rev Mol Cell Biol. 2009;10: 778–790. doi:10.1038/nrm2786

37. Somlyo AP, Somlyo AV. Signal transduction and regulation in smooth muscle. Nature. 1994;372: 231–236. doi:10.1038/372231a0

38. Ono K, Ono S. Two distinct myosin II populations coordinate ovulatory contraction of the myoepithelial sheath in the Caenorhabditis elegans somatic gonad. Molecular Biology of the Cell. American Society for Cell Biology; 2016;27: 1131–1142. doi:10.1091/mbc.E15-09-0648

39. Zou W, Shen A, Dong X, Tugizova M, Xiang YK, Shen K. A multi-protein receptor-ligand complex underlies combinatorial dendrite guidance choices in C. elegans. eLife. eLife Sciences Publications Limited; 2016;5: e18345. doi:10.7554/eLife.18345

40. Díaz-Balzac CA, Rahman M, Lazaro-Pena MI, Martin Hernandez LA, Salzberg Y, Aguirre-Chen C, et al. Muscle- and Skin-Derived Cues Jointly Orchestrate Patterning of Somatosensory Dendrites. Curr Biol. 2016;26: 2379–2387. doi:10.1016/j.cub.2016.07.008

41. Liu OW, Shen K. The transmembrane LRR protein DMA-1 promotes dendrite branching and growth in C. elegans. Nat Neurosci. 2011. doi:10.1038/nn.2978

42. Salzberg Y, Díaz-Balzac CA, Ramirez-Suarez NJ, Attreed M, Tecle E, Desbois M, et al. Skin-Derived Cues Control Arborization of Sensory Dendritesin Caenorhabditis elegans. Cell. Elsevier Inc; 2013;155: 308–320. doi:10.1016/j.cell.2013.08.058

43. Dong X, Liu OW, Howell AS, Shen K. An Extracellular Adhesion Molecule Complex Patterns Dendritic Branching and Morphogenesis. Cell. 2013;155: 296–307. doi:10.1016/j.cell.2013.08.059

44. Smith CJ, O'Brien T, Chatzigeorgiou M, Spencer WC, Feingold-Link E, Husson SJ, et al. Sensory Neuron Fates Are Distinguishedby a Transcriptional Switchthat Regulates Dendrite Branch Stabilization. Neuron. Elsevier Inc; 2013;79: 266–280. doi:10.1016/j.neuron.2013.05.009

45. Schroeder NE, Androwski RJ, Rashid A, Lee H, Lee J, Barr MM. Dauer-Specific Dendrite Arborization in C. elegans Is Regulated by KPC-1/Furin. Current Biology. Elsevier Ltd; 2013;23: 1527–1535. doi:10.1016/j.cub.2013.06.058

786 46. Salzberg Y, Ramirez-Suarez NJ, Bülow HE. The Proprotein Convertase KPC-
787    1/Furin Controls Branching and Self-avoidance of Sensory Dendrites in
788    Caenorhabditis elegans. Chisholm AD, editor. PLoS Genetics. 2014;10: e1004657–
789    13. doi:10.1371/journal.pgen.1004657

790 47. Dong X, Chiu H, Park YJ, Zou W, Zou Y, Özkan E, et al. Precise regulation of the
791    guidance receptor DMA-1 by KPC-1/Furin instructs dendritic branching decisions.
792    eLife. eLife Sciences Publications Limited; 2016;5: 308. doi:10.7554/eLife.11008

793 48. Sun LO, Jiang Z, Rivlin-Etzion M, Hand R, Brady CM, Matsuoka RL, et al. On and
794    off retinal circuit assembly by divergent molecular mechanisms. Science. 2013;342:
795    1241974–1241974. doi:10.1126/science.1241974

796 49. Pollitt AY, Insall RH. WASP and SCAR/WAVE proteins: the drivers of actin
797    assembly. J Cell Sci. The Company of Biologists Ltd; 2009;122: 2575–2578.
798    doi:10.1242/jcs.023879

799 50. Castellano F, Le Clainche C, Patin D, Carlier MF, Chavrier P. A WASp-VASP
800    complex regulates actin polymerization at the plasma membrane. EMBO J. 2001;20:
801    5603–5614. doi:10.1093/emboj/20.20.5603

802 51. Chen XJ, Squarr AJ, Stephan R, Chen B, Higgins TE, Barry DJ, et al. Ena/VASP
803    proteins cooperate with the WAVE complex to regulate the actin cytoskeleton.
804    Developmental Cell. 2014;30: 569–584. doi:10.1016/j.devcel.2014.08.001

805 52. Havrylenko S, Noguera P, Abou-Ghali M, Manzi J, Faqir F, Lamora A, et al.
806    WAVE binds Ena/VASP for enhanced Arp2/3 complex-based actin assembly.
807    Molecular Biology of the Cell. American Society for Cell Biology; 2015;26: 55–65.
808    doi:10.1091/mbc.E14-07-1200

809 53. Colavita A, Culotti JG. Suppressors of ectopic UNC-5 growth cone steering identify
810    eight genes involved in axon guidance in Caenorhabditis elegans. Developmental
811    Biology. 1998;194: 72–85. doi:10.1006/dbio.1997.8790

812 54. Bashaw GJ, Kidd T, Murray D, Pawson T, Goodman CS. Repulsive axon guidance:
813    Abelson and Enabled play opposing roles downstream of the roundabout receptor.
814    Cell. 2000;101: 703–715.

815 55. McConnell RE, Edward van Veen J, Vidaki M, Kwiatkowski AV, Meyer AS,
816    Gertler FB. A requirement for filopodia extension toward Slit during Robo-mediated
817    axon repulsion. The Journal of Cell Biology. 2016;213: 261–274.
818    doi:10.1083/jcb.201509062

819 56. Hung R-J, Terman JR. Extracellular inhibitors, repellents, and
820    semaphorin/plexin/MICAL-mediated actin filament disassembly. Cytoskeleton
821    (Hoboken). John Wiley & Sons, Inc; 2011;68: 415–433. doi:10.1002/cm.20527

57.   Lin C, Espreafico E, Mooseker M, Forscher P. Myosin drives retrograde F-actin
flow in neuronal growth cones. Neuron. 1996;16: 769–782.

58.   Mitchison TJ, Cramer LP. Actin-based cell motility and cell locomotion. Cell.
1996;84: 371–379.

59.   Medeiros NA, Burnette DT, Forscher P. Myosin II functions in actin-bundle
turnover in neuronal growth cones. Nat Cell Biol. 2006;8: 215–226.
doi:10.1038/ncb1367

60.   Murray A, Naeem A, Barnes SH, Drescher U, Guthrie S. Slit and Netrin-1 guide
cranial motor axon pathfinding via Rho-kinase, myosin light chain kinase and
myosin II. Neural Dev. 2010;5: 16. doi:10.1186/1749-8104-5-16

61.   Brown JA, Wysolmerski RB, Bridgman PC. Dorsal root ganglion neurons react to
semaphorin 3A application through a biphasic response that requires multiple
myosin II isoforms. Molecular Biology of the Cell. 2009;20: 1167–1179.
doi:10.1091/mbc.E08-01-0065

62.   Brown JA, Bridgman PC. Disruption of the cytoskeleton during Semaphorin 3A
induced growth cone collapse correlates with differences in actin organization and
associated binding proteins. Devel Neurobio. 2009;69: 633–646.
doi:10.1002/dneu.20732

63.   Gallo G. RhoA-kinase coordinates F-actin organization and myosin II activity
during semaphorin-3A-induced axon retraction. J Cell Sci. 2006;119: 3413–3423.
doi:10.1242/jcs.03084

64.   Fenix AM, Taneja N, Buttler CA, Lewis J, Van Engelenburg SB, Ohi R, et al.
Expansion and concatenation of non-muscle myosin IIA filaments drive cellular
contractile system formation during interphase and mitosis. Molecular Biology of
the Cell. 2016;27: 1465–1478. doi:10.1091/mbc.E15-10-0725

65.   Albeg A, Smith CJ, Chatzigeorgiou M, Feitelson DG, Hall DH, Schafer WR, et al.
C. elegans multi-dendritic sensory neurons: morphology and function. Mol Cell
Neurosci. 2011;46: 308–317. doi:10.1016/j.mcn.2010.10.001

66.   Shutova MS, Spessott WA, Giraudo CG, Svitkina T. Endogenous species of
mammalian nonmuscle myosin IIA and IIB include activated monomers and
heteropolymers. Curr Biol. 2014;24: 1958–1968. doi:10.1016/j.cub.2014.07.070

67.   Shutova M, Yang C, Vasiliev JM, Svitkina T. Functions of nonmuscle myosin II in
assembly of the cellular contractile system. Parsons M, editor. PLoS ONE. Public
Library of Science; 2012;7: e40814. doi:10.1371/journal.pone.0040814

68.   Nithianandam V, Chien C-T. Actin blobs prefigure dendrite branching sites. The
Journal of Cell Biology. 2018;217: 3731–3746. doi:10.1083/jcb.201711136

69.   Luo L, Wen Q, Ren J, Hendricks M, Gershow M, Qin Y, et al. Dynamic encoding of
859   perception, memory, and movement in a C. elegans chemotaxis circuit. Neuron.
860   2014;82: 1115–1128. doi:10.1016/j.neuron.2014.05.010

 FIGURE LEGENDS

 Figure 1: UNC-6/Netrin signaling mediates sister dendrite self-avoidance.

(A-B) Representative images and tracings of PVD sensory neurons in wild-type (A) and *unc-6* (B)

L4-stage larvae. PVD morphology was visualized with PVD::GFP or PVD::mCherry markers.

PVD cell body marked with an asterisk. (A) 1°, 2°, 3° and 4° dendritic branches and single axon

(blue arrowhead) are denoted. (B) Arrowheads point to overlaps between adjacent 3° dendrites

that failed to self-avoid in *unc-6* mutants. Scale bars are 5 μm. C) Summary of self-avoidance in

3° PVD dendrites denoting growth, contact and retraction. (D) Model of self-avoidance

mechanism involving reciprocal contact-dependent repulsion mediated by UNC-6/Netrin and its

receptors, UNC-40/DCC and UNC-5[9].

Figure 2: Constitutively active UNC-5 drives hyper-retraction of 3° PVD dendrites.

(A-D) Representative images and tracings of 3° dendrites anterior to PVD cell body in (A) wild

type (red) and (B) PVD::MYR::UNC-5 (blue) (Late L3-early L4 animals) (C) Gaussian fit to

distribution of lengths of PVD::mCherry-labeled 3° dendrites showing shorter average length for

MYR::UNC-5 (n = 156, 7 animals) *versus* wild type (n = 159, 8 animals), p = 0.03, Kolmogorov-

Smirnov statistical test. (D) Schematics of intact, wild-type UNC-5 receptor vs membrane-

associated, constitutively-activated myristylated-UNC-4 (MYR::MYR-5). (E-F) Dynamicity of 3°

dendrites in wild-type vs MYR::UNC-5-expressing PVD neurons labeled with PVD::GFP. (E)

Schematic (top) depicts displacement of the tip of a 3° dendrite relative to a fiduciary mark (vertical

dashed red line). Scale bar is 10 μm. Representative plots of displacement for wild-type (red) and

MYR::UNC-5 (blue) 3° dendrites over time. (F) The dynamic saltatory growth and retraction of

3° dendrites in the wild type vs the relative quiescence of MYR::UNC-5-expressing PVD neurons

is correlated with elevated standard deviation of tip movement from a fiduciary point in wild-type

vs MYR::UNC-5 expressing PVD neurons (p = 0.04 from Student's t-test. n = 4). (G) Self-

avoidance defects (percentage of overlapping 3° branches) are rare (< 5%) in wild type and

MYR::UNC-5. The *unc-6* self-avoidance defect is rescued by PVD expression of MYR::UNC-5.

Error bars denote mean and M. ***p<0.0001. n ≥ 15. from ANOVA with Tukey's posthoc test.

Scored in a PVD::mCherry background.

Figure 3. Mutations in genes that promote actin polymerization disrupt PVD dendrite self-

avoidance.

(A) Representative images (top) and schematics (bottom) of the actin cytoskeleton (Green) labeled

with PVD::ACT-1::GFP (A) and microtubules (Magenta) marked with *pdes-2*::TBA-1::mCherry.

PVD cell bodies (asterisks) and axons (arrowheads) are noted. (B) Representative images of

PVD::GFP for wild type, *unc-34*, *mig-10*, *unc-73*, *wsp-1* and *arx-5RNAi*. Asterisks denote PVD

cell bodies and arrowheads point to overlaps between adjacent PVD 3° dendrites. (C-E) Self-

avoidance defects are plotted in a PVD::GFP background (mean and SEM) as the percentage of

overlapping 3° dendrites for each genotype. ***p<0.001, *p<0.05, ANOVA with Tukey's posthoc

test, n ≥ 15 animals for all backgrounds. (C) Self-avoidance defects in *unc-73*, *mig-10* and *unc-*

34/Ena/VASP exhibit significant self-avoidance defects. *unc-34/Ena/VASP* mutants are rescued by

PVD   expression   of   UNC-34   (PVD::mCherry::UNC-34).   Myristyolated   UNC-5

(PVD::MYR::UNC-5) fails to rescue the self-avoidance defects of an *unc-34* mutant. (D) PVD

self-avoidance defects for *arx-5RNAi*, *unc-6*, *wsp-1* and *unc-6; wsp-1* mutants and compared to

wild-type and E.V. *(Empty Vector) RNAi controls. ***p<0.001, NS = Not Significant, ANOVA

with Tukey's posthoc test, n ≥ 15 animals. Scale bar is 10 μm.

Figure 4. Localization and trafficking of PVD::UNC-34::mCherry puncta in PVD.

(A-B) Leica iSIM images of PVD neurons (late L3 larvae) labeled with PVD::LifeAct::GFP

(green) and PVD::mCherry::UNC-34 (Magenta). Merge shown at top right. Insets depict punctate

mCherry::UNC-34 in 1°, 2° and 3° PVD dendrites. (B) Note punctate mCherry::UNC-34 near

tips (white arrowheads) of 3° dendrites Scale bars are 5 μm. (C-D) Kymographs of

PVD::mCherry::UNC-34 in PVD 3° dendrite captured in the Leica iSIM microscope. Moving

(green arrowheads) vs stationary (red arrowheads) mCherry::UNC-34 puncta are denoted. The

distal end of the 3° dendrite is indicated by an asterisk and a red dashed arrow line points toward

the PVD cell soma. Scale bar is 10 μm.

Figure 5. The actin cytoskeleton is highly dynamic in PVD dendrites.

(A) Representative images of PVD::LifeAct::GFP and PVD::mCherry labeled 2° dendrite

undergoing outgrowth through time (late L3 larvae). (B) Fluorescent intensity was measured from

regions of PVD dendrites undergoing outgrowth vs no-outgrowth (see Materials and Methods).

***$p<0.001$, N = 118 measurements from 6 dendrites. (C) Representative image showing enriched

LifeACT::GFP signal in distal, growing PVD dendrites labeled with PVD::mCherry. (D) Rapid

changes in the normalized fluorescence intensity of PVD::LifeAct::GFP versus the cytoplasmic

marker, PVD::mCherry, were quantified as the standard deviation of the normalized fluorescent

intensity measured at the tips of growing dendrites in a wild-type background. See Movie S3 and

Materials and Methods) N=22. (E) Normalized fluorescence intensity measurements of

PVD::LifeAct::GFP (green) vs cytoplasmic PVD::mCherry (red) at the tip of a growing late L3

stage PVD 3°dendrite from a representative time lapse movie

Figure 6: (A-D) Fluorescent intensity traces in wild-type PVD dendrites during 3° dendrite contact based self-avoidance. Fluorescent intensity measurements of PVD::LifeAct::GFP and PVD::mCherry were acquired from a 1μm region at the tip of the growing and contacting left and right 3° dendrites (Late L3-L4 animals). Measurements were normalized against the maximun intensity during a ~10 minute interval that includes at least one contact event. The cytoplasmic mCherry signal was used to measure the gap between the left and right 3° dendrites undergoing contact and plotted against time. The period of contact (vertical shading) corresponds to a minimum value for the gap (0-0.14μm) between contacting left and right 3° dendrites. (E) Representation of 3° dendrites with color schemes to indicate left and right 3° dendrites and regions quantified. (F) Measurements from contact and non-contact (control) regions of 3° dendrites undergoing self-avoidance did not detect a significant increase in LifeAct::GFP at contact between the tips of adjacent 3° dendrites. Normalized LifeAct::GFP fluorescence intensity values from these regions (see Materials and Methods) were plotted vs time, N = 8, NS = Not Significant, 2-way ANOVA with Bonferroni correction.

Figure 7. NMY-1/non-muscle myosin II mediates dendrite self-avoidance.

(A) Representative images (left) and tracings (right) of wild-type and *nmy-1(sb113)* labeled with PVD::GFP. PVD cell soma (asterisks) and overlapping 3° dendrites (arrowheads) are noted. The PDE neuron is marked with an asterisk and is adjacent to the PVD cell-body. (B) NMY-1 fused to GFP via a linker peptide (PVD::GFP::NMY-1) (top) expressed in PVD with PVD::mCherry (bottom) rescues self-avoidance defects (C) of *nmy-1(sb113)*. Scale bars are 5μm. (D) Quantification of self-avoidance defects in wild type and *nmy-1*. Self-avoidance defects for *nmy-*

1 mutants and *nmy-1* cell-specific RNAi (csRNAi) increase the percentage of overlapping 3°

branches. PVD expression of GFP-labeled NMY-1 (PVD::GFP::NMY-1) rescues the *nmy-*

1(sb113) self-avoidance defect. Scatter plots with mean and SEM, ***p<0.01, One way ANOVA

with Tukey's posthoc test. N = 15-20 animals per genotype. All defects were quantified in a

PVD::GFP background. Scale bar is 5 μm.

Figure 8. Phosphomimetic activation of myosin regulatory light chain, MLC-4, shortens 3°

PVD dendrites and rescues the *unc-6* dendritic self-avoidance defect.

(A) Phosphorylation of Regulatory Light Chain (RLC) induces a conformational change that

activates the myosin protein complex composed of myosin heavy chain (MHC), essential light

chain (ELC) and RLC. (B) Threonine and serine phosphorylation sites (red) in conserved regions

of *C. elegans*, *Drosophila* and human myosin RLCs. (C) Phosphomimetic construct,

PVD::GFP::MLC-4-DD, fused to GFP with a linker peptide for expression with PVD promoter.

(D-E) Representative images (top) and schematics (bottom) of PVD morphology anterior to cell

body (asterisk) denoting 3° dendrites for (D) wild type (red) and (E) MLC-4DD (blue) (L3 stage

larvae). Wild-type image also used in Figure 2A. Scale bars are 10 μm. 3° dendrite length (F) is

shorter and gaps between 3° dendrites (G) is wider for MLC-4DD versus wild type. Gaussian fit

was plotted with bin center values (μm) on the X-axis and the number of values that fall under

each bin center on the Y axis, ***p<0.001, Kolmogorov-Smirnov statistical test for cumulative

distributions of PVD 3° dendrite lengths and length of gaps in MLC-4DD (n = 141, 6 animals)

versus wild type (n = 156, 8 animals). (J) PVD expression of MLC-4DD rescues the *unc-6* mutant

self-avoidance defect, ***p<0.0001, One way ANOVA with Tukey's posthoc test. Mean and

SEM are shown. n ≥9 animals per genotype. Quantified with PVD::mCherry. Scale bar is 10 μm.

Figure 9. Opposing pathways regulate dendrite outgrowth and retraction.

Representative images and schematics of PVD in (A) *kpc-1/kpc-1,* (B) *kpc-1/+; unc-6/+* and (C)

kpc-1 +/+ dma-1; unc-6/+. PVD soma (asterisks) and overlapping 3° dendrites (arrowheads) are

noted. (D) Quantification of self-avoidance defects for each genotype. Note that all mutant alleles

are recessive but that *kpc-1/+; unc-6/+* double heterozygotes show a strong self-avoidance defect

that is suppressed in *kpc-1 +/+ dma-1; unc-6/+* triples. ***$p<0.0001$, one way ANOVA with

Tukey's posthoc test. Mean and SEM are indicated. N >7 animals per genotype. PVD is labeled

with *ser2prom3*::GFP for all images. Scale bar is 10 μm. Mutant alleles were: *kpc-1(xr58), unc-*

6(ev400) and *dma-1(wy686).* (E) Model of opposing pathways that promote actin polymerization

to define dendrite length. KPC-1 antagonizes surface expression of DMA-1 that interacts with an

epidermal receptor composed of SAX-7/CAM, MNR-1 and LECT-2 and functions with HPO-30

to promote actin polymerization and dendrite outgrowth. UNC-6/Netrin interacts with UNC-5 to

activate UNC-34/Ena/VASP which functions with WSP-1/WASP and the Arp2/3 complex to drive

F-actin assembly for NMY-1/myosin-dependent dendrite retraction in the self-avoidance response.

1007

1008

1009

1010

1011

1012

 SUPPORTING INFORMATION.

 Imaging and quantification of self-avoiding 3° dendrites: L3 animals of strain

 CLP928 *twnEx382[Pser2.3::LifeAct::EGFP, Pser2.3::mCherry, Pgcy-8::gfp]* [34] were

 immobilized for time-lapse imaging with 1μl of agarose beads (Polybead carboxylate

 microspheres, Polysciences, Catalog # 15913-10, 0.05 μm) and mounted on a 10% agarose pad.

 Z-stacks of PVD dendrites (0.5 um depth, 5 slices) were imaged at 35 second intervals with a 40X

 objective on a Nikon spinning disk confocal microscope. Contact between the 3° dendrites was

 established as a function of Max Feret (maximum distance between 2 points). Max Feret (in pixels)

 was measured in the mCherry channel (cystosolic marker) with NIS elements and contact defined

 as by the time-point at which the Max-Feret approaches zero. LifeAct::GFP and mCherry signals

 were measured for a ROI of 2 μm at the tips of the left and right dendrites in a single Z plane. Left

 and right fluorescence intensities were summed for comparison to a control ROI (2 μm). The

 change in fluorescence intensity (ΔF) for each contact and control region was calculated as the

 difference between the signal at contact (F_t) and the fluorescence value at two time points prior to

 contact (F_{t-2}) normalized to F_{t-2} or $\Delta F = (F_t - F_{t-2})/F_{t-2}$ [34]. These values were computed for

 contact and control ROIs for LifeAct::GFP and mCherry. Statistical tests used for comparison

 between the datasets (LifeAct::GFP contact vs control, mCherry contact vs control, LifeAct::GFP

 contact vs. mCherry contact) were 2-way ANOVA with Bonferroni correction and multiple t-tests

 with a False Discovery Rate of 1%. Both these tests yielded similar results.

 Supplemental Figure 1. Mutations in *unc-34/Ena/VASP* and *unc-73/Trio* disrupt PVD

 dendrite morphology.

(A) Quantification of 1° dendrite outgrowth and (B) number of 2° dendrites in wild-type, *unc-34*,

unc-73, *wsp-1* and PVD::mCherry::UNC-34;*unc-34*. Defective 1° dendrites show altered

alignment and/or extension defects. Mutations in *unc-34* and *unc-73* but not *wsp-1* exhibit defects

in 1° dendrite outgrowth and fewer 2° dendrites. Note that cell-autonomous expression of UNC-

34::mCherry does not rescue 1° and 2° dendritic defects but does restore self-avoidance (Figure

3C). (C) Representative images of PVD morphological defects in an *unc-34* mutant. Note

premature termination of 1° dendrites (red arrowheads). Scale bars are 10 μm. Mutant alleles are

unc-34(gm104), unc-73(rh40), wsp-1 (gm324).

1044
Supplemental figure 2. (A) Schematic (left) of adjacent 3° dendrites and representative time

lapse series (35 sec intervals) of PVD::mCherry and PVD::LifeAct::GFP in 3° dendrites showing

self-avoidance response. Point of contact is indicated by an asterisk and period of contact denoted

with dashed blue outline. Scale bar is 10 μm. (B) Schematic depicting ROIs for fluorescence

intensity measurements at the tips of adjacent (left vs right) 3° dendrites that undergo contact and

at an adjacent non-contact (control) region. (C-E) Graphical representation of the change in

fluorescence intensity at each time point vs that of the t = -2 time interval before contact at t = 0;

measurements were normalized to th[34] (See Materials and Methods). Normalized fluorescence

intensity values were plotted against time (min) for (C) LifeAct::GFP contact vs control, (D)

LifeAct::GFP contact vs mCherry contact, (E) mCherry contact vs mCherry control. The values at

each time-point were compared by 2-way Anova with Bonferonni correction, NS (Not Significant)

N=14.

1057

1058 **Movie S1: Time-lapse video representing the dynamic movement of 3° dendrites in a wild-**

1059 **type background.** Images of PVD::GFP were recorded (late L3-early L4) at 15 sec intervals with

1060 a 100X objective in a Nikon spinning disk microscope (see Materials and Methods). The entire

1061 length of the video is 240 seconds. The movie was edited using NIS elements software with Bayes

1062 advanced denoising. Green arrowheads denote the tips of 3° dendrites. Scale bar is 25 μm.

1063

Movie S2: Time-lapse video of 3° PVD dendrites in a MYR::UNC-5 background. Images of

PVD::GFP in the MYR::UNC-5 strain were recorded (late L3-early L4) at 15 sec intervals with a

100X objective in a Nikon spinning disk microscope (see Materials and Methods). The entire

length of the video is 12 minutes. Boxes denote tips of 3° dendrites showing limited outgrowth**.**

The movie was edited using NIS elements software with Bayes advanced denoising. Scale bar is

10μm.

1070

1071 **Movie S3: Time-lapse video showing the dynamicity of LifeAct::GFP during in 3° PVD**

1072 **dendrites**. Images of PVD::mCherry (left) and PVD::LifeAct::GFP (right) and were collected at

1073 1.3 min intervals with a 100 X objective in a Nikon spinning disk microscope (see Materials and

1074 Methods). The entire length of the video is 60 minutes. The movie was edited using NIS elements

1075 with automatic deconvolution. Scale bar is 5 μm. Spherical objects are autofluorescent granules.

1076

Movie S4: Time-lapse video demonstrating the rapid accumulation of LifeAct::GFP after

contact between 3° sister dendrites. Images of PVD::mCherry (left) and PVD::LifeAct::GFP

(right) and were collected at 1.3 min intervals with a 100 X objective? in a Nikon spinning disk

microscope (see Materials and Methods). The entire length of the video is 60 minutes. The movie

was edited using NIS elements software with automatic deconvolution. Arrows track the moving

3º dendrite undergoing contact and retraction. Scale bar is 5 μm.

1083

Movie S5: Time-lapse video demonstrating the rapid accumulation of LifeAct::GFP after

contact between 3° sister dendrites. Images of PVD::LifeAct::GFP were captured at 15 second

intervals using a 100X objective in a Nikon TIRF microscope for a total of 25 minutes at the L3-

L4 stage (see Materials and Methods). The movie was edited using NIS elements with Bayes

advanced denoising. The bottom panel depicts a fluorescence intensity profile through time

(bottom) for the ROI (green). Scale bar is 5 μm.

1090

1091   **Movie S6: Time-Lapse video demonstrating the trafficking of UNC-34::mCherry puncta in**

1092   **PVD 3 dendrites.** Images of PVD::mCherry::UNC-34 were captured continuously on a single

1093   frame using a 100X objective in a iSIM microscope for 5 minutes (See Materials and Methods).

1094   The movie was edited using a Metamorph software and deconvolved using Fiji software. Scale bar

1095   is 5 μm.

1096

1097

1098

1099
1100

1101

Figure 1: PVD 3° dendrites undergo self-avoidance in an UNC-6/Netrin dependent manner

A. wild-type

B. *unc-6*

C. outgrowth → contact → retraction

UNC-6 expressing cells

UNC-6
UNC-5
UNC-40

Figure 2: Constitutively active UNC-5 drives hyper-retraction of 3º PVD dendrites.

A. Wild-type

B. MYR::UNC-5

C. Length of 3° dendrites (2µm bins)

D.

Wild-type MYR::UNC-5

E.

F.

G.

Figure 3: Mutations in genes that promote actin polymerization disrupt PVD dendrite self-avoidance.

Figure 4: Localization and trafficking of PVD::UNC-34::mCherry puncta in PVD.

Figure 5: The actin cytoskeleton is highly dynamic in PVD dendrites

Figure 6: Dynamic changes in actin polymerization during contact based retraction

Figure 7: NMY-1/non-muscle myosin II mediates dendrite self-avoidance.

Figure 8: Phosphomimetic activation of myosin regulatory light chain, MLC-4, shortens 3º PVD dendrites and rescues the *unc-6* dendritic self-avoidance defect.

Figure 9: Opposing pathways regulate dendrite outgrowth and retraction.