Systemic loss of Sarm1 protects Schwann cells from chemotoxicity by delaying axon degeneration

Weili Tian*, Tim Czopka§ and Hernán López-Schier*

* Sensory Biology & Organogenesis, Helmholtz Zentrum Munich, Germany

§ Institute of Neuronal Cell Biology, Technical University of Munich, Germany

Correspondence to: hernan.lopez-schier@helmholtz-muenchen.de

Abbreviated title: Sarm1 and glioprotection

Keywords: Wallerian degeneration, Sarm1, Schwann cells, regeneration

ABSTRACT

Protecting the nervous system from chronic effects of physical and chemical stress is a pressing clinical challenge. The obligate pro-degenerative protein Sarm1 is essential for Wallerian axon degeneration. Thus, blocking Sarm1 function is emerging as a promising neuroprotective strategy with therapeutic relevance. Yet, the conditions that will most benefit from inhibiting Sarm1 remain undefined. Here we combine genome engineering, pharmacology and high-resolution intravital videmicroscopy in zebrafish to show that genetic elimination of Sarm1 increases Schwann-cell resistance to toxicity by diverse chemotherapeutic agents after axonal injury. Synthetic degradation of Sarm1-deficient axons reversed this effect, suggesting that glioprotection is a non-autonomous effect of delayed axon degeneration. Moreover, loss of Sarm1 does not affect macrophage recruitment to nerve-wound microenvironment, injury resolution, or neural-circuit repair. These findings anticipate that interventions aimed at inhibiting Sarm1 can counter heightened glial vulnerability to chemical stressors and may be an effective strategy to reduce chronic consequences of neurotrauma.

INTRODUCTION

The peripheral nerves that communicate skin, muscle and sensory organs with the

brain must maintain functionality throughout life despite frequent stress and

trauma (1-6). Loss of integrity of peripheral neurons and associated cells, including

glia, is a common occurrence in severe neurological dysfunctions that include

weakness, pain and loss of sensation (7). Glial loss leads to nerve demyelination,

defasciculation and neuronal death. Nerve injury triggers axon fragmentation and

degeneration. In turn, this induces the dedifferentiation of associated glial cells,

which enhances glial role in nerve repair. However, dedifferentiation also makes

glia vulnerable to degeneration after protracted denervation (8-12). Damaged axons

undergo Wallerian degeneration (13), which is triggered by the evolutionary ancient

pro-degenerative protein Sarm1 (Sterile Alpha and TIR Motif containing 1) (14-17).

Sarm1 is a modular protein that contains two sterile-alpha motifs (SAM), one

mitochondrial association (MT) and one TIR domain (14). Nerve injury activates

Sarm1 by TIR dimerization, which is sufficient to induce the axon degeneration (18,

19). Mechanistically, Sarm1 activation results in the loss of nicotinamide adenine

dinucleotide (NAD+) in damaged axons (20-22). The TIR domain acts as a

NADase (23), explaining why over-expression of the NAD+ synthesizing enzyme

NMNAT1 inhibits the degradation of severed axons by sustaining high levels of

NAD+ downstream of Sarm1 (24). Moreover, the loss of Sarm1 blocks the

degeneration of injured axons, and forced activation of Sarm1 induces axon

destruction in the absence of injury. Therefore, Sarm1 is a hierarchical regulator of

a signaling pathway that is necessary and sufficient for axonal degradation (22).

Upon peripheral-nerve injury, glial Schwann cells acquire a specialized function

that promotes the clearance of axon fragments ahead of axonal re-growth from the

proximal stump (25). Expeditious axonal regeneration is important because

sustained Schwann-cell denervation leads to protracted loss of glial terminal

phenotype and eventual death. In turn, glial loss impairs regenerating nerve

myelination and circuit repair, transforming acute neuropathies into irreversible

chronic neurological dysfunction. Therefore, inhibiting or delaying axon

destruction has been hypothesized as an effective strategy to counter heightened

Schwann-cell vulnerability to additional stressors that include metabolic imbalance

and drugs (26, 27). This idea has sparked intense efforts to identify specific

molecular inhibitors of axon degeneration for clinical applications (28, 29). As a

discrete hierarchical factor that is essential for axon degeneration, Sarm1 is an ideal

target for pharmacological interventions (18). However, the conditions that would

most benefit from inhibiting Sarm1 systemically are yet to be defined (30, 31). Here

we address the above issue using the powerful genetics of zebrafish, a small

vertebrate whose nervous system is anatomically simple but functionally similar to

that of mammals (32-34). Crucially, the zebrafish larva is ideal to study Schwann-

cell biology in the natural context of the behaving animal (35-42). By characterizing

loss-of-function mutations in Sarm1, we provide novel insights into the cellular basis

of axon-glia interactions, and provide data that encourage the quest to identify and

develop Sarm1 inhibitors for clinical applications.

RESULTS

Identification and mutagenesis of Sarm1 in zebrafish

The amino-acid sequence of Sarm1 is well conserved across species (15, 43). To

identify Sarm1 orthologs in zebrafish, we scanned publicly accessible genomic data

(***Danio rerio*** reference genome assembly version GRCz11) by a BLAST search

using the TIR domain, which is present in all known Sarm1 proteins (15, 44). This

exploration yielded a single candidate ***locus*** in chromosome 15. No other part of

the zebrafish genome appears to harbor Sarm1 paralogs. The genomic structure of

the putative zebrafish Sarm1 reminisces that of other species, containing 8 exons

that code for a protein of 713 amino acids, with the typical N-terminal auto-

inhibitory domain, two central SAM multimerization domains and a C-terminal

TIR degeneration domain (Figure 1A). Similar to ***D. melanogaster***, however, ***D.***

rerio Sarm1 lacks an obvious mitochondria-targeting sequence (MT). To test

whether the identified gene produces a protein with the expected functional role,

we used CRISPR/Cas9-mediated genome modification to generate loss-of-function

mutations in Sarm1. By targeting exon 1, we obtained germ-line transmission of

two alleles: ***sarm1^hzm13*** and ***sarm1^hzm14*** (Supplementary Figure 1A-B). The hzm13

allele introduces an 11-base deletion and T/C mutation, resulting in a frame shift

and premature stop codon. hzm14 is a 7-base deletion and AG/GA mutation that

also generates a frame shift and premature stop codon. Analysis of protein extracts

from wild-type embryos by Western-blot using an antibody to Sarm1 revealed a

single band of approximately 80kDa, which agrees with the expected size of the

full-length protein (Supplementary Figure 1C). This band was absent in protein

extracts from homozygous ***sarm1ʰᶻᵐ¹³*** zebrafish embryos. Of note, because this

antibody recognizes an epitope in the C-terminus of Sarm1, it does not allow to

discriminate between the expression of a truncated protein lacking all the domains

with known function, and the complete absence of Sarm1 induced by nonsense-

mediated mRNA decay. Homozygous ***sarm1ʰᶻᵐ¹³*** mutants display no overt

anatomical defects (Supplementary Figure 1D-E), are viable and develop into fertile

adults. Furthermore, a simple assay for sensorimotor function that consists of

eliciting the escape response after tactile stimuli showed that the displacement

distance and the average acceleration were no different between wild type and

Sarm1 mutants (Supplementary Figure 1F-G) (45).

Functional conservation of Sarm1 in axon degeneration

Although Sarm1 has already been extensively studied in neurons of ***Drosophila***

and mice, we deemed necessary to assess the effects of systemic loss of Sarm1 on

neuronal and non-neuronal cells in larval zebrafish. The reason is that no single

study has addressed Sarm1 function holistically, in a single organism, by intravital

high-resolution microscopy. To this end, we used several parameters of nervous-

system structure and function by combining ***sarm1ʰᶻᵐ¹³*** with transgenes expressing

fluorescent markers in the sensory neurons of the mechanosensory lateral line, as

well as in their associated Schwann cells (46, 47). The lateral-line system is ideal for

such ***in vivo*** studies under normal and altered conditions (48-51). It combines the

organization of a typical vertebrate sensory system with the amenability for

controlled experimental interventions that include microsurgery, pharmacology and

optogenetics (41, 42, 52, 53). We found that loss of Sarm1 does not affect the development and structure of the lateral-line sensory pathway (Supplementary Figure 1H-N). For a more detailed analysis, we fluorescently marked mitochondria in sensory neurons by expressing the mitochondria targeting sequence of the Cytochrome-C oxidase subunit 8A fused to mCherry (54). We chose to study mitochondria because they are dynamic organelles that distribute throughout the neuron by active transport mediated by molecular motors that move along microtubule tracks. In axons, microtubules are polarized such that their plus ends are directed toward the axon terminals, in turn orienting the movement of distinct molecular motors in the antero- and retro-grade directions (55). Intra-axonal movement direction reflects mitochondrial fitness because stressed organelles are biased in the retrograde direction (56, 57). Therefore, axonal mitochondria represent an optimal proxy for neuronal polarization, intracellular dynamics and overall health. We found a qualitatively similar number, density and distribution of mitochondria in the peripheral axons of wild-type and Sarm1-mutant animals. Kymographic analysis revealed a majority of static large mitochondrial groups, and some smaller fragments moving substantial distances at constant velocity in the anterograde and retrograde directions (Figure 1B-C). Importantly, quantifications showed no significant differences in the number and spatial distribution of axonal mitochondria (Figure 1D), or movement velocity and direction between wild-type and Sarm1-mutant specimens (Figure 1E).

Next, we addressed functional conservation of Sarm1 in zebrafish using a previously established bioassay of neurotrauma *in vivo*, which employs laser-mediated severing of individualized axons using single-neuron fluorescent-protein expression and high-resolution microscopy (41,58). To precisely cut axons, we focused an ultraviolet laser beam to a discrete region of the peripheral nerve. Upon severing, the distal axon segment quickly degenerated in wild-type specimens, whereas the proximal segment that stays associated with the neuronal perikaryon remained viable (Figure 1F). Kinetic analysis shows that Sarm1-deficient severed axons remained stable for over 5 days, whereas severed wild-type axons degraded within 8 hours (Figure 1G). The degeneration-resistant phenotype of Sarm1 mutants was eliminated by the transgenic introduction of a fluorescent-tagged full-length Sarm1 in mutant neurons (Figure 1H), demonstrating functional conservation of zebrafish Sarm1.

Loss of Sarm1 does not affect focal damage resolution

Physical injury often results in inflammatory responses that recruit immune cells to the wound. After nerve injury, injury-mediated production of reactive oxygen species (ROS) as well as chemoattractants produced by Schwann cells recruits inflammatory cells, including macrophages. Studies *ex vivo* using mammalian cultured neurons showed that Sarm1 acts downstream of mitochondrial ROS generation. Therefore, we decided to test *in vivo* if the loss of Sarm1 or the absence of Schwann cells would affect macrophage recruitment to the site of injury. To this end, we generated a transgenic line expressing membrane-targeted EGFP

under the control of the macrophage-specific promoter Mfap4. We combined this line with Tg[SILL:mCherry] and mutations in Sarm1 or Erbb2. Loss of Erbb2 in zebrafish impairs Schwann-cell migration along lateral-line axons and leads to nerve unmyelination and defasciculation (53, 59). We injured nerves with a laser and assessed macrophage behavior at high resolution. We found that the onset of recruitment and number of macrophages at the wound did not differ between wild-type specimens and Sarm1 or Erbb2 mutants (Figure 2A and Supplementary Movies 1-3). Macrophages arrived from various locations and moved in a qualitatively indistinguishable manner along the proximal and distal part of the axons in wild-type and mutant specimens. Quantitative imaging showed that the retention time of macrophages at the proximal side of the wound was unaltered in Sarm1 or Erbb2 mutants, although on average retention was marginally lengthened in the absence of Schwann cells (Figure 2B). In wild type, Sarm1 and Erbb2 mutants, macrophages engulfed debris locally at the injury site, and did not appear to be involved in the degradation of the distal part of the severed axons (Supplementary Movies 1-3). We found, however, a significant increase in the size of the engulfed debris by macrophages in Erbb2-mutant animals (Figure 2C). These results reveal that the loss of Sarm1 does not affect focal damage resolution by macrophages. Also, importantly, they indicate that macrophages are recruited to the wound independently of the Schwann cells. We did notice discrete and highly mobile axon fragments that were not associated with EGFP(+) cells in wild-type and Erbb2-mutant animals, but not in Sarm1-deficient fish.

Synthetic degradation of severed Sarm1-deficient axons

Calcium (Ca^{2+}) regulation in neurons is critical for homeostasis because sustained elevation of cytosolic calcium leads to axonal and neuronal degeneration. Often, this occurs because mitochondria release apoptosis-inducing factors and proteases in a calcium-dependent manner (60). Studies of mammalian neurons *in vitro* and of zebrafish have shown that neuronal damage triggers two waves of elevation of axoplasmic calcium (Ca^{2+}) (21, 52). The first occurs nearly immediately after injury and decays rapidly, whereas the second has slower onset and decay. Loss of Sarm1 prevents Ca^{2+} elevation in severed axons (21). We sought to further test the functional conservation of Sarm1 in zebrafish *in vivo* by monitoring Ca^{2+} dynamics in lateralis sensory axons at high-resolution before and after injury. We measured Ca^{2+} dynamics *in toto* at high-resolution before and after axon injury using the genetically-encoded ultrasensitive Ca^{2+} sensor GCaMP7a, to find that intact wild-type and Sarm1-mutant axons show undetectable levels of axoplasmic fluorescence above background (Figure 3A). Upon severing, fluorescent signal in wild-type axons distal segments increased immediately and subsequently decayed with a near constant slope, whereas fluorescence remained nearly undetectable in Sarm1-deficient distal axon segments (Figure 3B). Next, we examined the Ca^{2+} levels in mitochondria and the endoplasmic reticulum (ER) using genetically encoded vital sensors, respectively Mito-RGECO and ER-GCaMP3 (Figure 3C-F). We selected these organelles because Ca^{2+} release from the axonal ER activates the mitochondrial permeability transition pore to trigger axonal degeneration (61, 62). We found that both mitochondrial and reticular Ca^{2+} levels increased equally after

axon severing in severed wild-type and Sarm1-/- axons. These results reveal that

loss of Sarm1 attenuates calcium influx to the axoplasm, but not Ca^{2+} uptake in

mitochondria or the ER.

Because the second Ca^{2+} wave is responsible for the activation of the serine-

threonine protease Calpain, which in turn facilitates axonal fragmentation by

cleaving microtubules and neurofilaments (63), we decided to monitor Ca^{2+} levels in

lateralis sensory axons 2, 4, 8 and 12 hours-post-injury (hpi). In severed wild-type

axons, the second wave of axoplasmic Ca^{2+} starts 4 hpi in coincidence with axon

fragmentation, and remains elevated in axonal debris up until 8 hpi (Figure 4A).

Note that the temporal resolution of these images does not allow the resolution of

degeneration before axonal regeneration. By contrast, the second wave of

axoplasmic Ca^{2+} does not occur in Sarm1-mutant axons (Figure 4B). Thus, we

hypothesized that forcing a sustained elevation of axoplasmic Ca^{2+} will be sufficient

to trigger the degradation of severed Sarm1-deficient axons. To test this prediction,

we transgenically expressed the rat transient receptor potential cation channel

subfamily V member 1 (TRPV1) fused to tagRFP in lateralis afferent neurons of

Sarm1-mutant zebrafish. Expression of mCherry alone in neurons served as control.

TRPV1 is non-selective cation channel that exhibits a high divalent selectivity, and

whose activation produces an influx of Ca^{2+} into cells (64). Rat TRPV1 is activated

by temperatures above 43°C or by the vanilloid capsaicin. Importantly, this

TRPV1 is inactive at the temperature used to maintain zebrafish (28°C), and

zebrafish TRPV1 orthologs are insensitive to capsaicin (65). Therefore, rat TRPV1

expressed in zebrafish offers a tunable tool to elevate axoplasmic Ca^{2+} with

excellent temporal resolution. We severed TRPV1-expressing and mCherry-

expressing lateralis axons, and 2 hours later a vehicle solution or vehicle + capsaicin

were added in the water holding the fish (Supplementary Fgure 2A-B). Samples

were inspected 90 minutes later. Severed Sarm1-deficient axons not expressing

TRPV1 did not fragment in presence of capsaicin, or TRPV1-expressing axons

bathed in ethanol solution. As hypothesized, we found that Sarm1-deficient

TRPV1-expressing axon segments readily degraded in the presence of capsaicin

(Supplementary Figure 2C-F). These results demonstrate for the first time *in vivo*

that elevation of axoplasmic Ca^{2+} downstream of Sarm1 is sufficient to trigger axon

degradation.

Schwann cells are not essential to maintain Sarm1-/- axons

The Schwann cells support, fasciculate and myelinate sensory axons in vertebrates

(66). We reasoned that delayed axon degeneration by loss of Sarm1 might impact

the interaction between axons and Schwann cells. To assess this interaction *in vivo*,

we combined the *sarm1hzm13* allele with the triple transgenic line

Tg[gSAGFF202A ; UAS-GFP ; SILL:mCherry] to highlight the Schwann cells

with green fluorescence and the lateralis afferent neurons with red fluorescence (53,

67). Using high-resolution intravital microscopy, we ascertained that Schwann cells

develop normally and fasciculate sensory axons in Sarm1-deficient zebrafish (Figure

5A). Upon severing, wild-type axons were quickly cleared by the Schwann cells

through engulfment of axonal fragments and intracellular degradation of debris

(Figure 5B and Supplementary Movie 4). Thus, we asked if Schwann cells are

necessary for the maintenance of severed Sarm1-deficient axons by generating a

double mutant zebrafish line concurrently deficient for Erbb2 and Sarm1. In

Erbb2-deficient specimens, the distal portion of the severed axons fragmented and

were cleared (Figure 6A-B,D) but with a significant delay compared to wild-type

specimens (Figure 6A,D). By contrast, in Erbb2/Sarm1 double mutants, severed

axons did not fragment or degrade, identically to fish lacking only Sarm1 (Figure

6C,D). Thus, axon maintenance in Sarm1 mutants occurs independently of the

Schwann cells.

When an injury generates a gap in the glia, wound-adjacent Schwann cells actively

move and extend cellular projections resembling filopodia to quickly reconstitute a

continuous glial scaffold (53). Although continuous glia are not necessary for the re-

growth of the proximal axon stump, it prevents regenerating growth cones from

straying, in turn facilitating end-organ *de novo* innervation and circuit

reconstitution (39, 41). Thus, we decided to interrogate Schwann-cell behavior

immediately after axon severing and during axon regeneration combining

Tg[gSAGFF202A;UAS-GFP ; SILL:mCherry] double transgene with the Sarm1

mutation. As expected, we found that in wild-type animals, the Schwann cells

adjacent to the wound quickly extended filopodia to close the gap in the glial

scaffold ahead of axonal regeneration (Supplementary Movie 5). Re-growing fibers

then followed these sub-cellular glial bridges to reconstitute the nerve, suffering

mild defasciculation restricted to a small area within the injury (Figure 6E and

Supplementary Figure 3). In stark contrast, injury-adjacent Schwann cells did not

migrate or produce filopodia-like projections in Sarm1 mutants (Supplementary

Movie 6). Nonetheless, regenerating axons eventually negotiated the persistent

larger gap to grow along the distal glial scaffold. However, the reforming nerves

presented more pronounced local defasciculation (Figure 6F and Supplementary

Figure 4).

Loss of Sarm1 does not alter Schwann-cell phenotype

Denervated Schwann cells undergo partial dedifferentiation from myelinating to a

progenitor-like state as revealed by loss of expression of terminal-phenotype

markers, including myelin and myelin-associated proteins (25, 68, 69). The loss of

terminal phenotype promotes Schwann-cell proliferation and migration, which

enhances their regenerative function (70). We hypothesized that because Schwann-

cell denervation does not occur in Sarm1-mutants, distal Schwann cells may not

dedifferentiate. This would explain their lack of phagocytic and protrusive activities

after axon transection. Following this rationale, we immunostained samples with an

antibody to Claudin-k, which localizes to the junctions between mature Schwann

cells and is downregulated in denervated glia (41, 71). In wild-type specimens,

Claudin-k remained strongly expressed along the entire length of the lateralis

afferent nerve up to 6 hours after nerve injury (hpi), suggesting that distal Schwann

cells remain mature during distal-axon fragmentation (Figure 7A-B). Beginning at

10 hpi, however, distal Schwann cells had markedly less Claudin-k than proximal

cells. Finally, 24 hpi Claudin-k was conspicuously absent from distal Schwann cells

even after axons regeneration had commenced. By contrast, Claudin-k remained strongly expressed after axon severing in Sarm1-mutant animals during the same period, with no apparent difference between Schwann cells located at either side of the wound (Figure 7A,C). Next, we assessed myelination using the 6D2 monoclonal antibody, which recognizes a carbohydrate epitope in the piscine P0-like myelin glycoproteins IP1 and IP2 (72, 73). As with Claudin-k, 6D2 labeling faded in Schwann cells distal to the injury in wild-type specimens (Figure 8A). Yet, 6D2 labeling remained unchanged in Sarm1 mutants (Figure 8B). We obtained congruent results when addressing myelination directly in living specimens by using a transgenic line expressing membrane-targeted EGFP under the control of the myelin binding protein (Mbp:EGFP-CAAX, green) (Figure 8C-D) (74). Interestingly, forced degradation of Sarm1-deficient severed axons expressing the rat TRPV1 channel after capsaicin treatment induced associated Schwann cells to dedifferentiate (Supplementary Figure 2G). Together, these results indicate that Sarm1-deficient sensory axons are maintained independently of Schwann-cell support, and that the clearance the severed axons is not necessary for regenerating axon growth, pathfinding, myelination and re-innervation of sensory organs. In addition, they reveal that Schwann cells distal to the injury do not de-differentiate in Sarm1-mutant specimens.

Loss of Sarm1 protects Schwann cells from chemical toxins

Many chemotherapeutic agents invariably cause peripheral neurotoxicity, leading to permanent neuronal dysfunction (75-79). Under conditions of denervation,

Schwann cells become strongly susceptible to chemotoxicity (80). We wondered if the protracted maintenance of severed axons in Sarm1 mutants may suppress glial vulnerability. To test this idea, we treated wild-type and Sarm1-mutant zebrafish with several chemical compounds that are under clinical trials or used as first-line treatment for common cancers in humans. First, we used 10-Hydroxycamptothecin (10-HCT), which is extremely toxic to denervated glia (82), and counted Schwann cells using the fluorescence transgenic marker *Tg[gSAGFF202A ; UAS:EGFP]*, which is ideal for this experiment because the intensity of green fluorescence does not vary in denervated Schwann cells and, therefore, is independent of the maturity of these glia (41). We confirmed that 10-HCT does not affects in Schwann cells associated to viable axon (Figure 9A-B). However, it significantly reduced the number of Schwann cells in 10-HCT-treated wild-type animals after axon severing. By contrast, the number of distal Schwann cells was only marginally affected in Sarm1-mutant specimens (Figure 9B). Platinum-based, taxanes and some alkaloids are effective chemotherapeutic agents used as standards-of-care for various human malignancies, despite their severe neuropathic effects that include glial destruction (81). To address their effect on Schwann cell, we treated wild-type or Sarm1 mutant zebrafish with cisplatin, oxaliplatin, paclitaxel, docetaxel and vincristine. We found that upon nerve transection, all these drugs invariably killed injury-distal Schwann cells in wild-type specimens but not in Sarm1 mutants (Figure 9C). Importantly, none of these drugs affected Schwann cells associated with intact axons, suggesting that axons protect Schwann cells from chemical stress. To confirm this prediction, we forced the degradation of severed axons in Sarm1-deficient zebrafish treated

with 10-HCT or vincristine. To this end, we repeated the used of capsaicin to activate rat TRPV1 expressed in a sub-set of lateralis neurons in homozygous mutant fish, in which Sarm1 is absent from every cell, including Schwann cells. This experiment revealed that Schwann cells lacking Sarm1 again become vulnerable to chemotoxicity once severed axons were synthetically eliminated (Figure 9D), confirming that Sarm1-mediated glioprotection is non-autonomous and depends upon the presence of non-degradable axons.

DISCUSSION

The sensory neurons that innervate skin, sensory receptors, joints and muscle communicate peripheral information to the brain, enabling animals to perform the essential activities of daily life. Using zebrafish and a battery of tests that include subcellular structural characterization of sensory neurons and associated Schwann cells, neuronal-function, and behavioral assays, we offer a comprehensive and integrated analysis of neuronal and glial response to injury, as well as on the consequences of blocking axon degeneration systemically. It has been well established that the absence of Sarm1 in **Drosophila** and the mouse prevents the degradation of damaged axons (15, 83). Although we were not predicting any differences for Sarm1 in zebrafish, assaying functional conservation in our experimental neuronal pathway is not dispensable because some neurotraumatic conditions lead to neuronal loss in the absence of Sarm1. Loss of Sarm1 improves functional recovery after traumatic brain injury in mice, and inhibits vincristine-mediated neurotoxicity (84, 85). Sarm1-deficient mice are viable (19, 86). We found

that chronic and systemic loss of Sarm1 is compatible with zebrafish viability and

sensorineural function, and demonstrate that the long-term maintenance of non-

degradable axon fragments has no detrimental effect on the repair of a sensory

circuit. Furthermore, regenerating axons fasciculate and myelinate normally,

indicating that they do not compete with non-degradable axons segments for

exiting glia. Therefore, we conclude that the axonal destructive and nerve

reconstructive processes occur in parallel. In addition, these data suggest that a

competitive balance between axon degradation and regeneration does not appear

to have shaped the evolution of Sarm1 and, by extension, Wallerian degeneration.

Focusing on Schwann cells, we show that glial cells adjacent to nerve injury in

Sarm1 mutants behave dramatically different than those of wild-type specimens.

Specifically, Schwann cells in Sarm1 mutants do not migrate or extend projections

to bridge the gap in the glial scaffold, indicating that these cells do not directly sense

missing intercellular contacts or glial discontinuity. Instead, our findings suggest

alternative scenarios. One is that the degradation of axons releases signals that

induce Schwann cell to change behavior. In mice, for example, nerve damage

promotes mesenchymal behavior of Schwann cells surrounding the wound via

TGF-β signaling, which drives collective Schwann-cell migration across the wound

(87). Although we did not observe Schwann-cell migration, signals derived from

injured axons may promote wound-adjacent Schwann cells to extend projections to

bridge the gap in a similar manner. Interestingly, we observed that filopodia-like

structures emerged from Schwann cells at both sides of the injury in wild-type

animals, suggesting that repair-inducing signals are likely diffusible, affecting glial

cells independently of their association with axons. It remains to be determined if

the source of such signals is the damaged axons, the denervated Schwann cells, or

other cells in the wound microenvironment.

Upon nerve injury, activated macrophages rapidly accumulate around the wound

and contribute to Wallerian degeneration and to axonal regeneration. Specifically,

classical pro-inflammatory M1-type phagocytic macrophages remove axonal and

myelin debris, whereas anti-inflammatory M2-type macrophages modulate

Schwann-cell activity and promote axon regeneration (96). Immediately after nerve

injury, Schwann cells release chemokines that attract or retain macrophages at the

wound before axon fragmentation. Importantly, pharmacological activation of

macrophage recruitment to the wound microenvironment enhances axon

regeneration, whereas decrease of macrophage infiltration inhibits axon

regeneration. Reactive oxygen species (ROS) are conserved early wound signals

across species. Injury-mediated ROS production recruits inflammatory cells,

including macrophages. Primary sources of ROS after nerve damage are the

mitochondria and cellular NADPH oxidases. Studies in *ex vivo* cultured neurons

indicate that Sarm1 acts downstream of mitochondrial ROS generation. Also,

previous studies have characterized Wallerian axon degeneration and regeneration

in Erbb2 mutant specimens, but did not address effects on the wound

microenvironment and the behavior of macrophages (93-94). Accordingly, tested *in*

vivo if the loss of Sarm1 as well as the absence of Schwann cells would affect

macrophage recruitment, to find that the onset of recruitment, the retention time

and number of macrophages at the wound did not differ between wild-type

specimens and Sarm1 or Erbb2 mutants. Also, as in wild-type specimens,

macrophages in Sarm1 and Erbb2 mutants engulfed debris locally, which were

larger within macrophages in Ebbr2 mutants. Thus, the loss of Sarm1 does not

affect focal damage resolution by macrophages, which are recruited and activated

independently of the Schwann cells. The above findings suggest that activated

phagocytic cells other than macrophages and Schwann cells, possibly neutrophils

(97), engulf and clear distal axon fragments after damage, and that this process does

not occur in Sarm1-deficient animals.

In conclusion, current mechanistic details about Sarm1 function derive from

independent observations extrapolated from a wide variety of experimental systems.

This has made it difficult to synthesize findings and reconcile some conflicting

observations. Here, we have exploited a powerful *in vivo* genetic system to

comprehensively study the consequence of systemic loss of Sarm1 from the sub-

cellular to the organismal level in a single vertebrate system. By generating

zebrafish carrying loss-of-function mutations in Sarm1, we confirmed and deepen

previous findings from ***Drosophila***, the mouse and cultured cells. Several novel

insights derive from the work presented here. First, that loss of Sarm1 is well

tolerated by the animal. Second, that neuronal-circuit repair is not contingent upon

rapid clearance of damaged axons. Third, that Schwann cells are not necessary for

maintenance of severed axons *in vivo*. Fourth, that after the axotomy Schwann

cells distal to the cut site don't dedifferentiate in Sarm1-mutants ***in vivo***. Fifth, that the protracted maintenance of transected axons dramatically improves Schwann-cell tolerance to chemotoxicity after nerve injury after neurotrauma. This scenario is likely to occur in the clinical setting because nerve trauma is an inescapable consequence of surgical interventions, chronic metabolic dysfunction including diabetes, and pharmacological treatments such as antibiotics and anæsthesia that increase cellular stress (81). Therefore, these findings are of obvious pathophysiological significance. Crucially, these data lend strong support to the idea that direct interventions to systemically inhibit axon degradation are promising strategies to reduce chronic consequences of neurotrauma. Because TIR domain dimerization is necessary and sufficient to degrade NAD+, it renders Sarm1 amenable to inhibition by small molecules, as it has been demonstrated for the TIR domain of TLR2 (95). Our findings encourage the development of Sarm1 inhibitors for therapeutic applications (18, 29).

METHODS

Zebrafish strains and husbandry

Zebrafish (**Danio rerio**) were maintained in a centralized facility in accordance to guidelines by the Ethical Committee of Animal Experimentation of the Helmholtz Zentrum München, the German Animal Welfare act Tierschutzgesetz §11, Abs. 1, Nr. 1, Haltungserlaubnis, to European Union animal welfare, and to protocols number Gz.:55.2-1-54-2532-202-2014 and Gz.:55.2-2532.Vet_02-17-187 from the "Regierung von Oberbayern", Germany. The transgenic lines Tg[UAS:EGFP], Tg[HGn39D] and Tg[SILL:mCherry] (67), Tg[gSAGFF202A] (41), Tg[UAS:GCaMP7a] (88), Tg[mbpa:tgRFP-CAAX]^tum102Tg (also known as Tg[MBP:tgRFP]) (89) and Erbb2 mutants (41, 59) have been previously published. The Tg[Sarm1-/-] was generated by CRISPR/Cas9 mediated mutagenesis.

Sarm1 mutagenesis

We used CRISPR/Cas9-mediated genome modification to generate mutations in exon 1 of Sarm1. Cas9 mRNA and sgRNAs were co-injected into one-cell stage embryos. Cas9 mRNA was generated **in vitro** from PmeI linearized CAS9 vector (pMLM3613) using the Ambion™ mMESSAGE mMACHINE T7 Kit. The mRNA was purified with the RNA easy kit (Qiagen). To generate the sgRNAs, the target exon was sequenced and the sequence information was used to design oligonucleotides for the sgRNA guide vector (pDR274) using the on-line tool "ZiFiT Targeter software package" (http://zifit.partners.org**/**) (90). The sgRNA sequence for exon 1 of Sarm1 is 5'-GGGACTTGGAAGAGACCCGC-3'. The annealed oligonucleotide was cloned into the BsaI-digested pDR274 vector using T4 ligase (NEB M0202). Resulting clones were sequenced to verify correctness, and then linearized with DraI. The purified linearized DNA fragment was employed to generate the sgRNA using the T7 MEGAscript kit (Ambion™). By out-crossing adult fish resulting from injection, we obtained germ-line transmission of two independent alleles: **sarm1^hzm13** and **sarm1^hzm14**. For genotyping mutant carriers, we used the primers: Forward: 5'-GATTTGCCGTTATCTCTCCA-3' and Reverse: 5'-TCAAGCAGTTTGGCAGACTC-3'.

DNA constructs

The DNA constructs SILL:mCherry, SILL:Gal4 (67), UAS:Synapsin1-GFP (53) and mbp:EGFP-CAAX have been previously described. The vectors pMLM3613 (42251) and pDR272 (42250) were purchased from Addgene. The plasmids UAS:TRPV1-tagRFP, the coding sequence of rat TRPV1 containing the E600K mutation fused to tagRFP was synthesized by Genecat and the expression construct was generated using Tol2 kit. The constructs SILL:mito-mCherry, SILL:Sarm1-v2a-mCherry, SILL:Kaede, SILL:mitoRGECO and SILL:erGCaMP3 were generated using Tol2 kit. Mitochondria in lateralis neurons were marked by expressing the mitochondria targeting sequence from the Cytochrome-C oxidase subunit 8A fused to mCherry. The plasmids containing mito-RGECO and er-GCaMP3 (92) were a gift of D.W. Raible (University of Washington).

Antibodies and immunostaining

Whole-mount immunostaining was performed as following. First, samples of zebrafish embryos or larvae were fixed by immersion in ice-cold 4% paraformaldehyde diluted into phosphate-buffered saline buffer containing 0.2% Tween20 (PBST), and incubated overnight at 4°C. Then, the samples were washed at room temperature (RT) with PBST three times, 10 minutes per wash, and subsequently blocked in 10% bovine serum albumin (BSA), also at RT for 1 hour. Next, the samples were incubated in primary antibodies at 4°C overnight. Next, the samples were washed in PBST for 2 hours, changing to fresh buffer every 30 minutes. Finally, they were incubated with secondary antibodies at 4°C overnight. Primary antibodies and concentrations: mouse anti-Acetylated tubulin, 1:1000 (Sigma T7251); rat anti-Claudin-k, 1:500 (gift from T. Becker, University of Edinburgh, U.K.) (71); mouse 6D2, 1:5 (gift from Dr. G. Jeserich, University of Osnabrück, Germany) (73). Secondary antibodies used were at the following concentrations: donkey anti-Mouse Alexa Fluor® 555, 1:200, Abcam ab150106; donkey anti-Rat IgG H&L (Alexa Fluor® 647) pre-adsorbed, 1:200, Abcam ab150155). Samples were washed in PBST for 30 minutes and mounted in

Vectashield one day before microscopic examination. Imaging of fixed samples was

done with a laser-scanning confocal microscope (LSM 510, Carl Zeiss).

Intravital microscopy

For videomicroscopy, larvae were anesthetized with MS-222 (0.013% M/V) in

Danieau's and mounted in 0.8% low melting-point agarose on 35mm glass-bottom

Petri dishes. Samples were gently pressed against the glass using a hair-loop glued

to the tip of a glass pipette, as previously described. The agarose dome was

immersed in Danieau's with MS-222. Images of cells were acquired using a

spinning-disc microscope with a 40× air objective at 28.5°C (58). Z-stacks were set

to 0.8-1.2 μm intervals. Time intervals were 10 minute or 15 minute per stack. The

resulting raw data were processed, assembled and analyzed with ImageJ.

Western blot assay

Wild-type and mutant larvae were anesthetized and killed 5 days-post-fertilization.

Samples were homogenized in ice-cold RIPA buffer (with protease inhibitor

cocktail from Roche (Cat.04693159001). After homogenization, the samples were

incubated on ice for 30 minutes for further lysis. The resulting lysate was

centrifuged at 1200 rpm at 4°C, and the supernatant was taken for the BCA assay.

The supernatant was diluted in loading buffer and boiled at 99°C for 5 minutes.

Next, the samples were run in SDS-PAGE and transferred onto a PVDF

membrane. After blocking the membrane in 5% skimmed milk (diluted in PBST)

for 1 hour, the membrane was incubated with the primary antibody (rabbit anti-

Sarm1, 1:500, ANASPEC 55381; Mouse anti-β-Tubulin, 1:2000, Sigma T5168) at

4°C overnight. The next day, the membrane was washed and incubated with HRP-

labeled secondary antibody (Peroxidase-Affini Pure Goat Anti-Mouse IgG (H+L),

1:10000, Jackson Immuno Research 115035003; Peroxidase-Affini Pure Goat Anti-

Rabbit IgG (H+L), 1:10000, Jackson ImmunoResearch 115035144) for 1 hour.

Images were acquired by developing the membrane with ECL (Pierce™ ECL

Western Blotting Substrate, Thermo Fisher, 32109).

Laser microsurgery

To mark lateralis sensory neurons individually, DNA of the SILL:mCherry construct was injected into eggs of Tg[HGn39D], Tg[HGn39D; Sarm1-/-], Tg[SILL:mCherry; gSAGFF202A; UAS:EGFP] or Tg[SILL:mCherry ; gSAGFF202A ; UAS:EGFP ; Sarm1-/-] zebrafish. Resulting larvae were selected according to red-fluorescence in lateralis neurons. Selected samples were mounted into agarose as described above, and their peripheral axons were targeted an ultraviolet laser (350nm) using the iLasPulse system (Roper scientific AS, Evry, France), as described previously. The laser beam was delivered using a 63x water-immersion objective (50, 53, 58). The laser pulses were calibrated and applied to the target area until a clear gap in the axons was visible. The samples were observed again 1 hour post-injury (hpi) to confirm complete axon transection.

Quantification of mitochondrial density and motility

To analyze mitochondria in sensory axons, we generated kymographs of mito-mCherry fluorescent spots using the Multi-Kymograph tool of the Fiji software (http://fiji.sc). The movement of mitochondria was determined by the slope of the lines drawn over time, and the direction of movement was determined by the moving mitochondria crossing the time line (vertical) in the kymographs. The data were analyzed with Python scripts and the Graphpad Prism software.

Calcium imaging

For calcium imaging in lateralis neurons, Tg[SILL:Gal4; UAS:GCaMP7a] double-transgenic larvae were anesthetized and mounted in 0.8% low melting-point agarose on a 35mm glass-bottom Petri dish. Imaging was acquired through a 63x water-immersion objective with an exposure time of 400 milliseconds. Laser-mediated axon transection was done after the 4^{th} imaging of the time series. Next, live videomicroscopy was done for 2 minutes at a frame rate of 400 milliseconds at 28.5°C. The raw data were analyzed with ImageJ. To quantify the calcium signal, the images were processed to ImageJ. The region of interest (ROI) was selected and measured the value with time point. GCaMP or RGECO intensity changes were calculated as follows: $\Delta F/F0 = (F-F0)/F0$, where F0 is the value of the fluorescent

signal before axons were transected, and F is the value of the fluorescent signal with

time point after axon severing (92).

Chemogenetics

For chemogenetic experiments, we co-injected the SILL:Gal4 with UAS:mCherry

or UAS:ratTRPV1-tagRFP into the Tg[UAS:GCaMP7a] or Tg[HGn39D; Sarm1-

/-]. The positive larva with SILL:Gal4; UAS:ratTRPV1-tagRFP; UAS:GCaMP7a

expression was used to activate TRPV1 channels in zebrafish by incubation in 5μM

capsaicin (Sigma, M2028) and subsequent live imaging for 1hour of the mounted

and anesthetized embryo. Images were acquired through a 63x water-immersion

objective with an exposure time of 400 milliseconds. For experiments with the

Sarm1 mutant larvae, the HGn39D with SILL:mCherry or ratTRPV1-tagRFP

positive animals were laser axotomized. The larvae were treated with 10 μM

capsaicin or ethanol (1:1000, v/v) 2h after transection. 1.5h after capsaicin

treatment, images were taken by spinning disc microscopy.

Behavioral assays

For the touch-mediated escape response, 2dpf embryos were gently dechorionated

and kept in Danieau's solution at 28°C for at least 1 hour. Embryos were place into

a flat uncovered Petri dish containing Danieau's and were recorded with a high-

speed camera (NX4 series, Imaging solution, GmbH). Video recording was

launched and a randomly chosen embryo was touched with a blunt glass needle

until it evoked a reaction. Recordings were done under white-light illumination

over 150 seconds at a rate of 200 frames per second (fps). The swimming

trajectories were obtained with 3D Particle Tracker plugin, ImageJ software. The

further quantification and statistics were using Python.

Statistics and Reproducibility

The Student's t test (two tailed), ANOVA test, T-test (Tukey test) and Wilcoxon

rank sum test were applied using Python scripts. Error bars in all figures are

standard errors of the mean (SEM).

REFERENCES

1. Ameh V, and Crane S. Nerve injury following shoulder dislocation: the emergency physician's perspective. *Eur J Emerg Med.* 2006;13(4):233-5.
2. Heneka MT, Carson MJ, El Khoury J, Landreth GE, et al. Neuroinflammation in Alzheimer's disease. *Lancet Neurol.* 2015;14(4):388-405.
3. Missios S, Bekelis K, and Spinner RJ. Traumatic peripheral nerve injuries in children: epidemiology and socioeconomics. *J Neurosurg Pediatr.* 2014;14:688-94.
4. Rowe RK, Ellis GI, Harrison JL, Bachstetter AD, et al. Diffuse traumatic brain injury induces prolonged immune dysregulation and potentiates hyperalgesia following a peripheral immune challenge. *Mol Pain.* 2016;12.
5. Tzekov R, Phifer J, Myers A, Mouzon B, and Crawford F. Inflammatory changes in optic nerve after closed-head repeated traumatic brain injury: Preliminary study. *Brain Inj.* 2016;30(12):1428-35.
6. Ziogas NK, and Koliatsos VE. Primary Traumatic Axonopathy in Mice Subjected to Impact Acceleration: A Reappraisal of Pathology and Mechanisms with High-Resolution Anatomical Methods. *J Neurosci.* 2018; 38:4031-47.
7. Roselli F, and Caroni P. From intrinsic firing properties to selective neuronal vulnerability in neurodegenerative diseases. *Neuron.* 2015;85(5):901-10.
8. Cattin AL, and Lloyd AC. The multicellular complexity of peripheral nerve regeneration. *Curr Opin Neurobiol.* 2016;39:38-46.
9. Parrinello S, Napoli I, Ribeiro S, Wingfield Digby P, Fedorova M, Parkinson DB, et al. EphB signaling directs peripheral nerve regeneration through Sox2-dependent Schwann cell sorting. *Cell.* 2010;143(1):145-55.
10. Ziegenfuss JS, Doherty J, and Freeman MR. Distinct molecular pathways mediate glial activation and engulfment of axonal debris after axotomy. *Nat Neurosci.* 2012;15(7):979-87.
11. Harrisingh MC, Pérez-Nadales E, Parkinson DB, Malcolm DS, Mudge AW, and Lloyd AC. The Ras/Raf/ERK signalling pathway drives Schwann cell dedifferentiation. *EMBO J.* 2004;23(15):3061-71.
12. Scherer SS. The biology and pathobiology of Schwann cells. *Curr Opin Neurol.* 1997;10(5):386-97.
13. Waller A. Experiments on the Section of the Glossopharyngeal and Hypoglossal Nerves of the Frog, and Observations of the Alterations Produced Thereby in the Structure of Their Primitive Fibres. *Philosophical Transactions of the Royal Society of London.* 1850;140:423-9
14. Gerdts J, Summers DW, Milbrandt J, and DiAntonio A. Axon Self-Destruction: New Links among SARM1, MAPKs, and NAD+ Metabolism. *Neuron.* 2016;89(3):449-60.
15. Osterloh JM, Yang J, Rooney TM, Fox AN, Adalbert R, Powell EH, et al. dSarm/Sarm1 is required for activation of an injury-induced axon death pathway. *Science.* 2012;337(6093):481-4.

16. Walker LJ, Summers DW, Sasaki Y, Brace EJ, Milbrandt J, and DiAntonio A.
MAPK signaling promotes axonal degeneration by speeding the turnover of the
axonal maintenance factor NMNAT2. *Elife.* 2017;6.
17. Yang J, Wu Z, Renier N, Simon DJ, et al. Pathological axonal death through a
MAPK cascade that triggers a local energy deficit. *Cell.* 2015;160:161-76.
18. Essuman K, Summers DW, Sasaki Y, Mao X, DiAntonio A, and Milbrandt J.
The SARM1 Toll/Interleukin-1 Receptor Domain Possesses Intrinsic NAD(+)
Cleavage Activity that Promotes Pathological Axonal Degeneration. *Neuron.*
2017;93(6):1334-43 e5.
19. Gerdts J, Summers DW, Sasaki Y, DiAntonio A, and Milbrandt J. Sarm1-
mediated axon degeneration requires both SAM and TIR interactions. *J
Neurosci.* 2013;33(33):13569-80.
20. Gerdts J, Brace EJ, Sasaki Y, DiAntonio A, and Milbrandt J. SARM1
activation triggers axon degeneration locally via NAD(+) destruction. *Science.*
2015;348(6233):453-7.
21. Loreto A, Di Stefano M, Gering M, and Conforti L. Wallerian Degeneration Is
Executed by an NMN-SARM1-Dependent Late Ca(2+) Influx but Only
Modestly Influenced by Mitochondria. *Cell Rep.* 2015;13(11):2539-52.
22. Summers DW, Gibson DA, DiAntonio A, and Milbrandt J. SARM1-specific
motifs in the TIR domain enable NAD+ loss and regulate injury-induced
SARM1 activation. *Proc Natl Acad Sci U S A.* 2016;113(41):E6271-E80.
23. Essuman K, Summers DW, Sasaki Y, Mao X, Yim AKY, DiAntonio A, et al.
TIR Domain Proteins Are an Ancient Family of NAD(+)-Consuming Enzymes.
Curr Biol. 2018;28(3):421-30 e4.
24. Sasaki Y, Nakagawa T, Mao X, DiAntonio A, and Milbrandt J. NMNAT1
inhibits axon degeneration via blockade of SARM1-mediated NAD(+)
depletion. *Elife.* 2016;5.
25. Jessen KR, Mirsky R, and Lloyd AC. Schwann Cells: Development and Role
in Nerve Repair. *Cold Spring Harb Perspect Biol.* 2015;7(7):a020487.
26. Jortner BS. Mechanisms of toxic injury in the peripheral nervous system:
neuropathologic considerations. *Toxicol Pathol.* 2000;28(1):54-69.
27. Morell P, and Toews AD. Schwann cells as targets for neurotoxicants.
Neurotoxicology. 1996;17(3-4):685-95.
28. Koeppen AH. Wallerian degeneration: history and clinical significance. *J
Neurol Sci.* 2004;220(1-2):115-7.
29. Simon DJ, and Watkins TA. Therapeutic opportunities and pitfalls in the
treatment of axon degeneration. *Curr Opin Neurol.* 2018;31(6):693-701.
30. Schuldiner O, and Yaron A. Mechanisms of developmental neurite pruning.
Cell Mol Life Sci. 2015;72(1):101-19.
31. Omura T, Omura K, Tedeschi A, Riva P, et al. Robust Axonal Regeneration
Occurs in the Injured CAST/Ei Mouse CNS. *Neuron.* 2015;86(5):1215-27.
32. D'Rozario M, Monk KR, and Petersen SC. Analysis of myelinated axon
formation in zebrafish. *Methods Cell Biol.* 2017;138:383-414.
33. Friedrich RW, Genoud C, and Wanner AA. Analyzing the structure and
function of neuronal circuits in zebrafish. *Front Neural Circuits.* 2013;7:71.

34.   Friedrich RW, Jacobson GA, and Zhu P. Circuit neuroscience in zebrafish.
677   ***Curr Biol***. 2010;20(8):R371-81.
678   35.   Ackerman SD, Garcia C, Piao X, Gutmann DH, and Monk KR. The adhesion
679   GPCR Gpr56 regulates oligodendrocyte development via interactions with
680   Galpha12/13 and RhoA. ***Nat Commun***. 2015;6:6122.
681   36.   Bin JM, and Lyons DA. Imaging Myelination In Vivo Using Transparent
682   Animal Models. ***Brain Plast***. 2016;2(1):3-29.
683   37.   Fontenas L, and Kucenas S. Livin' On The Edge: glia shape nervous system
684   transition zones. ***Curr Opin Neurobiol***. 2017;47:44-51.
685   38.   Ohki Y, Wenninger-Weinzierl A, Hruscha A, Asakawa K, Haass C, et al.
686   Glycine-alanine dipeptide repeat protein contributes to toxicity in a zebrafish
687   model of C9orf72 associated neurodegeneration. ***Mol Neurodegener***.
688   2017;12(1):6.
689   39.   Rosenberg AF, Isaacman-Beck J, Franzini-Armstrong C, and Granato M.
690   Schwann cells and deleted in colorectal carcinoma direct regenerating motor
691   axons towards their original path. ***J Neurosci***. 2014;34(44):14668-81.
692   40.   Wehner D, Tsarouchas TM, Michael A, Haase C, Weidinger G, Reimer MM,
693   et al. Wnt signaling controls pro-regenerative Collagen XII in functional spinal
694   cord regeneration in zebrafish. ***Nat Commun***. 2017;8(1):126.
695   41.   Xiao Y, Faucherre A, Pola-Morell L, Heddleston JM, Liu TL, Chew TL, et al.
696   High-resolution live imaging reveals axon-glia interactions during peripheral
697   nerve injury and repair in zebrafish. ***Dis Model Mech***. 2015;8(6):553-64.
698   42.   O'Donnell KC, Vargas ME, and Sagasti A. WldS and PGC-1alpha regulate
699   mitochondrial transport and oxidation state after axonal injury. ***J Neurosci***.
700   2013;33(37):14778-90.
701   43.   Conforti L, Gilley J, and Coleman MP. Wallerian degeneration: an emerging
702   axon death pathway linking injury and disease. ***Nat Rev Neurosci***.
703   2014;15:394.
704   44.   Howe K, Clark MD, Torroja CF, Torrance J, Berthelot C, Muffato M, et al.
705   The zebrafish reference genome sequence and its relationship to the human
706   genome. ***Nature***. 2013;496(7446):498-503.
707   45.   Sztal TE, Ruparelia AA, Williams C, and Bryson-Richardson RJ. Using
708   Touch-evoked Response and Locomotion Assays to Assess Muscle
709   Performance and Function in Zebrafish. ***J Vis Exp***. 2016(116).
710   46.   Ghysen A, and Dambly-Chaudière C. The lateral line microcosmos. ***Genes***
711   ***Dev***. 2007;21(17):2118-30.
712   47.   Pujol-Martí J, and López-Schier H. Developmental and architectural principles
713   of the lateral-line neural map. ***Front Neural Circuits***. 2013;7:47.
714   48.   Drerup CM, and Nechiporuk AV. In vivo analysis of axonal transport in
715   zebrafish. ***Methods Cell Biol***. 2016;131:311-29.
716   49.   Haehnel-Taguchi M, Akanyeti O, and Liao JC. Behavior, Electrophysiology,
717   and Robotics Experiments to Study Lateral Line Sensing in Fishes. ***Integr***
718   ***Comp Biol***. 2018;58(5):874-83.
719   50.   Pujol-Martí J, Faucherre A, Aziz-Bose R, Asgharsharghi A, Colombelli J,
720   Trapani JG, et al. Converging axons collectively initiate and maintain synaptic

721   selectivity in a constantly remodeling sensory organ. ***Curr. Biol.***
722   2014;24(24):2968-74.

51.   Olt J, Ordoobadi AJ, Marcotti W, and Trapani JG. Physiological recordings
from the zebrafish lateral line. ***Methods Cell Biol.*** 2016;133:253-79.

52.   Vargas ME, Yamagishi Y, Tessier-Lavigne M, and Sagasti A. Live Imaging of
Calcium Dynamics during Axon Degeneration Reveals Two Functionally
Distinct Phases of Calcium Influx. ***J Neurosci.*** 2015;35(45):15026-38.

53.   Xiao Y, Tian W, and Lopez-Schier H. Optogenetic stimulation of neuronal
repair. ***Curr Biol.*** 2015;25(22):R1068-9.

54.   Kim MJ, Kang KH, Kim CH, and Choi SY. Real-time imaging of
mitochondria in transgenic zebrafish expressing mitochondrially targeted GFP.
Biotechniques. 2008;45(3):331-4.

55.   Mandal A, Pinter K, and Drerup CM. Analyzing Neuronal Mitochondria ***in***
vivo Using Fluorescent Reporters in Zebrafish. ***Front Cell Dev Biol.***
2018;6:144.

56.   Lin MY, Cheng XT, Tammineni P, Xie Y, Zhou B, Cai Q, et al. Releasing
Syntaphilin Removes Stressed Mitochondria from Axons Independent of
Mitophagy under Pathophysiological Conditions. ***Neuron.*** 2017;94:595-610 e6.

57.   Miller KE, and Sheetz MP. Axonal mitochondrial transport and potential are
correlated. ***J Cell Sci.*** 2004;117(Pt 13):2791-804.

58.   Xiao Y, and López-Schier H. Studying Axonal Regeneration by Laser
Microsurgery and High-Resolution Videomicroscopy. ***Methods Mol Biol.***
2016;1451:271-80.

59.   Lyons DA, Pogoda HM, Voas MG, Woods IG, Diamond B, Nix R, et al.
erbb3 and erbb2 are essential for schwann cell migration and myelination in
zebrafish. ***Curr Biol.*** 2005;15(6):513-24.

60.   Ozaki T, Yamashita T, and Ishiguro S. Mitochondrial m-calpain plays a role in
the release of truncated apoptosis-inducing factor from the mitochondria.
Biochim Biophys Acta. 2009;1793(12):1848-59.

61.   Court FA, and Coleman MP. Mitochondria as a central sensor for axonal
degenerative stimuli. ***Trends Neurosci.*** 2012;35(6):364-72.

62.   Villegas R, Martinez NW, Lillo J, Pihan P, Hernandez D, Twiss JL, et al.
Calcium release from intra-axonal endoplasmic reticulum leads to axon
degeneration through mitochondrial dysfunction. ***J Neurosci.*** 2014;34:7179.

63.   Park JY, Jang SY, Shin YK, Suh DJ, and Park HT. Calcium-dependent
proteasome activation is required for axonal neurofilament degradation.
Neural Regen Res. 2013;8(36):3401-9.

64.   Caterina MJ, Schumacher MA, Tominaga M, Rosen TA, Levine JD, and
Julius D. The capsaicin receptor: a heat-activated ion channel in the pain
pathway. ***Nature.*** 1997;389(6653):816-24.

65.   Gau P, Poon J, Ufret-Vincenty C, Snelson CD, Gordon SE, Raible DW, et al.
The zebrafish ortholog of TRPV1 is required for heat-induced locomotion. ***J***
Neurosci. 2013;33(12):5249-60.

66.   Brosius LA, and Barres BA. Contrasting the glial response to axon injury in the
central and peripheral nervous systems. ***Dev. Cell.*** 2014;28(1):7-17.

67.   Faucherre A, Pujol-Martí J, Kawakami K, and López-Schier H. Afferent
neurons of the zebrafish lateral line are strict selectors of hair-cell orientation.
PLoS One. 2009;4(2):e4477.
68.   Sasaki Y, Hackett AR, Kim S, Strickland A, and Milbrandt J. Dysregulation of
NAD(+) Metabolism Induces a Schwann Cell Dedifferentiation Program. *J
Neurosci.* 2018;38(29):6546-62.
69.   Soto J, and Monje PV. Axon contact-driven Schwann cell dedifferentiation.
Glia. 2017;65(6):864-82.
70.   Carr MJ, and Johnston AP. Schwann cells as drivers of tissue repair and
regeneration. *Curr Opin Neurobiol.* 2017;47:52-7.
71.   Munzel EJ, Schaefer K, Obirei B, Kremmer E, Burton EA, Kuscha V, et al.
Claudin k is specifically expressed in cells that form myelin during development
of the nervous system and regeneration of the optic nerve in adult zebrafish.
Glia. 2012;60(2):253-70.
72.   López-Schier H, and Hudspeth AJ. Supernumerary neuromasts in the
posterior lateral line of zebrafish lacking peripheral glia. *Proc Natl Acad Sci
U.S.A.* 2005;102(5):1496-501.
73.   Bastmeyer M, Jeserich G, and Stuermer CA. Similarities and differences
between fish oligodendrocytes and Schwann cells in vitro. *Glia.* 1994;11:300.
74.   Czopka T, Ffrench-Constant C, and Lyons DA. Individual oligodendrocytes
have only a few hours in which to generate new myelin sheaths in vivo. *Dev
Cell.* 2013;25(6):599-609.
75.   Cavaletti G, Alberti P, and Marmiroli P. Chemotherapy-induced peripheral
neurotoxicity in the era of pharmacogenomics. *Lancet Oncol.* 2011;12:1151.
76.   Kaiser J, Bledowski C, and Dietrich J. Neural correlates of chemotherapy-
related cognitive impairment. *Cortex.* 2014;54:33-50.
77.   McLeary F, Davis A, Rudrawar S, Perkins A, and Anoopkumar-Dukie S.
Mechanisms underlying select chemotherapeutic-agent-induced
neuroinflammation and subsequent neurodegeneration. *Eur J Pharmacol.*
2018;842:49-56.
78.   Fukuda Y, Li Y, and Segal RA. A Mechanistic Understanding of Axon
Degeneration in Chemotherapy-Induced Peripheral Neuropathy. *Front
Neurosci.* 2017;11:481.
79.   Gornstein E, and Schwarz TL. The paradox of paclitaxel neurotoxicity:
Mechanisms and unanswered questions. *Neuropharmacology.* 2014;76:175.
80.   Imai S, Koyanagi M, Azimi Z, Nakazato Y, Matsumoto M, Ogihara T, et al.
Taxanes and platinum derivatives impair Schwann cells via distinct
mechanisms. *Sci Rep.* 2017;7(1):5947.
81.   Yang S, Abrahams MS, Hurn PD, Grafe MR, Kirsch JR. Local anesthetic
Schwann cell toxicity is time and concentration dependent. *Reg Anesth Pain
Med.* 2011;36:444-51.
82.   Bremer J, Skinner J, and Granato M. A small molecule screen identifies *in
vivo* modulators of peripheral nerve regeneration in zebrafish. *PLoS One.*
2017;12(6):e0178854.

83.   Gilley J, Ribchester RR, and Coleman MP. Sarm1 Deletion, but Not Wld(S),
Confers Lifelong Rescue in a Mouse Model of Severe Axonopathy. ***Cell Rep.***
2017;21:10-6.
84.   Henninger N, Bouley J, Sikoglu EM, An J, Moore CM, King JA, et al.
Attenuated traumatic axonal injury and improved functional outcome after
traumatic brain injury in mice lacking Sarm1. ***Brain.*** 2016;139(Pt 4):1094-105.
85.   Geisler S, Doan RA, Strickland A, Huang X, Milbrandt J, and DiAntonio A.
Prevention of vincristine-induced peripheral neuropathy by genetic deletion of
SARM1 in mice. ***Brain.*** 2016;139(Pt 12):3092-108.
86.   Turkiew E, Falconer D, Reed N, and Hoke A. Deletion of Sarm1 gene is
neuroprotective in two models of peripheral neuropathy. ***J Peripher Nerv***
Syst. 2017;22(3):162-71.
87.   Clements MP, Byrne E, Camarillo Guerrero LF, Cattin AL, Zakka L, Ashraf A,
et al. The Wound Microenvironment Reprograms Schwann Cells to Invasive
Mesenchymal-like Cells to Drive Peripheral Nerve Regeneration. ***Neuron.***
2017;96(1):98-114 e7.
88.   Muto A, Ohkura M, Abe G, Nakai J, and Kawakami K. Real-time
visualization of neuronal activity during perception. ***Curr Biol.*** 2013;23:307.
89.   Auer F, Vagionitis S, and Czopka T. Evidence for Myelin Sheath Remodeling
in the CNS Revealed by ***In Vivo*** Imaging. ***Curr Biol.*** 2018;28(4):549-59 e3.
90.   Wright DA, Thibodeau-Beganny S, Sander JD, Winfrey RJ, Hirsh AS,
Eichtinger M, et al. Standardized reagents and protocols for engineering zinc
finger nucleases by modular assembly. ***Nat Protoc.*** 2006;1(3):1637-52.
91.   Czopka T, and Lyons DA. Individual oligodendrocytes have only a few hours
in which to generate new myelin sheaths ***in vivo***. ***Dev. Cell.*** 2013;25(6):599.
92.   Esterberg R, Linbo T, Pickett SB, Wu P, Ou HC, Rubel EW, et al.
Mitochondrial calcium uptake underlies ROS generation during
aminoglycoside-induced hair cell death. ***J Clin Invest.*** 2016;126(9):3556-66.
93.   Atanasoski, S., Scherer, S. S., Sirkowski, E., Leone, D., Garratt, A. N.,
Birchmeier, C., & Suter, U. ErbB2 signaling in Schwann cells is mostly
dispensable for maintenance of myelinated peripheral nerves and proliferation
of adult Schwann cells after injury. ***J. Neurosci.*** 2006; 26: 2124-31.
94.   Han, S. B., Kim, H., Lee, H., Grove, M., Smith, G. M., & Son, Y. J. Postinjury
induction of activated ErbB2 selectively hyperactivates denervated Schwann
cells and promotes robust dorsal root axon regeneration. ***J. Neurosci.*** 2017;
37: 10955–10970.
95.   Mistry P, Laird MH, Schwarz RS, Greene S, Dyson T, Snyder GA, et al.
Inhibition of TLR2 signaling by small molecule inhibitors targeting a pocket
within the TLR2 TIR domain. ***Proc.Natl.Acad.Sci.U.S.A.*** 2015; 112:5455-60.
96.   Kigerl KA, Gensel JC, Ankeny DP, Alexander JK, Donnelly DJ, Popovich PG.
Identification of two distinct macrophage subsets with divergent effects causing
either neurotoxicity or regeneration in the injured mouse spinal cord. ***J.***
Neurosci. 2009; 29(43):13435-44.

97.   Lindborg JA, Mack M, Zigmond RE. Neutrophils Are Critical for Myelin
Removal in a Peripheral Nerve Injury Model of Wallerian Degeneration. ***J.***
Neurosci. 2017; 37(43):10258-10277

FIGURE LEGENDS

Figure 1. a) Structure Sarm1 indicating alignment of the Sarm1 functional

domains from different species (not at scale). **b)** Confocal image of axonal

mitochondria marked with mito-mCherry in wild-type and Sarm1-/-. Red arrows

point to prominent mitochondrial groups in axons. **c)** Upper panels, kymographs

from videomicroscopic recording of axonal mitochondria in wild type (H) (left

panel) and Sarm1-/- (I) (right panel). Lower panels show color-coded traces of

moving mitochondria in anterograde (green) and retrograde (red) directions, taken

from the kymographs shown in the upper panels. **d)** Density of mitochondria in

5dpf wild type and Sarm1-/-, Error bar = SEM; n.s. = not significant, p value from

Student's t test, n=25 (WT), n=19 (Sarm1-/-). **e)** Mobility of the mitochondria in

5dpf wild type and Sarm1-/-. Circles show the anterograde and triangles the

retrograde movement of the mitochondria. p value from one-way ANOVA, wild

type n=26, Sarm1-/- n=26. **f)** Time-lapse images of axonal degeneration of GFP-

labeled lateralis sensory neuron in wild-type (left) and Sarm1-/- larvae (right). hpi =

hour post injury, scale bar=50μm, white asterisk indicates the regrowing axons

from the proximal stump. **g)** Quantification of the time from axon transection to

fragmentation in wild type (n=13) and Sarm1-/- (n=13). **h)** Shows a recue

experiment in which the expression of a functional Sarm1 in sensory neurons of

Sarm1-mutant fish (right hand side panels) suffices to degrade severed axons,

similarly to wild-type fish (left hand side panels). Middle panels show non-

degradable Sarm-1 deficient severed axons.

Figure 2. a) Quantification of the number of macrophages recruited to the injury site and adjacent axon segments. **b)** Quantification of the time macrophages interact with axon segments. **c)** Quantification of the size of debris within macrophages. One-way ANOVA was conducted firstly, then P values for T-test in between two individual group.

Figure 3. a) Confocal image of a single lateralis sensory axon expressing the green-fluorescent calcium sensor GCaMP7a in wild type (left column) and Sarm1-/- fish (right column). Rows show that same samples before laser-mediated transection (top), immediately after transection (middle) and 60 seconds after transection (bottom). In A, C and E, white arrowheads indicate the position of the axon, specifically when signal-to-background is low. Scale bar 20μm. **b)** Shows quantification of the first wave of axoplasmic calcium. Data are shown as mean ± SEM; p from one-way ANOVA, wild type n = 16, Sarm1-/- 16. **c)** Shows a confocal image of lateralis sensory axons expressing the red-fluorescent calcium sensor RGECO in wild type (left column) and Sarm1-/- fish (right column). Rows show that same samples before laser-mediated transection (top), immediately after transection (middle), and 60 seconds after transection (bottom). **d)** Quantification mitochondrial calcium influx shows the strong and nearly identical elevation and decay in wild type and Sarm1-/- immediately after the cuts. Data are shown as mean ± SEM; p from one-way ANOVA, wild type n = 16, Sarm1-/- 16. **e)** Shows a confocal image of lateralis sensory axons expressing the green-fluorescent calcium sensor CCaMP3 targeted to the endoplasmic reticulum (ER) in wild type (left column) and Sarm1-/- fish (right column). Rows show that same samples before laser-mediated transection (top), immediately after transection (middle), and 60 seconds after transection (bottom). **f)** Quantification ER calcium influx shows strong and statistically equal elevation and decay in wild type and Sarm1-/- after the cuts. Data are shown as mean ± SEM; p from one-way ANOVA, wild type n = 16, Sarm1-/- 16.

Figure 4. a) Confocal image of a lateralis sensory axons expressing the mCherry (red) and green-fluorescent calcium sensor GCaMP7a (green) in wild type fish. Rows show same axons 2, 4, 8 and 12 hours after transection (hours-post-injury = hpi). **b)** Confocal image of lateralis sensory axons expressing the mCherry (red) and green-fluorescent calcium sensor GCaMP7a (green) in Sarm1-/- fish. Scale bar 20µm.

Figure 5. a) Confocal images of a double transgenic 5dpf larva showing Schwann cells marked by expression of GFP (green) under the control of the Tg[gSAGFF202A] Gal4 driver, and lateralis afferent neurons marked by expression of mCherry under the control of the SILL enhancer (red). Wild type (top), Sarm1 mutants (bottom). Scale bar 20µm. **b)** Images show the indicated time points after axon transection (hours post injury = hpi) from a videomicroscopic recording of Schwann cells (green) and their interaction with axons (red) in wild-type and Sarm1-/-. White arrowheads indicate Schwann cells engulfing axonal debris in the wild type. A white arrow indicates degradation-resistant axon segment in Sarm1-/-. Please, note that the proximal axon stump in Sarm1-/- is not visible in these images because it is outside the focal plane.

Figure 6. a-c) Images of mCherry-expressing (red) transected axons in wild type (A), Erbb2-/- mutants (B), and Sarm1-/- ; ErBb2-/- double mutants (C). Scale bar 100µm. **d)** Quantification of transected axon fragmentation in Erbb2-/- and Sarm1-/-; ErBb2-/-. Error bar = SEM, p value from One-way ANOVA test, n = 15 (each group). **e)** Image of a from Supplemental Movie 1, showing the discrete local defasciculation of regenerated the sensory fiber (red) and the bridging of the glial gap by Schwann cells (green) in a wild-type specimen. **f)** Equivalent experiment, taken from Supplemental Movie 2, showing a more pronounced local defasciculation of the regenerated the sensory fiber (red) in a Sarm1-mutant specimen. Note that and the bridging of the glial gap does not occur.

Figure 7. a-c) Schematic model of the confocal imaging locations on severed axons (A). Confocal images of wild type (B) and Sarm1-/- (C) specimens in 6hpi, 10hpi and 24hpi. hpi = hour post injury. The specimens were the Tg[gSAGFF202A;UAS:EGFP; SILL:mCherry] lines and were stained with Claudin-k (magenta) antibody. Scale bar 50μm.

Figure 8. a-b) Confocal images of wild type (A) and Sarm1-/- (B) specimens expressing EGFP in sensory neurons of the lateral line (green) and stained with the monoclonal antibody 6D2. Stainings were performed at indicated time points after axons severing (hpi). The arrows point to the cutting sites. Scale bar is 50μm. **c)** Live imaging of the *Tg[Mbp-EGFP; SILL:mCherry]* after severing. The arrows indicated the fragmented axons and the arrowheads the fragmented myelin. Scale bar 20μm. **d)** Live imaging of the Sarm1-/- in *Tg[Mbp-EGFP; SILL:mCherry]* after severing. Arrows indicate regrowing axons, and arrowheads indicate the juxtaposition between the regrowing axons. Scale bar 20μm.

Figure 9. a) Confocal images showing Schwann cells (green) and lateralis sensory axons (red) in a control specimen (in which axons were not transected), in a specimen 48 hours after axon transection, and in specimens treated with 10-HCT (10-Hydroxycamptothecin). Left column is wild type and right column shows Sarm1-/-. In all cases, the concentration of 10-HCT in water was 40μm. Scale bar 100μm. **b)** Quantification of the Schwann cells from A). Data are shown as mean ± SEM. ** means $p < 0.01$, two-way ANOVA, n = 8 (each group), followed by T-test for two individual group. **c)** Quantification of Schwann cells of WT, WT severed, Sarm1-/- and Sarm1-/- severed with the treatment of the indicated chemical compounds for 48 hours. Concentrations: Paclitaxel 40μm, Docetaxel 0.1μm, Oxaliplatin 500μm, Cisplatin 50μm, Vincristine 50μm. Data are shown as mean ± SEM. * means $p < 0.05$; ** means $p < 0.01$, three-way ANOVA, n = 8 (each group), followed by T-test for two individual group. **d)** Quantification of Schwann cells after axon severing, in specimens treated with 10-HCT or Vincristine. Left bar

974   group is wild type. Right bar group is Sarm1-/-, and Sarm1-/- with synthetically

975   eliminated axon segments. Two-way ANOVA, followed by T-test for two

976   individual groups.

a

0h

6h

24h

48h

b erbb2-/-

0h

6h

24h

48h

c Sarm1-/- + erbb2-/-

0h

6h

24h

48h

d Fragmentation time / h

P<0.001

> 5 d   > 5 d

WT   Sarm1-/-   erbb2-/-   Sarm1-/-; erbb2-/-

e wild type

Axons Glia

f Sarm1-/-

Axons Glia

a

6 hpi	10 hpi	24 hpi

b

Glia

Axons

Cldn-K

Merge

wild type

C

Glia

Axons

Cldn-K

Merge

Sarm1-/-