1 Title: Man against machine: Do fungal fruitbodies and eDNA give similar
2 biodiversity assessments across broad environmental gradients?
3
4 Tobias Guldberg Frøslev^{*1,2}, Rasmus Kjølner¹, Hans Henrik Bruun¹, Rasmus
5 Ejrnæs⁴, Anders Johannes Hansen², Thomas Læssøe³, Jacob Heilmann-
6 Clausen³

7
8 *1) Department of Biology, University of Copenhagen, Universitetsparken 15,*
9 *DK-2100 Copenhagen Ø, Denmark, Denmark.*

10 *2) Natural History Museum of Denmark, University of Copenhagen, Øster*
11 *Voldgade 5-7, DK-1350 Copenhagen K, Denmark.*

12 *3) Department of Biology /Natural History Museum of Denmark,*
13 *Universitetsparken 15, DK-2100 Copenhagen Ø, Denmark, Denmark.*

14 *4) Department of Bioscience, University of Aarhus, Grenåvej 14, DK-8410*
15 *Rønde, Denmark*

16 *Corresponding author: tobiasgf@snm.ku.dk

18 **Abstract**

19 The majority of Earths biodiversity is unknown. This is particularly true for the
20 vast part of soil biodiversity, which rarely can be observed directly.

21 Metabarcoding of DNA extracted from the environment (eDNA) has become
22 state-of-the-art in assessing soil biodiversity. Also for fungal community
23 profiling eDNA is seen as an attractive alternative to classical surveying based
24 on fruitbodies. However, it is unknown whether eDNA-metabarcoding
25 provides a representative sample of fungal diversity and census of threatened

species. Therefore conservation planning and assessment are still based on fruitbody inventories. Based on a dataset of unprecedented width and depth, representing both soil eDNA-metabarcoding and expert inventorying of fungal fruitbodies, we document for the first time the validity of eDNA as practical inventory method and measure of conservation value for fungi. Fruitbody data identified fewer species in total and per site, and had larger variance in site richness. Focusing on macrofungi – the class Agaricomycetes, and in turn the order Agaricales – metrics of total richness and compositional similarity converged between the methods. eDNA was suboptimal for recording the non-soil dwelling fungi. β -diversity was similar between methods, but more variation in community composition could be explained by environmental predictors in eDNA data. The fruitbody survey was slightly better in finding red-listed species. We find a better correspondence between biodiversity indices derived from fungal fruitbodies and DNA-based approaches than indicated in earlier studies. We argue that (historical) fungal community data based on fruitbody forays – with careful selection of taxonomic groups – may be interpreted together with modern DNA-based approaches.

Keywords: fungal fruitbodies, environmental DNA, metabarcoding, species richness, community composition, red-listed species, environmental gradients

1. Introduction

1.1 methods for inventorying fungi

For decades, inventory and identification of fungal fruitbodies were – together with isolation and culturing – the only way to assess fungal communities

(Hueck, 1953; Lange, 1948; Kjølner and Struwe, 1980; Rayner and Todd, 1980; Tyler, 1985; Schmit and Lodge, 2005). Since the 1990s, these methods have been supplemented with DNA-based methods, e.g. sequencing root samples to identify mycorrhizal fungi (Gardes and Bruns, 1996; Helgason *et al.*, 1998) or sequencing of cloned PCR products from soil/litter samples (Schadt *et al.*, 2003; O'Brien *et al.*, 2005; Taylor *et al.* 2014) – methods that allow for a more targeted study of some compartments, but are difficult to apply to ecosystem-wide inventories of large sampling sites. More recently, massive parallel sequencing of environmental DNA (eDNA) – now known as eDNA-metabarcoding (Taberlet *et al.*, 2012) – has gained ground in studies of fungal communities (e.g. Schmidt *et al.*, 2013; Pellissier *et al.*, 2014; Tedersoo *et al.*, 2014; Barnes *et al.*, 2016), and allows for such wide inventories. . In this study we compare a thorough fruitbody inventory with eDNA-metabarcoding for ecosystem-wide inventorying of the fungal community. Fruitbody surveys and eDNA-based methods both have their strengths and limitations and may be seen as complementary, rather than competing approaches (Truong *et al.*, 2017).

1.2 Fruitbody inventorying

Fruitbody surveys are low tech but laborious, requiring life-long expert taxonomic skills if thorough and reproducible data are to be achieved (Newton *et al.*, 2003). However, many fungi do not produce fruitbodies and are systematically omitted. Other taxa are likely to be under-sampled, as they are rarely fruiting, or produce very small, inconspicuous, short-lived or below-ground fruitbodies (Taylor and Finlay, 2003; Löhmus 2009; van der Linde *et*

al., 2012). Fruitbody formation and duration are highly influenced by (local variations in) season and weather conditions, which may hamper comparisons of sites, unless sampling is repeated over several years (Newton *et al.*, 2003; O'Dell *et al.*, 2004).

1.3 eDNA-metabarcoding

eDNA-metabarcoding is low tech when it comes to field sampling, but require high tech lab facilities and advanced post sequencing bioinformatics. It provides a broader taxonomic sample of the fungal community of not only the sexually reproducing, fruitbody forming fungi. Sampling of soil eDNA is less dependent on seasonality and climatic variation. Also, the majority of fungal biodiversity has yet to be described (Hibbett *et al.*, 2011) and a large proportion of available barcode references lack proper annotation (Hibbett *et al.*, 2011; Hibbett *et al.*, 2016; Nilsson *et al.*, 2016; Yahr *et al.*, 2016). This limits ecological interpretation of detected community differences in relation to guild structure, trait space or taxonomic composition. Furthermore, when sampling for eDNA only a tiny fraction of a particular site surface area can be sampled, even with an intensive design. Hence, the sample representativity depends on the heterogeneity of species distributions within habitats and the size of mycelia (Lilleskov *et al.*, 2004) – factors not easily assessed. This is a potential caveat, especially for detection of rare species – e.g. red-listed taxa – important for nature conservation (van der Linde *et al.*, 2012). Although eDNA-metabarcoding has been shown to successfully identify red-listed species (Geml *et al.*, 2014; van der Linde *et al.*, 2012), these may be more easily detected as fruitbodies, which may be targeted by trained experts over

large study areas in relatively short time, particularly for species with long-lived fruitbodies (e.g. perennial polypores; Runnel *et al.*, 2015).

1.4 Fruitbodies versus eDNA-metabarcoding

Several studies detect a limited overlap in communities between fruitbody surveys and DNA-based approaches on a habitat scale (Gardes and Bruns, 1996; Dahlberg *et al.*, 1997; Jonsson *et al.*, 1999; Porter *et al.*, 2008; Geml *et al.*, 2009; Fischer *et al.*, 2012; Baptista *et al.*, 2015). However, it remains an open question whether key community metrics nonetheless correlate along environmental gradients, so that results from either method can be used as proxy for the other. In the context of nature conservation and monitoring, it would be attractive if eDNA-metabarcoding can be proven to detect target species (e.g. red-listed species) – for which e.g. historical data of decline is known, or where monitoring programs are already running – independent of optimal fruiting seasons and availability of taxonomic expertise. Finally, it would be valuable if historical data based on fruitbody surveys hold valid information on fungal communities, and may be compared or combined with modern eDNA-metabarcoding for inferring of temporal change.

1.5 Approach and expectations

In this study, we compare richness, community composition and community-environment relation in a large ecological space using two parallel data sets, a thorough fruitbody inventory and data obtained by eDNA-metabarcoding of soil. All data were gathered from the same 130 40x40 m sample plots in Denmark and taken over the same 2-3 year period.

Overall, we expected eDNA-metabarcoding to detect more species than fruitbody sampling. However, we expected the fruitbody survey to detect more red-listed species, due to the targeted survey across the whole study area and the fact that most red-listed species produce conspicuous fruitbodies. We expected eDNA-metabarcoding to provide stronger correlation with environmental gradients, due to the expected better coverage of taxonomic diversity. We expected comparability between the two approaches to be highest for community composition, and lowest for red-listed species detection, as previous studies have indicated stochastic variation in noisy data to affect richness estimates more than community composition (Abrego *et al.*, 2016; Lekberg *et al.*, 2014). Finally, we expected higher correspondence between fruitbody and eDNA-metabarcoding data, when the former were restricted to species recorded at soil level, and when both were restricted to Agaricomycetes and Agaricales.

2. Materials and methods

2.1 Study sites

130 sites of 40 × 40 m spread out across Denmark were studied. The study sites covered an ecospace spanning the major environmental gradients of terrestrial ecosystems, i.e. soil moisture, soil fertility and successional stage (Brunbjerg *et al.* 2017b). The 130 sites were selected by stratified random sampling to represent 24 environmental strata (habitat types). Six habitat types were cultivated: three types of fields (rotational, grass leys, set aside) and three types of forest plantations (beech, oak, spruce). The remaining 18

strata were natural habitat types, constituting all factorial combinations of:
fertile and infertile; dry, moist and wet; open, tall herb/scrub and forest. We
replicated these 24 strata in each of five geographical regions across
Denmark. We further included a subset of 10 perceived biodiversity hotspots,
two within each region. This study was part of the Danish biodiversity study,
Biowide, and an elaborate description of design and data collection is
available in Brunbjerg et al. (2017a).

2.2 Fruitbody survey

Each site was visited twice during the main fungal fruiting season in 2014
(August - early September and October - early November) and once during
the main fruiting season in 2015 (late August - October), focussing on all
groups of Basidiomycota and Ascomycota, but excluding non-stromatic
Pyrenomycetes and Discomycetes with fruitbodies regularly smaller than 1
mm. Most woody debris was turned over to locate e.g. corticioid fungi, but no
structured attempts to find hypogeous fungi were conducted, although a few
were found by chance. In sites with tall and dense herbaceous vegetation,
regular inspections were carried out in kneeling position. A site visit lasted
approximately 1 hour, in very open monotonous sites sometimes less, e.g. in
newly ploughed arable sites. All visits were led by expert field mycologist
Thomas Læssøe typically accompanied by one helper. Numerous samples
were taken back to a mobile lab for immediate microscopic investigation, and
more interesting or critical material was dried as voucher material and in part
deposited at the fungal herbarium (C) of the National History Museum of

Denmark. Some specimens difficult to identify were forwarded to external experts.

2.3 Environmental variables

A complete inventory of vascular plants was done for each site. Ellenberg Indicator Values, EIV (Ellenberg *et al.*, 1991) reflect plant species' abiotic optimum and have often been used in vegetation studies to describe local conditions (Diekmann, 2003). Mean Ellenberg Indicator Values were calculated based on the plant lists for each site for the light conditions (EIV.L), soil nutrient status (EIV.N) and soil moisture (EIV.F). Ellenberg values together with measured variables (see supplementary methods) for precipitation, soil pH, soil organic matter content, soil carbon content, soil phosphorous and light were used in the models to explain community structure.

2.4 Sequence data

Soil was collected from all sites followed by DNA-extraction and sequencing with primers targeting fungi as described elsewhere (Brunbjerg *et al.*, 2017a). For each site, 81 soil samples (each sample was approximately 5 cm diam. and 15 cm depth) were collected in a virtual grid with samples 4 m apart using a simple gardening tool. For each site, a large bulk soil sample was constructed by thorough mixing of 81 single soil samples, and subjected to DNA extraction with the MoBio PowerMax kit. The fungal ITS2 region was amplified using primers gITS7 (Ihrmark *et al.*, 2012) and ITS4 (White *et al.*, 1990). Libraries were MiSeq sequenced (Illumina Inc., San Diego, CA, USA),

at the Danish National Sequencing Centre using two 250 bp PE runs. OTU tables (species-site table) were constructed, aiming for a definition of OTUs (Operational Taxonomic Units) that approximates species level delimitation. This was achieved by an initial processing with DADA2 (Callahan *et al.*, 2016) to identify exact amplicon sequence variants including removal of chimeras, followed by ITS extraction with ITSx (Bengtsson-Palme *et al.*, 2013) and subsequent clustering with VSEARCH (Rognes *et al.*, 2016) at 98.5 % – the consensus clustering level used to delimit species hypotheses (SHs) in the UNITE database (Kõljalg *et al.*, 2014), and subsequent post-clustering curation using LULU (Frøslev *et al.*, 2017) to eliminate remaining redundant sequences. Taxonomic assignment of the OTUs was done using the 2017 UNITE general FASTA release (<http://dx.doi.org/10.15156/BIO/587475>). Sequence data is available from DataDryad, and files documenting the analyses from GitHub [[Data will be made available before publication. Until then it will be available by contacting the first author]].

For the more descriptive analyses, we used full fruitbody data. For some more direct comparisons, we restricted the fruitbody data to species collected at the soil surface for a more qualified comparison, as it was evident that only a small proportion of the non-soil fungi were registered by the soil-based eDNA-metabarcoding. Furthermore both datasets were filtered to obtain two increasingly taxonomically focussed subsets – Agaricomycetes and Agaricales. eDNA-metabarcoding and fruitbody data were then assessed for correspondence in a set of biodiversity metrics. Species composition was the

focus of the study, and as biological abundance is difficult to assess with either method, presence/absence data was used for all analyses.

2.5 Overlap between methods

The frequency of each species/OTU across the 130 sites was assessed for the full datasets, and the proportion of species/OTUs recorded with both methods or only as fruitbody or OTU was assessed. As incomplete and insufficiently annotated DNA reference data exacerbate the discrepancies between fruitbody and metabarcoding data, some focussed analyses were performed on only the species recorded with both methods ('coinciding species').

2.6 Species richness and sampling effort

OTU richness was not greatly influenced by sequencing depth, Spearman rank $r = 0.98$ between OTU count based on rarefied data (10,000 reads per sample) and full data, and OTU richness measures were thus estimated from the full (not rarefied) data. Species accumulation was assessed for all datasets, and the variation of the recorded site richness per site was assessed by calculation of the relative standard deviation of richness for each method. Pearson correlation was used to test for correspondence between estimates of species richness and OTU richness across the 130 sites.

2.7 Red-listed species

As a measure of conservation value we used the count of red listed fungal species in the IUCN categories from near threatened to critically endangered

on the official Danish red list (IUCN 2012, Wind & Pihl 2010). We assessed both the total number of red-listed species identified with either method, as well as the correspondence of site wise counts of red-listed species.

2.9 Community composition

Community dissimilarity was estimated with the Sørensen dissimilarity metric using the `vegdist` function in `vegan`. Five out of the 130 sites had less than 4 observed fruitbody species and were removed prior to analyses of community dissimilarity. Correlation between community dissimilarity measures based on different datasets were tested with the Mantel test (method = "pearson", 999 permutations) and Procrustes test (999 permutations) using the functions in `vegan`. Community turnover along gradients (assessed as dissimilarity) was tested for correlation with environmental distance using the `bioenv` function in `vegan`. Here Sørensen distance is used for community dissimilarity and Euclidean distance for environmental dissimilarity, and we allowed up to four explanatory variables to be selected.

3 Results

3.1 Overlap between methods

The fruitbody survey recorded fewer species than the eDNA-metabarcoding approach (Fig. 1a, Supplementary Fig. 1). The fruitbody survey included 8,793 observations (a species in a site), and recorded 1,751 species (1,358 Agaricomycetes of which 847 belonged to Agaricales). The eDNA-metabarcoding included total 30,668 observations (an OTU in a site), and

275 recorded 8,110 OTUs (2,521 Agaricomycetes and 1,293 Agaricales). 1,288
 276 (74 %) of the fruitbody species were recorded as fruitbodies only, while 463
 277 (26 %) were found also as OTUs. 7517 (93 %) of the OTUs were found with
 278 eDNA-metabarcoding only, while 593 (7 %) were recorded also as fruitbodies
 279 (i.e. had species name annotations corresponding to the 463 species
 280 mentioned above). For these 463 coinciding species there was a tendency
 281 towards pairwise correspondence of species and OTU frequency (Fig. 1a and
 282 1c). Four coinciding species were common as fruitbodies, but rare as OTUs.
 283 Three of these (*Mycena speirea*, *Clitopilus hobsonii*, *Mollisia cinerea*) are
 284 normally observed on woody or herbaceous substrates, and the last (*Galerina*
 285 *vittiformis*) is associated with bryophytes. The top ten most frequent coinciding
 286 OTUs were less frequent as fruitbodies – all common soil fungi, except
 287 *Ganoderma applanatum*, a wood decomposer not generally perceived as a
 288 soil fungus. Community dissimilarity estimates based on the 463 coinciding
 289 species resulted in corresponding composition estimates (mantel r-statistic
 290 0.63, and 0.83 correlation in a symmetric procrustes rotation). Site OTU
 291 richness and fruitbody species richness was highly correlated when
 292 considering only the 463 coinciding species, $r = 0.81$ with a slope close to 1
 293 (Fig. 1d).
 294 394 (37%) of the 1,067 soil fruitbody species were also registered as OTUs,
 295 whereas only 69 (10%) of the 684 non-soil fruitbody species were also
 296 registered as OTUs. Per site, an average of 6.3% of the soil fruitbody species
 297 were captured as OTUs, but only 0.18% of the non-soil fruitbody species. As it
 298 was evident that the soil eDNA captured little of the non-soil fungi, we made

comparison analyses of site richness and red-list recording both using the full fruitbody data but also on fruitbody data excluded the non-soil species.

3.2 Overall species richness and sampling effort

Although the exact logging of expenses was not part of the project, we estimate that the costs of the two approaches would be approximately equal, if repeated with the focussed aim of monitoring. The fruitbody survey included 3 x 10,000 km driving, and four months of salary (three months of collecting, one month of identification) – excluding the aid from volunteers in the fruitbody survey, whereas the eDNA-metabarcoding included 10,000 km driving, approximately 6,000 USD lab consumables, and 3 months salary (1 month collecting, 2 months lab work and bioinformatics). Species accumulation curves did not reach an asymptote for any of the datasets after sampling of the 130 sites (Fig. 2a). This was most pronounced for the full eDNA-metabarcoding dataset and least for the non-soil Agaricales fruitbody dataset. eDNA-metabarcoding became increasingly similar to fruitbody data with narrowed taxonomic focus. The variation in site species/OTU richness across the 130 sites was lowest for eDNA-metabarcoding, and highest for fruitbody data (markedly higher for non-soil fungi), but more similar with a narrowed taxonomic focus (Fig. 2b).

3.3 Richness correlation between methods.

Soil fruitbody species richness of the 130 study sites (Fig. 3) ranged from 0-115 (0-111 Agaricomycetes and 0-79 Agaricales), while OTU richness ranged from 66-476 (11-157 Agaricomycetes, 6-87 for Agaricales). Correlation

between site species richness and OTU richness (Fig. 3) was moderate for the full datasets ($r = 0.43$), but strong when restricted to Agaricomycetes ($r = 0.64$) and Agaricales ($r = 0.58$). Correlations became even stronger ($r = 0.74 - 0.81$) when only considering the 463 'coinciding species' – species recorded with both methods (Fig. 1d, Supplementary Fig. 1d and 1h). OTU richness based on only Agaricomycetes or Agaricales were strongly correlated with OTU richness based on the full data and ($r = 0.74$ and $r = 0.8$, respectively, Supplementary Fig. 2).

3.4 Overall taxonomic composition

Taxonomic composition of eDNA-metabarcoding and fruitbody data became increasingly similar when going from full data to Agaricomycetes and Agaricales (Fig. 4, Supplementary Fig. 3, Supplementary Tables 1-4). Fruitbody data was heavily skewed towards Basidiomycota (90 %), whereas the eDNA-metabarcoding was composed of 39 % Ascomycota, 40 % Basidiomycota, and 20 % species from other phyla (Fig. 4a, Supplementary table 2). However, the relative proportions and absolute frequencies of taxa progressively converged when focussing on Agaricomycetes (Fig. 4b, Supplementary Fig. 3c) and Agaricales (Fig. 4c, Supplementary Fig. 3d). The non-soil fruitbody data was less dominated by Agaricomycetes and Agaricales than the soil-fungi data. All phyla and classes (except Dacrymycetes and Atractiellomycetes) were represented by more species/OTUs in the eDNA-metabarcoding than in the fruitbody data (Fig. 4, Supplementary Fig. 3ab). A few Agaricomycetes orders (Polyporales, Hymenochaetales, Auriculariales, Gomphales) were represented by more species in the fruitbody data than in

the eDNA-metabarcoding (Supplementary Fig. 3c). Almost all Agaricales genera were detected by both methods and with roughly similar species numbers.

3.5 Red-listed species

The soil surface fruitbody survey recorded more red-listed species than the eDNA-metabarcoding (Fig. 5, Supplementary Table 5). 100 red-listed species were recorded as fruitbodies on the soil surface (144 including the non-soil fungi), whereas 63 red-listed species were found as OTUs. 26 red-listed species were recorded with both methods, 37 red-listed species was detected as OTUs only, and 74 as fruitbodies only. Only one red-listed species from the non-soil part of the fruitbody data (*Ganoderma pfeifferi*) was also detected as OTU. When restricting the comparison of red-listed species to species present in the molecular reference database the figures for soil fruitbodies (71 species) was almost equal to the figures for OTUs (Fig. 5). When the 130 sites were grouped into categories with 0, 1-2 or 3 or more red-listed species recorded as fruitbodies or as OTUs, there was a good correspondence between the two methods (Supplementary Fig. 4).

3.6 Community – environment relation

Mantel tests showed very similar and strong correlations between community dissimilarity measures from fruitbody and eDNA-metabarcoding data, mantel $r = 0.67$. In this case no improvements were achieved by increased taxonomic overlap (Mantel- $r = 0.68$ and 0.62 for Agaricomycetes and Agaricales respectively), or when restricted to soil-fungi (Mantel- $r = 0.68$, 0.71 and 0.64 for full data, Agaricomycetes and Agaricales respectively). These correlations

were corroborated by procrustes analyses with correlation coefficients of 0.87, 0.87 and 0.83, and 0.87, 0.87, 0.83 for soil-fungi for the same comparisons (all p-values < 0.01). Environmental variables explained more of the community dissimilarity for eDNA-metabarcoding data than for the survey data, and the amount of explained variation was largest for the taxonomically more inclusive datasets (Fig. 6) with the maximum explained variation for the full eDNA-metabarcoding dataset (0.66) and the lowest for the soil-fruitbody Agaricales (0.50). Adding a fourth explanatory variable did not increase the amount of explained variation for most datasets. Based on all subsets of the fruitbody data, the best three explanatory variables for community composition were mean Ellenberg soil nutrient status (EIV.N), mean Ellenberg light indicator value (EIV.L) and soil phosphorous, whereas mean Ellenberg soil nutrient status (EIV.N), mean Ellenberg soil moisture (EIV.F) and soil pH were the best for the DNA metabarcoding.

4. Discussion

More species (OTUs) were detected by eDNA-metabarcoding than by the classic fruitbody survey. This could mainly be attributed to the detection of groups, which always go undetected in a fruitbody survey, e.g. diverse groups of moulds and yeasts. The fruitbody survey data was strongly dominated by fruitbody-forming basidiomycetes. In general, there was a relatively poor correlation for richness measures and taxonomic composition between the two full datasets, but increased strength of correlation when narrowing the focus to Agaricomycetes and subsequently to just Agaricales. Similarly, excluding wood-inhabiting and other non-soil fungi improved the

correspondence between the datasets, showing that these largely go undetected in soil-based eDNA sampling. The fruitbody survey identified more red-listed species, but the difference was less pronounced than anticipated, and results were almost similar when delimited to soil-dwelling fungi only.

4.1 Taxonomic composition similar for macrofungi

The taxonomic composition was remarkably similar between eDNA-metabarcoding and fruitbody data when focussing on the Agaricomycetes, and even more pronouncedly the Agaricales. Many of the major discrepancies align with expectations – i.e. taxonomically difficult groups like *Inocybe* (Larsson et al 2009, Ryberg et al. 2008) and *Cortinarius* (Frøslev et al. 2007) were markedly more species rich as eDNA OTUs than as well-delimited species identified from fruitbodies. Approximately half of the Agaricales species recorded as fruitbodies were also found as OTUs and vice versa (Supplementary Fig. 1e). Considering the very similar proportions of Agaricales genera between the methods, it can be assumed that a large part of non-overlapping species can be explained by incomplete DNA reference data and different taxonomic concepts in handbooks for species identification of fruitbodies compared to sequence databases. An effort to expand and curate DNA reference databases is hence essential to improve future DNA-based ecological studies as already suggested by other researchers (Hibbett *et al.*, 2011; Hibbett *et al.*, 2016; Nilsson *et al.*, 2016; Yahr *et al.*, 2016).

4.2 Soil DNA captures soil-fungi

In this study, we extracted DNA from soil samples. Although DNA from non-soil fungi may be expected to be present in the soil, it has not earlier been tested to what extend soil DNA can be used to register fungi not having their active growing life-stages within the soil, such as wood decomposing fungi. Several non-soil fungi were detected in this study, but they were observed in much fewer sites than soil-fungi when comparing to the corresponding fruitbody data. In fact, the few higher taxa that were more speciose in the fruitbody data, were primarily non-soil taxa like *Polyporales*, *Hymenochaetales*, *Crepidotus*, etc. It was, however, interesting to note that *Ganoderma applanatum* (a wood decaying polypore) was found as OTU in 51 of 130 sites, including several sites with no trees, suggesting the species to be abundantly present in the spore bank. Although soil sampling catches fungi associated with above ground carbon sources, our results indicate that these are heavily undersampled. Studies indicate, that this part compose a major proportion of the total funga (Unterseher *et al.*, 2011; Arnold and Lutzoni, 2007; Arnold, 2007), so to get a more complete estimate of the total fungal community, DNA-based methods will need to include sampling of above ground structures.

4.3 Detection of red-listed species

The fruitbody survey registered more red-listed species in total and average per site. However, when adjusting for red-listed species not present in the DNA reference database (476 of the 656 Danish red-listed species were present with sequence data in UNITE), and omitting red-listed species associated with dead wood and other non-soil resources, the eDNA-

metabarcoding approach performed almost as good. However, the methods partly recorded different red-listed species, indicating that fruitbody surveys and eDNA-metabarcoding could be used complementarily to get a more reliable assessment of local conservation value, which to some degree conflicts with the findings of Runnel et al. (2015) that found fruitbody surveys to be superior to eDNA-based sampling of redlisted wood-inhabiting polypores at stand scale. The detection of red-listed species from environmental DNA samples must be expected to increase as sequence databases become more complete and well-annotated.

4.4 Species turnover comparable

Our results showed that community composition estimated from DNA-metabarcoding data correlated well with estimates based on fruitbody data. This correlation did not change much after narrowing taxonomic focus to Agaricomycetes/Agaricales, indicating that all approaches are suitable for describing fungal communities and species turnover along environmental gradients. However, eDNA-metabarcoding outperformed fruitbody data when it came to correlation with environmental gradients expressed by independent environmental variables for all subsets of data. Further, it appears that the wider soil fungal community is more predictable than the fruitbody community. This could be caused by fruitbodies constituting a more stochastic subset of the total fungi, or alternatively that Agaricomycetes and Agaricales depend on less easily measured properties of the environment. Most species of Agaricomycetes and Agaricales produce billions of spores that are effectively dispersed (Peay & Bruns 2014) – which is not the case for some of the other

main groups of fungi in this study (Money 2016). Hence the detected community of these by eDNA may in part be a signal from the spore bank. The spore bank community has been shown to have relatively low correlation with the active community of the same taxa for pine associated ectomycorrhizal fungi (Glassman et al 2015), and thus, the lower correlation seen in our study may potentially be caused by a similar discrepancy. The lower performance of fruitbody survey data likely also indicates that fruitbody formation is more sensitive to e.g. unpredictable variation in weather conditions.

As seen from the taxonomic composition the eDNA-metabarcoding has a much higher proportion of Ascomycota and other phyla of ‘micro-fungi’, but also a relatively lower proportion of non-soil Ascomycota and Basidiomycota. DNA-metabarcoding thus targets a community with a larger proportion of micro-fungi (possibly also due to PCR amplification biases), which must be assumed to be more dependent on soil composition and humidity, whereas the fruitbody data targets a community of macrofungi with a larger dependence on the vegetation and above ground conditions. This is reflected in light being among the best explanatory variables for the fruitbody data, and soil moisture and pH for the eDNA-metabarcoding data.

4.5 Sampling efficiency/depth.

The results obtained in this study reflect the exact sampling protocols for both fruitbody survey and eDNA sampling, as well as the bioinformatics processing of the sequence data. The fruitbody survey included three visits to each site, and it is obvious that more sampling visits continuously will add to

the species list, and may be necessary to get a fully representative sample (Halme and Kotiaho, 2012; Newton *et al.*, 2003; Straatsma *et al.*, 2015) (but see Abrego *et al.* 2016). The (eDNA) soil sampling method included the mixing of 81 soil cores and thus several kilos of soil for each site, and was uniquely large compared to previous studies (e.g. Porter *et al.*, 2008; Geml *et al.*, 2010; Baptista *et al.*, 2015; Pellissier *et al.*, 2014; Geml *et al.*, 2009; Schmidt *et al.*, 2013). However, it still covered only 0.01 % of the soil surface of the 40 x 40 m sites, and of the approximately 5-20 kg soil sampled from each plot only 4 g of soil was used for DNA extraction. Also, we made no attempt to maximize coverage of visible variation at the sites but sampled completely systematically. Hence, both sampling approaches could be both up- and down-scaled for applications in practice. A study in Switzerland (Straatsma *et al.*, 2015) recorded fruitbodies on a weekly basis over 21 years, and identified 101 species on average per year (408 species in total) in a forest study area close to ours in size (1,500 vs 1,600 m²). Although, their total number exceeds the site average of 68 fruitbody species (and 236 OTUs) in our study, their yearly average of 101 is only slightly higher than the average (93.5) of our forest/plantation sites after three 1 h visits, and we predict that it would require much further effort to get a significantly larger average species number for the fruitbody data. Presently, there is little knowledge on which parameters are most important for getting a representative sample with the eDNA-metabarcoding approach. We expect that extracting and sequencing the 81 soil cores separately or sequencing many sub-samples of the bulk sample would increase the number of detected OTUs, but this would also pose a marked increase of lab consumables and

processing time. As many fungal mycelia must be assumed to be restricted in size and/or time, and as our bulk sample only covers 0.01 % of the soil surface, we expect that additional bulk samples – done at the same time or at another time of the year – would be the most cost-efficient way of capturing a larger sample of the real fungal community.

4.6 Practical applications

Both approaches represent surveys that are realistic to perform within the limits of standard surveys and research studies, and expenses were roughly comparable. For the full data, eDNA-metabarcoding resulted in more species (OTU) observations (30,668) than the fruitbody survey (8,793), whereas the numbers were more similar for Agaricomycetes (9,091 vs. 7,214) and Agaricales (4,507 vs. 4,334). In our data, fruitbody richness was a relatively weak predictor of total fungal richness as assessed with DNA metabarcoding, but was a relatively good predictor of the richness of fruitbody forming fungi (Agaricomycetes and Agaricales). This indicates that species richness of these groups may be assessed interchangeably with eDNA-metabarcoding or as fruitbodies.

For detection of red-listed species, eDNA-metabarcoding performed much better than expected, but still we would recommend a manual search for fruitbodies in all cases where larger areas need to be surveyed e.g. for conservation value, as also suggested for wood-inhabiting fungi (Runnel *et al.*, 2015). However, eDNA-metabarcoding may supply valuable information, in cases of poor fruiting conditions, or in more targeted plot-based monitoring programmes. Fruitbody and eDNA-metabarcoding data result in comparable

measures of species turnover, and thus, our results indicate that data may be combined for example to evaluate long time series including historical fruitbody data and future DNA-based surveys.

Acknowledgements

Karl-Henrik Larsson, Leif Örstadius and Viacheslav Spirin are thanked for identification of critical fungal collections, Anne Storgaard, Annegrete Eriksen, David Boertmann, Erik Arnfred Thomsen, Hanne Petra Katballe, Jette Anitha Hansen, Leif Wegge Laursen, Thomas Kehlet, Tom Smidth, Torbjørn Borgen are thanked for assistance in the field during fruitbody inventories. JHC acknowledge the Danish National Research Foundation for funding for the Center for Macroecology, Evolution and Climate, grant no. DNRF96.

Conflicts of interest

All authors declare no conflicting interests.

Authors' contributions

All authors designed the study, TL collected the fruitbody data, TGF collected and analyzed the molecular data, TGF carried out the statistical analyses, TGF, JHC and RK wrote first draft, all authors contributed to revising the paper.

Data Accessibility

Sequence data is available from DataDryad, and files documenting the analyses from GitHub. Fruitbody data is deposited in GBIF. [[more information will be added before publication]].

References

- Abrego N, Halme P, Purhonen J, Ovaskainen O. (2016). Fruit body based inventories in wood-inhabiting fungi: Should we replicate in space or time? *Fungal Ecol* **20**: 225–232.
- Arnold AE. (2007). Understanding the diversity of foliar endophytic fungi: progress, challenges, and frontiers. *Fungal Biol Rev* **21**: 51–66.
- Arnold AE, Lutzoni F. (2007). Diversity and host range of foliar fungal endophytes: are tropical leaves biodiversity hotspots? *Ecology* **88**: 541–549.
- Baptista P, Reis F, Pereira E, Tavares RM, Santos PM, Richard F, *et al.* (2015). Soil DNA pyrosequencing and fruitbody surveys reveal contrasting diversity for various fungal ecological guilds in chestnut orchards. *Environ Microbiol Rep* **7**: 946–954.
- Barnes CJ, Maldonado C, Frøslev TG, Antonelli A, Rønsted N. (2016). Unexpectedly high beta-diversity of root-associated fungal communities in the Bolivian Andes. *Front Microbiol* **7**: 1–13.
- Bengtsson-Palme J, Ryberg M, Hartmann M, Branco S, Wang Z, Godhe A, *et al.* (2013). Improved software detection and extraction of ITS1 and ITS2 from ribosomal ITS sequences of fungi and other eukaryotes for analysis of environmental sequencing data. *Methods Ecol Evol* **4**: 914–919.
- Brunbjerg AK, Bruun HH, Brøndum L, Classen AT, Fog K, Frøslev TG, *et al.*

596 (2017b). A systematic survey of regional multitaxon biodiversity:
 597 evaluating strategies and coverage. *bioRxiv*. e-pub ahead of print, doi:
 598 <http://dx.doi.org/10.1101/158030>.

599 Brunbjerg AK, Bruun HH, Moeslund JE, Sadler JP, Svenning JC, Ejrnæs R.
 600 (2017a). Ecospace: A unified framework for understanding variation in
 601 terrestrial biodiversity. *Basic Appl Ecol* **18**: 86-94.

602 Callahan BJ, McMurdie PJ, Holmes SP. (2017). Exact sequence variants
 603 should replace operational taxonomic units in marker-gene data analysis.
 604 *ISME J* **11**: 2639–2643.

605 Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP.
 606 (2016). DADA2: High-resolution sample inference from Illumina amplicon
 607 data. *Nat Methods* **13**: 581–583.

608 Dickie IA. (2010). Insidious effects of sequencing errors on perceived diversity
 609 in molecular surveys. *New Phytol* **188**: 916–918.

610 Diekmann M. (2003). Species indicator values as an important tool in applied
 611 plant ecology - a review. *Basic Appl Ecol* **4**: 493–506.

612 Ellenberg H, Weber HE, Düll R, Wirth V, Werner W, Paulißen D. (1991).
 613 Zeigerwerte von Pflanzen in Mitteleuropa. *Scr Geobot* **18**: 1–248.

614 Fischer AL, Moncalvo J-M, Klironomos JN, Malcolm JR. (2012). Fruiting body
 615 and molecular rDNA sampling of fungi in woody debris from logged and
 616 unlogged boreal forests in northeastern Ontario. *Écoscience* **19**: 374–
 617 390.

618 Frøslev TG, Jeppesen TS, Læssøe T, Kjøller R (2007). Molecular
 619 phylogenetics and delimitation of species in *Cortinarius* section
 620 *Calochroi* (Basidiomycota, Agaricales) in Europe. *Molecular*

621 phylogenetics and evolution, 44(1), 217-227.

622 Frøslev TG, Kjøller R, Bruun HH, Ejrnæs R, Brunbjerg AK, Pietroni C, et al.

623 (2017). Algorithm for post-clustering curation of DNA amplicon data

624 yields reliable biodiversity estimates. *Nat Commun* **8**(1): 1188.

625

626 Gardes M, Bruns TD. (1996). Community structure of ectomycorrhizal fungi in

627 a *Pinus muricata* forest: above- and below-ground views. *Can J Bot* **74**:

628 1572–1583.

629 Geml J, Gravendeel B, Van Der Gaag KJ, Neilen M, Lammers Y, Raes N, et

630 al. (2014). The contribution of DNA metabarcoding to fungal

631 conservation: Diversity assessment, habitat partitioning and mapping

632 red-listed fungi in protected coastal *Salix repens* communities in the

633 Netherlands. *PLoS One* **9**(6): e99852.

634 Geml J, Laursen GA, Herriott IC, McFarland JM, Booth MG, Lennon N, et al.

635 (2010). Phylogenetic and ecological analyses of soil and sporocarp DNA

636 sequences reveal high diversity and strong habitat partitioning in the

637 boreal ectomycorrhizal genus *Russula* (Russulales; Basidiomycota).

638 *New Phytol* **187**: 494–507.

639 Geml J, Laursen GA, Timling I, McFarland JM, Booth MG, Lennon N, et al.

640 (2009). Molecular phylogenetic biodiversity assessment of arctic and

641 boreal ectomycorrhizal *Lactarius* Pers. (Russulales; Basidiomycota) in

642 Alaska, based on soil and sporocarp DNA. *Mol Ecol* **18**: 2213–2227.

643 Glassman SI, Peay KG, Talbot JM, Smith DP, Chung JA, Taylor JW, ... &

644 Bruns TD (2015). A continental view of pine-associated ectomycorrhizal

645 fungal spore banks: a quiescent functional guild with a strong

646 biogeographic pattern. *New Phytologist*, 205, 1619-1631.

647 Halme P, Kotiaho JS. (2012). The importance of timing and number of
648 surveys in fungal biodiversity research. *Biodivers Conserv* **21**: 205–219.

649 Helgason T, Daniell TJ, Husband R, Fitter AH, Young JPW. (1998). Ploughing
650 up the wood-wide web? *Nature* **394**: 431.

651 Hibbett DS, Ohman A, Glotzer D, Nuhn M, Kirk P, Nilsson RH. (2011).
652 Progress in molecular and morphological taxon discovery in Fungi and
653 options for formal classification of environmental sequences. *Fungal Biol*
654 *Rev* **25**: 38–47.

655 Hibbett D, Abarenkov K, Kõljalg U, Öpik M, Chai B, Cole J, *et al.* (2016).
656 Sequence-based classification and identification of Fungi. *Mycologia*
657 **108**: 1049–1068.

658 Hueck HJ. (1953). Myco-sociological methods of investigation. *Plant Ecol* **4**:
659 84–101.

660 Ihrmark K, Bödeker ITM, Cruz-Martinez K, Friberg H, Kubartova A, Schenck J,
661 *et al.* (2012). New primers to amplify the fungal ITS2 region - evaluation
662 by 454-sequencing of artificial and natural communities. *FEMS Microbiol*
663 *Ecol* **82**: 666–677.

664 IUCN. (2012). IUCN Red List Categories and Criteria: Version 3.1. Second
665 edition. Gland, Switzerland and Cambridge, UK: IUCN. iv + 32pp.

666 Jonsson T, Kokalj S, Finlay R, Erland S. (1999). Ectomycorrhizal community
667 structure in a limed spruce forest. *Mycol Res* **103**: 501–508.

668 Kjøller A, Struwe S. (1980). Microfungi of decomposing red alder leaves and
669 their substrate utilization. *Soil Biol Biochem* **12**: 425–431.

670 Kõljalg U, Nilsson RH, Abarenkov K, Tedersoo L, Taylor AFS, Bahram M.

671 (2014). Towards a unified paradigm for sequence-based identification of
672 fungi. *Mol Ecol* **22**: 5271–5277.

673 Lange M. (1948). The agarics of Maglemose: a study in the ecology of the
674 agarics. *Dansk Bot Arkiv* **13**(1): 1-141.

675 Larsson E, Ryberg M, Moreau PA, Mathiesen ÅD, Jacobsson S. (2009).
676 Taxonomy and evolutionary relationships within species of section
677 Rimosae (Inocybe) based on ITS, LSU and mtSSU sequence data.
678 *Persoonia*, 23, 86.

679 Lekberg Y, Gibbons SM, Rosendahl S. (2014). Will different OTU delineation
680 methods change interpretation of arbuscular mycorrhizal fungal
681 community patterns? *New Phytol* **202**: 1101–1104.

682 Lilleskov EA, Bruns TD, Horton TR, Taylor DL, Grogan P. (2004). Detection of
683 forest stand-level spatial structure in ectomycorrhizal fungal communities.
684 *FEMS Microbiol Ecol* **49**: 319–332.

685 van der Linde S, Holden E, Parkin PI, Alexander IJ, Anderson IC. (2012). Now
686 you see it, now you don't: The challenge of detecting, monitoring and
687 conserving ectomycorrhizal fungi. *Fungal Ecol* **5**: 633–640.

688 Lõhmus A. (2009). Factors of species-specific detectability in conservation
689 assessments of poorly studied taxa: the case of polypore fungi. *Biol*
690 *Conserv* **142**: 2792-2796.

691 Money NP. (2016) Spore Production, Discharge, and Dispersal. In:
692 Watkinson SC, Boddy L, Money NP (eds.) *The Fungi*. Academic Press, pp.
693 67-97.
694

695 Newton AC, Davy LM, Holden E, Silverside A, Watling R, Ward SD. (2003).
696 Status, distribution and definition of mycologically important grasslands
697 in Scotland. *Biol Conserv* **111**: 11–23.

698 Nilsson RH, Wurzbacher C, Bahram M, R. M. Coimbra V, Larsson E,
699 Tedersoo L, *et al.* (2016). Top 50 most wanted fungi. *MycoKeys* **12**: 29–
700 40.

701 O'Brien HE, Parrent JL, Jackson JA, Moncalvo J, Vilgalys R. (2005). Fungal
702 Community Analysis by Large-Scale Sequencing of Environmental
703 Samples. *Appl Environ Microbiol* **71**: 5544–5550.

704 O'Dell TE, Lodge DJ, Mueller GM. (2004). Approaches to sampling
705 macrofungi. In: Mueller GM, Bills GF, Foster MS (eds) *Biodiversity of*
706 *Fungi: Inventory and Monitoring Methods*. Elsevier, pp 163–168.

707 Peay KG, Bruns TD. (2014). Spore dispersal of basidiomycete fungi at the
708 landscape scale is driven by stochastic and deterministic processes and
709 generates variability in plant–fungal interactions. *New Phytologist*, 204(1),
710 180–191.

711 Pellissier L, Niculita-Hirzel H, Dubuis A, Pagni M, Guex N, Ndiribe C, *et al.*
712 (2014). Soil fungal communities of grasslands are environmentally
713 structured at a regional scale in the Alps. *Mol Ecol* **23**: 4274–4290.

714 Porter TM, Skillman JE, Moncalvo JM. (2008). Fruiting body and soil rDNA
715 sampling detects complementary assemblage of Agaricomycotina
716 (Basidiomycota, Fungi) in a hemlock-dominated forest plot in southern
717 Ontario. *Mol Ecol* **17**: 3037–3050.

718 Rayner AD, Todd NK. (1980). Population and community structure and
719 dynamics of fungi in decaying wood. *Adv Bot Res* **7**: 333–420.

720 Rognes T, Flouri T, Nichols B, Quince C, Mahé F. (2016). VSEARCH: a
721 versatile open source tool for metagenomics. *PeerJ* **4**: e2584.

722 Runnel K, Tamm H, Lõhmus A. (2015). Surveying wood-inhabiting fungi: Most
723 molecularly detected polypore species form fruit-bodies within short
724 distances. *Fungal Ecol* **18**: 93–99.

725 Ryberg M, Nilsson RH, Kristiansson E, Töpel M, Jacobsson S, Larsson E.
726 (2008). Mining metadata from unidentified ITS sequences in GenBank: a
727 case study in *Inocybe* (Basidiomycota). *BMC Evolutionary Biology*, 8(1),
728 50.

729 Schadt CW, Martin AP, Lipson DA, Schmidt SK. (2003). Seasonal dynamics
730 of previously unknown fungal lineages in tundra soils. *Science* **301**:
731 1359–1361.

732 Schmidt P-A, Bálint M, Greshake B, Bandow C, Römbke J, Schmitt I. (2013).
733 Illumina metabarcoding of a soil fungal community. *Soil Biol Biochem* **65**:
734 128–132.

735 Schmit JP, Lodge DJ. (2005). Classical methods and modern analysis for
736 studying fungal diversity. In: Dighton J, White JF, Oudemans P (eds).
737 *The Fungal Community: its Organization and Role in the Ecosystem*,
738 *third ed.* CRC Press, pp 193–214.

739 Smith DP, Peay KG. (2014). Sequence depth, not PCR replication, improves
740 ecological inference from next generation DNA sequencing. *PLoS One*
741 **9**(2): e90234.

742 Straatsma G, Egli S, Straatsma G, Ayer F, Egli S. (2015). Species richness,
743 abundance, and phenology of fungal fruit bodies over 21 years in a
744 Swiss forest plot. *Mycol Res* **105**: 515–523.

745 Taberlet P, Coissac E, Pompanon F, Brochmann C, Willerslev E. (2012).
746 Towards next-generation biodiversity assessment using DNA
747 metabarcoding. *Mol Ecol* **21**: 2045–2050.

748 Taylor AFS, Finlay RD. (2003). Effects of liming and ash application on below
749 ground ectomycorrhizal community structure in two Norway spruce
750 forests. *Water, Air, Soil Pollut Focus* **3**: 63–76.

751 Taylor DL, Hollingsworth TN, McFarland JW, Lennon NJ, Nusbaum C, Ruess
752 RW. (2014). A first comprehensive census of fungi in soil reveals both
753 hyperdiversity and fine-scale niche partitioning. *Ecological*
754 *Monographs* **84**(1): 3–20.

755 Tedersoo L, Bahram M, Põlme S, Kõljalg U, Yorou NS, Wijesundera R, *et al.*
756 (2014). Global diversity and geography of soil fungi. *Science* **346**(6213):
757 1256688.

758 Truong C, Mujic AB, Healy R, Kuhar F, Furci G, Torres D, *et al.* (2017). How
759 to know the fungi: combining field inventories and DNA-barcoding to
760 document fungal diversity. *New Phytol* **214**: 913–919.

761 Tyler G. (1985). Macrofungal flora of Swedish beech forest related to soil
762 organic matter and acidity characteristics. *For Ecol Manage* **10**: 13–29.

763 Unterseher M, Jumpponen A, Öpik M, Tedersoo L, Moora M, Dormann CF, *et*
764 *al.* (2011). Species abundance distributions and richness estimations in
765 fungal metagenomics - Lessons learned from community ecology. *Mol*
766 *Ecol* **20**: 275–285.

767 White TJ, Bruns TD, Lee SB, Taylor JW. (1990). Amplification and direct
768 sequencing of fungal ribosomal RNA Genes for phylogenetics. In: *PCR -*
769 *Protocols and Applications - A Laboratory Manual, Publisher: Academic*

770 *Press*. pp 315–322.

771 Wind P and Pihl S. (2010). The Danish Red List. - The National

772 Environmental Research Institute, Aarhus University [2004].

773 redlist.dmu.dk (updated April 2010).

774 Yahr R, Schoch CL, Dentinger BTM. (2016). Scaling up discovery of hidden

775 diversity in fungi: impacts of barcoding approaches. *Philos Trans R Soc*

776 *London B Biol Sci* **371**: 1–11.

Figure 1. Frequency of species (and OTUs) among the 130 sampling sites. a) Frequency of species sorted by decreasing frequency, and grouped by species recorded with both methods or only as OTU or as fruitbody, y-axis indicates the number of sampling sites (of 130) in which a species was recorded, number of species and number of observations (a species/OTU in a site) are indicated for each group. b) Top 10 most frequent species recorded with either method. c) scatterplot of fruitbody-based frequency vs DNA frequency of the 463 species recorded by both methods. d) Species richness of the 130 sites as recorded with fruitbodies or OTUs for the 463 species recorded with both methods.

Figure 2. Sampling effort and richness variation. a) Cumulative species richness when sampling the 130 sites for the full data, Agaricomycetes and Agaricales. b) Relative standard deviation (RSD) of site species/OTU richness across the 130 sampling sites for the full data, Agaricomycetes and Agaricales.

Figure 3. Correlation between site fruitbody species richness and OTU richness. Blue lines represent the linear regression of OTU richness against species richness, while the dotted line shows the identity line ($x=y$). Correlations are shown for the taxonomic subsets (all taxa, Agaricomycetes, Agaricales). Fruitbody data is restricted to species registered at the soil surface.

803

804 Figure 4. Taxonomic composition. Plots show the number of species (and OTUs) assigned to
805 higher taxa. Composition is shown for OTUs, fruitbody (soil) and fruitbody (non-soil). Left plot
806 in each panel shows the absolute richness (number of species) within different taxa, right plot
807 shows the relative richness. a) Number of species in each class for full datasets. b) Number
808 of species in each order in Agaricomycetes. c) Number of species in each genus of

809 Agaricales. Most frequent taxa for each dataset is shown for all datasets, the rest are pooled
810 in the category 'Other'.
811
812

813

814 Figure 5. Number of red-listed species recorded. Total number of red-listed species found as
815 fruitbodies (restricted to soil fungi), and OTUs. Taxa present in the DNA reference database
816 (and thus possible to identify with both methods) are indicated in green, and taxa not present
817 in the DNA reference database (and thus not possible to identify with DNA) are indicated in
818 red.

819

820

821

822 Figure 6. Proportion variation of community dissimilarity. X-axis shows the number of
823 explanatory environmental variables selected by the model, and y-axis shows the total
824 amount of explained variation. Colours indicate the dataset (OTU, fruitbody (all data) and
825 fruitbody (soil species), line type indicates the three taxonomic subsets (All taxa,
826 Agaricomycetes and Agaricales).