1   **The loci of behavioral evolution: evidence that *Fas2* and *tilB***

2   **underlie differences in pupation site choice behavior between**

3   ***Drosophila melanogaster* and *D. simulans***

5   **Alison Pischedda [§,*], Michael P. Shahandeh [§,†], and Thomas L. Turner [†]**

6
[*]Department of Biology, Barnard College, Columbia University, New York NY 10027
[†]Department of Ecology, Evolution, and Marine Biology, University of California Santa Barbara
[§]Denotes co-first authors

Short title: Mapping *Drosophila* behavioral divergence

Key words

Pupation, behavior, genetics, evolution, *Drosophila*

Corresponding authors:

Alison Pischedda
Department of Biology
Barnard College, Columbia University
3009 Broadway
New York, NY 10027
apischedda@barnard.edu
22

23   Michael P. Shahandeh
24   Department of Ecology, Evolution and Marine Biology
25   University of California, Santa Barbara
26   Santa Barbara, CA 93106-9620
27   mshahandeh@ucsb.edu

Abstract

The recent boom in genotype-phenotype studies has led to a greater understanding of the genetic architecture of a variety of traits. Among these traits, however, behaviors are still lacking, perhaps because they are complex and environmentally sensitive phenotypes, making them difficult to measure reliably for association studies. Here, we aim to fill this gap in knowledge with the results of a genetic screen for a complex behavioral difference, pupation site choice, between *Drosophila melanogaster* and *D. simulans*. In this study, we demonstrate a significant contribution of the X chromosome to the difference in pupation site choice behavior between these species. Using a panel of X-chromosome deletions, we screened the majority of the X chromosome for causal loci, and identified two regions that explain a large proportion of the X-effect. We then used gene disruptions and RNAi to demonstrate the substantial effects of a single gene within each region: *Fas2* and *tilB*. Finally, we show that differences in *tilB* expression correlate with the differences in pupation site choice behavior between species. Our results suggest that even complex, environmentally sensitive behaviors may evolve through changes to loci with large phenotypic effects.

Introduction

The behaviors of closely related species can be remarkably different, and these differences can have important biological consequences. Behavioral evolution in insects has major impacts on crop decimation and disease vectoring (Karageorgi et al., 2017; McElhany et al., 1995). For example, one subspecies of the yellow fever mosquito, *Aedes aegypti*, prefers to bite humans, while another closely related species prefers other animals (McBride et al., 2014; Peterson, 1977). Understanding insect preferences therefore presents a major inroad to effective disease and pest management (Zhou et al., 2010). Behaviors are also critical to the creation and maintenance of biodiversity, as host, habitat, and mating preference behaviors are often key players in speciation and local adaptation (Coyne & Orr, 1997, 2004).

Despite the importance of behavioral traits, we know little about the genetic basis of their evolution. GePheBase, the most extensive compilation of natural genetic variants associated with trait differences, currently catalogs over 1800 associations, of which only 22 are for behavior (Martin & Orgogozo, 2013). From these 22, and others in the literature, it is clear that individual genes can sometimes have large effects on evolved differences in behavior (Leary et al., 2012; McGrath et al., 2011; Prince et al., 2017). It also clear that changes to sensory receptors in the peripheral nervous system can explain dramatic shifts in behavior (Cande et al., 2013; Leary et al., 2012; McBride et al., 2014). With so few studies, however, it is difficult to conclude how frequently we expect single loci to have large effects or how often sensory receptors explain species differences, due to pervasive ascertainment bias (Rockman, 2012). Indeed, a recent study discovered an exception to this emerging pattern: the evolution of a central neural circuit, rather than a peripheral sensory neuron, explains differences in mating behavior in two *Drosophila* species (Seeholzer et al., 2018).

The current lack of genetic studies mapping behavioral variation presumably arises from

the fact that behaviors are difficult to measure reliably and repeatedly. Behavioral phenotypes

often integrate multiple cues, are sometimes context dependent, and can be innate or learned,

making it difficult to exclude environmentally induced variation. To better understand the

genetic basis of behavioral evolution, we therefore need more case studies, with a focused effort

on "metamodel" systems with documented behavioral differences between closely related

species (Kopp, 2009).

Flies in the genus *Drosophila* are well poised to address these challenges. In *Drosophila*,

hundreds of genetically identical individuals from variable wild-caught strains can be reared in a

common environment, isolated at the beginning of their adult life stage, and repeatedly assayed

for a trait of interest. Such a design significantly reduces the potential for environmentally

induced variation to obscure genetic differences in behavior. Additionally, there are many

Drosophila species with large, characterized differences in a variety of complex behaviors

(Orgogozo & Stern, 2009). Undeniably, work comparing the repeated evolution of

morphological traits across closely related species of *Drosophila* has significantly advanced our

understanding of the general patterns linking genotype and phenotype for developmental traits

(Kittelmann et al., 2017; Rebeiz & Williams, 2017; Sucena et al., 2003). Studies that investigate

genetic complexity in *Drosophila*, where fine-mapping and functional follow-up are possible,

should make similar progress for behaviors.

In the present study, we seek to address the lack of behavioral association studies using

the *Drosophila* model system. Here, we investigate the difference in pupation site preference

between two *Drosophila* species: *D. melanogaster* and *D. simulans*. When the larval stages of

these species are ready to metamorphose into adults, they first enter a pupal stage. The pupa,

which lasts for a number of days, is immobile and therefore vulnerable to parasitism, predation,

desiccation, and disease (Markow, 1981a). Before pupating, larvae enter a "wandering" stage,

where they search for an appropriate pupation site (Riedl et al., 2007; Sokolowski et al., 1984).

Depending on the strain and species, larvae vary from pupating directly on their larval food

source to traveling more than 40 cm away from it (Stamps et al., 2005). This behavior has been

extensively studied, and is exquisitely sensitive to environmental conditions— individuals alter

their behavior in response to light, moisture, pH, the presence of other species, parasitism, and

more (Hodge & Caslaw, 1998; Markow, 1981b; Sameoto & Miller, 1968a; Seyahooei et al.,

2009). Despite this environmentally induced variation, the effects of genotype on preference are

considerable. Within species, strains and populations often differ in how far they travel from

their food source before pupating, although the most consistent experimentally demonstrated

differences are between species (Markow, 1979; Vandal et al., 2008). Interestingly, differences

in pupation site choice behavior between species do not correspond to their taxonomic

classification (Shivanna et al., 1996). For example, *D. melanogaster* and *D. simulans* shared a

common ancestor 2-3 million years ago (Lachaise & Silvain, 2004), and are extremely similar in

terms of their ecology, morphology, and physiology (Parsons, 1975). Previous work shows,

however, that they differ markedly in terms of pupation site choice, with *D. simulans* pupating

closer to the larval food source, on average (Markow, 1979, 1981a). This difference is not due to

laboratory adaptation, as freshly collected individuals show the same pattern (Markow, 1979,

1981a). These species are frequently collected in the same microhabitats, and their differences in

pupation site choice behavior have been postulated to be a form of niche partitioning (Markow,

1979, 1981a). Supporting this hypothesis, pupation site choice responds to density dependent

selection in the laboratory (Mueller & Sweet, 1986), and provides a potential increase in

competitive ability between species ovipositing in the same media (Arthur & Middlecote, 1984).

Here, we investigate the genetic basis of this difference in pupation site choice between *D.*

melanogaster and *D. simulans*. Despite substantial reproductive isolation, *D. melanogaster*

females can be persuaded to mate with *D. simulans* males in the lab, and vigorous female F1

hybrids result from the cross. Males are usually inviable, but we use a *D. simulans* hybrid male

rescue strain (Brideau et al., 2006a; Watanabe, 1979) to circumvent this challenge, and show that

a significant proportion of the species difference in pupation behavior can be mapped to the X

chromosome, consistent with findings using other *Drosophila* species (Erezyilmaz & Stern,

2013). Still, these hybrids remain sterile, so genetic dissection using a backcross mapping

population is not possible. Instead, we use a widely available set of transgenic *D. melanogaster*

deficiency lines to screen a substantial portion of the X chromosome (Cook et al., 2012; Ryder et

al., 2004). We use these lines to create hybrid females that lack large, overlapping portions of the

X chromosome from *D. melanogaster*, and therefore express only *D. simulans* alleles in those

regions. We identify two loci with large effects on pupation behavior. We then employ genetic

knockouts of candidate loci within these regions to demonstrate their effects, and use RNAi

knockdown to further test the role of two genes, *touch insensitive larvae B* (*tilB*) and *Fasciclin 2*

(*Fas2*). Finally, we use real time qRT-PCR to test for species-level differences in gene

expression of *tilB* in larvae, and show that *tilB* is more highly expressed in *D. melanogaster* than

in *D. simulans*.

Methods

General fly maintenance

Unless otherwise stated, we maintained all fly strains for these experiments in 20 mm diameter vials containing standard cornmeal-molasses-yeast medium at 25°C under a 12h:12h light/dark cycle at 50% relative humidity. Under these conditions, we established non-overlapping two-week lifecycles as follows. For all stocks, except LH_M and *Lhr* (see below), we transferred male and female adult flies into fresh vials containing food media supplemented with live yeast on the surface for 1-3 days, at which point the flies were discarded. 14 days later (after all progeny had eclosed), we again transferred adult flies into fresh vials for 1-3 days to begin the next generation. We maintained LH_M and *Lhr* identically, except we additionally regulated density by transferring only 10 males and 10 females to begin the next generation.

Characterizing pupation behavior for *D. melanogaster* and *D. simulans*

We measured pupation behavior for 11 *D. melanogaster* and 12 *D. simulans* strains collected from various locations throughout the world (Table S1). The 11 *D. melanogaster* strains included 10 of the "founder" wild-type inbred lines of the Drosophila Synthetic Population Resource (King et al., 2012; flyrils.org), and a single wild-type line created from the LH_M laboratory-adapted population (Rice et al., 2005). The 12 *D. simulans* strains included 11 wild-type strains and a single strain carrying *Lethal hybrid rescue* (*Lhr*), a mutation that restores viability in *D. melanogaster*/*D. simulans* hybrid males.

To measure pupation behavior, we placed 10 males and 10 females from a specific line (both 3-5 days old) into half pint bottles and allowed females to oviposit overnight on a 10 mm

diameter petri dish filled with food medium that was placed in the opening of the bottle. In total,

we set up 5 bottles for each line. The following morning, we transferred 100 eggs from the petri

dishes into vials containing food medium that were lined with an acetate sleeve on which the

larvae could pupate. In total, we set up 5-8 vials per line. Vials were held at 25°C for 8 days, at

which time the liner was removed and the locations of the pupae were recorded (8 days was long

enough for almost all larvae to pupate without any flies eclosing). A pupa was considered "on"

the food if it was within 1 cm of the food surface, while all pupae that were further than 1 cm

from the food surface were considered "off" the food. For each vial, we calculated the proportion

of pupae on the food surface. For comparisons between species, our unit of replication was the

mean proportion of pupae on the food surface from each line (i.e. N=11 for *D. melanogaster* and

N=12 for *D. simulans*).

Crossing *D. melanogaster* with *D. simulans*

For all crosses below, we created F1 hybrids between *D. melanogaster* and *D. simulans* using the

following protocol. *D. simulans* males were collected as virgins within 6 hours of eclosion and

held at room temperature in groups of 20 in vials containing food medium for 3-4 days. To set up

crosses, we collected young *D. melanogaster* virgin females within 2-3 hours of eclosion, and

combined 8-12 of these females with 20 *D. simulans* males in vials containing food medium

supplemented with an *ad lib* amount of live yeast on the surface. We then pushed a long foam

plug down into the vial, leaving approximately 1 cm of space above the food surface. We held

flies under these conditions for 3 days, at which time they were transferred from these "cross

vials" into "pupation vials" that contained food medium with no added yeast, and were lined

with an acetate sleeve on which the larvae could pupate. We always set up crosses using *D.*

melanogaster females and *D. simulans* males, because crosses in the opposite direction were

never successful.

Measuring pupation behavior in F1 hybrids

To create F1 hybrid males and females, we used *D. simulans* males from the *Lethal hybrid*

rescue (*Lhr*) strain (Watanabe, 1979)*.* The *Lhr* mutation restores viability in F1 hybrid males,

which are usually lethal (Brideau et al., 2006b). To create F1 hybrid females and F1 males with a

D. melanogaster X chromosome ("melX" males), we crossed wild-type females from our LH_M

strain (provided by William Rice) to *Lhr D. simulans* males (Fig 1A). Because we were unable to

successfully cross *D. simulans* females to *D. melanogaster* males, we created F1 hybrid males

with the *D. simulans* X chromosome ("simX" males) by crossing *D. melanogaster* LH_M females

that carry a compound X chromosome (*C(1)DX y f*) (Rice et al., 2005) to *D. simulans Lhr* males.

The background of this strain is identical to that of LH_M, with the exception of the compound X

chromosome. The compound X in these females ensured that the X chromosome was transmitted

from *D. simulans* fathers to their F1 hybrid sons (Fig 1B). For each direction of the cross, we

combined *D. melanogaster* females with *D. simulans* males, as described above. This crossing

scheme ensures that all maternal inheritance (cytoplasmic and mitochondrial) in the reciprocal

male hybrid crosses originates from the *D. melanogaster* parent. Thus, these hybrids have an

identical background with the exception of the sex chromosomes, and any differences we

observe between melX and simX males are directly attributable to their different sex

chromosomes.

After 3 days in the cross vial, we transferred males and females into pupation vials for 24

hours, at which time the flies were removed. While screening hybrid pupation behavior, we also

concurrently screened pupation behavior for the parental *D. melanogaster* strain (LH$_M$) and the

parental *D. simulans* strain (*Lhr*) for comparison. Parental strain cross vials contained only a

moderate amount of yeast, were set up with only 5 males and 5 females (pure species crosses

produce more offspring), and did not have a plug pushed down into the vial, but were otherwise

treated identically to the hybrid crosses. In total, we set up 30-33 vials per treatment.

All pupation vials were held at 25°C for 8 days, at which time the liner was removed. We

removed any remaining larvae, and cut the liner at a point 1 cm above the food surface. The

portion of the liner that contained pupae within 1 cm of the food surface was returned to the

original vial (the "on vial"), while the portion of the liner with pupae further off of the food

surface was placed in another vial containing food medium (the "off vial"). The flies that eclosed

were sexed and counted 7 days later (15 days post-egg); all flies that eclosed in the "on vial"

were considered flies that pupated on the food surface, while all flies that eclosed in the "off vial"

were considered flies that pupated off the food surface. We then calculated the proportion of

individuals that pupated on the food for each type of individual (genotype and sex).

To assess the validity of using hybrid behavior to map interspecific differences, we set up

an additional experiment to ensure that hybrids behaved typically with respect to pupation site

choice behavior. This is a potential concern, as hybrids between *D. melanogaster* and *D.*

simulans are known to differ from either parent in a variety of traits (Barbash & Ashburner, 2003;

Sturtevant, 1920; Takano, 1998). For a subset of pupation vials, we calculated the average

pupation height of males and females from each strain as follows. Instead of dividing the

pupation liner into "on" food and "off" food sections, we cut the liner at 8 intervals, each spaced

1 cm from the last, starting at the food surface. These portions of the liner were ranked from 1

(on the food surface) to 8 (the furthest from the food surface), and were transferred to separate

vials containing food medium. 7 days later the flies that eclosed were counted, and we used the

rankings to calculate a mean pupation height, such that a higher value indicates a farther distance

from the food. We performed this experiment for our *D. simulans* strain (*Lhr*), our reciprocal

hybrid crosses (melX and simX in an LH_M background), the *D. melanogaster* strain w^{1118}, and

melX hybrids in a w^{1118} background (w^{1118} is the background strain for the majority of

deficiencies we used in our screen).

Fig 1. Crossing schemes to generate reciprocal X chromosome hybrid males and deficiency/balancer hybrid females. A. Crossing wild-type *D. melanogaster* females (LH$_M$, shown in white) to *D. simulans* males (*Lhr*, shown in grey) produces hybrid males with a *D. melanogaster* X chromosome (melX) and hybrid females. **B.** Crossing a compound X C(1)DX LH$_M$ female to *Lhr* males produces hybrid males with a *D. simulans* X chromosome (simX). Females of this cross would inherit two *D. melanogaster* X chromosomes and a *D. simulans* Y chromosome, but are inviable. **A and B.** Note that the background of the reciprocal male hybrids resulting from each cross (melX and simX) is an identical combination of *Lhr* and LH$_M$ with the exception of the sex chromosomes. **C.** Crossing *D. melanogaster* X chromosome deficiency lines, which have a balancer X chromosome with a dominant visible marker (DVM) and an X chromosome with a large deletion, to *Lhr* produces deficiency hybrid females, balancer females, balancer males, and deficiency males (*mostly dead due to large deletions on a hemizygous chromosome with some deficiency lines being exceptions).

Mapping hybrid pupation behavior using the Bloomington Deficiency Kit

Because we found a significant effect of the X chromosome on the difference in pupation site

choice behavior between *D. simulans* and *D. melanogaster*, we devised a crossing scheme using

molecularly engineered chromosomal deficiencies to screen the X chromosome for loci

contributing to this difference. These deficiencies are part of the Bloomington Deficiency Kit

(Cook et al., 2012), available from the Bloomington Drosophila Stock Center (BDSC). We

assayed a total of 90 deficiency strains covering 87% of the X chromosome (Table S2). We

restricted our deficiency screen to lines from the BSC, Exelixis, and DrosDel sets to control for

strain background effects while also maximizing chromosome coverage.

To set up crosses, we collected deficiency females as young virgins (2-3 hours after

eclosing) and crossed them to *D. simulans* males from the *Lhr* strain. After 3 days in the cross

vial, we transferred males and females into pupation vials for 24-48 h, at which time the flies

were removed. We then divided the pupations vials into "on" and "off" vials as we did for our F1

hybrids (above).

These crosses produced two types of hybrid female that were heterozygous for *D.*

melanogaster/*D. simulans* at each autosome (Fig 1C). The deficiency hybrid females contain the

D. melanogaster deficiency X chromosome and a *D. simulans* X chromosome, making them

hemizygous for a segment of the X chromosome. At this locus, these hybrid females only

express *D. simulans* alleles. The balancer hybrid females contained the *D. melanogaster* balancer

X chromosome (marked with a dominant visible marker) and a *D. simulans* X chromosome.

These females are heterozygous for *D. melanogaster*/*D. simulans* over the entirety of the X

chromosome, and thus express both *D. simulans* and *D. melanogaster* alleles. Although the

deficiency hybrids are our flies of interest, the balancer hybrids provide an experimental control,

as these females developed in the same environment as our experimental flies. As a result, we

calculated the proportion of deficiency hybrid females that pupated on the food and the

proportion of balancer hybrid females that pupated on the food. We then used these measures to

calculate a "pupation index" as the proportion of deficiency hybrids pupating on the food divided

by the proportion of balancer females pupating on the food. To increase the accuracy of our

estimates, we only included pupation vials in our analysis that yielded at least 10 of each type of

female. For each deficiency hybrid strain we measured, we report the median pupation index of

all replicates, because there were often high-scoring outliers that significantly skewed the mean

pupation index. These outliers almost always had abnormally high pupation indices, so focusing

on median values makes our findings more conservative.

Any deficiency hybrid cross with a median pupation index greater than 1 indicates that

more deficiency females pupated on the food compared to balancer females, potentially because

the deficiency includes *D. melanogaster* genetic variation that is involved in pupation site choice

behavior. Alternatively, simply creating flies that are hemizygous at a locus on the X

chromosome may result in a variety of pleiotropic effects that make larvae less likely to climb up

the vial. To test for this, when a deficiency hybrid cross showed a pupation index significantly

greater than 1 (Table S2), we crossed that *D. melanogaster* deficiency strain to a *D.*

melanogaster wild-type strain (T.4). If these *D. melanogaster* deficiency crosses displayed the

same pattern, we considered the effect of the deficiency on pupation behavior to be a byproduct

of deleting a large portion of the X chromosome, rather than revealing recessive *D. simulans*

variation, and discarded them. If instead the pupation index for the *D. melanogaster* cross was

significantly lower than the pupation index for the *D. simulans* cross, we pursued that deficiency

for further validation. To ensure that this pattern is not a result of epistasis from the hemizygous

region in a hybrid background, we further crossed these deficiencies to an additional *D.*

melanogaster (T.1) and *D. simulans* (Mex180) strain to test for background-specific effects.

Testing candidate genes in deficiency regions using gene knockouts and RNAi knockdown

For regions of interest identified by our deficiency screen, we ordered transgenic knockouts for

any genes available within the region at the time. Our deficiency screen identified two regions of

interest. The first is the overlap of Df(1)BSC869 and DF(1)ED6720, excluding the region

covered by DF(1)ED6727, which did not have a pupation index greater than 1 (Fig 4A, Table

S2). The resulting region of interest spans X:4,204,351 - 4,325,174 (Fig 4C). Within the first

region, there are 23 genes, of which 20 are protein coding (Table S3A). According to

modENCODE expression data, 15 of those 20 protein coding genes are expressed in larvae,

while only 6 are also expressed in the larval nervous system, which we would expect for genes

regulating behavior (Graveley et al., 2011). Five of these six genes are well described. At the

time of assay, only three of the five characterized genes within this region had non-lethal verified

loss-of-function alleles available: *Fas2* (Fasciclin2), *mei9* (meiotic 9), and *norpA* (no receptor

potential A). We tested knockouts of each for an effect on pupation site choice behavior. We

screened two *Fas2* knockouts, the mutant allele *Fas2^{eb112}* (*Fas2^{eb112}*/FM7c; Grenningloh et al.,

1991; provided by Brian McCabe), and a p-element insertion allele, *Fas2^{G0293}* (former BDSC

Stock 11850; full genotype: *w^{67c23}P{lacW}fas2^{G0293}*/FM7c). It is important to note that the nature

of the lesion is uncharacterized for *Fas2^{G0293}* (although our behavioral data below suggest that it

is indeed a loss of function allele, as it behaves indistinguishably from the verified knockout

Fas2^{eb112}). We additionally screened the *mei-9^{A1}* mutant allele (*w^{1} mei-9^{A1}*/FM7h; BDSC stock

#6792) and the *norpA^{36}* mutant allele (*w* norpA^{36}*; BDSC stock #6792). Because *norpA^{36}* is not

held over a balancer, we also set up crosses using a *norpA* rescue strain created in the same

background (*w* norpA³⁶*; *P{w[+mC]=ninaE.norpA.E}2*; BDSC stock # 52276) as a control. It is

worth noting that while *norpA* is expressed in the larval nervous system, it is only partially

contained within this region, and is also largely deleted by another deficiency (Df(1)ED6727;

Table S2) that did not display a pupation index significantly greater than 1 or 0.88 (the grand

mean for all deficiency hybrids).

The second region of interest identified by our deficiency screen was the region deleted

by DF(1)Exel6255 (X:21,519,203 – 22,517,665; Fig 4D). Within this region are 28 genes, of

which 22 are protein coding (Table S3B). Of the 22 protein coding genes, 14 are expressed in

larvae – 13 of which have some expression in the larval nervous system (Graveley et al., 2011).

Of these, 7 are described. We obtained knockout strains for both of the characterized genes

expressed in the larval nervous system that had verified loss-of-function alleles available at the

time: *tilB* (touch insensitive larva B) and *wap* (wings apart). We screened two *tilB* mutant alleles,

tilB¹ and *tilB²* (*y w tilB¹ᐟ²* /FM4; Kernan et al., 1994; provided by Daniel Eberl), and the *wap²*

mutant allele (*wap²*/FM6; BDSC stock # 8133).

Like the deficiency strains, each of our gene disruptions (with the exception of *norpA*) is

held over a balancer chromosome with a visible marker. To measure the pupation behavior of

hybrids containing knockout copies of these *D. melanogaster* genes, we crossed each *D.*

melanogaster knockout strain to *Lhr* using the previously described methods, and calculated the

pupation index as the proportion of knockout females on food / the proportion of balancer

females on the food. As for our deficiency screen, any hybrid knockouts with a median PI greater

than 1 for the hybrid cross suggest that the knockout gene may be involved in pupation site

choice. We also crossed each *D. melanogaster* knockout strain to a wild-type *D. melanogaster*

(T.4) strain to ensure that hybrid knockouts with a pupation index greater than 1 are not simply

an artifact of being hemizygous for this particular gene. We crossed knockout strains that

displayed the pattern we expect for a gene involved in pupation site choice (i.e. a pupation index

significantly greater than 1 when crossed to *D. simulans*, which is also significantly greater than

the pupation index when crossed to *D. melanogaster*) to an additional *D. simulans* (Mex180) and

D. melanogaster (T.1) wild-type strain for verification. For *norpA*, we crossed the mutant and

control strains to *Lhr*, and compared hybrid *norpA* mutant females to hybrid *norpA* rescue

females. We additionally crossed these strains to the *D. melanogaster* T.4 strain.

Our knockout screen identified two genes that appear to be involved in pupation site

choice: *tilB* and *Fas2*. We further tested the effects of these genes on pupation behavior using

RNAi knockdown in *D. melanogaster*. We used the elav-Gal4 driver (P{w[+mC]=GAL4-

elav.L}2/CyO; BDSC #8765), which expresses Gal4 throughout the nervous system. We drove

down the expression of *tilB* and *Fas2* throughout the nervous system by crossing elav-Gal4

virgin females to UAS-*tilB* (BDSC #29391: y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.JF03324}attP2)

and UAS-*Fas2* (BDSC #34084: y[1] sc[*] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.HMS01098}attP2)

males, respectively. The resulting flies express a gene-specific hairpin RNA throughout the

nervous system, causing the degradation of mRNA, and thus, reduced expression of that gene

(Perkins et al., 2015; Zeng et al., 2015). As experimental controls for each RNAi cross, we also

crossed elav-Gal4 virgin females to the RNAi background stock (*y v; attP2, y+* (y[1] v[1] ;

P{y[+t7.7]=CaryP}attP2; BDSC stock # 36303) and the Gal4-1 stock (containing a hairpin RNA

targeting Gal4 in VALIUM20; BDSC stock # 35784). Together, these controls allow us to

account for the effect of both the Gal4 mutation and general expression of hairpin RNA

throughout the nervous system. Any differences we detect between these controls and our RNAi

crosses must therefore be due to the expression of the gene-specific (*tilB* or *Fas2*) hairpin RNA.

We set up pupation vials using the methods described above, and for each cross, we calculated

the proportion of RNAi (or control) flies on the food (removing any data points with fewer than

20 experimental flies). If more RNAi flies pupate on the food in the experimental cross (in which

the expression of the gene is driven down) compared to the control crosses (in which gene

expression is unaffected), this provides further support for that gene's involvement in pupation

site choice. For these vials, we additionally determined the average pupation height of flies from

each cross using the methods described above for F1 hybrids; if RNAi flies pupate closer to the

food than flies from the control crosses, this suggests that gene may be involved in pupation site

choice.

Testing *tilB* for species-specific differences in larval transcript expression

We selected two each of our 11 *D. melanogaster* and 12 *D. simulans* strains to test for larval

stage-specific expression differences of candidate genes using real time qRT-PCR – one extreme

and one average. For *D. simulans*, we selected Geo288 and Per005 (Table S1), because Geo288

has the highest proportion of pupae on the food of the *D. simulans* strains and Per005 is closest

to the species mean (Fig 2). For *D. melanogaster*, we selected CA1 and T.4 (Table S1) because

T.4 has the lowest proportion of pupae on the food of the *D. melanogaster* strains, and CA1 is

closer to the species mean (Fig 2).

To harvest larvae from these strains, we allowed adult females to oviposit in standard

vials containing food media between the hours of 8 AM and 12 PM over two consecutive days.

120 hours after the final oviposition day, we floated larvae out of the food media using a 20%

sucrose in water solution, sucked them up using a transfer pipet, briefly rinsed them with DI

water on cheesecloth, and snap froze them using liquid nitrogen (Nichols et al., 2012). In this

way, we collected 20-30 mg of larvae from two developmental time points: 96 and 120 hours

following oviposition. These time points approximate early wandering and late wandering larval

stages. We chose these time points because they are presumably when pupation site choice

occurs, and because the larvae are large enough for many to be harvested at once using the above

methods. We extracted mRNA using the Qiagen RNeasy Plus Mini Kit, and prepared cDNA

using the Promega Verso kit.

To quantify transcript abundance, we designed primers that span a single intron near the

3' end of *tilB* (Huggett et al., 2005). Additionally, we used primers for the gene *RpL32* as an

internal control (Al-Atia et al., 1985; Chertemps et al., 2007). *Fas2* is a complex gene with

multiple splice forms, so we were unsuccessful in designing general primers that would amplify

all transcripts in both species. For this reason, we did not include *Fas2* in these experiments. A

full list of primers and transcripts can be found in Table S4A. For each stage and strain, we

prepared two to three biological replicates, which we then amplified in two technical replicates

for 40 rounds of qPCR. Using *RpL32* transcript number as an internal control, we calculated

relative transcript abundance while correcting for species differences in primer efficiency. We

estimated primer efficiency differences by serially diluting gDNA from each of our *D.*

melanogaster and *D. simulans* strains, performing qPCR, and using a standard curve to calculate

adjusted amplification factors (Table S4B). To ensure that we were amplifying cDNA made from

RNA, and not gDNA contamination, we performed gel electrophoresis on our cDNA samples to

ensure we only visualized the short, intron-less, band.

Estimating effect sizes

We used our results to estimate how much of the difference in pupation site preference

between *D. melanogaster* and *D. simulans* can be attributed to: *i*) the X chromosome, *ii*) our

deficiencies of interest (Df(1)BSC869/DF(1)ED6720 and Df(1)Exel6255), and *iii*) *Fas2* and *tilB*.

We first calculated the "species difference ratio" (using the data from our parental/hybrid screen

in Fig 3) by dividing the median proportion of males on the food for *D. simulans* by the median

proportion of males on the food for *D. melanogaster* males (species difference ratio = 5.27). We

then calculated an "X effect ratio" by dividing the median proportion of simX males on the food

by the median proportion of melX males on the food. To determine how much of this species

difference can be attributed to the X chromosome, we divided the "X effect ratio" by the

"species difference ratio".

We then estimated the effect size of our deficiencies by calculating "deficiency effect

ratios" (the median pupation index when the deficiency was crossed to the *D. simulans Lhr*

strain/the median pupation index when the deficiency was crossed to the *D. melanogaster* T.4

strain). Because Df(1)BSC869 and DF(1)ED6720 overlap, we used their mean deficiency effect

ratio to estimate the effect size of the overlapping region. To determine how much of the X effect

difference can be attributed to each deficiency region, we divided the "deficiency effect ratio" by

the "X effect ratio".

Finally, we estimated the effect size of our two identified candidate genes, *Fas2* and *tilB*.

For each allele we tested, we calculated a "knockout effect ratio" (the median pupation index

when the knockout was crossed to the *D. simulans Lhr* strain/the median pupation index when

the knockout was crossed to the *D. melanogaster* T.4 strain), and then we used the average

knockout effect ratio of the two alleles for each gene (i.e. *Fas2*: average of *Fas2^{eb112}* and

Fas2^{G0293}; *tilB*: average of *tilB^1* and *tilB^2*). We then determined the contribution of each gene to

its deficiency effect size, and to the overall X effect, by dividing the "knockout effect ratio" by

the "deficiency effect ratio" and the "X effect ratio", respectively.

For all effect size estimates, we calculated bootstrapped 95% confidence intervals using

100,000 bootstraps.

Data availability

All relevant data supporting the findings of this study are represented in the figures and

supplementary tables.

Results:

1. Differences in pupation behavior between *D. melanogaster* and *D. simulans*

We found significant variation among the 11 *D. melanogaster* and 12 *D. simulans* strains in the

proportion of individuals that pupated on the food surface (Fig 2; *D. melanogaster* Wilcoxon test:

χ^2= 42.69, df = 10, p<0.0001; *D. simulans* Wilcoxon test: χ^2= 56.34, df = 11, p<0.0001). When

we tested for a species difference in pupation behavior, we found that our *D. simulans* lines had

significantly more pupae on the food surface compared to our *D. melanogaster* lines, on average

(Fig 2; Wilcoxon test: χ^2= 8.37, df = 1, p = 0.0038).

Although we controlled egg density to characterize species differences in pupation

behavior, there were viability differences among our surveyed lines. As a result, we had

significant variation in the total number of pupae in each vial for our *D. melanogaster* lines

(ANOVA: $F_{10,65}$=5.58, p<0.0001) and our *D. simulans* lines (ANOVA: $F_{11,66}$=8.01, p<0.0001).

Previous studies have found that larval density correlates with pupation height in *D.*

melanogaster (Sokal et al., 1960). To control for differences in density, we also performed the

same analyses above using the residuals from a regression between number of pupae in the vial

and proportion of pupae on the food. None of our findings changed using this analysis (Fig S1),

indicating that our results were not affected by variation in larval density.

Although most of our *D. melanogaster* lines have been in the lab since the 1950s and

1960s, we found no difference in the proportion of pupae on the food between *D. simulans*

strains collected in the 1950s/1960s and those collected in the 2000s (Table S1; Wilcoxon test:

χ^2= 0.24, df=1, p=0.62), indicating that the species difference we report here is unlikely to be an

artifact of laboratory adaptation in our surveyed strains.

Fig 2. Pupation behavior differences between *D. melanogaster* and *D. simulans*.
The mean proportion of individuals in a vial that pupated on the surface of the food for 11 *D.*
melanogaster lines and 12 *D. simulans* lines described in Table S1. Error bars denote the 95%
confidence interval around each individual mean (N = 5-8). The dashed horizontal lines indicate the grand
mean for each species. The boxes surrounding the dashed lines denote the 95% confidence interval
around the grand mean.

2. Differences in pupation behavior have a significant X effect

We found significant differences among F1 hybrid genotypes when we screened hybrids

alongside their parental strains (Fig 3; Full model Wilcoxon test: χ^2= 144.57, df= 6, p<0.0001).

Specifically, a significantly higher proportion of F1 hybrid males pupated on the food when they

had inherited a *D. simulans* X chromosome (simX males) compared to a *D. melanogaster* X

chromosome (melX males), indicating that this species divergence in pupation behavior has a

significant X effect (p<0.0001 after correcting for multiple comparisons; Fig 3). This is

supported by the fact that the proportion of individuals that pupated on the food was not

significantly different between melX hybrid males and *D. melanogaster* males, or between simX

hybrid males and *D. simulans* males. All of these findings are unchanged when we control for

density effects (Fig S2).

Additionally, we found no difference in the proportion of F1 hybrid females and melX

hybrid males (p=0.70) that pupated on the food, while F1 hybrid females pupated on the food

significantly less often compared to simX hybrid males (p<0.0001 after correcting for multiple

comparisons; Fig 3). The fact that F1 hybrid females behave identically to melX hybrid males

indicates that the variation in pupation behavior on the *D. melanogaster* X chromosome (i.e.

fewer pupae on the food) is dominant to the pupation behavior on the *D. simulans* X

chromosome (i.e. more pupae on the food), because F1 hybrid females have one X chromosome

from each species.

The fact that hybrid males and females with a *D. melanogaster* X chromosome behave

indistinguishably from their *D. melanogaster* parent strain, and hybrid males with a *D. simulans*

X chromosome behave indistinguishably from their *D. simulans* parent strain (Fig 3), suggests

that hybrid pupation site choice behavior falls well within the typical range exhibited by either

parent strain. We found a similar pattern when we compared the average pupation height of F1

hybrids and their parents: F1 females and melX males did not have significantly different

pupation heights compared to their *D. melanogaster* parents, and simX males did not differ from

their *D. simulans* parental strain (Fig S3, Table S5). This is important to note, as it implies that

hybrids, particularly simX hybrids, are not just unable to climb the vial walls due to

developmental inconsistencies caused by intrinsic incompatibilities. If anything, simX hybrids

displayed slightly higher proportions of individuals climbing the vial walls (and pupated slightly

further from the food) compared to their parent strain, underscoring their vigor.

Fig 3. Pupation site choice behavior for *D. melanogaster*, *D. simulans*, and their F1 hybrids. The
proportion of individuals that pupated on the food surface for males and females from both species and
their F1 hybrids. *D. melanogaster* males and females were taken from the LH$_M$ strain, while *D. simulans*
males and females were taken from the *Lhr* strain. F1 hybrids resulted from a cross between these two
strains. The "melX" hybrid males have the *D. melanogaster* X chromosome, and the "simX" hybrid males
have the *D. simulans* X chromosome. Both hybrids have *D. melanogaster* cytoplasmic inheritance. Box
plots display the median (bold bar), interquartile range (box), and full extent of the data set excluding
outliers (whiskers). Bars labeled with different letters are significantly different from one another after
sequential Bonferroni correction for multiple comparisons (p < 0.0001 after correction, N = 26-33).

3. A deficiency screen of the X chromosome identifies two regions of interest

We found significant variation in pupation index among the deficiency hybrid crosses

(Kruskal-Wallis Test: χ^2=336.90, df=89, p<0.0001; Fig 4A). A pupation index greater than 1

indicates that more deficiency hybrids pupated on the food than balancer hybrids. This suggests

that the *D. melanogaster* deficiency region may be revealing recessive *D. simulans* genetic

variation that causes the deficiency hybrids to pupate on the food surface. However, the average

pupation index across all 90 deficiency hybrid crosses was 0.88 (Fig 4A), which was

significantly lower that our expected mean of 1 (Wilcoxon test: p<0.0001). As a result, we

compared the pupation index for all deficiencies to our expected value of 1 and to the grand

mean pupation index for these lines (0.88). Six deficiencies had pupation indices significantly

greater than one: (Df(1)ED411, Df(1)BSC869, Df(1)ED6720, Df(1)ED6906, Df(1)BSC530, and

Df(1)Exel6255). Three of these, (Df(1)BSC869, Df(1)ED6906, and Df(1)Exel6255), remained

significant after sequential Bonferroni correction for multiple comparisons (Fig 4A; Table S2).

Because the other 3 deficiencies, (Df(1)ED411, Df(1)ED6720, and Df(1)BSC530), had pupation

indices significantly greater than 0.88 after sequential Bonferroni correction (Table S2), we

included them in our list of potential deficiencies of interest.

To ensure that the deficient region actually reveals *D. simulans* variation contributing to

pupation site choice behavior, rather than creating lines that behave abnormally due to the

extended hemizygosity within the deficiency region, we crossed each of the six significant

deficiencies listed above to the T.4 wild-type *D. melanogaster* strain. For two of the six

deficiency strains, Df(1)ED411 and Df(1)BSC530, we found no difference in the pupation index

when crossed to *D. melanogaster* compared to the pupation index when crossed to the *Lhr* strain

(Wilcoxon tests; Df(1)ED411: n= 29-36, χ^2=1.74, p=0.19, Fig S4A; Df(1)BSC530: n=18-22,

χ^2=1.47, p=0.23, Fig S4C), and Df(1)ED6906 had a significantly higher pupation index when

crossed to *D. melanogaster* (n= 26-33, χ^2=9.42, p=0.002; Fig S4B). These results suggest that

the phenotypes of these three lines are a result of their hemizygosity within the deficiency region,

as the deficient *D. melanogaster* females still pupate on the food more often than the balancer

females. Note, this effect is unlikely to be driven by the balancer females, as the same balancers

occur in many of the other deficiency lines we screened.

When we crossed the remaining three deficiency strains, Df(1)BSC869, Df(1)ED6720,

and Df(1)Exel6255, to the T.4 *D. melanogaster* strain, we found a pupation index significantly

lower than the index we calculated when crossing to *Lhr* (Wilcoxon tests; Df(1)BSC869: n= 51,

χ^2=4.35, p=0.037; Df(1)BSC6720: n= 30-35, χ^2=7.90, p=0.0049; Df(1)Exel6255: n= 55-57,

χ^2=4.68, p=0.0306; Fig 4B), and no different than one (Df(1)BSC869: p=0.46; Df(1)BSC6720:

p=0.10; Df(1)Exel6255: p=0.50). This result suggests that these deficient regions do reveal

recessive *D. simulans* variation affecting pupation site choice behavior in hybrids, but have no

effect when made hemizygous in *D. melanogaster.* Two of these three deficiencies overlap:

Df(1)BSC869 and Df(1)ED6720 (Fig 4A). To further confirm that this pattern is not unique to

these specific *D. melanogaster* and *D. simulans* wild-type strains, we crossed one of the

overlapping deficiency strains, Df(1)BSC869, and Df(1)Exel6255 to another *D. melanogaster*

wild-type strain (T.1) and another *D. simulans* wild-type strain (Mex180). The pattern remained

consistent for both of these deficiencies: when crossed to *D. melanogaster,* the pupation index

was significantly lower than when crossed to *D. simulans* (Fig 4B; Wilcoxon tests: Df(1)BSC869:

n=50-51, χ^2=7.78, p=0.0053; Df(1)Exel6255: n= 23, χ^2=7.74, p=0.0054).

One deficiency strain, Df(1)ED7265, had a pupation index significantly lower than the

grand mean of all the lines (0.88) following Bonferroni correction for multiple comparisons

(Table S2). This suggests the potential for a region of the *D. simulans* X chromosome with

transgressive effects—that is, a locus that causes *D. simulans* larvae to pupate farther from the

food surface. While this is certainly interesting, we did not pursue this region because its effect is

contrary to the species-wide difference we found. This locus may be interesting for further

investigation, however, as it could contribute to the significant variation in pupation behavior we

observed among *D. simulans* strains (Fig 2).

Fig 4. Hybrid deficiency screen of the X chromosome identifies two regions of interest. A. The
median pupation index for each of the 90 deficiency hybrid crosses (y-axis) is plotted by the physical map
distance each engineered deletion spans along the X chromosome (x-axis). Deficiencies with a pupation
index significantly greater than 1 (solid line) after correction for multiple comparisons are denoted by an
asterisk (significance levels for all deficiencies are listed in Table S2). Deficiencies with a median
pupation index significantly greater than 0.88 (the grand mean, dashed line) after correction for multiple
comparisons are denoted with a cross. All of these lines also had pupation indices significantly greater
than 1 before correcting for multiple comparisons. The two regions we pursued for candidate gene
validation are highlighted in light grey. Note, we did not pursue the remaining three significant deficiency
strains because they showed similar pupation indices when crossed to *D. melanogaster* (Fig S4). **B.** The
pupation indices of the deficiencies from the grey highlighted areas in Part A are shown for the original
hybrid cross (*Lhr*) and for a cross to the T.4 *D. melanogaster* strain. BSC869 and Exel6255 were
additionally crossed to *D. simulans* strain Mex180 and *D. melanogaster* strain T.1. Asterisks denote a
significant difference between pupation indices of deficiency strains crossed to *D. melanogaster* and *D.
simulans* (* = p < 0.05, ** = p < 0.01). **C.** The region uncovered by the overlap of deficiencies BSC869
and ED6720 and the 23 genes contained within it. Direction of the gene transcript is denoted with an
arrow. The three genes with available disruption strains *(Fas2, mei9,* and *norpA*) are labelled. **D.** The
region uncovered by Exel6255 and the 28 genes contained within it. Direction of the gene transcript is
denoted with an arrow. The two genes with available disruption strains *(tilB* and *wap)* are labelled.

4. Gene knockouts and RNAi knockdown suggest that *Fas2* is involved in divergent pupation behavior

We tested knockouts of three candidate genes in our first region of interest: the overlap of
Df(1)BSC869 and DF(1)ED6720, excluding the region covered by DF(1)ED6727 (Fig 4C). We
found no significant difference in the pupation index obtained when we crossed the *mei-9^{A1}*
mutant allele to the *Lhr D. simulans* strain and the T.4 *D. melanogaster* strain (Wilcoxon test: n=
51-52, χ^2=0.37, p=0.54; Fig S5A), indicating that *mei9* is unlikely to be involved in pupation site
choice. Similarly, we found no significant differences in the pupation behavior of female hybrids
containing the *norpA^{36}* mutant allele and female hybrids containing a *norpA* rescue allele
(Wilcoxon test: n= 22-24, χ^2=3.10, p=0.08 Fig S6), with the knockout hybrids actually having a
slightly lower proportion of flies pupating on the food. These data suggest that gene knockouts
with expression in the larval nervous system do not, in general, increase the number of hybrids
pupating on the food.

In contrast, when we crossed the mutant allele $Fas2^{eb112}$ to *D. simulans* (*Lhr*) and *D.*

melanogaster (T.4), we found that $Fas2^{eb112}$ hybrids had a significantly higher pupation index

than the *D. melanogaster* knockouts (Wilcoxon test: n= 50-51, χ^2= 6.97, p= 0.0083; Fig 5A),

suggesting that *Fas2* may be involved in pupation site choice. To ensure this pattern is not

unique to these strains, we crossed $Fas2^{eb112}$ to additional *D. simulans* (Mex180) and *D.*

melanogaster (T.1) wild-type strains. We again found the same pattern: the pupation index for

knockout hybrids was significantly higher than for *D. melanogaster* knockouts (Wilcoxon test:

n= 52-53, χ^2= 27.2, p<0.0001; Fig 5A). As further verification, we tested a second *Fas2* strain: a

p-element insertion allele, $Fas2^{G0293}$, and similarly found that the pupation index for $Fas2^{G0293}$

hybrids (crossed to *Lhr*) was significantly higher than that for *D. melanogaster* knockouts

(crossed to T.4; Wilcoxon test: n= 52-53, χ^2= 9.73, p=0.0018; Fig 5B). Although the nature of

the lesion is uncharacterized for $Fas2^{G0293}$, these results suggest that it is indeed a loss of

function allele, as it behaves indistinguishably from the verified knockout $Fas2^{eb112}$. We used the

consensus combined p-value test (Rice, 1990), which tests the combined effect of independent

tests of the same hypothesis, to look at the overall pattern for $Fas2^{eb112}$, and *Fas2* as a whole (i.e.

including results from both $Fas2^{eb112}$ and $Fas2^{G0293}$), and found a strongly significant pattern of

higher pupation indices for hybrid crosses compared to *D. melanogaster* crosses ($Fas2^{eb112}$: p=

3.59×10^{-6}; *Fas2*: p= 2.35×10^{-8}).

Next, we used RNAi with the elav-Gal4 driver to reduce expression of *Fas2* throughout

the nervous system in *D. melanogaster*. We found that a significantly higher proportion of *Fas2*

RNAi flies pupated on the food compared to the control flies from either the background

(Wilcoxon test: n= 42-48, p<0.0001 after sequential Bonferroni correction) or Gal4-1 cross

(Wilcoxon test: n= 42-43, p<0.0001 after sequential Bonferroni correction; Fig 5C); these results

are unchanged when we control for density effects (Fig S7A). Similarly, when we compared

pupation height, we found that RNAi flies pupated significantly closer to the food compared to

both the background (Wilcoxon test: n=42-48, p<0.0001 after sequential Bonferroni correction)

and Gal-4 crosses (Wilcoxon test: n= 42-43, p<0.0001 after sequential Bonferroni correction; Fig

S8A), providing further evidence for *Fas2*'s role in pupation site choice.

Fig 5. Knockouts and RNAi knockdown confirm the role of *Fas2* in evolved differences in pupation site choice. A. The pupation indices of the *Fas2eb112* gene disruption are shown for the comparison between the original hybrid cross (*Lhr*) and a cross to the *D. melanogaster* T.4 strain. Also shown is the comparison for crosses to the *D. simulans* strain Mex180 and the *D. melanogaster* strain T.1. **B.** The pupation indices for a second gene disruption, *Fas2G0293*, are shown for the original hybrid cross (*Lhr*) and a *D. melanogaster* strain (T.4). For A and B, asterisks denote significance (** = p < 0.01, **** = p<0.0001; N = 50-53). **C.** The results of pan-neuronal knockdown of the *Fas2* transcript via RNAi. The proportion of RNAi/control individuals found on the food are shown for the control cross (elav-Gal4 driver crossed to the RNAi background stock), the Gal-4 hairpin RNA cross (Gal-4), and *Fas2* RNAi cross. For C, asterisks denote significance after correcting for multiple comparisons (**** = p<0.0001; N = 42-48).

5. Gene knockouts and RNAi knockdown suggest that *tilB* is involved in divergent pupation

behavior

We tested knockouts of two candidate genes in our second region of interest: DF(1)Exel6255

(Fig 4D). We found no significant difference in the pupation index obtained when we crossed the

wap² mutant allele to the *Lhr D. simulans* strain and the T.4 *D. melanogaster* strain (Wilcoxon

test: n= 51-55, χ^2=0.32, p=0.57; Fig S5B), indicating that *wap* is unlikely to be involved in

pupation site choice. In contrast, when we crossed the *tilB¹* and *tilB²* mutant alleles to *D.*

simulans (*Lhr*) and *D. melanogaster* (T.4), we found that the *tilB* knockout hybrids had

significantly higher pupation indices than the *D. melanogaster* knockouts for both alleles

(Wilcoxon tests; *tilB¹*: n= 56, χ^2= 6.61, p= 0.0101; Fig 6A; *tilB²*: n= 57, χ^2= 6.61, p= 0.0101; Fig

6B). To test whether this difference is consistent in other backgrounds, we crossed both the *tilB¹*

and *tilB²* mutant alleles to additional *D. simulans* (Mex180) and *D. melanogaster* (T.1) wild-type

strains. We had difficulty crossing our *tilB* strains to the Mex180 strain, so our sample sizes for

these crosses are smaller, but there is a nonsignificant trend towards a higher pupation index for

the knockout hybrids compared to the *D. melanogaster* hybrids for both alleles (Wilcoxon tests;

tilB¹: n=36-37, χ^2= 2.80, p= 0.0943, Fig 6A; *tilB²*: n=12-18, χ^2= 3.44, p= 0.0638, Fig 6B). We

used the combined consensus p-value test (Rice, 1990) to look at the overall pattern for *tilB¹*,

tilB², and *tilB* as a whole (i.e. including results from both *tilB¹* and *tilB²*), and found a strongly

significant pattern of higher pupation indices for hybrid crosses compared to *D. melanogaster*

crosses (*tilB¹*: p= 0.0022; *tilB²*: p= 0.0037; *tilB*: p= 2.47 x 10^{-5}).

As for *Fas2* above, we then used RNAi with the elav-Gal4 driver to reduce expression of

tilB throughout the nervous system in *D. melanogaster*. We found that a significantly higher

proportion of RNAi flies pupated on the food compared to the control flies from either the

background (Wilcoxon test: n= 46-47, p<0.01 after sequential Bonferroni correction) or Gal4-1

cross (Wilcoxon test: n= 41-46, p<0.01 after sequential Bonferroni correction; Fig 6C); these

findings are consistent when we control for density effects (Fig S7B). In addition, when we

compared pupation height, we found that RNAi flies pupated significantly closer to the food

compared to both the background (Wilcoxon test: n=46-47, p<0.0001 after sequential Bonferroni

correction) and Gal-4 crosses (Wilcoxon test: n= 46-47, p<0.0001 after sequential Bonferroni

correction; Fig S8B), providing additional support for *tilB*'s role in pupation site choice.

Fig 6. Knockouts and RNAi knockdown confirm the role of *tilB* in evolved differences in pupation site choice. A. The pupation indices of the *tilB1* gene disruption are shown for the comparison between the original hybrid cross (*Lhr*) and a cross to the *D. melanogaster* T.4 strain. Also shown is the comparison for crosses to the *D. simulans* strain Mex180 and the *D. melanogaster* strain T.1. **B.** The pupation indices for a second gene disruption, *tilB2*, are shown for the original hybrid cross (*Lhr*) and the cross to *D. melanogaster* strain T.4, and for crosses to the *D. simulans* strain Mex180, and the *D. melanogaster* strain T.1. For A and B, asterisks denote significance (* = p < 0.05, ** = p = 0.01; N = 41-47). **C.** The results of pan-neuronal knockdown of the *tilB* transcript via RNAi. The proportion of RNAi/control individuals found on the food are shown for the control cross (elav-Gal4 driver crossed to the RNAi background stock), the Gal-4 hairpin RNA cross (Gal-4), and *tilB* RNAi cross. For C, asterisks denote significance after correcting for multiple comparisons (* = p < 0.05, ** = p < 0.01; N = 41-47).

6. *tilB* is expressed more highly in *D. melanogaster* strains

We performed qRT-PCR to quantify relative *tilB* transcript abundance for two strains of *D. simulans* (Per005 and Geo288) and two strains of *D. melanogaster* (CA1 and T.4). For the two *D. simulans* strains and the Geo288 *D. melanogaster* strain, we collected larvae from two stages of larval development (96 and 120 hours following oviposition). For the other *D. melanogaster* strain (T.4), we were only able to obtain enough larval tissue at 96 hours following oviposition, due to low fecundity. Because the relative transcript abundance data had a skewed distribution, we used the reciprocal root transformation to normalize the species and overall distributions (Shapiro-Wilk tests for normality: all p >0.34). We then analyzed the relative transcript abundance of *tilB* using a nested ANOVA with the following factors: species, strain nested within species, larval age (96 and 120 hours), and the interaction between species and larval age. The interaction term between species and larval age was not significant (p=0.41), so we removed it from the model. We found that larvae from the 120-hour sampling period had significantly lower *tilB* expression than 96-hour larvae ($F_{1,14}$= 6.16, p= 0.026). While we did not detect any significant differences between the 2 strains from the same species ($F_{2,14}$=2.39, p=0.13), we found a significantly higher average relative amount of *tilB* transcript in *D. melanogaster* larvae compared to *D. simulans* larvae ($F_{1,14}$= 9.74, p= 0.0075; Fig 7).

Because we performed qRT-PCR on an extreme strain and a strain closer to the average for each species, these four lines represent a continuum of pupation site choice behavior, with T.4 (*D. melanogaster*) having the lowest proportion of pupae on the food, followed by CA1 (*D. melanogaster*), then Per005 (*D. simulans*), and last, Geo288 (*D. simulans*) having the highest proportion of pupae on the food (Fig 2). These four strains follow an identical pattern for *tilB*

gene expression, with T.4 having the highest relative transcript abundance, and Geo288 having

the lowest (Fig 7). Although it is not possible to detect a significant effect with a sample size of 4,

this suggests that *tilB* gene expression may be negatively correlated with the proportion of pupae

on the food (Spearman's rank correlation: $r_s = -1$, $p < 0.10$).

Fig 7. Relative differences in *tilB* transcript abundance between species. The relative abundance of
tilB transcript detected by qRT-PCR in *D. melanogaster* and *D. simulans.* Each data point represents the
average of two technical replicates for a single biological replicate. Relative transcript abundance is
significantly higher in *D. melanogaster* strains on average ($p < 0.01$). Light grey squares (T.4) and
triangles (CA1) represent the two *D. melanogaster* strains measured, while dark grey diamonds (Per005)
and circles (Geo288) represent the two *D. simulans* strains. The dashed lines depict the average
expression across both time points for each strain. The black circles represent the species-wide mean,
and error bars depict the 95% confidence interval surrounding the mean. Note that the exact rank of
relative *tilB* expression is perfectly inversely correlated to the pupation indices of each strain (i.e., the
strain with the highest *tilB* expression has the lowest proportion of pupae on the food, and so on).

7. Effect sizes

We used the results above to estimate how much of the difference in pupation site preference

between *D. melanogaster* and *D. simulans* can be attributed to the X chromosome, and found

that the X accounts for approximately 55.6% (95% CI= 31.4%-80.2%) of the difference. We then

estimated the effect size of our deficiencies of interest, and found that the overlap of

Df(1)BSC869 and DF(1)ED6720 explains approximately 44.1% (95% CI= 33.5%-74/5%) of the

X chromosome effect, and Df(1)Exel6255 explains approximately 52.3% (95% CI= 33.7%-

85.9%) of the X effect. Finally, we estimated the effect size of our two identified candidate genes,

Fas2 and *tilB*. We found that *Fas2* explains approximately 93% (95% CI= 78.5%-134.9%) of the

pupation difference attributed to the overlap between Df(1)BSC869 and DF(1)ED6720, and 41%

(95% CI= 33.8%-76.5%) of the pupation difference attributed to the X chromosome. Similarly,

we found the *tilB* explains approximately 87.2% (95% CI= 66.2%-135.9%) of the pupation

difference attributed to Df(1)Exel6255, and 45.6% (95% CI= 35.5%-76.1%) of the pupation

difference attributed to the X chromosome.

It is important to note that our effect size estimate for the X chromosome may be an

overestimate, as calculating effect sizes using only reciprocal hybrids does not account for

potential transgressive autosomal effects (Mittleman et al., 2017). While subject to the same

potential effects, the effect size of the deficiencies and individual gene knockouts were

calculated as proportions of the total X effect (and also of the deficiency effect for gene

knockouts), and thus represent a more accurate estimate of their overall contribution, regardless

of the total X effect. Still, these estimates could be affected by variation in the different *D.*

melanogaster backgrounds in which the deficiencies and knockout strains were made.

Discussion

A species-level difference in pupation site choice behavior

Our initial survey of pupation site choice behavior in *D. melanogaster* and *D. simulans* expands

upon a previously reported interspecific difference (Markow, 1979). Consistent with these

previous results, on average, *D. simulans* strains had a greater proportion of flies pupating on the

food surface. However, we used 11 *D. melanogaster* and 12 *D. simulans* strains sourced from

around the globe to demonstrate this difference. While the species difference holds when

comparing the grand mean of all strains for each species, there is substantial variation within

species. This variation is so significant that the species' distributions overlap, with some *D.*

simulans strains, like Mex180 and Cal006, more closely resembling *D. melanogaster* strains in

pupation behavior (Fig 2). These documented differences in pupation behavior among

Drosophila species, in combination with our understanding of the environmental variables that

affect this behavior within species (Hodge & Caslaw, 1998; Sameoto & Miller, 1968b;

Seyahooei et al., 2009; Sokal et al., 1960), will be useful in identifying the selection pressures (if

any) that affect the evolution of this trait. Differences in pupation site choice behavior may be a

form a niche partitioning where species co-occur, as has been previously suggested (Arthur &

Middlecote, 1984). Alternatively, pupation site choice may be an adaptive response to parasite or

parasitoid presence (Kraaijeveld & Godfray, 2003). A globally sourced panel of lines with

significant variation, such as we describe, provides an inroad for studies comparing pupation

behavior to differences in the ecology of each collection site, such that we can better understand

the ultimate causes of this behavioral evolution.

The genetic architecture of pupation site choice behavioral evolution

Using hybrid crosses in the same background that controlled for maternal inheritance, we were

able to estimate the effect of the X chromosome on pupation behavior. We found no evidence

that hybrid behavior was atypical relative to either parent strain, suggesting these measurements

are indeed reliable. We found a significant effect of the X, in that reciprocal hybrid males pupate

in similar locations as the X-donating parent. We calculate that this chromosome explains 55.6%

(95% CI= 31.4%-80.2%) of the total phenotypic difference between parent strains. Although the

exact contribution of the X chromosome reported here may be an overestimate due to

transgressive autosomal effects that cannot be detected in a hybrid background (Mittleman et al.,

2017), a similar X-effect has been detected for pupation behavior when comparing *D. simulans*

and *D. sechellia* (Erezyilmaz & Stern, 2013).

We also found that hybrid females, which inherit one X chromosome from each parent,

pupate like the *D. melanogaster* parent strain. This suggests that *D. melanogaster* alleles are

dominant to *D. simulans* alleles. The dominance of *D. melanogaster* alleles makes it possible to

use engineered deletions, available in *D. melanogaster* strains, to map regions containing

recessive *D. simulans* variation affecting pupation behavior (Cook et al., 2012).

The three significant deficiencies identify two regions of the X chromosome with

detectable effects on pupation behavior: one spans X:4,204,351 – 4,325,174 and explains ~44.1%

of the X-effect, and the other spans X:21,519,203 – 22,517,665 and explains ~52.3% of the X-

effect. These regions contain 23 and 28 genes, respectively. Our analysis of the available gene

knockouts within these regions points to two genes: *tilB* and *Fas2* (see below). We calculate that

tilB and *Fas2* explain the majority of the effect of their respective deficiency regions (point

estimate = 87.2%, 95% CI= 66.2%-135.9%; point estimate = 93%, 95% CI= 78.5%-134.9%,

respectively). Taken together, our results suggest that a substantial share of the difference in

pupation behavior between *D. simulans* and *D. melanogaster* can be attributed to just two genes.

tilB* and *Fas2*: loci of evolution for divergent pupation behavior between *D. simulans* and *D.

melanogaster

We have presented substantial evidence for a role of both *tilB* (*touch insensitive larva B*) and

Fas2 (*Fasciclin* 2) in the divergence of pupation behavior among these species. For each locus,

we have shown that two independent knockouts replicate the pattern of the regions identified by

the deficiency screen: hybrid females hemizygous for the *D. simulans* locus pupate on the food

surface significantly more often than females with both the *D. simulans* and *D. melanogaster* loci.

We then used RNAi knockdown of each gene in *D. melanogaster* to show that reduced

expression of *tilB* and *Fas2* transcripts results in a more *D. simulans*-like pupation site choice

behavior. To directly test for differences in expression of *tilB*, we performed qRT-PCR during

two larval developmental time points. Congruent with our RNAi knockdown results, we find that

mean relative transcript abundance is significantly higher in *D. melanogaster* larvae than in *D.*

simulans larvae. Interestingly, we surveyed two lines per species, representing a continuum of

pupation site choice behavior, and found that the mean proportion of pupae on the food surface

for these lines was perfectly negatively correlated with *tilB* gene expression. Although these

results are only for 4 lines, they suggest that lower *tilB* gene expression may be associated with a

higher proportion of larvae pupating on the food surface.

While our present study does not present a functional analysis of the *D. melanogaster* or

D. simulans Fas2 or *tilB* alleles, we can use the *D. melanogaster* annotation of each gene to

speculate about their role in the evolution of pupation behavior. *Fas2* is a large gene, spanning

over 70,000 base pairs, with expression peaking during the larval wandering stage (L3)

(Graveley et al., 2011). It is also complex, with seven transcripts composed of various

combinations of 16 exons. Broadly, *Fas2* functions as a neuronal recognition molecule, and is

involved in patterning the larval nervous system (Grenningloh et al., 1991). Expression of *Fas2*

is critical for synapse formation and growth at the larval neuromuscular junction (Davis et al.,

1997; Schuster et al., 1996) and is also important for patterning of the larval mushroom body

(Kurusu et al., 2002). With expression in both the central and peripheral nervous system, it is

possible that differences at the *Fas2* locus differentially wire the *D. melanogaster* and *D.*

simulans brains, altering how larvae perceive or interpret stimuli. Whether these differences are a

result of evolution of the protein sequence, and/or spatial or temporal differences in transcript

expression remains to be determined.

Unlike *Fas2*, *tilB* is a short gene, spanning just over 1,700 base pairs, with a single

transcript composed of 5 exons. *tilB* is also expressed in wandering larvae and pupae, though it

shows higher expression in testes of adult males, due to its role in developing sperm flagella

(Graveley et al., 2011). In fact, *tilB* is associated with ciliary motility (Kavlie et al., 2010), and is

a part of the mechanosensory transduction machinery (Göpfert & Robert, 2003). Mutant *tilB*

larvae display normal locomotor activity, but have a reduced withdrawal response to physical

disturbance (Kernan et al., 1994). Consistent with this finding, our data suggest that changes in

tilB expression could potentially result in differences in peripheral sensory perception between *D.*

melanogaster and *D. simulans* larvae, ultimately influencing larval pupation site choice behavior.

A more precise functional analysis of *tilB* is necessary to test this hypothesis.

Standards of evidence and the challenges of interspecific mapping

Above, we have discussed the results of our interspecific deficiency screen of the

Drosophila X chromosome. This technique has long been employed to map morphological traits,

physiological traits, hybrid incompatibility loci, (Barbash et al., 2003; Bour et al., 2000; Cattani

& Presgraves, 2012; Cote et al., 1986; Konopka & Benzer, 1971; Pardy et al., 2018; Sawamura

et al., 2004), and behaviors (Fanara et al., 2002; Laturney & Moehring, 2012; Moehring &

Mackay, 2004). Nonetheless, these screens have been criticized due to their susceptibility to

epistatic interactions that can produce false positives (Anholt & Mackay, 2004) and for the

imperfect comparison of deficiency chromosomes to balancer chromosomes (Stern, 2014). These

problems may be exacerbated in a hybrid background. These issues are potentially reflected by

the global average pupation index for these deficiencies being 0.88, rather than 1 (Fig 4A),

indicating that, on average, more balancer hybrids pupated on the food compared to deficiency

hybrids.

We attempted to exclude the possibility of false positives in our initial deficiency screen

in two ways. First, we crossed each deficiency with a significant pupation index to *D.*

melanogaster to test for deleterious effects of the deficiencies themselves. Indeed, 3 of 6

significant deficiencies produced similar results when crossed to *D. melanogaster*, and were

discarded. Next, to ensure the patterns shown by the remaining three were not due to epistasis

among strains, we crossed each to a second *D. simulans* and *D. melanogaster* strain and found

consistent results. Taken together, these results suggest that these three deficiencies are likely

revealing recessive *D. simulans* variation, rather than background epistatic interactions. It should

be noted, however, that species-wide epistasis cannot be conclusively ruled out; nonetheless,

such a result is still biologically relevant and meaningful.

The results of our gene disruption tests further enforce our deficiency screen results,

again, with multiple controls to limit the possibility of false positives and/or strain-specific

epistatic effects. We tested *D. melanogaster* gene disruption strains in a hybrid background for

two or three potential candidate genes that are expressed in the larval nervous system from each

region identified by our deficiency screen. For each region, only a single candidate gene (*Fas2*

and *tilB*) replicated the deficiency effect, suggesting that disrupting larval nervous system genes

in a hybrid background does not generally interfere with pupation behavior. Further, we crossed

our *tilB* and *Fas2* gene disruption strains to multiple *D. melanogaster* and *D. simulans* strains,

and found no evidence of strain-specific epistatic interactions. Finally, we tested two knockouts

of each candidate gene to ensure that our results were not specific to the gene aberration.

In addition to our hybrid mapping experiments, we have shown that a reduction in *Fas2*

and *tilB* gene expression via RNAi knockdown in a *D. melanogaster* background leads to a more

D. simulans-like phenotype. This test is performed in a pure *D. melanogaster* background,

eliminating the potential for hybrid background epistasis, and the results are consistent with our

gene disruption screen. Additionally, these results hold true when calculating both the proportion

of flies pupating on the food surface, and the average pupation height. This result is even further

corroborated for *tilB*, which shows higher relative expression in *D. melanogaster* compared to *D.*

simulans strains (Figure 7). Taken together, our hybrid gene disruption screens and gene

expression studies in *D. melanogaster* make an excellent case for a role of *tilB* and *Fas2* in

divergent pupation site choice behavior.

Higher standards of evidence have been suggested for mapping in *Drosophila* – the

reciprocal hemizygosity test being the gold standard (Stern, 2014). In this study, we performed

one half of this test (hybrid females with a disrupted *D. melanogaster* X chromosome and an

intact *D. simulans* X chromosome). Unfortunately, the reciprocal half of the test (hybrid females

with a disrupted *D. simulans* X chromosome and an intact *D. melanogaster* X chromosome) is

not possible in this case for several reasons. The hybrid cross is only successful in one direction

(*D. simulans* males crossed to *D. melanogaster* females), so we would need to use *D. simulans*

gene disruption males to create the reciprocal cross. This is not an option, however, as *tilB*

disruptions are male sterile, and *Fas2* disruptions are male lethal.

A test that could potentially support our findings would be the addition of dominant *D.*

melanogaster Fas2 and tilB alleles to a *D. simulans* X background to recover the *D.*

melanogaster behavior of pupating off the food surface. We previously performed a similar

experiment in males using *D. melanogaster* Y-linked X duplication strains (Table S6) (Cook et

al., 2010). Unfortunately, making hybrid males that have two broad segments of the X

chromosome created flies that pupated almost entirely on the surface of the food (Fig S9). A

similar experiment could be performed in females using third chromosome-linked X duplication

strains (Venken et al., 2010), however this is likely to produce a similar effect. Further, these

experiments are prone to the same potential epistatic interactions as the deficiencies.

While we have presented multiple lines of evidence, all consistently supporting a role for

tilB and *Fas2* in divergent pupation site choice behavior, these results must be considered in light

of the above caveats. We have taken measures to address these caveats as completely as possible,

and present our results with a high standard of evidence. Our study highlights the difficulty of

interspecific mapping in producing conclusive results, and underscores a need for transgenic

tools to be developed in non-model *Drosophila* species.

Areas for future research

Our results highlight two main areas for further research. First, while we now have a

better understanding of the genetic underpinnings of pupation site choice behavior evolution, it is

unclear why these differences evolved in the first place. The variation we recorded among

globally sourced *D. melanogaster* and *D. simulans* strains provides a valuable tool with which to

pursue this line of inquiry. Second, while we have provided evidence for two specific genes

involved in the evolution of this phenotype, we have little understanding of their function in

pupation site choice behavior, or how exactly these functions have evolved. The results of our

expression study suggest a role of expression differences for *tilB*, but further functional follow-

up is necessary to identify the precise molecular underpinnings of pupation site choice

preference.

915 Acknowledgments

We would like to thank Alexandra Lamoureux, Susanne Tilk, Nastaran Hosseinipour, Sayeh

Akhavan, Wesley Cochrane and Veronica Cochrane for their assistance in the collection of

pupation preference data. We also thank Daniel Eberl, Stuart Macdonald, Brian McCabe, and

William Rice for providing fly stocks used for this study. This work would not have been

possible without the community-supported resources available at the Cornell *Drosophila* Stock

Center (CSBR 1820594), the Bloomington *Drosophila* stock center (NIH P400D018537), or

FlyBase (flybase.org; Gramates et al., 2017). This work was funded by the National Institutes of

Health (R01 GM098614).

Literature Cited

Al-Atia, G. R., Fruscoloni, P., & Jacobs-Lorena, M. (1985). Translational Regulation of mRNAs for Ribosomal Proteins during Early Drosophila Development†. *Biochemistry*, *24*(21), 5798–5803. https://doi.org/10.1021/bi00342a017

Anholt, R. R. H., & Mackay, T. F. C. (2004, November 1). Quantitative genetic analyses of complex behaviours in Drosophila. *Nature Reviews Genetics*. Nature Publishing Group. https://doi.org/10.1038/nrg1472

Arthur, W., & Middlecote, J. (1984). Evolution of pupation site and interspecific competitive ability in Drosophila hydei. *Biological Journal of the Linnean Society*, *23*(4), 343–352. https://doi.org/10.1111/j.1095-8312.1984.tb00150.x

Barbash, D. A., & Ashburner, M. (2003). A novel system of fertility rescue in Drosophila hybrids reveals a link between hybrid lethality and female sterility. *Genetics*, *163*(1), 217–226.

Barbash, D. A., Siino, D. F., Tarone, A. M., & Roote, J. (2003). A rapidly evolving MYB-related protein causes species isolation in Drosophila. *Proceedings of the National Academy of Sciences of the United States of America*, *100*(9), 5302–7. https://doi.org/10.1073/pnas.0836927100

Bour, B. A., Chakravarti, M., West, J. M., & Abmayr, S. M. (2000). Drosophila SNS, a member of the immunoglobulin superfamily that is essential for myoblast fusion. *Genes & Development*, *14*(12), 1498–511. https://doi.org/10.1101/GAD.14.12.1498

Brideau, N. J., Flores, H. A., Wang, J., Maheshwari, S., Wang, X., & Barbash, D. A. (2006a). Two Dobzhansky-Muller genes interact to cause hybrid lethality in Drosophila. *Science*, *314*(5803).

Brideau, N. J., Flores, H. A., Wang, J., Maheshwari, S., Wang, X., & Barbash, D. A. (2006b).

Two Dobzhansky-Muller Genes interact to cause hybrid lethality in Drosophila. *Science*,

314(5803), 1292–1295. https://doi.org/10.1126/science.1133953

Cande, J., Prud'homme, B., & Gompel, N. (2013). Smells like evolution: The role of

chemoreceptor evolution in behavioral change. *Current Opinion in Neurobiology*.

https://doi.org/10.1016/j.conb.2012.07.008

Cattani, V. M., & Presgraves, D. C. (2012). Incompatibility between X chromosome factor and

pericentric heterochromatic region causes lethality in hybrids between drosophila

melanogaster and its sibling species. *Genetics*, *191*(2), 549–559.

https://doi.org/10.1534/genetics.112.139683

Chertemps, T., Duportets, L., Labeur, C., Ueda, R., Takahashi, K., Saigo, K., & Wicker-Thomas,

C. (2007). A female-biased expressed elongase involved in long-chain hydrocarbon

biosynthesis and courtship behavior in Drosophila melanogaster. *Proceedings of the*

National Academy of Sciences of the United States of America, *104*(11), 4273–8.

https://doi.org/10.1073/pnas.0608142104

Cook, R. K., Christensen, S. J., Deal, J. A., Coburn, R. A., Deal, M. E., Gresens, J. M., … Cook,

K. R. (2012). The generation of chromosomal deletions to provide extensive coverage and

subdivision of the Drosophila melanogaster genome. *Genome Biology*, *13*(3), R21.

https://doi.org/10.1186/gb-2012-13-3-r21

Cook, R. K., Deal, M. E., Deal, J. A., Garton, R. D., Brown, C. A., Ward, M. E., … Cook, K. R.

(2010). A new resource for characterizing X-linked genes in Drosophila melanogaster:

Systematic coverage and subdivision of the X chromosome with nested, Y-linked

duplications. *Genetics*, *186*(4), 1095–1109. https://doi.org/10.1534/genetics.110.123265

Cote, B., Bender, W., Curtis, D., & Chovnickt, A. (1986). Molecular mapping of the rosy locus

in Drosophila melanogaster. *Genetics*, *112*, 769–783.

Coyne, J. A., & Orr, H. A. (1997). Patterns of Speciation in Drosophila Revisited. *Evolution*,

51(1), 295–303.

Coyne, J. A., & Orr, H. A. (2004). *Speciation*. Sinauer Associates. Retrieved from

https://books.google.com/books/about/Speciation.html?id=2Y9rQgAACAAJ

Davis, G. W., Schuster, C. M., & Goodman, C. S. (1997). Genetic analysis of the mechanisms

controlling target selection: Target- derived Fasciclin II regulates the pattern of synapse

formation. *Neuron*, *19*(3), 561–573. https://doi.org/10.1016/S0896-6273(00)80372-4

Erezyilmaz, D. F., & Stern, D. L. (2013). Pupariation site preference within and between

drosophila sibling species. *Evolution*, *67*(9), 2714–2727. https://doi.org/10.1111/evo.12146

Fanara, J. J., Robinson, K. O., Rollmann, S. M., Anholt, R. R. H., & Mackay, T. F. C. (2002).

Vanaso is a candidate quantitative trait gene for Drosophila olfactory behavior. *Genetics*,

162(3), 1321–1328. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/12454076

Göpfert, M. C., & Robert, D. (2003). Motion generation by Drosophila mechanosensory neurons.

Proceedings of the National Academy of Sciences of the United States of America, *100*(9),

5514–5519. https://doi.org/10.1073/pnas.0737564100

Gramates, L. S., Marygold, S. J., Dos Santos, G., Urbano, J. M., Antonazzo, G., Matthews, B.

B., … Baker, P. (2017). FlyBase at 25: Looking to the future. *Nucleic Acids Research*,

45(D1), D663–D671. https://doi.org/10.1093/nar/gkw1016

Graveley, B. R., Brooks, A. N., Carlson, J. W., Duff, M. O., Landolin, J. M., Yang, L., …

Celniker, S. E. (2011). supplements - The developmental transcriptome of Drosophila

melanogaster. *Nature*, *471*(7339), 473–9. https://doi.org/10.1038/nature09715

Grenningloh, G., Jay Rehm, E., & Goodman, C. S. (1991). Genetic analysis of growth cone

guidance in drosophila: Fasciclin II functions as a neuronal recognition molecule. *Cell*,

67(1), 45–57. https://doi.org/10.1016/0092-8674(91)90571-F

Hodge, S., & Caslaw, P. (1998). The Effect of Resource pH on Pupation Height in Drosophila

(Diptera: Drosophilidae). *Journal of Insect Behavior*, *11*(1), 47–57.

https://doi.org/10.1023/A:1020814516158

Huggett, J., Dheda, K., Bustin, S., & Zumla, A. (2005, June 7). Real-time RT-PCR normalisation;

strategies and considerations. *Genes and Immunity*. Nature Publishing Group.

https://doi.org/10.1038/sj.gene.6364190

Karageorgi, M., Bräcker, L. B., Lebreton, S., Minervino, C., Cavey, M., Siju, K. P., …

Prud'homme, B. (2017). Evolution of Multiple Sensory Systems Drives Novel Egg-Laying

Behavior in the Fruit Pest Drosophila suzukii. *Current Biology*, *27*(6), 847–853.

https://doi.org/10.1016/j.cub.2017.01.055

Kavlie, R. G., Kernan, M. J., & Eberl, D. F. (2010). Hearing in drosophila requires TilB, a

conserved protein associated with ciliary motility. *Genetics*, *185*(1), 177–188.

https://doi.org/10.1534/genetics.110.114009

Kernan, M., Cowan, D., & Zuker, C. (1994). Genetic dissection of mechanosensory transduction:

Mechanoreception-defective mutations of drosophila. *Neuron*, *12*(6), 1195–1206.

https://doi.org/10.1016/0896-6273(94)90437-5

King, E. G., Merkes, C. M., McNeil, C. L., Hoofer, S. R., Sen, S., Broman, K. W., … Macdonald,

S. J. (2012). Genetic dissection of a model complex trait using the Drosophila Synthetic

Population Resource. *Genome Research*, *22*(8), 1558–1566.

https://doi.org/10.1101/gr.134031.111

Kittelmann, S., Buffry, A. D., Almudi, I., Yoth, M., Sabaris, G., Franziska, A., … Pueyo-, J.

(2017). Gene regulatory network architecture in different developmental contexts influences

the genetic basis of morphological evolution. *PLOS Genetics*, *14*(5), 1–20.

https://doi.org/10.1371/journal.pgen.1007375

Konopka, R. J., & Benzer, S. (1971). Clock mutants of Drosophila melanogaster. *Proceedings of*

the National Academy of Sciences of the United States of America, *68*(9), 2112–6.

https://doi.org/10.1073/pnas.68.9.2112

Kopp, A. (2009). Metamodels and phylogenetic replication: A systematic approach to the

evolution of developmental pathways. *Evolution*, *63*(11), 2771–2789.

https://doi.org/10.1111/j.1558-5646.2009.00761.x

Kraaijeveld, A. R., & Godfray, H. C. J. (2003). Potential life-history costs of parasitoid

avoidance in Drosophila melanogaster. *Evolutionary Ecology Research*, *5*(8), 1251–1261.

Retrieved from http://www.evolutionary-ecology.com/issues/v05n08/kkar1615.pdf

Kurusu, M., Awasaki, T., Masuda-Nakagawa, L. M., Kawauchi, H., Ito, K., & Furukubo-

Tokunaga, K. (2002). Embryonic and larval development of the Drosophila mushroom

bodies: concentric layer subdivisions and the role of fasciclin II. *Development (Cambridge,*

England), *129*(2), 409–419. Retrieved from

http://www.ncbi.nlm.nih.gov/pubmed/11807033

Lachaise, D., & Silvain, J. F. (2004). How two Afrotropical endemics made two cosmopolitan

human commensals: The Drosophila melanogaster-D. simulans palaeogeographic riddle. In

Genetica (Vol. 120, pp. 17–39). https://doi.org/10.1023/B:GENE.0000017627.27537.ef

Laturney, M., & Moehring, A. J. (2012). Fine-scale genetic analysis of species-specific female

preference in Drosophila simulans. *Journal of Evolutionary Biology*, *25*(9), 1718–1731.

https://doi.org/10.1111/j.1420-9101.2012.02550.x

Leary, G. P., Allen, J. E., Bunger, P. L., Luginbill, J. B., Linn, C. E., Macallister, I. E., …

Wanner, K. W. (2012). Single mutation to a sex pheromone receptor provides adaptive

specificity between closely related moth species. *Proceedings of the National Academy of*

Sciences, *109*(35), 14081–14086. https://doi.org/10.1073/pnas.1204661109

Markow, T. A. (1979). A survey of intra- and interspecific variation for pupation height in

Drosophila. *Behavior Genetics*, *9*(3), 209–217. https://doi.org/10.1007/BF01071301

Markow, T. A. (1981a). Light-dependent pupation site preferences in Drosophila: Behavior of

adult visual mutants. *Behavioral and Neural Biology*, *31*(3), 348–353.

https://doi.org/10.1016/S0163-1047(81)91409-6

Markow, T. A. (1981b). Light-dependent pupation site preferences in Drosophila: Behavior of

adult visual mutants. *Behavioral and Neural Biology*, *31*(3), 348–353.

https://doi.org/10.1016/S0163-1047(81)91409-6

Martin, A., & Orgogozo, V. (2013). The loci of repeated evolution: A catalog of genetic hotspots

of phenotypic variation. *Evolution*, *67*(5), 1235–1250. https://doi.org/10.1111/evo.12081

McBride, C. S., Baier, F., Omondi, A. B., Spitzer, S. A., Lutomiah, J., Sang, R., … Vosshall, L.

B. (2014). Evolution of mosquito preference for humans linked to an odorant receptor.

Nature, *515*(7526), 222–227. https://doi.org/10.1038/nature13964

McElhany, P., Real, L. A., & Power, A. G. (1995). Vector preference and disease dynamics: A

study of barley yellow dwarf virus. *Ecology*, *76*(2), 444–457.

https://doi.org/10.2307/1941203

McGrath, P. T., Xu, Y., Ailion, M., Garrison, J. L., Butcher, R. A., & Bargmann, C. I. (2011).

Parallel evolution of domesticated Caenorhabditis species targets pheromone receptor genes.

Nature, *477*(7364), 321–325. https://doi.org/10.1038/nature10378

Mittleman, B. E., Manzano-Winkler, B., Hall, J. B., Korunes, K. L., & Noor, M. A. F. (2017).

The large X-effect on secondary sexual characters and the genetics of variation in sex comb

tooth number in Drosophila subobscura. *Ecology and Evolution*, *7*(2), 533–540.

https://doi.org/10.1002/ece3.2634

Moehring, A. J., & Mackay, T. F. C. (2004). The Quantitative Genetic Basis of Male Mating

Behavior in Drosophila melanogaster. *Genetics*, *167*(3).

Mueller, L. D., & Sweet, V. F. (1986). Density-dependent natural selection in Drosophila:

evolution of pupation height. *Evolution*, *40*(6), 1354–1356. https://doi.org/10.2307/2408963

Nichols, C. D., Becnel, J., & Pandey, U. B. (2012). Methods to Assay Drosophila Behavior. *J.*

Vis. Exp, (61), 3795–103791. https://doi.org/10.3791/3795

Orgogozo, V., & Stern, D. L. (2009, April 15). How different are recently diverged species?:

More than 150 phenotypic differences have been reported for the D. melanogaster species

subgroup. *Fly*. Taylor & Francis. https://doi.org/10.4161/fly.8836

Pardy, J. A., Rundle, H. D., Bernards, M. A., & Moehring, A. J. (2018). The genetic basis of

female pheromone differences between Drosophila melanogaster and D. simulans. *Heredity*,

1. https://doi.org/10.1038/s41437-018-0080-3

Parsons, P. A. (1975). The Comparative Evolutionary Biology of the Sibling Species, Drosophila

melanogaster and D. Simulans. *The Quarterly Review of Biology*, *50*(2), 151–169.

https://doi.org/10.1086/408437

Perkins, L. A., Holderbaum, L., Tao, R., Hu, Y., Sopko, R., McCall, K., … Perrimon, N. (2015).

The transgenic RNAi project at Harvard medical school: Resources and validation. *Genetics*,

201(3), 843–852. https://doi.org/10.1534/genetics.115.180208

Peterson, J. (1977). *Behavioral Differences in Two Sub-species of Aedes Aegypti (L.)(diptera,*

Culicidae) in East Africa. University of Notre Dame.

Prince, D. J., O'Rourke, S. M., Thompson, T. Q., Ali, O. A., Lyman, H. S., Saglam, I. K., …

Miller, M. R. (2017). The evolutionary basis of premature migration in Pacific salmon

highlights the utility of genomics for informing conservation. *Science Advances*, *3*(8),

e1603198. https://doi.org/10.1126/sciadv.1603198

Rebeiz, M., & Williams, T. M. (2017, February 1). Using Drosophila pigmentation traits to study

the mechanisms of cis-regulatory evolution. *Current Opinion in Insect Science*. Elsevier.

https://doi.org/10.1016/j.cois.2016.10.002

Rice, W. R. (1990). A Consensus Combined P-Value Test and the Family-Wide Significance of

Component Tests. *Biometrics*, *46*(2), 303. https://doi.org/10.2307/2531435

Rice, W. R., Linder, J. E., Friberg, U., Lew, T. A., Morrow, E. H., & Stewart, A. D. (2005).

Inter-locus antagonistic coevolution as an engine of speciation: Assessment with hemiclonal

analysis. *Proceedings of the National Academy of Sciences*, *102*(Supplement 1), 6527–6534.

https://doi.org/10.1073/pnas.0501889102

Riedl, C. A. L., Sokolowski, M. B., Bodner, T., & Goldberg, R. G. (2007). Genetic and

Behavioral Analysis of Natural Variation in Drosophila melanogaster Pupation Position. *Fly*,

1(February), 23–32. https://doi.org/10.4161/fly.3830

Rockman, M. V. (2012). The QTN program and the alleles that matter for evolution: All that's

gold does not glitter. *Evolution*. https://doi.org/10.1111/j.1558-5646.2011.01486.x

Ryder, E., Blows, F., Ashburner, M., Bautista-Llacer, R., Coulson, D., Drummond, J., … Russell,

S. (2004). The DrosDel collection: A set of P-element insertions for generating custom

chromosomal aberrations in Drosophila melanogaster. *Genetics*, *167*(2), 797–813.

1110  https://doi.org/10.1534/genetics.104.026658

1111 Sameoto, D. D., & Miller, R. S. (1968a). Selection of Pupation Site by Drosophila Melanogaster

1112  and D. Simulans. *Ecology*, *49*(1), 177–180. https://doi.org/10.2307/1933580

1113 Sameoto, D. D., & Miller, R. S. (1968b). Selection of Pupation Site by Drosophila Melanogaster

1114  and D. Simulans. *Ecology*, *49*(1), 177–180. https://doi.org/10.2307/1933580

1115 Sawamura, K., Roote, J., Wu, C. I., & Yamamoto, M. T. (2004). Genetic Complexity Underlying

1116  Hybrid Male Sterility in Drosophila. *Genetics*, *166*(2), 789–796.

1117  https://doi.org/10.1534/genetics.166.2.789

1118 Schuster, C. M., Davis, G. W., Fetter, R. D., & Goodman, C. S. (1996). Genetic Dissection of

1119  Structural and Functional Components of Synaptic Plasticity. I. Fasciclin II Controls

1120  Synaptic Stabilization and Growth. *Neuron*, *17*(4), 641–654. https://doi.org/10.1016/S0896-

1121  6273(00)80197-X

1122 Seeholzer, L. F., Seppo, M., Stern, D. L., & Ruta, V. (2018). Evolution of a central neural circuit

1123  underlies Drosophila mate preferences. *Nature*, *559*(7715), 564–569.

1124  https://doi.org/10.1038/s41586-018-0322-9

1125 Seyahooei, M. A., Kraaijeveld-Smit, F. J. L., Kraaijeveld, K., Crooijmans, J. B. M., Van Dooren,

1126  T. J. M., & Van Alphen, J. J. M. (2009). Closely related parasitoids induce different

1127  pupation and foraging responses in Drosophila larvae. *Oikos*, *118*(8), 1148–1157.

1128  https://doi.org/10.1111/j.1600-0706.2009.17508.x

1129 Shivanna, N., Murthy, G. S. S., & Ramesh, S. R. (1996). Larval pupation site preference and its

1130  relationship to the glue proteins in a few species of Drosophila. *Genome*, *1*(1), 105–111.

1131  https://doi.org/10.1139/g96-014

1132 Sokal, R. R., Ehrlich, P. R., Hunter, P. E., & Schlager, G. (1960). Some Factors Affecting

Pupation Site of Drosophila. *Annals of the Entomological Society of America*, *53*(2), 174–

182. Retrieved from

http://www.ingentaconnect.com/content/esa/aesa/1960/00000053/00000002/art00006

Sokolowski, M. B., Kent, C., & Wong, J. (1984). Drosophila larval foraging behaviour:

Developmental stages. *Animal Behaviour*, *32*(3), 645–651. https://doi.org/10.1016/S0003-

3472(84)80139-6

Stamps, J., Buechner, M., Alexander, K., Davis, J., & Zuniga, N. (2005). Genotypic differences

in space use and movement patterns in Drosophila melanogaster. *Animal Behaviour*, *70*(3),

609–618. https://doi.org/10.1016/j.anbehav.2004.11.018

Stern, D. L. (2014). Identification of loci that cause phenotypic variation in diverse species with

the reciprocal hemizygosity test. *Trends in Genetics*.

https://doi.org/10.1016/j.tig.2014.09.006

Sturtevant, A. H. (1920). Genetic Studies on DROSOPHILA SIMULANS. I. Introduction.

Hybrids with DROSOPHILA MELANOGASTER. *Genetics*, *5*(5), 488–500. Retrieved from

http://www.ncbi.nlm.nih.gov/pubmed/17245951

Sucena, E., Delon, I., Jones, I., Payre, F., & Stern, D. L. (2003). Regulatory evolution of

shavenbaby/ovo underlies multiple cases of morphological parallelism. *Nature*, *424*(6951),

935–938. https://doi.org/10.1038/nature01768

Takano, T. S. (1998). Loss of notum macrochaetae as an interspecific hybrid anomaly between

Drosophila melanogaster and D. simulans. *Genetics*, *149*(3), 1435–1450.

Vandal, N. B., Siddalingamurthy, G. S., & Shivanna, N. (2008). Larval pupation site preference

on fruit in different species of Drosophila. *Entomological Research*, *38*(3), 188–194.

https://doi.org/10.1111/j.1748-5967.2008.00163.x

Venken, K. J. T., Popodi, E., Holtzman, S. L., Schulze, K. L., Park, S., Carlson, J. W., …

Kaufman, T. C. (2010). A molecularly defined duplication set for the X chromosome of

Drosophila melanogaster. *Genetics*, *186*(4), 1111–1125.

https://doi.org/10.1534/genetics.110.121285

Watanabe, T. K. (1979). A gene that rescues the lethal hybrids between Drosophila melanogaster

and D. simulans. *The Japanese Journal of Genetics*, *54*(5), 325–331.

https://doi.org/10.1266/jjg.54.325

Zeng, X., Han, L., Singh, S. R., Liu, H., Neumüller, R. A., Yan, D., … Hou, S. X. (2015).

Genome-wide RNAi Screen Identifies Networks Involved in Intestinal Stem Cell

Regulation in Drosophila. *Cell Reports*, *10*(7), 1226–1238.

https://doi.org/10.1016/j.celrep.2015.01.051

Zhou, J. J., Field, L. M., & He, X. L. (2010). Insect odorant-binding proteins: Do they offer an

alternative pest control strategy? *Outlooks on Pest Management*, *21*(1), 31–34.

https://doi.org/10.1564/21feb08

1170

1171

 Supplementary Figures and Captions:

1173

Fig S1. Pupation behavior differences between _D. melanogaster_ and _D. simulans_ after controlling
for density. The mean residuals from a regression between the proportion of pupae on the surface of the
food and the total number of pupae in the vial for 11 _D. melanogaster_ lines and 12 _D. simulans_ lines
described in Table S1. Error bars denote the 95% confidence interval around each individual mean (N =
5-8). The dashed horizontal lines indicate the grand mean for each species. The boxes surrounding the
dashed lines denote the 95% confidence interval around the grand mean.

Fig S2. Pupation behavior for _D. melanogaster_, _D. simulans_, and their F1 hybrids after controlling for density. The mean residuals from a regression between the proportion of individuals in a vial that pupated on the surface of the food and the total number of individuals in the vial are shown for males and females from both species and their F1 hybrids. _D. melanogaster_ males and females were taken from the LH$_M$ strain, while _D. simulans_ males and females were taken from the _Lhr_ strain. F1 hybrids resulted from a cross between these two strains. The "melX" hybrid males have the _D. melanogaster_ X chromosome, and the "simX" hybrid males have the _D. simulans_ X chromosome. Both hybrids have _D. melanogaster_ cytoplasmic inheritance. Box plots labeled with different letters are significantly different from one another after sequential Bonferroni adjustment for multiple comparisons (p < 0.0001, N = 26-33).

1190

Fig S3. Pupation behavior for *D. melanogaster*, *D. simulans*, and their F1 hybrids, measured as average pupation height within a vial. A. The average pupation height is shown for males and females from both species and their F1 hybrids. Higher pupation heights indicate individuals pupated further from the food. **B.** The residuals from a regression between the average pupation height of individuals in a vial and the total number of individuals in that vial are shown for males and females from both species and their F1 hybrids. The results from pairwise statistical tests can be found in Table S5.

1197

Fig S4. Pupation site choice behavior for additional significant hybrid deficiencies. Pupation
indices for **A.** Df(1)ED411 (N=29-36), **B.** Df(1)ED6906 (N=26-33), and **C.** Df(1)BSC530 (N=18-22). Each
of the three deficiencies displayed a median pupation index significantly greater than the average (0.88)
when crossed to *D. simulans* (*Lhr*) after correcting for multiple comparisons. However, when each was
crossed to *D. melanogaster* (T.4), the pupation index was still significantly elevated, indicating that the
behavior of these flies is impacted by the deficient region in general, rather than the *Lhr* genotype it
reveals.

1205

Fig S5. Pupation behavior for knockout strains crossed to both *D. melanogaster* and *D. simulans*.
A. The pupation indices of a *mei9* gene disruption crossed to *D. melanogaster* (T.4, N = 52) and *D. simulans* (*Lhr*, N = 51). **B.** The pupation indices of a *wap* gene disruption crossed to *D. melanogaster* (T.4, N = 51) and *D. simulans* (*Lhr*, N = 55).

1210

1211

Fig S6. Pupation behavior for norpA knockouts and control. A. The proportion of females that
pupated on the food is shown for the *norpA* mutant allele (norpA) and the *norpA* rescue allele (norpA
rescue) when crossed to *D. simulans* (*Lhr*). There was no significant difference in pupation behavior
between the knockout and the rescue allele (p=0.08). **B.** Pupation behavior for *norpA* knockout and *norpA*
rescue allele female hybrids (crossed to *Lhr*) after controlling for density. Vials were adjusted for density
using the residuals from a regression between the proportion of females in a vial that pupated on the
surface of the food and the total number of individuals in the vial. There was no significant difference
between the knockout and rescue allele after controlling for density (p=0.52). **C.** The proportion of
females that pupated on the food when the *norpA* mutant allele (dark grey, triangle) and the *norpA* rescue
allele (light grey, circle) were crossed to *D. melanogaster* (T.4) and *D. simulans* (*Lhr*). Shown are the least
square means (± 1 SE) from an ANOVA with "species", "*norpA* allele" and their interaction as fixed effects.
The actual analysis was conducted using residuals to control for density, and the "*norpA* allele" x "species"
interaction term was removed from the model as it was not significant (p=0.37).

1225

Fig S7. Pupation site choice behavior for *Fas2* and *tilB* RNAi crosses after controlling for density.
A. The results of pan-neuronal knock down of the *Fas2* transcript via RNAi. Shown are the mean residuals from a regression between the proportion of individuals that pupated on the surface of the food and the total number of individuals in the vial for the control cross, the Gal-4 line, and RNA interference cross. Asterisks denote significance after correcting for multiple comparisons (* = p < 0.05, ** = p < 0.01, ** = p<0.001; N = 42-48). **B.** The results of pan-neuronal knockdown of the *tilB* transcript via RNAi. Shown are the mean residuals from a regression between the proportion of individuals that pupated on the surface of the food and the total number of individuals in the vial for the control cross, the Gal-4 line, and RNA interference cross. Asterisks denote significance after correcting for multiple comparisons (* = p < 0.05, ** = p < 0.01; N = 41-47).

1236

Fig S8. Pupation site choice behavior for *Fas2* and *tilB* RNAi crosses, measured as average pupation height within a vial. Higher pupation heights indicate individuals pupated further from the food. **A.** The results of pan-neuronal knockdown of the *Fas2* transcript via RNAi. The average pupation height of individuals found on the food are shown for the control cross, the Gal-4 line, and RNA interference cross. Asterisks denote significance after correcting for multiple comparisons (**** = $p < 0.0001$; N = 42-48). **B.** The results of pan-neuronal knockdown of the *tilB* transcript via RNAi. The average pupation height of individuals found on the food are shown for the control cross, the Gal-4 line, and RNA interference cross. Asterisks denote significance after correcting for multiple comparisons (*** = $p < 0.0001$; N = 46-45).

Fig S9. Duplication Screen. The results of a duplication hybrid screen. Each duplication hybrid was created by crossing a DP(1;Y) *D. melanogaster* female, which carries a unique segment of the X chromosome attached to the Y chromosome in an XXY background, to an *Lhr D. simulans* male. The resulting hybrid males have a complete *D. simulans* X chromosome in addition to 1 of 22 segments of the *D. melanogaster* X chromosome translocated to the Y chromosome. Females are largely inviable. In this way, we tested for dominant *D. melanogaster* X loci underlying the interspecific pupation difference. We scored pupation behavior following the same methods used for deficiency and knockout hybrids, and measured the proportion of pupae on the food for each duplication hybrid. Shown for comparison are the values for males from the *D. simulans* parent strain (*Lhr*) and hybrid males with a *D, simulans* X chromosome (simX). Interestingly, all of the duplication hybrids had significantly more males pupating on the food surface than either their *D. simulans* parent strain or typical simX hybrids that are not heterozygous for a region of the X. From this, we conclude that making males heterozygous over broad regions of the X chromosome results in behavioral or developmental inconsistencies that leave larvae unable to climb the vial walls to pupate. Information about these Y-linked X duplications can be found in Table S5.