Ultra-Efficient Short Read Sequencing of T Cell Receptor Repertoires

Janelle M. Montagne[1], Xuwen Alice Zheng[1], Iago Pinal-Fernandez[2,3], Jose C. Milisenda[4], Lisa Christopher-Stine[5], Thomas E. Lloyd[3], Andrew L. Mammen[2,3], & H. Benjamin Larman[1,*]

[1]Division of Immunology, Pathology Department, Johns Hopkins University School of Medicine, Baltimore, MD, USA.

[2]Muscle Disease Unit, Laboratory of Muscle Stem Cells and Gene Regulations, National Institute of Arthritis and Musculoskeletal and Skin Diseases, National Institutes of Health, Bethesda, MD, USA.

[3]Department of Neurology, Johns Hopkins University School of Medicine, Baltimore, MD, USA.

[4]Internal Medicine Department, Hospital Clinic, Universitat de Barcelona, Barcelona, Spain and Centro de Investigación Médica en Red Enfermedades Raras.

[5]Division of Rheumatology, Johns Hopkins University School of Medicine, Baltimore, MD, USA.

Abstract:

T cell receptor (TCR) repertoire sequencing is increasingly employed to characterize adaptive

immune responses. However, current TCR sequencing methodologies are complex and

expensive, limiting the scale of feasible studies. Here we present Framework Region 3

AmplifiKation sequencing (FR3AK-seq), a simplified multiplex PCR-based approach for the ultra-

efficient analysis of TCR complementarity determining region 3 (CDR3) repertoires. By using

minimal primer sets targeting a conserved region adjacent to CDR3, undistorted amplicons are

analyzed via short read, single-end sequencing. We find that FR3AK-seq is sensitive and

quantitative, performing comparably to two industry standards. FR3AK-seq was utilized to quickly

and inexpensively characterize the T cell infiltrates of muscle biopsies obtained from 145 patients

with idiopathic inflammatory myopathies and controls. A cluster of related TCRs was identified in

samples from patients with sporadic inclusion body myositis, suggesting the presence of a shared

antigen-driven response. The ease and minimal cost of FR3AK-seq removes critical barriers to

routine, large-scale TCR CDR3 repertoire analyses.

Introduction:

T cell receptor (TCR) repertoire analysis has emerged as a powerful tool for examining adaptive immune responses. TCR repertoires, generated by the process of V(D)J recombination, encompass the T cell clones within a given individual or sample. The unique TCR of each clone defines its antigen specificity, and TCRs can be associated with distinct cellular phenotypes and tissue occupancy. Notably, TCR repertoires encompass dynamic populations of cells that represent past and current immune exposures and reactivities.[1] Development of next generation sequencing (NGS)-based technologies over the last decade has enabled the unprecedented analysis of TCR repertoires.[2–5]

TCR recognition requires antigen processing and presentation of peptide epitopes on autologous major histocompatibility complex (MHC) molecules. Analysis of TCR repertoires both within and between individuals can be indicative of disease status, prior infections or immunizations, and individual-specific attributes of epitope selection.[6–12] Clonally expanded or tissue-infiltrating T cell clones provide evidence of antigen-directed immune responses; integrating TCR repertoire analyses with phenotypic measurements (e.g. single cell transcriptional profiling[13] or flow cytometric sorting of T cell subpopulations[14,15]) can further illuminate qualitative aspects of T cell responses (**Fig 1a**).

A variety of methods have been used to generate TCR libraries for NGS, including multiplex PCR, 5'-RACE, and target enrichment.[2–5,16,17] Generally, these methodologies prioritize sequencing of the hypervariable complementarity determining region 3 (CDR3) of the TCR beta (TCRB) chain. The TCRB CDR3 harbors the greatest amount of sequence diversity, confers most of the antigen specificity to a TCR, and can be used as a unique surrogate for T cell clonal identity.[18–20] Despite their utility, however, current repertoire sequencing approaches remain complex and expensive. These limitations have greatly hindered the utilization of TCR repertoire-based analyses, particularly for studies involving large numbers of samples. We therefore sought to develop a

simple, quantitative, multiplex PCR-based approach that prioritizes the ultra-efficient analysis of

TCRB CDR3 sequences.

The sequence diversity of the TCRB chain variable domain, which in humans is encoded by >120

unique TCRBV alleles, necessitates the use of complex primer pools for multiplex PCR

amplification. Primer competition and differential amplification efficiencies distort clonal amplicon

abundance, confounding quantification.[21] Sophisticated techniques have been developed to

computationally correct amplification bias, such as the use of spike-in library standards or the

incorporation of unique molecular identifiers (UMIs).[22–24] Adaptive Biotechnologies has become

a current industry leader by offering a spike-in standard corrected, multiplex PCR-based assay.

At a cost of $500-$1100 per sample, however, the scale of feasible studies using this assay has

been greatly constrained. Unique molecular identifier (UMI) based approaches, including the

Immunoverse assay (ArcherDX), provide quantitation and enhanced sequence accuracy by

resolving amplification and sequencing errors.[23–28] However, UMI approaches typically require

multi-day protocols, complicated analytical pipelines, and deeper sequencing to obtain sufficient

sampling of distorted libraries.[17,29] By designing maximally compact primer sets and a streamlined

workflow, we have essentially eliminated PCR amplification bias while maintaining high sequence

level accuracy. Lower sequencing depth requirements, coupled with CDR3 analysis via single-

end, short-read (100 nucleotide) sequencing, dramatically reduces the cost associated with this

approach to TCR repertoire analysis.

Here, we benchmark our simplified multiplex PCR-based assay, which we call Framework Region

3 AmplifiKation Sequencing (FR3AK-seq), against two different industry standards for quantitative

TCR repertoire sequencing: Adaptive Biotechnologies' multiplex PCR-based hsTCRB

immunoSEQ assay and ArcherDX's UMI-based Immunoverse HS TCR assay. We found that

FR3AK-seq data, which was analyzed using open-source software, was in excellent agreement

with both. Finally, we illustrate the utility of FR3AK-seq to characterize T cell responses within

biopsied muscle tissue from a cohort of 145 patients with idiopathic inflammatory myopathies and

controls.

Results:

Minimal primer sets targeting the TCRB framework region 3 (FR3) efficiently amplify CDR3

sequences

We hypothesized that minimizing the number of potentially competing V-gene primers, while also

minimizing amplicon length, would maximize the efficiency of TCRB CDR3 region amplification.

To identify candidate primer binding sequences upstream and proximal to CDR3, we constructed

a phylogenetic tree of all functional TCRB FR3 regions available from the Immunogenetics

(IMGT/GENE-DB) reference directory.[30] This revealed a high degree of sequence conservation

within the 3' terminus of FR3 (**Fig 1b**). This homology may reflect a critical role for these

sequences in the optimal presentation of the CDR3 loop. The 3' 20 nucleotides of these IMGT

FR3s were therefore extracted and used as target primer binding sequences. Remarkably, the

128 functional TCRBV alleles reported by IMGT are represented by only 47 distinct 3'-terminal

20-mers. Beyond their sequence homology, additional advantages of targeting these sequences

include minimal CDR3 amplicon length (and in turn optimal amplification efficiency) and the ability

to sequence using short (100 nucleotide) single end reads initiated proximal and upstream of

CDR3. We refer to this approach to TCRB CDR3 analysis as FR3 AmplifiKation sequencing, or

"FR3AK-seq".

We developed a greedy algorithm to automate the design of three candidate sets of primers, each

targeting the 47 distinct TCRB FR3 3' 20-mers (**Fig S1**): one primer set lacks any universal bases

or mismatches ("Full" set), one primer set lacks universal bases but allows a single mismatch to

occur ("1MM" set), and one primer set contains parsimonious incorporation of up to 3 universal

(inosine) bases and allows one mismatch ("Compact" set). No mismatches were permitted within

the five nucleotides of the 3' end of the primers, as a precaution to minimize interference with

polymerase extension.[31–33] The Full set consists of 47 primers (equal to the number of unique

FR3 3' 20-mers), the 1MM set consists of 34 primers, and the Compact set consists of 20 primers

(**Table S1**). We calculated the expected efficiency of each primer to amplify each FR3 sequence;

these efficiencies are visualized in the form of a clustered heatmap (**Fig 1c**). Importantly, each of

the 47 unique 3' FR3 20 mers are predicted to be amplified efficiently by at least one primer in

each set.

Figure 1: Utility of T cell receptor beta (TCRB) chain sequencing and FR3AK-seq multiplex PCR primer design. **a.** TCRB chain sequencing can be used to (i) identify related CDR3s that may share antigen specificity, (ii) detect expanded clones, (iii) identify tissue infiltrating clones, and (iv) link CDR3 sequences with T cell phenotypes. **b**. A schematic of the TCRB chain. A neighbor joining phylogenetic tree of FR3s extracted from the 128 functional human TCRBV alleles in the IMGT/GENE-DB reference directory and multiple sequence alignment from a dominant FR3 sequence cluster (red box on tree) identifies homology within the 3' ~20 nucleotides (yellow box). **c.** Clustered heatmaps for the Full, 1MM, and Compact primer sets showing primer amplification efficiencies for each primer across each of the 47 unique 3' 20 nucleotides of the human TCRBV FR3 region. Inosines were considered exact matches for the Compact set. Y axis: 47 unique TCRBV FR3s. X axis: primers. The order of the primers in each heatmap can be found in **Table S1**. The order of unique TCRBV FR3s is identical to the order of the Full primers for all heatmaps.

TCRB FR3 primers amplify CDR3 sequences with minimal bias

Each primer was tested separately and within its set for the ability to produce amplicon from peripheral blood mononuclear cell (PBMC) cDNA. Agarose gel electrophoresis revealed a PCR product at the expected size of ~220 base pairs (**Fig S2a**). Importantly, incorporation of the inosine base did not prevent amplification by the KAPA Fast HotStart *Taq* DNA Polymerase. While lower PCR primer annealing temperature might increase off-target priming, it may also reduce unwanted bias in amplification from primers containing a single nucleotide mismatch.[34] We therefore utilized gradient PCR to assess the effect of annealing temperature on the number of unique CDR3 sequences detected using the Compact primer set. As expected, more unique CDR3 clones were recovered at lower annealing temperatures (**Fig S2b**), and a similar trend was observed for the 1MM primer set. The annealing temperature that produced the maximum number of unique clones was identified for each set as follows: Full: 52°C, 1MM: 47°C, Compact: 42°C. Use of a hemi-nested RT-PCR strategy (**Fig S3**) resulted in no apparent amplification of non-TCRB sequences, even at the lowest PCR annealing temperature.

We next quantified PCR amplification bias inherent to the three primer sets. cDNA first underwent 20 cycles of PCR. An aliquot of this product was diluted 2^{10}-fold and then subjected to 10 more cycles of PCR; sequencing was then performed on both amplicons and the resulting data sets compared to each other. In this experiment, significant per-cycle amplification bias would manifest as discordance. However, the concordance remains high for all three primer sets, suggesting that FR3AK-seq primers negligibly bias the CDR3 repertoire during cycles of PCR amplification ($\rho = 0.997$ for the Full primer set and $\rho = 0.998$ for the 1MM primer set, **Fig 2a**; $\rho = 0.997$ for the Compact primer set, **Fig S2c**).

We then performed rarefaction analysis to determine the relationship between RNA input amount, sequencing depth, and the number of unique clones detected using FR3AK-seq. We found that diversity discovery is saturated at ~1×10^6 reads for 100 ng of purified peripheral blood T cell RNA

(**Fig 2b**), providing the rationale for RNA input amounts and sequencing depths used in

subsequent analyses. We also assessed the sensitivity of FR3AK-seq to detect rare clones by

spiking varying amounts of monoclonal Jurkat RNA (from 1 up to 10,000 cellular equivalents) into

a background of 400 ng PBMC RNA (~20% T cells). We found that FR3AK-seq detected the

Jurkat CDR3 sequence at all input amounts at the expected abundances, while the abundance

of the top three PBMC T cell clones remained constant across the dilution series (**Fig 2c**). The

Jurkat sequence abundance was used to calculate cellular equivalents for the top three PBMC

clones in each sample. Notably, the coefficient of variation for these cellular equivalents was

below 1% in the 1,000 Jurkat cell equivalent spike-in samples (CV = 0.897%). This provided the

basis for the subsequent use of 1,000 cellular equivalents of spike-in cells.

Figure 2: FR3AK-seq multiplex PCR performance. **a.** MA plots of 2^{10} dilution experiments for the Full and 1MM primer sets demonstrated high linear concordance (Full: 0.997, 1MM: 0.998, Compact in **Fig S2c**), indicating negligible PCR bias. Clones with ≥10 counts in either dataset were compared. Insets indicate number of clones above or below Y axis limits (set to 2 and –2 for visualization). **b.** Rarefaction analysis using T cell RNA purified from PBMCs showed the relationship between RNA input, number of clones detected, and sequencing depth using FR3AK-seq. **c.** The spiked-in Jurkat CDR3 sequence was detected at the expected abundances using FR3AK-seq (performed in triplicate). Estimated abundances of the top three T cell clones were consistent among triplicates and across spiked-in samples. 95% confidence intervals for regression lines and coefficients of variation for the cellular equivalents calculated for the T cell clones are shown.

FR3AK-seq performs comparably to a multiplex PCR-based industry standard

T cell RNA from two donors, A and B, as well as a third sample comprised of 90% Donor A RNA and 10% Donor B RNA (sample "C"), were used to generate cDNA for benchmarking studies against the current multiplex PCR-based industry standard, the immunoSEQ hsTCRB "Deep Resolution" sequencing service offered by Adaptive Biotechnologies. Separate cDNA aliquots were also subjected to FR3AK-seq analysis using each of the three primer sets. The open source software MiXCR v2.1.11 was used to define and quantify CDR3 sequences from both assays for direct comparison.[35] The technical reproducibility of the immunoSEQ assay was quantified by comparing the abundances of Donor A's unique CDR3 sequences against their corresponding abundances in sample C (concordance correlation coefficient, ρ = 0.989, left panel **Fig 3a**). The same analysis was applied to the FR3AK-seq data sets, which were generated using each of the three FR3 primer sets. Equally high measures of internal concordance were observed (ρ = 0.990 for the 1MM primer set, right panel **Fig 3a;** ρ = 0.990 for the Full primer set and ρ = 0.987 for the Compact primer set, **Fig S4a**). Importantly, the concordance between immunoSEQ and FR3AK-seq measurements of CDR3 abundance was also very high (ρ = 0.975, **Fig 3b**, **Fig S4b**), providing further evidence that FR3AK-seq amplifies CDR3s with negligible bias. The majority of even moderately abundant immunoSEQ-defined CDR3 sequences were also detected using any of the three FR3 primer sets (**Fig 3c**). Additionally, the 9:1 mixture composed of cells from Donor A and Donor B (sample C) was used to determine quantification accuracy of the FR3 primer sets relative to the immunoSEQ assay (**Fig 3d**, **Fig S4c**). These data indicate that all three FR3 primer sets provide CDR3 quantification comparable to the industry standard immunoSEQ assay, with the 1MM primer set performing most accurately.

Figure 3: FR3AK-seq performs comparably to the multiplex PCR-based immunoSEQ assay from Adaptive Biotechnologies. **a.** MA plots comparing technical replicates of the immunoSEQ and 1MM FR3AK-seq assays. **b.** MA plot comparing the 1MM FR3AK-seq assay against the immunoSEQ assay. Clones with ≥10 counts in either dataset were included. **c.** A Venn diagram shows overlapping and non-overlapping CDR3 sequences detected using immunoSEQ and all three FR3AK-seq primer sets. CDR3 sequences were included in this analysis if they had a clone count ≥10 in any of the four data sets. **d.** Relative frequencies of Donor A (black points) and Donor B (red points) in sample C determined using the immunoSEQ assay or the 1MM FR3AK-seq assay. Insets in **a** and **b** indicate number of clones above or below Y axis limits (set to 2 and -2 for visualization).

FR3AK-seq performs comparably to a unique molecular identifier (UMI)-based industry standard

To assess whether FR3AK-seq achieves results comparable to UMI-based methodologies, we compared the FR3AK-seq 1MM and immunoSEQ datasets to one obtained using ArcherDX's UMI-based Immunoverse HS TCR assay. PBMC RNA from the same samples A, B, and C described above were subjected to Immunoverse analysis for comparison with the FR3AK-seq and immunoSEQ datasets. The technical reproducibility of the Immunoverse assay was similar to that of both immunoSEQ and FR3AK-seq (ρ = 0.967, **Fig 4a**). Additionally, concordance between FR3AK-seq and Immunoverse was high (ρ = 0.900, upper panel **Fig 4b**). Relatively high concordance was also observed when comparing immunoSEQ to Immunoverse (ρ = 0.886, lower panel **Fig 4b**). Crucially, the majority of clones detected by the Immunoverse assay were also detected by both immunoSEQ and FR3AK-seq (**Fig 4c**). These data demonstrate that FR3AK-seq has a high sensitivity for detecting clones of the correct sequence (4352/4403, 98.8% of total clones detected by Immunoverse). Strikingly, a large number (2803) of clones were detected by both immunoSEQ and FR3AK-seq but not Immunoverse. Rarefaction analysis of the FR3AK-seq and Immunoverse datasets explained this difference in clonal detection sensitivity, in that FR3AK-seq sampled more clones at all sequencing depths (**Fig 4d**). Although FR3AK-seq achieves better coverage of the repertoire than Immunoverse across sequencing depths, we wanted to quantify how many of the FR3AK-seq detected clones may be the result of false discovery. To address this question, we performed FR3AK-seq 1MM analysis of 1,000 cellular equivalents of Jurkat cell RNA in triplicate. Since the Jurkat cell line is monoclonal, the frequencies of subdominant clones, presumably generated by PCR or sequencing errors, can be used to assess the rate of false positive discovery. 99.98% of MiXCR-defined CDR3s mapped to the correct reported Jurkat TCRB sequence in each sample. We found an average of 50 non-Jurkat clones across triplicates, of which only one was moderately abundant ($\geq$10 clone counts) in each. This secondary clone

was the same between replicates and represented on average 0.0067% of total MiXCR-defined

CDR3s. Using default MiXCR settings, FR3AK-seq analysis is therefore associated with a very

low rate of false discovery due to PCR or sequencing errors. Modification of MiXCR thresholds

can be used to adjust the stringency with which these subdominant clones are grouped. We also

compared the CDR3 lengths of clones detected using each sequencing platform, and found that

FR3AK-seq's short (100 nucleotide) reads did not bias the distribution of CDR3 lengths recovered

(**Fig 4e**).

Figure 4: FR3AK-seq performs comparably to the UMI-based ArcherDX Immunoverse assay. **a.** MA plot comparing Immunoverse technical replicates. **b.** MA plots comparing the 1MM FR3AK-seq assay against the Immunoverse assay (top panel) and Adaptive Biotechnologies' immunoSEQ assay against the Immunoverse assay (bottom panel). Clones with ≥10 counts in either dataset were included (using pre-deduplicated counts for the Immunoverse dataset). **c.** A Venn diagram shows overlapping and non-overlapping CDR3 sequences detected using Immunoverse, immunoSEQ, and the 1MM FR3AK-seq primer set. CDR3 sequences were included in this analysis if they had a clone count of ≥10 in any of the three data sets. **d.** Rarefaction analysis comparing the 1MM FR3AK-seq assay to the Immunoverse assay using 100 ng of T cell RNA (~250,000 cells) for 1MM FR3AK-seq analysis and 200 ng of T cell RNA (~500,000 cells) for Immunoverse analysis. 1MM: ~1.2x10^6 reads, Immunoverse: ~1.4x10^6 reads. **e.** Violin plots comparing CDR3 lengths captured by the short-read 1MM FR3AK-seq versus longer-read immunoSEQ and Immunoverse assays. Insets in **a** and **b** indicate number of clones above or below Y axis limits (set to 2 and -2 for visualization).

FR3AK-seq enables inexpensive cohort-scale repertoire studies

Muscle biopsies from 145 patients with idiopathic inflammatory myopathies (IIMs, 124) as well as healthy controls (9) and non-IIM controls (12) were analyzed using the 1MM primer set at a total cost of ~$20 per sample (**Table S2**). IIM patients included those with dermatomyositis (DM, 40), immune-mediated necrotizing myopathy (IMNM, 49), inclusion body myositis (IBM, 14), and anti-synthetase syndrome (ASyS, 21) (**Table S3**). For absolute T cell number estimation, 1,000 cell equivalents of RNA derived from a clonal tumor infiltrating T cell line[36] were spiked into each sample after RNA purification. Aggregated CD4 and CD8 sequencing reads from RNA-seq analysis of the same samples (performed separately and reported elsewhere) tightly correlated with FR3AK-seq based estimates of cellular equivalents (**Fig 5a**). In subsequent analyses, CDR3 sequences present at levels at or below one cell equivalent per biopsy were considered 'bystander' T cell clones, unlikely to be involved in the disease process. By this metric, the number of non-bystander T cells were elevated in each disease subgroup versus the controls (**Fig 5b**). In agreement with previous immunohistochemical analyses, IBM biopsies contained particularly high levels of T cell infiltrates.[37–39] We determined this to be the case both in terms of absolute T cell number and clonal diversity (**Fig 5b, Fig S5a**). No distinct patterns of J gene usage were found among patient subgroups (**Fig S6a**), and CDR3 hydrophobicity and lengths were similar between patient subgroups (**Fig S6b-c**).

The open source software GLIPH (grouping of lymphocyte interactions by paratope hotspots)[10] was next used to search for clusters of related CDR3 sequences within the aggregated set of non-bystander CDR3s (**Table S4**). 143 clusters composed of sequences contributed by at least 3 individuals were identified. We first examined clusters that had a disproportionate number of sequences contributed by single individuals. **Fig 5c** provides an example of a cluster dominated by related CDR3 sequences contributed by a single ASyS patient. Compared to CDR3 sequences from the same patient which were not in any cluster, the clonal abundances of these CDR3s were

significantly increased, suggestive of a polyclonal, shared antigen-driven expansion. These

results suggest that GLIPH is able to cluster related CDR3 sequences that have been detected

and quantified using FR3AK-seq.

We next determined whether any of the GLIPH clusters were composed of sequences contributed

disproportionately by individuals from one of the IIM subgroups. Indeed, we found a GLIPH cluster

that was composed exclusively of sequences contributed by IBM patients. Specifically, 5 of the

12 IBM patients with non-bystander T cells contributed at least one sequence to this cluster (p =

1.4×10^{-6}, **Fig 5d**). Remarkably, all CDR3 sequences in this IBM cluster are linked to a single J-

chain (TRBJ2-2*01), providing additional support for shared antigen recognition**.** Taken together,

this study demonstrates the utility of FR3AK-seq in probing T cell responses at cohort scale.

Figure 5: FR3AK-seq detects related T cell responses in subsets of patients with idiopathic inflammatory myopathies. **a.** Scatterplot of estimated T cell counts per patient versus CD4 and CD8 expression levels obtained from bulk mRNA-seq. Each patient data point is colored according to their IIM association. Pearson correlation coefficient (r) is shown. **b.** Distributions of cell counts per patient for each IIM and controls. Kruskal-Wallis p-value is provided. **c.** A CDR3 cluster enriched for sequences from a single patient with ASyS was identified using GLIPH software (p = 1.4×10^{-7}). The amino acid sequence logo of the CDR3 region is shown above the violin plot. The clones within this cluster are more expanded than corresponding unclustered clones from the same patient. Mann-Whitney p = 0.04. Cell counts were set to a maximum of 20 for visualization. **d.** An IBM-exclusive cluster was also identified using GLIPH, encompassing 5 of 12 IBM patients who had non-bystander T cell infiltrates (41.7%, p=1.4×10^{-6}). Multiple sequence alignment (using MUSCLE) is shown for the corresponding MiXCR-defined CDR3s and the J chain allele (TRBJ2-2*01).

Discussion

Although the utility of TCR repertoire sequencing to characterize T cell responses has been well established, the lack of a streamlined, quantitative, inexpensive, and non-proprietary assay has limited the scope and scale of feasible studies. By rationally designing minimal sets of primers that target the 3' terminus of FR3, we have developed a multiplex PCR-based approach for ultra-efficient library preparation and sequencing of TCRB CDR3 repertoires. By minimizing amplification bias (via reduction of primer number and amplicon length), the resulting sequencing libraries quantitatively capture clonal abundance distributions with an accuracy comparable to both multiplex PCR-based and UMI-based industry standards. While the library preparation and sequencing strategy presented here already bring the per-sample cost to ~$20 (**Table S2**)**,** we expect that alternative approaches to library preparation, reduced sequencing depth, and declining sequencing costs will likely enable another ~10-fold reduction in per-sample cost. In addition to reducing the overall cost of obtaining CDR3 sequences, we have developed the FR3AK-seq workflow to require minimal expertise, effort, and time. Quantitative sequencing libraries can easily be prepared within a single day. Comparisons of protocols and cost between FR3AK-seq, immunoSEQ, and Immunoverse are summarized in **Table S5**.

Our approach to TCRB repertoire analysis should be generalizable to additional immune receptor types, as well as to those of non-human species (see **Table S6** for *Mus musculus* TCRB primers). Although these mouse primers have not been as extensively validated as the human set, they have been confirmed to efficiently amplify CDR3 sequences from mouse tissues. We have additionally used the same FR3 primer design and analysis principles to target human TCR alpha, gamma, and delta repertoires, although these primers have not yet been validated (**Table S7**). We have also designed a set of TCRB J region reverse primers (**Table S8**) that perform comparably to our constant region reverse primer using cDNA templates ($\rho = 0.949$, **Fig S7**). These will be potentially useful for analysis of FFPE samples which suffer from extensive RNA

degradation. Interestingly, during the preparation of this manuscript, the Euroclonality Group

(creators of the BIOMED2 primer set[40]) published a new set of TCR primers for next-generation

sequencing, including some that are designed in the 5' portion of FR3,[41] providing additional

validation for an FR3-targeted approach to TCR repertoire sequencing.

V gene usage is often of interest, for example to detect usage skewing and for functional studies

involving TCR cloning. FR3AK-seq prioritizes small amplicons to reduce both PCR bias and cost.

Therefore, by design, amplicons do not contain easily recoverable V gene sequence information.

However, it is theoretically possible to make inferences on each CDR3's associated V genes,

based on the pattern of FR3 primer usage. We expect these inferences to be more accurate for

more abundant clones. When analysis of V gene usage is desired, complementary techniques

(e.g. 5'RACE) can be parsimoniously utilized to capture this information. One may wish to use

FR3AK-seq to track unique CDR3 sequences of interest over time, across tissues, and/or after

FACS analysis – these FR3AK-seq detected CDR3 sequences can then be readily associated

with a V gene by merging data sets.

To demonstrate the use of FR3AK-seq for efficient cohort scale analysis of TCRB CDR3

repertoires, we characterized the muscle-infiltrating T cells present within muscle biopsies

obtained from 145 inflammatory muscle disease patients and controls. TCRB CDR3 sequence

clustering indicated both donor and disease specific antigen-driven T cell responses. Importantly,

the majority of T cell clones detected in IBM muscle biopsies were not hyperexpanded, in contrast

to the hypothesis that infiltrating IBM T cells originate from a clonal T cell large granular

lymphocytic leukemia (TLGLL) (**Fig S5b**).[42,43] Future studies will utilize FR3AK-seq in combination

with *in vitro* stimulation to characterize the antigenic determinants of these clonal clusters.

Methods:

FR3 primer design

FR3 sequences from all functional human TCRB V alleles were downloaded from the IMGT/LIGM-DB reference directory (non-functional and pseudogenes were excluded).[30] Multiple sequence alignment (MSA) was performed using MUSCLE and UPGMA (Unweighted Pair Group Method with Arithmetic Mean) and neighbor-joining phylogenetic trees were generated via the R package msa.[44] The 3' 20 nucleotides of the TCRB chain FR3s were subsequently used to design three sets of primers as described in the text. We automated design of the 1MM and the Compact primer sets with an algorithm outlined in **Fig S1**, available on Github (https://github.com/jmmontagne/FR3AK-seq). Human TCRB primer sequences can be found in **Table S1**.

PBMC preparation, T cell purification, RNA extraction, and cDNA synthesis

PBMCs from Donors A and B were isolated by Ficoll-Paque (GE Healthcare, Pittsburgh, PA) gradient centrifugation and cryopreserved. T cells were purified from thawed PBMCs using the EasySep Human T cell enrichment kit (STEMCELL Technologies, Cambridge, MA). RNA was extracted from purified human T cells using the RNeasy Plus Minikit (Qiagen, Germantown, MD). 0.5 μg Donor B RNA was mixed with 4.5 μg Donor A RNA to create sample C for quantitation experiments. 4 μg each of Donor A, Donor B, and sample C RNA was reverse transcribed using a TCRB chain constant region reverse primer with Superscript III First-Strand Synthesis System (Invitrogen, Waltham, MA). cDNA was column purified with the Oligo Clean and Concentrator Kit (Zymo Research, Irvine, CA). 100 ng of purified cDNA from each sample was sent to Adaptive Biotechnologies (Seattle, WA) for the immunoSEQ hsTCRb Deep sequencing service or used for FR3AK-seq PCR. For the ArcherDX (Boulder, CO) Immunoverse analysis, RNA was extracted from matched Donor A and B PBMCs and mixed at the same 9:1 ratio to create sample C. 800

363   ng of RNA from each sample was input into the Immunoverse assay. The concordance between

364   FR3AK-seq and Immunoverse remained high despite the use of PBMC RNA rather than T cell

365   RNA for the Immunoverse assay. All primer sequences are provided in **Table S1**.

Polymerase Chain Reaction (PCR) and sequencing library preparation

100 ng TCRB chain cDNA was used as template for PCR with KAPA2G Fast Multiplex Mix (Kapa

Biosystems, Wilmington, MA). Forward TCRB FR3 primers from the Full, 1MM, or Compact

primer sets were used with a single nested TCR beta chain constant region reverse primer at 0.2

μM per primer. Samples underwent 30 cycles of PCR for visualization by agarose gel

electrophoresis, or 20 cycles for NGS library preparation ("PCR1"). We compared the

performance of Herculase II Fusion DNA polymerase (Agilent, Santa Clara, CA) and KAPA2G

Fast Multiplex Mix for PCR1 and found that both enzymes perform comparably (ρ = 0.966, **Fig**

S8). Herculase II is therefore an acceptable enzyme alternative for PCR1 if desired.

20 cycles of PCR2 were performed on PCR1 product (2 μL of PCR1 added to 18 μL of PCR2

master mix, which contained 0.25 μM each i5 and i7 sample barcoding primers) to incorporate

sample barcodes and Illumina sequencing adaptors using Herculase II Fusion DNA Polymerase.

Equal volumes of barcoded PCR2 products (5 μL each) were pooled and PCR column purified

using QIAquick PCR Purification Kit (Qiagen). Libraries were quantified using KAPA Library

Quantification Kit for Illumina Platforms (Kapa Biosystems). Barcoding primer sequences can be

found in **Supplementary File 1**. See **Fig S3** for a schematic of our sequencing strategy.

Sequencing and CDR3 analysis

Sequencing was performed on an Illumina NextSeq 500 (immunoSEQ benchmarking

comparisons), MiSeq (Immunoverse samples), or HiSeq 2500 (IBM muscle biopsy analysis).

CDR3s were identified and quantified using MiXCR v2.1.11 software[35] with default parameters

except as noted in **Table S9.** Data obtained from Adaptive Biotechnologies' immunoSEQ assay

were re-analyzed using MiXCR v2.1.11 and identical parameters for comparison to our own data.

For reanalysis, full nucleotide sequences for each CDR3 from the Adaptive Biotechnologies

ImmunoSEQ dataset were expanded to repeat as many times as indicated by the corresponding

"count" column. This file, with each clonal nucleotide sequence represented as many times as its

"count" column, was converted to FASTA format for compatibility with MiXCR. The sequences

obtained from ArcherDX's Immunoverse assay were expanded to repeat as many times as their

pre-deduplicated clone counts and then matched back to their deduplicated clone counts after

reanalysis with MiXCR v2.1.11. Clone count cutoffs for each analysis are as described in the

figure legends and refer to the pre-deduplicated counts for the Immunoverse assay (Note:

deduplicated clone counts were used for all comparisons, while pre-deduplicated counts were

only used to filter the dataset to $\geq$ 10 counts). Information regarding the total number of clones,

the number of singletons, and the number of clones occurring at $\geq$10 counts for FR3AK-seq,

immunoSEQ, and Immunoverse can be found in **Table S10**.

Repertoire analysis using muscle biopsies from patients with idiopathic inflammatory

myopathies (IIMs)

RNA isolated from muscle biopsies taken from 145 IIM patients or controls was reverse

transcribed, PCR amplified, sequenced, and analyzed as described above. Healthy control

muscle biopsies were obtained from healthy volunteers at the Skeletal Muscle Biobank of the

University of Kentucky, while non-IIM controls were obtained for clinical purposes at the Johns

Hopkins Neuromuscular Pathology Laboratory and were normal after pathologic evaluation. We

added 1,000 cell equivalents of RNA from a clonal tumor infiltrating T cell line as a spike-in to

each sample.[36] Sequencing of 6 spike-in only samples provided the CDR3 sequences to remove

from downstream analyses of samples containing muscle biopsy RNA. The spike-in counts were

used for cell number quantification within patient samples, which distinguished 'bystander' versus

'non-bystander' T cell clones. One non-IIM control muscle biopsy was sequenced in duplicate,

making the total number of samples 146.

CDR3 sequences obtained from these samples were analyzed for disease-specific clusters using

the GLIPH 1.0 group discovery algorithm (see **Table S9** for parameters).[10] To determine the

statistical significance of each cluster, we performed chi-square analysis on all clusters with at

least three contributing individuals, followed by Benjamini-Hochberg multiple comparison

correction. This analysis was performed for both the number of sequences contributed by each

disease subgroup to each cluster, as well as the number of patients in each disease group

represented within each cluster. P-values ≤0.05 after multiple test correction were considered

statistically significant.

Author Contributions:

Conceptualization and experimental design: J.M.M., H.B.L.; performing experiments: J.M.M.; data

analysis and software: J.M.M., X.A.Z.; human specimen procurement and analysis: I.P-F., J.C.M.,

L.C-S., T.E.L., A.L.M., H.B.L.; data interpretations: J.M.M., X.A.Z., T.E.L., A.L.M., H.B.L.; paper

writing: J.M.M., H.B.L. All authors reviewed the paper.

Competing Interests:

H. Benjamin Larman is a consultant for Tscan Therapeutics, which seeks to develop T cell

receptor based cellular therapies.

Acknowledgements:

The authors would like to thank Dr. Erik Wright for his advice on primer design, Dr. Suzanne L.

Topalian for the TIL1558 cell line, and Dr. Maximilian F. Konig for his review of the manuscript.

We would also like to thank the Johns Hopkins Medical Institute Deep Sequencing and Microarray

Core Facility and the Johns Hopkins Institute of Genetic Medicine Genetics Resources Core

Facility. This work was supported by NIH grant U24AI118633 and a Prostate Cancer Foundation

Young Investigator Award.

References:

1. Attaf, M., Huseby, E. & Sewell, A. K. αβ T cell receptors as predictors of health and disease. *Cell. Mol. Immunol.* **12**, 391–399 (2015).
2. Freeman, J. D., Warren, R. L., Webb, J. R., Nelson, B. H. & Holt, R. A. Profiling the T-cell receptor beta-chain repertoire by massively parallel sequencing. *Genome Res.* **19**, 1817–1824 (2009).
3. Robins, H. S. *et al.* Comprehensive assessment of T-cell receptor beta-chain diversity in alphabeta T cells. *Blood* **114**, 4099–4107 (2009).
4. Weinstein, J. A., Jiang, N., White, R. A., Fisher, D. S. & Quake, S. R. High-throughput sequencing of the zebrafish antibody repertoire. *Science* **324**, 807–810 (2009).
5. Boyd, S. D. *et al.* Measurement and clinical monitoring of human lymphocyte clonality by massively parallel VDJ pyrosequencing. *Sci. Transl. Med.* **1**, 12ra23 (2009).
6. Muraro, P. A. *et al.* T cell repertoire following autologous stem cell transplantation for multiple sclerosis. *J. Clin. Invest.* **124**, 1168–1172 (2014).
7. Gros, A. *et al.* PD-1 identifies the patient-specific CD8+ tumor-reactive repertoire infiltrating human tumors. *J. Clin. Invest.* **124**, 2246–2259 (2014).
8. Beausang, J. F. *et al.* T cell receptor sequencing of early-stage breast cancer tumors identifies altered clonal structure of the T cell repertoire. *Proc. Natl. Acad. Sci. U. S. A.* **114**, E10409–E10417 (2017).
9. Emerson, R. O. *et al.* Immunosequencing identifies signatures of cytomegalovirus exposure history and HLA-mediated effects on the T cell repertoire. *Nat. Genet.* **49**, 659–665 (2017).
10. Glanville, J. *et al.* Identifying specificity groups in the T cell receptor repertoire. *Nature* **547**, 94–98 (2017).
11. Dash, P. *et al.* Quantifiable predictive features define epitope-specific T cell receptor repertoires. *Nature* **547**, 89–93 (2017).
12. DeWitt, W. S. *et al.* Human T cell receptor occurrence patterns encode immune history, genetic background, and receptor specificity. *eLife* **7**, (2018).
13. Han, A., Glanville, J., Hansmann, L. & Davis, M. M. Linking T-cell receptor sequence to functional phenotype at the single-cell level. *Nat. Biotechnol.* **32**, 684–692 (2014).
14. Morris, H. *et al.* Tracking donor-reactive T cells: Evidence for clonal deletion in tolerant kidney transplant patients. *Sci. Transl. Med.* **7**, 272ra10 (2015).
15. Theil, A. *et al.* T cell receptor repertoires after adoptive transfer of expanded allogeneic regulatory T cells. *Clin. Exp. Immunol.* **187**, 316–324 (2017).
16. Mamedov, I. Z. *et al.* Quantitative tracking of T cell clones after haematopoietic stem cell transplantation. *EMBO Mol. Med.* **3**, 201–207 (2011).
17. Rosati, E. *et al.* Overview of methodologies for T-cell receptor repertoire analysis. *BMC Biotechnol.* **17**, 61 (2017).
18. Rudolph, M. G., Stanfield, R. L. & Wilson, I. A. How TCRs bind MHCs, peptides, and coreceptors. *Annu. Rev. Immunol.* **24**, 419–466 (2006).
19. Turner, S. J., Doherty, P. C., McCluskey, J. & Rossjohn, J. Structural determinants of T-cell receptor bias in immunity. *Nat. Rev. Immunol.* **6**, 883–894 (2006).
20. Miles, J. J., Douek, D. C. & Price, D. A. Bias in the αβ T-cell repertoire: implications for disease pathogenesis and vaccination. *Immunol. Cell Biol.* **89**, 375–387 (2011).
21. Kebschull, J. M. & Zador, A. M. Sources of PCR-induced distortions in high-throughput sequencing data sets. *Nucleic Acids Res.* **43**, e143 (2015).
22. Carlson, C. S. *et al.* Using synthetic templates to design an unbiased multiplex PCR assay. *Nat. Commun.* **4**, 2680 (2013).
23. Shugay, M. *et al.* Towards error-free profiling of immune repertoires. *Nat. Methods* **11**, 653–655 (2014).

24. Peng, Q., Vijaya Satya, R., Lewis, M., Randad, P. & Wang, Y. Reducing amplification
artifacts in high multiplex amplicon sequencing by using molecular barcodes. *BMC*
Genomics **16**, 589 (2015).
25. Vander Heiden, J. A. *et al.* pRESTO: a toolkit for processing high-throughput sequencing
raw reads of lymphocyte receptor repertoires. *Bioinformatics* **30**, 1930–1932 (2014).
26. Shugay, M. *et al.* VDJtools: Unifying Post-analysis of T Cell Receptor Repertoires. *PLoS*
Comput. Biol. **11**, (2015).
27. Oakes, T. *et al.* Quantitative Characterization of the T Cell Receptor Repertoire of Naïve
and Memory Subsets Using an Integrated Experimental and Computational Pipeline Which
Is Robust, Economical, and Versatile. *Front. Immunol.* **8**, (2017).
28. Ma, K.-Y. *et al.* Immune Repertoire Sequencing Using Molecular Identifiers Enables
Accurate Clonality Discovery and Clone Size Quantification. *Front. Immunol.* **9**, 33 (2018).
29. Egorov, E. S. *et al.* Quantitative profiling of immune repertoires for minor lymphocyte
counts using unique molecular identifiers. *J. Immunol. Baltim. Md 1950* **194**, 6155–6163
(2015).
30. Giudicelli, V., Chaume, D. & Lefranc, M.-P. IMGT/GENE-DB: a comprehensive database
for human and mouse immunoglobulin and T cell receptor genes. *Nucleic Acids Res.* **33**,
D256-261 (2005).
31. Simsek, M. & Adnan, H. Effect of single mismatches at 3'-end of primers on polymerase
chain reaction. *J. Sci. Res. Med. Sci.* **2**, 11–14 (2000).
32. Wu, J.-H., Hong, P.-Y. & Liu, W.-T. Quantitative effects of position and type of single
mismatch on single base primer extension. *J. Microbiol. Methods* **77**, 267–275 (2009).
33. Wright, E. S. *et al.* Exploiting extension bias in polymerase chain reaction to improve primer
specificity in ensembles of nearly identical DNA templates. *Environ. Microbiol.* **16**, 1354–
1365 (2014).
34. Ishii, K. & Fukui, M. Optimization of annealing temperature to reduce bias caused by a
primer mismatch in multitemplate PCR. *Appl. Environ. Microbiol.* **67**, 3753–3755 (2001).
35. Bolotin, D. A. *et al.* MiXCR: software for comprehensive adaptive immunity profiling. *Nat.*
Methods **12**, 380–381 (2015).
36. Deng, L. *et al.* Structural basis for the recognition of mutant self by a tumor-specific, MHC
class II-restricted T cell receptor. *Nat. Immunol.* **8**, 398–408 (2007).
37. Arahata, K. & Engel, A. G. Monoclonal antibody analysis of mononuclear cells in
myopathies. I: Quantitation of subsets according to diagnosis and sites of accumulation and
demonstration and counts of muscle fibers invaded by T cells. *Ann. Neurol.* **16**, 193–208
(1984).
38. Fyhr, I. M., Moslemi, A. R., Lindberg, C. & Oldfors, A. T cell receptor beta-chain repertoire
in inclusion body myositis. *J. Neuroimmunol.* **91**, 129–134 (1998).
39. Cavazzana, I., Fredi, M., Selmi, C., Tincani, A. & Franceschini, F. The Clinical and
Histological Spectrum of Idiopathic Inflammatory Myopathies. *Clin. Rev. Allergy Immunol.*
52, 88–98 (2017).
40. van Dongen, J. J. M. *et al.* Design and standardization of PCR primers and protocols for
detection of clonal immunoglobulin and T-cell receptor gene recombinations in suspect
lymphoproliferations: Report of the BIOMED-2 Concerted Action BMH4-CT98-3936.
Leukemia **17**, 2257–2317 (2003).
41. Brüggemann, M. *et al.* Standardized next-generation sequencing of immunoglobulin and T-
cell receptor gene recombinations for MRD marker identification in acute lymphoblastic
leukaemia; a EuroClonality-NGS validation study. *Leukemia* **33**, 2241–2253 (2019).
42. Greenberg, S. A., Pinkus, J. L., Amato, A. A., Kristensen, T. & Dorfman, D. M. Association
of inclusion body myositis with T cell large granular lymphocytic leukaemia. *Brain* **139**,
1348–1360 (2016).

43. Greenberg, S. A. *et al.* Highly differentiated cytotoxic T cells in inclusion body myositis.
Brain J. Neurol. (2019) doi:10.1093/brain/awz207.
44. Bodenhofer, U., Bonatesta, E., Horejš-Kainrath, C. & Hochreiter, S. msa: an R package for
multiple sequence alignment. *Bioinforma. Oxf. Engl.* **31**, 3997–3999 (2015).
544
545

SUPPLEMENTARY FIGURES:

a

b

START → input unique FR3 20 mer FASTA

Compact primer set: algorithm

number of remaining unique FR3 20 mers > 0 —FALSE→ END

TRUE

calculate nt frequency at each position of remaining unique FR3 20 mers

determine max nt frequency & set matching primer position to that nt (tie: set most 5' nt)

exactly 16/20 primer nts determined

FALSE

TRUE

max nt identical between FR3s at next position —FALSE→ > 16/20 primer nts determined —TRUE→ remaining primer nts set to inosine

FALSE

set matching primer position to max nt, remaining primer nts set to inosine

TRUE

set matching primer position to that nt

< 20/20 primer nts determined

TRUE

FALSE

mismatch in 3'-5 nts between primer & unique FR3

TRUE

keep unique FR3 in next primer design cycle

FALSE

1 mismatch or less between primer & unique FR3 —TRUE→ remove unique FR3 in next primer design cycle

FALSE

Figure S1: FR3AK-seq primer design algorithm schematic. **a.** 1MM primer set design algorithm. Input is a

FASTA formatted file containing each unique FR3 20-mer nucleotide (nt) sequence. Nt frequencies at each

of the 20 FR3 positions were calculated. The position with the maximum nt frequency in the unique FR3

20-mers was used and the corresponding position in the primer was set to that nt. In the case of ties, the

most 5' position was set first. Once all 20 nts of the primer were determined, unique FR3 20-mers were

removed from the next round of primer design only if they: i) did not have a mismatch in the most 3'-5 nts

(as described in the text) and ii) had 1 mismatch or an exact match to the designed primer. The primer

design loop continued until no unique FR3 20-mer sequences remained. **b.** Compact primer set design

algorithm. Primers were designed as in **a** until 16/20 primer nts were determined. At this point, if the position

with the maximum nt frequency in the unique FR3 20-mers was identical to the number of remaining FR3

20-mers, then these nts were set to these positions. If they were not and exactly 16/20 nts in the primer

were already determined, then the nt with the maximum frequency was set to the corresponding primer

position and the remaining 3 nt positions in the primer were set to inosines. If they were not and >16/20 nts

in the primer were already determined, then the remaining nt positions were set to inosines. In this way, we

allowed for up to 3 inosines and 1 mismatch. Once all 20 nts of the primer were determined, the remaining

design was identical to **a**, where we considered each inosine a match to any nt in the FR3 regardless of its

position in the primer.

Figure S2: FR3AK-seq multiplex PCR performance. **a.** Individual primers from all three sets produce

amplicons of the expected size from human PBMC cDNA (~220 bp). Asterisks indicate inosine-containing

primers. Order of primers can be found in **Table S1**. **b.** The number of clones detected versus annealing

temperature using the Compact primer set is shown. A similar trend was observed for the 1MM primer set.

To account for differences in number of reads between samples, FASTQ sequencing files were trimmed to

identical line numbers before analysis. **c.** MA plot of 2^{10} dilution experiment for the Compact primer set

demonstrated high linear concordance (ρ=0.997), indicating negligible PCR bias. Clones with ≥10 counts

in either dataset were compared. Insets indicate number of clones above or below Y axis limits (set to 2

and −2 for visualization).

Figure S3: FR3AK-seq dual index sequencing strategy. **a.** Reverse transcription using a TCRB constant

region primer generates cDNA for use as template in **b.** PCR1 using our forward TCRB FR3 primers and a

hemi-nested TCRB constant region reverse primer. FR3 primers have 5' ad1.2 adapter sequences and the

constant region primer has a 5' ad_min adapter sequence. **c.** PCR2 using PCR1 product as a template and

indexed ad1.2 and ad_min primers, including P5 and P7 adapters for Illumina sequencing. Samples are

mixed upon completion of PCR2 prior to **d.** Next Generation Sequencing (NGS): Illumina 100-base single-

end reads along with index 5 (i5) and index 7 (i7) reads. Dual indexes are used for demultiplexing reads

from each sample and can be found in **Supplementary File S1**. No index read mismatches were permitted

during sample demultiplexing. Primer and adapter sequences can be found in **Table S1**.

Figure S4: FR3AK-seq Full and Compact primer sets perform comparably to immunoSEQ. **a.** MA plots showing technical replicates of the Full (left panel) and Compact (right panel) FR3AK-seq assays. **b.** MA plot comparing the Full (left panel) and Compact (right panel) FR3AK-seq assay against the immunoSEQ assay. Clones with ≥10 counts in either dataset were included. **c.** Relative frequencies of Donor A (black points) and Donor B (red points) in sample C determined using the Full (left panel) and Compact (right panel) FR3AK-seq assay. Insets in **a** and **b** indicate number of clones above or below Y axis limits (set to 2 and −2 for visualization)

Figure S5: a. Distributions of unique clones per patient for each IIM subgroup and controls. Kruskal-Wallis

p-value is shown. **b.** Clonal space homeostasis for each IBM donor with at least one clone present at >2 T

cell equivalents in their muscle biopsy. Criteria for clone size are shown.

Figure S6: Summary statistics of FR3AK-seq TCRB repertoires from each IIM subgroup and controls. **a.**

Stacked bar plot of J gene usage for each IIM subgroup and controls. **b.** Grand Average of Hydropathy

(GRAVY) for CDR3s from each IIM subgroup and controls. **c.** CDR3 length comparisons between each IIM

subgroup and controls.

Figure S7: MA plot demonstrating the performance of the FR3AK-seq constant region reverse PCR1 primer as compared to a set of J region reverse primers. cDNA was used as a starting material along with the 1MM forward primer set with PCR conditions as described in the Methods section. CDR3 sequences were included in this analysis if they had a clone count of ≥10 in either data set. J region primers can be found in **Table S8**.

Figure S8: MA plot demonstrates that Herculase II Fusion DNA polymerase performs comparably to KAPA 2G Fast Multiplex Mix in PCR1. CDR3 sequences were included in this analysis if they had a clone count of ≥10 in either data set.

618　**SUPPLEMENTARY TABLES:**

Table S1: FR3AK-seq human TCRB primers			
TCRB constant region primers			
Sequence		**Step**	
accagtgtggccttttgggtgtg		Reverse transcription	
cttttgggtgtgggagatctctg		Reverse primer: PCRI	
Adapter sequence for TCRB constant region PCRI primer (add to 5' end)			
Sequence		**Adapter name**	
ctggagttcagacgtgtgctcttccgatct		ad_min	
Adapter sequence for FR3 primers (add to 5' end)			
Sequence		**Adapter name**	
ggagctgtcgttcactc		AD1.2	
Full: TCRB FR3 primers			
Primer Sequence	**IMGT Accession | Associated V allele**	**Unique FR3 & Heatmap order**	**Gel order**
actccgccgtgtatctctgt	X61441|TRBV7-8*02 M11953|TRBV7-8*01 M27384|TRBV7-8*03	1	25
actctgccgtgtatctctgt	X74843|TRBV7-3*04	2	20
actcggccgtgtatctctgt	X58797|TRBV11-3*02 X61442|TRBV7-2*01 L36190|TRBV7-2*02 X58796|TRBV11-2*03 M27387|TRBV7-2*04 U03115|TRBV11-3*01 L36092|TRBV11-2*01 U07975|TRBV7-2*03 M33235|TRBV11-2*02 MGT000021|TRBV11-2*04 AB305924|TRBV11-3*04	3	47
actcagccgtgtatctctgt	M13550|TRBV7-3*05 X61440|TRBV7-3*01	4	4
actcagctgtgtatctctgt	L36092|TRBV7-4*01	5	36
actcagccgtgtatctccgt	M97943|TRBV7-3*02	6	45
actcagccgcgtatctccgt	AF009660|TRBV7-3*03	7	15
actcggccatgtatcgctgt	X58806|TRBV7-6*02 L36092|TRBV7-6*01	8	23
actcagccatgtatcgctgt	L36092|TRBV7-7*01 X57607|TRBV7-7*02	9	37
actcggccatgtatctctgt	M33233|TRBV11-1*01 L14854|TRBV7-9*07 M27385|TRBV7-9*05 M14261|TRBV7-9*04 AF009663|TRBV7-9*03 M62377|TRBV11-3*03 L36092|TRBV7-9*01	10	26

	M15564|TRBV7-9*02 X74844|TRBV7-9*06		
actcagccgtgtatctctac	U03115|TRBV17*01 IMGT000021|TRBV17*02	11	34
actcagccctgtatctctgc	U07977|TRBV4-1*01 M13855|TRBV4-1*02	12	5
acttggctgtgtatctctgt	X61444|TRBV7-1*01	13	33
actcggccctgtatctctgc	X58812|TRBV4-3*02 L06888|TRBV4-3*03 X57616|TRBV4-3*04 U07978|TRBV4-3*01	14	27
actcggccctgtatctctgt	X58811|TRBV4-2*02 U07975|TRBV4-2*01 X57611|TRBV5-5*02 X57615|TRBV5-4*02 X58803|TRBV5-8*02 L36092|TRBV5-8*01 X58804|TRBV5-4*04 L36092|TRBV5-5*01 L36092|TRBV5-4*01 AF009660|TRBV5-3*02 X58801|TRBV5-5*03 S50547|TRBV5-4*03 X61439|TRBV5-3*01	15	1
actcggccctctatctctgt	L36092|TRBV5-6*01 L36092|TRBV5-7*01	16	18
actcggccctttatctttgc	L36092|TRBV5-1*01 M14271|TRBV5-1*02	17	40
acacggcactgtatctctgc	L36092|TRBV23-1*01	18	39
actcagctgtgtacttctgt	K02546|TRBV12-4*01 X07192|TRBV12-3*01 M14264|TRBV12-4*02	19	31
actcagctgtgtatttttgt	X07223|TRBV12-5*01	20	14
actcagctttgtatttctgt	L36092|TRBV9*01 M27380|TRBV9*03 AF009660|TRBV9*02	21	19
actctgctgtgtatttctgt	L06889|TRBV3-1*02 U07977|TRBV3-1*01	22	28
actcagccatgtacttctgt	M64351|TRBV2*03 L36092|TRBV2*01 M62379|TRBV2*02	23	44
actcagccctgtacttctgt	U03115|TRBV13*01 M62378|TRBV13*02	24	9
actctggcttctatctctgt	L36092|TRBV30*01 M13554|TRBV30*04 L06893|TRBV30*05 Z13967|TRBV30*02	25	22
acgcagccatgtacctgtgt	X58800|TRBV15*02 IMGT000021|TRBV15*04	26	43
acacagccatgtacctgtgt	U03115|TRBV15*01	27	3

acgcagccatgtaccagtgt	M62376|TRBV15*03	28	42
attcggcagcttatttctgt	L36092|TRBV18*01	29	16
attctggagtttatttctgt	X57722|TRBV14*02 X06154|TRBV14*01	30	46
attcagcagtgtattttgt	L26054|TRBV16*03 L26231|TRBV16*01	31	7
agacatctgtgtacttctgt	AF009662|TRBV6-6*02 X58815|TRBV6-6*03 L33102|TRBV10-3*04 L36092|TRBV6-6*01 U17047|TRBV10-3*02 U03115|TRBV10-3*01 AF009660|TRBV6-4*02 X61446|TRBV6-1*01 L06892|TRBV6-6*05 X74848|TRBV6-6*04 L33101|TRBV10-3*03 X61653|TRBV6-4*01 L36092|TRBV6-5*01	32	29
agacatctgtgtacttgtgt	L36092|TRBV6-8*01	33	21
aaacatctgtgtacttctgt	U07978|TRBV6-3*01 X61445|TRBV6-2*01	34	24
agacatctgtatacttctgt	X61447|TRBV6-9*01	35	30
agacttctgtttacttctgt	L36092|TRBV6-7*01	36	10
agacctctgtacttctgt	L36092|TRBV27*01	37	6
agacatctatgtacctctgt	U08314|TRBV28*01	38	38
agacagctctttacttctgt	M11951|TRBV24-1*01 IMGT000021|TRBV24-1*02	39	8
agacatctgtatatttctgc	L36092|TRBV10-1*01 AF009660|TRBV10-1*02	40	35
agacatctgtgtatttctgc	L36092|TRBV10-2*01 IMGT000021|TRBV10-2*03	41	32
atacctctcagtacctctgt	L36092|TRBV25-1*01	42	41
cgacagctttctatctctgt	U48259|TRBV19*02 M97725|TRBV19*03 L36092|TRBV19*01	43	2
gcttctacatctgcagtgct	X72719|TRBV20-1*02	44	17
acagcagcttctacatctgc	D13088|TRBV20-1*06 X57604|TRBV20-1*05 M14263|TRBV20-1*04 M11955|TRBV20-1*01 X74852|TRBV20-1*07 M11954|TRBV20-1*03	45	12
acagcagcatatatctctgc	L36092|TRBV29-1*01 X04926|TRBV29-1*03 M13847|TRBV29-1*02	46	13
cccgctcccagacatctgtg	U17048|TRBV10-2*02	47	11
1MM: TCRB FR3 primers			

Primer Sequence	Heatmap Order	Gel Order
actcggccgtgtatctctgt	1	25
actcagctgtgtatctctgt	2	19
actcagccgcgtatctccgt	3	9
actcagccatgtatcgctgt	4	20
actcagccgtgtatctctac	5	26
actcagccctgtatctctgc	6	1
acttggctgtgtatctctgt	7	18
actcggccctctatctctgt	8	11
actcggccctttatctttgc	9	22
acacggcactgtatctctgc	10	30
actcagctgtgtacttctgt	11	16
actcagctgtgtatttttgt	12	28
actcagctttgtatttctgt	13	12
actctgctgtgtatttctgt	14	14
actcagccctgtacttctgt	15	5
actctggcttctatctctgt	16	27
acgcagccatgtacctgtgt	17	24
acgcagccatgtaccagtgt	18	23
attcggcagcttatttctgt	19	10
attctggagtttatttctgt	20	29
attcagcagtgtatttttgt	21	3
agacatctgtgtacttctgt	22	15
agacatctgtgtacttgtgt	23	13
agacttctgtttacttctgt	24	6
agacctctctgtacttctgt	25	2
agacatctatgtacctctgt	26	21
agacagctctttacttctgt	27	4
agacatctgtgtatttctgc	28	17
atacctctcagtacctctgt	29	34
cgacagctttctatctctgt	30	33
gcttctacatctgcagtgct	31	31
acagcagcttctacatctgc	32	7
acagcagcatatatctctgc	33	8
cccgctcccagacatctgtg	34	32

Compact: TCRB FR3 primers		
Primer Sequence	**Heatmap Order**	**Gel Order**
actcigciitgtatctctgt	1	1
actcagccgigtatctccgt	2	7
actcggccatgtatcgctgt	3	11
actcagccgtgtatctctac	4	10
actcigccctitatctitgc	5	5
acacggcactgtatctctgc	6	16
actcagctgtgtatttttgt	7	14
actctggcttctatctctgt	8	12
acicagccatgtacciitgt	9	4
attcggcagiitatttitgt	10	8

Primer Sequence		
attctggagtttatttctgt	11	15
agacatctitgtacititgt	12	3
agaciictitttacttctgt	13	13
actcagciitgtacttctgt	14	2
agacatctgtitatttctgc	15	6
atacctctcagtacctctgt	16	20
cgacagctttctatctctgt	17	19
gcttctacatctgcagtgct	18	17
acagcagcttitaiitctgc	19	9
cccgctcccagacatctgtg	20	18

PCR2 primers	
Primer Sequence: X refers to barcode nucleotides	**Adapters**
caagcagaagacggcatacgagatXXXXXXXXgtgactggagttcagacgtg tgctcttccgatct	P7_ad_min
aatgatacggcgaccaccgagatctacacXXXXXXXXacactctttccgatc tggagctgtcgttcactc	P5_AD1.2

619   **Table S1:** Human TCRB FR3AK-seq primers and adapter sequences.

Table S2: FR3AK-seq cost per sample			
Step	**Item**	**Price**	**Cost per sample**
Reverse Transcription	Invitrogen Superscript III first-strand synthesis system	$467.00 (50 reactions)	$9.34
cDNA clean up	Zymo Research Oligo Clean & Concentrator	$423.00 (384 preps)	$1.10
PCR1	KAPA2G Fast Multiplex Mix	$295.88 (6.25 mL/ 624-20μL reactions)	$0.47
PCR2	Herculase II Fusion DNA Polymerase	$475.00 (400 uL, 2000-20 uL reactions)	$0.24
Reverse Transcription, 1MM PCR1, PCR2	Primers	$204.75 (36 primers, 6250 reactions PCR1/RT) $11.38 (2 primers, 5000 reactions PCR2)	$0.03 $0.002
Plastics	Pipet tips, tubes, etc.	$1.00/sample	$1.00
Sequencing (HiSeq 2500)	100 nt single-end Illumina reads, 213 samples at ~1x10^6 reads per sample	$1600.00 (100bp, per lane)	$7.51
TOTAL			**$19.69**

620   **Table S2:** Cost per sample for FR3AK-seq. HiSeq cost calculations were based on the Johns Hopkins

621   Genetics Resources Core Facility High Throughput Sequencing Center (internal pricing) at an estimated

622   1×10^6 reads per sample. Note that these calculations do not include overhead, personnel costs, or time

623   spent by the researcher.

Table S3: Idiopathic Inflammatory Myopathy and Control Muscle Biopsy Cohort						
	Controls (n=21)	DM (n=40)	ASyS (n=21)	IMNM (n=49)	IBM (n=14)	Total (n=145)
Female	71% (15)	53% (9)	89% (8)	67% (18)	54% (7)	66% (57)
White	82% (9)	71% (12)	44% (4)	56% (15)	83% (10)	66% (50)
Black	18% (2)	18% (3)	44% (4)	37% (10)	17% (2)	28% (21)
Other race	0% (0)	12% (2)	11% (1)	7% (2)	0% (0)	7% (5)
Age	50.6 (13.9)	59.7 (13.4)	46.2 (8.4)	51.8 (13.4)	66.1 (11.6)**	54.6 (14.0)

624   * $p<0.05$,** $p<0.01$,*** $p<0.001$
625   **Table S3:** Description of the IIM muscle biopsy cohort. Chi-squared or Fisher´s exact tests were used to

626   compare the dichotomous variables of each one of the clinical groups with the rest. Student's T test was

627   used to compare the age of each one of the clinical groups with the rest.

Table S4: GLIPH summary	
Input IIM FR3AK-seq data	
Number of samples with non-bystander T cell clones	122 (23 excluded)
Number of non-bystander T cell clones included in analysis	11351
Number of bystander T cell clones	278121
GLIPH Output	
Total number of GLIPH clusters	527
Total number of GLIPH clusters with at least 3 individuals contributing a T cell clone	143
Number of non-bystander T cell clones clustered by GLIPH	2125

628   **Table S4:** Summary of IIM and control muscle biopsy FR3AK-seq data used in GLIPH analysis. Bystander

629   clones refer to those appearing at a frequency of less than or equal to 1 cellular equivalent.

Table S5: Comparisons between platforms				
Platform	**ImmunoSEQ hsTCRB service (Adaptive Biotechnologies)**	**ImmunoSEQ hsTCRB kit (Adaptive Biotechnologies)**	**Immunoverse (ArcherDX)**	**FR3AK-seq**
Method	Multiplex PCR with bias correction	Multiplex PCR with bias correction	UMI	Multiplex PCR
Time	• <1 day to prepare sample and ship to company • Processed data within ~2-3 weeks	• <1 day to prepare sample • ~1 day to obtain processed data	• ~2 days for library preparation • ~1 day to obtain processed data	• ~1 day for library preparation • ~1 day to obtain processed data
Hands-on user steps	• Reverse transcription (if using cDNA)	• 3 step library preparation (if using cDNA) • Sequencing • Upload fastq files to immunoSEQ analyzer for bias correction	• 9 step library preparation • Sequencing • Upload fastq files to ArcherDX portal for deduplication analysis	• 3 step library preparation • Sequencing • MiXCR analysis
Cost per sample	Reverse transcription	$9.87	Kit	$629.00	Kit	$240.00	$19.69 (see **Table S2**)
	Deep Sequencing Service	$1100	MiSeq	$44.00	MiSeq	$44.00	
	TOTAL: $1109.87		**TOTAL**: $642.02		**TOTAL:** $284.00		**TOTAL:** $19.69

635 **Table S5:** Comparisons between Adaptive Biotechnologies' immunoSEQ assay (both service and kit),

636 ArcherDX's Immunoverse assay, and FR3AK-seq. All prices are approximate. MiSeq cost calculations were

637 based on the Johns Hopkins Transcriptomics and Microarray Core (JHU internal pricing) at an estimated

638 $1x10^6$ reads per sample.

Table S6: FR3AK-seq *Mus musculus* TCRB primers	
***Mus musculus* TCRB constant region primers**	
Sequence	**Step**
tggctcaaacaaggagaccttgg	Reverse transcription
ggtggagtcacatttctcagatcc	Reverse primer: PCRI
Full: *Mus musculus* TCRB FR3 primers	
Primer Sequence	**IMGT Accession | Associated V allele**
acagcagtatctatttctgt	X16695|TRBV30*01
agatgaccgttttttctctgt	X01643|TRBV19*02
actctgctgtgtatctctgt	AJ249819|TRBV4*02

	X56725|TRBV4*01
agacatctttgtacttctgt	AE000663|TRBV13-1*02 M15618|TRBV13-1*01
aaggcagaaccttgtactgc	X01642|TRBV1*02
agatggccgttttctctgt	AE000664|TRBV19*01 AJ249821|TRBV19*03
actctggcttctacctctgt	X03277|TRBV31*01 M26418|TRBV31*02
actcagctgtctattttgt	AJ249820|TRBV5*05 M20878|TRBV5*04 X00438|TRBV5*02 X02779|TRBV5*03 AE000663|TRBV5*01
acagaggcttatatctctgt	AE000664|TRBV20*01 M11859|TRBV20*02
attcagctgtgtacttctgt	X16691|TRBV21*01
actcagcggtgtatctttgt	X14388|TRBV16*04 M15459|TRBV16*03 M13670|TRBV16*02 L29434|TRBV16*01
actcagcactgtacttgtgc	AE000664|TRBV23*01
agggcagaaccttgtactgc	AE000663|TRBV1*01
agacagctgtatatttctgt	M15616|TRBV13-3*01
actcagctgtgtacttctgt	X03865|TRBV3*02 AE000663|TRBV3*01
actcagcactgtacctctgt	L48997|TRBV24*04 M16203|TRBV24*03 AJ249822|TRBV26*02 K02548|TRBV26*01 M61184|TRBV24*01
actcagctgtgtatctgtgt	AE000664|TRBV15*01
agacatcagtgtacttctgt	AJ250103|TRBV13-2*05 M26417|TRBV13-2*03 AJ249823|TRBV13-2*04 M15617|TRBV13-2*01 M27350|TRBV13-2*02
attctgccatgtacctctgt	AE000664|TRBV17*01
agacatctgtgtacttctgt	X00696|TRBV29*02 AE000664|TRBV29*01
actcggccacatacttctgt	AE000663|TRBV2*01
actctgccgtgtacttctgt	M15613|TRBV12-2*01 X02782|TRBV12-2*02
actctgctatgtacttctgt	M30881|TRBV12-1*02 M15614|TRBV12-1*01
acacagccacctatctctgt	AE000664|TRBV14*01

640 **Table S6:** *Mus musculus* TCRB FR3AK-seq primers. Adapter sequences can be found in **Table S1**.

Table S7: FR3AK-seq human TCRA, TCRG, and TCRD primers	
human TCRA: constant region primers	
Sequence	**Step**
cggtgaataggcagacagacttg	Reverse transcription
cagacttgtcactggatttagag	Reverse primer: PCRI
Full: human TCRA FR3 primers	
Primer Sequence	**IMGT Accession | Associated V allele**
atgctgctgtgtactactgc	AE000660|TRAV26-2*01
actcagccacctacttctgt	M17660|TRAV23/DV6*02 AE000660|TRAV23/DV6*01 M97704|TRAV23/DV6*03
actcagctatctacttctgt	AE000659|TRAV5*01
atgctgctgagtacttctgt	AE000659|TRAV8-3*01 AE000660|TRAV8-7*01 M35617|TRAV8-3*02
atgcgtctgagtacttctgt	L06885|TRAV8-3*03
atacaggccactacctctgt	X04957|TRAV27*02
actccgctgtgtacttctgt	AE000659|TRAV9-1*01
actcagccacctacctctgt	X58736|TRAV21*02 AE000660|TRAV21*01
atacaggcctctacctctgt	AE000660|TRAV27*01 D13075|TRAV27*03
actctgccgtgtactactgc	AE000660|TRAV18*01
actcagctgtctacttttgt	M17658|TRAV13-2*02 AE000659|TRAV13-2*01
actcagcaatgtatttctgt	AE000659|TRAV14/DV4*02 M21624|TRAV14/DV4*03
atgtaggaacctacttctgt	AE000660|TRAV25*01
attcagccacctatttctgt	AE000659|TRAV7*01
atgcaggcatctacctctgt	AE000660|TRAV34*01
acacagctgagtacttctgt	AE000659|TRAV8-1*01
attcagccacatacctctgt	M17661|TRAV24*02
atgtaggcatctacttctgt	AE000660|TRAV35*01 X58738|TRAV35*02
acgcggctgagtacttctgt	M12959|TRAV8-4*04 X63455|TRAV8-4*05 M17650|TRAV8-2*02 M12423|TRAV8-4*02 D13077|TRAV8-4*03 AE000659|TRAV8-2*01 AE000659|TRAV8-4*01 K02777|TRAV8-4*06
actcaggcgtttatttctgt	AE000660|TRAV22*01
actcggccgtctacctctgt	Z46643|TRAV36/DV7*04 X58767|TRAV36/DV7*03 X61070|TRAV36/DV7*02
acactgcgatgtatttctgt	AE000661|TRAV38-1*01 L06880|TRAV38-1*04 M95394|TRAV38-1*03

	M64355|TRAV38-1*02
actcagcaatgtacttctgt	M21626|TRAV14/DV4*01
actcagcagtatacttctgt	AE000660|TRAV19*01
actcagccgtgtactactgt	X73521|TRAV40*01
acacggctgagtacttctgt	X02850|TRAV8-6*01 AE000659|TRAV8-6*02
actcagccacatacctctgt	AE000660|TRAV24*01
actcagctacctacctctgt	AE000659|TRAV6*01 X58747|TRAV6*02 Z49060|TRAV6*03 Y10410|TRAV6*05 Y10409|TRAV6*04
actctgcagtgtacttctgt	S81645|TRAV29/DV5*02 AE000660|TRAV29/DV5*01
acactgctgtgtactattgc	AE000660|TRAV26-1*01 M27371|TRAV26-1*02 L06886|TRAV26-1*03
actcggccatctacctctgt	AE000660|TRAV36/DV7*01
actctgcctcttacttctgc	X04939|TRAV1-1*02 AE000658|TRAV1-1*01
tctctgccacctacttctgt	AE000661|TRAV39*01
actctgcctcttacctctgt	AE000658|TRAV1-2*01
acactgctgtgtactactgc	AE000658|TRAV4*01
actccgctttgtacttctgt	AE000658|TRAV3*01
actcagccacttatctctgt	X70305|TRAV20*04 S60789|TRAV20*03 AE000660|TRAV20*01 X68696|TRAV20*02
atgctgctgtttactactgt	AE000658|TRAV2*01 M17659|TRAV2*02
acccggctgagtacttctgt	M17665|TRAV8-4*07
attcagccacctacctctgt	AE000659|TRAV12-2*01 AE000659|TRAV12-1*01 M17657|TRAV12-1*02 M17656|TRAV12-3*02 M81774|TRAV12-2*02 X04946|TRAV12-2*03 AE000659|TRAV12-3*01
actctgccgtctacatctgt	AE000661|TRAV41*01
actcaggaacctacttctgc	L06883|TRAV30*03 AE000660|TRAV30*01 X58768|TRAV30*02
atgccgcgatgtatttctgt	AE000661|TRAV38-2/DV8*01
attcagcctcctacatctgt	AE000659|TRAV10*01
actcggctgtctacttctgt	AE000659|TRAV13-1*01 X04954|TRAV13-1*02
acactgcttcttacttctgt	AE000660|TRAV17*01
actcagcggtgtacttctgt	L06882|TRAV9-2*04 X58745|TRAV9-2*02 L06881|TRAV9-2*03

	AE000659|TRAV9-2*01
actcagccatgtattactgt	AE000659|TRAV16*01

1MM: human TCRA FR3 primers
actccgctgtgtacttctgt
actcagcggtgtacttctgt
actcggccgtctacctctgt
actcagccacctacctctgt
acgcggctgagtacttctgt
actcagcagtatacttctgt
actcagctatctacttctgt
actctgccgtctacatctgt
actcagcaatgtacttctgt
actctgccgtgtactactgc
acactgcttcttacttctgt
attcagccacctatttctgt
actctgcctcttacctctgt
actcggctgtctacttctgt
atacaggcctctacctctgt
actcagccgtgtactactgt
acactgcgatgtatttctgt
atgctgctgtgtactactgc
attcagcctcctacatctgt
actctgcagtgtacttctgt
atgccgcgatgtatttctgt
atgctgctgtttactactgt
actcagccacttatctctgt
acactgctgtgtactactgc
atgcaggcatctacctctgt
actcagccatgtattactgt
actctgcctcttacttctgc
atgtaggcatctacttctgt
acactgctgtgtactattgc
actcaggcgtttatttctgt
actcagctgtctacttttgt
actcaggaacctacttctgc
atgcgtctgagtacttctgt
atgctgctgagtacttctgt
atgtaggaacctacttctgt
acacagctgagtacttctgt
tctctgccacctacttctgt
attcagccacatacctctgt

Compact: human TCRA FR3 primers
actcigcigtitacttctgt
actcagccaiitacitctgt
acicigctgagtacttctgt
actcagccitgtaitictgt
actctgccgiitactictgc
aticaggciictacctctgt
aiictgctgtgtactaitgc

attcagccicctaiitctgt	
acactgcgatgtatttctgt	
atgtaggiaictacttctgt	
actcaggcgtttatttctgt	
actctgcctcttacctctgt	
actcagctgtctactttgt	
atgccgcgatgtatttctgt	
actcaggaacctacttctgc	
atgcgtctgagtacttctgt	
acactgcttcttacttctgt	
tctctgccacctacttctgt	
atgctgctgtttactactgt	

human TCRG: constant region primers	
Sequence	**Step**
tatgttccagccttctggag	Reverse transcription
gccttctggagctttgtttcagc gccttctggagttttgtttcagc	Reverse primers: PCRI

Full: human TCRG FR3 primers	
Primer Sequence	**IMGT Accession | Associated V allele**
atgaggtggtgtaccactgt	Y11227|TRGV11*01 AF159056|TRGV11*02
attctgggttctattactgt	M12949|TRGV1*01
actttggggtctattactgt	IMGT000011|TRGV2*03
actctggggtctattactgt	M27337|TRGV2*02 M13429|TRGV2*01 M13434|TRGV8*01
acatagctacctactactgt	X15274|TRGV9*02 X07205|TRGV9*01
acatggccgtttactactgt	X74798|TRGV10*02 X07206|TRGV10*01
attctggggtctattactgt	X13355|TRGV5*01 X04038|TRGV3*02 M13430|TRGV3*01
actctggagtctattactgt	X13354|TRGV4*02 X15272|TRGV4*01

1MM: human TCRG FR3 primers	
actctggggtctattactgt	
atgaggtggtgtaccactgt	
acatggccgtttactactgt	
acatagctacctactactgt	
attctgggttctattactgt	

Compact: human TCRG FR3 primers	
aititggigtctattactgt	
acatagctacctactactgt	
atgaggtggtgtaccactgt	
acatggccgtttactactgt	

human TCRD: constant region primers	
Sequence	**Step**

Primer Sequence	IMGT Accession / Description
gaattccttcaccagacaagcgac	Reverse transcription
ccagacaagcgacatttgttcc	Reverse primer: PCRI
Full: human TCRD FR3 primers	

Primer Sequence	IMGT Accession | Associated V allele
attcagcaaagtactttgt	M22198|TRDV1*01
atgccgcgatgtatttctgt	AE000661|TRAV38-2/DV8*01
actctgcagtgtacttctgt	S81645|TRAV29/DV5*02 AE000660|TRAV29/DV5*01
actcggccatctacctctgt	AE000660|TRAV36/DV7*01
actcagccacctacttctgt	M97704|TRAV23/DV6*03 Y10411|TRAV23/DV6*04 AE000660|TRAV23/DV6*01 M17660|TRAV23/DV6*02
actcagcaatgtatttctgt	AE000659|TRAV14/DV4*02 M21624|TRAV14/DV4*03
actcggccgtctacctctgt	Z46643|TRAV36/DV7*04 X58767|TRAV36/DV7*03 X61070|TRAV36/DV7*02
acagtgccacttactactgt	M23326|TRDV3*01 X15261|TRDV3*02
atgaagggtcttactactgt	X15207|TRDV2*01 Y13426|TRDV2*02 AE000661|TRDV2*03
actcagcaatgtacttctgt	M21626|TRAV14/DV4*01
1MM: human TCRD FR3 primers	
actcggccgtctacctctgt	
actcagcaatgtacttctgt	
actctgcagtgtacttctgt	
atgccgcgatgtatttctgt	
atgaagggtcttactactgt	
actcagccacctacttctgt	
acagtgccacttactactgt	
attcagcaaagtactttgt	
Compact: human TCRD FR3 primers	
actcggccitctacctctgt	
actcagcaaiitaittctgt	
atgccgcgatgtatttctgt	
actctgcagtgtacttctgt	
atgaagggtcttactactgt	
acagtgccacttactactgt	
attcagcaaagtactttgt	

Table S7: Human TCRA, TCRG, and TCRD FR3AK-seq primers. Adapter sequences can be found in

Table S1

Table S8: FR3AK-seq human TCRB J region primers	
Primer Sequence	**IMGT Accession | Associated J allele**
ccagcactgagagccgggtc	X02987|TRBJ2-4*01
ccagaaccaggagtcctccg	X02987|TRBJ2-2P*01
cgagcaccaggagccgcgtg	X02987|TRBJ2-5*01
ccagtacggtcagcctagag	X02987|TRBJ2-2*01
cgagcactgtcagccgggtg	X02987|TRBJ2-3*01
ctgtgaccgtgagcctggtg	X02987|TRBJ2-7*02 M14159|TRBJ2-7*01
ctacaacggttaacctggtc	K02545|TRBJ1-2*01
ctacaacagtgagccaactt	M14158|TRBJ1-3*01
ctgtcacagtgagcctggtc	M14158|TRBJ1-6*01 L36092|TRBJ1-6*02
ctacaactgtgagtctggtg	K02545|TRBJ1-1*01
ctaggatggagagtcgagtc	M14158|TRBJ1-5*01
ccaagacagagagctgggtt	M14158|TRBJ1-4*01
ccagcacggtcagcctgctg	X02987|TRBJ2-6*01
ctagcacggtgagccgtgtc	X02987|TRBJ2-1*01

646   **Table S8:** Human TRBJ FR3AK-seq primers. Adapter sequences can be found in **Table S1**.

Table S9: Software parameters	
MiXCR v2.1.11	
Parameter	**Rationale**
assemble -OaddReadsCountOnClustering=true	makes error correction algorithms to combine clone abundancies (all "children" clone counts added to head clone)
assemble --index	For IIM samples only; links reads in fastq file to CDR3s
exportClones --chains TRB	exports TCRB chain sequences only
GLIPH 1.0	
Algorithm step **--inputs required**	**Parameters** **Input file(s) column names**
1) ./gliph/bin/gliph-group-discovery.pl --tcr TCR_TABLE	default TCR_TABLE = only CDR3b sequences
2) ./gliph/bin/gliph-group-scoring.pl --convergence_file TCR_TABLE-convergence-groups.txt --clone_annotations FULL_TCR_TABLE --motif_pval_file TCR_TABLE.minp.ove10.txt	default TCR_TABLE-convergence-groups.txt = from step 1 FULL_TCR_TABLE = CDR3b, Patient, Counts TCR_TABLE.minp.ove10.txt Counts = from step 1

659 **Table S9:** MiXCR and GLIPH parameters used in analysis. Default parameters were used except as noted.

660 Only CDR3b sequences were used with the gliph-group-discovery algorithm. The gliph-group-scoring

661 algorithm was used to match patient and counts with original CDR3b sequences.

Table S10: Summary of Donor A clones across platforms			
Platform	**ImmunoSEQ hsTCRB service (Adaptive Biotechnologies)**	**Immunoverse (ArcherDX)**	**FR3AK-seq**
Total clones detected	394797	67274	362288
Number of singletons (clone count = 1)	345880	20570 (pre-deduplication) 31668 (post-deduplication)	198635
Number of clones present at ≥ 10 counts	2702	10689 (pre-deduplication) 1080 (post-deduplication)	7189

663 **Table S10:** Summary of Donor A CDR3 sequences detected across platforms.