Gut microbiome response to a modern Paleolithic diet in a Western lifestyle context

Barone M[a], Turroni S[a], Rampelli S[a], Soverini M[a], D'Amico F[a], Biagi E[a], Brigidi P[a], Troiani E[b]

Candela M[a]

[a] Unit of Microbial Ecology of Health, Department of Pharmacy and Biotechnology, University of

Bologna, Italy; [b] Primary Care Unit and Territorial Health, Social Security Institute, Republic of

San Marino.

CONTACT: Silvia Turroni silvia.turroni@unibo.it Unit of Microbial Ecology of Health,

Department of Pharmacy and Biotechnology, University of Bologna, via Belmeloro 6, 40124, Italy

ABSTRACT

The progressive reduction of gut microbiome (GM) biodiversity along human evolutionary history

has been found to be particularly exacerbated in Western urban compared to traditional rural

populations, and supposed to contribute to the increasing incidence of chronic non-communicable

diseases. Together with sanitation, antibiotics and C-section, the Western diets, low in microbiota-

accessible carbohydrates (MACs) while rich in industrialized and processed foods, are considered

one of the leading causes of this shrinkage. However, significant questions remain unanswered,

especially whether high-MAC low-processed diets may be sufficient to recover GM diversity in

Western urban populations. Here, we profiled the GM structure of urban Italian subjects adhering to

the modern Paleolithic diet (MPD), a dietary pattern featured by high consumption of MACs and

low-to-zero intake of refined sugars and processed foods, and compared data with other Italian

individuals following a Mediterranean Diet (MD), as well as worldwide traditional hunter-gatherer

populations from previous publications. Notwithstanding a strong geography effect on the GM

structure, our results show an unexpectedly high degree of GM biodiversity in MPD subjects, which

well approximates that of traditional populations. Increasing the consumption of MACs at the

expense of refined sugars, and minimizing the intake of processed foods, both hallmarks of the

MPD, could be the key to rewild the Western microbiota, counteracting the loss of GM diversity

and thus restoring evolutionarily important functionality to our gut for improved human health.

Keywords: microbiome, Paleolithic diet, Western lifestyle, MACs, xenobiotics.

INTRODUCTION

In order to understand the specificities of the human microbiome assembly, extensive meta-analyses

of human and non-human primate microbiomes have been recently carried out.[1, 2] This comparative

approach has highlighted the reduction of individual biodiversity as one of the distinctive features

of the human gut microbiome (GM).[1] Interestingly, this hallmark has been found to be exacerbated

in Western urban populations, which show even more marked compression of personal diversity

than traditional and rural counterparts.[3-6] Consistent with the disappearing microbiota hypothesis,[7]

the dramatic shrinkage of individual GM diversity in Western urban populations depicts a

maladaptive microbiome state that has been supposed to contribute to the rising incidence of

chronic non-communicable diseases, such as obesity, diabetes, asthma and inflammatory bowel

disease.[8-11] Consequently, in recent years, a large body of research has been devoted to

understanding the mechanisms leading to the diversity loss in the Western urban GM. It is in this

scenario that the multiple-hit hypothesis has been advanced.[8] According to this theory, the

progressive reduction of human GM diversity has occurred at multiple stages along the recent

transition to modern urban societies, and several aspects typical of the urbanization process - such

as sanitation, antibiotics, C-section and Western diet - have been pointed out as contributing factors.

In particular, the reduction in quantity and diversity of Microbiota-Accessible Carbohydrates

(MACs) in the diet has been considered one of the leading causes of the disappearing GM in

Western urban populations.[8] Recently defined, MACs include fermentable fibers that - indigestible

by the host - become available as an energy source for a specific GM fraction enriched in

Carbohydrate Active Enzymes (CAZymes).[8, 12] Moreover, food additives, emulsifiers and

xenobiotics – ubiquitous in industrially processed foods – have recently been shown as important

additional drivers of GM diversity shrinkage.[13]

All currently available studies exploring the disappearing GM are based on the comparison between

Western urban and traditional rural populations.[3-6, 14-16] Consistently, the observed GM differences

are likely to be the result of the combined action of several covariates in addition to the diet – i.e.

ethnicity, geographical origin, climate, subsistence, medication, hygiene and life sharing – and do

not allow to weight the importance of the single determinants. Indeed, to the best of our knowledge,

no study has been specifically designed to dissect the role of MACs deprivation and xenobiotics

exposure as driving factors forcing the compression of GM diversity in Western urban populations.

In the last few years, the Modern Paleolithic Diet (MPD), with high intake of vegetables, fruit, nuts,

eggs, fish and lean meat, while excluding grains, dairy products, salt and refined sugar, has attracted

substantial public attention in the Western world because of its potential multiple health benefits.[17-

20] Being enriched in MACs and completely excluding industrially processed food, the MPD pattern

represents a model of Western urban diet ideal for disentangling the impact of MACs and food

xenobiotics on the human GM.

In the present work, we profiled the GM structure of 15 Italian subjects following the MPD and

compared it with that of Italian individuals largely adhering to the Mediterranean Diet (MD) from

our previous works.[5, 21] Notwithstanding the small sample size, our GM exploratory study gave us

the unique opportunity to assess to what extent the consumption of high amounts of MACs along

with the exclusion of industrially processed food, may counteract the GM diversity reduction as

observed in Western urban populations. Indeed, the comparison between MPD and Western diets in

subjects living in the same country allows excluding the impact of confounding drivers of GM

variation, such as geography, ethnicity, medication, hygiene and subsistence.[14, 15, 21] In order to

extend the GM comparison at the meta-population level, we expanded our meta-analysis by

including publically available microbiome data from traditional hunting and gathering populations

showing high GM diversity, such as the Hadza from Tanzania, from our previous publication,[5] the

Matses from Peru,[6] and the Inuit from the Canadian Arctic.[22]

According to our data, increased consumption of unprocessed foods and dietary MACs, as observed

in MPD individuals, is sufficient, even in a Western context, to recover the levels of GM diversity

typically found in traditional rural populations. Although the mechanisms underlying the human-

microbiome interactions are still far from being fully understood, the possibility of rewilding the

modern microbiota through the diet could be the key to restore evolutionarily important

functionality to the gut, ultimately improving our health.

RESULTS

Diet and gut microbiome structure in Italian adults following the modern Paleolithic diet

Fifteen healthy individuals who have been following the MPD for at least one year were recruited

from different urban areas across Italy (Lombardia, Piemonte, Emilia-Romagna, Toscana, Umbria,

Lazio, Campania, Molise, Puglia and Calabria regions). Specifically, 3 female and 12 male adults,

with an average age of 39.2 years (range, 26 – 57) and average Body Mass Index (BMI) of 22.1

kg·m^{-2} (range, 19.4 – 25.7), were enrolled in our cohort (**Table S1**).

The MPD adopted by the 15 subjects is mainly based on the consumption of unprocessed foods,

with high intake of vegetables, fruit, nuts, eggs, fish and lean meat, while excluding grains, dairy

products, salt and refined sugar. The daily total calorie intake, as well as that of macro- and micro-

nutrients, assessed through 7-day weighted food intake records (7D-WRs), are reported in **Table**

S2. The average daily energy intake of the enrolled cohort is 1,843.45 kcal (range, 1,563 – 2,186

kcal). The percentage of macronutrients is distributed as follows: fat, 51.02%; protein, 30.14%;

carbohydrate, 18.84% (**Figure 1A**). With regard to lipids, 51.65% of total calories are from

monounsaturated fatty acids (MUFAs), 30.93% from saturated fatty acids (SFAs) and 17.42% from

polyunsaturated fatty acids (PUFAs) (**Figure 1B**). The average daily fiber intake is 14.64 g/1000

kcal.

The GM structure of MPD Italian adults was profiled through 16S rRNA gene sequencing of fecal

DNA. A total of 864,439 high-quality reads (mean ± sd, 57.6 ± 19.7; range, 25,142 – 95,924) were

generated and clustered in 7,483 OTUs. The phyla Firmicutes (relative abundance, mean ± sem,

65.1 ± 2.1 %) and Bacteroidetes (24.6 ± 2.2 %) dominate the gut microbial ecosystem, with

Proteobacteria (4.4 ± 1.6 %), Actinobacteria (3.4 ± 0.8%) and Verrucomicrobia (1.2 ± 0.5 %) as

minor components. At family level, *Ruminococcaceae* (26.7 ± 1.7 %), *Lachnospiraceae* (18.7 ± 1.4

%), *Bacteroidaceae* (13.7 ± 1.8 %) and *Prevotellaceae* (7.4 ± 2.4 %) are the dominant GM

constituents. The most abundant (≥ 5%) bacterial genera are *Bacteroides*, *Prevotella*, and

Faecalibacterium, while *Coprococcus*, *Ruminococcus*, *Blautia*, *Lachnospira*,

Phascolarctobacterium, *Streptococcus*, *Roseburia*, *Akkermansia*, *Oscillospira* and *[Eubacterium]*

represent minor components of the microbial ecosystem (range 4.4 ± 0.7 % - 1.0 ± 0.4 %) (**Figure

2**).

Gut microbiome diversity in MPD Italian adults and comparison with other Western urban

populations and traditional communities

In order to investigate whether the adherence to the MPD is sufficient to promote a more diverse

GM ecosystem - even in a Western urban context - we compared the GM diversity between the 15

MPD subjects and 143 urban Italians with different level of adherence to the MD, whose GM

composition was described in De Filippis *et al.*[21] and Schnorr *et al.*[5] Moreover, to extend the meta-

analysis to a global level, the GM structural profiles of the following traditional hunter-gatherer

populations were included: 27 Hadza from Tanzania,[5] 25 Matses from Peru[6] and 21 Inuit from

Canada.[22] According to our findings, significant differences in the GM biodiversity among the

study groups were detected (Simpson index, P-value = 2.6×10^{-15}; Shannon index, P-value = 2.2

$\times 10^{-16}$; Kruskal-Wallis test) (**Figure 3**). Interestingly, the GM diversity observed for MPD

subjects far exceeds that of urban Italians adhering to the MD (Simpson index, P-value = 2.5×10^{-7};

Shannon index, P-value = 6.1×10^{-9}; Wilcoxon test), and is even greater than that observed in

Matses (P-value = 0.0082; 0.0039) and Inuit (P-value = 0.00075; 0.0027). On the other hand, no

significant difference was found between MPD individuals and the Hadza (P-value = 0.39; P-value

= 0.26).

The PCoA based on Bray-Curtis distances was next used to assess overall genus-level

compositional differences in the GM structure between study groups. Our data show clear

separation of GM profiles by provenance and, within the Italian cohort, by dietary pattern (P-value

$< 1 \times 10^{-5}$, permutation test with pseudo-F ratios) (**Figure 4A**). Interestingly, MPD subjects show a

low level of interpersonal GM variation (Bray-Curtis distances, mean ± SD, 0.42 ± 0.095),

approximating that observed for the Hadza (0.36 ± 0.092) (**Figure 4B**). Finally, to identify the

bacterial drivers with a statistically significant contribution (permutation correlation test, P-value <

0.001) to the sample ordination, we superimposed the genus relative abundance on the PCoA plot

(**Figure S1**). According to our data, the microorganisms characterizing the Italian cohort are

Bacteroides, *Collinsella*, *Coprococcus* and *Blautia*. The genera *Clostridium, Prevotella*,

[Prevotella], *Catenibacterium* and *Oscillospira* were found to be associated with Hadza and

Matses, while *Sutterella* and *Parabacteroides* with Inuit.

DISCUSSION

Herein we compared the GM compositional structure and diversity of 15 urban Italian adults

adhering to the MPD with previously published data from 143 urban Italian adults largely adhering

to the MD[5, 21] and 73 traditional hunter-gatherers, including 27 Hadza from Tanzania,[5] 25 Matses

from Peru[6] and 21 Inuit from Canada.[22] According to our findings, the study groups segregate by

geographical origin, with a further separation within the Italian cohort reflecting the diet pattern

(MPD vs MD). This provenance-dependent effect on the human GM structure probably involves

the concomitant action of several covariates, which concur in shaping the GM structure, such as

geography, ethnicity, lifestyle and dietary habits. The Italian origin of the GM seems to be defined

by a higher abundance of *Bacteroides*, *Collinsella*, *Coprococcus* and *Blautia*, bacterial genera

commonly found within Western microbiomes.[3-6] According to the literature, the separation due to

geography seems to be less evident among the traditional populations, with Matses and Hadza

sharing a high abundance of *Prevotella*.[5, 6]

These data confirm recent findings that demonstrate the predominance of host location and

ethnicity, with respect to diet, as determinants of human GM variation.[14, 15] However, despite the

overall Western-like configuration, the MPD-associated GM structure stands out from that of

Italians adhering to the MD for a much higher degree of biodiversity, which well approximates that

observed in traditional hunter-gatherers. Since the Italian subjects of our cohort share the

provenance and all that it entails, including the lifestyle, it can be hypothesized that the MPD-

associated bloom in GM diversity is accounted for by the peculiarities of the MPD compared to the

MD. Though the two diets are similar in many respects – i.e. high intake of fruit, vegetables, fish

and nuts, as well as low glycemic load – the MPD is in fact distinguished by: (i) greater

consumption of MACs, mainly from wild plant foods; (ii) total exclusion of industrially processed

products; (iii) higher intake of unsaturated fatty acids, especially MUFAs, from olive oil, nuts and

meat; (iv) no consumption of foods containing refined sugars.[17-20] It is, therefore, tempting to

speculate that these MPD distinctive features may be sufficient to support the consolidation of a

highly diversified GM layout, thus counteracting the loss of GM biodiversity, typically observed in

Western urban populations as compared to traditional communities.[3-6]

In conclusion, our findings shed some light on the possibility of recovering GM diversity in

Western urban populations through diet. Increasing the consumption of MACs at the expense of

refined sugars, and minimizing the intake of processed foods, both hallmarks of the MPD, could

indeed act synergistically in facilitating the regain of an eubiotic level of GM diversity. Moreover,

the high intake of MUFAs, as found in the MPD, suggests that these fatty acids could play a role in

supporting the increase in GM diversity, which is worthy of being further explored in larger

cohorts.

MATERIALS AND METHODS

Subjects and sample collection

Fifteen healthy individuals following a MPD for at least one year were recruited from different

urban areas across Italy (Lombardia, Piemonte, Emilia-Romagna, Toscana, Umbria, Lazio,

Campania, Molise, Puglia and Calabria regions). Anthropometric data and stool samples were

collected from each participant who had not taken antibiotics in the previous 3 months. Fecal

samples were immediately frozen at -20°C and then delivered to the laboratory of the Microbial

Ecology of Health Unit (Dept. Pharmacy and Biotechnology, University of Bologna, Bologna,

Italy) where they were stored at -80°C until processing. Each subject was asked to fill in a 7-day

weighted food intake record (7D-WR), with the total food and beverage consumption, as previously

described.[23] Daily total calorie intake as well as that of macro- and micro-nutrients were assessed

through the MètaDieta® software version 3.7 (METEDA). Written informed consent was obtained

from all volunteers. All work was approved by the Ethics Committee of the Sant'Orsola-Malpighi

Hospital, University of Bologna (ref. number, 118/2015/U/Tess).

Microbial DNA extraction

Total bacterial DNA was extracted from each stool sample using the DNeasy Blood and Tissue kit

(QIAGEN) with the modifications previously described by Biagi et al.[24] In brief, 250 mg of fecal

samples were suspended in 1 ml of lysis buffer (500 mM NaCl, 50 mM Tris-HCl pH 8, 50 mM

EDTA, 4% (w/v) SDS), added with four 3-mm glass beads and 0.5 g of 0.1-mm zirconia beads

(BioSpec Products) and homogenized using a FastPrep instrument (MP Biomedicals) with three

bead-beating steps at 5.5 movements/sec for 1 min, and 5-min incubation in ice between treatments.

After incubation at 95°C for 15 min, stool particles were pelleted by centrifugation at 14,000 rpm

for 5 min. Nucleic acids were precipitated by adding 260 µl of 10 M ammonium acetate and one

volume of isopropanol. The pellets were then washed with 70% ethanol and suspended in TE

buffer. RNA was removed by treatment with 2 µl of DNase-free RNase (10 mg/ml) at 37°C for 15

min. Protein removal and column-based DNA purification were performed following the

manufacturer's instructions (QIAGEN). DNA was quantified with the NanoDrop ND-1000

spectrophotometer (NanoDrop Technologies).

16S rRNA gene sequencing

For each sample, the V3-V4 region of the 16S rRNA gene was amplified using the S-D-Bact-0341-

b-S-17/S-D-Bact-0785-a-A-21 primers[25] with Illumina overhang adapter sequences. PCR reactions

were performed in a final volume of 25 µl, containing 12.5 ng of genomic DNA, 200 nM of each

primer, and 2X KAPA HiFi HotStart ReadyMix (Kapa Biosystems, Roche), in a Thermal Cycler T

(Biometra GmbH) with the following gradient: 3 min at 95°C for the initial denaturation, 25 cycles

of denaturation at 95°C for 30 sec, annealing at 55°C for 30 sec and extension at 72°C for 30 sec,

and a final extension step at 72°C for 5 min. PCR products of around 460 bp were purified using a

magnetic bead-based system (Agencourt AMPure XP; Beckman Coulter) and sequenced on

Illumina MiSeq platform with the 2×250 bp paired-end protocol, according to the manufacturer's

guidelines (Illumina). Briefly, each indexed library was prepared by limited-cycle PCR using

Nextera technology, and further purified as described above. The libraries were subsequently

pooled at equimolar concentrations, denatured with 0.2 N NaOH, and diluted to 6 pM before

loading onto the MiSeq flow cell. Sequencing reads were deposited in MG-RAST (ID: …).

Bioinformatics and statistics

Raw sequences were processed using a pipeline that combines PANDAseq[26] and QIIME.[27] The UCLUST software[28] was used to bin high-quality reads into operational taxonomic units (OTUs) at 0.97 similarity threshold through an open-reference strategy. Taxonomy was assigned through the RDP classifier, using the Greengenes database as a reference (release May 2013). Chimera filtering was performed by using ChimeraSlayer.[29] All singleton OTUs were discarded.

16S rRNA gene sequencing data of our cohort were compared with publicly available data from the following previous studies: De Filippis *et al.*[21] (127 Italians; NCBI Sequence Read Archive (SRA) accession number: SRP042234), Schnorr *et al.*[5] (16 Italians and 27 Hadza hunter-gatherers from Tanzania; MG-RAST ID: 7058), Obregon-Tito *et al.*[6] (25 Matses hunter-gatherers from Peru; NCBI SRA: PRJNA268964), and Girard *et al.*[22] (21 Inuit from the Canadian Arctic; Qiita ID: 10439). Alpha diversity was assessed using the Shannon and Simpson indices. Beta diversity was evaluated using the Bray-Curtis dissimilarity measure. All statistical analysis was performed in R 3.3.2, using R Studio 1.0.44 and the libraries vegan, made4 and stats. The significance of data separation in the Principal Coordinates Analysis (PCoA) of Bray-Curtis distances was tested using a permutation test with pseudo-*F* ratios (function Adonis of vegan). Superimposition of bacterial genera on the PCoA plot was performed using the envfit function of vegan. Wilcoxon test was used to assess significant differences between groups (for intra- and inter-individual diversity), while Kruskal–Wallis test was used for multiple comparisons. P-values were corrected for false discovery rate (FDR, Benjamini-Hochberg) and P-values ≤ 0.05 were considered statistically significant.

ACKNOWLEDGMENTS

We would like to thank, and Jessica Baldini and Sara Quercia from Wellmicro S.r.l. (Bologna, Italy) for the skillful technical assistance in processing 7D-WRs and sequencing fecal samples, as well as the Paleo Meeting Organizing Committee (Italy).

REFERENCES

266   1.   Moeller AH, Li Y, Mpoudi Ngole E, Ahuka-Mundeke S, Lonsdorf EV, Pusey AE, *et al*. Rapid changes in the gut microbiome during human evolution. Proc Natl Acad Sci U S A 2014; 111:16431-5.

2.   Davenport ER, Sanders JG, Song SJ, Amato KR, Clark AG, Knight R. The human microbiome in evolution. BMC Biol 2017; 15:127.

3.   De Filippo C, Cavalieri D, Di Paola M, Ramazzotti M, Poullet JB, Massart S, *et al.* Impact of diet in shaping gut microbiota revealed by a comparative study in children from Europe and rural Africa. Proc Natl Acad Sci U S A 2010; 107:14691-6.

4.   Yatsunenko T, Rey FE, Manary MJ, Trehan I, Dominguez-Bello MG, Contreras M, *et al*. Human gut microbiome viewed across age and geography. Nature 2012; 486:222-7.

5.   Schnorr SL, Candela M, Rampelli S, Centanni M, Consolandi C, Basaglia G, *et al*. Gut microbiome of the Hadza hunter-gatherers. Nat Commun 2014; 5:3654.

6.   Obregon-Tito AJ, Tito RY, Metcalf J, Sankaranarayanan K, Clemente JC, Ursell LK, *et al.* Subsistence strategies in traditional societies distinguish gut microbiomes. Nat Commun 2015; 6:6505.

7.   Blaser MJ. The theory of disappearing microbiota and the epidemics of chronic diseases. Nat Rev Immunol 2017; 17:461-3.

8.   Sonnenburg ED, Sonnenburg JL. Starving our microbial self: the deleterious consequences of a diet deficient in microbiota-accessible carbohydrates. Cell Metab 2014; 20:779-86.

9.   Mosca A, Leclerc M, Hugot JP. Gut microbiota diversity and human diseases: should we reintroduce key predators in our ecosystem? Front Microbiol 2016; 7:455.

10.   Zuo T, Kamm MA, Colombel JF, Ng SC. Urbanization and the gut microbiota in health and

inflammatory bowel disease. Nat Rev Gastroenterol Hepatol 2018; 15:440-52.

11.   Cani PD, Jordan BF. Gut microbiota-mediated inflammation in obesity: a link with

gastrointestinal cancer. Nat Rev Gastroenterol Hepatol 2018. [Epub ahead of print]

12.   El Kaoutari A, Armougom F, Gordon JI, Raoult D, Henrissat B. The abundance and variety

of carbohydrate-active enzymes in the human gut microbiota. Nat Rev Microbiol 2013; 11:497-504.

13.   Danchin A. Bacteria in the ageing gut: did the taming of fire promote a long human

lifespan? Environ Microbiol 2018. [Epub ahead of print]

14.   He Y, Wu W, Zheng HM, Li P, McDonald D, Sheng HF, *et al*. Regional variation limits

applications of healthy gut microbiome reference ranges and disease models. Nat Med 2018;

24(10):1532-1535.

15.   Deschasaux M, Bouter KE, Prodan A, Levin E, Groen AK, Herrema H, *et al*. Depicting the

composition of gut microbiota in a population with varied ethnic origins but shared geography. Nat

Med 2018; 24(10):1526-1531.

16.   Ayeni FA, Biagi E, Rampelli S, Fiori J, Soverini M, Audu HJ, *et al*. Infant and adult gut

microbiome and metabolome in rural Bassa and urban settlers from Nigeria. Cell Rep 2018;

23:3056-67.

17.   Lindeberg S, Jonsson T, Granfeldt Y, Borgstrand E, Soffman J, Sjostrom K, *et al*. A

Palaeolithic diet improves glucose tolerance more than a Mediterranean-like diet in individuals with

ischaemic heart disease. Diabetologia 2007; 50:1795-807.

18.   Jonsson T, Granfeldt Y, Ahren B, Branell UC, Palsson G, Hansson A, *et al*. Beneficial

effects of a Paleolithic diet on cardiovascular risk factors in type 2 diabetes: a randomized cross-

over pilot study. Cardiovasc Diabetol 2009; 8:35.

19.   Whalen KA, McCullough ML, Flanders WD, Hartman TJ, Judd S, Bostick RM. Paleolithic

and Mediterranean diet pattern scores are inversely associated with biomarkers of inflammation and

oxidative balance in adults. J Nutr 2016; 146:1217-26.

20. Otten J, Stomby A, Waling M, Isaksson A, Soderstrom I, Ryberg M, *et al.* A heterogeneous

response of liver and skeletal muscle fat to the combination of a Paleolithic diet and exercise in

obese individuals with type 2 diabetes: a randomised controlled trial. Diabetologia 2018; 61:1548-

59.

21. De Filippis F, Pellegrini N, Vannini L, Jeffery IB, La Storia A, Laghi L, et al. High-level

adherence to a Mediterranean diet beneficially impacts the gut microbiota and associated

metabolome. Gut 2016; 65:1812-21.

22. Girard C, Tromas N, Amyot M, Shapiro BJ. Gut microbiome of the Canadian Arctic Inuit.

mSphere 2017; 2.

23. Dall'Asta C, Scarlato AP, Galaverna G, Brighenti F, Pellegrini N. Dietary exposure to

fumonisins and evaluation of nutrient intake in a group of adult celiac patients on a gluten-free diet.

Mol Nutr Food Res 2012; 56:632-40.

24. Biagi E, Franceschi C, Rampelli S, Severgnini M, Ostan R, Turroni S, *et al*. Gut microbiota

and extreme longevity. Curr Biol 2016; 26:1480-5.

25. Klindworth A, Pruesse E, Schweer T, Peplies J, Quast C, Horn M, *et al*. Evaluation of

general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based

diversity studies. Nucleic Acids Res 2013; 41:e1.

26. Masella AP, Bartram AK, Truszkowski JM, Brown DG, Neufeld JD. PANDAseq: paired-

end assembler for illumina sequences. BMC Bioinformatics 2012; 13:31.

27. Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, *et al*.

QIIME allows analysis of high-throughput community sequencing data. Nat Methods 2010; 7:335-

6.

28. Edgar RC. Search and clustering orders of magnitude faster than BLAST. Bioinformatics

2010; 26:2460-1.

29.   Haas BJ, Gevers D, Earl AM, Feldgarden M, Ward DV, Giannoukos G, *et al*. Chimeric 16S

rRNA sequence formation and detection in Sanger and 454-pyrosequenced PCR amplicons.

Genome Res 2011; 21:494-504.

FIGURE LEGENDS

Figure 1. Macronutrient composition of the modern Paleolithic diet. (A) Bar plots of the

percent caloric contribution of fat, protein and carbohydrate per subject, based upon weighted food

intake records over 7 days. The pie chart shows the summary of the average macronutrient intake

for the entire cohort. (B) Pie chart of the lipid type summary. PUFAs: polyunsaturated fatty acids;

MUFAs: monounsaturated fatty acids; SFAs: saturated fatty acids.

Figure 2. Phylogenetic structure of the gut microbiome of Italian adults adhering to the

modern Paleolithic diet. Bar plots of the genus-level composition of the gut microbiome of the

enrolled subjects. The pie chart shows the average relative abundance of bacterial genera. Only

bacterial genera with relative abundance > 0.5 % are shown. * = Unclassified.

Figure 3. The gut microbiome of Italian subjects following the modern Paleolithic diet shows

intermediate biodiversity between Western urban and traditional populations. Box and scatter

plots showing the alpha diversity values, measured with Simpson and Shannon indices, for each

study population (i.e. urban Italians adhering to the modern Paleolithic diet from the present study,

urban Italians adhering to the Mediterranean diet,[21] Hadza from Tanzania,[5] Matses from Peru,[6] and

Inuit from Canadian Arctic[22]). Different letters in the box plots indicate significant differences (P-

value < 0.05, Wilcoxon test). MPD = Modern Paleolithic Diet; MD = Mediterranean Diet.

Figure 4. Beta diversity of the fecal microbiome of Italian subjects following the modern

Paleolithic diet compared with other Western urban populations and traditional

communities. (A) The PCoA plot shows the Bray-Curtis distances between the genus-level

microbiota profiles of urban Italians adhering to the modern Paleolithic diet from the present study,

urban Italians adhering to the Mediterranean diet,[21] Hadza from Tanzania,[5] Matses from Peru,[6] and

Inuit from Canadian Arctic.[22] A significant segregation among study populations was found (P-

value $< 1 \times 10^{-5}$; permutation test with pseudo-F ratios). (B) Boxplots represent the interpersonal

variation, in terms of Bray-Curtis distances between genus-level microbiota profiles. Different

letters in the boxplots indicate significant differences (P-value < 0.05, Wilcoxon test). MPD =

Modern Paleolithic Diet; MD = Mediterranean Diet.

SUPPLEMENTAL ONLINE MATERIAL

Figure S1. Superimposition of the genus relative abundance on the PCoA plot. Arrows represent

the direction of significant correlations (permutation correlation test, P-value < 0.001). MPD =

Modern Paleolithic Diet; MD = Mediterranean Diet.

Table S1. Anthropometric data of the enrolled cohort.

Table S2. MPD macro- and micro-nutrients summary based on MétaDieta output.

Relative Abundance - Genus level

Legend:

- Other genera
- RF39*
- [Eubacterium]
- Catenibacterium
- Erysipelotrichaceae*
- Phascolarctobacterium
- Dialister
- Ruminococcus
- Oscillospira
- Faecalibacterium
- Ruminococcaceae*
- Ruminococcaceae;Other
- [Ruminococcus]
- Roseburia
- Lachnospira
- Dorea
- Coprococcus
- Blautia
- Lachnospiraceae*
- Lachnospiraceae;Other
- Clostridium
- Clostridiaceae*
- Clostridiales;Other;*
- Clostridiales;Other;Other
- Streptococcus
- S24-7
- Rikenellaceae
- Prevotella
- Parabacteroides
- Bacteroides
- Other*