Rare variants imputation in admixed populations: Comparison across reference panels and bioinformatics tools.

Sanjeev Sariya [a,b], Joseph Lee[a,b], Richard Mayeux [a,b,c.], Badri N. Vardarajan [a,b], Dolly Reyes-Dumeyer [a,b], Jennifer Manly, Adam Brickman, Rafael Lantigua[d], Martin Medrano[e], Ivonne Z. Jimenez-Velazquez[f] and Giuseppe Tosto[a,b,c]

   a. Taub Institute for Research on Alzheimer's Disease and the Aging Brain, College of Physicians and Surgeons, Columbia University. 630 West 168th Street, New York, NY 10032.

   b. The Gertrude H. Sergievsky Center, College of Physicians and Surgeons, Columbia University. 630 West 168th Street, New York, NY 10032.

   c. Department of Neurology, College of Physicians and Surgeons, Columbia University and the New York Presbyterian Hospital. 710 West 168th Street, New York, NY 10032

   d. Medicine College of Physicians and Surgeons, and The Department of Epidemiology, School of Public Health, Columbia University, New York, NY, USA

   e. School of Medicine, Pontificia Universidad Catolica Madre y Maestra, Santiago, Dominican Republic

   f. Department of Medicine, Geriatrics Program, School of Medicine, University of Puerto Rico, San Juan, Puerto Rico

Word counts: Abstract 344; Main Text 3,793; Tables: 4; Figures: 2; References: 35

Correspondence:
Giuseppe Tosto MD PhD
Taub Institute for Research on
Alzheimer's Disease and the Aging Brain
630 West 168th Street
New York, NY 10032
Tel: +1 212 305 9274
Email: gt2260@cumc.columbia.edu

Abstract.

Background: Imputation has become a standard approach in genome-wide association studies (GWAS) to infer *in silico* untyped markers. Although feasibility for common variants imputation is well established, we aimed to assess rare and ultra-rare variants' imputation in an admixed Caribbean Hispanic population (CH).

Methods: We evaluated imputation accuracy in CH (N=1,000), focusing on rare (0.1% ≤minor allele frequency (MAF) ≤ 1%) and ultra-rare (MAF < 0.1%) variants. We used two reference panels, the Haplotype Reference Consortium (HRC; N=27,165) and 1000 Genome Project (1000G phase 3; N=2,504) and multiple phasing (SHAPEIT, Eagle2) and imputation algorithms (IMPUTE2, MACH-Admix). To assess imputation quality, we reported: a) high-quality variant counts according to imputation tools' internal indexes (e.g. IMPUTE2 "info"≥80%). b) Wilcoxon Signed-Rank Test comparing imputation quality for genotyped variants that were masked and imputed; c) Cohen's kappa coefficient to test agreement between imputed and whole-exome sequencing (WES) variants; d) imputation of G206A mutation in the *PSEN1* (ultra-rare in the general population an more frequent in CH) followed by confirmation genotyping. We also tested ancestry proportion (European, African and Native American) against WES-imputation mismatches in a Poisson regression fashion.

Results: SHAPEIT2 retrieved higher percentage of imputed high-quality variants than Eagle2 (rare: 51.02% vs. 48.60%; ultra-rare 0.66% vs 0.65%, Wilcoxon p-value < 0.001). SHAPEIT-IMPUTE2 employing HRC outperformed 1000G (64.50% vs. 59.17%; 1.69% vs 0.75% for high-quality rare and ultra-rare variants, respectively; Wilcoxon p-value < 0.001). SHAPEIT-IMPUTE2 outperformed MaCH-Admix. Compared to 1000G, HRC-imputation retrieved a

higher number of high-quality rare and ultra-rare variants, despite showing lower agreement

between imputed and WES variants (e.g. rare: 98.86% for HRC vs. 99.02% for 1000G). High

Kappa ($K = 0.99$) was observed for both reference panels. Twelve G206A mutation carriers were

imputed and all validated by confirmation genotyping. African ancestry was associated with

higher imputation errors for uncommon and rare variants (p-value < 1e-05).

Conclusion: Reference panels with larger numbers of haplotypes can improve imputation quality

for rare and ultra-rare variants in admixed populations such as CH. Ethnic composition is an

important predictor of imputation accuracy, with higher African ancestry associated with poorer

imputation accuracy.

Keywords: Rare variants, Imputation accuracy, Admixed ancestry, GWAS, 1000Genomes, HRC

Introduction

Genome-wide association studies (GWASs) are a major tool to identify common variants

associated with complex diseases. GWAS can include 550K to over 2M Single Nucleotide

Polymorphisms (SNPs) (Ha et al., 2014) to cover the human genome evenly. Although GWAS

has shown to be a robust method to identify disease loci of interest, they rarely point to a causal

coding variant. In fact, microarray SNP chips for GWAS are optimally designed to uncover

common variants, often associated with small effect sizes mostly located in intronic and

intergenic regions. The focus of genetic investigations has since shifted toward rarer alleles with

larger effect sizes (Gibson, 2012). With the changing paradigm, imputation of rare variants has

become an important topic to enhance the genome coverage in GWAS. Imputation is a process

of inferring untyped SNP markers in the discovery population by using densely typed SNPs in

external reference panel(s). These 'in silico' markers increase the coverage of association tests

while conducting genome-wide association analysis. In addition, the large number of SNPs

facilitates meta-analysis when merging data from different study cohorts.

The quality of imputation essentially depends on two parameters: available reference datasets

and algorithms that employ those reference datasets. Previous studies have shown that

imputation quality depends on how well reference panels reflect the study population. To

respond to the needs, the 1000 Genome project (1000G) now in its third phase release, has

proven to be one of the most frequently used reference panels (Genomes Project et al., 2015).

Using these composite reference panels, a number of studies (Pei et al., 2010; Howie et al., 2012;

Verma et al., 2014; Liu et al., 2015) have compared imputation accuracy using different

imputation tools and algorithms, although the results are equivocal. Few studies (Browning and

Browning, 2009; Zheng et al., 2012; Zheng et al., 2015) assessed the impact of reference panel

size and input data's features - such as density of SNPs - to impute rare variants, suggesting

larger size of reference panels work better. Surakka and colleagues (Surakka et al., 2016)

assessed accuracy of imputed SNPs by evaluating rate of false polymorphisms in a Finnish

population using global reference panels – Haplotype Reference Consortium (HRC) release 1,

1000G phase 1 and a local reference panel. They concluded that higher false positive rate was

observed in imputation from global reference panels compared to imputation performed using a

local panel. Other studies (Huang et al., 2015; Das et al., 2016) found accuracy of imputation

increases with number of haplotypes, specifically for variants with $MAF \leq 0.5\%$, suggesting

larger sized reference panel for SNP imputation. In Hispanic populations, Nelson and colleagues

(Nelson et al., 2016) compared 1000G phase 1 (N=1,092) with 1000G phase 3 (N=2,504),

concluding that phase 3 with more than a 2-fold increase in sample size improved accuracy for

variants with $MAF <1\%$. On the other hand, Nagy and colleagues (Nagy et al., 2017) showed

that HRC reference panel provides new insight for novel variants particularly for rare variants in

a family-based Scottish study cohort. Aforementioned studies highlighted the need of a larger

sized reference panel to improve imputation quality. Herzig and colleagues (Herzig et al., 2018)

assessed tools for haplotype phasing and their impact on imputation in a population isolate of

Campora in southern Italy, and showed that SHAPEIT2, SHAPEIT3 and EAGLE2 were highly

accurate in phasing; MINIMAC3, IMPUTE4 and IMPUTE2 were found to be reliable for

imputation. Roshyara and colleagues (Roshyara et al., 2014) compared MaCH-Admix,

IMPUTE2, MACH, MACH-Minimac in different ethnicities by evaluating accuracy of correctly

imputed SNPs; MaCH-Minimac outperformed SHAPEIT-IMPUTE2 in subsamples of different

ethnic groups. These studies demonstrated how employed imputation algorithm determines

quality of inferred SNPs.

However, no study to our knowledge has evaluated reference panels in tandem with different

imputation algorithms to assess imputation quality of inferred SNPs based on MAF in a three-

way admixed population. Based on these findings, we assessed imputation quality, focusing on

rare and ultra-rare variants, in a large dataset of Caribbean Hispanics (CH) leveraging available

GWAS and sequencing data available for our cohort.

Methods

We will refer SNPs with MAF between 1-5% as "uncommon," 0.1-1% as "rare," and ≤0.1% as "ultra-rare." We considered SNPs with IMPUTE-Info metric ≥0.40 as "good-quality" and ≥0.80 as "high-quality."

GWAS samples and genotyping. We studied a total of 1,000 Caribbean Hispanics: 719 individuals from Estudio Familiar Investigar Genetica de Alzheimer (EFIGA), a study of familial LOAD; and 281 individuals from the multiethnic longitudinal cohort, Washington Heights, Inwood, Columbia Aging Project (WHICAP). The information on study design, recruitment and GWAS methods for the EFIGA and WHICAP study was previously described in Tosto, G., et al (Tosto et al., 2015).

GWAS quality control (QC). Genotyped data underwent quality control using PLINK (v1.90b4.9 64-bit) (Purcell et al., 2007)**.** Briefly, we excluded SNPs with missing rate ≥5% followed by exclusion of SNPs with MAF ≤ 1%. We then removed SNPs with P-value < 1e-6 for Hardy-Weinberg Equilibrium. Samples with missing call rate ≥5% were also excluded from analysis.

Global Ancestry estimation and selection of "true Hispanics". Prior to imputation, we estimated global ancestry using the ADMIXTURE (v.1.3.0) software (Alexander et al., 2009; Zhou et al., 2011). We conducted supervised admixture analyses using three reference populations: African Yoruba (YRI) and non-Hispanic white of European Ancestry (CEU) from the HAPMAP project as representative of African and European ancestral populations; and eight Surui, 21 Maya, 14 Karitiana, 14 Pima and seven Colombian individuals from the Human Genome Diversity Project (HGDP) were used to represent native American ancestry (Li et al., 2008). We used ~80,000

autosomal SNPs that were: I) genotyped in all three datasets (Caribbean Hispanics, 1000G and

HGDP); II) common (i.e. MAF >5 %); and III) in linkage equilibrium. Supervised admixture

analyses with the three reference populations (YRI, CEU, and Native Americans) revealed that

European lineage accounted for most the ancestral origins (59%), followed by African (33%) and

native American ancestry (8%). We then selected only individuals with at least 1% of all three

ancestral populations.

Reference panels. HRC reference panel contained over 39M SNPs from 27,165 individuals who

participated in 17 different studies (**Table 1).** The data were downloaded from the Wellcome

Trust Sanger Institute (WTSI). The panel contained 27,165 individuals from seventeen different

studies.

1000G phase 3 reference panel contained over 81M SNPs from 2,504 individuals

(https://mathgen.stats.ox.ac.uk/impute/1000GP_Phase3.tgz). It includes 26 ethnic groups, with

most variants rare, approximately 64 million had MAF <0.5%; approximately 12 million had a

MAF between 0.5% and 5%; and approximately 8 million have MAF >5%. In order to perform

imputation with MaCH-Admix, 1000G Phase 3 pre-formatted data were downloaded from

ftp://yunlianon:anon@rc-ns-

ftp.its.unc.edu/ALL.phase3_v5.shapeit2_mvncall_integrated.noSingleton.tgz that contained over

47M SNPs. The subsequent analyses were restricted to autosomal chromosomes, only.

Phasing and Imputation Procedures. We compared SHAPEIT2 (Delaneau et al., 2013) and Eagle2

(Loh et al., 2016) by phasing and then imputing (see next section) a single chromosome

(chromosome 21), using both reference panels. We refer to SHAPEIT2 as SHAPEIT when used

in tandem with IMPUTE2 for the remainder of paper.

Imputation was carried out using two bioinformatics tools: IMPUTE2 (Howie et al., 2009) and

MaCH-Admix. For both, imputation quality ranged from 0 to 1, with 0 indicating complete

uncertainty in imputed genotypes, and 1 indicating no uncertainty in imputed genotypes.

<u>IMPUTE2 (version 2.3.2)</u>. IMPUTE2 uses an MCMC algorithm to integrate over the space of

possible phase reconstructions for observed data genotypes. We conducted imputation in non-

overlapping 1MB chunk regions; chunk coordinates were specified using the "–*int*" option.

Other options were used with default parameters (**Supplementary section 1**). Briefly, we used a

default 250KB buffer region to avoid quality deterioration on the ends of chunk region. "-Ne"

value as 2000 suggested for robust imputation which scales linkage disequilibrium and

recombination error rate. <u>MaCH-Admix</u>. We used MaCH-Admix because it uses a method

based on IBS matching in a piecewise manner. The method breaks genomic region under

investigation into small pieces and finds reference haplotypes that best represent every small

piece, for each target individual separately. MaCH-Admix imputes in three steps: phasing,

estimation of model parameter that includes error rare and recombination rate and lastly,

haplotype-based imputation. MaCH-Admix (version Beta 2.0.185) was run on default parameters

of 30 rounds, 100 states (--autoFlip flag). Details can be found in supplementary file (**section 1**).

We initially compared performance between MaCH-Admix and IMPUTE2 using the 1000G

reference panel for Chromosome 21 only. We then proceeded to impute all remaining

chromosomes with the tool that performed better.

Imputation Performance Metrics. Different imputation tools were used to assess quality of

imputed SNPs with metrics based on the tool employed. IMPUTE2 uses "info" parameter to

report imputation quality that measures relative statistical information about SNP allele

frequency from imputed data. It reflects the information in imputed genotypes relative to the

information if only the allele frequency were known. Info metric is often used to filter poorly

imputed SNPs from IMPUTE2 and is reported for all imputed SNPs. In addition, IMPUTE2 uses

also an internal metric known as R^2 that measures squared correlation between input (genotyped)

and imputed (masked) SNP. R^2 is reported for input (genotyped) SNPs only, where genotyped

SNPs have been masked internally and imputed.

MaCH-Admix uses *Rsq* to report imputation quality. The R^2 metric is also known as variance

ratio, calculated as proportion of empirically observed variance (based on the imputation) to the

expected binomial variance p(1−p), where p is the minor allele frequency. Despite quality

measures from IMPUTE2 and MaCH-Admix being highly correlated (Marchini and Howie,

2010), we calculated a *r2hat* score to generate a single common metric to assess imputation

quality across the software (Hancock et al., 2012) (v109,

http://www.unc.edu/~yunmli/tgz/r2_hat.v109.tgz).

We compared performance of MaCH-Admix and SHAPEIT-IMPUTE2 by: a) Reporting raw

SNP counts based on quality (MaCH-Admix "Rsq" and IMPUTE2 "info"); b) Comparing *r2hat*

for overlapping imputed SNPs from both tools; c) Conducting a Wilcoxon Signed-Rank Test (R

v3.4.2) on *r2hat* value of overlapping SNPs.

We compared performance of Eagle2 and SHAPEIT2 phasing tools in tandem with IMPUTE2 as

imputation tools across reference panels by: a) Comparing IMPUTE2 R^2: b) Conducting a

Wilcoxon Signed-Rank Test on R^2 value; c) Reporting raw counts of imputed SNPs based on

their imputation quality (e.g. IMPUTE2 "info") and stratified by MAF bins (e.g. common, rare,

ultra-rare).

Agreement between Imputed and Sequence data. To further test the quality of imputation -

without relying on software's internal metrics (i.e. "info" and R^2) -   we calculated genotyped

concordance between imputed and WES data using the VCF-compare (v0.1.14-12-gcdb80b8)

tool (Danecek et al., 2011). First, we converted posterior probabilities obtained through

imputation into genotype data using PLINK software (v1.90b4.9) by applying a threshold of 0.9

(**supplementary section 1**), such that SNPs that failed on this criterion were left uncalled. For

example, an imputed SNP with P(G=0,1,2)= (0.01,0.9,0.09) would be called as a '1'

(heterozygous), whereas an imputed SNP with P(G=0,1,2) = (0.2, 0.6, 0.2) would be left

uncalled. We restricted the comparison to overlapping SNPs between HRC, 1000G reference

panels and whole-exome sequencing (WES) data for Chromosome 14 only, on SNPs with 0%

missingess (plink --missing flag) in WES data. We also assessed variants' agreement according

to different MAF bins for high-quality ("info" ≥0.8) SNPs. The output resulted in number of

variant "mismatches", i.e. the count of allele not matching between imputed and sequenced

variants per individual. To measure interrater reliability we computed Cohen's kappa coefficient

(McHugh, 2012) for both the panels against WES data. Kappa coefficient ≤ 0 indicates no

agreement, 0.01–0.20 as none to slight, 0.21–0.40 as fair, 0.41– 0.60 as moderate, 0.61–0.80 as

substantial, and 0.81–1.00 as almost perfect agreement.

Effects of Ancestry on Imputation Quality. To assess how ancestry affected imputation quality,

we conducted a Poisson regression using R. We used percentage of global ancestry (European

(CEU), Native (NAT) and African (YRI) as predictors, and total number of mismatches as main

outcome; analyses were restricted to high-quality SNPs only.

Imputation of G206A Mutation in PSEN1. To evaluate imputation performance for rare variants

in an admixed population, we examined a founder mutation, p.Gly206Ala (G206A - rs63750082)

in the *PSEN1* gene (PSEN1-G206A) (Athan et al., 2001; Lee et al., 2015). The PSEN1-G206A

mutation is rare variant observed primarily in Puerto Ricans with familial early onset

Alzheimer's disease (EOAD), but it is rare Puerto Ricans with LOAD. It was extremely rare in

populations with ancestries other than Puerto Rican (Arnold et al., 2013). It also has a highly

variable age at onset of EOAD associated with this mutation (range: 40-73 years). The mutation

was present in the 1000G phase 3 reference panel with an allele frequency of 0.001, but was

absent in the HRC reference panel. To verify whether individuals who were found to carry the

PSEN1-G206A mutation based on imputed genotypes, they were either whole genome

sequenced or genotyped using the KASP genotyping technology by LGC genomics

(https://www.lgcgroup.com), which uses allele-specific PCR for SNP calling. Agreement

between imputed and genotype data for the PSEN1-G206A mutation was then assessed.

Results

Comparison of Phasing Tools: Eagle2 vs. SHAPEIT2. To select the optimal tool for phasing, we

compared SHAPEIT2 with Eagle2 using Chromosome 21 with 13,066 genotyped SNPs by

performing subsequent imputation with IMPUTE2 on phased outputs, and using both reference

panels. We found SHAPEIT2 better than Eagle2 when evaluated for mean R^2 and the imputation

SNP "info" quality using both the reference panels. For instance, using the 1000G, we observed

higher mean R^2 for data phased with SHAPEIT2 as compared to Eagle2 (0.92 vs. 0.91; Wilcoxon

p-value < 0.001). When HRC panel was employed, mean R^2 of 0.89 was observed for

SHAPEIT2 against 0.85 for Eagle2 (Wilcoxon signed-rank test p-value < 0.001).

SNP count comparison details can be found in **Supplementary Table 1-2**. Regardless of the

reference panel employed, we observed higher percentage of "high-quality" rare and ultra-rare

SNPs for SHAPEIT-IMPUTE2 than Eagle2-IMPUTE2. For instance, 1000G-imputation

retrieved 51.02% of rare SNPs were of "high-quality" using SHAPEIT-IMPUTE2 against

48.38% with Eagle2-IMPUTE2. Detailed comparisons for different MAF bins and quality

threshold can be found in **Supplementary Section 2**. We found Eagle2 faster than SHAPEIT2

when computation times were compared; for instance, with HRC reference panel Eagle2 was ~6

times faster than SHAPEIT2 (**Supplementary Table 3**). We then imputed the remaining

chromosomes on phased output from SHAPEIT2. Comparison of phasing tools by assessing

switch error rate was beyond the scope of this paper due to limited resources, for e.g., availability

of phased reference panel for an admixed population.

MaCH-Admix vs. IMPUTE2. We found SHAPEIT-IMPUTE2 performed better than MaCH-

Admix. There were 1,104,648, and 653,791 SNPs (1000G) for SHAPEIT-IMPUTE2 and MaCH-

Admix respectively for chromosome 21. We imputed 1,104,648 (100%) and 646,594 (98.89%)

SNPs for SHAPEIT-IMPUTE2 and MaCH-Admix respectively; 549,091 SNPs were

overlapping. For SHAPEIT-IMPUTE2 we observed 446,591 (Info $\geq$ 0.40) bi-allelic SNPs, in

contrast with 598,943 (Rsq $\geq$ 0.30) from MaCH-Admix (**Supplementary Table 4**). SNP counts

for different MAF bins based on platform-specific quality index can be found in **Supplementary**

Table 5.When the two outputs were compared in terms of *r2hat*, SHAPEIT-IMPUTE2 showed

an average r2hat of 0.62 against 0.36 from MaCH-Admix (Wilcoxon signed-rank test p-value <

0.001). Also, MaCH-Admix was 109 times slower than IMPUTE2. (**Supplementary Table 6),**

thus, comparison between different panels using MaCH-Admix were excluded due to limited

resources. For the remaining of this manuscript, we focused on imputation employing SHAPEIT-

IMPUTE2, only.

Comparison between HRC and 1000G using SHAPEIT-IMPUTE2. Using SHAPEIT-IMPUTE2,

we imputed 81,240,392 and 38,532,090 SNPs across 22 chromosomes with 1000G and HRC

reference panels, respectively (**Table 2).**

Overall, we observed slightly higher mean R^2 with 1000G than with HRC panel (0.94 vs. 0.92;

Wilcoxon p-value< 0.001). Nevertheless, when the analyses were restricted to only "good-" and

"high-quality" SNPs, HRC consistently performed better: 60.82% of HRC-imputed SNPs were

"good-quality" and 48.87% were "high-quality" (Wilcoxon signed-rank test p-value < 0.001). On

the contrary, 40.32% of 1000G imputed SNPs were "good-quality" and 30.11% were "high-

quality". Further, we evaluated performance for uncommon, rare and ultra-rare SNPs. For "good-

" and "high-quality" SNPs, HRC outperformed 1000G. For example, HRC panel produced

62.85% of "high-quality" rare SNPs, whereas 1000G had 53.83% (**Table 3**). When average

imputation "Info" quality was evaluated, we observed SNPs with MAF ≤5% had higher average

info from HRC-imputation than with 1000G (Wilcoxon p-value< 0.001) (**Figure 1**).

Next, we restricted our analyses to *overlapping* SNPs across both reference panels only, based on

their chromosome and position mapping, reference and non-reference alleles. For "good-"and

"high-quality" SNPs, both panels performed similarly (**Table 2**). When restricted to uncommon,

rare and ultra-rare SNPs, we observed higher percentage of "good-" and "high-quality" SNPs

with HRC panel as compared to 1000G reference panel (**Table 3**). Specifically, we found 7.44%

of HRC-imputed ultra-rare SNPs were "good-quality" vs. 4.95% with the 1000G. For "high-

quality" SNPs, we found 1.69% of HRC-imputed were ultra-rare SNPs vs. 0.75% 1000G-

imputed ultra-rare SNPs. Further, Wilcoxon test was computed on info value of "high-quality"

ultra-rare SNPs (2,972) present in both overlapping 1000G and HRC panel (P-value < 0.001.).

Complete list of counts and percentages across reference panels, MAF bins and quality score can

be found in **Table 3**.

The case of G206A and the effect of chromosomal chunk size on imputation quality. Using

1000G reference panel, 12 individuals were likely G206A carriers. SNP rs63750082 was

imputed with an IMPUTE2 "info" score of 0.48 using 1MB as chromosomal region. When we

increased the chunk size to 5MB, IMPUTE-Info score drastically improved to 0.94 (**Figure 2**).

SNP rs63750082 is absent from HRC panel therefore no imputation was achieved. Those

patients labeled as mutation carriers according to imputation were then genotyped: all 12 were

confirmed to be G206A carriers, therefore achieving a perfect imputation prediction (100%

agreement) for that specific SNP.

Genotype Concordance and Kappa Coefficient. Out of the 1,000 individuals included in our

study, 262 had whole exome sequencing (WES) data available (Raghavan et al., 2018). We had

14,157 overlapping SNPs in WES, HRC and 1000G reference panels with 0% missingness in

WES data on chromosome 14; SNPs imputed with each reference panel were compared against

WES data separately. When concordance was evaluated, HRC panel performed slightly poorer,

despite showing higher number of high-quality variants than 1000G (**Table 4.a**). With 1000G,

we observed 3,542 rare and 35 ultra-rare high-quality SNPs; across 262 samples, we counted

1,245 ((1,245/(3,542*262))*100= 0.13%) and 10 (0.10%) mismatches for rare and ultra-rare

respectively. Using HRC, we retrieved 3,759 rare and 93 ultra-rare high-quality variants.

Mismatch count of 2,439 (0.24%) and 32 (0.13%) was observed for rare and ultra-rare variants

respectively in HRC panel. Details about pipeline can be found in **supplementary section 3**.

Next, we computed Cohen's kappa coefficient (K) for 14,157 imputed SNPs common in WES

and the two reference panels. For 1000G-imputation, we observed Kappa (K) of ~0.99 for both

rare and ultra-rare "high-quality" variants. For both rare and ultra-rare high-quality variants in

HRC panel, we observed a K of 0.99 (**Table 4.a).** Details about pipeline can be found in

supplementary section 4.

Effects of Ancestry on Imputation Quality. We evaluated the influence of individual ancestral

component separately on SNP mismatches for Chromosome 14 on 262 individuals. For both

reference panels we found that higher African ancestry (YRI) was associated with higher number

of mismatches (**Supplementary table 7**). For instance, with 1000G reference panel, for rare

variants (Info ≥0.80), we observed an estimate of 1.46 (P-value<0.001) for YRI component

(indicating that for each unit increase in YRI ancestry, it results in 1.46 additional mismatches).

Details on confidence intervals and robust standard errors can be found in **supplementary file**

(Table 7 and Section 5). We did not observe significant effect of ancestry on high-quality ultra-

rare variants in both panels.

Discussion

This study examined imputation performances in a cohort Caribbean Hispanics, focusing on

uncommon, rare and ultra-rare variant, by comparing different phasing and imputation tools, as

well as evaluating the effects of different reference panels. Overall, uncommon and rare variants

can be imputed in this population, characterized by a unique genetic background. Caribbean

Hispanics are admixed with 59% of their genetic component from European, 32% African, and

8% Native American ancestry (Tosto et al., 2015). Due to their genetic makeup and unique

linkage disequilibrium patterns, admixed populations offer unique opportunity in studying

complex diseases. First, disease prevalence varies across ethnic groups (Igartua et al., 2015) and

certain admixed populations show higher incidence rates and prevalence (e.g. Alzheimer,

diabetes etc.) or lower ones (e.g. multiple sclerosis). Second, variants that are ethnicity specific

may explain the higher prevalence of the disease of interest in the admixed groups.

In the present study, we examined multiple parameters of imputation using the Caribbean

Hispanics as an example with the idea that our findings can be applied to rare variant imputation

in other studies of admixed populations. First, we found that imputation using SHAPEIT-

IMPUTE2 phasing generated better results than Eagle2-IMPUTE2, and SHAPEIT-IMPUTE2 is

superior to MaCH-Admix in terms of imputation performances and process time.

Using SHAPEIT-IMPUTE2, 1000G SNPs outnumbered HRC panel because of the higher

number of SNPs included in the reference panel itself. However, when we restricted our analyses

to overlapping "good-" and "high-quality" SNPs (i.e. those variants that most likely would be

included in association analyses), HRC-imputation outperformed 1000G with higher. The

superior performance of HRC over 1000G was confirmed also when we focused on uncommon,

rare and ultra-rare SNPs only. Our findings confirm data in literature, i.e. reference panels with

higher number haplotypes perform better in different scenarios.

Overall, the higher quality of imputation for rare and ultra-rare variants was also confirmed when

we tested results against sequencing data. Finally, higher YRI global ancestry was found to

significantly impair SNP imputation, suggesting that imputation quality decreases with increased

African ancestry.

Lastly, SHAPEIT-IMPUTE2 with 1000G reference panel was successful identifying G206A

mutation carriers. We also noticed that imputation quality drastically improved when imputation

was conducted using large (5MB) chunk size as compared to small (1MB) chunks. This seems to

contradict previous observation: Zhang et al (Zhang et al., 2011) studied the effect of window

size on imputation carried out in an African-American population.   They concluded that window

size of 1MB could be considered acceptable. Possible explanations for these different results

might be the more complex admixture of CH compare to AA (three-way vs. two-way admixed)

and a more complex LD pattern for the G206A region. Ultimately, we recommend a window

size of 5MB chromosomal region to achieve high-quality imputation of SNPs with MAF ≤1% in

CH.

This work has limitations. First, we could carry out the comparison between the two reference

panels only for overlapping variants, limiting our observation to a subset of the variants included

in the 1000G. This is a result of the HRC composition, which includes several studies (e.g.the

1000G project) and comprises only a consensus number of variants. Second, we tested the

agreement between imputed and sequenced variants in a subset of individuals that had both

GWAS and WES available.

Acknowledgments

We thank the EFIGA study participants and the EFIGA research and support staff for their

contributions to this study. This study was supported by funding from the National Institute on

Aging [R21AG054832 (GT); 5R37AG015473 and RF1AG015473 (RM); R56 AG051876 and

R01 AG058918 (JL)] and the BrightFocus Foundation (A2015633S (JL)).

Conflict of Interest Statement

The authors declare no conflict of interests.

References.

Alexander, D.H., Novembre, J., and Lange, K. (2009). Fast model-based estimation of ancestry in unrelated individuals. *Genome Res* 19(9)**,** 1655-1664. doi: 10.1101/gr.094052.109.

Arnold, S.E., Vega, I.E., Karlawish, J.H., Wolk, D.A., Nunez, J., Negron, M., et al. (2013). Frequency and clinicopathological characteristics of presenilin 1 Gly206Ala mutation in Puerto Rican Hispanics with dementia. *J Alzheimers Dis* 33(4)**,** 1089-1095. doi: 10.3233/JAD-2012-121570.

Athan, E.S., Williamson, J., Ciappa, A., Santana, V., Romas, S.N., Lee, J.H., et al. (2001). A founder mutation in presenilin 1 causing early-onset Alzheimer disease in unrelated Caribbean Hispanic families. *JAMA* 286(18)**,** 2257-2263.

Browning, B.L., and Browning, S.R. (2009). A unified approach to genotype imputation and haplotype-phase inference for large data sets of trios and unrelated individuals. *Am J Hum Genet* 84(2)**,** 210-223. doi: 10.1016/j.ajhg.2009.01.005.

Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A., et al. (2011). The variant call format and VCFtools. *Bioinformatics* 27(15)**,** 2156-2158. doi: 10.1093/bioinformatics/btr330.

Das, S., Forer, L., Schonherr, S., Sidore, C., Locke, A.E., Kwong, A., et al. (2016). Next-generation genotype imputation service and methods. *Nat Genet* 48(10)**,** 1284-1287. doi: 10.1038/ng.3656.

Delaneau, O., Zagury, J.F., and Marchini, J. (2013). Improved whole-chromosome phasing for disease and population genetic studies. *Nat Methods* 10(1)**,** 5-6. doi: 10.1038/nmeth.2307.

Genomes Project, C., Auton, A., Brooks, L.D., Durbin, R.M., Garrison, E.P., Kang, H.M., et al. (2015). A global reference for human genetic variation. *Nature* 526(7571)**,** 68-74. doi: 10.1038/nature15393.

Gibson, G. (2012). Rare and common variants: twenty arguments. *Nat Rev Genet* 13(2)**,** 135-145. doi: 10.1038/nrg3118.

Ha, N.T., Freytag, S., and Bickeboeller, H. (2014). Coverage and efficiency in current SNP chips. *Eur J Hum Genet* 22(9)**,** 1124-1130. doi: 10.1038/ejhg.2013.304.

Hancock, D.B., Levy, J.L., Gaddis, N.C., Bierut, L.J., Saccone, N.L., Page, G.P., et al. (2012). Assessment of genotype imputation performance using 1000 Genomes in African American studies. *PLoS One* 7(11)**,** e50610. doi: 10.1371/journal.pone.0050610.

Herzig, A.F., Nutile, T., Babron, M.C., Ciullo, M., Bellenguez, C., and Leutenegger, A.L. (2018). Strategies for phasing and imputation in a population isolate. *Genet Epidemiol* 42(2)**,** 201-213. doi: 10.1002/gepi.22109.

Howie, B., Fuchsberger, C., Stephens, M., Marchini, J., and Abecasis, G.R. (2012). Fast and accurate genotype imputation in genome-wide association studies through pre-phasing. *Nat Genet* 44(8)**,** 955-959. doi: 10.1038/ng.2354.

Howie, B.N., Donnelly, P., and Marchini, J. (2009). A flexible and accurate genotype imputation method for the next generation of genome-wide association studies. *PLoS Genet* 5(6)**,** e1000529. doi: 10.1371/journal.pgen.1000529.

Huang, J., Howie, B., McCarthy, S., Memari, Y., Walter, K., Min, J.L., et al. (2015). Improved imputation of low-frequency and rare variants using the UK10K haplotype reference panel. *Nat Commun* 6**,** 8111. doi: 10.1038/ncomms9111.

Igartua, C., Myers, R.A., Mathias, R.A., Pino-Yanes, M., Eng, C., Graves, P.E., et al. (2015).
Ethnic-specific associations of rare and low-frequency DNA sequence variants with
asthma. *Nat Commun* 6**,** 5965. doi: 10.1038/ncomms6965.
Lee, J.H., Cheng, R., Vardarajan, B., Lantigua, R., Reyes-Dumeyer, D., Ortmann, W., et al.
(2015). Genetic Modifiers of Age at Onset in Carriers of the G206A Mutation in PSEN1
With Familial Alzheimer Disease Among Caribbean Hispanics. *JAMA Neurol* 72(9)**,**
1043-1051. doi: 10.1001/jamaneurol.2015.1424.
Li, J.Z., Absher, D.M., Tang, H., Southwick, A.M., Casto, A.M., Ramachandran, S., et al.
(2008). Worldwide human relationships inferred from genome-wide patterns of variation.
Science 319(5866)**,** 1100-1104. doi: 10.1126/science.1153717.
Liu, Q., Cirulli, E.T., Han, Y., Yao, S., Liu, S., and Zhu, Q. (2015). Systematic assessment of
imputation performance using the 1000 Genomes reference panels. *Brief Bioinform*
16(4)**,** 549-562. doi: 10.1093/bib/bbu035.
Loh, P.R., Danecek, P., Palamara, P.F., Fuchsberger, C., Y, A.R., H, K.F., et al. (2016).
Reference-based phasing using the Haplotype Reference Consortium panel. *Nat Genet*
48(11)**,** 1443-1448. doi: 10.1038/ng.3679.
Marchini, J., and Howie, B. (2010). Genotype imputation for genome-wide association studies.
Nat Rev Genet 11(7)**,** 499-511. doi: 10.1038/nrg2796.
McHugh, M.L. (2012). Interrater reliability: the kappa statistic. *Biochem Med (Zagreb)* 22(3)**,**
276-282.
Nagy, R., Boutin, T.S., Marten, J., Huffman, J.E., Kerr, S.M., Campbell, A., et al. (2017).
Exploration of haplotype research consortium imputation for genome-wide association
studies in 20,032 Generation Scotland participants. *Genome Med* 9(1)**,** 23. doi:
10.1186/s13073-017-0414-4.
Nelson, S.C., Stilp, A.M., Papanicolaou, G.J., Taylor, K.D., Rotter, J.I., Thornton, T.A., et al.
(2016). Improved imputation accuracy in Hispanic/Latino populations with larger and
more diverse reference panels: applications in the Hispanic Community Health
Study/Study of Latinos (HCHS/SOL). *Hum Mol Genet* 25(15)**,** 3245-3254. doi:
10.1093/hmg/ddw174.
Pei, Y.F., Zhang, L., Li, J., and Deng, H.W. (2010). Analyses and comparison of imputation-
based association methods. *PLoS One* 5(5)**,** e10827. doi: 10.1371/journal.pone.0010827.
Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M.A., Bender, D., et al. (2007).
PLINK: a tool set for whole-genome association and population-based linkage analyses.
Am J Hum Genet 81(3)**,** 559-575. doi: 10.1086/519795.
Raghavan, N.S., Brickman, A.M., Andrews, H., Manly, J.J., Schupf, N., Lantigua, R., et al.
(2018). Whole-exome sequencing in 20,197 persons for rare variants in Alzheimer's
disease. *Ann Clin Transl Neurol* 5(7)**,** 832-842. doi: 10.1002/acn3.582.
Roshyara, N.R., Kirsten, H., Horn, K., Ahnert, P., and Scholz, M. (2014). Impact of pre-
imputation SNP-filtering on genotype imputation results. *BMC Genet* 15**,** 88. doi:
10.1186/s12863-014-0088-5.
Surakka, I., Sarin, A.-P., Ruotsalainen, S.E., Durbin, R., Salomaa, V., Daly, M., et al. (2016).
The rate of false polymorphisms introduced when imputing genotypes from global
imputation panels. *bioRxiv*. doi: 10.1101/080770.
Tosto, G., Fu, H., Vardarajan, B.N., Lee, J.H., Cheng, R., Reyes-Dumeyer, D., et al. (2015). F-
box/LRR-repeat protein 7 is genetically associated with Alzheimer's disease. *Ann Clin*
Transl Neurol 2(8)**,** 810-820. doi: 10.1002/acn3.223.

Verma, S.S., de Andrade, M., Tromp, G., Kuivaniemi, H., Pugh, E., Namjou-Khales, B., et al.
(2014). Imputation and quality control steps for combining multiple genome-wide
datasets. *Front Genet* 5**,** 370. doi: 10.3389/fgene.2014.00370.
Zhang, B., Zhi, D., Zhang, K., Gao, G., Limdi, N.N., and Liu, N. (2011). Practical Consideration
of Genotype Imputation: Sample Size, Window Size, Reference Choice, and Untyped
Rate. *Stat Interface* 4(3)**,** 339-352.
Zheng, H.F., Ladouceur, M., Greenwood, C.M., and Richards, J.B. (2012). Effect of genome-
wide genotyping and reference panels on rare variants imputation. *J Genet Genomics*
39(10)**,** 545-550. doi: 10.1016/j.jgg.2012.07.002.
Zheng, H.F., Rong, J.J., Liu, M., Han, F., Zhang, X.W., Richards, J.B., et al. (2015).
Performance of genotype imputation for low frequency and rare variants from the 1000
genomes. *PLoS One* 10(1)**,** e0116487. doi: 10.1371/journal.pone.0116487.
Zhou, H., Alexander, D., and Lange, K. (2011). A quasi-Newton acceleration for high-
dimensional optimization algorithms. *Stat Comput* 21(2)**,** 261-273. doi: 10.1007/s11222-
009-9166-3.

Table 1: SNP counts in HRC and 1000G reference panel.

Reference Panel	Individuals	Autosomal variants	Bi-allelic SNPs	Multi-allelic SNPs
1000G Phase 3	2,504	81,706,022	7,781,8332	388,7690
HRC	27,165*	39,131,600	39,131,600	NA

*For Chromosome 1, the number of individuals were 22,691

527 **Table 2**: Type of imputed SNPs across reference panels.

Reference Panel	Multi-allelic SNPs			Bi-allelic SNPs			Total SNPs		
	Total SNPs	Info* ≥0.40 (%)	Info ≥0.80 (%)	Total SNPs	Info ≥0.40 (%)	Info ≥0.80 (%)	Total SNPs	Info ≥0.40 (%)	Info ≥0.80 (%)
All SNPs									
1000G	3,319,815	2,586,342 (77.90)	2,061,295 (62.09)	77,920,577	31,423,926 (40.32)	23,468,086 (30.11)	81,240,392	31,423,926 (41.86)	25,529,381 (31.42)
HRC	NA	NA	NA	38,532,090	23,436,980 (60.82)	18,833,790 (48.87)	38,532,090	23,436,980 (60.82)	18,833,790 (48.79)
SNPs overlapping HRC & 1000G									
1000G	NA	NA	NA	30,090,251	22,631,112 (75.21)	18,408,585 (61.17)	30,090,251	22,631,112 (75.21)	18,408,585 (61.17)
HRC	NA	NA	NA	30,090,251	22,438,268 (74.56)	18,395,036 (61.13)	30,090,251	22,438,268 (74.56)	18,395,036 (61.13)

532   **Table 3**. SNP Counts for all Bi-allelic uncommon, rare and ultra-rare SNPs.

MAF	1000G			HRC		
	Info ≥0	Info ≥0.40 (%)	Info ≥0.80 (%)	Info ≥0	Info ≥0.40 (%)	Info ≥0.80 (%)
All SNPs						
[1% - 5%]	6,025,281	5,989,223 (98.90)	5,441,982 (90.31)	5,434,996	5,421,257 (99.84)	5,061,904 (93.13)
[0.1% - 1%)	20,249,058	16,881,286 (83.36)	10,901,789 (53.83)	11,780,671	10,931,924 (92.79)	7,404,808 (62.85)
[0 – 0.1%)	44,562,205	1,490,434 (3.34)	242,717 (0.544)	15,055,433	828,256 (5.50)	174,673 (1.16)
SNPs overlapping HRC & 1000G						
[1% - 5%]	5,624,956	5,604,308 (99.63)	5,148,285 (91.52)	5,396,207	5,385,364 (99.79)	5,037,187 (93.34)
[0.1% - 1%)	11,875,603	10,442,603 (87.93)	7,027,312 (59.17)	10,945,899	10,268,136 (93.80)	7,060,908 (64.50)
[0 – 0.1%)	6,314,479	312,967 (4.95)	47,614 (0.75)	7,519,807	560,043 (7.44)	127,423 (1.69)

Table 4.a Comparison for mismatch counts and Kappa (*K*) for HRC and 1000G using WES data
on Chromosome 14.

MAF	1000G				HRC			
	Info ≥0.80				Info ≥0.80			
	SNP	Total SNPs in all persons[*]	Mismatch	Kappa (K)	SNP	Total SNPs in all persons[*]	Mismatch	Kappa (K)
[1% - 5%]	2,354	610,550	7,397 (1.22%)	0.99	2,264	587,961	8,963 (1.52%)	0.99
[0.1% - 1%)	3,542	926,109	1,245 (0.13%)	0.99	3,759	982,734	2,439 (0.24%)	0.99
[0 – 0.1%)	35	9,163	10 (0.10%)	0.99	93	24,348	32 (0.13%)	0.99

*Less value than 262*SNP because imputed with poor posterior probability failed to be
converted from .gen to plink format.

538   **Figure 1**: Comparison of average info quality between HRC and 1000G reference panel for all

539   chromosomes

541   **Figure 2:** Comparison of average info on CHR14: 70-75MB (5MB) vs 73-74MB (1MB) region

Comparison of average info between HRC and 1000G