1   # Low acetylcholine during early sleep is important for motor memory consolidation

2   Samsoon Inayat[1,3], Qandeel[1,3], Mojtaba Nazariahangarkolaee[1], Surjeet Singh[1], Bruce L. McNaughton[1,2],

3   Ian Q. Whishaw[1], and Majid H. Mohajerani[1,4]

4   1. Canadian Centre for Behavioural Neuroscience, University of Lethbridge, Lethbridge, Alberta,

5    Canada

6   2. Center for the Neurobiology of Learning and Memory, University of California, Irvine,

7    California, USA

8   3. These authors contributed equally

9   4. Corresponding author: mohajerani@uleth.ca

11   **Abstract:** Sleep rejuvenates the body and plays a role in consolidating newly encoded memory for long-

12   term storage. Evidence suggests that early sleep, featuring reduced cholinergic tone and slow-wave

13   activity, is important for consolidating episodic memory. The relationship between procedural memory

14   consolidation, early sleep, and acetylcholine level has not been investigated. We hypothesized that

15   increasing acetylcholine levels during early sleep would disrupt the consolidation of procedural memory

16   as measured by rotarod and skilled reaching tasks. Separate groups of adult mice were trained on the tasks

17   and after training, were given physostigmine, an acetylcholinesterase inhibitor, to increase cholinergic

18   tone in subsequent sleep. Post sleep testing showed that physostigmine impaired motor performance.

19   Home-cage video monitoring and electrophysiology revealed that physostigmine disrupted sleep structure

20   in the first hour after injection, delayed slow-wave sleep onset, and reduced slow-wave power in the

21   hippocampus and cortex. The impaired motor performance with physostigmine, however, was not due to

22   its effects on sleep structure, as one hour of sleep deprivation after training did not impair performance.

23   A reduction in cholinergic tone using chemogenetic inactivation of cholinergic neurons during early sleep

24   also did not affect rotarod performance. Finally, administration of agonists and antagonists of muscarinic

25   and nicotinic acetylcholine receptors revealed that it was the activation of muscarinic receptors during

26   early sleep that impaired rotarod performance. The experiments support the idea that the increased slow

27   wave activity of early sleep and associated inactivation of muscarinic receptors due to reduced

28   acetylcholine contribute to procedural memory consolidation.

Significance statement: Altered acetylcholine levels or modified expression of acetylcholine receptors exist in mental/neurological disorders such as Alzheimer's, Parkinson's, and Huntington disease and schizophrenia. Our study contributes to the understanding of the role of low acetylcholine levels and increased slow-wave EEG sleep for procedural memory consolidation. We show that increasing acetylcholine levels during early sleep dominated by slow waves disrupts motor memory consolidation. An important implication of our study is the importance of sleep onset to the timing of the administration of therapeutic drugs (existing and in development) for diseases that alter acetylcholine levels and impair memory.

INTRODUCTION

Memory formation consists of two processes; encoding which involves stimulus-induced gene-expression within cells and consolidation which happens through formation and strengthening of synaptic connections (Ebbinghaus 1885-Republished Translation 1964, Jenkins and Dallenbach 1924, McGaugh 2000, Dudai 2004, Frankland and Bontempi 2005, Diekelmann and Born 2010, Clopath 2012, Rasch and Born 2013, Yang, Lai et al. 2014, Dudai, Karni et al. 2015, Chambers 2017). Memory consolidation can happen in both awake and sleep states, can last from seconds to years, and might involve different processes for episodic, memory for facts and experiences, and procedural, memory for perceptual conditioning and motor skill (Clopath 2012, Rasch and Born 2013, Chambers 2017). A growing interest is on the sleep effects on memory consolidation. Procedural and episodic memories might be consolidated during rapid eye movement (REM) or non-REM (NREM) phases/stages of sleep respectively (Yaroush, Sullivan et al. 1971, Plihal and Born 1997), or through cyclic succession of NREM and REM (Giuditta, Ambrosini et al. 1995, Ambrosini and Giuditta 2001). Reactivation of memories during sleep (Wilson and McNaughton 1994, Kudrimoti, Barnes et al. 1999, Sutherland and McNaughton 2000, Kali and Dayan 2004, Euston, Tatsuno et al. 2007, Buhry, Azizi et al. 2011, Tononi and Cirelli 2014, Yang, Lai et al. 2014), formation and strengthening of spines during NREM and REM phases of sleep, and synapse pruning during REM sleep might also play a role (Yang, Lai et al. 2014, Li, Ma et al. 2017). The synaptic homeostasis theory of memory consolidation suggests a mechanism through which synaptic strengthening could occur. It proposes that low levels of neuromodulators such as acetylcholine, dopamine, serotonin, during NREM sleep permit overall synaptic downscaling while allowing recently activated synapses to be relatively more active thus improving signal-to-noise and thereby providing a mechanism for synaptic

stabilization, memory replay, and memory consolidation (Tononi and Cirelli 2003, Tononi and Cirelli

2014).

Consistent with the synaptic homeostasis theory, low levels of acetylcholine (ACh) allow synaptic

pruning/formation and episodic memory consolidation during NREM sleep (Hasselmo 1999, Walker

2009, Alger, Chambers et al. 2015, Miyamoto, Hirai et al. 2017). Whether procedural memory

consolidation is similarly enhanced by low ACh levels is uncertain. Studies on the effects of lesions that

lower cortical ACh on motor learning have produced mixed results (Conner, Culberson et al. 2003;

Gharbawie and Whishaw 2003). Other evidence suggests that increasing cholinergic tone in humans

during post-learning NREM does not affect consolidation of procedural memories (Gais and Born 2004).

Nevertheless, disrupting cholinergic levels through sleep deprivation has also been shown to affect motor

memory. For example, reactivation of motor task-specific neurons during NREM sleep is involved in

forming new synapses after motor learning in mice but consolidation is impaired in sleep deprived animals

(Yang, Lai et al. 2014). Furthermore, there is a proposed relationship between the amount of slow-wave

activity (SWA) during NREM sleep and motor learning. For example, humans learning a reaching task

display local increases in SWA in the parietal cortex (Landsness, Ferrarelli et al. 2011) whereas arm

immobilization induces a local decrease in SWA (Huber, Ghilardi et al. 2006). SWA also significantly

improves visual texture discrimination skills (Gais, Plihal et al. 2000). In mice, performance of a skilled

forelimb reaching for food task increased local SWA in motor cortex (Hanlon, Faraguna et al. 2009). In

sum, this evidence points to a role for decreased cholinergic tone during NREM sleep in procedural

memory consolidation, but a more definitive conclusion requires a direct test of the idea.

In the present study, we investigated the role of cholinergic activity during early NREM sleep in the

consolidation of procedural memory of the motor skills of mice performing a rotarod task and a skilled

reaching task. We described the effects of acetylcholine levels on sleep structure and phases, we related

acetylcholine levels in post-training sleep to subsequent motor performance, and we investigated the role

of muscarinic and nicotinic acetylcholine receptors in motor memory consolidation. In sum, we propose

that low acetylcholine levels during SWS are necessary for the consolidation of procedural memory.

RESULTS

Physostigmine injection after a motor learning task impairs the consolidation of motor memories

Two separate groups of mice were trained using two different motor tasks (rotarod and skilled forelimb

reaching). The rotarod task was used because learning takes place quickly, in 10-20 trials within a single

training session, allowing assessment of the effects of manipulations during relatively immediate post-

training sleep (Yang, Lai et al. 2014). In contrast, skilled forelimb reach task was used because training

takes place slowly in multiple training sessions (each with 20 trials) spread over 2-6 days. Thus, it involves

multiple cycles of encoding and post-training sleep. For both tasks, we investigated the effect of post-

training and pre-sleep injections of physostigmine, which is an acetylcholinesterase inhibitor and prevents

the breakdown of ACh thus increasing cholinergic tone (Aquilonius and Hartvig 1986, Pubchem 2018).

Rotarod Task Results

In the rotarod task (Fig. 1A), mice learn to maintain balance on an elevated rotating drum to prevent

themselves from falling. The maximum speed they attain just before they fall is used as a measure of their

learning during training and performance at a later time. On day 1 (training day) mice were given 10 trials

of rotarod early in the morning (8:30am – 10:00am) when sleep pressure is high. They gradually learnt

the task as speeds increased with trials (Fig. 1B - left). The increase in speeds was much faster in trials 1,

2 and 3 and became gradually faster in successive trials for 58 mice in the baseline group ($F_{9,513} = 24.52$,

$p_{gg} = 5.97e-27$, ANOVA-RM, effect for trials, p_{gg} is Greenhouse-Geisser corrected p value – see methods).

Post hoc comparisons with Tukey-Kramer's correction revealed that speeds in trials 2 and 3 were

significantly higher than in trial 1 ($p = 0.010$ and $p = 1.103e-5$ respectively) indicating fast learning.

Furthermore, speeds in trials 7-10 were significantly higher than in trial 3 ($p<0.006$ for all pairwise

comparisons) suggesting continual learning in subsequent trials. At the end of the training sessions, saline

or physostigmine was immediately administered in two groups of mice, and they were returned to their

home cages allowing them to rest or sleep.

To assess whether mice retained motor memories acquired on the training day, we gave 10 trials of rotarod

training on day 5 (probe day) around the same time of light/dark cycle on which initial training had

occurred (Fig. 1A). We compared speeds in the first trial of the probe day with speeds in the last trial of

the training day. For mice in the saline group, the two sets of speeds were similar (Student's t-test one-

tailed), indicating retention of motor memory formed on the training day. For mice in the physostigmine

group, speeds in the first trial of the probe day were significantly lower than speeds in the last trial of the

training day in the baseline group ($t_{70} = 2.50$, $p =0.007$, Student's t-test one-tailed) suggesting impaired

retention of motor memory compared to the saline group (Fig. 1C). We also compared average speeds

(over animals) for all trials within the experimental groups using a one-way ANOVA, which revealed a

significant difference among groups ($F_{3,568} = 52.01$, $p =1.03e-29$). The average speed of saline group was

significantly greater than that of the baseline group, a composite of all mice given preliminary training on

Day 1, (Fig. 1D, p=3.768e-9, post hoc Tukey's correction) as well as physostigmine group (p=3.768e-9,

post hoc Tukey's correction). There was no significant difference between average speeds of

physostigmine and the baseline group. These results thus suggest that the learned motor memories for

mice in the saline group were consolidated better than for mice in the physostigmine group.

Figure 1. Increasing acetylcholine levels using physostigmine (physo) in early sleep impairs the
consolidation of motor memories induced by the rotarod task. **(A)** Experimental setup and behavioral
protocol. In the rotarod task, animals learn to run and keep balance on an elevated rotating drum to prevent
themselves from falling. Mice were trained on Day 1 (training day) and retested on Day 5 (probe day). **(B)**
Average speeds (over animals) attained on the rotarod versus trials for training and probe days. N indicates
number of animals in each group. After fast learning in the first half of trials on training day, mice
gradually reached a plateau in performance. *** indicates significant effect of trials on speed ($F_{9,513}$ =
24.52, p_{gg} = 5.97e-27, ANOVA-RM). On probe day, performance of mice in saline and physo-late groups
further improved significantly with trials ($F_{9,117}$ = 3.10, p_{gg} = 0.02 for saline, blue *, and $F_{9,45}$ = 5.17, p_{gg}
= 0.01 for physo-late, green *, ANOVA-RM). The performance of mice in the physo group did not
improve significantly with trials ($F_{9,117}$ = 1.92, p_{gg} = 0.13, ANOVA-RM), however, there was an initial

improvement in the first three trials ($F_{2,26} = 6.37$, $p_{gg} = 0.01$, ANOVA-RM with p = 0.03 for comparison
of trials 3 and 1 with post-hoc Tukey-Kramer correction, red *). **(C)** Average speeds in trial 1 of physo
group were significantly smaller than those in last trial of baseline group suggesting impairment in
consolidation ($t_{70} = 2.50$, p = 0.007, one-tailed Student's t-test). **(D)** Between groups comparison of
average speeds over animals using one-way ANOVA ($F_{3,568} = 52.01$, p = 1.03e-29). Error bars represent
SEM. p_{gg} is Greenhouse-Geisser corrected p-value. ANOVA-RM is one-way ANOVA with repeated
measures.
143

Although the half-life of physostigmine is less than 1 hr in rodents (Somani 1989), but physostigmine also

produces bradycardia, salivation, and tremors. There was a possibility that these non-mnemonic effects of

physostigmine affected performance on the probe day. In a separate cohort of mice (n=6), physostigmine

was injected 24 hours after the last trial ended on the training day (physo-late group). Rotarod speeds of

these mice in the first trial on the probe day were similar to baseline speeds in the last trial of the training

day (Fig. 1B). In addition, average speed (over animals) in all trials was significantly greater compared to

baseline and physostigmine groups (Fig. 1D. p = 3.768e-9 for both comparisons, one-way ANOVA with

post hoc Tukey's correction). These results suggest that motor memory induced by the rotarod task is

consolidated after the first sleep session post training.

Skilled Forelimb Reach Task Results

The single pellet reaching task (Fig. 2A) requires animals to coordinated movements of the arm, wrist,

and hand in order to reach through a slot to obtain a food pellet. Mice were trained/tested for 10 days and

on each day one group was injected saline and a second group injected with physostigmine after training

(Fig. 2A). Mice injected with physostigmine were slower in learning the task, as suggested by a smaller

number of animals learning the task within the first 5 days compared to those injected with saline (Fig.

2B, an association between substance injected and success was observed on days 3 and 4, $\chi^2(1) = 10.8315$,

p=0.00099 and $\chi^2(1) = 4.492$ p=0.0341 respectively, Chi square independence test). A successful animal

here was defined as one which completed 20 trials on that particular day with at least one successful reach.

Furthermore, we compared the average percentages (over animals) of successful trials over days (Fig. 2C)

which increased with time, showing that mice in both groups did learn (for saline group, $F_{9,81} = 31.49$, p_{gg}

= 2.69e-8, for physo group, $F_{9,90} = 14.21$, $p_{gg} = 1.27e-7$, ANOVA-RM, effect of days). The average

percentages for the physostigmine group were significantly different than those for saline group ($F_{9,171} =$

2.81, $p_{gg} = 0.0169$, ANOVA-RM for group-days interactions) with significantly smaller values for

physostigmine on days 3,7, and 8 (p=0.041, p=0.009, and p=0.011 respectively, post hoc tukey-kramer

correction, group by days). These results show that the administration of physostigmine immediately after

reach training impaired performance.

Figure 2. Increasing acetylcholine levels using physostigmine in early sleep impairs the consolidation
of motor memories induced by the skilled forelimb reach task. (A) Experimental setup and behavioral
protocol. Animals learn to reach through a slot and grasp a food pellet placed on a platform. Mice were
trained/tested for 10 days. They were filmed on 9[th] day and reaches were scored from videos. **(B)**
Percentage of successful animal vs days. A "successful" animal completed 20 trials with at least one
successful reach and grasp. Firm line is fitted curve (sigmoidal). An association between substance
injected and success was observed for days 3 and 4 ($\chi^2(1) = 10.8315$, $p<0.001$ and $\chi^2(1) = 4.492$ $p<0.05$
respectively, Chi square independence test). **(C)** Percent single reach successes (averaged over animals)

vs days. Reaching accuracy improved over the training period for both groups (p_{gg} = 2.69e-8 and 1.27e-7
for saline and physostigmine groups respectively). However, reaching accuracy of physostigmine group
was significantly different from that of saline group ($F_{9,171}$ = 2.81, p_{gg} = 0.0169, ANOVA-RM, group-
days interaction) and smaller on days 3,7, and 8 (p=0.041, p=0.009, and p=0.011 respectively, post hoc
tukey-kramer correction, group by days). **(D)** Quality of reaches was assessed from video recordings done
on day 9 for 12 components of reach and grasp motion. A higher score indicates inferior performance.
Although group-component interaction was not significant for saline and physostigmine groups ($F_{11,209}$ =
1.83, p_{gg} = 0.129, ANOVA-RM), the grasp scores were significantly higher for physo group (p = 0.002,
post hoc tukey-kramer correction, black asterisk) as well as scores for fore-feet, lift, and elbow-in (t_{19} = -
1.89, p = 0.037, t_{19} = -2.08, p = 0.025, t_{19} = -1.84, p = 0.040 respectively, magenta asterisks, one-tailed
Student's t-test). **(E)** Average component scores (over animals) for the physostigmine group were
significantly larger than those for saline group (t_{754} = -4.711, p = 1.464e-6, one-tailed Student's t-test). All
Data are presented as Mean±SEM.
192

To assess the effect of physostigmine injections on the quality of reaches, we scored reaching movements

from the video recordings on day 9 by giving a score to each of 12 components of movements that make

up the reaching act (Whishaw, Whishaw et al. 2008, Whishaw, Faraji et al. 2017). The first three

successful reaches were scored and for a given component movement, a score of 0, 0.5, and 1 was assigned

for good, impaired, and abnormal performance respectively. A higher score would thus indicate inferior

performance. There was no significant difference between the scores for physostigmine and saline groups

determined by one-way ANOVA with repeated measures ($F_{11,209}$ = 1.83, p_{gg} = 0.129, group-components

interaction). However, post-hoc tukey-kramer correction revealed a higher mean score for the grasp

component for physostigmine (p = 0.002). Mean scores for the fore-feet, lift and elbow-in components

were also significantly larger for the physostigmine group compared to saline group revealed with one-

tailed student's t-test (t_{19} = -1.89, p = 0.037, t_{19} = -2.08, p = 0.025, t_{19} = -1.84, p = 0.040 respectively).

Finally, average component scores (over components and animals) were higher for the physostigmine

group (Fig. 1E, t_{754} = -4.711, p = 1.464e-6, one-tailed Student's t-test). These results show that as mice in

the physostigmine group acquired the task, their motor performance was impaired compared to the saline

group. Larger scores in the physostigmine group for grasp, forefeet, lift, and elbow-in components could

indicate impaired consolidation for associated memories compared to the saline group.

Physostigmine injection after rotarod training altered sleep structure

To investigate how physostigmine injection post-training impaired subsequent motor performance, we

asked whether physostigmine injection altered post-training activity or sleep. Separate cohorts of mice

were injected with saline or physostigmine after rotarod training, and returned to their home cages. Their

behavior was monitored using a video monitoring system developed in house (Singh, Bermudez-Contreras

et al. 2018). From the video frames, locomotion (Fig. 3A) was quantified by counting the number of pixels

that changed between consecutive video frames (Singh, Bermudez-Contreras et al. 2018). Motion profiles

were then transformed to motion/rest traces using the mean motion profile as threshold. Mice in the

physostigmine group had increased activity levels and showed delayed onset of rest/sleep. For each

mouse, the cumulative history of motion/rest periods was determined. The average cumulative duration

of motion for mice in the saline group was significantly right-shifted compared to mice in the

physostigmine group (Fig. 3B, p = 9.533e-143, one-tailed two-sample Kolmogorov-Smirnov test) and the

immobility periods of mice in the saline group were longer. Mice in the physostigmine group were more

active and had shorter episodes of rest. These results thus suggest that physostigmine injections altered

sleep/activity in the mice.

Figure 3. Quantification of motion after rotarod task in home cage using video monitoring. There is larger

motion in animals after physostigmine injection. (**A**) Representative raw traces of rms value of change in number

of pixels in video frames versus time. Blue and red traces are for mice injected with saline and physostigmine

(physo) respectively. Mean of the whole trace was used as a threshold to classify motion and rest periods shown in

black traces below each raw trace. (**B**) Average cumulative motion versus time. For each mouse, the cumulative

sum was determined from motion/rest traces. Shaded regions show SEM. The mean curve for saline group is

significantly right shifted compared to physo group (p = 9.533e-143, two-sample Kolmogrov-Smirnov test)

suggesting delayed motion after injections in mice in saline group.

Physostigmine delays slow wave sleep onset and reduces slow wave sleep power

To further explore how physostigmine might alter sleep structure, we used electrophysiology to monitor

EEG in one group of mice (N = 5). We implanted electrodes in the trapezius muscle (in the neck),

hippocampus

Figure 4. Increasing acetylcholine levels using physostigmine (physo) reduces slow-wave sleep and slow-wave power in the first hour of sleep post injection. (A) Representative hippocampal (HPC) and cortical (CTX – from M2, secondary motor area) local field potentials (LFP), electromyography (EMG) signal, hypnogram, and spectrogram. In the hypnogram, RS, SWS, and W represent states of animals i.e. REM sleep, slow-wave sleep, and wake periods respectively, which were scored from EMG traces and spectrogram of hippocampal and cortical LFPs.

PSD is power spectral density. Note that the hypnogram for physostigmine shows more fragmented sleep structure
compared to that for saline. **(B)** Cumulative duration vs time of recording for awake, SWS, and REM sleep states.
Difference in states between animals injected with physostigmine and saline lies mostly in the first hour post
injection. Mean cumulative curves for all states were significantly shifted between animals injected with saline and
physostigmine (p < 0.0001 for all comparisons, Kolmogorov-Smirnov test) **(C)** Comparison of sleep structure
between saline and physo groups in the first hour after injection. Waking activity in physo group was higher ($t_4 = -$
4.14, p = 0.014) while SWS was reduced ($t_4 = 4.32$, p = 0.012) – paired Student's t-test. **(D)** SWS latency increased
in physo group compared to saline group ($t_4 = -3.43$, p = 0.013, paired Student's t-test one-tailed). **(E)** Slow wave
power during SWS in the first hour of recording reduced in physo group compared to saline group in HPC and
cortical areas (M2 – Secondary motor area, LP-S1 – Lip primary sensory area, M1 – primary motor area, and RS –
retrosplenial cortex) revealed with one-tailed - paired Student's t-test. **(F)** Number of thalamo-cortical spindles
identified from M2 cortical LFP during the SWS period were smaller in physo group compared to saline. All Data
are presented as Mean±SEM.

(HPC), primary and secondary motor cortices (M1 and M2 respectively), cortical lip area (LP), and the
retrosplenial cortex (RS). Local field potentials (LFPs) were recorded after injecting saline and on a
subsequent day after injecting physostigmine (Fig. 4A). Mice were injected at 8:30am and signals were
analysed for the first four hours after injection. In both hippocampal and cortical spectrograms, we
observed visible reduction in power spectral density (PSD) in slow wave band within the first hour after
physostigmine injections (Fig. 4A). For further comparisons, we first determined sleep structure by
finding durations of wakefulness, REM sleep, and SWS using EMG signals and spectrogram of
hippocampal and cortical (from M2) LFPs (Fig. 4A and 4B). Compared to saline injections, sleep structure
after physostigmine injections was altered in the first hour of recording. After the first hour, cumulative
duration curves for saline and physostigmine injections were parallel for awake, SWS, and REM sleep
conditions (Fig. 4B) suggesting similar dynamics. After physostigmine injections mice had delayed onset
of SWS (Fig. 4A and 4B) and time spent awake was increased (Fig. 4C, $t_4 = -4.14$, p = 0.014, paired
Student's t-test one-tailed) while the SWS period was shorter (Fig. 4C, $t_4 = 4.32$, p = 0.012). REM sleep
was minimal during the first hour and not different for physostigmine and saline injections. SWS latency
was longer after physostigmine (31.17 ± 3.26 mins for physo vs 16.70 ± 2.76 mins for saline, Fig. 4D, $t_4 = -$
3.43, p = 0.013) and slow-wave power was reduced in the hippocampus (Fig. 4D, $t_4 = 4.00$, p = 0.008)
and cortex, (M2 - $t_4 = 3.79$, p = 0.0006, LP- $t_4 = 7.11$, p = 0.001, M1- $t_2 = 5.34$, p = 0.017, and RS- $t_2 =$
4.34, p = 0.025). The duration of thalamocortical spindles from M2 was significantly reduced after
physostigmine injections within the first hour (Fig. 4F) but the spindle occurrence frequency was similar.
These results thus suggest that physostigmine injection delay the onset of SWS sleep while reducing its
quality (lower slow wave power).

Figure 5. 1-hr sleep deprivation post rotarod training does not impair motor memories induced by the
rotarod task. (A) Control animals were put in home cages at 10am while sleep deprived animals were kept awake
and put in home cages at 11am **(B)** Average speeds (over animals) attained on the rotarod versus trials. N indicates
number of animals in each group. Average speeds of mice in baseline group improved significantly with trials ($F_{9,135}$
= 3.23, p_{gg} = 0.0094, ANOVA-RM, effect of trials). **(C)** Average speeds (over animals) of all trials for control and
sleep deprived groups were significantly larger than those of baseline group (p < 0.0005 for both control and sleep
deprived groups, one-way ANOVA with post-hoc Tukey's correction, $F_{2,317}$ = 17.55, p = 5.90e-8). **(D)** Average
cumulative motion versus time. Sleep deprived animals had smaller motion (p < 0.0005, two-sample Kolmogrov-
Smirnov test). Shaded regions show SEM.

One-hour sleep deprivation after rotarod training does not impair consolidation of motor memories
induced by the rotarod task

One might argue that the observed differences in rotarod performance were due to the sleep disruptive
effects of physostigmine rather than to increasing acetylcholine levels *per se* (Figs. 3 and 4). To investigate
this potentially confounding factor, we performed another set of experiments in a separate cohort of
animals in which one group of animals was sleep deprived for the first hour post rotarod training while
animals in the control group were put in home cages right after training to allow them to sleep (Fig. 5A).
As before, all animals in this set of experiments were combined to obtain "baseline" performance on
training day (Fig. 5B, left). The average speed (over animals) increased with trials ($F_{9,135}$ = 3.23, p_{gg} =
0.0094, ANOVA-RM, effect of trials) with a significant increase from trial 1 to trials 3, 4, and 10 (p =

0.045, 0.012, and 0.049 respectively, post-hoc Tukey-Kramer correction). The increase in speeds with

trials for control and sleep deprived groups were not significant on the probe day. The average speeds

over trials and animals for both control and sleep deprived groups were significantly greater than the

baseline (Fig. 5C, $p < 0.0005$ for both control and sleep deprived groups, one-way ANOVA with post-hoc

Tukey's correction, $F_{2,137} = 17.55$, $p = 5.90e-8$), however. We also quantified motion of animals post-

training with home-cage video recording (Fig. 5D) and observed that sleep deprived animals had lower

overall motion ($p < 0.0005$, two-sample Kolmogrov-Smirnov test) suggesting that they perhaps slept more

in shorter duration compared to controls. These results suggest that one-hour sleep deprivation post motor

training does not impair the consolidation of motor memories.

Chemogenetic inactivation of cholinergic neurons during early sleep does not impair the

consolidation of motor memories

Alternatively to increasing acetylcholine levels using physostigmine, here, we investigated the effect of

further reducing acetylcholine levels during early sleep on motor memory consolidation using transgenic

Chat-Cre::CAG-hM4Di mice (Sternson & Roth , 2014; Madisen et al., 2010). These mice express

inhibitory (hM4Di) DREADDs (Zhu, Aryal et al. 2016) within cholinergic neurons, and following the

administration of the drug clozapine-*N*-oxide (CNO), presynaptic release of neurotransmitter

acetylcholine in cholinergic neurons decreases. Since CNO exerts its maximum effects an hour to an hour-

and-half after the administration, we injected CNO just before the rotarod training in mice belonging to

the Chat-Cre::CAG-hM4Di CNO group (Fig. 6A). In another group of Chat-Cre::CAG-hM4Di mice,

saline was injected to establish a control group. As for C57 mice, all Chat-Cre::CAG-hM4Di mice for the

training day were combined to form a baseline group. Mice learnt the task on the training day (Fig. 6B,

$F_{9,144} = 3.72$, $p_{gg} = 0.005$, ANOVA-RM, effect of trials) but their learning curve was not as steep as that

for the C57 mice (compare with Fig. 1B). Speeds in the first trial on the probe day were similar to speeds

in the last trial of training day for both CNO and saline groups (one-tailed Student's t-test), however.

Furthermore, average speeds (over animals) for all trials in both CNO and saline groups were significantly

higher than the average speeds for the baseline group on the training day (Fig. 6C, $p < 0.001$ for both CNO

and Saline, one-way ANOVA with post-hoc Tukey's correction, $F_{2,339} = 31.27$, $p = 3.482e-13$), suggesting

that mice improved further on the probe day. The average speeds in the CNO group, however, were smaller

than those in saline group (Fig. 6C, $p = 0.0006$). These results suggest that the action of CNO on transgenic

mice during early sleep does not appreciably affect rotarod performance and has a small effect on motor

memory consolidation compared to physostigmine injections.

Figure 6. Decreasing acetylcholine levels in early sleep does not impair motor memories induced by the
rotarod task. (A) Chat-Cre::CAG-hM4Di transgenic animals were injected with either saline or clozapine-*N*-oxide
(CNO) before starting rotarod training because CNO exerts its maximum effect an hour to an hour and a half after
administration. **(B)** Average speeds (over animals) attained on the rotarod versus trials. N indicates number of
animals in each group. For training day, saline and CNO groups were combined to form the baseline group. Speeds
improved with trials for mice in baseline group ($F_{9,144} = 3.72$, $p_{gg} = 0.005$, ANOVA-RM, effect of trials, black **)
and saline group ($F_{9,63} = 3.40$, $p_{gg} = 0.029$, blue *, ANOVA-RM). **(C)** Average speeds (over animals) of all trials
in both CNO and saline groups were significantly higher than those in baseline group ($p < 0.001$ for both CNO and
Saline, one-way ANOVA with post-hoc Tukey's correction, $F_{2,339} = 31.27$, $p = 3.482e-13$).

Activation of muscarinic ACh receptors during early sleep impairs motor memory consolidation

In earlier experiments, post-training administration of physostigmine which indirectly activates both

muscarinic and nicotinic ACh receptors impaired post-sleep motor performance. Here, we investigated

the individual involvement of ACh receptors in motor memory consolidation. Based on the anatomical

distribution of ACh receptors in the brain, we hypothesized that, similar to previous studies, the muscarinic

ACh receptors would play a larger role in motor memory consolidation as compared to nicotinic ACh
receptors (Power, Vazdarjanova et al. 2003, Brown 2010, Hut and Van der Zee 2011). To activate
muscarinic and nicotinic ACh receptors selectively, the respective agonists oxotremorine and nicotine
were injected immediately after rotarod training in two separate groups of mice (behavioral protocol 1,
Fig. 1A). For both groups, speeds in the first trials on the probe day (Fig. 7) were similar to speeds in the
last trials on the training day (one-tailed Student's t-test) demonstrating performance retention. The
average speeds (over animals) for all trials for the nicotine group were significantly faster compared to
the baseline group (Fig. 7B, p = 0.0004 one-way ANOVA with post-hoc Tukey's correction, $F_{4,611} = 25.97$,
p = 6.823e-20) but slower than the speeds in the saline group (Fig. 7B, p = 0.002). For the oxotremorine
group, average speeds were similar to those in the baseline group and were significantly smaller than those
in the saline group (Fig. 7B, p = 1.272e-8, one-way ANOVA with post hoc Tukey's correction). These
results suggest that oxotremorine has a larger effect on the consolidation of motor memories as compared
to nicotine.

Figure 7. Activation of muscarinic acetylcholine (ACh) receptors during early sleep impairs motor memory
consolidation. (A) Average speeds (over animals) attained on the rotarod versus trials. N indicates number of
animals in each group. Baseline and saline groups are same as in Fig. 1. Average speeds of mice in groups
administered with oxotremorine (oxo) and nicotine (nico) did not significantly improve with trials on probe day
(NS, ANOVA-RM) but those of mice in scopolamine-mecamylamine (sco+mec) group improved significantly ($F_{9,63}$
= 3.67, p_{gg} = 0.024, ANOVA-RM, effect of trials) **(B)** The average speeds (over animals) of all trials for nico and
sco+mec groups but not oxo group were significantly larger than those for the baseline group (p < 0.0005 for both,
one-way ANOVA with post-hoc Tukey's correction, $F_{4,611} = 25.97$, p = 6.823e-20).

As a supporting experiment, we used a combination of two drugs, scopolamine and mecamylamine for

combined blockade of muscarinic and nicotinic acetylcholine receptors. Scopolamine and mecamylamine

were given immediately after rotarod training (See protocol 1 in Fig. 1A). The speeds obtained by the

mice in the first trial on the probe day were not significantly different from the last trial on the training

day (one-tailed Student's t-test). There was an increase in speeds on the probe day (Fig. 7A, $F_{9,63} = 3.67$,

$p_{gg} = 0.024$, ANOVA-RM, effect of trials). Furthermore, the average speeds (over animals) of all trials

were significantly higher than those for baseline and oxotremorine groups (Fig. 7B, $p = 3.307e\text{-}6$ and $p =$

0.009 respectively, one-way ANOVA with post-hoc Tukey's correction, $F_{4,611} = 25.97$, $p = 6.823e\text{-}20$).

Combined blockade of ACh receptors therefore did not alter performance on the rotarod task.

Table 1 shows the significance results of combined one-way ANOVA (with post-hoc tukey's corrections

for p values) for all groups of C57 mice (N = 58) used in the rotarod task. The results although slightly

different from those reported above when one-way ANOVA was done on subsets of groups, the main

conclusions still hold. Significance results from combined ANOVA strongly support the conclusion that

reduced muscarinic activity is associated with improved retention.

Table 1 – Significance results from One-way ANOVA. Post hoc Tukey's Correction. *, **, and *** for p<0.05, p<0.01, and p<0.001 respectively. ns is not significant

	Baseline	Saline	Physo	Physo-late	Sco+Mec	Oxo	Nico
Baseline		***	ns	***	***	ns	**
Saline			***	ns	ns	***	*
Physo				***	***	ns	***
Physo-late					*	***	***
Sco+Mec						*	ns
Oxo							ns
Nico							

Abbreviations: Physo – Physostigmine, Sco+Mec – Scopolamine + Mecamylamine, Physo-late – Physostigmine administered 24 hrs later, Oxo – Oxotremorine, Nico – Nicotine

DISCUSSION

We investigated how altering ACh levels during early sleep affects procedural memory

consolidation of two motor tasks. Supporting previous studies of episodic memory (Hasselmo 1999, Gais

and Born 2004, Hasselmo and McGaughy 2004), we find that low levels of ACh during NREM sleep

enhances procedural memory consolidation. Physostigmine, which is expected to increase ACh levels,

given immediately after training, delayed the onset of NREM sleep, reduced slow-wave power in the

cortex and hippocampus), and impaired post-sleep motor performance. Physostigmine given 24 hrs after

training, after first sleep session was over, did not affect motor performance. To rule out that sleep

disruption by physostigmine did not cause impaired motor performance in the initial experiments,

supporting experiments confirmed that one-hour sleep deprivation after training does not affect motor

performance and CNO injection into transgenic mice, which might be expected to decrease acetylcholine

levels, did not affect motor performance. By using agonists and antagonists of muscarinic and nicotinic

ACh receptors, we found that only activation of muscarinic receptors during early sleep impairs motor

performance. Taken together, these experiments are consistent with the idea that inactivation of

muscarinic receptors due to natural lowering of ACh levels during early sleep contributes to the

consolidation of procedural memory.

It is well known that motor skills show the greatest improvement if intermittent training is given

(Karni, Meyer et al. 1998, Buitrago, Ringer et al. 2004, Peters, Chen et al. 2014). It is suggested that this

pattern in which daily improvement is preserved and then enhanced on successive days of training is

enabled by a post training period of consolidation. Our results from animals injected with saline after

training in both rotarod and reach tasks reflect a similar pattern. For the rotarod task, the improvement

obtained on the first day was preserved and augmented on the second day. Similarly, for the skilled reach

task there was a day to day improvement in performance which eventually plateaued after a number of

days of training. The expectation underlying the experiments was that post training procedural memory

consolidation is important for this daily improvement in motor performance.

We obtained support for the idea that lower ACh levels during early sleep has a positive effect in

enhancing motor performance suggesting a positive role for lower ACh levels in procedural memory

consolidation. First, we used physostigmine to increase ACh levels in post training sleep. We did not

measure ACh levels directly but inferred from sleep pattern changes that cholinergic tone was increased.

In the first hour of sleep, post injection, SWS sleep onset was delayed and there was a reduction in slow-

wave power, consistent with half-life of physostigmine in rodents which is less than an hour (Somani

1989). Previous studies have reported similar changes in sleep patterns following manipulation of

cholinergic tone (Sitaram, Wyatt et al. 1976, Vanderwolf, Raithby et al. 1993, Dringenberg and

Vanderwolf 1996, Dringenberg and Vanderwolf 1997, Gais and Born 2004, Anaclet, Pedersen et al. 2015).

We then asked whether impairment in motor performance resulting from physostigmine injections was

due either to an increase in acetylcholine levels or to sleep disruption. We ruled out the later possibility

by showing that one hour of sleep deprivation post training does not impair motor performance. This is

consistent with previous studies showing that 3-7 hours (greater than one hour) of sleep deprivation
impairs motor memory consolidation (Gais, Plihal et al. 2000, Yang, Lai et al. 2014, Nagai, de Vivo et al.
2017) – but see Nagai et al (2017) who showed that sleep deprivation only affected motor performance
on a complex wheel task but not on the rotarod task. We also showed that the first sleep session after
motor learning is crucial for memory consolidation in-line with studies of effects of motor learning on
SWS, supporting the importance of first hour of sleep post training (Huber, Ghilardi et al. 2006,
Aeschbach, Cutler et al. 2008, Hanlon, Faraguna et al. 2009, Landsness, Crupi et al. 2009, Landsness,
Ferrarelli et al. 2011). Taken together, these results suggest that impairment in motor performance that we
observed with physostigmine injections was due to an increase in ACh levels in the first hour of sleep post
training. We therefore conclude that reduced ACh levels during early sleep are crucial for motor memory
consolidation parallel to a previous study for declarative memory (Gais and Born 2004). Moreover, we
propose that as for declarative memory, ACh levels might function as a switch between brain modes of
encoding and consolidation respectively (Hasselmo 1999, Hasselmo and McGaughy 2004, Rasch, Born
et al. 2006). We hypothesize that in a future experiment, blocking the action of acetylcholine during
wakefulness will impede new motor learning but will enhance motor memory consolidation.

Next, we asked whether further reducing ACh levels during early sleep would enhance motor
memory consolidation. Since, reducing ACh levels pharmacologically in wild type mice is a challenging
manipulation, we used a chemogenetic approach with transgenic mice to inactivate cholinergic neurons
during early sleep. Again, we did not measure ACh levels directly but inferred from behavior that levels
were reduced or they remained unaffected. The rotarod speeds of mice in CNO group were significantly
higher than those in baseline group but they were also significantly lower than those in saline group
suggesting an effect of CNO injection. However, undesired effects of CNO might have adversely affected
motor performance in comparison with saline group (Gomez, Bonaventura et al. 2017, Mahler and Aston-
Jones 2018, Manvich, Webster et al. 2018). These results suggest that further reducing ACh levels does
not affect motor memory consolidation as does increasing ACh levels. There are mixed findings about
whether selective ACh reduction affects motor performance (Conner, Culberson et al. 2003, Gharbawie
and Whishaw 2003, Ramanathan, Tuszynski et al. 2009, Conner, Kulczycki et al. 2010). Assuming that
CNO had the intended action of reducing acetylcholine levels, we conclude that such acute manipulation
does not adversely affect motor memory consolidation. These results also support our earlier conclusion
regarding the requirement of low acetylcholine levels for memory consolidation.

Finally, we investigated which ACh receptors might play a crucial role in motor memory consolidation.
Post-sleep motor performance was reduced by oxotremorine, a muscarinic acetylcholine receptor agonist,
suggesting that activation of muscarinic receptors is undesirable for memory consolidation during early
sleep. There was also no significant difference between oxotremorine and nicotine groups. In natural early
sleep therefore, reduction in ACh on muscarinic receptors is related to motor memory consolidation. Other
work has also shown that muscarinic acetylcholine receptors are more involved in cognitive functioning
(Hut and Van der Zee 2011, Zwart, Reed et al. 2018) and their activation causes increased REM like EEG
whereas their blockade produces slow wave activity (Vanderwolf, Raithby et al. 1993, Dringenberg and
Vanderwolf 1996).

The findings of the present study are relevant to contemporary theories of how sleep contributes
to motor memory consolidation. Our findings are consistent with the synaptic homeostatic theory of sleep
that slow wave sleep (SWS) facilitates memory consolidation by initiating downscaling of synaptic
strengths to bring the synaptic load impacting each neuron to baseline levels, thus increasing signal to
noise ratio of recently activated synapses (Tononi and Cirelli 2003). It also argues that low levels of ACh
present during SWS might mediate the synaptic downscaling process (Tononi and Cirelli 2006).
Consistent with this theory, when we increased ACh levels during early sleep, consolidation was impaired.
Our findings are not inconsistent with the report that it is the cyclic succession of SWS and REM is
important for memory consolidation (Giuditta, Ambrosini et al. 1995, Ambrosini and Giuditta 2001, Rasch
and Born 2013, Gais and Schonauer 2017). This finding suggests that SWS causes selective weakening
of non-adaptive memories (noise) and strengthening of adaptive memories (useful for survival) while
REM sleep is involved in integration of adaptive memories into existing knowledge. In our study,
physostigmine likely altered sleep structure disrupting the cyclic succession of SWS and REM and thus
causing impaired consolidation in this way. The present results are not consistent with the dual process
theory that advocates that SWS supports the consolidation of episodic memories while REM sleep
facilitates procedural memory consolidation (Stickgold, Whidbee et al. 2000, Louie and Wilson 2001,
Mednick, Nakayama et al. 2003, Peigneux, Laureys et al. 2003, Gais and Born 2004, Mascetti, Muto et
al. 2013). We find that low ACh levels during SWS facilitate motor memory consolidation. Thus, our
findings are consistent with numerous other studies (see above) that are inconsistent with this theory.
Perhaps the REM sleep dependent consolidation of procedural memories is dependent on the type of task
(Chambers 2017).

In conclusion, our results suggest that downregulated ACh levels during SWS are associated with
motor memory consolidation. We used pharmacological manipulation for altering ACh levels, but future
studies might use more refined methods. For example, by activating or silencing cholinergic neurons
optogenetically with a feedback system to detect sleep states, one could clamp ACh levels in the desired
brain region and study their role in memory consolidation using behavioral measures, electrophysiology
or optical imaging (Mohajerani, Chan et al. 2013). Furthermore, measurements to observe synaptic and
biochemical changes during sleep would allow the study of structural correlates of memory consolidation.

METHODS

All experiments were performed in accordance with Canadian Council of Animal Care and were approved
by the University of Lethbridge Animal Welfare Committee.

Experimental Animals

Adult wild-type (WT) mice (C57 Bl/6J, Jackson laboratories), N=110, and transgenic mice (Chat-
Cre::CAG-hM4Di), N=17, 3-6 months old, both male and female, and weighing 20-30g were used in this
study. Transgenic mice had inhibitory (hM4Di) DREADDs in cholinergic neurons, which were activated
by the inert molecule clozapine-*N*-oxide (CNO) to reduce release of acetylcholine. For generating these
mice, CAG-hM4Di mice (B6N.129-Gt(ROSA)26Sortm1(CAG-CHRM4*,-mCitrine) Ute/J) were crossed
with Chat-Cre mice (B6;129S6-Chattm2(cre)Lowl/J) to enable expression of CAG hM4Di in cholinergic
neurons of the brain (Stachniak, Ghosh et al. 2014, Vardy, Robinson et al. 2015). All mice were housed
up to four per cage and were provided food and water ad libitum. They were kept in a controlled
temperature (22°C), humidity, and light with a 12:12 light/dark cycle. All testing and training was
performed during the light phase of the cycle at the same time each day.

Motor Learning Tasks

Rotarod Task

In the rotarod task, mice learn to balance while running on an elevated rotating drum whose speed is
gradually increasing (Fig. 1A). As they learn to keep their balance they can stay on the rotating drum for
longer times at higher speeds of the drum. Eighty four C57/BL6 and 17 Chat-Cre::CAG-hM4Dimice were
used and were randomly divided into control and experimental groups. We used the modified version of

the rotarod as described by shiotuski et al, (Shiotsuki, Yoshimi et al. 2010) to emphasize the learning

aspect of the test. A four lane rotarod with automatic timers and fall sensors (Med Associates Inc.) with a

test chamber (74 cm / 84 cm / 50 cm) was used. The Diameter of drum was 7cm and was covered with

Duct tape to prevent mouse from gripping on to the surface. Animals were placed on the drum 3 mins

before the start of each session to habituate them. An accelerating rotarod paradigm was used in which

the rotation speed gradually increased from 4-40 rpm over the course of 5 min. The time latency and

rotation speed were automatically recorded by the photo sensors when the animal was unable to keep up

with the increasing speed and fell. The rotarod training consisted of one session (1-1.5h) with 10 trials and

an inter trial interval of 3-5 mins. Performance was measured as average speed animals achieved during

the training session. To evaluate long-term memory, 10 trials were repeated 5 days after the initial session

and speeds were compared. All sessions were performed at the same time 8:00-10:00 am in the morning

and animals were returned to home cages to allow them to sleep.

Skilled-Forelimb Reach Task

The skilled forelimb reaching task was established by Whishaw et al (Whishaw 2000) and is used to study

fine/precise co-ordinated movements of forelimb, arm, hand and digits as they work together to retrieve

food. For the skilled forelimb reach task 21 C57/BL6 mice were used and were divided into control and

drug groups by a third person making sure the experimenter was blind to the two groups. The animals

were placed on food restriction 3-4 days before the beginning of training. Prior to food restriction, mice

were weighed three days to obtain an average pre-restriction weight followed by a daily monitoring during

the food restriction period to maintain 85% of the average weight.

The animals were placed in a clear plexiglass testing chamber (20 cm long, 9 cm wide, and 20 cm high)

with a slit (1 cm wide) located in the center of the front wall. A 3cm wide shelf with two grooves/notches

at each side of the slot was mounted 1cm above the floor and outside of the front wall. The grooves were

curved out on the shelf to place 14mg food pellets. All mice were handled daily and were habituated to

the testing chamber for a week. Two days before training, the mice were given 10 pellets inside the

chamber and then 10 pellets very close to the slit on shelf before moving the pellet further on the groove.

The training/testing began on the day the mice were presented with food pellets in the grooves located on

the shelf. The training/testing consisted of 10 days with one session each day consisting of 20 trials.

"Reach" was scored when the animal extended its forelimb through the slot. A "success" was scored if

the animal grasped the food, retracted the paw and successfully brought the pellet back to its mouth and

consumed it in a single reach. On each day, immediately after training/testing and before sleep all mice

received either drug or saline and were allowed to sleep. On 9[th] day of training/testing all mice were filmed

with a Panasonic HDC-SDT750 camera at 60 frames per second with an exposure rate of 1ms.

Illumination for filming was obtained from a two-arm cold light source (Nikon Inc.). Quality of reaches

was assessed from the videos by the scoring method developed by Whishaw (Whishaw 2000).

Components of reach related to, hind feet, fore feet, sniff, lift, elbow in, advance, pronation, grasp,

supination 1, supination 2, release, and replace were scored. 0, 0.5, and 1 scores were given for good,

impaired, and abnormal reaches respectively. A higher score thus indicated inferior performance. On each

day of testing the order of the animals to be tested was randomized and the experimenter was blind to the

two groups.

Drugs and Solutions

The drugs Physostigmine Salicylate, Scopolamine Hydrobromide, Mecamylamine Hydrochloride,

Oxotremorine M, and Nicotine hydrogen tartrate salt were used in this study. All drugs were obtained

from Sigma-aldrich and dissolved in saline for administration. Saline was injected in control animals.

Physostigmine was first dissolved in 100% ethyl alcohol to make a stock solution of 10mg/ml and later

diluted 1000 times in saline to obtain a final concentration of 0.01mg/ml. All Drugs were prepared just

before use or diluted from a stock solution. The drugs were administered intraperitoneally. For rotarod

task, mice were assigned to six groups. Each group received one of the drugs; physostigmine (P group),

Oxotremorine (Oxo group), Nicotine (Nico Group), Scopolamine and Mecamylamine (SM group), and

Saline (S group). For skilled reach task, mice were assigned to two groups with one receiving

physostigmine and the other saline.

Pharmacological Treatment.

Physostigmine group (Physo group)

To increase cholinergic tone during SWS 14 mice were administered with physostigmine (an acetylcholine

esterase inhibitor which indirectly stimulates both muscarinic and nicotinic receptors) intraperitoneally

right after the first rotarod training session (Gais and Born 2004, Jafari-Sabet, Jafari-Sabet et al. 2016,

Dhingra and Soni 2018). Since it was a single dose study an acute dosage of physostigmine 0.1mg/kg was

used. After the training session and drug administration, the animals were transferred to their home cages

and allowed to sleep. The elimination half-life of physostigmine is documented to be 1.5-2 hours in

humans (Hartvig, Wiklund et al. 1986) and less than 1 hour in rodents (Somani 1989) enough to stay in

the system for the course of early SWS. No side effects were observed. Probe session was done on fifth

day. In the control condition, 14 mice received saline (S group).

Physostigmine-late Group (Physo-late 24hrs)

Next, we wanted to determine whether or not the performance of the animals on probe day was affected

because of the side-effects of physostigmine injected on Day 1. We therefore trained 6 mice on the rotarod

on Day 1 but instead of giving the drug right after the rotarod session and before sleep, we injected with

same dose of physostigmine 24 hours later. Retesting was done similarly on Day 5 as with Physo group.

Scopolamine/mecamylamine group (Sco+Meca group).

To block cholinergic transmission, we administered a combination of the muscarinic receptor antagonist

scopolamine (0.4mg/kg) and the nicotinic receptor antagonist mecamylamine (3mg/kg) intraperitoneally

to 8 mice. Previous studies have shown that these drugs if not given in combination do not produce any

effect on declarative memory encoding and consolidation (Rasch, Born et al. 2006). Therefore, both

substances were administered simultaneously, after the end of the rotarod session on Day 1. Animals

were then transferred to their home cages and allowed to sleep. We chose relatively low doses to avoid

strong side effects of cholinergic blockade and to ensure that the substances had largely washed out at the

time of probe testing. The half-life in plasma has been estimated at 4.5 ± 1.7 hour for scopolamine (Putcha,

Cintron et al. 1989) and 10.1 ± 2 hour for mecamylamine (Young, Shytle et al. 2001).

Oxotremorine and nicotine Groups (Oxo and Nico respectively)

To observe ACh receptor specificity for mediation of motor memory consolidation, we administered either

oxotremorine (0.01mg/kg) or Nicotine (2mg/kg), a selective muscarinic and nicotinic ACh receptor

agonist respectively, to two separate groups of animals (oxotremorine n=8, nicotine n=8) after the rotarod

task. The doses were chosen based on past studies that examined the effect of nicotine and oxotremorine

on rodents (Power, Vazdarjanova et al. 2003). The half-life has been reported as 100 minutes for

oxotremorine and 1.5-2 hours for nicotine (Moyer, Charlson et al. 2002, Marchand, Brossard et al. 2017).

Animals were then transferred to their home cages and were allowed to sleep. Probe session was done on
Day 5.

Baseline group

598   All 58 WT mice trained on Day 1 were combined to form one baseline group. The performance of mice
599   in other groups mentioned above was compared to the baseline group.

Transgenic Groups

17 Transgenic mice, 9 in CNO group (Tg-CNO) and 8 in saline group (Tg-Saline) were used. The dosage
of CNO used was 5mg/kg (MacLaren, Browne et al. 2016). CNO was obtained from the Thermo-Fisher
Scientific and was dissolved in 10% DMSO then diluted to a final concentration of 0.5 mg/ml CNO with
saline solution. As the onset of the action of CNO begins 1-hour post administration, it was injected just
before the rotarod training session so that the effect of drug will start in the post-learning sleep after the
training. Probe session was done on Day 5 as explained above.

Motion quantification with filming in home cage

In a separate cohort of mice (n = 10), sleep recording was done by filming animals in their home cages
after the rotarod training session to observe their motion for quantification of post-learning activity. Five
mice were given saline while the rest received physostigmine. A Pi camera connected to a Rasp-berry pi
computer was attached to the roof of the lid of the home cage and 2.5 hours long videos were recorded
post training (Singh, Bermudez-Contreras et al. 2018). Motion estimation was done in Matlab$^{®}$
(Mathworks, Natick, MA) by first identifying difference of consecutive frames and then calculating the
mean difference over pixels to get an approximation of the overall change in the values of pixels from one
frame to the other. The accuracy of motion estimation was verified with manual observation.

Sleep quantification using electrophysiology

Animal Surgeries

Four adult C57BL6J mice and one Chat-Cre::CAG-hM4Di were anesthetized with isoflurane (2.5%
induction, 1-1.5% maintenance) and implanted with cortical, hippocampal, and muscular electrodes using
aseptic techniques. For cortical and hippocampal recording of local field potentials, bipolar (tip separation

= 0.6 mm) and monopolar electrodes made from Teflon-coated stainless-steel wire (bare diameter 50.8

µm) were implanted in cortical areas and pyramidal layer of CA1 hippocampal region according to the

following coordinates in mm: primary motor cortex (M1): AP: 1.5, ML: 1.7, DV: 1.5, secondary motor

cortex (M2): AP: 1.7, ML: 0.6, DV:1.1 mm, Lip cortical sensory area (LP-S1): AP: 0.75, ML: 3.0, DV:

1.6, retrosplenial cortex (RS): AP: -2.5, ML: 0.6, DV: 1.1, and hippocampus (HPC): AP: -2.5, ML: 2.0,

DV:1.1 mm. For EMG, a multistranded Teflon-coated stainless-steel wire (gauge 40) was implanted into

the neck musculature using a 25 gauge needle. The reference and ground screws were placed on the skull

over the cerebellum. The other end of electrode wires were clamped between two receptacle connectors

(Mill-Max Mfg. Corp.) and the headpiece was secured to the skull using metabond and dental cement.

Electrophysiology

Animals were allowed to recover for 10 days after surgery and then habituated for 5 to 7 days in the

recording setup. On baseline days, animals were injected with saline at 8:25 am and moved to the

recording setup where their sleep activity was recorded from 8:30 am for 4 hours using a motorized

commutator (NeuroTek Inc.). Baseline recording was repeated for three days. On fourth day, same steps

were repeated except animals were injected with physostigmine (dosage 0.1 mg/kg). Local filed potentials

and EMG activty were amplified, filtered (0.1-4000 Hz) and digitized at 16 kHz using a Digital Lynx SX

Electrophysiology System (Neuralynx, Inc.) and the data were recorded and stored on a local PC using

Cheetah software (Neuralynx, Inc.). A Pi camera was used to record animal's behaviour during the

recording.

Data Analysis

All the analyses were performed offline. Sleep scoring was performed in 6-sec long epochs. For that, raw

EMG activity was filtered between 90 Hz and 1000 Hz and then rectified and integrated using 4-sec

moving windows. This signal was thresholded to detect periods of immobility. Slow wave power (0.5 to

4 Hz) of the cortical LFP was calculated in each epoch using taper spectral analysis and thresholded using

values between 0.04 and 0.1 mV^2/Hz (depending on the animal) to detect SWA in cortical recording.

When the animal was immobile and cortical LFP showed SWA, the epoch was scored as SWS. For

detecting REM sleep, the ratio of theta power (6 to 9 Hz) to the total power of hippocampal LFP was

calculated in each epoch using taper spectral analysis and when this ratio was above 0.4-0.6 (different for

individual animals) and EMG showed immobility, the state was scored as REM sleep. All other epochs

were considered as waking state. These scorings were further confirmed using video recording. To

investigate the effect of physostigmine on sleep, duration of each behavioral state in each recording session

was normalized to the recording duration and then compared between physostigmine and saline. SWS

latency was calculated as the time between the onset of recording and the onset of first SWS epoch longer

than 20 seconds. Moreover, slow wave power (0.5 to 4 Hz) of cortical signal was averaged across all SWS

epochs in the first hour of recording and then compared between physostigmine and saline. For saline

injection, sleep structure, SWS latency and SW power were averaged across three baseline days for each

animal.

For detecting spindle events, following Philips et. al., M2 cortical signal was filtered between 8 and 18

Hz and then rectified (Phillips, Bartsch et al. 2012). The envelope of the rectified signal was then

calculated using cubic interpolation of its maxima. Putative spindle events were detected where the

envelope signal was higher than 3.5 standard deviations (SD) from its mean. Onset and end of spindle

events was determined where the envelope signal was higher than 2.5 SD from its mean. Spindles shorter

than 500msec or longer than 2sec were rejected. Spindles that occurred within 100 msec of each other

were combined to be considered a single event.

Sleep deprivation

666   Animals were observed for signs of sleep such as little motion and closed eyes and were gently moved

667   with hand to prevent them from sleeping. Place of their nest was also changed sometimes to keep them

668   awake for a little longer period before they would begin to sleep again. Occasionally, their cage was lifted

669   and put down to keep them awake. However, animals were not disturbed when they were already awake.

Statistical Analysis

Statistical analysis of behavioral data was done in SPSS (Version 22) and verified in Matlab R2016 while

all other statistical analysis was done in Matlab R2016. All data are presented as mean $\pm$ the standard error

of mean (S.E.M). Statistical tests, paired and two sample Student's t-test (both two and one tailed), one-

way analysis of variance (ANOVA), and one-way ANOVA with repeated measures (ANOVA-RM) were

used. Statistical test are reported in the text along with p values. For ANOVA-RM, Greenhouse-Geisser

corrected p value (p_{gg}) is reported for tests in which Mauchly's criteria of sphericity failed (Mauchly 1940,

Geisser and Greenhouse 1958, Huynh and Feldt 1976). An alpha value of 0.05 was used for determining

significance.

FUNDING

This study was supported by research fund provided by the Canadian Institutes of Health Research (CIHR)

Grant# 390930, Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery

Grant #40352, Alberta Innovates (CAIP Chair) Grant #43568, Alberta Alzheimer Research Program Grant

PAZ15010 and PAZ17010, and Alzheimer Society of Canada Grant# 43674 to MHM; and NSERC

Discovery Grant RGPIN-2017-03857 and NSF Grant 1631465 to BLM.

ACKNOWLEDGEMENTS

We thank you Dr. Gerlinde Metz for providing us with the apparatus for performing the rotarod task. We

thank Behroo Mirza Agha for her help with reach task training and data analyses. We thank Jessica Kuntz

for her help with reach task data analyses. We thank Di Shao and the University of Lethbridge animal care

staff for animal husbandry. Finally, we thank Drs. Masami Tatsuno and Bryan Kolb for useful discussions

regarding the preparation of this manuscript.

AUTHOR CONTRIBUTIONS

Q, SI, IQW, BLM and MHM designed the study. Q and SI performed all rotarod and reach task

experiments. Q, SI, and IQW analyzed behavioral data. SI and SS collected and analyzed home cage video

recording data. For electrophysiology experiments, MN performed animal surgeries for implanting

electrodes, SI, Q, and MN collected data, and MN and SI analyzed data. MHM participated in all data

analyses. SI, Q, IQW, and MHM wrote the manuscript, which all authors commented on and edited. IQW

and MHM supervised the study.

SUBMISSION & CORRESPONDENCE INFORMATION

Dr. Majid Mohajerani, mohajerani@uleth.ca

WORK PERFORMED

All the work was performed at the Canadian Center for Behavioral Neuroscience, University of Lethbridge, Alberta, Canada

DISCLOSURE STATEMENT

None.

REFERENCES

1. Aeschbach, D., A. J. Cutler and J. M. Ronda (2008). "A role for non-rapid-eye-movement sleep homeostasis in perceptual learning." J Neurosci **28**(11): 2766-2772.
2. Alger, S. E., A. M. Chambers, T. Cunningham and J. D. Payne (2015). "The role of sleep in human declarative memory consolidation." Curr Top Behav Neurosci **25**: 269-306.
3. Ambrosini, M. V. and A. Giuditta (2001). "Learning and sleep: the sequential hypothesis." Sleep Med Rev **5**(6): 477-490.
4. Anaclet, C., N. P. Pedersen, L. L. Ferrari, A. Venner, C. E. Bass, E. Arrigoni and P. M. Fuller (2015). "Basal forebrain control of wakefulness and cortical rhythms." Nat Commun **6**: 8744.
5. Aquilonius, S. M. and P. Hartvig (1986). "Clinical pharmacokinetics of cholinesterase inhibitors." Clin Pharmacokinet **11**(3): 236-249.
6. Brown, D. A. (2010). "Muscarinic acetylcholine receptors (mAChRs) in the nervous system: some functions and mechanisms." J Mol Neurosci **41**(3): 340-346.
7. Buhry, L., A. H. Azizi and S. Cheng (2011). "Reactivation, replay, and preplay: how it might all fit together." Neural Plast **2011**: 203462.
8. Buitrago, M. M., T. Ringer, J. B. Schulz, J. Dichgans and A. R. Luft (2004). "Characterization of motor skill and instrumental learning time scales in a skilled reaching task in rat." Behav Brain Res **155**(2): 249-256.
9. Chambers, A. M. (2017). "The role of sleep in cognitive processing: focusing on memory consolidation." Wiley Interdiscip Rev Cogn Sci **8**(3).
10. Clopath, C. (2012). Synaptic consolidation: an approach to long-term learning. Cogn Neurodyn. **6:** 251-257.
11. Conner, J. M., A. Culberson, C. Packowski, A. A. Chiba and M. H. Tuszynski (2003). "Lesions of the Basal forebrain cholinergic system impair task acquisition and abolish cortical plasticity associated with motor skill learning." Neuron **38**(5): 819-829.
12. Conner, J. M., M. Kulczycki and M. H. Tuszynski (2010). "Unique contributions of distinct cholinergic projections to motor cortical plasticity and learning." Cereb Cortex **20**(11): 2739-2748.
13. Dhingra, D. and K. Soni (2018). "Behavioral and biochemical evidences for nootropic activity of boldine in young and aged mice." Biomed Pharmacother **97**: 895-904.
14. Diekelmann, S. and J. Born (2010). "The memory function of sleep." Nat Rev Neurosci **11**(2): 114-126.
15. Dringenberg, H. C. and C. H. Vanderwolf (1996). "5-Hydroxytryptamine (5-HT) agonists: effects on neocortical slow wave activity after combined muscarinic and serotonergic blockade." Brain Res **728**(2): 181-187.
16. Dringenberg, H. C. and C. H. Vanderwolf (1997). "Neocortical activation: modulation by multiple pathways acting on central cholinergic and serotonergic systems." Exp Brain Res **116**(1): 160-174.

17. Dudai, Y. (2004). "The neurobiology of consolidations, or, how stable is the engram?" <u>Annu Rev Psychol</u>
55: 51-86.
18. Dudai, Y., A. Karni and J. Born (2015). "The Consolidation and Transformation of Memory." <u>Neuron</u> **88**(1):
20-32.
19. Ebbinghaus, H. (1885-Republished Translation 1964). <u>Memory: A contribution to experimental</u>
<u>psychology</u>. Oxford, England, Dover.
20. Euston, D. R., M. Tatsuno and B. L. McNaughton (2007). "Fast-forward playback of recent memory
sequences in prefrontal cortex during sleep." <u>Science</u> **318**(5853): 1147-1150.
21. Frankland, P. W. and B. Bontempi (2005). "The organization of recent and remote memories." <u>Nat Rev</u>
<u>Neurosci</u> **6**(2): 119-130.
22. Gais, S. and J. Born (2004). "Low acetylcholine during slow-wave sleep is critical for declarative memory
consolidation." <u>Proc Natl Acad Sci U S A</u> **101**(7): 2140-2144.
23. Gais, S., W. Plihal, U. Wagner and J. Born (2000). "Early sleep triggers memory for early visual
discrimination skills." <u>Nat Neurosci</u> **3**(12): 1335-1339.
24. Gais, S. and M. Schonauer (2017). "Untangling a Cholinergic Pathway from Wakefulness to Memory."
<u>Neuron</u> **94**(4): 696-698.
25. Geisser, S. and S. W. Greenhouse (1958). "An Extension of Box's Results on the Use of the F Distribution
in Multivariate Analysis." <u>The Annals of Mathematical Statistics</u> **29**(3): 885-891.
26. Gharbawie, O. A. and I. Q. Whishaw (2003). "Cholinergic and serotonergic neocortical projection lesions
given singly or in combination cause only mild impairments on tests of skilled movement in rats:
evaluation of a model of dementia." <u>Brain Res</u> **970**(1-2): 97-109.
27. Giuditta, A., M. V. Ambrosini, P. Montagnese, P. Mandile, M. Cotugno, G. Grassi Zucconi and S. Vescia
(1995). "The sequential hypothesis of the function of sleep." <u>Behav Brain Res</u> **69**(1-2): 157-166.
28. Gomez, J. L., J. Bonaventura, W. Lesniak, W. B. Mathews, P. Sysa-Shah, L. A. Rodriguez, R. J. Ellis, C. T. Richie,
B. K. Harvey, R. F. Dannals, M. G. Pomper, A. Bonci and M. Michaelides (2017). "Chemogenetics revealed:
DREADD occupancy and activation via converted clozapine." <u>Science</u> **357**(6350): 503-507.
29. Hanlon, E. C., U. Faraguna, V. V. Vyazovskiy, G. Tononi and C. Cirelli (2009). "Effects of skilled training on
sleep slow wave activity and cortical gene expression in the rat." <u>Sleep</u> **32**(6): 719-729.
30. Hartvig, P., L. Wiklund and B. Lindstrom (1986). "Pharmacokinetics of physostigmine after intravenous,
intramuscular and subcutaneous administration in surgical patients." <u>Acta Anaesthesiol Scand</u> **30**(2):
177-182.
31. Hasselmo, M. E. (1999). "Neuromodulation: acetylcholine and memory consolidation." <u>Trends Cogn Sci</u>
3(9): 351-359.
32. Hasselmo, M. E. and J. McGaughy (2004). "High acetylcholine levels set circuit dynamics for attention and
encoding and low acetylcholine levels set dynamics for consolidation." <u>Prog Brain Res</u> **145**: 207-231.
33. Huber, R., M. F. Ghilardi, M. Massimini, F. Ferrarelli, B. A. Riedner, M. J. Peterson and G. Tononi (2006).
"Arm immobilization causes cortical plastic changes and locally decreases sleep slow wave activity." <u>Nat</u>
<u>Neurosci</u> **9**(9): 1169-1176.
34. Hut, R. A. and E. A. Van der Zee (2011). "The cholinergic system, circadian rhythmicity, and time memory."
<u>Behav Brain Res</u> **221**(2): 466-480.
35. Huynh, H. and L. S. Feldt (1976). "Estimation of the Box Correction for Degrees of Freedom from Sample
Data in Randomized Block and Split-Plot Designs." <u>Journal of Educational Statistics</u> **1**(1): 69-82.
36. Jafari-Sabet, M., A. R. Jafari-Sabet and A. Dizaji-Ghadim (2016). "Tramadol state-dependent memory:
involvement of dorsal hippocampal muscarinic acetylcholine receptors." <u>Behav Pharmacol</u> **27**(5): 470-
478.
37. Jenkins, J. G. and K. M. Dallenbach (1924). "Obliviscence during Sleep and Waking." <u>The American Journal</u>
<u>of Psychology</u> **35**(4): 605-612.
38. Kali, S. and P. Dayan (2004). "Off-line replay maintains declarative memories in a model of hippocampal-
neocortical interactions." <u>Nat Neurosci</u> **7**(3): 286-294.

39. Karni, A., G. Meyer, C. Rey-Hipolito, P. Jezzard, M. M. Adams, R. Turner and L. G. Ungerleider (1998). "The
acquisition of skilled motor performance: fast and slow experience-driven changes in primary motor
cortex." Proc Natl Acad Sci U S A **95**(3): 861-868.
40. Kudrimoti, H. S., C. A. Barnes and B. L. McNaughton (1999). "Reactivation of hippocampal cell assemblies:
effects of behavioral state, experience, and EEG dynamics." J Neurosci **19**(10): 4090-4101.
41. Landsness, E. C., D. Crupi, B. K. Hulse, M. J. Peterson, R. Huber, H. Ansari, M. Coen, C. Cirelli, R. M. Benca,
M. F. Ghilardi and G. Tononi (2009). "Sleep-dependent improvement in visuomotor learning: a causal role
for slow waves." Sleep **32**(10): 1273-1284.
42. Landsness, E. C., F. Ferrarelli, S. Sarasso, M. R. Goldstein, B. A. Riedner, C. Cirelli, B. Perfetti, C. Moisello, M.
F. Ghilardi and G. Tononi (2011). "Electrophysiological traces of visuomotor learning and their
renormalization after sleep." Clin Neurophysiol **122**(12): 2418-2425.
43. Li, W., L. Ma, G. Yang and W. B. Gan (2017). "REM sleep selectively prunes and maintains new synapses
in development and learning." Nat Neurosci **20**(3): 427-437.
44. Louie, K. and M. A. Wilson (2001). "Temporally structured replay of awake hippocampal ensemble
activity during rapid eye movement sleep." Neuron **29**(1): 145-156.
45. MacLaren, D. A., R. W. Browne, J. K. Shaw, S. Krishnan Radhakrishnan, P. Khare, R. A. Espana and S. D.
Clark (2016). "Clozapine N-Oxide Administration Produces Behavioral Effects in Long-Evans Rats:
Implications for Designing DREADD Experiments." eNeuro **3**(5).
46. Mahler, S. V. and G. Aston-Jones (2018). "CNO Evil? Considerations for the Use of DREADDs in Behavioral
Neuroscience." Neuropsychopharmacology **43**(5): 934.
47. Manvich, D. F., K. A. Webster, S. L. Foster, M. S. Farrell, J. C. Ritchie, J. H. Porter and D. Weinshenker (2018).
"The DREADD agonist clozapine N-oxide (CNO) is reverse-metabolized to clozapine and produces
clozapine-like interoceptive stimulus effects in rats and mice." Sci Rep **8**(1): 3840.
48. Marchand, M., P. Brossard, H. Merdjan, N. Lama, R. Weitkunat and F. Ludicke (2017). "Nicotine Population
Pharmacokinetics in Healthy Adult Smokers: A Retrospective Analysis." Eur J Drug Metab Pharmacokinet
42(6): 943-954.
49. Mascetti, L., V. Muto, L. Matarazzo, A. Foret, E. Ziegler, G. Albouy, V. Sterpenich, C. Schmidt, C. Degueldre,
Y. Leclercq, C. Phillips, A. Luxen, G. Vandewalle, R. Vogels, P. Maquet and E. Balteau (2013). "The impact
of visual perceptual learning on sleep and local slow-wave initiation." J Neurosci **33**(8): 3323-3331.
50. Mauchly, J. W. (1940). "Significance Test for Sphericity of a Normal n-Variate Distribution." Ann. Math.
Statist. **11**(2): 204-209.
51. McGaugh, J. L. (2000). "Memory--a century of consolidation." Science **287**(5451): 248-251.
52. Mednick, S., K. Nakayama and R. Stickgold (2003). "Sleep-dependent learning: a nap is as good as a night."
Nat Neurosci **6**(7): 697-698.
53. Miyamoto, D., D. Hirai and M. Murayama (2017). "The Roles of Cortical Slow Waves in Synaptic Plasticity
and Memory Consolidation." Front Neural Circuits **11**: 92.
54. Mohajerani, M. H., A. W. Chan, M. Mohsenvand, J. LeDue, R. Liu, D. A. McVea, J. D. Boyd, Y. T. Wang, M.
Reimers and T. H. Murphy (2013). "Spontaneous cortical activity alternates between motifs defined by
regional axonal projections." Nat Neurosci **16**(10): 1426-1435.
55. Moyer, T. P., J. R. Charlson, R. J. Enger, L. C. Dale, J. O. Ebbert, D. R. Schroeder and R. D. Hurt (2002).
"Simultaneous analysis of nicotine, nicotine metabolites, and tobacco alkaloids in serum or urine by
tandem mass spectrometry, with clinically relevant metabolic profiles." Clin Chem **48**(9): 1460-1471.
56. Nagai, H., L. de Vivo, M. Bellesi, M. F. Ghilardi, G. Tononi and C. Cirelli (2017). "Sleep Consolidates Motor
Learning of Complex Movement Sequences in Mice." Sleep **40**(2).
57. Peigneux, P., S. Laureys, S. Fuchs, A. Destrebecqz, F. Collette, X. Delbeuck, C. Phillips, J. Aerts, G. Del Fiore,
C. Degueldre, A. Luxen, A. Cleeremans and P. Maquet (2003). "Learned material content and acquisition
level modulate cerebral reactivation during posttraining rapid-eye-movements sleep." Neuroimage
20(1): 125-134.
58. Peters, A. J., S. X. Chen and T. Komiyama (2014). "Emergence of reproducible spatiotemporal activity
during motor learning." Nature **510**(7504): 263-267.

59. Phillips, K., U. Bartsch, A. McCarthy, D. Edgar, M. Tricklebank, K. Wafford and M. Jones (2012). Decoupling
of Sleep-Dependent Cortical and Hippocampal Interactions in a Neurodevelopmental Model of
Schizophrenia. Neuron. **76:** 526-533.
60. Plihal, W. and J. Born (1997). "Effects of early and late nocturnal sleep on declarative and procedural
memory." J Cogn Neurosci **9**(4): 534-547.
61. Power, A. E., A. Vazdarjanova and J. L. McGaugh (2003). "Muscarinic cholinergic influences in memory
consolidation." Neurobiol Learn Mem **80**(3): 178-193.
62. Pubchem. (2018). "Physostigmine." from https://www.ncbi.nlm.nih.gov/pubmed/.
63. Putcha, L., N. M. Cintron, J. Tsui, J. M. Vanderploeg and W. G. Kramer (1989). "Pharmacokinetics and oral
bioavailability of scopolamine in normal subjects." Pharm Res **6**(6): 481-485.
64. Ramanathan, D., M. H. Tuszynski and J. M. Conner (2009). "The basal forebrain cholinergic system is
required specifically for behaviorally mediated cortical map plasticity." J Neurosci **29**(18): 5992-6000.
65. Rasch and J. Born (2013). "About sleep's role in memory." Physiol Rev **93**(2): 681-766.
66. Rasch, B. and J. Born (2013). "About sleep's role in memory." Physiol Rev **93**(2): 681-766.
67. Rasch, B. H., J. Born and S. Gais (2006). "Combined blockade of cholinergic receptors shifts the brain from
stimulus encoding to memory consolidation." J Cogn Neurosci **18**(5): 793-802.
68. Shiotsuki, H., K. Yoshimi, Y. Shimo, M. Funayama, Y. Takamatsu, K. Ikeda, R. Takahashi, S. Kitazawa and N.
Hattori (2010). "A rotarod test for evaluation of motor skill learning." J Neurosci Methods **189**(2): 180-
185.
69. Singh, S., E. Bermudez-Contreras, M. Nazari, R. J. Sutherland and M. Mohajerani (2018). "Low-Cost
Solution for Rodent Home-Cage Behaviour Monitoring." bioRxiv.
70. Sitaram, N., R. J. Wyatt, S. Dawson and J. C. Gillin (1976). "REM sleep induction by physostigmine infusion
during sleep." Science **191**(4233): 1281-1283.
71. Somani, S. M. (1989). "Pharmacokinetics and pharmacodynamics of physostigmine in the rat after oral
administration." Biopharm Drug Dispos **10**(2): 187-203.
72. Stachniak, T. J., A. Ghosh and S. M. Sternson (2014). "Chemogenetic synaptic silencing of neural circuits
localizes a hypothalamus-->midbrain pathway for feeding behavior." Neuron **82**(4): 797-808.
73. Stickgold, R., D. Whidbee, B. Schirmer, V. Patel and J. A. Hobson (2000). "Visual discrimination task
improvement: A multi-step process occurring during sleep." J Cogn Neurosci **12**(2): 246-254.
74. Sutherland, G. R. and B. McNaughton (2000). "Memory trace reactivation in hippocampal and neocortical
neuronal ensembles." Curr Opin Neurobiol **10**(2): 180-186.
75. Tononi, G. and C. Cirelli (2003). "Sleep and synaptic homeostasis: a hypothesis." Brain Res Bull **62**(2):
143-150.
76. Tononi, G. and C. Cirelli (2006). "Sleep function and synaptic homeostasis." Sleep Med Rev **10**(1): 49-62.
77. Tononi, G. and C. Cirelli (2014). "Sleep and the price of plasticity: from synaptic and cellular homeostasis
to memory consolidation and integration." Neuron **81**(1): 12-34.
78. Vanderwolf, C. H., A. Raithby, M. Snider, C. Cristi and C. Tanner (1993). "Effects of some cholinergic
agonists on neocortical slow wave activity in rats with basal forebrain lesions." Brain Res Bull **31**(5):
515-521.
79. Vardy, E., J. E. Robinson, C. Li, R. H. Olsen, J. F. DiBerto, P. M. Giguere, F. M. Sassano, X. P. Huang, H. Zhu, D.
J. Urban, K. L. White, J. E. Rittiner, N. A. Crowley, K. E. Pleil, C. M. Mazzone, P. D. Mosier, J. Song, T. L. Kash,
C. J. Malanga, M. J. Krashes and B. L. Roth (2015). "A New DREADD Facilitates the Multiplexed
Chemogenetic Interrogation of Behavior." Neuron **86**(4): 936-946.
80. Walker, M. P. (2009). "The Role of Slow Wave Sleep in Memory Processing." J Clin Sleep Med **5**(2 Suppl):
S20-26.
81. Whishaw, I. Q. (2000). "Loss of the innate cortical engram for action patterns used in skilled reaching and
the development of behavioral compensation following motor cortex lesions in the rat."
Neuropharmacology **39**(5): 788-805.

82. Whishaw, I. Q., J. Faraji, J. Kuntz, B. Mirza Agha, M. Patel, G. A. S. Metz and M. H. Mohajerani (2017).
"Organization of the reach and grasp in head-fixed vs freely-moving mice provides support for multiple
motor channel theory of neocortical organization." Exp Brain Res **235**(6): 1919-1932.
83. Whishaw, I. Q., P. Whishaw and B. Gorny (2008). "The structure of skilled forelimb reaching in the rat: a
movement rating scale." J Vis Exp(18).
84. Wilson, M. A. and B. L. McNaughton (1994). "Reactivation of hippocampal ensemble memories during
sleep." Science **265**(5172): 676-679.
85. Yang, G., C. S. Lai, J. Cichon, L. Ma, W. Li and W. B. Gan (2014). "Sleep promotes branch-specific formation
of dendritic spines after learning." Science **344**(6188): 1173-1178.
86. Yaroush, R., M. J. Sullivan and B. R. Ekstrand (1971). "Effect of sleep on memory. II. Differential effect of
the first and second half of the night." J Exp Psychol **88**(3): 361-366.
87. Young, J. M., R. D. Shytle, P. R. Sanberg and T. P. George (2001). "Mecamylamine: new therapeutic uses
and toxicity/risk profile." Clin Ther **23**(4): 532-565.
88. Zhu, H., D. K. Aryal, R. H. Olsen, D. J. Urban, A. Swearingen, S. Forbes, B. L. Roth and U. Hochgeschwender
(2016). "Cre-dependent DREADD (Designer Receptors Exclusively Activated by Designer Drugs) mice."
Genesis **54**(8): 439-446.
89. Zwart, R., H. Reed and E. Sher (2018). "Oxotremorine-M potentiates NMDA receptors by muscarinic
receptor dependent and independent mechanisms." Biochem Biophys Res Commun **495**(1): 481-486.

911