Unpredictable effects of the genetic background of transgenic lines in physiological quantitative traits

Amalia Evangelou*, Anastasia Ignatiou*, Chloi Antoniou*, Sofia Kalanidou*, Sotiroulla Hadjimatheou, Sotiroulla Ellina, Rafaella Athanasiou, Myrofora Panagi, Yiorgos Apidianakis[#], Chrysoula Pitsouli[#]

University of Cyprus, Department of Biological Sciences, P.O. Box 20537, Nicosia, Cyprus

* These authors contributed equally

[#] Correspondence should be addressed to Chrysoula Pitsouli (pitsouli@ucy.ac.cy) or Yiorgos Apidianakis (apidiana@ucy.ac.cy)

ABSTRACT

Most physiology, fitness and disease phenotypes are considered complex traits since they exhibit continuous variation in natural populations. The use of homozygous inbred strains allows the assessment of the impact of genetic composition on complex phenotypes. Nevertheless, to understand gene function, we often combine transgenic lines to assess the phenotype of interest in a heterozygous state. Here, using inbred wild-type *Drosophila* strains, we assessed the phenotypic variation of six physiology and fitness traits: female fecundity, survival and intestinal mitosis upon oral infection, defecation rate and fecal pH upon oral infection, and terminal tracheal cell branching in hypoxia. Indeed, we found continuous variation in all traits tested. To estimate the extent to which small genetic background differences affected these traits, we assessed the effects of commonly used *Drosophila UAS-RNAi* transgenic strains and their backcrossed isogenized counterparts in the same assays. We tested 20 non-isogenic strains (10 KK and 10 GD) from the Vienna *Drosophila* Resource Center and their isogenic counterparts without Gal4 induction and found that the different phenotypes were variably sensitive to the genetic background. For example, survival upon infection and female fecundity exhibited differences in 50% and 40% of the tested isogenic vs. non-isogenic pairs, respectively, whereas all other phenotypes were affected in only 10-25% of the cases. We also observed that the genetic background affects Gal4 transgenes unpredictably. Thus, depending on the trait of interest, the genetic background of commonly used transgenic strains needs to be considered carefully during experimentation.

INTRODUCTION

Most physiological, morphological, biochemical, behavioral, fitness and disease phenotypes exhibit continuous variation among natural populations and are therefore considered as complex or quantitative traits. This continuity in variation is attributed to a combination of genetic composition, environmental effects and gene-environment interactions (Mackay, 2009). To assess the impact of environmental factors on distinct but constant genetic backgrounds to tease out the effect of the genotype from that of the environment, inbred lines can be used. Naturally-occurring phenotypic variation in distinct inbred backgrounds allows the implementation of genome-wide association (GWA) and the identification of key genetic variants impinging on the phenotype under study. The *Drosophila* Genetics Reference Panel (DGRP) collection corresponds to a set of >200 inbred sequenced fruit fly strains, which facilitate GWA studies (Mackay et al., 2012; Huang et al., 2014,). GWAS analyses using the DGRP collection have identified significant genomic associations with a variety of fitness and physiology traits ranging from lifespan and sleep to abdominal pigmentation (Durham et al., 2014; Ivanov et al., 2015; Harbison et al., 2013; Dembeck et al., 2015; Sunaga et al., 2016).

The natural variation of the wild-type genetic background not only affects complex traits, but it often also affects the expressivity and penetrance of different mutations. Genetic background effects have been observed in all model organisms tested, including fruit flies, mice, nematodes, yeast, plants and bacteria (Chandler et al., 2013). These effects potentially led to contradictory results across studies that investigated the role of genes affecting longevity, such as *Indy* and *sir-2*, whereby the original papers that implicated the genes in increased longevity (Rogina et al., 2000; Rogina et al., 2004) could not be replicated in subsequent studies performed in different wild-type genetic backgrounds (Toivonen et al., 2007; Burnett et al., 2011; Viswanathan and Guarente, 2011). Furthermore, extensive studies assessing the phenotype of various alleles of the gene *sd* in *Drosophila*, which affect wing size, have shown that the same allele is able to generate a continuum of phenotypes when introduced in different wild-type genetic backgrounds (Chari and Dworkin, 2013; Dowrkin et al., 2009). In addition, it has been shown that the genetic background can also affect RNAi phenotypes. For example, injection of RNAi constructs in different wild-type genetic backgrounds of *Tribolium* led to distinct phenotypes, which were attributed not to off-target effects or differences in the function of the RNAi machinery, but to maternal effects (Kitzman et al., 2013). In conclusion, such phenotypic inconsistencies underscore the importance of controlling for the genetic background in similar studies.

Backcrossing is an established breeding scheme where a characteristic (i.e. a trait, a gene, a locus or a chromosome segment) is introgressed from a donor parent into the genomic background of a recurrent parent. Progeny of successive generations are selected for the characteristic of interest and subsequently backcrossed to the recurrent parent. As backcrossed generations accumulate, the proportion of the genome of the donor parent tends to zero, except of the part hosting the characteristic of interest. Backcrossing and introgression have long been used in breeding improvement programs, but backcrossing is also particularly useful in dissecting the genetic architecture of quantitative traits, because it can isolate a gene or chromosomal locus of interest in different genetic backgrounds (Hospital, 2005). Isogenic (or congenic) strains derive from backcrossing schemes and contain nearly identical genomes; namely, the genome of the recurrent parent except from the small part that is selected to differ (Kooke et al., 2012). Isogenic strains incorporating specific

transgenes (with P or other transposable element ends or phage ends) in different genetic backgrounds can be easily generated in *Drosophila* using phenotypic markers of transgenesis (i.e. eye color, fluorescence) as a tool for selection.

Here, we used *Drosophila melanogaster* to assess the effects of the genetic background in six different fitness and physiology traits (female fecundity, survival and intestinal stem cell mitosis upon oral bacterial infection, defecation rate and fecal pH upon oral bacterial infection, and tracheal branching upon hypoxia). First, we assessed natural phenotypic variation of these traits using the DGRP collection. Then, we modified the genetic background of *UAS-RNAi* transgenic lines and compared the acquired phenotypes in isogenic vs non-isogenic lines. Finally, we modified the genetic background of Gal4 and assessed its phenotypic consequences. We found that the genetic background affects different traits to different extents and we propose ways to control for such effects in future experiments.

RESULTS

The genetic background plays a key role to the phenotype of complex traits. To understand the complexity of various quantitative fitness and physiology traits of interest, we screened the DGRP collection of wild-type inbred isogenic strains (Mackay et al., 2012) for female fecundity, survival upon oral bacterial infection, intestinal physiology upon oral bacterial infection (intestinal stem cell mitosis-mediated regeneration, defecation rate and fecal pH) and cellular response to hypoxia (terminal tracheal branching). All traits were assessed in *Drosophila* adults, except tracheal branching, which was assessed in third instar larvae. Z-score analysis underscored the continuity of the observed phenotypes for each trait (Figure 1). The standard deviation range of the screened DGRP strains, which indicates the phenotypic range, for each screen equaled 5.57, 5.24, 5.43, 4.28, 4.55 and 4.85 for fecundity, survival, intestinal mitosis, defecation rate, fecal pH and branching, respectively. Genome wide association (GWA) analysis of each screen via the DGRP2 algorithm (http://dgrp2.gnets.ncsu.edu/) identified 36, 44, 95, 32, 18 and 21 variants associated with fecundity, survival, intestinal mitosis, defecation rate, fecal pH and branching, respectively. To assess the correlation between the six DGRP screens, we performed pair-wise correlation analyses. No significant correlation was observed between screens (Figure 2). A weak correlation was found between the defecation rate and fecal pH (R^2=0.1345, p=0.1) (Fig. 2L) with all other comparisons having essentially no correlation (R^2<0.04) (Fig. 2A-K, M-O).

The genetic background of UAS-RNAi transgenes (without Gal4) affects complex traits to different extents. Since the effect of the homozygous wild-type genotype produces distinct quantitative phenotypes in all traits studied, we devised a way to assess the effects of heterozygosis in the above traits. The use of heterozygotes is necessary in most genetic experiments utilizing the classical genetic tools available in *Drosophila*, such as the Gal4-UAS system (Brand and Perrimon, 1993) and the ever-increasing number of *UAS-RNAi* lines targeting practically any gene of the fly genome. Initially, to avoid the effect of overexpression by the Gal4, we decided to assess the effect of genetic background on the *UAS-RNAi* insertion alone. All *UAS-RNAi* lines were generated in an isogenic *w[1118]* host strain (VDRC) and the balancer stocks used to map or stabilize insertions were also isogenic to the VDRC *w[1118]* strain. Therefore, and to maintain the insertions as closely as possible to their original genotype, we decided to introgress the *UAS-RNAi* insertions of twenty randomly-selected VDRC lines in our laboratory *w[1118]* strain for further experiments. Ten strains from the VDRC KK library, generated via site-specific *φχ31* recombination (Markstein et al., 2008), and ten strains from

the VDRC GD library (Dietzl et al., 2007), generated via P-element random transposition (Spradling and Rubin, 1982), targeting in pairs the same gene were used. To generate "isogenic" *UAS-RNAi* strains, we backcrossed each VDRC KK and GD strain (from now on called "non-isogenic") to our laboratory w^{1118} strain for at least six generations. During backcrossing the fraction of the donor parent (VDRC line) halves after every backcross: 50%, 25%, 12.5% etc., in F1, 1st backcross, 2nd backcross etc., respectively. The average proportion of the recurrent parental genome after each backcross increases and can be calculated by the formula: $(2^{(b+1)} - 1)/2^{(b+1)}$ or $1 - (1/2)^{(b+1)}$, where b equals the number of backcrosses assuming an infinite population. Thus, the proportion of the donor genome is given as $(1/2)^{(b+1)}$ (Kooke et al., 2012). Assuming six generations of backcrossing, the recurrent parental genome (our laboratory w^{1118}) equals 99.21% and the donor genome (VDRC w^{1118}) 0.79%.

The homozygous non-isogenic (VDRC) and isogenic (VDRC in our w^{1118} strain) lines were subsequently crossed to w^{1118}, the heterozygous progeny of the crosses were collected and assessed for eight fitness and physiology traits. In addition to the traits studied in the DGRP screening, we also tested intestinal mitosis during homeostasis (without infection) and tracheal branching in normoxia (21% O_2). GD and KK non-isogenic and isogenic lines were assessed in pairs and those producing significantly different phenotypes were quantified (Figure 3). In this scheme, the progeny of the non-isogenic cross to our laboratory w^{1118} contain 50% of our laboratory w^{1118} genome and 50% of the VDRC w^{1118} genome, whereas the progeny of the isogenic cross to our laboratory w^{1118} contain approximately 100% of our laboratory w^{1118} genome. Therefore, the genome of the progeny of the two crosses differs by up to 50%. Theoretically, since the VDRC lines were generated in a w^{1118} genetic background, we expected not to find significant differences between the original VDRC strains (non-isogenic) and the introgressed lines (isogenic). Strikingly, we observed different impact of isogenization in the various traits and this was independent of the type of transgene (GD, inserted in a random genomic location; or KK, inserted in a specific genomic location); the most affected being survival and fecundity, with 50% and 40% of the isogenic vs. non-isogenic pairs producing significantly different phenotypes (Figure 3A-D), respectively, and the least affected being tracheal branching in normoxia and hypoxia with only 10% of the isogenic vs. non-isogenic pairs producing significantly different phenotypes (Figure 3M-P).

Specifically, 4 out of 10 GD (2 up, 2 down) and 4 out of 10 KK (2 up, 2 down) isogenic vs. non-isogenic pairs (40% in total) differed in fecundity (Figure 3A-B); 5 out of 10 GD (4 down, 1 up) and 5 out of 10 KK (4 down, 1 up) isogenic vs. non-isogenic pairs (50% in total) differed in survival upon pathogenic oral bacterial infection (Figure 3C-D); 2 out of 10 GD (1 up, 1 down) and 3 out of 10 KK (3 down) isogenic vs. non-isogenic pairs (25% in total) differed in intestinal mitosis in homeostatic conditions (Figure 3E-F); 4 out of 10 GD (1 down, 3 up) and 1 out of 10 KK (1 down) isogenic vs. non-isogenic pairs (25% in total) differed in intestinal mitosis upon oral pathogenic infection (Figure 3G-H); 2 out of 10 GD (2 down) and 2 out of 10 KK (1 up, 1 down) isogenic vs. non-isogenic pairs (20% in total) differed in defecation rate, i.e. number of fecal spots upon bacterial infection (Figure 3I-J); 2 out of 10 GD (2 up) and 2 out of 10 KK (2 down) isogenic vs. non-isogenic pairs (20% in total) differed in fecal pH upon bacterial infection (Figure K-L); 0 out of 10 GD and 2 out of 10 KK (2 down) isogenic vs. non-isogenic pairs (10% in total) differed in tracheal branching in normoxia (Figure 3M-N); and 0 out of 10 GD and 2 out of 10 KK (2 down) isogenic vs. non-isogenic pairs (10% in total) differed in tracheal branching in hypoxia (Figure 3O-P). A summary of these data is presented in Table 1. In conclusion, the genetic background affects to a different extent each complex trait.

Further analysis of the differences of isogenic vs. non-isogenic pairs in the various assays indicated that, although statistically significant, the phenotype might differ by a small or large percentage, as reflected by the fold-change difference between the genotypes. For example, for the total of 10 isogenic/non-isogenic pairs exhibiting differences in survival upon infection, the percentage of phenotypic difference ranges from 16% (13326/GD iso/non-iso pair) to 81.8% (101365/KK iso/non-iso pair). Since the biggest difference between an isogenic/non-isogenic pair might be considered more important, we assessed the intensity of the observed phenotypic differences by calculating the percentage of the pairs presented with more than 25% and more than 50% difference (Table 2). Interestingly, we noticed that the more sensitive the trait to the genotype, the more the percent difference between isogenic/non-isogenic pairs (Figure 4A). For the six traits that we had performed the DGRP screening, we assessed the degree of correlation between the z-score range (indicating the phenotype range in wild-type inbred strains) and the percentage of isogenic/non-isogenic pairs that presented a phenotype in total, as well as the percentage of isogenic/non-isogenic pairs that exhibited more that 25% and 50% difference in phenotype (Figure 4B-D). We found a significant correlation (R^2=0.8053, p=0.05) between the z-score range of the screen and the fraction of isogenic/non-isogenic pairs presented with more than 25% phenotypic difference (Figure 4C) and this did not improve further when we implemented a 50% cutoff (R^2=0.7797, p=0.06) (Figure 4D). Therefore, the traits affected more by the genotype are presented with an increased number of phenotypically-different isogenic/non-isogenic pairs; and the fraction of those pairs with more than 25% difference correlates with the sensitivity of the trait to the genotype.

The genetic background affects Gal4 activity. To assess the effect of the genetic background on Gal4 activity per se, we introgressed two transgenes: *UAS-srcGFP* and *Dl-Gal4* in a subset of DGRP strains. *Dl-Gal4>GFP* allows the visualization of intestinal stem cells in the adult *Drosophila* intestine (Zeng et al., 2010). In addition, it provides a way to assess the generation of dysplastic stem cell clusters (Biteau et al., 2008) upon genetic perturbations (e.g. by driving expression of UAS lines impinging on mitosis and cell differentiation), during aging or bacterial challenge. After six backcrossing generations to each DGRP strain, the flies were isogenic in the DGRP background while heterozygous for the *UAS-srcGFP* and *Dl-Gal4* transgenes. We assessed dysplastic cluster formation (clusters of more than 5 cell-large *Dl-Gal4>GFP* groups) in these strains by visualizing the GFP in dissected midguts. We found that there was variability in cluster formation in the different homozygous DGRP genotypes (Figure 5A). Then, to assess the effect of overexpression with Gal4, we used an RNAi line targeting *apc* (*UAS-apc^RNAi*), which, when induced promotes intestinal stem cell proliferation (Cordero et al., 2012). Although the phenotypic range was similar in the homozygous lines and the uninduced *UAS-apc^RNAi* heterozygotes, no correlation was observed between the genotypes (Figure 4D). Thus, making the DGRP strains heterozygous with the same strain (progeny contain 50% of the DGRP genome and 50% of the *UAS-apc^RNAi* genome), and reducing the genetic background by half did not produce predictable results. In addition, when the *UAS-apc^RNAi* progeny were brought to 29°C for induction of the transgene (3 days at 25°C followed by 2 days at 29°C with *Pseudomonas* feeding) and simultaneously infected with bacteria to boost mitosis, no correlation in cluster formation was observed between the uninduced and induced state of the same genotype (Figure 5B-D). Thus, the genetic background matters even when the *UAS-RNAi* is induced. In conclusion, the homozygous DGRP effect in cluster formation is not maintained upon crossing, and even when the Gal4-UAS is activated in these

flies and infection is added to enhance the phenotype, the effects were not predictable based on the heterozygous genotype.

A modulator of terminal branching lies on the 4th chromosome. A strong genetic background effect from the 4th chromosome was observed during assessment of branching in the isogenic/non-isogenic experiment. Specifically, since terminal branching needed to be assessed in larvae instead of adults, and because the KK VDRC lines had a deep red eye color as heterozygotes, initially, we kept the isogenized KK strains as heterozygotes and we kept selecting subsequent generations by eye color. Because the use of KK heterozygotes would prevent us from being certain for the genotype of the progeny larvae upon crossing to w^{1118}, we used a compound balancer stock to ensure that the genetic composition of the progeny would be the desired (Figure 6A). By pre-crossing the heterozygous isogenic KK lines to *yw/Y; T(2;3)/eyFLP2* males, we acquired males with balanced X, 2nd and 3rd chromosomes (but not 4th) and when these were crossed to w^{1118} females, the genotype of the progeny was selected to be isogenic KK, except for half of the cases of the 4th chromosome (which was derived from the compound balancer stock). Impressively, we noticed that when these KK [via T(2;3)] isogenic strains were crossed to our laboratory w^{1118}, their progeny differed from those derived from crosses of the non-isogenic strains to our laboratory w^{1118} in 40% of the cases (4 out of 10 iso/non-iso pairs; 1 down and 3 up in isogenic) in normoxia and in 100% of the cases (10 out of 10 iso/non-iso pairs; all 10 up in isogenic) in hypoxia (Figure 6B-C). Thus, the 4th chromosome of the compound balancer stock imposed a strong enhancing effect on branching.

DISCUSSION

Genetic background effects have been observed in various model organisms ranging from bacteria to mice (Chandler et al., 2013). The genetic background of wild-type inbred strains introduces natural variation in various fitness, physiology and disease phenotypes. In this study, the effects of the genetic background on different quantitative traits were measured. Not surprisingly, we found that all traits studied (female fecundity, survival upon pathogenic infection, intestinal mitosis upon pathogenic infection, defecation rate and fecal pH upon pathogenic infection and terminal branching in hypoxia) produced variable phenotypes when assessed in inbred homozygous DGRP strains (Figure 1). Based on the genomic information of the DGRP strains, we have identified genomic variants (SNPs, Insertions, Deletions) with a potential role on these phenotypes. Genetic characterization of these variants will be presented elsewhere.

In addition, we observed significant and variable genetic background effects when we assessed transgenic *UAS-RNAi* and *Gal4* constructs in different wild type genetic backgrounds. Specifically, although the VDRC *UAS-RNAi* lines were generated in an isogenic w^{1118} host strain and, whenever necessary, the balancers used were isogenic to the host strain (Dietzl et al., 2007), when the constructs were introgressed in our laboratory w^{1118} strain, they produced distinct phenotypes in various traits when compared to the original (we call them "non-isogenic") VDRC lines. This would mean that either the original lines changed over time or that the two w^{1118} strains (VDRC and ours) were not the same, because they are maintained in different environments. We did not observe major differences between the GD and KK insertions in terms of the number of strains affected by isogenization, although the first were generated in random genomic locations using P-element transformation, and thus cannot be

controlled for position effects. Interestingly, the genetic background effects were obvious in *UAS-RNAi* lines even in the absence of induction with a driver Gal4. Differences in Gal4-dependent phenotypes were also observed, in inbred strains as well as upon crossing to UAS and environmental challenge (Figure 5).

Moreover, we found a strong effect from the 4th chromosome in terminal branching, when assessing isogenic genotypes through balancing via a compound balancer strain and a genetic scheme ensuring that X, 2 and 3 chromosomes derived from the KK isogenic, but the 4th chromosome was 50% that of the KK isogenic and 50% that of the balancer stock (Figure 6A). Specifically, the compound balancer 4th chromosome increased branching. Since the 4th chromosome is small, largely heterochromatic and encompasses approximately 100 genes (Riddle and Elgin, 2018), it would be interesting to assess if any of these plays a role in terminal branching. Initial Gene Ontology (GO) analysis of the 4th chromosome genes, indicates that three of them (*ci*, an effector of the Hh pathway, *pan*, an effector of the Wg pathway, and *zyx*, a cytoskeletal protein) function in the trachea and *zyx* is involved in terminal branching morphology and tracheal air filling (Merabet et al., 2005; Renfranz et al., 2010).

Since the genetic toolkit of *Drosophila* continuously expands, we tend to use more and more transgenic lines (especially *UAS-RNAi*, that allows tissue-specific knockdown, when combined with *Gal4*) to understand the function of genes of interest (Mohr and Perrimon, 2012; Kaya-Çopur and Schnorrer, 2016). Depending on the phenotype we aim to study, we should be aware that the genetic background matters more or less. For example, traits like survival and fecundity are seriously affected by the genetic background, whereas tracheal branching is much less affected. So, what should one do to ensure that the phenotypes produced in their screen are not because of background effects? Considering the findings of this study, we propose the following guidelines:

1. Isogenization of all transgenic lines used in the same genetic background is ideal and will allow direct comparisons of strains, but it is time consuming. Some researchers would recommend to use more than one wild-type background for isogenization. This would be feasible and desirable, when your study focuses on a particular gene.
2. If isogenization is not an option (e.g. in cases of large-scale screening of *UAS-RNAi* lines), consider only big differences in your assays. We noticed that the intensity of the phenotypic difference (i.e. more that 25% difference) correlated with the variation of the phenotype (z-score range) when assessing wild-type flies.
3. Know your assay and the range of phenotypic differences expected. A small-scale pilot experiment will provide essential information of the phenotypic range. For example, terminal tracheal branching has a narrow phenotypic range and thus smaller phenotypic differences might be significant.
4. Repeat experiments independently three times to account for the impact of gene-to-environment variation.
5. Use more than one way to validate your results, e.g. when assessing RNAi phenotypes, multiple genetically-different *UAS-RNAi* lines targeting the same gene should be used.

MATERIALS AND METHODS

Drosophila maintenance: All strains and crosses were maintained on standard agar cornmeal fly food in a 12-hour light-dark cycle in a temperature-controlled incubator (Fitotron) at 25ºC (unless specified otherwise) with 65% humidity. For the DGRP screening for fecal spot number

and pH measurements, prior to infection, female mated flies were aged for 4 days at 25°C in bottles changed daily containing fly food supplemented with 50 ug/ml of the broad-range antibiotic Rifampicin, which does not kill PA14, but can eliminate most of the microorganisms present in the intestine of the flies. Rifampicin was also used in screening of the isogenic/non-isogenic pairs (Figure 3) for all assays except branching.

Drosophila Strains: The *Drosophila* Genetics Reference Panel (DGRP) collection of wild type inbred sequenced strains (Mackay et al., 2012; Huang et al., 2014) was used in all screens assessing physiological traits (fecundity, survival, intestinal regeneration, fecal motility, fecal spots, tracheal branching). The Vienna *Drosophila* Research Center (VDRC) UAS-RNAi lines used in this study are the following: 32863/GD and 106488/KK targeting *CG42318/app*, 45054/GD and 108473/KK targeting *CG8348/Dh44*, 19172/GD and 103958/KK targeting *CG17330/jhamt*, 13550/GD and 110349/KK targeting *CG13654*, 14641/GD and 101680/KK targeting *CG43902*, 14747/GD and 102995/KK targeting *CG33344/CCAP-R*, 37261/GD and 107921/KK targeting *CG32058/Ir67c*, 24275/GD and 110170/KK targeting *CG8282/Snx6*, 24746/GD and 103712/KK targeting *CG3326/Fign*, 13326/GD and 101365/KK targeting *CG43946/Glut1*. Other stocks used in this study were the following (source and/or stock center numbers in parentheses): *UAS-apc^{RNAi}* (VDRC, #1333), *UAS-srcGFP* (BDSC, #5432), *Dl-Gal4* (Zeng et al., 2010), *yw; ey-FLP2/T(2;3)CyO;TM6B,Tb,Hu* (BDSC, #8204).

Isogenization of transgenic lines: The VDRC lines, considered "non-isogenic", were backcrossed to the laboratory *w^{1118}* strain for at least 6 generations to produce the "isogenic" lines. For the GD lines, the transgenes were homozygosed following the backcrossing based on the intensity of their eye color. For the KK lines, due to the intensity of the red eye color, it was safer to maintain the transgenes in a heterozygous state (unless specified otherwise).

Fecundity measurement assays: Female flies were allowed to mate for 2 days at 25°C. Subsequently, males are removed from the cultures and females are left to lay eggs for 24 hours in a square bottle with 50 ml of food. Females are flipped to a new bottle for egg-laying every day for the next 3 days. Average fecundity is calculated as the number of progeny produced by the females in the 4 bottles divided by the number of females.

Pseudomonas aeruginosa infection: A 3 ml overnight culture of *Pseudomonas aeruginosa* PA14 was diluted 1:100 in LB (Luria Bertani medium) to prepare an overday 3 ml culture. When the culture reached $OD_{600}=3$, it was used to prepare the infection mix (5 ml per vial: 3.5 ml ddH$_2$O, 1 ml 20% sucrose and 0.5 ml PA14 $OD_{600}=3$) or sucrose mix (5 ml per vial control: 4 ml ddH$_2$O and 1 ml 20% sucrose). 5 ml of the infection or control mix was used to soak a cotton ball in a narrow fly vial, which was subsequently plugged with a dry cotton ball. After 4-5 hours starvation in empty vials, the flies were put in the infection vials and incubated at 25°C with the cotton plug facing down.

For the fecal spot and pH measurements, flies were fed concentrated bacteria of $OD_{600}=50$ (when the overday culture reached $OD_{600}=2$, it was concentrated 25 times by centrifugation and resuspension of the bacterial pellet in 4% sucrose). Specifically, each feeding vial plugged with a cotton ball contained 5 ml agar gel (3% w/v agar in H$_2$O) on top of which a Whatmann disc with 200 ul of the infection mix was placed. 25 mated starved young female flies were allowed to feed for 15 hours at 25°C on infection mix (or 4% sucrose for control) in each vial and subsequently their fecal spots were assessed for numbers and pH.

Survival assays: Young female flies were subjected to intestinal infection with the virulent *Pseudomonas aeruginosa* strain PA14. The percentage of dead flies was calculated daily as

the (number of dead flies per vial/total number of flies) x 100 until all flies are dead in each vial. LT50% (lethal time 50%), the time when 50% of the flies were dead was used as an indicator of survival for comparisons.

Intestinal regeneration measurement: Female adult flies (4-7 days old) were subjected to oral infection with the pathogenic strain of *Pseudomonas aeruginosa* PA14. Five days upon infection initiation, intestinal mitosis was measured by staining the intestines with the Rabbit-anti-pH3 antibody (Millipore) followed by secondary detection with Donkey-anti-Rabbit Alexa555 (Invitrogen), as described previously (Apidianakis et al., 2009). Mitotic cells were counted in whole midguts under the fluorescent Zeiss Axioscope A1. At least 10 midguts per genotype were used for mitotic index calculations per genotype.

Fecal number and pH measurements: Infected female flies were starved in empty vials for 5 hours and then placed in vials with cotton balls impregnated with 5 ml 4% sucrose with 0.5% w/v bromophenol blue (BPB) pH=7. BPB is a pH-sensitive dye, which is yellow in pH<3 and turns blue in pH>4.7. 50 flies were split in 3 vials with BPB and were allowed to feed for 5 hours (vials were placed in the incubator with the cotton plug facing down). 10 flies from each vial were moved to a petri dish containing a sterile cotton ball impregnated with 2.5 ml of the BPB solution and were left to poo for 20 hours. Then, the flies were removed from the plates and the total number of fecal spots was measured in the 3 plates. After counting the fecal spots, 10 spots from 2 separate locations of each plate (6 samples) were collected in an microtube in 25 ul ddH_2O pH=5.5 and their pH was measured with a fine tip pH meter. All experiments were performed at 25°C.

Terminal tracheal branching: Vials containing larvae were capped with net during development to allow quick gas exchange and were subjected to hypoxia (5% O_2) for 4 hours in a controlled chamber in the 25°C incubator. For control experiments, larvae were maintained in normoxia (21% O_2) in the 25°C incubator. Late L3 larvae were aligned on a microscope slide (dorsal side up) and heat-killed by placing the slide on a 70°C hot plate for 3-5 sec (Ghabrial and Krasnow, 2006). Subsequently, and in less than one hour upon heat-killing, the two terminal tracheal cells of the second tracheal metamere were assessed for the number of their terminal branches and photographed under bright-field on a Zeiss Axioscope A1.

Dysplastic cluster assays: Dysplastic clusters encompassing more than 5 cells were counted in adult *Drosophila* midguts of flies expressing GFP in intestinal stem cells via *Dl-Gal4>GFP*. *UAS-srcGFP* and *Dl-Gal4* were introgressed together in 22 DGRP lines. After six backcrossing generations, the transgenes were maintained in the DGRP backgrounds by selection of GFP expression and were kept in a heterozygous state. Baseline cluster formation was measured in 5-7 days old adult females maintained at 25°C. For *apc^RNAi* experiments, the DGRP (*Dl-Gal4>GFP*) isogenic strains were crossed to *UAS-apc^RNAi* at 25°C and the adult female progeny were aged for 3 days at 25°C before dysplastic cluster measurement for the uninduced state (the fact that no increase of cluster number was observed confirmed that *apc^RNAi* was not induced in these conditions). For *apc^RNAi* induction, the adult female progeny were aged for 3 days at 25°C before feeding with *P. aeruginosa* PA14 for 2 days at 29°C, where *apc^RNAi* is induced concomitantly with infection, which boosts the regenerative ability of the gut. Dysplastic clusters were measured under a fluorescent Axioscope A.1 microscope (Zeiss) in dissected midguts (Apidianakis et al, 2009) upon fixation and staining with rabbit- or chicken-anti-GFP (Invitrogen) and mouse-anti-Prospero (DSHB) followed by secondary detection with anti-rabbit or anti-chicken Alexa 488 and anti-mouse Alexa 555 (Invitrogen).

Statistical analysis: Normalized z-scores for each DGRP screen were calculated by the formula $z=(x-\mu)/\sigma$, where x is the mean observed value for each genotype normalized to the set average (each DGRP screen was performed in multiple sets of at least 10-20 genotypes), μ is the normalized population mean and σ is the standard deviation of the population. For hypoxia branching, z-score was calculated from the observed genotype means without normalization of the values, because the genotypes assessed per set were few.

For the isogenic/non-isogenic pair screening, initially each experiment was performed in duplicate and if both times the result had the same trend (at least once significant based on Student's t-test), the experiment was repeated for a third time. If 2 out of 3 times the result was statistically significant with the same trend, the difference was considered significant (noted with p-values in Figures 3 and 6, * $p\leq0.05$, ** $p\leq0.01$ and *** $p\leq0.001$).

For survival assays the p-value was calculated with the Kaplan Meier estimator (MedCalc statistical software). For all other assays, p-values were calculated with the Student's t-Test in Excel (Office package). For correlation graphs, trendlines and R^2 values were calculated in Excel and p-values were calculated by Pearson statistics.

FIGURE LEGENDS

Figure 1: Z-score analysis of the DGRP phenotypes of six complex traits. (**A**) Female fecundity, (**B**) Survival upon oral infection with _P. aeruginosa_, (**C**) Defecation rate upon oral infection with _P. aeruginosa_, (**D**) Fecal pH upon oral infection with _P. aeruginosa_, (**E**) Intestinal mitosis upon oral infection with _P. aeruginosa_, and (**F**) Terminal tracheal branching in hypoxia.

Figure 2: Pair-wise correlation analyses of the six DGRP screens. (**A**) Survival vs intestinal mitosis upon oral infection with _P. aeruginosa_. (**B**) Branching in hypoxia vs intestinal mitosis upon oral infection with _P. aeruginosa_. (**C**) Fecundity vs intestinal mitosis upon oral infection with _P. aeruginosa_. (**D**) Defecation rate upon oral infection with _P. aeruginosa_ vs intestinal mitosis upon oral infection with _P. aeruginosa_. (**E**) Fecal pH upon oral infection with _P. aeruginosa_ vs intestinal mitosis upon oral infection with _P. aeruginosa_. (**F**) Survival upon oral infection with _P. aeruginosa_ vs branching in hypoxia. (**G**) Fecundity vs branching in hypoxia. (**H**) Fecundity vs survival upon oral infection with _P. aeruginosa_. (**I**) Defecation rate upon oral infection with _P. aeruginosa_ vs survival upon oral infection with _P. aeruginosa_. (**J**) Defecation rate upon oral infection with _P. aeruginosa_ vs fecundity. (**K**) Defecation rate upon oral infection with _P. aeruginosa_ vs branching in hypoxia. (**L**) Fecal pH upon oral infection with _P. aeruginosa_ vs defecation rate upon oral infection with _P. aeruginosa_. (**M**) Fecal pH upon oral infection with _P. aeruginosa_ vs survival upon oral infection with _P. aeruginosa_. (**N**) Fecal pH upon oral infection with _P. aeruginosa_ vs branching in hypoxia. (**O**) Fecal pH upon oral infection with _P. aeruginosa_ vs fecundity. Trendlines are indicated in red and the R^2 values are shown in all graphs.

Figure 3: Isogenic vs non-isogenic comparisons of twenty _UAS-RNAi_ transgenic lines in the absence of Gal4 for eight traits. (**A-B**) Fecundity in GD and KK lines. (**C-D**) Survival upon oral infection with _P. aeruginosa_ in GD and KK lines. (**E-F**) Midgut mitosis in uninfected conditions in GD and KK lines. (**G-H**) Midgut mitosis upon oral infection with _P. aeruginosa_ in GD and KK lines. (**I-J**) Defecation rate upon oral infection with _P. aeruginosa_ in GD and KK lines. (**K-L**) Fecal pH upon oral infection with _P. aeruginosa_ in GD and KK lines. (**M-N**) Tracheal branching in normoxia in GD and KK lines. (**O-P**) Tracheal branching in hypoxia in GD and KK lines.

Significant pair-wise comparisons are indicated with brackets, whereby * p≤0.05, ** p≤0.01 and *** p≤0.001.

Figure 4: The phenotypic range correlates with the intensity of the difference between isogenic vs non-isogenic phenotypes. (**A**) Percentage of isogenic vs non-isogenic pairs presented with phenotypic differences and subcategories indicating the intensity of the observed differences, more than 25% difference between isogenic/non-isogenic and more than 50% difference between isogenic/non-isogenic. (**B**) Correlation of z-score range and the percentage of isogenic/non-isogenic pairs that differed. (**C**) Correlation of z-score range and the fraction of the total isogenic/non-isogenic pairs that differed by more than 25%. (**D**) Correlation of z-score range and the fraction of the total isogenic/non-isogenic pairs that differed by more than 50%. Trendlines are indicated in red and the R^2 values are shown in all graphs.

Figure 5: Gal4 is affected by the genetic background. (**A**) Dysplastic cluster formation in 22 DGRP lines carrying one copy of the *Dl-Gal4>GFP* transgenes. (**B**) Dysplastic cluster formation in the progeny of 22 DGRP lines carrying one copy of the *Dl-Gal4>GFP* transgenes crossed to *UAS-apc^{RNAi}* in uninduced conditions. (**C**) Dysplastic cluster formation in 22 DGRP lines carrying one copy of the *Dl-Gal4>GFP* transgenes crossed to *UAS-apc^{RNAi}* in induced conditions plus oral infection with *P. aeruginosa*. (**D**) Correlation of homozygous DGRP (*Dl-Gal4>GFP*) strains and heterozygous DGRP *Dl-Gal4>GFP/UAS-apc^{RNAi}*. (**E**) Correlation of heterozygous DGRP *Dl-Gal4>GFP/UAS-apc^{RNAi}* uninduced and induced with oral *P. aeruginosa* infection.

Figure 6: A modulator of terminal tracheal branching lies on the 4th chromosome. (**A**) Crossing scheme used to ensure that larvae assessed for terminal branching carried half of the KK isogenic genotype. Chromosomes X, 2 and 3, but not 4, were always derived from the KK isogenic parent. (**B**) Isogenic via T(2;3) vs non-isogenic comparisons for tracheal branching in normoxia. (**C**) Isogenic via T(2;3) vs non-isogenic comparisons for tracheal branching in hypoxia. Significant pair-wise comparisons are indicated with brackets, whereby * p≤0.05, ** p≤0.01 and *** p≤0.001.

ACKNOWLEDGMENTS

The authors would like to thank Maria Glykainou, Patapios Patapiou and Styliani Irakleous for their help with DGRP screening. We also thank the VDRC and BDSC for the *UAS-RNAi* and DGRP fly strains used in this study, respectively, the DSHB for antibodies, and Steven Hou for the *Dl-Gal4* stock. Work in the Pitsouli and Apidianakis laboratories was supported by the Marie Curie GIG FP7 - People and the Fondation Santé.

REFERENCES

Apidianakis Y, Pitsouli C, Perrimon N, Rahme L. (2009). Synergy between bacterial infection and genetic predisposition in intestinal dysplasia. *Proc Natl Acad Sci USA*. 106(49):20883-8.

Biteau B, Hochmuth CE, Jasper H. (2008). JNK activity in somatic stem cells causes loss of tissue homeostasis in the aging *Drosophila* gut. *Cell Stem Cell*. 3(4):442-55.

Brand AH, Perrimon N. (1993). Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. *Development*. 118(2):401-15.

Burnett C, Valentini S, Cabreiro F, Goss M, Somogyvári M, Piper MD, Hoddinott M, Sutphin GL, Leko V, McElwee JJ, Vazquez-Manrique RP, Orfila AM, Ackerman D, Au C, Vinti G, Riesen M, Howard K, Neri C, Bedalov A, Kaeberlein M, Soti C, Partridge L, Gems D. (2011). Absence of effects of Sir2 overexpression on lifespan in *C. elegans* and *Drosophila*. *Nature*. 477(7365):482-5.

Chandler CH, Chari S, Dworkin I. (2013). Does your gene need a background check? How genetic background impacts the analysis of mutations, genes, and evolution. *Trends Genet*. 29(6):358-66.

Chari S, Dworkin I. (2013). The conditional nature of genetic interactions: the consequences of wild-type backgrounds on mutational interactions in a genome-wide modifier screen. *PLoS Genet.* 9(8):e1003661.

Cordero JB, Stefanatos RK, Myant K, Vidal M, Sansom OJ. (2012). Non-autonomous crosstalk between the Jak/Stat and Egfr pathways mediates Apc1-driven intestinal stem cell hyperplasia in the *Drosophila* adult midgut. *Development*. 139(24):4524-35.

Dembeck LM, Huang W, Magwire MM, Lawrence F, Lyman RF, Mackay TF. (2015). Genetic Architecture of Abdominal Pigmentation in *Drosophila melanogaster*. *PLoS Genet.* 11(5):e1005163.

Dietzl G, Chen D, Schnorrer F, Su KC, Barinova Y, Fellner M, Gasser B, Kinsey K, Oppel S, Scheiblauer S, Couto A, Marra V, Keleman K, Dickson BJ. (2007). A genome-wide transgenic RNAi library for conditional gene inactivation in *Drosophila*. *Nature*. 448(7150):151-6.

Durham MF, Magwire MM, Stone EA, Leips J. (2014). Genome-wide analysis in *Drosophila* reveals age-specific effects of SNPs on fitness traits. *Nat Commun.* 5:4338.

Dworkin I, Kennerly E, Tack D, Hutchinson J, Brown J, Mahaffey J, Gibson G. (2009). Genomic consequences of background effects on scalloped mutant expressivity in the wing of *Drosophila melanogaster*. *Genetics*. 181(3):1065-76.

Ghabrial AS, Krasnow MA. (2006). Social interactions among epithelial cells during tracheal branching morphogenesis. *Nature*. 441(7094):746-9.

Harbison ST, McCoy LJ, Mackay TF. (2013). Genome-wide association study of sleep in *Drosophila melanogaster. BMC Genomics.* 14:281.

Hospital F. (2005). Selection in backcross programmes. *Philos Trans R Soc Lond B Biol Sci*. 360(1459):1503-11.

Huang W, Massouras A, Inoue Y, Peiffer J, Ràmia M, Tarone AM, Turlapati L, Zichner T, Zhu D, Lyman RF, Magwire MM, Blankenburg K, Carbone MA, Chang K, Ellis LL, Fernandez S, Han Y, Highnam G, Hjelmen CE, Jack JR, Javaid M, Jayaseelan J, Kalra D, Lee S, Lewis L, Munidasa M,

Ongeri F, Patel S, Perales L, Perez A, Pu L, Rollmann SM, Ruth R, Saada N, Warner C, Williams A, Wu YQ, Yamamoto A, Zhang Y, Zhu Y, Anholt RR, Korbel JO, Mittelman D, Muzny DM, Gibbs RA, Barbadilla A, Johnston JS, Stone EA, Richards S, Deplancke B, Mackay TF. (2014). Natural variation in genome architecture among 205 *Drosophila melanogaster* Genetic Reference Panel lines. *Genome Res.* 24(7):1193-208.

Ivanov DK, Escott-Price V, Ziehm M, Magwire MM, Mackay TF, Partridge L, Thornton JM. (2015). Longevity GWAS Using the *Drosophila* Genetic Reference Panel. *J Gerontol A Biol Sci Med Sci.* 70(12):1470-8.

Kaya-Çopur A, Schnorrer F. (2016). A Guide to Genome-Wide In Vivo RNAi Applications in *Drosophila*. *Methods Mol Biol.* 1478:117-143.

Kitzmann P, Schwirz J, Schmitt-Engel C, Bucher G. (2013). RNAi phenotypes are influenced by the genetic background of the injected strain. *BMC Genomics.* 14:5.

Kooke R, Wijnker E, Keurentjes JJ. (2012). Backcross populations and near isogenic lines. Methods Mol Biol. 871:3-16.

Mackay TF. (2010). Mutations and quantitative genetic variation: lessons from *Drosophila*. *Philos Trans R Soc Lond B Biol Sci.* 365(1544):1229-39.

Mackay TF, Richards S, Stone EA, Barbadilla A, Ayroles JF, Zhu D, Casillas S, Han Y, Magwire MM, Cridland JM, Richardson MF, Anholt RR, Barrón M, Bess C, Blankenburg KP, Carbone MA, Castellano D, Chaboub L, Duncan L, Harris Z, Javaid M, Jayaseelan JC, Jhangiani SN, Jordan KW, Lara F, Lawrence F, Lee SL, Librado P, Linheiro RS, Lyman RF, Mackey AJ, Munidasa M, Muzny DM, Nazareth L, Newsham I, Perales L, Pu LL, Qu C, Ràmia M, Reid JG, Rollmann SM, Rozas J, Saada N, Turlapati L, Worley KC, Wu YQ, Yamamoto A, Zhu Y, Bergman CM, Thornton KR, Mittelman D, Gibbs RA. (2012) The *Drosophila melanogaster* Genetic Reference Panel. *Nature*. 482(7384):173-8.

Markstein M, Pitsouli C, Villalta C, Celniker SE, Perrimon N. (2008). Exploiting position effects and the gypsy retrovirus insulator to engineer precisely expressed transgenes. *Nat Genet.* 40(4):476-83.

Merabet S, Hombria JC, Hu N, Pradel J, Graba Y. (2005). Hox-controlled reorganisation of intrasegmental patterning cues underlies *Drosophila* posterior spiracle organogenesis. *Development*. 132(13):3093-102.

Mohr SE, Perrimon N. (2012). RNAi screening: new approaches, understandings, and organisms. *Wiley Interdiscip Rev RNA*. 3(2):145-58.

Renfranz PJ, Blankman E, Beckerle MC. (2010). The cytoskeletal regulator zyxin is required for viability in *Drosophila melanogaster*. *Anat Rec (Hoboken)*. 293(9):1455-69.

Riddle NC, Elgin SCR. (2018). The *Drosophila* Dot Chromosome: Where Genes Flourish Amidst Repeats. *Genetics*. 210(3):757-772.

Rogina B, Reenan RA, Nilsen SP, Helfand SL. (2000). Extended life-span conferred by cotransporter gene mutations in *Drosophila*. *Science*. 290(5499):2137-40.

Rogina B, Helfand SL. (2004). Sir2 mediates longevity in the fly through a pathway related to calorie restriction. *Proc Natl Acad Sci USA.* 101(45):15998-6003.

Spradling AC, Rubin GM. (1982). Transposition of cloned P elements into *Drosophila* germ line chromosomes. *Science*. 218(4570):341-7.

Sunaga S, Akiyama N, Miyagi R, Takahashi A. (2016). Factors underlying natural variation in body pigmentation of *Drosophila melanogaster*. *Genes Genet Syst.* 91(2):127-137.

Toivonen JM, Walker GA, Martinez-Diaz P, Bjedov I, Driege Y, Jacobs HT, Gems D, Partridge L. (2007). No influence of Indy on lifespan in *Drosophila* after correction for genetic and cytoplasmic background effects. *PLoS Genet*. 3(6):e95.

Viswanathan M, Guarente L. (2011). Regulation of *Caenorhabditis elegans* lifespan by *sir-2.1* transgenes. *Nature*. 477(7365):E1-2.

Zeng X, Chauhan C, Hou SX. (2010) Characterization of midgut stem cell- and enteroblast-specific Gal4 lines in *Drosophila*. *Genesis*. 48(10):607-11. doi: 10.1002/dvg.20661.

	Isogenic vs non-isogenic pairs with difference (20 total) (% of total)	GD pairs (10 total) / KK pairs (10 total)	Up in GD isogenic / Down in GD isogenic / Up in KK isogenic / Down in KK isogenic
Fecundity	8 (40%)	4	2 (Up in GD isogenic)
			2 (Down in GD isogenic)
		4	2 (Up in KK isogenic)
			2 (Down in KK isogenic)
Survival upon oral infection	10 (50%)	5	1 (Up in GD isogenic)
			4 (Down in GD isogenic)
		5	1 (Up in KK isogenic)
			4 (Down in KK isogenic)
Midgut mitosis in baseline conditions	5 (25%)	2	1 (Up in GD isogenic)
			1 (Down in GD isogenic)
		3	0 (Up in KK isogenic)
			3 (Down in KK isogenic)
Midgut mitosis upon oral infection	5 (25%)	4	1 (Up in GD isogenic)
			3 (Down in GD isogenic)
		1	0 (Up in KK isogenic)
			1 (Down in KK isogenic)
Defecation rate upon oral infection	4 (20%)	2	0 (Up in GD isogenic)
			2 (Down in GD isogenic)
		2	1 (Up in KK isogenic)
			1 (Down in KK isogenic)
Fecal pH upon oral infection	4 (20%)	2	2 (Up in GD isogenic)
			0 (Down in GD isogenic)
		2	0 (Up in KK isogenic)
			2 (Down in KK isogenic)
Terminal branching in normoxia	2 (10%)	0	0 (Up in GD isogenic)
			0 (Down in GD isogenic)
		2	0 (Up in KK isogenic)
			2 (Down in KK isogenic)
Terminal branching in hypoxia	2 (20%)	0	0 (Up in GD isogenic)
			0 (Down in GD isogenic)
		2	0 (Up in KK isogenic)
			2 (Down in KK isogenic)

Table 1: Isogenic and non-isogenic GD and KK transgenic *UAS-RNAi* lines present variable differences in various phenotypic traits in the absence of Gal4. The last column compares the isogenic and non-isogenic strains with regards to the direction of the phenotype (up: increased, and down: decreased in the isogenic when compared to the non-isogenic).

	Isogenic vs non-isogenic pairs with difference (20 total) (% of total)	Isogenic vs non-isogenic pairs with >25% difference (% of total)	Isogenic vs non-isogenic pairs with >50% difference (% of total)
Fecundity	8 (40%)	7 (35%)	5 (25%)
Survival upon oral infection	10 (50%)	7 (35%)	2 (10%)
Midgut mitosis in baseline conditions	5 (25%)	5 (25%)	5 (25%)
Midgut mitosis upon oral infection	5 (25%)	5 (25%)	2 (10%)
Defecation rate upon oral infection	4 (20%)	1 (5%)	0 (0%)
Fecal pH upon oral infection	4 (20%)	0 (0%)	0 (0%)
Terminal branching in normoxia	2 (10%)	0 (0%)	0 (0%)
Terminal branching in hypoxia	2 (20%)	0 (0%)	0 (0%)

Table 2: The intensity of the phenotypic differences in comparisons of isogenic and non-isogenic GD and KK transgenic *UAS-RNAi* lines varies in the absence of Gal4.

Figure 1

Figure 2

Figure 3 (part 1)

Figure 3 (part 2)

Figure 4

Figure 5

Figure 6