Title: Development of oocytes in a post-spawning Japanese eel *Anguilla japonica*

Temminck & Schlegel in captivity

Running title: Developing oocytes in a post-spawning eel

Authors: Satoru Tanaka[1], Akihiro Okamura[1], Naomi Mikawa[1], Yoshiaki Yamada[1],

Noriyuki Horie[1], Tomoko Utoh[1], Huan Zhang[1,2], Hideo P. Oka[1,3] & Katsumi

Tsukamoto[1,4]

Affiliations:

[1]IRAGO Institute Co., Ltd., Tahara, Aichi 441-3605, Japan

[2]Present adress: Department of Marine Sciences, University of Connecticut, CT 06340,

USA

[3]Present address: Hamamatsu, Shizuoka 431-0102, Japan

[4] Department of Aquatic Bioscience, Graduate School of Agricultural and Life Sciences,

The University of Tokyo, 1-1-1 Yayoi, Bunkyo-ku, Tokyo, 113-8657 Japan

Correspondence: A. Okamura, IRAGO Institute Co., Ltd., Tahara, Aichi 441-3605,

Japan; e-mail: aokamura@irago.co.jp

Abstract

It is generally believed that parent freshwater eels (*Anguilla* sp.) die soon after spawning

on the assumption that eels are a semelparous (or monocyclic) fish (spawn once at the

last stage of life) like Pacific salmonids.   However, we observed for the first time a

post-spawning female *Anguilla japonica* again possessed developing oocytes reaching

the final maturation stage in captivity five months after the last spawning even without

hormonal treatment.   Here we describe information on this female about its biological

characteristics including gonadal histology and endocrine profiles. The data suggest that

lowering water temperature for a period of time is one of the important factors

influencing spontaneous gonadal development in this specimen. We also discuss the

possibility of induced multiple spawning of this species in captivity.

Key words: semelparity, anguillid eel, artificial maturation, spontaneous maturation,

water temperature

Freshwater eels (*Anguilla* sp.) are a food resource of worldwide importance, but they

are becoming globally endangered, especially in three temperate species European eel

Anguilla anguilla L., American eel *A. rostrata* (Lesueur, 1817) and *A. japonica*

Temminck & Schlegel (Casselman 2003; Dekker 2009; Tsukamoto, Aoyama & Miller

2009).   To conserve these eels, techniques for artificial maturation of parent eels by

hormonal administration and production of glass eels as seedlings for aquaculture

industries have been intensively studied over the past four decades (Yamamoto &

Yamauchi 1974; Tanaka 2003; Pedersen 2003; Oliveira & Hable 2010).

38   Usually, artificially matured parent eels by hormonal treatments are ordinarily

discarded after spawning on the assumption that eels are a semelparous (or

monocyclic) fish (spawn once at the last stage of life) like Pacific salmonids

(*Oncorhynchus* sp.) (Wallace & Selman 1981; Murua & Saborido-Rey 2003).

However, we observed for the first time a post-spawning female *A. japonica* again

possessed developing oocytes reaching the final maturation stage in captivity five

months after the last spawning even without hormonal treatment.   Here we describe

information on this female about its biological characteristics including gonadal

histology and endocrine profiles, and we discuss the possibility of induced multiple

spawning of this species in captivity.

48   A female silver eel of 67 cm in total length (TL) and 445 g in body weight

(BW) was captured by a set net in Mikawa Bay (34°N, 137°E) on 19 November 1996

(day 0).   Estimated age by the number of the otolith annuli was eight.   After

acclimatization to an indoor tank (10,000 l) of sea water (34.5 practical salinity unit

(PSU)) maintained at 18-20 °C for 2 weeks (day 13), the female was artificially

induced to maturation by repeated injection of salmon pituitary extract (SPE, 20 mg kg

BW[-1], once a week) for 6 weeks (Ohta, Kagawa, Tanaka, Okuzawa, Iinuma & Hirose

1997; Kagawa 2003) (Fig. 1).   Some males (200–300 g BW), which were purchased

from a commercial dealer, were also injected repeatedly with human chorionic

gonadotropin (Sankyo, Tokyo, Japan) at a dose of 200 IU per fish once a week at 20 °C

until full maturation (Ohta et al. 1997). Then the female was finally injected with

luteinizing hormone-releasing hormone (LH-RH, 1 mg kg BW^{-1}) (Satoh, Yamamori &

Hibiya 1992), and set in the 10,000-L tank together with some matured males (day 58).

Two days later (day 60), the female spontaneously released approximately 140,000

eggs based on calculation by the subtraction of BW after spawning (406 g) from that

before spawning (493 g).   Fertilization rate were about 38.9% and the diameters of

fertilized eggs was 1.55-1.65 mm, similar to *A. japonica* fertilized eggs collected at the

southern part of the Mariana Ridge (1.49–1.71 mm; n = 12) (Yoshinaga, Miller,

Yokouchi, Otake, Kimura, Aoyama, Watanabe, Shinoda, Oya, Miyazaki, Zenimoto,

Sudo, Takahashi, Ahn, Manabe, Hagihara, Morioka, Itakura, Machida, Ban, Shiozaki,

Ai & Tsukamoto 2011).   About 40,000 larvae hatched and survived until 10 days after

hatching without feeding.   The number of eggs released by the female was

considerably smaller than the estimated fecundity of this species ranged from 5 to 10

million eggs for female silver eels (35.7 to 92.4 cm TL) captured at Mikawa Bay

(Matsui 1972), suggesting that almost all oocytes possibly remained to be spawned in

the abdomen.   However, the abdomen gradually shrunk, and the genital pore was

completely closed soon after spawning.

Two days after spawning (day 62), the female was moved to an outdoor tank

(2,000 l) of seawater (31.5 PSU) and kept for 134 days without diets (Fig. 1).   The

water temperature of the tank was once decreased to 10.7°C on day 122 and increased

gradually to 14.3 °C until day 196.   On day 196, we found that the body of the

specimen was markedly thinned compared to the initial body, but the abdomen was

somewhat swelling and pliable despite the long-term starvation.   Therefore, in

anticipation of second spawning, the female was again moved to the indoor tank of

water temperature at 22°C and kept for 22 days (Fig. 1).   However, spawning did not

occur, so we dissected the female after heavy anesthetization on day 218 to observe the

gonads macroscopically and microscopically according to the procedure described in

Yamamoto, Omori & Yamauchi (1974).   The BW of the specimen was decreased to

245 g, about 60% of the BW just after spawning, on day 218 (Fig. 1).   Although we

have no detailed data on during the starvation, there is the possibility that the female

once attained a weight of less than 245 g and the weight increased again as the result of

oocyte hydration.

The female had apparently advanced ovaries with gonad-somatic index (GSI =

100 GW BW^{-1}; GW: gonad weight) value of 15.4 compared with those of migrating

female eels caught at the coastal waters of Mikawa Bay (range: 0.2-4.3, n=286) (Utoh,

Mikawa, Okamura, Yamada, Akazawa, Tanaka, Horie & Oka 2004), whereas their GSI

values can reach 17.8-51.4 (n=20) at their spawning by artificial maturation (Okamura,

Yamada, Horie, Utoh, Mikawa, Tanaka & Tsukamoto 2008).   Part of this high GSI

value is probably due to a loss of body weight, however, the thickness of ovaries was

enlarged compared with those found in migrating females captured at coastal waters

(Utoh *et al*. 2004) and its colour was yellow/white (Fig. 2a) and contained healthy

oocytes (*ca*. 700 μm in mean diameter).   Histology of the ovaries showed that each

oocyte included a migrating germinal vesicle but there were no large lipid droplets

visible (Fig. 2b), corresponding to those of the early migratory nucleus stage with

oocyte diameters of 600-700 μm found in artificially matured Japanese eels

(Yamamoto *et al*. 1974).   However, the number and size of yolk globule in these

oocytes seemed to be somewhat smaller (Fig. 2b), compared with those often found in

artificially matured eels at the same stage (Adachi, Ijiri, Kazeto & Yamauchi 2003).

Some of degenerating oocytes were also observed (Fig. 2b).   Furthermore, the GSI

value of this female is similar to those of post-spawning female Japanese eels caught at

the spawning area in the western North Pacific (9.0-13.4, n=3)(Tsukamoto, Chow,

Otake, Kurogi, Mochioka, Miller, Aoyama, Kimura, Watanabe, Yoshinaga, Shinoda,

Kuroki, Oya, Watanabe, Hata, Ijiri. Kazeto. Nomura, & Tanaka 2011).   These

post-spawning eels had also relatively advanced ovaries including oocytes at the

mid-vitellogenic stage with oocyte diameters of 350-600 μm.

Estradiol-17β (E2) and vitellogenin levels in the serum were also measured by

the methods described elsewhere (Asahina, Kambegawa & Higashi 1995; Mikawa,

Utoh & Oka, 2001; Utoh, Horie, Mikawa, Okamura, Yamada, Akazawa, Tanaka & Oka

2005).   Serum E_2 levels detected in this female (0.069 ng ml^{-1}) was apparently lower

than those usually found in artificially matured eels at the migratory nucleus stages

(range: 9.4-12.3 ng ml^{-1}, n=75) (Matsubara, Lokman, Kazeto, Adachi & Yamauchi

2005).   This low E2 levels, however, may be not strange at the post vitellogenic

stage, because higher levels of E2 often detected in artificially matured eels at the final

maturation stage are considered to be a peculiar feature induced by SPE injections

(Chiba, Iwatsuki, Hayami, Hara & Yamauchi 1994; Ijiri, Kazeto, Takeda, Chiba,

Adachi & Yamauchi 1995).   Generally, serum E2 levels in other teleosts increase in

parallel with the progress of vitellogenesis and the decrease at the final maturation

stage (Fostier; Jalabert; Billard; Breton & Zohar 1983), suggesting that low E2 levels

found in the present female may reflect a normal endocrine profile which appears in

wild eels under natural conditions.

Meanwhile, vitellogenin, which is the precursor of yolk protein synthesized in

liver in response to E_2, was detected at 2.52 mg ml^{-1} in the serum, which was a level

included in the range found in wild silver eels at the early- to mid-vitellogenic stages

caught in Mikawa Bay (range: 0.1-6.5 mg ml^{-1}, n=144) (Mikawa *et al*. 2001), but was

apparently lower than those found in artificially matured female silver eels (2.5-10 mg

ml^{-1} n=20) (Sato, Kawazoe, Suzuki & Aida 2000).   This levels of vitellogenin found

in the present female are possibly the reminder of circulating vitellogenin, because it

should be no longer necessary at this stage.

136   These reproductive characteristics suggest that this female had been ready to

undergo the final maturation phase.   Then, why did the female possess developing

oocytes five months after the previous spawning?   One possibility is that developed

oocytes were retained in the female's abdomen just as it was for five months.   This

possibility is suggested by the calculation that the number of eggs spawned by the

female (140, 000 eggs) was considerably smaller than the usual fecundity of this

species (5 to 10 million eggs).   However, Changchun, Yujun, Zhengfeng, Yiqiang &

Yejin (1980) observed that remnant oocytes in post-spawning Japanese eels were

usually absorbed by at least two months after spawning in captivity.   Therefore, we

think that this possibility is low.   Another possibility is that these oocytes newly

developed after the previous spawning.   The external color of ovaries in the female on

day 218 (Fig. 1a) appeared to be fresh and healthy, which were considerably differed

from those with a lot of ecchymoses usually found in females just after spawning in

captivity (Mikawa, unpublished data), suggesting the de novo development of oocytes

in this female.   Because the serum levels of the salmon gonadotropin (GTH) injected

into eels rapidly decrease to considerable low levels by four days and undetectable

levels by 24 days after the last injection (Sato, Kawazoe, Suzuki & Aida 1996; 2000),

the influence of residual activity of SPE might be no longer available soon after

spawning.   If so, we suppose that the development of these oocytes had been

controlled by endogenous hormones.

Generally, it has been known that eels do not sexually mature in captivity

without exogenous hormonal assistance, because their endogenous hormonal systems

do not work under captive conditions, probably due to a lack of appropriate

environmental triggers for maturation (Adachi *et al.* 2003; Palstra, Ginneken & van

den Thillart 2009).   Perhaps, relatively lower water temperature for five months

experienced by the present female is likely one of important factors influencing its

gonadal development.   Tracking experiment of silver *A. anguilla* revealed that they

undergo diel vertical migration during the spawning migration between 300 and 500 m

in depth where temperatures were lower than 13°C (Tesch 1989).   Another tracking

study showed that silver *A. dieffenbachii* Gray occasionally dove to around 6°C

(Jellyman & Tsukamoto 2010).   A laboratory experiment also indicated that a variable

thermal regime that increased from lower (10 °C) to higher temperatures (14 and

17 °C) induced a faster gonadal development, higher gonadotropin β-subunits (fshβ

and lhβ) expression and higher serum E2 levels during vitellogenesis in European

female eels injected with carp pituitary (Pérez, Peñaranda, Dufour, Baloche, Palstra,

van den Thillart & Asturiano 2011).   In a marine eel, *Conger myriaster* (Brevoort),

gonadal development completes without hormonal treatment in captivity at 6°C and

they spawn several days after an abrupt temperature elevation to 10 °C (Utoh & Horie

2011).   Therefore, it is possible that relatively lower water temperatures (11-14°C) for

five months stimulated pituitary hormonal axis and induced spontaneous oocyte

development.   The abrupt elevation of water temperature from 14 to 22°C (Fig. 1),

however, might be unfavorable for the female reaching final maturation phase.   Satoh

et al. (1992) and Dou, Yamada, Okamura, Shinoda, Tanaka & Tsukamoto (2007)

showed that spontaneous spawning of artificial matured *A. japonica* is efficiently

induced by an elevation of water temperature from 18 or 20 to 22°C.　However, the

present female did not spawn after the temperature elevation from 14 to 22°C.　It is

possible that the timing or width of temperature elevation was not adequate for the

female in the present study.　These inadequate temperature conditions during the final

maturation phase were likely reflected in a relatively small number of yolk globules in

oocytes and some of degenerating oocytes found in this female (Fig. 2b).

Previously, Changchun *et al*. (1980) also observed histologically a number of

newly developed oocytes at the pre-vitellogenic stage in post-spawning Japanese eels

(n=23) two years after spawning and they did not die but ingested diets and regained

their BW after spawning.　Horie *et al*. (unpublished data) observed that a number of

post-spawning *A. japonica* females repeatedly spawned (at least 3 times) in captivity

two or three weeks following previous spawning events when they were continually

received hormonal administration.　These evidences suggest that eel ovary is possibly

polycyclic and the type of ovarian development of organization in this species might

be 'group-synchronous' rather than 'synchronous' (Wallace & Selman 1981; Murua &

Saborido-Rey 2003).

In conclusion, the post-spawning female Japanese eel possessed almost

matured oocytes five months after spawning, suggesting that captive eels have the

potential of multiple spawning.　In addition, lower water temperatures for a period of

time may be one of the important factors influencing spontaneous gonadal

development in this species, which may lead to establishing a technique to obtain

efficiently eggs or to induce naturally eel maturation without exogenous hormones.

References

References

Adachi, S., Ijiri, S., Kazeto, Y. & Yamauchi, K. (2003) Oogenesis in the Japanese Eel,

Anguilla japonica. In: *Eel Biology* (ed. by K. Aida, K. Tsukamoto & K.

Yamauchi), pp. 301-317. Springer-Verlag, Tokyo.

Asahina, K., Kambegawa , A. & Higashi, T. (1995) Development of a microtiter plate

enzyme-linked immunosorbent assay for17α, 20β-21-trihydroxy-4-pregnen-3-

one, a teleost gonadal steroid. *Fisheries science* **61**, 491–494.

Casselman, J. M. (2003) Dynamics of resources of American eel, *Anguilla rostrata*:

declining abundance in the 1999s. In: *Eel Biology* (ed. by K. Aida, K.

Tsukamoto & K. Yamauchi), pp. 255–274. Springer-Verlag, Tokyo.

Changchun, Z., Yujun, T., Zhengfeng, S., Yiqiang, W. & Yejin, Y. (1980) The

possibility of resuming the reproductive capacity in spawned eels.

Oceanologia et Limnologia Sinica **11**, 241-246.

Chiba, H., Iwatsuki, K., Hayami, K., Hara, A. & Yamauchi, K. (1994) Changes in

serum steroid hormones and vitellogenin levels in cultured female European

eels, *Anguilla anguilla* during artificially induced ovarian development.

Journal of the World Aquaculture Society **25**,553-560.

Dekker, W. (2009) A conceptual management framework for the restoration of the

declining European eel stock. *American Fisheries Society Symposium* **58**,

3-19.

Dou, S.Z., Yamada, Y., Okamura, A., Shinoda, A., Tanaka, S. & Tsukamoto, K. (2007)

Temperature influence on the spawning performance of artificially-matured

Japanese eel, *Anguilla japonica*, in captivity. *Environmental Biology of Fishes*.

82, 151–64.

Fostier, A., Jalabert, B., Billard, B., Breton, B. & Zohar, Y. (1983) The gonadal steroids,

In: *Fish Physiology, volume IX, part A* (ed. by W.S. Hoar, D.J. Randall & E.M.

Donaldson), pp. 277-372. Academic Press, New York.

Ijiri, S., Kazeto, Y., Takeda, N., Chiba, H., Adachi, S. & Yamauchi, K. (1995) Changes

in serum steroid hormones and steroidogenic ability of ovarian follicles during

artificial maturation of cultivated Japanese eel, *Anguilla japonica.*

Aquaculture **135**, 3–16.

Jellyman, D. & Tsukamoto, K. (2010) Vertical migrations may control maturation in

migrating female *Anguilla dieffenbachii. Marine Ecology Progress Series* **404**,

241–247

Kagawa, H. (2003) Artificial induction of oocyte maturation and ovulation. In: *Eel*

Biology (ed. by K. Aida, K. Tsukamoto & K. Yamauchi), pp. 401-414.

Springer-Verlag, Tokyo, Japan.

Matsubara, H., Lokman, P. M., Kazeto, Y., Adachi, S. & Yamauchi, K. (2005) Serum

steroid profiles in artificially maturing female Japanese eel, *Anguilla japonica.*

Aquaculture **243**, 393–402.

Matsui, I. (1972) *Unagigaku (Eel biology).* Kouseisha-Kouseikaku, Tokyo, Japan, 283

pp.

Mikawa, N., Utoh, T. & Oka, H. P. (2001) Annual change in plasma vitellogenin levels

and liver in the female Japanese eel, *Anguilla japonica* appeared in Mikawa

Bay. In: *Proceedings of the international symposium: Advances in eel biology*

(ed. by K. Aida, K. Tsukamoto & K. Yamauchi), pp.207-209. The University

of Tokyo, Tokyo.

Murua, H. & F. Saborido-Rey, F. (2003) Female reproductive strategies of marine fish

species of the North Atlantic. *Journal of Northwest Atlantic Fishery Science*

33, 23–31.

Ohta, H., Kagawa, H., Tanaka, H., Okuzawa, K., Iinuma, N. & Hirose, K. (1997)

Artificial induction of maturation and fertilization in the Japanese eel,

Anguilla japonica. Fish Physiology and Biochemistry **17**, 163-169.

Okamura, A., Yamada, Y., Horie, N., Utoh, T., Mikawa, N., Tanaka, S. & Tsukamoto K.

(2008) Effects of silvering state on induced maturation and spawning in wild

female Japanese eel *Anguilla japonica. Fisheries Science* **74**, 642–8.

Oliveira, K. & Hable, W. E. (2010) Artificial maturation, fertilization, and early

development of the American eel (*Anguilla rostrata*). *Canadian Journal of*

Zoology **88**, 1121-1128

Palstra, A., van Ginneken, V. & van den Thillart, G. (2009) Effects of swimming on

silvering and maturation of the European eel, *Anguilla anguilla* L. In:

Spawning Migration of the European Eel (ed. by G. van den Thillart, S.

Dufour & J.C. Rankin), pp. 229-251. Springer, New York.

Pedersen, B.H. (2003) Induced sexual maturation of the European eel *Anguilla*

anguilla and fertilisation of the eggs. *Aquaculture* **224**, 323-338.

Pérez, L., Peñaranda, D.S., Dufour, S., Baloche, S., Palstra, A.P., van den Thillart

G.E.E.J.M. & Asturiano J.F. (2011) Influence of temperature regime on

endocrine parameters and vitellogenesis during experimental maturation of

European eel (*Anguilla anguilla*) females. *General and Comparative*

Endocrinology **174**:51–9.

Sato, N., Kawazoe, I., Suzuki, Y. & Aida K. (1996) Use of an emulsion prepared with

lipophilized gelatin for the induction of ovarian maturation in the Japanese eel

Anguilla japonica. Fisheries science **62**, 806-814.

Sato, N., Kawazoe, I., Suzuki, Y. & Aida, K. (2000) Adequate interval of sGTH

276  administration to induce vitellogenesis, and induction of ovulation by

277  simultaneous administration of sGTH and 17a-hydroxyprogesterone in the

278  Japanese eel. *Fisheries science* **66**, 644-654.

279 Satoh, H., Yamamori, K. & Hibiya, T. (1992) Induced spawning of the Japanese eel.

280  *Nippon Suisan Gakkaishi* **58**, 825-832.

281 Tanaka, H. (2003) Techniques for larval rearing. In: *Eel biology* (ed. by K. Aida, K.

282  Tsukamoto & K. Yamauchi), pp. 427-434. Springer-Verlag, Tokyo.

283 Tesch, F. W. (1989) Changes in swimming depth and direction of silver eels (*Anguilla*

284  *anguilla*) from the continental shelf to the deep sea. *Aquatic Living Resources*

285  **2**, 9-20.

286 Tsukamoto K., Aoyama J. & Miller M. J. (2009) Present status of the Japanese eel:

287  resources and recent research. *American Fisheries Society Symposium* **58**,

288  21–35.

289 Tsukamoto, K., Chow, S., Otake, T., Kurogi, H., Mochioka, N., Miller, M.J., Aoyama,

290  J., Kimura, S., Watanabe, S., Yoshinaga, T., Shinoda, A., Kuroki, M., Oya, M.,

291  Watanabe, T., Hata, K., Ijiri. S., Kazeto. Y., Nomura, K. & Tanaka, H. (2011)

292  Oceanic spawning ecology of freshwater eels in the western North Pacific.

293  *Nature Communications* **2**, 179. doi:10.1038/ncomms1174

294 Utoh, T. & Horie, N. (2011) Studies on the technique of seed production of the

295  common Japanese conger *Conger myriaster*. *Nippon Suisan Gakkaishi* **77**,

296  775-778.

297 Utoh, T., Mikawa, N., Okamura, A., Yamada, Y., Akazawa, A., Tanaka, S. Horie, N.,

298  Akazawa, A & Oka, H. P. (2004) Ovarian morphology of the Japanese eel in

299  Mikawa Bay. *Journal of Fish Biology*. **64**, 502-513.

300 Utoh, T., Horie, N., Mikawa, N., Okamura, A., Yamada, Y., Akazawa, A., Tanaka, S. &

301    Oka, H. P. (2005). Annual changes in ovarian development and plasma

302    estradiol-17β level in reared female common Japanese conger, *Conger*

303    *myriaster. Fisheries Science* **71**, 38-47.

304 Wallace, R. A. & Selman, K. (1981) Cellular and dynamic aspects of oocyte growth in

305    teleosts. *American Zoologist* **21**, 325-343.

306 Yamamoto, K., Omori, M. & Yamauchi, K. (1974) Oogenesis of the Japanese eel.

307    *Bulletin of the Japanese Society for the Science of Fish* **40**, 9-15.

308 Yamamoto, K. & Yamauchi, K. (1974) Sexual maturation of Japanese eel and

309    production of eel larvae in the aquarium. *Nature* **251**, 220-222.

310 Yoshinaga, T., Miller, M.J., Yokouchi, K., Otake, T., Kimura, S., Aoyama, J., Watanabe,

311    S., Shinoda, A., Oya, M., Miyazaki, S., Zenimoto, K., Sudo, R., Takahashi, T.,

312    Ahn, H., Manabe, R., Hagihara S., Morioka, H., Itakura, H., Machida, M., Ban,

313    K., Shiozaki, M., Ai, B. & Tsukamoto K. (2011) Genetic identification and

314    morphology of naturally spawned eggs of the Japanese eel *Anguilla japonica*

315    collected in the western North Pacific. *Fisheries Science* **77**, 983–92.

Figure legends

Figure 1   Changes in body weight (dotted line) of the *Anguilla japonica* female,

which was once induced spawning and then reared for about 5 months, and water

temperature experienced by the female (solid line). Arrows indicate the timings of

hormonal injection.

Figure 2   Macroscopic image of ovaries (a) and histological image of oocytes (b) in

the *Anguilla japonica* female reared for about 5 months after the last spawning.   G,

gut; L, liver; O, ovary; S, swim bladder; U, urogenital pore; d, oocyte atresia; n

nucleus.

° C

FIG. 1.

FIG. 2.