Intraspecific variation promotes trait clustering and species coexistence through higher-

order interactions.

Gaurav Baruah[1] and Robert John[2]

[1]Department of Evolutionary Biology and Environmental studies, University of Zurich,

Winterthurerstrasse 30, 4055 Zurich

[2]Department of Biological Sciences, Indian Institute of Science Education and Research,

Kolkata, Mohanpur 741246, India.

Corresponding author: Gaurav Baruah

Email: piyushgkb@gmail.com

Keywords: higher-order interactions, intraspecific variation, eco-evolutionary dynamics,

species coexistence, trait clustering, stability

Abstract

Within species variation is widespread in nature. Ecological and evolutionary effects of such

variation have been suggested to be detrimental for species coexistence. Community level

studies of species coexistence have largely focused on competitive interactions that are

characteristically pairwise, although, species interactions could involve combinations of

higher-order. Mechanistic ecological models that included higher-order interactions have

indicated the stabilizing effect of such interactions. It is however unknown how evolutionary

dynamics could affect species coexistence in a community dictated by both pairwise and

higher-order interactions. Also unknown are the effects of individual variation on species

coexistence in a community where higher order interactions are pervasive. In this study, we

explore the effects of intraspecific variation on patterns of species coexistence in a

competitive community dictated by both pairwise and higher-order interactions. We found

that higher-order interactions greatly stabilize species coexistence across different levels of

strength in competition. Surprisingly, high intraspecific variation promoted species

coexistence, which was prominent at high levels of strength in competition. Further, species

coexistence promoted by higher levels of individual variation were also robust to external

environmental perturbation. In addition to that, species' traits tend to cluster together as

individual variation in the community increased. Our results indicated that individual

variation can significantly promote species coexistence by reducing trait divergence and

consequently attenuating the inhibitory effect of dominant species through higher-order

interactions.

Introduction

Explanations for multi-species coexistence in ecological communities have largely been

sought at the species level by emphasizing average differences among species driven by

competitive interactions or life history trade-offs (Clark 2010a; Violle et al. 2012; Kraft,

Godoy, and Levine 2015; Letten, Ke, and Fukami 2017; Valladares et al. 2015; Gravel,

Guichard, and Hochberg 2011; Wittmann and Fukami 2018). These differences among

species along multiple ecological dimensions could minimise niche overlap and allow for

long-term species coexistence (Clark et al. 2010; Barabás, J Michalska-Smith, and Allesina

2016; Barabas and D'Andrea 2016). However, species appear to compete for only a small

number of limiting resources giving rise to the paradox of large numbers of coexisting

species on apparently a small number of limiting resources (Hutchinson 1961; Letten et al.

2018; Shoresh, Hegreness, and Kishony 2008). It has also become apparent that many species

coexist despite little difference in demographic or resource-based niches, at least along the

few dimensions that have been measured (Condit et al. 2006). So, although there are strong

theoretical arguments that average differences among species can account for species

coexistence, adequate empirical support has rarely been found.

Classical competition models of coexistence consider interactions among species pairs and

such models require parameter trade-offs to stabilize communities or to limit the strength of

competition among coexisting species in accordance with the competitive exclusion principle

(Barabás and Meszéna 2009; Barabás, J Michalska-Smith, and Allesina 2016). The

implausibility of highly structured competitive relationships in species-rich communities has

prompted models of coexistence based on ecological equivalence rather than life historical

differences (Hubbell 2006; Rosindell, Hubbell, and Etienne 2011; Segura et al. 2011).

Theoretical studies with competition models further show that any stability achieved through

pairwise competitive interactions can be disrupted by random interactions among species

(Bairey, Kelsic, and Kishony 2016; Barabás, J Michalska-Smith, and Allesina 2016; Allesina

and Levine 2011). The number of coexisting species then declines inversely with the strength

of interactions among species pairs (Bairey, Kelsic, and Kishony 2016). Interaction strength

therefore places an upper bound on the numbers of coexisting species, implying that strong

pairwise competitive interactions alone cannot promote species coexistence in a large

community.

Interactions among species are not always constrained to species pairs and can involve

higher-order combinations (Wilson 1992; Mayfield and Stouffer 2017; Bairey, Kelsic, and

Kishony 2016; Grilli et al. 2017; Terry, Morris, and Bonsall 2017) , where interactions

between a species pair is modulated by one or more other species (Fig. 1). In an ecological

system where pairwise interactions structure communities, indirect or higher-order effects

may alter these interactions and restructure communities (Terhorst et al. 2018; Levine et al.

2017). For example, a species that is a superior competitor for a given resource can inhibit an

inferior competitor for the same resource, but a third species may modulate the strength of

this inhibition without affecting either of the two competitors directly (Bairey, Kelsic, and

Kishony 2016). Such attenuation of the pairwise inhibitory effect can be density-mediated or

trait-mediated, and can lead to qualitatively different community dynamics compared to pure

pairwise interactions. The importance of such higher-order interactions has been recognised

(Levine et al. 2017), but the singular focus of coexistence studies on average species level

differences has meant that few investigations have been undertaken.

A further consequence of the focus on testing differences in species-level averages is that

within-species or individual level variation has largely been ignored (Siefert 2012; Hart et al.

2016). Observations that variation within species exceeds the differences in species-level

averages have prompted much theoretical and empirical research (Hart et al. 2016; Barabás, J

Michalska-Smith, and Allesina 2016; Hausch, Vamosi, and Fox 2018; Barabas and D'Andrea

2016). Intraspecific variation can have both ecological and evolutionary effects on

competitive interactions, which ultimately determine patterns of species coexistence. For

example, intraspecific trait variation can hamper species coexistence by increasing

competitive ability, niche overlap and even-spacing among species (Barabas and D'Andrea

2016), or by altering competitive outcomes through non-linear averaging of performances

(Hart et al. 2016). There is equally compelling evidence that intraspecific variation promotes

species coexistence, mainly through disruption of interspecific competitive abilities and

obscuring the effect of strongly competitive individuals in a community (Clark 2010b;

Bolnick et al. 2011). Experimental work has shown that intraspecific variation although

allows a community to be resilient to invaders, creates the opportunity for competitive

exclusion among strong competitors (Hausch, Vamosi, and Fox 2018). These contrasting

findings indicate the need for further investigations, particularly given that high levels of

intraspecific trait variation within communities appears to be more a rule than an exception.

We notice that the importance of higher-order species interactions and of intraspecific

variation on species coexistence had been investigated separately. The effect of intraspecific

trait variation and eco-evolutionary dynamics on structuring large communities where both

pairwise and higher-order interactions dominate a community is unknown. Purely pairwise

interactions in a community lead to even trait spacing when intraspecific variation is high.

Consequently, due to high intraspecific variation, competitive exclusion of inferior species in

a large community becomes inevitable (Fig. 1). However, a community dominated by both

pairwise and higher-order interactions could lead to less even spacing of species in a trait axis

and might lead to trait clustering. This could be because with high intraspecific variation

present in the community, higher-order interactions could significantly alleviate and stabilize

the negative pairwise interactions that lead to distinct spacing in the first place.

Here in this study, we examine the importance of higher order interactions and intraspecific

variation in structuring species coexistence and trait patterning. We do this using a modified

Lotka-Volterra modelling approach, where dynamics of the whole community is mediated

both by pairwise competitive interactions as well as higher-order three-way interactions.

Specifically, we model a one-dimensional quantitative trait that contributes to the competitive

ability of species interacting in the community. We show that in the presence of higher-order

interactions, high intraspecific variation across different levels of strength in competition

leads to significantly greater numbers of species coexisting in a community than when

individual variation is low. We show analytically and with model simulations that

intraspecific variation not only contributes to species coexistence, but also stabilizes the

community to external perturbation. In addition, our analyses reveal that intraspecific

variation in a community where higher-order interactions dictates dynamics leads to stable

trait clustering. Our study links the recent ecological studies of higher-order interactions with

eco-evolutionary dynamics and intraspecific variation.

2. Methods and Models

2.1 Community model with pairwise interactions

In our community model, we consider species competing with each other in a one-

dimensional trait axis, where a species' competitive ability is determined by a one-

dimensional quantitative trait z. Individuals of a species vary along the competitive trait z of

interest such that the distribution of the primary trait z is normally distributed with mean u_i

for species i and variation given by σ_i^2. Under such conditions, the dynamics of a species i is

given by Lotka-Volterra equations as (Barabas and D'Andrea 2016):

$$\frac{dN_i}{dt} = N_i(t)\left(b_i(t) - \sum_{j}^{s} \alpha_{ij}(t)N_j(t) \right) \quad (1)$$

139   And the dynamics of the mean competitive trait u_i is given by:

$$\frac{du_i}{dt} = h_i^2 u_i(t)\left(\overline{b_i(t)} - \sum_{j}^{s} \beta_{ij}(t)N_j(t) \right) \quad (2)$$

Where $\alpha_{ij}(t)$ describes the pairwise competition coefficient of species i with species j at any

time t. This competition coefficient derives directly from Gaussian competition kernel (See

appendix 2). If the two species are similar to each other in terms of their average trait value u,

then competition between them is stronger than when they are farther apart in the trait axis;

h_i^2 is the heritability of species i, $b_i(t)$ describes the growth rate of the species i in the

absence of any competition which is determined by where they lie in the trait axis z; $\overline{b_i(t)}$

describes the growth of the trait and $\beta_{ij}(t)$ quantifies the evolutionary pressure on the trait z

of species *i* due to competition with the species *j* in the community (this has been derived in

Barabas et al, 2016).

2.2 Community model with higher-order interactions

The above equations 1 and 2, captures the eco-evolutionary dynamics of a multispecies

community where pairwise interactions dominate community dynamics. It is still plausible

that a community could exhibit higher-order interactions than just between pairs of species.

In extension to the above model, we include density-mediated three-way higher-order

interactions where density of a third species influences pairwise competitive interactions.

Under these circumstances, the equations become (see appendix 2):

$\frac{dN_i}{dt} = N_i(t)\left(b_i(t) - \sum_j^S \alpha_{ij}(t)N_j(t) - \sum_j^S \sum_k^S \epsilon_{ijk}(t)N_j(t)N_k(t)\right).$ (3)

And the dynamics of the competitive trait u_i is given by:

$\frac{du_i}{dt} = h_i^2 \sigma_i^2 \frac{\partial}{\partial u_i}\left(\frac{1}{N_i}\frac{dN_i}{dt}\right),$

$= h_i^2 \sigma_i^2 u_i(t)\left(\overline{b_i(t)} - \sum_j^S \beta_{ij}(t)N_j(t) - \sum_j^S \sum_k^S \gamma_{ijk}(t)N_j(t)N_k(t)\right),$ (4)

where $\epsilon_{ijk}(t)$ gives the 3-way interactions in the sense that strictly pairwise trait-based

interactions are also affected by the presence of a third species *k*; γ_{ijk} denotes 3-way

interactions affecting evolutionary dynamics of mean trait *u* for species *i*. Similar to the

pairwise Gaussian interaction kernel, the three way interaction remains Gaussian with a third

species *k* influencing the interaction between the two species *i* and *j* given as (see appendix

2):

$$\epsilon_{ijk}(t) = N_k(t) \frac{w}{\sqrt{2\sigma_i^2 + 2\sigma_j^2 + w^2}} e^{-\frac{\left(u_i(t) - u_j(t)\right)^2}{2\sigma_i^2 + 2\sigma_j^2 + w^2}},$$

And, $\gamma_{ijk}(t)$ can be written as (appendix 2):

$$\gamma_{ijk}(t) = - N_k(t) \frac{2w\left(u_i(t) - u_j(t)\right)}{\left(2\sigma_i^2 + 2\sigma_j^2 + w^2\right)^{\frac{3}{2}}} e^{-\frac{\left(u_i(t) - u_j(t)\right)^2}{2\sigma_i^2 + 2\sigma_j^2 + w^2}},$$

Where, $\epsilon_{ijk}(t)$ and $\gamma_{ijk}(t)$ are three-dimensional tensors of size (S x S x S), where S is the

total number of species in the community. σ_i^2 and σ_j^2 are the intraspecific trait variation for

species i and species j respectively; w^2 is the width of the competition kernel which is

Gaussian (see appendix 2); $u_i(t)$ is the average trait value of species i and $u_j(t)$ is the

average trait value for species j. Thereby, eco-evolutionary dynamics in this purely

competitive community is dominated not only by pairwise trait-based competition but also by

three way higher-order interactions. In such a case, eco-evolutionary dynamics might deviate

from dynamics dominated by purely pairwise competitive coefficients as in (Barabas and

D'Andrea 2016). For details of the formulation see appendix 1-2.

2.3 Species coexistence in higher-order competition models with and without intraspecific

variation

Using the three-way interactions community model (see section 2.2 above), we assess the

influence of intraspecific trait variation on species coexistence. We examine analytically and

compare species richness in this multispecies community model with and without

intraspecific variation. For mathematical simplicity, in this section, we assume that

intraspecific variation is same for all the species in the community such that $\sigma_i^2 = \sigma_j^2 = \sigma^2$.

Based on strictly pairwise and three-way interactions in a diverse community, Bairey et al.

(2016) derived an upped bound for species richness. Accordingly, a diverse multispecies

community with pairwise as well as three-way interactions will follow (appendix 3):

$S = \frac{1-\epsilon_{ijk}}{\alpha_{ij}}.$

Hence ratio of species richness with and without intraspecific variation (see appendix 3) will

follow:

$$\frac{S_{var}}{S} = \frac{\left(\frac{1-\epsilon_{ijk}}{\alpha_{ij}}\right)}{\frac{1-\epsilon'_{ijk}}{\alpha'_{ij}}} \quad , \tag{5}$$

Where $\epsilon'_{ijk}, \alpha'_{ij}$ are three way and pairwise interactions without intraspecific variation, *i.e.*,

$\sigma_i^2 = \sigma_j^2 = 0$ and S_{var} and S are species richness in the community with and without

intraspecific variation respectively. We analyse the results from simulations of our model

with this derived analytical solution of species richness, with and without intraspecific

variation (see Results).

2.4 Simulations of the community model with higher-order interactions

We assessed the effect of different levels of intraspecific trait variation on community

structure and species coexistence using data generated from simulations of our community

model. We simulated both trait dynamics and population dynamics resulting from equations

(3) and (4). Initial community size for the start of each simulation was 40. All the 40 species

were randomly given an initial trait value within -0.5 to 0.5 in the trait axis. Outside this trait

regime, fitness value of a species will be extreme and growth rate will be negative.

Effectively, this strict criterion qualitatively means that outside this trait boundary resource

acquisition by a species is too low to survive and have positive growth rate. We carried out

45 replicate simulations for each level of intraspecific variation. We also simultaneously

tested the influence of the width of the competition kernel, which signifies the strength of

pairwise interaction, using a full factorial design where all possible combinations of

intraspecific variation and strength in competition width were tested for their influence on

species coexistence. In all our simulations, heritability h_i^2 of the trait for all species was fixed

at 0.1.

We evolved our community for a maximum of 1×10^4 time points, but we concluded

each simulation when the community had reached a stable state. We assumed that the

community attained a stable state if the ratio of minimum value of the entropy of the

community given by, $- \sum N_i \log (N_i)$, at two different time points, 500 units apart ($\Delta t =$

500), remains bounded within 10^{-5}. This condition was checked when the community had

evolved for more than 5×10^3 time points. If this condition was not met, we kept the

simulation going for another 5×10^3 time points before checking for the same condition. This

condition was however met at almost every simulation indicating the tendency for

convergence toward stable species density values.

2.4.1 Levels of width of the competition kernel and intraspecific variation

The width of the competition kernel *w*, (see appendix 2) was varied from 0.2 through 0.45

with increments of 0.05. For each *w*, three different levels of intraspecific variation were

tested in a fully factorial manner (6 different *w* values $\times$ 3 different σ^2 values $\times$ 45

replicates). Specifically, for each *w*, intraspecific variation for each of the 40 species in the

community was randomly sampled from a uniform distribution with three different levels: a)

low variation: $\sigma^2 = [\, 0.0006, 0.003]$; b) intermediate variation: $\sigma^2 = [\, 0.003, 0.009]$; and c)

high variation: $\sigma^2 = [\, 0.01, 0.05]$ (See Table 1, for parameters used).

2.5 Trait clustering:

Theoretical models have suggested that species coexisting together tend to spread more

evenly along a trait axis than expected (Barabas and D'Andrea 2016; D'Andrea and Ostling

2016). However, empirical studies have shown that it is possible for species clusters to

emerge along a trait axis (Segura et al. 2011; Vergnon, van Nes, and Scheffer 2012). Here,

we use a quantitative metric to evaluate the effect of intraspecific variation on the patterning

of traits in the trait-axis. We measured trait similarity between species coexisting together by

measuring the coefficient of variation (CV) of adjacent trait means (D'Andrea and Ostling

2016). High values of CV would indicate clustering of trait means of species in the trait axis

while lower CV values would indicate even spacing of traits.

2.6 Stability and robustness measures of species coexistence

Stability of our community model with higher-order interactions was measured by calculating

the Jacobian at equilibrium. Specifically, the Jacobian of our dynamical system at a given

point is (see appendix 3):

$$J_{ij} = -\delta_{ij}N_i + (-1)N_i\alpha_{ij} + (-1)N_i \sum_k N_k(\gamma_{ijk} + \gamma_{ikj})$$

$$= N_i\left(-\delta_{ij} - \alpha_{ij} - \sum_k N_k(\gamma_{ijk} + \gamma_{ikj})\right)$$

where, δ_{ij} is the Kronecker delta. At equilibrium it is possible that all the species coexist, but

for the community to be locally stable, the eigenvalues of the Jacobian at that equilibrium

point must all have negative eigenvalues. Thereafter, we measured the average robustness of

the community by taking the geometric mean of the absolute values of the eigenvalues of the

Jacobian (May 1973) (see appendix 3). Average community robustness measures the mean

response of the community to environmental perturbation (Barabas and D'Andrea 2016).

Specifically, this quantity measures the average return times in response to environmental

perturbation for each of the species in the community. For each replicate simulation of each

level of intraspecific variation, we calculated the average community robustness as the

measure to evaluate how intraspecific variation affected robustness of species coexistence.

Here, high values of average community robustness indicate lower stability.

3. Results

3.1 Analytical solution for the three-way competition model with and without intraspecific variation

We found that communities with higher intraspecific variation resulted in greater numbers of coexisting species than with communities that had no intraspecific variation (Fig. 2). At low levels of intraspecific variation, the ratio of species richness with and without intraspecific variation was around 1. But as intraspecific variation increased, the ratio of $\frac{S_{var}}{S}$ also increased significantly, showing that variation within species led to greater numbers of coexisting species than without intraspecific variation.

3.2 Effect of intraspecific variation and strength in competition on species coexistence

We found that, with increases in intraspecific variation, the numbers of coexisting species increased. At low levels of competition *w*, the effect of intraspecific variation on species coexistence was minimal, particularly for *w* = 0.2 and *w* = 0.25. But as the intensity of competition increased, we observed intraspecific variation had a stabilizing effect on species coexistence. At high levels of competition *w*, high intraspecific variation allowed a greater number of species to coexist in the trait axis (Fig. 3, Fig. 4).

3.3 Trait clustering

We measured trait clustering by quantifying coefficient of variation in the trait axis around a species' neighbourhood. We found that with increased intraspecific variation, coefficient of variation increased, indicating that traits tend to cluster together. Particularly, this result was evident only at high levels of intraspecific variation across all intensities of competition (Fig. 5).

3.4 Robustness of species coexistence

With increases in intraspecific variation, average robustness of the community increased. The

community became robust to external perturbation with increasing intraspecific trait variation

when compared with a community where intraspecific variation was low (Fig. 6).

4. Discussion

The importance and the consequence of intraspecific variation in community ecology is

intensely debated (Clark et al. 2010; Clark 2010b; Violle et al. 2012), with contrasting

findings being reported. Some studies have found that ecological and evolutionary

consequences of individual variation tend to weaken species coexistence (Hart et al. 2016;

Barabas and D'Andrea 2016). The nature of competitive interactions however appears to be

critical in determining the role of intraspecific variation. In competition models, purely

pairwise interactions place upper bounds on the numbers of coexisting species, decreasing

with increases in intensity of interactions, but including higher-order interactions leads to

qualitatively different dynamics (Bairey, Kelsic, and Kishony 2016; Mayfield and Stouffer

2017; Grilli et al. 2017). We investigated how intraspecific variation influences coexistence

in communities with both pairwise and higher-order interactions and found strong evidence

for stabilizing effects of intraspecific variation for species coexistence.

The assumption that pairwise interactions between species are sufficient to describe

competition in a community is ubiquitous in coexistence theory (Levine et al. 2017). Strong

competition (e.g., for shared limiting resources) between pairs of species would drive species

apart in niche space, structure communities, and maintain diversity. However, there is little

evidence that the observed species-level differences in mean demographic rates or resource

use are sufficient to explain species coexistence. Species may of course differ along many

dimensions that are either unmeasured or unseen, and this may be evident in the high levels

of intraspecific variation that is generally found (Clark 2010). Consistent with strong

arguments from other studies that intraspecific variation contributes to maintaining diversity

(Clark 2010), we found a strong stabilizing effect of intraspecific variation in communities

structured by pairwise and higher order competitive interactions.

In mechanistic models of competition where the underlying biology is modelled explicitly,

higher-order interactions can emerge subsequently in the process (Abrams 1983). Where

higher-order interactions have been explicitly modelled in phenomenological ecological

models, they act as a stabilizing factor in maintaining species diversity (Bairey, Kelsic, and

Kishony 2016; Grilli et al. 2017). We modelled the evolution of a trait that dictates

competitive ability between species and introduced higher order competitive interactions

where pairwise interactions were modulated by the density of a third species. Consistent with

earlier studies on the role of higher-order interactions (Bairey, Kelsic, and Kishony 2016;

Wilson 1992; Grilli et al. 2017) we found that such interactions greatly stabilize the dynamics

of species in the community. Expectedly, purely pairwise interactions led to lower numbers

of coexisting species as the strength of pairwise competitive interactions increased (Bairey,

Kelsic, and Kishony 2016). When we introduced three-way interactions, the dynamics of the

community quickly reached a stable equilibrium (Fig. 2, Fig. 3, Fig. 4). A strong competitor

in the trait axis can significantly affect the growth of inferior competitor. This results in a

disproportionately higher abundance for the dominant competitor compared to the

competitively inferior species. However, our results suggested that with the introduction of

three-way interactions, this dominance of the competitively superior species is significantly

reduced due to the presence of the third similar species leading to proportionately similar

densities for all the three species (Fig.1, Fig. 2). Earlier studies have studied the impact of

higher-order interactions from an ecological perspective, where the evolutionary side of

things was largely ignored. Our eco-evolutionary model that included higher-order

interactions led to stable coexistence of all distinct phenotypes, particularly when strength in

competition was low. With increases in the strength of pairwise competition, higher heritable

individual variation in the phenotypes stabilized ecological dynamics and led to higher

number of species coexisting. Higher-order interactions that could emerge in species-rich

competitive systems have not been well explored in the context of species coexistence

(Saavedra et al. 2017). Although, empirical studies on quantifying higher-order interactions

in field systems is exceedingly difficult (Mayfield and Stouffer 2017), ignoring such

interactions would limit fundamental understanding of the mechanisms behind species

coexistence in complex communities.

Our results show that greater levels of intraspecific variation can lead to higher species

richness but this effect was more prominent when pairwise competition was strong (w >0.25)

(Fig. 3, Fig. 4). Earlier studies have indicated that the numbers of species coexisting in eco-

evolutionary models incorporating purely pairwise interactions are always less than the

number of species coexisting in the absence of evolutionary dynamics (Edwards et al. 2018).

With sufficient intraspecific variation, a species can evolve into an uninvasible phenotype

that can lead to significant increases in its density. Consequently, the species with uninvasible

phenotypes could easily displace other species in the community (Edwards et al. 2018;

Barabas and D'Andrea 2016). However, with the incorporation of three-way higher order

interactions, the increases in density of superior species with sufficient intraspecific variation

is significantly limited, leading to higher number of coexisting species. With purely pairwise

interactions, eco-evolutionary models with higher intraspecific trait variation would lead to

greater overlap in the trait axis and species would limit other species more than they limit

themselves. .Consequently, the number of species coexisting with high intraspecific variation

should decrease substantially. With just pairwise interactions pervasive in a community, the

optimal number of species that could coexist scales inversely to the strength of pairwise

competition (Bairey, Kelsic, and Kishony 2016). With three way interactions, however, the

optimal number of species that could coexist in a community scales as $\frac{1-\epsilon_{ijk}}{\alpha_{ij}}$, such that

incorporation of higher order three way interactions would lead to increases in the number of

species that could coexist (Bairey, Kelsic, and Kishony 2016) when compared to a

community with purely pairwise interactions. Further, with higher intraspecific variation, the

strength in the three-way interaction strength decreases substantially that causes a rise in the

number of species that could coexist (Fig. 4, see appendix 3).

Clustering of species in a trait axes has been documented in nature particularly in aquatic

beetles, and freshwater algae (Scheffer and van Nes 2006; Holling 1992; Vergnon, van Nes,

and Scheffer 2012). Trait variation within populations in a community is a widespread

phenomenon in nature, however the implication of such trait variation on patterning of traits

is still debated. Trait patterning varies widely, often conforming to even spacing or

sometimes displaying extensive overlap (Vergnon, van Nes, and Scheffer 2012; Siefert

2012). In our eco-evolutionary model, where competition between species is includes both

pairwise and three-way interactions, increases in trait variation led to significant of trait

clustering (Fig. 5). Lotka-Volterra models dominated by pairwise interactions generally

support the idea that species tend to distribute more evenly along a trait axis than expected by

neutral evolution for the given trait (Barabas and D'Andrea 2016; Barabás, Meszéna, and

Ostling 2012). This is mostly because of the underlying competition kernel. Usually the

competition kernel is formulated in a way that species with similar phenotypes compete more

than species with dissimilar phenotypes. In such a case, naturally, when two species, for

instance, are placed in a trait axis (see Fig. 1) in way that they are very similar, initially there

would be strong pairwise competitive interactions between the species. In the presence of

evolutionary dynamics, both species would displace themselves in order to minimize the

negative effect of competition on each other's fitness. This leads to less of a trait overlap and

more of a trait divergence. However, the scenario changes when higher order interactions

come into play, such that the trait divergence due to strong pairwise competition is stabilized,

and species evolve to have more overlap in the trait axis. In other words, with the addition of

three-way higher-order interactions, the even spacing is decreased because the third species

attenuates the inhibitory or the displacing effect of the dominant species in the pairwise

interaction community, thereby maintaining stable coexistence and more trait overlap (and

thus overhauling the 'limiting similarity principle') (Bairey, Kelsic, and Kishony 2016).

When high-intraspecific variation is introduced to such a community, this pattern of trait

clustering becomes more evident as species tend to converge in the trait axis (Tobias et al.

2014).

Our modelling results suggest that higher intraspecific variation leads to robust and

stable species coexistence (Fig. 6). This means, that with higher intraspecific trait variation,

communities become more stable to external environmental perturbation (Barabas and

D'Andrea 2016). Local stability of a community at eco-evolutionary equilibrium will be

guaranteed if the community matrix or the Jacobian has all negative eigenvalues. This

however is not guaranteed in all dynamical systems at equilibrium (Barabás, Meszéna, and

Ostling 2012). Our results suggest that's high intraspecific variation in a community

dominated by pairwise as well as higher-order interactions, are significantly more robust and

stable than communities with low intraspecific variation. This stabilizing effect of

intraspecific variation is due to the fact that traits of species evolved into locations in the trait

axis that was advantageous to average community robustness. Moreover, with high

intraspecific variation, species could quickly recover and evolve to new trait means after an

external perturbation. Consequently, high variation and evolutionary dynamics could greatly

stabilize community responses to external perturbation (Dakos et al. 2018). Similar studies

also reported the stabilizing effect of higher intraspecific variation on community robustness

(Barabas and D'Andrea 2016) . Further earlier studies have examined the effect of

intraspecific variation on robustness of species coexistence in a competitive community

dominated by pairwise interactions (Barabas and D'Andrea 2016; Barabás, J Michalska-

Smith, and Allesina 2016). These studies suggest that intraspecific variation however does

promote species coexistence in a community with purely pairwise interactions. With the

introduction of higher-order interactions in a community, the number of species that could

coexist in a large community increases substantially (Grilli et al. 2017), where diversity

scales differently with different order of interactions (Bairey, Kelsic, and Kishony 2016).

Incorporating three-way higher order interactions alongside pairwise interactions in a

competitive community, we showed that intraspecific trait variation could significantly

stabilize and promote species coexistence in a large community.

In conclusion, we showed that intraspecific variation could promote species coexistence in a

competitive community provided pairwise competitive interactions and three-way higher

order interactions contributed to the dynamics of the species and trait patterning. Our work

demonstrates the importance of within species variation in a classical competition framework

that focuses on species-level differences. We show that intraspecific variation promotes

species coexistence, particularly when competitive interactions include higher-order

interactions and are of higher intensity. Furthermore, high intraspecific variation not only

promotes stable and robust species coexistence but also leads to trait convergence, a non-

intuitive result that has been found in other studies. An important next step would thus be to

characterize higher order interactions as well as individual variation in relation to capturing

variation in fitness in a diverse species community.

Author contributions

GB conceived the study, developed the model framework and did the analyses with feedback

from RJC; GB and RJC both contributed equally to writing the manuscript.

References

Abrams, Peter A. 1983. "Arguments in Favor of Higher Order Interactions." *The American*

Naturalist 121 (6): 887–91. https://doi.org/10.1086/284111.

Allesina, Stefano, and Jonathan M. Levine. 2011. "A Competitive Network Theory of

Species Diversity." *Proceedings of the National Academy of Sciences* 108 (14): 5638–

5642.

Bairey, Eyal, Eric D. Kelsic, and Roy Kishony. 2016. "High-Order Species Interactions

Shape Ecosystem Diversity." *Nature Communications* 7 (August): 12285.

https://doi.org/10.1038/ncomms12285.

Barabas, Gyorgy, and Rafael D'Andrea. 2016. "The Effect of Intraspecific Variation and

Heritability on Community Pattern and Robustness." *Ecology Letters*. Blackwell

Publishing Ltd.

Barabás, György, Matthew J Michalska-Smith, and Stefano Allesina. 2016. "The Effect of

Intra- and Interspecific Competition on Coexistence in Multispecies Communities." *The*

American Naturalist 188 (1): E1–12. https://doi.org/10.1086/686901.

Barabás, György, and Géza Meszéna. 2009. "When the Exception Becomes the Rule: The

Disappearance of Limiting Similarity in the Lotka–Volterra Model." *Journal of*

Theoretical Biology 258 (1): 89–94. https://doi.org/10.1016/J.JTBI.2008.12.033.

Barabás, György, Géza Meszéna, and Annette Ostling. 2012. "Community Robustness and

Limiting Similarity in Periodic Environments." *Theoretical Ecology* 5 (2): 265–82.

https://doi.org/10.1007/s12080-011-0127-z.

Bolnick, Daniel I, Priyanga Amarasekare, Márcio S Araújo, Reinhard Bürger, Jonathan M

Levine, Mark Novak, Volker H W Rudolf, Sebastian J Schreiber, Mark C Urban, and

David A Vasseur. 2011. "Why Intraspecific Trait Variation Matters in Community

Ecology." *Trends in Ecology & Evolution* 26 (4): 183–92.

https://doi.org/10.1016/j.tree.2011.01.009.

Clark, James S., David Bell, Chengjin Chu, Benoit Courbaud, Michael Dietze, Michelle

Hersh, Janneke Hillerislambers, et al. 2010. "High-Dimensional Coexistence Based on

Individual Variation: A Synthesis of Evidence." *Ecological Monographs* 80 (4): 569–

608.

Clark, James S. 2010a. "Individuals and the Variation Needed for High Species Diversity in

Forest Trees." *Science (New York, N.Y.)* 327 (5969): 1129–32.

https://doi.org/10.1126/science.1183506.

———. 2010b. "Individuals and the Variation Needed for High Species Diversity in Forest

Trees." *Science* 327 (5969): 1129–32.

http://www.sciencemag.org/content/327/5969/1129.abstract.

Condit, Richard, Peter Ashton, Sarayudh Bunyavejchewin, H S Dattaraja, Stuart Davies,

Shameema Esufali, Corneille Ewango, et al. 2006. "The Importance of Demographic

Niches to Tree Diversity." *Science (New York, N.Y.)* 313 (5783): 98–101.

https://doi.org/10.1126/science.1124712.

D'Andrea, Rafael, and Annette Ostling. 2016. "Challenges in Linking Trait Patterns to Niche

Differentiation." *Oikos* 125 (10): 1369–85. https://doi.org/10.1111/oik.02979.

Dakos, Vasilis, Blake Matthews, Andrew Hendry, Jonathan Levine, Nicolas Loeuille, Jon

Norberg, Patrik Nosil, Marten Scheffer, and Luc De Meester. 2018. "Ecosystem Tipping

Points in an Evolving World." *BioRxiv*, October, 447227.

https://doi.org/10.1101/447227.

Edwards, Kyle F., Colin T. Kremer, Elizabeth T. Miller, Matthew M. Osmond, Elena

Litchman, and Christopher A. Klausmeier. 2018. "Evolutionarily Stable Communities:

A Framework for Understanding the Role of Trait Evolution in the Maintenance of

Diversity." Edited by Lutz Becks. *Ecology Letters*, September.

https://doi.org/10.1111/ele.13142.

Gravel, Dominique, Frédéric Guichard, and Michael E. Hochberg. 2011. "Species

Coexistence in a Variable World." *Ecology Letters*.

Grilli, Jacopo, György Barabás, Matthew J. Michalska-Smith, and Stefano Allesina. 2017.

"Higher-Order Interactions Stabilize Dynamics in Competitive Network Models."

Nature 548 (7666): 210. https://doi.org/10.1038/nature23273.

Hart, Simon P., Sebastian J. Schreiber, Jonathan M. Levine, and Tim Coulson. 2016. "How

Variation between Individuals Affects Species Coexistence." *Ecology Letters* 19 (8):

825–38.

Hausch, Stephen, Steven M. Vamosi, and Jeremy W. Fox. 2018. "Effects of Intraspecific

Phenotypic Variation on Species Coexistence." *Ecology* 99 (6): 1453–62.

https://doi.org/10.1002/ecy.2346.

Holling, C. S. 1992. "Cross-Scale Morphology, Geometry, and Dynamics of Ecosystems."

Ecological Monographs 62 (4): 447–502. https://doi.org/10.2307/2937313.

Hubbell, Stephen P. 2006. "Neutral Theory and the Evolution of Ecological Equivalence."

Ecology 87 (6): 1387–98. http://dx.doi.org/10.1890%2F0012-

9658%282006%2987%5B1387%3ANTATEO%5D2.0.CO%3B2.

Hutchinson, G . E . 1961. "The Paradox of the Plankton Author." *The American Naturalist* 95

(882): 137–45. https://doi.org/10.1086/282171.

Kraft, Nathan J B, Oscar Godoy, and Jonathan M Levine. 2015. "Plant Functional Traits and

the Multidimensional Nature of Species Coexistence." *Proceedings of the National*

Academy of Sciences of the United States of America 112 (3): 797–802.

https://doi.org/10.1073/pnas.1413650112.

Letten, Andrew D., Manpreet K. Dhami, Po-Ju Ke, and Tadashi Fukami. 2018. "Species

Coexistence through Simultaneous Fluctuation-Dependent Mechanisms." *Proceedings of the National Academy of Sciences* 115 (26): 6745–50. https://doi.org/10.1073/pnas.1801846115.

Letten, Andrew D., Po-Ju Ke, and Tadashi Fukami. 2017. "Linking Modern Coexistence Theory and Contemporary Niche Theory." *Ecological Monographs* 87 (2): 161–77. https://doi.org/10.1002/ecm.1242.

Levine, Jonathan M., Jordi Bascompte, Peter B. Adler, and Stefano Allesina. 2017. "Beyond Pairwise Mechanisms of Species Coexistence in Complex Communities." *Nature* 546 (7656): 56–64. https://doi.org/10.1038/nature22898.

May, Robert M. 1973. "Qualitative Stability in Model Ecosystems." *Ecology* 54 (3): 638–41. https://doi.org/10.2307/1935352.

Mayfield, Margaret M., and Daniel B. Stouffer. 2017. "Higher-Order Interactions Capture Unexplained Complexity in Diverse Communities." *Nature Ecology & Evolution* 1 (3): 0062. https://doi.org/10.1038/s41559-016-0062.

Rosindell, James, Stephen P. Hubbell, and Rampal S. Etienne. 2011. "The Unified Neutral Theory of Biodiversity and Biogeography at Age Ten." *Trends in Ecology and Evolution* 26 (7): 340–48. https://doi.org/10.1016/j.tree.2011.03.024.

Saavedra, Serguei, Rudolf P. Rohr, Jordi Bascompte, Oscar Godoy, Nathan J. B. Kraft, and Jonathan M. Levine. 2017. "A Structural Approach for Understanding Multispecies Coexistence." *Ecological Monographs* 87 (3): 470–86. https://doi.org/10.1002/ecm.1263.

Scheffer, Marten, and Egbert H. van Nes. 2006. "Self-Organized Similarity, the Evolutionary Emergence of Groups of Similar Species." *Proceedings of the National Academy of Sciences of the United States of America* 103 (16): 6230–35. https://doi.org/10.1073/pnas.0508024103.

Segura, Angel M, Danilo Calliari, Carla Kruk, Daniel Conde, Sylvia Bonilla, and Hugo Fort.

   2011. "Emergent Neutrality Drives Phytoplankton Species Coexistence." *Proceedings.*

   Biological Sciences 278 (1716): 2355–61. https://doi.org/10.1098/rspb.2010.2464.

Shoresh, Noam, Matthew Hegreness, and Roy Kishony. 2008. "Evolution Exacerbates the

   Paradox of the Plankton." *Proceedings of the National Academy of Sciences of the*

   United States of America 105 (34): 12365–69.

Siefert, Andrew. 2012. "Incorporating Intraspecific Variation in Tests of Trait-Based

   Community Assembly." *Oecologia* 170 (3): 767–75. https://doi.org/10.1007/s00442-

   012-2351-7.

Terhorst, Casey P, Peter C Zee, Katy D Heath, Thomas E Miller, Abigail I Pastore, Swati

   Patel, Sebastian J Schreiber, Michael J Wade, and Matthew R Walsh. n.d. "Evolution in

   a Community Context: Trait Responses to Multiple Species Interactions*." Accessed

   July 2, 2018. https://doi.org/10.1086/695835.

Terry, J. Christopher D., Rebecca J. Morris, and Michael B. Bonsall. 2017. "Trophic

   Interaction Modifications: An Empirical and Theoretical Framework." Edited by James

   Grover. *Ecology Letters* 20 (10): 1219–30. https://doi.org/10.1111/ele.12824.

Tobias, Joseph A., Charlie K. Cornwallis, Elizabeth P. Derryberry, Santiago Claramunt,

   Robb T. Brumfield, and Nathalie Seddon. 2014. "Species Coexistence and the Dynamics

   of Phenotypic Evolution in Adaptive Radiation." *Nature* 506 (7488): 359–63.

   https://doi.org/10.1038/nature12874.

Valladares, Fernando, Cristina C. Bastias, Oscar Godoy, Elena Granda, and Adrián Escudero.

   2015. "Species Coexistence in a Changing World." *Frontiers in Plant Science* 6

   (October): 866. https://doi.org/10.3389/fpls.2015.00866.

Vergnon, Remi, Egbert H. van Nes, and Marten Scheffer. 2012. "Emergent Neutrality Leads

   to Multimodal Species Abundance Distributions." *Nature Communications* 3 (1): 663.

https://doi.org/10.1038/ncomms1663.

Violle, Cyrille, Brian J. Enquist, Brian J. McGill, Lin Jiang, Cécile H. Albert, Catherine

Hulshof, Vincent Jung, and Julie Messier. 2012. "The Return of the Variance:

Intraspecific Variability in Community Ecology." *Trends in Ecology & Evolution* 27

(4): 244–52. https://doi.org/10.1016/J.TREE.2011.11.014.

Wilson, David Sloan. 1992. "Complex Interactions in Metacommunities, with Implications

for Biodiversity and Higher Levels of Selection." *Ecology* 73 (6): 1984–2000.

https://doi.org/10.2307/1941449.

Wittmann, Meike J., and Tadashi Fukami. 2018. "Eco-Evolutionary Buffering: Rapid

Evolution Facilitates Regional Species Coexistence despite Local Priority Effects." *The*

American Naturalist 191 (6): E000–E000. https://doi.org/10.1086/697187.

Table 1: List of parameters, variables and functions of the model and their respective

description and values.

Parameters/variables/functions	Description	Value
N_i	Density of species i	….
θ		0.5
b_i	Growth rate of species i	$\frac{1}{\sqrt{2\pi}\sigma_i}\left[\,\mathrm{erf}\left(\frac{\theta-u_i}{\sqrt{2}\sigma_i}\right)+\mathrm{erf}\left(\frac{\theta+u_i}{\sqrt{2}\sigma_i}\right)\right]$ Where erf is the error function
α_{ij}	Competition coefficient between	$\frac{w}{\sqrt{2\sigma_i^2+2\sigma_j^2+w^2}}\exp\left(-\frac{\left(u_i-u_j\right)^2}{2\sigma_i^2+2\sigma_j^2+w^2}\right)$

	species *i* and *j*	
ϵ_{ijk}	Three-way interaction : competition between species *i* and *j* is influenced by species *k* (a tensor of size SxSxS)	As given in equation
u_i	Mean trait value of species *i*	Can take values in the range from -0.5 to 0.5. Initial mean trait values are randomly assigned uniformly to species in the range [-0.5, 0.5]
$\bar{b}_\iota$		$\dfrac{1}{\sqrt{2\pi}\sigma_i}\left[\exp\left(\dfrac{-(\theta + u_i)^2}{2\sigma_i^2}\right) - \exp\left(\dfrac{-(\theta - u_i)^2}{2\sigma_i^2}\right)\right]$
β_{ij}	Evolutionary pressure by species *j* on species *i*'s mean trait value u_i	$\dfrac{2w(u_j - u_i)}{\sqrt{\left(2\sigma_i^2 + 2\sigma_j^2 + w^2\right)^3}}\exp\left(-\dfrac{\left(u_i - u_j\right)^2}{2\sigma_i^2 + 2\sigma_j^2 + w^2}\right)$
γ_{ijk}	Evolutionary pressure on	As given in equation

	trait value of species i due to three-way interactions, such that pairwise competition between two species i and j is modulated by the third species k, (a tensor of size SxSxS)	
σ_i^2, σ_j^2	Trait variance of species i, j;	Three levels- a) High variation [0.01,0.05] ; b) medium variation [0.003,0.009]; c) low variation [0.0006,0.003] Species trait variance are uniformly sampled for each replicate from the ranges given above
S	Total number of starting species in the	40

	community	
w	Width of the competition kernel signifying the strength in competition	Various- 0.2,0.25,0.3,0.35,0.4,0.45

List of figures:

Figure 1. An example image of how intraspecific trait variation in communities with (a) pairwise

competitive interactions (black arrows) and (b), (c) higher-order interactions (dashed-arrows) can

affect trait patterning and coexistence. (a) Three different species (Red, blue and orange) are spaced

along a trait axis with high variation. Interactions between the three species are inherently pairwise.

Starting with purely pairwise competitive communities' initial high intraspecific variation will

ultimately eventually lead (1) to competitive exclusion of the red species. In addition, the other two

remaining species (blue and orange) will space themselves far apart minimizing trait overlap and

leading to the emergence of what is called the 'limiting similarity' principle. However, with the

introduction of higher-order interactions, low levels of intraspecific variation (b) will also lead to (2)

species minimizing trait overlap but leading to all species coexisting. However, (c) high intraspecific

variation and with higher-order interactions will lead to (3) more trait overlap as well as coexistence

of all the three species.

Figure 2. Example dynamics of trait and species density over time for three levels of intraspecific

variation for a particular $w = 0.3$. (A) High intraspecific variation; (B) Medium intraspecific variation;

(c) low intraspecific variation. Different colors represent different species.

Figure 3. Effect of intraspecific trait variation on species richness. Boxplots denote the total number

of species that coexisted at the end of the simulations for different levels of competition denoted by *w*

levels. Grey colour indicates low individual variation; red colours represent medium intraspecific

variation; brown colour denotes high level of intraspecific variation.

Figure 4.   Effect of intraspecific trait variation on ratio of species richness with and without

intraspecific variation. Red color dots denotes the analytical prediction of the model as given by

equation (5). Black dots represent replicate model simulations for each level of intraspecific variance.

Note that there is a shift in the ratio of species richness with to without intraspecific variation (ITV)

once intraspecific trait variation increases above 0.01. This denotes that number of species coexisting

increases significantly after intraspecific variation increases above the threshold of 0.01.   This

simulation and analytical prediction were shown for $w = 0.4$.

Figure 5. Effect of intraspecific variation on trait convergence for different levels of strength in competition, *w*. Note that trait convergence increases substantially with increasing levels of intraspecific variation across different *w*. Grey colour indicates low individual variation; red colours represent medium intraspecific variation, brown colour denotes high level of intraspecific variation.

621   Figure 6. Effect of intraspecific variation on average community robustness for different levels of

622   competition strength, *w*. Note that lower value of average robustness signifies greater community

623   stability. With increases in intraspecific variation, average community robustness increased

624   significantly across different levels of strength in competition, *w*. This indicates that communities

625   where intraspecific variation is high, they are more stable to external perturbations.

626