Nucleosome positioning stability is a significant modulator of germline

mutation rate variation across the human genome

Cai Li[1,#] and Nicholas M. Luscombe[1,2,3]

[1] The Francis Crick Institute, 1 Midland Road, London, NW1 1AT, UK

[2] Okinawa Institute of Science & Technology Graduate University, Okinawa,
904-0495, Japan

[3] UCL Genetics Institute, University College London, Gower Street, London,
WC1E 6BT, UK

Correspondence: cai.li@crick.ac.uk

Abstract

Understanding the patterns and genesis of germline *de novo* mutations is important

for studying genome evolution and human diseases. Nucleosome organization is

suggested to be a contributing factor to mutation rate variation across the genome.

However, the small number of published *de novo* mutations and the low resolution of

earlier nucleosome maps limited our understanding of how nucleosome organization

affects germline mutation rates in the human genome. Here, we systematically

investigated the relationship between nucleosome organization and fine-scale

mutation rate variation by analyzing >300,000 *de novo* mutations from whole-

genome trio sequencing and high-resolution nucleosome maps in human. We found

that *de novo* mutation rates are elevated around strong, translationally stable

nucleosomes, a previously under-appreciated aspect. We confirmed this observation

having controlled for local sequence context and other potential confounding factors.

Analysis of the underlying mutational processes suggests that the increased mutation

rates around strong nucleosomes are shaped by a combination of low-fidelity

replication, frequent DNA damage and insufficient/error-prone repair in these regions.

Interestingly, strong nucleosomes are preferentially located in young SINE/LINE

elements, implying frequent nucleosome re-positioning (i.e. shifting of dyad position)

and their contribution to hypermutation at new retrotransposons during evolution.

These findings provide novel insights into how chromatin organization affects

germline mutation rates and have important implications in human genetics and

genome evolution.

1 Introduction

Germline *de novo* mutations, which arise during gametogenesis and can be passed to offspring, are the primary source of genetic variation, contributing substantially to biological diversity and evolution. *De novo* mutations are also thought to play significant roles in early-onset genetic disorders such as intellectual disability, autism spectrum disorder, and other developmental diseases (Veltman and Brunner 2012; Acuna-Hidalgo et al. 2016). Thus, investigating the patterns and genesis of *de novo* mutations in the germline is important for understanding genome evolution and human diseases.

It is well known that germline and somatic mutation rates vary across the human genome over several scales of resolution, from nucleotide- to whole chromosome levels (Hodgkinson and Eyre-Walker 2011; Segurel et al. 2014). Previous studies of germline and somatic mutations (mainly cancer genomic variants) revealed several factors linked to mutation rate variation across the human genome, including local sequence context (Michaelson et al. 2012), replication timing (Stamatoyannopoulos et al. 2009), recombination rate (Francioli et al. 2015), DNA accessibility (Sabarinathan et al. 2016) and histone modifications (Michaelson et al. 2012; Schuster-Bockler and Lehner 2012). Despite these findings, however, two recent studies (Terekhanova et al. 2017; Smith et al. 2018) reported that less than 40% of local mutation rate variation at 100Kb to 1Mb resolution can be explained by the genomic features identified so far, implying that unknown but important factors remain to be found. Moreover, the underlying mutational processes causing the observed mutation rate variation remain poorly understood. A few recent studies pointed to the contribution of error-prone replicative processes (Harris and Nielsen 2014; Lujan et al. 2014; Reijns et al. 2015; Seplyarskiy et al. 2017; Seplyarskiy et al. 2018) and differential DNA repair efficiency (Supek and Lehner 2015; Perera et al. 2016; Sabarinathan et al. 2016; Frigola et al. 2017). Despite these advances, it remains a big challenge to understand the mutational mechanisms associated with mutation rate variation, particularly in the germline.

Chromatin organization has been reported to contribute to regional variation in mutation rates (Makova and Hardison 2015) because of structural constraints on the mutability of genomic sequences. In particular, nucleosome positioning was suggested as a significant factor in humans and other eukaryotes (Sasaki et al. 2009; Tolstorukov et al. 2011; Chen et al. 2012; Michaelson et al. 2012; Lujan et al. 2014; Pich et al. 2018). Studies in different lineages (Sasaki et al. 2009; Tolstorukov et al.

2011; Lujan et al. 2014) reported increased substitution rates around the centers of nucleosomal sequences and increased insertion/deletion rates in nucleosome-free DNA. However, there are also disagreements between published studies. For example, Michaelson et al. (2012) suggested that high nucleosome occupancy tends to suppress *de novo* mutations, but Smith et al. (2018) found that a comparative analysis using datasets from different studies resulted in opposing conclusions, likely affected by different variant calling strategies. Due to the limited availability of germline *de novo* mutations for humans, most previous studies focused on coarse-grained mutation rate variation (e.g. ≥1kb windows), or used within-species polymorphisms and inter-species divergence for analysis. The drawback of using polymorphism and divergence data is that they might be affected by natural selection and other evolutionary processes such as GC-biased gene conversion. In addition, because of the limitation of available nucleosome maps, some previous studies treated all annotated nucleosomes equally, ignoring the diverse contexts in which they form. Therefore, the effects of nucleosome organization on germline mutation rate variation, particularly at high resolution remain to be elucidated. Here we take advantage of the rapid increase in the number of germline *de novo* mutation datasets and better understanding of nucleosome organization in the human genome to perform a systematic analysis of this topic.

2 Results

2.1 Datasets used for analysis

We used germline *de novo* mutations identified in multiple large-scale trio sequencing projects; these total >300,000 single-nucleotide variants (SNVs) and >30,000 short insertions/deletions (INDELs) (**Fig. 1a, Supplementary Fig.1a**), after removing those in genomic regions that could confound downstream analysis (see Methods). Most of these *de novo* mutations originated from three studies (Jonsson et al. 2017; Turner et al. 2017a; Yuen et al. 2017), each presenting about 100,000 mutations. In addition to *de novo* mutations from trio sequencing, we also examined extremely rare variants (allele frequency ≤ 0.0001) from the gnomAD database (Lek et al. 2016), which are thought to be subject to limited selection and non-adaptive evolutionary processes, and so can be approximated to *de novo* mutations (Carlson et al. 2018).

Fig. 1 *De novo* mutations in different nucleosome contexts. (**a**) Germline *de novo* mutation data used for analysis. Numbers of *de novo* mutations after filtering out those in confounding regions are shown in the brackets. (**b**) Illustration of MNase-seq method and the nucleosome datasets used in this analysis. (**c**) Fold enrichment/depletion of *de novo* SNVs in different nucleosome contexts. 'Strong', translationally stable positioning; 'Rotational', rotationally but not translationally stable positioning; 'Others', the remaining genomic regions. On the left is the fold enrichment for three subgroups of strong nucleosomes with different stabilities. (**d**) Fold enrichment/depletion of *de novo* INDELs in different nucleosome contexts. (**e**) Fold enrichment/depletion of strong nucleosomes in different repeat sequence contexts. SINE, short interspersed nuclear element; LINE, long interspersed nuclear element; TR, tandem repeat; LTR, long terminal repeat. (**f**) Above, meta-profile of *de novo* SNV/INDEL densities, centered at dyads of strong nucleosomes. Below, the same meta-profile zoomed in on the middle nucleosome in the array.

Nucleosome positioning is described by the translational setting, which defines the location of the nucleosomal midpoint (also called 'dyad'), and the rotational setting, which defines the orientation of the DNA helix on the histone surface (Gaffney et al. 2012). Using MNase-seq measurements across seven lymphoblastoid cell lines, Gaffney et al. (2012) previously identified about a million 'strong' nucleosomes that adopt highly stable translational positions across the sequenced cell lines (covering 198Mb of the genome). We also considered rotationally stable nucleosomes identified in DNase-seq samples across 43 cell types (Winter et al. 2013), covering 892Mb of the genome. There is a ~50Mb overlap between regions bound by strong nucleosomes and rotationally stable nucleosomes.

We divided the human genome (sex chromosomes excluded) into three categories based on the type of nucleosome context (**Fig. 1b**): i) regions containing translationally stable ('strong') nucleosomes (198Mb); ii) regions with rotationally but not translationally stable nucleosomes (796Mb); and iii) all other non-N base genomic regions (1,703Mb). Although nucleosome positioning can vary between cell types, particularly in transcriptionally regulated regions, West et al. (2014) suggested that such variation is very limited overall. As the nucleosome maps were not derived from germ cells, we excluded nucleosomes that were found to differ in positioning between cell types (~23Mb; see Methods).

2.2 *De novo* mutations in different nucleosome contexts

We first checked the numbers of mutations in the different genomic regions defined above. Regions associated with strong nucleosomes have ~30% more *de novo* SNVs (**Fig. 1c**) and ~15% more *de novo* INDELs (**Fig. 1d**) relative to expectations, whereas regions containing rotationally stable nucleosomes are slightly depleted of both mutation types. These enrichments are also apparent for the extremely rare variants from gnomAD (**Supplementary Fig. 1b,c**), though effect sizes are smaller compared with *de novo* mutations; the smaller enrichment here could be due to the fact that highly mutable sites are under-represented among extremely rare variants owing to higher allele frequencies in the population (Harpak et al. 2016). When separating the one million strong nucleosomes to three subgroups based on the translational positioning stability scores defined by Gaffney et al., we found that the nucleosomes with higher stability exhibit higher mutation rates (**Fig. 1c,d**), suggesting the nucleosome translational positioning stability (termed positioning stability for short hereafter) is associated with the mutation rate variation across the genome.

We further noticed that ~80% of strong nucleosomes overlap with repetitive elements,
especially short interspersed nuclear elements (SINEs) and long interspersed
nuclear elements (LINEs) (**Fig. 1e**). Genetic variations in repetitive sequences are
difficult to detect because of poor mappability, leading to a high proportion of false
negatives (Lee and Schatz 2012). Thus the enrichments of *de novo* mutations and
rare polymorphisms in strong nucleosomal regions are probably underestimated.

Interesting mutation patterns also emerge at higher resolutions. Meta-profiles of
mutation densities relative to the nucleosomal dyads reveal increased SNV density
and reduced INDEL density compared with the flanking linker regions (**Fig. 1f**),
resembling previous observations based on polymorphism data and a much smaller
number of nucleosomes (Tolstorukov et al. 2011). Since many strong nucleosomes
are associated with Alu elements and LINE-1 (L1) elements (Alu and L1 are the most
abundant SINE and LINE families in the human genome, respectively), we separated
the strong nucleosomes into three groups: i) Alu-associated, ii) L1-associated and iii)
others. Alu-associated nucleosomes display increased SNV rates around the dyads
(**Supplementary Fig. 1d**), whereas non-Alu nucleosomes show increased SNV rates
~60bp from the dyads, close to the nucleosome edges. In contrast, the patterns of
INDEL densities are relatively similar among the different groups (**Supplementary
Fig. 1d**).

**2.3 Statistical tests confirm the significant association between nucleosome
positioning stability and mutation rate variation**

Local sequence contexts can affect mutation rates; for instance, CpG sites are
known to be highly mutable (**Supplementary Fig. 1d**). Other factors such as DNA
methylation, histone modification, chromatin accessibility, replication timing and
recombination rate have also been reported to affect mutation rates. Therefore, to
control for potential confounding factors, we used a logistic regression framework to
systematically assess the contribution of nucleosome positioning to mutation rate
variation.

We defined three variables to quantify nucleosomal properties relative to a specific
nucleotide position. The first two relate to the translational positioning of nearby
nucleosomes: i) d_{mean}, the mean distance between the focal site and the midpoints of
mapped MNase-seq fragments (we considered MNase-seq midpoints within ±95bp of
the focal site, corresponding to the average nucleosome repeat length reported by
Gaffney et al. 2012) and ii) d_{var}, the variance of these distances (**Fig. 2a**). A smaller
d_{mean} indicates that the site is closer to the nucleosome dyad and smaller d_{var}

represents more stable translational positioning around that site. Because the
relationship between d_{mean} and SNV rates is not monotonic nor linear (see **Fig. 1f,**
Supplementary Fig. 1d), we considered d_{mean} be a categorical variable by binning
values into five intervals (see Methods). The third variable is nucleosome occupancy,
calculated as the per-base MNase-seq fragment coverage after correcting for
sequencing and mapping biases (see Methods).

We also considered the local nucleotide sequence (5bp upstream and downstream of
the focal site), RNA expression in spermatogonial stem cells (SSCs), DNA
methylation (SSCs), chromatin accessibility (ATAC-seq in SSCs), eight histone
modifications (ENCODE ChIP-seq in ESCs), replication timing (Repli-seq in
GM12878) and recombination rates (HapMap project). We found that nucleosome
positioning stability (d_{var}) has a relatively weak but statistically significant correlation
with many of these factors (**Supplementary Fig. 2**), suggesting non-independence of
factors; therefore, it is important to control for them in assessing the contribution of
positioning stability to mutation rate variation.

We compared the full logistic regression model encompassing all variables with
reduced models missing individual variables using likelihood-ratio tests (**Fig. 2b,c**;
see Methods for more detail). As CpG sites are highly mutable and contribute to a
large fraction of mutations, we tested CpG and non-CpG sites separately (**Fig. 2b,c**).
We found that positioning stability (d_{var}) is significantly associated with mutation rate
variation in both CpG and non-CpG contexts while controlling for other factors.
Generally, the effects of positioning stability on mutation rates for A/T sites are
greater than for C/G sites, and the effects on SNV mutation rates more significant
than for INDELs. The lower significance for INDELs could partly be due to the fact
that there are fewer such *de novo* mutations available for modeling. More specifically,
among SNV mutation rates, we found highly significant contributions of positioning
stability for all mutation types in CpG and non-CpG contexts (**Supplementary Fig. 3**),
except for CpG>TpG mutations (adjusted p = 0.10). Notably, the regression
coefficients for d_{var} are always negative (**Supplementary Table 1**), indicating that
positioning stability is inversely associated with mutation rates, corroborating the
patterns observed in **Fig. 1**.

Fig. 2 Statistical tests for evaluating contribution of nucleosome organization to mutation rate variation. (**a**) Definition of two nucleosome positioning-related variables (d_{mean} and d_{var}) for a given site in the genome. (**b-c**) Statistical significance of each variable by performing likelihood ratio tests for assessing their contribution to mutation rate variation: **b** for *de novo* SNVs and **c** for *de novo* INDELs. For *de novo* SNVs, we performed tests for A/T sites, C/G sites in non-CpG context and C/G sites in CpG context separately. The red vertical lines represent the significance cut-off (0.05) for the adjusted p values (Bonferroni-Holm correction). 'us', upstream; 'ds', downstream.

The mean distance to the dyads, d_{mean}, also has statistically significant effects on SNV/INDEL mutation rates (**Fig. 2b,c**). The relationship with SNVs (but not INDELs) is non-linear, as evidenced by the loss of significance upon treating d_{mean} as a continuous variable (**Supplementary Fig. 4**). Nucleosome occupancy is also statistically significant; in contract to the positioning variables however, here the effect is much larger for INDELs than SNVs (**Fig. 2b,c;** INDELs, adjusted p = 5.8e-37; SNVs, adjusted p = 0.21, 1.6e-6, 2.2e-7).

In agreement with previous observations on rare variants (Carlson et al. 2018), local sequence context appears to be the biggest contributing factor for local mutation rate

variation (**Fig. 2b,c**). Contributions generally decrease with increasing distance to the surveyed site. Although the larger sequence context could still have influence on mutation rates, including more distant bases can lead to some issues such as overfitting in regression models. We appreciate that nucleosome positioning stability is at least partly determined by the occupied DNA sequence and thus its effects on mutation rates to some degree can be attributed to the associated sequence (this also applies to other reported factors such as replication timing). However, higher-order interactions among the long stretches of nucleotides which guide nucleosome positioning are difficult to model properly. We achieved similar statistical significance for positioning stability after including non-additive two-way interaction effects for ±5 nucleotides and the 7-mer mutability estimates from Carlson et al. (**Supplementary Fig. 5**).

Finally in addition to sequence context, we recapitulated the effects of previously reported factors (**Fig. 2b,c, Supplementary Fig. 3**). DNA methylation and H3K9me3 are two common epigenetic signatures that statistically significantly contribute to mutation rate variation (Schuster-Bockler and Lehner 2012). H3K4me1, H3K4me2, H3K4me3 and H3K27me3 are also statistically significant for specific mutation types. Replication timing has highly statistically significant effects on both SNV and INDEL mutation types. Recombination rate and open chromatin (measured by ATAC-seq) also have significant effects on many mutation types. However, transcription levels appear to have no effect on local mutation rates according to our result.

Taken together, our modeling analysis recapitulated previously reported factors affecting mutation rates and confirmed the importance of nucleosome positioning stability as another contributing factor.

2.4 Mutational processes associated with elevated mutability around strong nucleosomes

2.4.1 Mutational signature analysis

Next we sought to investigate the underlying mutational processes that explain the elevated mutation rates associated with strong nucleosome positioning. For this, we first used the MutationalPatterns software (Blokzijl et al. 2018) to calculate the contribution of different COSMIC mutational signatures to germline *de novo* SNVs. Three major signatures (Signatures 1, 5 and 16) explain 88% of all *de novo* SNVs (**Fig. 3a**); however, Signatures 3 (3.4%), 6 (14%), 7 (2.8%) and 12 (9.2%) are enriched only among SNVs associated with strong nucleosomes (**Fig. 3a**). There are

further differences between nucleosome-associated SNVs overlapping Alus, L1s and 'other' regions (**Fig. 3a);** for example, Signatures 6 and 12 make major contributions to Alu-associated SNVs (27% and 17% respectively) but less marked to 'Others' group (3.7% and 3.9%) and none to L1 group. Signature 3 is found in all three groups, but shows relatively higher contribution in the L1 group. The mutational signatures of L1-associated SNVs are similar to that of genome-wide background (**Fig. 3a**), which might be because (i) L1 elements make up a considerable fraction (>20%) of the whole genome and (ii) strong nucleosomes are under-detected in very young L1 elements owing to difficulties in mapping short MNase-seq reads (Alus are easier to map because they are smaller).

Fig. 3 Analysis of mutational processes related to strong nucleosomes. (a) Comparison of contribution of COSMIC mutational signatures in different sets of *de novo*

SNV data. Only signatures with non-zero values are shown. Asterisks indicate the
mutational signatures showing ≥2% increase relative to the background contribution (i.e.
'All SNVs'). On the right is brief information about aetiologies (from the COSMIC website)
of highlighted signatures. (**b**) Mutation profiles around strong nucleosomes for the
PCAWG cancer genomes harboring driver mutations in *POLE* and genes involved in
mismatch repair (MMR). Numbers of cancer genomes used are indicated in the brackets.
The relative escape ratios of MMR were estimated by comparing '*POLE*-mutated MMR
proficient' and '*POLE*-mutated MMR deficient' samples. The profiles are centered at the
annotated strong nucleosomes. (**c**) Mutation profiles around strong nucleosomes for
bMMRD cancer genomes and the estimated relative escape ratios of Pol ε or Pol δ. (**d**)
Signal of DSBs around strong nucleosomes based on the END-seq data. Only the strong
nucleosomes of high 75-mer mappability within ±500bp were considered. HU,
hydroxyurea.

Having assigned mutational signatures, we can infer possible underlying DNA
damage and repair mechanisms that explain the elevated local mutation rates (**Fig.
3a**). Signature 1, a common mutational process due to spontaneous deamination of
5-methylcytosine, displays increased contribution to the Alu-associated SNVs; this is
expected because young Alu elements harbor many CpG sites and are usually highly
methylated. For other suggested mutational processes in **Fig. 3a**, we took advantage
of multiple genomic datasets for further investigation (see below).

309 2.4.2 Mismatch repair

Signature 6, which contributes ~14% to strong nucleosome-associated SNVs, is
linked to mismatch repair (MMR). This is a major repair pathway active during DNA
replication: it mainly repairs mismatches and short INDELs introduced during DNA
synthesis that escaped polymerase proofreading. We assessed the impact of the
MMR pathway by analyzing somatic mutations from two sets of cancer genomes
(Campbell et al. 2017). The first possessed driver mutations in catalytic subunit gene
POLE of DNA polymerase ε (Pol ε, the major replicase for the leading strand) and
one or more of the core MMR genes (*MLH1*, *MSH2*, *MSH6*, *PMS1* and *PMS2*). The
second set comprised cancers with mutated *POLE* but intact MMR pathway.
Mutations in *POLE* in these cancers can lead to reduced base selectivity and/or
deficient proofreading during replication, producing unusually large numbers of
mutations (so called 'ultra-hypermutation') which facilitated our analysis. Because
detecting somatic mutations in tumors is more difficult than for ordinary individual re-
sequencing data and mutations in poor-mappability regions could be easily missed,
we focused on strong nucleosomes with high mappability (see Methods).

For samples with defective *POLE* and MMR, mutation rates around the strong nucleosome boundaries and adjacent linker DNA are much higher than surrounding regions (**Fig. 3b**). This suggests replication errors occur frequently around strong nucleosomes. We detected a similar pattern for samples with mutated *POLE* but proficient MMR. This implies that many errors introduced during error-prone replication of mutated *POLE* also escaped the repair by MMR at strong nucleosomes, which were then translated into somatic mutations.

By comparing the mutation rates in the two sets of *POLE*-mutated cancers, we were able to estimate the MMR escape ratio, which quantifies the relative amount of replication errors that remains unrepaired (see Methods for more detail). Strong-nucleosome regions (especially boundaries and adjacent linkers) have ~10% higher MMR escape ratio than the genome-wide background, suggesting that replication errors at these regions are more likely converted to mutations in daughter cells. Differences in the escape ratio in these regions cannot be ascribed to local base composition, as both C/G sites and A/T sites exhibit similar patterns (**Fig. 3b**), though A/T sites appear to have a higher ratio. Moreover, the apparent ~200-bp periodic patterns in escape ratios and mutation densities (**Fig. 3b**) are suggestive of associations with nucleosome positioning rather than sequence alone.

2.4.3 DNA polymerase fidelity

We also analyzed data from children with inherited biallelic mismatch repair deficiency (bMMRD) (Shlien et al. 2015), which included ultra-hypermutated genomes with mutations in *POLE* or the catalytic subunit gene *POLD1* of polymerase δ (Pol δ, the major replicase for the lagging strand). The bMMRD data can be used for investigating polymerase fidelity around strong nucleosomes. Here, we estimated Pol ε and Pol δ escape ratios using the same method as that for MMR escape ratios (**Fig. 3c**, see Methods). We found that strong nucleosomes have higher escape ratios for both polymerases relative to the genomic background (**Fig. 3c**), implying that replication polymerases have lower fidelity in these regions. These escape ratios are even higher than that for MMR (**Fig. 3b,c**), and A/T sites also have larger escape ratios than C/G sites (**Supplementary Fig. 6a**). In addition, the periodic pattern in the relative escape profiles (**Fig. 3c, Supplementary Fig. 6a**) suggests that nucleosome positioning contributes to the heterogeneity in replicase fidelity across the genome.

Signature 12 is another prominent mutational signature that contributes 9.2% to *de novo* SNVs associated with strong nucleosomes, but whose aetiology is currently unknown. By performing mutational signature analysis, we found that Signature 12

contributes 21.15%~21.99% to mutations in *POLD1*-mutated bMMRD samples
(**Supplementary Fig. 6b,c**), but much less for other bMMRD samples (0~2.88% for
POLE-mutated, and 3.32%~10.43% for *POLE/POLD1*-intact). This suggests that the
previously uncharacterized signature is likely associated with Pol δ, and that many *de*
novo mutations around strong nucleosomes arise from errors escaping Pol δ proof-
reading and post-replicative MMR during lagging strand replication. On the other
hand, Signature 10, the only signature known to be associated with Pol ε deficiency,
exhibits little contribution to strong nucleosome-associated SNVs (**Fig. 3a**). This
suggests that although both Pol ε and Pol δ have low fidelity around strong
nucleosomes (**Fig. 3c**), the majority of the replication errors that are eventually
converted to *de novo* mutations are derived from Pol δ.

Previous analysis in budding yeast suggested that Okazaki fragment-associated
mutagenesis during lagging strand replication contribute significantly to mutation rate
variation across the genome (Reijns et al. 2015). They found that in budding yeast
Okazaki junctions tend to have elevated mutation rates and are usually close to
nucleosome dyads. We tested this point in human by re-analyzing the OK-seq data
(Petryk et al. 2016), which sequenced the Okazaki fragments in human cells. We
found that, unlike that in yeast, Okazaki junctions in human appear to be relatively
more frequently located around the linker regions (**Supplementary Fig. 7**), rather
than the dyads, implying the mutagenic effects of Okazaki junctions in yeast are not
the same in human. The difference may be partly due to the reason that budding
yeast lacks the typical H1 histone found in human and other eukaryotes. However,
the very short reads (single-ended 50bp) of OK-seq data only allowed us to examine
the nucleosomes with high mappability with 50-mers (~10% of strong nucleosomes),
the distribution of Okazaki junctions in unexamined regions is unclear.

2.4.4 Double-strand breaks

Double-strand break (DSB) repair represented by Signature 12 is another potential
mechanism involved in strong nucleosome-associated mutations (**Fig. 3a**). A recent
study (Tubbs et al. 2018) investigating genome-wide DSBs using END-seq
suggested that poly(dA:dT) tracts are recurrent sites of replication-associated DSBs.
Our analysis of this data revealed a higher frequency of DSBs around strong
nucleosomes compared with genomic background (**Fig. 3d**). The trend holds for
experiments with and without hydroxyurea treatment (HU, a replicative stress-
inducing agent), suggesting that strong nucleosomes contribute to endogenous
replication stress and higher frequency of DSBs. It is notable that young Alu and L1

elements harbor prominent poly(dA:dT) tracts, which are enriched at the boundary and linker regions of strong nucleosomes (**Supplementary Fig. 8a**). The patterns of high DSB frequency still hold true when looking at strong nucleosomes associated with different repeats (**Supplementary Fig. 8b-c**). Yet, because the END-seq data was sequenced with single-ended 75bp reads and majority of young Alu and L1 elements can not be assessed with such short reads, we did not perform further analysis regarding this. Since DSB repair can be error-prone (Rodgers and McVey 2016), even using high-fidelity homologous recombination, frequent DSB-formation and subsequent error-prone repair likely contribute to the elevated mutation rates around strong nucleosomes.

2.5 Strong nucleosome positioning is mostly associated with young repeat elements and undergoes frequent turnover

~70% of strong nucleosomes are located in SINE and LINE retrotransposons (**Fig. 1e**). Upon examination of the subfamilies (**Fig. 4a,b**), we uncovered specially strong enrichments for evolutionarily young L1s (e.g. L1PA2 to L1PA11) and Alus (e.g. AluY to AluSx). Since younger repeats have poorer mappability, these observations probably underestimate the true enrichment. This may also explain why several of the youngest L1 subfamilies (L1PA2 to L1PA5) have lower enrichments relative to the slightly older subfamilies (**Fig. 4a**). It has long been reported that nucleosomes preferentially bind specific segments of Alu elements, supported by both *in vitro* and *in vivo* evidence (Englander et al. 1993; Englander and Howard 1995). Later studies confirmed preferential nucleosome positions within Alus (Salih et al. 2008; Tanaka et al. 2010). These studies also suggested that A-rich sequences and CpG dinucleotides in Alus likely play important roles in the unique positioning patterns. We recapitulated their findings using MNase-seq data of Gaffney et al. (**Fig. 4c**) with two hotspots of strong nucleosomes in young Alus, which fade away in older elements. Young Alu elements exhibit elevated *de novo* mutation rates (**Fig. 4c**), which could be partly contributed by the associated strong nucleosome positioning as evaluated in previous sections.

The weakening of positioning stability in older Alu elements is accompanied by reduced *de novo* mutation rates for both SNVs and INDELs, although the pattern in INDELs is less clear compared with SNVs (**Fig. 4c**). We hypothesize that the sequence changes in young Alu elements arising from *de novo* mutations might affect the stability of nucleosome positioning. The histone octamer is thought to preferentially bind DNA sequences presenting lower deformation energy costs

(Tolstorukov et al. 2008). Using the nuScore software (Tolstorukov et al. 2008) to
estimate deformation energies based on the DNA sequence and nucleosome core
particle structure, we indeed found that Alus exhibit lower deformation energies than
surrounding regions. Furthermore, the deformation energies increased in older
subfamilies, suggesting that the evolutionary changes in Alu sequences reduced their
nucleosome-binding affinity. We further analyzed the 3' end sequences of L1
elements harboring strong nucleosomes and observed similar patterns
(**Supplementary Fig. 9**).

Fig. 4 Strong nucleosomes are mostly associated with young LINE/SINE elements.
439

(**a**) Fold enrichment/depletion of strong nucleosomes in LINE/L1 subfamilies of different
ages (top 30 abundant subfamilies are shown), with dot sizes and colors depicting
numbers of associated strong nucleosomes and subfamily ages respectively. (**b**) Fold
enrichment/depletion of strong nucleosomes in SINE/Alu subfamilies of different ages. (**c**)
Densities of strong nucleosome dyads, *de novo* SNVs and *de novo* INDELs along the
Alu sequence and surrounding regions for Alu subfamilies of different ages, with the
average values for all Alu regions of different subfamilies in the barplots on the right.
Two major loci for strong nucleosomes are shown at the top. The deformation energy
estimated by nuScore is shown at the bottom panel. Alu elements of >=250bp were used
for plotting the meta-profiles and all elements were scaled to a 300bp region in the plots.
The structure of the Alu consensus sequence is shown at the top and bottom.

Several studies investigating genetic changes around nucleosomes suggested that
natural selection appears to preserve nucleosome positioning during evolution
(Prendergast and Semple 2011; Tolstorukov et al. 2011; Drillon et al. 2016), but they
had different opinions on how selection acts on the sequence. On the other hand,
another model (Warnecke et al. 2013) suggested that the observed sequence
divergence patterns around nucleosomes can be explained by frequent nucleosome
re-positioning after mutations, rather than by natural selection. These studies were
mainly based on human polymorphisms or inter-species divergence and so were
potentially confounded by mutation rate heterogeneity associated with nucleosomes
and other non-adaptive processes such as GC-biased gene conversion. The use of
germline *de novo* mutations helps resolve this debate. As we showed before, there is
considerable *de novo* mutation rate variation around strong nucleosomes (**Fig. 1f**,
Supplementary Fig. 1), which can not be ignored in any selection analysis.
Furthermore, analysis on strong nucleosomes associated with different SINE/LINE
subfamilies clearly showed that strong positioning is preferentially present in young
elements and the positioning stability decays substantially over time (**Fig. 4**), which
strongly supports the re-positioning model. Since LINE/SINE elements make up a
large majority of strong nucleosomes in the genome, selection for preserving
positioning is unlikely widespread as previously suggested, though it may happen at
some particular regions.

3 Discussion

While nucleosome organization is long known to be involved in DNA damage/repair
processes (Smerdon 1991), its effects on germline mutation rate variation across the
genome (particularly in higher eukaryotes) are poorly understood, largely due to the
limited available *de novo* mutations and poor-resolution of nucleosome mapping. Our
analysis combining several large-scale *de novo* mutation and nucleosome datasets
in human have provided several important insights into this topic.

One main finding in our work is that strong nucleosome translational positioning is
associated with elevated germline *de novo* mutation rates, which is also supported by
the analysis with extremely rare variants in polymorphism data. The direct use of *de*
novo mutations in the analysis avoided many confounding evolutionary factors in
some previous studies. Importantly, our statistical tests controlling for nucleosome
occupancy and other related factors confirmed the significant contribution of
nucleosome positioning stability to the mutation rate variation. While nucleosome
organization was reported to be associated with mutation rate variation in human,

previous research usually focused on nucleosome occupancy and positioning stability was rarely considered. Therefore we discovered a novel factor that significantly modulate the germline mutation rate variation, which we think is an important aspect for future related research.

Investigating mutational processes in the germline and somatic tissues remains challenging. Nevertheless, by analyzing published genomic datasets, we obtained several informative results regarding potential mutational processes associated with strong nucleosomes. Analysis based on the COSMIC mutational signatures suggested that MMR and DSB are likely involved in the elevated mutation rates around strong nucleosomes. Our further analysis with multiple independent datasets indeed supported MMR and DSB as contributing mutational mechanisms. In addition, analysis with the bMMRD cancers suggests that lower replicase fidelity of both Pol ε and Pol δ around strong nucleosomes also contributes to the elevated mutation rates. The underlying molecular mechanisms about how strong positioning affects replicase fidelity and associated repair (MMR or DSB repair) are still not clear. Based on the evidence from our analysis and previous studies (Li et al. 2009; Reijns et al. 2015; Tubbs et al. 2018), we speculate that strong nucleosomes may act as strong barriers which impair the performance of the replication and repair machineries in those regions. Strong nucleosome positioning could also affect other unexamined DNA damage/repair processes related to germline development, but many published genomic datasets about DNA damage/repair were generated with very short sequencing reads (e.g. <100bp), which hinder detailed analysis in highly repetitive regions. Improved sequencing strategies such as long-read sequencing should be beneficial for future related studies.

Interestingly, we found that strong nucleosomes are preferentially located in young LINE and SINE elements, two common types of retrotransposons in the human and other mammalian species. Due to their potentially deleterious effects, newly inserted retrotransposons are highly repressed by multiple regulatory mechanisms, such as DNA methylation and H3K9me3 (Slotkin and Martienssen 2007). Strong nucleosome positioning could be another layer of the repressive system. Strong positioning in young SINEs/LINEs may mask access to the transcription machinery. Furthermore, the contribution of strong nucleosomes to hypermutation in young SINEs/LINEs can reduce the retotransposition capacity more rapidly. Therefore, strong positioning and hypermutation in SINEs/LINEs might have facilitated the their spread across the genome and help produce large amounts of genetic variations (substrates for natural selection) for the host.

The decreasing numbers of strong nucleosomes in older LINE/SINE elements imply
widespread nucleosome re-positioning during evolution. Since nucleosome
positioning is affected by the surrounding sequences, the re-positioning probably
results from accumulated mutations. Our data disagree with the previous hypothesis
of widespread selection for maintaining nucleosome positioning in the human
genome (Prendergast and Semple 2011). Another reason for favoring the re-
positioning model is that most genomic regions do not employ strong positioning,
possibly due to its relatively high mutagenic potential.

Finally, we summarized our major findings in a proposed model in **Fig. 5**, which
demonstrates the relationship among nucleosome positioning, mutation rate variation,
retrotransposons and evolution. Given the importance of germline *de novo* mutations
in evolution and human diseases and the universal roles of nucleosomes in
eukaryotic genome organization and regulation, our work should have profound
implications in related research areas.

Fig. 5 Proposed model for how nucleosome positioning stability modulates
mutation rate variation and influences evolution. While the genome is generally
covered by weak nucleosomes, newly inserted SINE/LINE retrotransposons are
enriched with strong nucleosomes. The regions bound by strong nucleosomes are
subject to high mutation rates during germline development, which in turn reduce the
nucleosome binding affinity of the sequence and the transposition capacity of associated
retrotransposons.

Methods

Mutation datasets

De novo mutations identified in multiple large-scale trio sequencing project were downloaded from denovo-db v1.6.1 (Turner et al. 2017b). Seven studies with >1000 *de novo* mutations (Genome of the Netherlands 2014; Turner et al. 2016; Yuen et al. 2016; Jonsson et al. 2017; Turner et al. 2017a; Yuen et al. 2017; Werling et al. 2018) were considered in our analysis (**Supplementary Fig. 1a**). Extremely rare variants (derived allele frequency ≤ 0.0001) were obtained from Genome Aggregation Database (gnomAD, release 2.0.2) (Lek et al. 2016).

Nucleosome datasets

We used 1,037,801 strong nucleosomes (i.e. translationally stable nucleosomes) identified based on MNase-seq data of sequenced seven lymphoblastoid cell lines from Gaffney et al. (Gaffney et al. 2012). The original hg18-based coordinates of annotated nucleosomes were converted to hg19 using the 'liftOver' tool from UCSC genome browser. The rotationally stable nucleosomes identified based on 49 DNase-seq samples (43 distinct cell types) were from Winter et al. (Winter et al. 2013). We classified the human genome into three groups based on the nucleosome contexts (**Fig. 1b**): 1) regions covered by translationally stable ('strong') nucleosomes; 2) regions covered by rotationally but not stable translationally nucleosomes; and 3) the remaining genome regions. Chromosomes X and Y were excluded from analysis as some other datasets used in our work lacked data for these chromosomes. As the nucleosome maps we used were not derived from germ cells, for downstream analysis we excluded the genomic regions in which nucleosome positioning were found to differ between human embryonic stem cells and differentiated fibroblasts (West et al. 2014). Based on the positioning stability scores defined in Gaffney et al., we divided the one million strong nucleosomes into three categories of equal sizes with different levels of stability – 'high', 'middle' and 'low', which were used for analysis shown in **Fig. 1** and **Supplementary Fig. 1.**

Accounting for mappability

Sequencing read mappability can significantly affect variant calling results and other aligned read-depth based measurements (e.g. nucleosome occupancy). The sequencing reads for detecting *de novo* mutations used in our analysis were mainly 150bp paired-end reads, with fragment sizes ranging from 300-700bp (**Supplementary Fig. 1**). We used the Genome Mappability Analyzer (GMA) (Lee

and Schatz 2012) to generate the mappability scores for simulated PE150 reads with fragment sizes set to be 400bp. Only the regions with GMA mappability scores of >=90 (~2.6Gb) were considered for most analyses, unless specified otherwise. We did not use the mappability tracks from ENCODE for the de novo mutation data, because those tracks were only for single-ended reads. For some analyses, additional filtering were applied if other associated datasets suffered from more severe mappability issues. For measuring nucleosome occupancy, we used the method described in the Gaffney et al. to simulate paired-end 25bp reads matching the base compositions of MNase-seq data in the human genome, and then calculated per-base coverage depth by the simulated fragments. The 10bp-bin ratios between the MNase-seq read coverage and the simulated read coverage were used for measuring the occupancy.

Enrichment analysis for *de novo* mutations in different nucleosome contexts

The tool Genomic association tester (GAT) (Heger et al. 2013) was used to estimate the expected numbers of mutations in different contexts (sampling >=1000 times), which were then compared to the observed numbers. Low-mappability regions were excluded from analysis. A similar analysis was also done for the extremely rare variants of gnomAD. Analysis of meta-profiles along strong nucleosomes was done using deepTools (Ramirez et al. 2014).

Statistical modelling of the contribution of different factors to mutation rate variation

As described in the main text, for a given site, we defined two variables regarding the translational positioning of nearby nucleosomes (**Fig. 2a**):

$$d_{mean} = \frac{\sum_{i=1}^{n} d_i}{n}, 0 \leq d \leq 95,$$

$$d_{var} = \frac{\sum_{i=1}^{n}(d_i - d_{mean})^2}{n}$$

where *d* is the distance between a MNase-seq midpoint to the focal site, *0 ≤ d ≤ 95*. We considered MNase-seq midpoints within ±95bp of the focal site, because genome-wide nucleosome repeat length was estimated to be 191.4bp (Gaffney et al. 2012). Genomic sites without any MNase-seq midpoint within ±95bp were excluded from analysis. The measurements for nucleosome occupancy were 10bp-bin ratios between the MNase-seq read coverage and the simulated read coverage. We did not use the positioning score *S(i)* defined in Gaffney et al. to measure positioning stability

in our modelling analysis, because *S(i)* was designed for identifying the stable dyads
and so for non-dyad positions it does not represent the positioning stability properly.

RNA expression, DNA methylation and chromatin accessibility (ATAC-seq) data from
human spermatogonial stem cells was from Guo et al. (Guo et al. 2017). For the
RNA-seq and ATAC-seq data from Guo et al., because the genome-wide read signal
tracks were not available, we downloaded, processed and mapped the raw reads to
generate the genome-wide tracks. Since suitable data for histone modifications in
human germ cells was not available, we used the ChIP-seq data of human
embryonic stem cells from ENCODE (ENCODE Consortium 2012). Replication timing
data (Repli-seq of GM12878) was from ENCODE. The data of recombination rates
was from the HapMap project (International HapMap Consortium et al. 2007).

A binary logistic regression framework was used to assess the contribution of
different factors to mutation rate variation across the genome systematically. The
logistic regression model is described as below:

$$\mu = \Pr(y = 1) = \frac{\exp(\beta_0 + \beta_1 X_1 + \cdots + \beta_p X_p)}{1 + \exp(\beta_0 + \beta_1 X_1 + \cdots + \beta_p X_p)}$$

$$= \frac{\exp(\boldsymbol{X}\beta)}{1 + \exp(\boldsymbol{X}\beta)}$$

$$\text{logit}(\mu) = \log\left(\frac{\mu}{1-\mu}\right) = \boldsymbol{X}\beta$$

where $\mu = \Pr(y = 1)$ denotes the probability that a genomic site is mutated (for
testing individual SNV mutation types (e.g. A>T), μ is the probability that a site is
mutated to a specific nucleotide), $\boldsymbol{X}$ represents the observations for the considered
variables (categorical or continuous, e.g. d_{mean}, d_{var}, adjacent nucleotides, etc.), and β
is the vector of parameters to be estimated.

We used the Bayesian logistic regression model implemented in the 'bayesglm'
(Gelman et al. 2008) of the R package 'arm', which was reported to perform well in
handling the complete separation issue in logistic regression models (Gelman et al.
2008). The complete separation issue is common when one class is rare relative to
the other and (or) there are many regressors in a model. As we had only ~300,000
de novo mutations, the probability for a given site to be mutated in our data is
~1/10,000, which is a rare event.

Within the logistic regression framework, we compared the full model with all
considered variables to a reduced model without one specific variable by performing
likelihood-ratio tests in R ('anova' function) to evaluate the significance for each
variable. The resulting p values of a set of likelihood-ratio tests were adjusted for
multiple testing with Bonferroni-Holm correction.

To perform the regression analysis, we generated the data of all variables for the *de*
novo mutation sites and subsampled a fraction of the non-mutated sites as the
control sites. We did not use all the non-mutated sites in the genome as it would lead
to a large imbalance in the sizes of two classes ('mutated' and 'non-mutated') and
much larger computational burden. For *de novo* SNVs, we randomly generated
2,561,953 non-mutated sites (about 1/1000 of the accessible genome, about 10
times as many as *de novo* SNVs) and 256,337 non-mutated sites (about 1/10,000 of
the accessible genome, about 10 times as many as *de novo* INDELs) for INDELs.
For *de novo* INDELs, we used the INDELs of ≤5bp for regression analysis, because
long INDELs were rare and may have a high false-positive rate). For RNA expression,
DNA methylation, chromatin accessibility, replication timing, recombination rate and
histone modifications data, we used the average value of the ±10bp of a focal site for
the specific feature based on the genome-wide signal tracks. We also assessed
different window sizes (±5bp and ±20bp), which led to similar results.

For SNVs, we performed logistic regression tests for mutation types at A/T sites and
C/G sites separately and distinguished C/G sites in CpG and non-CpG contexts. We
also tested for nine individual SNV mutation types (three for A/T sites, three for C/G
sites at CpG contexts, and three for non-CpG contexts, **Supplementary Fig. 3**). The
regression coefficients for the full model of each test are given in **Supplementary
Table 1.**

Since the variable d_{mean} has a non-monotonic relationship with mutation rates, we
binned the values into five categories: [0,18], [19, 36], [37, 54], [55, 73] and [74, 95]
(first four bins implying nucleosome-bound regions, and the last bin implying close to
the linker). We also did tests when considering d_{mean} as a continuous variable
(**Supplementary Fig. 4)**.

In the regression models mentioned above, we did not consider the non-additive
effects of adjacent nucleotides (±5 bp). When we tried adding non-additive effects for
±5 nucleotides (considering only two-way interactions; taking a much longer running
time), we got similar results regarding the association of positioning stability (d_{var}) and
mutation rates (**Supplementary Fig. 5)**. We also tried using the 7-mer mutability

estimates from Carlson et al. (Carlson et al. 2018), which incorporated non-additive

effects among ±3 nucleotides, as predictors in the regression models and which gave

a similar significance for d_{var} (**Supplementary Fig. 5**).

Analysis of mutational processes

The contribution of COSMIC mutational signatures (Alexandrov et al. 2013) to

different sets of mutations (*de novo* SNVs and somatic mutations from bMMRD

samples) was predicted using the 'fit_to_signatures' function in the R package

'MutationalPatterns' (Blokzijl et al. 2018).

POLE mutated genomes from PCAWG project (Campbell et al. 2017) were used to

evaluate the differential MMR efficiency between strong and non-strong nucleosome

regions, We compared the mutation densities in cancer genomes with *POLE* mutated

and a deficient MMR to those with *POLE* mutated and a proficient MMR. The MMR

pathway was considered deficient if a driver mutation (annotated by the PCAWG

consortium) was found in one of five MMR core genes - *MLH1*, *MSH2*, *MSH6*, *PMS1*

and *PMS2*.

For a given bin (10bp-size) in the meta-profile, we calculated the relative MMR

escape ratio relative to genomic background around strong nucleosomes as

described in the following formula,

$$R_i^{escape} = \frac{\dfrac{m_i^{POLE^*, MMR^{WT}}}{m_i^{POLE^*, MMR^*}}}{\dfrac{\overline{m}^{POLE^*, MMR^{WT}}}{\overline{m}^{POLE^*, MMR^*}}}$$

where m_i is the mutation density for the *i*th bin, and $\overline{m}$ is the genome-wide average

mutation density of a specific sample group, estimated by simulating random

windows in the genome. A similar logic was used when evaluating relative escape

ratios of Pol ε (mutated POLE) and Pol δ (mutated POLD1) using the somatic

mutations generated by the bMMRD project (Shlien et al. 2015).

When analyzing PCAWG and bMMRD data, to account for potential mappability

issues, we focused on the highly mappable regions based on the CrgMapability

scores from ENCODE. We used CrgMapability scores here, which are more stringent

than GMA ones, because detecting somatic mutations in tumors is more difficult than

for ordinary individual re-sequencing data. We considered the strong nucleosomes

which have a 100mer CrgMapability score of 1 (meaning any 100-bp read from these

regions can be mapped uniquely in the genome) within ±800bp of the dyads. We

then simulated a same number of 1600bp-sized regions from the genome that satisfy the mappability requirement to calculate the background mutation density. Note that in theory the mappabiity issue in the relative escape ratios should be very small because the two sets of samples have the same mappability for a given bin and the ratio calculation normalizes the effects of different mappability among regions.

The raw reads of OK-seq data (Petryk et al. 2016) were downloaded from NCBI and mapped to the human genome. We kept only the uniquely mapped reads for inferring Okazaki junctions. The very 5' end sites of aligned reads (separating reads mapped to Watson and Crick strands) were considered putative Okazaki junction signals.

To investigate DSBs around strong nucleosmes, we downloaded the genome-wide tracks of human END-seq data (GSM3227951 and GSM3227952) (Tubbs et al. 2018). Because the reads of END-seq data were single-ended 75bp, we considered the strong nucleosomes which have a 75mer CrgMapability score of 1 within ±500bp of the strong nucleosome dyads for analysis.

Enrichment analysis for strong nucleosomes in different repeat contexts

GAT (Heger et al. 2013) was used to estimate the expected numbers of strong nucleosomes in different contexts (sampling >=1000 times), which were compared to the observed numbers. The annotations of repeat elements (Feb 2009, Repeat Library 20140131) were downloaded from RepeatMasker (Tempel 2012). We also did GAT analysis for LINE-1(L1) and Alu subfamilies of different ages. The age information of repeat families was from Giordano et al. (Giordano et al. 2007). For generating the MNase-seq midpoints along the repeat consensus sequences, we made use of the alignment information in the RepeatMasker result files ('hg19.fa.align.gz') and mapped the hg19-based coordinates to the coordinates in the consensus sequences. Nucleosome deformation energy scores of all sites in the human genome were estimated using nuScore (Tolstorukov et al. 2008). For the L1 analysis shown in **Supplementary Fig. 9**, we only considered the 3' end regions of L1 subfamilies, because 5' end regions of L1 elements are usually truncated in the genome and their subfamily identities are difficult to be determined.

References

Acuna-Hidalgo R, Veltman JA, Hoischen A. 2016. New insights into the generation and role of de novo mutations in health and disease. *Genome biology* **17**(1): 241.

Alexandrov LB, Nik-Zainal S, Wedge DC, Aparicio SA, Behjati S, Biankin AV, Bignell GR, Bolli N, Borg A, Borresen-Dale AL et al. 2013. Signatures of mutational processes in human cancer. *Nature* **500**(7463): 415-421.

Blokzijl F, Janssen R, van Boxtel R, Cuppen E. 2018. MutationalPatterns: comprehensive genome-wide analysis of mutational processes. *Genome medicine* **10**(1): 33.

Campbell PJ, Getz G, Stuart JM, Korbel JO, Stein LD. 2017. Pan-cancer analysis of whole genomes. *BioRxiv*: 162784.

Carlson J, Locke AE, Flickinger M, Zawistowski M, Levy S, Myers RM, Boehnke M, Kang HM, Scott LJ, Li JZ et al. 2018. Extremely rare variants reveal patterns of germline mutation rate heterogeneity in humans. *Nature communications* **9**(1): 3753.

Chen X, Chen Z, Chen H, Su Z, Yang J, Lin F, Shi S, He X. 2012. Nucleosomes suppress spontaneous mutations base-specifically in eukaryotes. *Science* **335**(6073): 1235-1238.

Drillon G, Audit B, Argoul F, Arneodo A. 2016. Evidence of selection for an accessible nucleosomal array in human. *BMC genomics* **17**: 526.

ENCODE Consortium. 2012. An integrated encyclopedia of DNA elements in the human genome. *Nature* **489**(7414): 57-74.

Englander EW, Howard BH. 1995. Nucleosome positioning by human Alu elements in chromatin. *The Journal of biological chemistry* **270**(17): 10091-10096.

Englander EW, Wolffe AP, Howard BH. 1993. Nucleosome interactions with a human Alu element. Transcriptional repression and effects of template methylation. *The Journal of biological chemistry* **268**(26): 19565-19573.

Francioli LC, Polak PP, Koren A, Menelaou A, Chun S, Renkens I, Genome of the Netherlands C, van Duijn CM, Swertz M, Wijmenga C et al. 2015. Genome-wide patterns and properties of de novo mutations in humans. *Nature genetics* **47**(7): 822-826.

Frigola J, Sabarinathan R, Mularoni L, Muinos F, Gonzalez-Perez A, Lopez-Bigas N. 2017. Reduced mutation rate in exons due to differential mismatch repair. *Nature genetics* **49**(12): 1684-1692.

Gaffney DJ, McVicker G, Pai AA, Fondufe-Mittendorf YN, Lewellen N, Michelini K, Widom J, Gilad Y, Pritchard JK. 2012. Controls of nucleosome positioning in the human genome. *PLoS genetics* **8**(11): e1003036.

Gelman A, Jakulin A, Pittau MG, Su Y-S. 2008. A weakly informative default prior distribution for logistic and other regression models. *The Annals of Applied Statistics* **2**(4): 1360-1383.

Genome of the Netherlands C. 2014. Whole-genome sequence variation, population structure and demographic history of the Dutch population. *Nature genetics* **46**(8): 818-825.

Giordano J, Ge Y, Gelfand Y, Abrusan G, Benson G, Warburton PE. 2007. Evolutionary history of mammalian transposons determined by genome-wide defragmentation. *PLoS computational biology* **3**(7): e137.

Guo J, Grow EJ, Yi C, Mlcochova H, Maher GJ, Lindskog C, Murphy PJ, Wike CL, Carrell DT, Goriely A et al. 2017. Chromatin and Single-Cell RNA-Seq Profiling Reveal Dynamic Signaling and Metabolic Transitions during Human Spermatogonial Stem Cell Development. *Cell stem cell* **21**(4): 533-546 e536.

Harpak A, Bhaskar A, Pritchard JK. 2016. Mutation Rate Variation is a Primary
Determinant of the Distribution of Allele Frequencies in Humans. *PLoS*
genetics **12**(12): e1006489.
Harris K, Nielsen R. 2014. Error-prone polymerase activity causes multinucleotide
mutations in humans. *Genome research* **24**(9): 1445-1454.
Heger A, Webber C, Goodson M, Ponting CP, Lunter G. 2013. GAT: a simulation
framework for testing the association of genomic intervals. *Bioinformatics*
29(16): 2046-2048.
Hodgkinson A, Eyre-Walker A. 2011. Variation in the mutation rate across
mammalian genomes. *Nature reviews Genetics* **12**(11): 756-766.
International HapMap Consortium Frazer KA Ballinger DG Cox DR Hinds DA Stuve LL
Gibbs RA Belmont JW Boudreau A Hardenbol P et al. 2007. A second
generation human haplotype map of over 3.1 million SNPs. *Nature*
449(7164): 851-861.
Jonsson H, Sulem P, Kehr B, Kristmundsdottir S, Zink F, Hjartarson E, Hardarson MT,
Hjorleifsson KE, Eggertsson HP, Gudjonsson SA et al. 2017. Parental influence
on human germline de novo mutations in 1,548 trios from Iceland. *Nature*
549(7673): 519-522.
Lee H, Schatz MC. 2012. Genomic dark matter: the reliability of short read mapping
illustrated by the genome mappability score. *Bioinformatics* **28**(16): 2097-
2105.
Lek M, Karczewski KJ, Minikel EV, Samocha KE, Banks E, Fennell T, O'Donnell-Luria
AH, Ware JS, Hill AJ, Cummings BB et al. 2016. Analysis of protein-coding
genetic variation in 60,706 humans. *Nature* **536**(7616): 285-291.
Li F, Tian L, Gu L, Li GM. 2009. Evidence that nucleosomes inhibit mismatch repair in
eukaryotic cells. *The Journal of biological chemistry* **284**(48): 33056-33061.
Lujan SA, Clausen AR, Clark AB, MacAlpine HK, MacAlpine DM, Malc EP, Mieczkowski
PA, Burkholder AB, Fargo DC, Gordenin DA et al. 2014. Heterogeneous
polymerase fidelity and mismatch repair bias genome variation and
composition. *Genome research* **24**(11): 1751-1764.
Makova KD, Hardison RC. 2015. The effects of chromatin organization on variation in
mutation rates in the genome. *Nature reviews Genetics* **16**(4): 213-223.
Michaelson JJ, Shi Y, Gujral M, Zheng H, Malhotra D, Jin X, Jian M, Liu G, Greer D,
Bhandari A et al. 2012. Whole-genome sequencing in autism identifies hot
spots for de novo germline mutation. *Cell* **151**(7): 1431-1442.
Perera D, Poulos RC, Shah A, Beck D, Pimanda JE, Wong JW. 2016. Differential DNA
repair underlies mutation hotspots at active promoters in cancer genomes.
Nature **532**(7598): 259-263.
Petryk N, Kahli M, d'Aubenton-Carafa Y, Jaszczyszyn Y, Shen Y, Silvain M, Thermes C,
Chen CL, Hyrien O. 2016. Replication landscape of the human genome.
Nature communications **7**: 10208.
Pich O, Muinos F, Sabarinathan R, Reyes-Salazar I, Gonzalez-Perez A, Lopez-Bigas N.
2018. Somatic and Germline Mutation Periodicity Follow the Orientation of the
DNA Minor Groove around Nucleosomes. *Cell* **175**(4): 1074-1087 e1018.
Prendergast JG, Semple CA. 2011. Widespread signatures of recent selection linked
to nucleosome positioning in the human lineage. *Genome research* **21**(11):
1777-1787.
Ramirez F, Dundar F, Diehl S, Gruning BA, Manke T. 2014. deepTools: a flexible
platform for exploring deep-sequencing data. *Nucleic acids research* **42**(Web
Server issue): W187-191.

Reijns MAM, Kemp H, Ding J, de Proce SM, Jackson AP, Taylor MS. 2015. Lagging-
strand replication shapes the mutational landscape of the genome. *Nature*
518(7540): 502-506.
Rodgers K, McVey M. 2016. Error-Prone Repair of DNA Double-Strand Breaks.
Journal of cellular physiology **231**(1): 15-24.
Sabarinathan R, Mularoni L, Deu-Pons J, Gonzalez-Perez A, Lopez-Bigas N. 2016.
Nucleotide excision repair is impaired by binding of transcription factors to
DNA. *Nature* **532**(7598): 264-267.
Salih F, Salih B, Kogan S, Trifonov EN. 2008. Epigenetic nucleosomes: Alu sequences
and CG as nucleosome positioning element. *Journal of biomolecular structure*
& dynamics **26**(1): 9-16.
Sasaki S, Mello CC, Shimada A, Nakatani Y, Hashimoto S, Ogawa M, Matsushima K,
Gu SG, Kasahara M, Ahsan B et al. 2009. Chromatin-associated periodicity in
genetic variation downstream of transcriptional start sites. *Science*
323(5912): 401-404.
Schuster-Bockler B, Lehner B. 2012. Chromatin organization is a major influence on
regional mutation rates in human cancer cells. *Nature* **488**(7412): 504-507.
Segurel L, Wyman MJ, Przeworski M. 2014. Determinants of mutation rate variation
in the human germline. *Annual review of genomics and human genetics* **15**:
47-70.
Seplyarskiy VB, Akkuratov EE, Akkuratova N, Andrianova MA, Nikolaev SI, Bazykin
GA, Adameyko I, Sunyaev SR. 2018. Error-prone bypass of DNA lesions
during lagging-strand replication is a common source of germline and cancer
mutations. *Nature genetics*.
Seplyarskiy VB, Andrianova MA, Bazykin GA. 2017. APOBEC3A/B-induced
mutagenesis is responsible for 20% of heritable mutations in the TpCpW
context. *Genome research* **27**(2): 175-184.
Shlien A, Campbell BB, de Borja R, Alexandrov LB, Merico D, Wedge D, Van Loo P,
Tarpey PS, Coupland P, Behjati S et al. 2015. Combined hereditary and
somatic mutations of replication error repair genes result in rapid onset of
ultra-hypermutated cancers. *Nature genetics* **47**(3): 257-262.
Slotkin RK, Martienssen R. 2007. Transposable elements and the epigenetic
regulation of the genome. *Nature reviews Genetics* **8**(4): 272-285.
Smerdon MJ. 1991. DNA repair and the role of chromatin structure. *Current opinion*
in cell biology **3**(3): 422-428.
Smith TCA, Arndt PF, Eyre-Walker A. 2018. Large scale variation in the rate of germ-
line de novo mutation, base composition, divergence and diversity in humans.
PLoS genetics **14**(3): e1007254.
Stamatoyannopoulos JA, Adzhubei I, Thurman RE, Kryukov GV, Mirkin SM, Sunyaev
SR. 2009. Human mutation rate associated with DNA replication timing.
Nature genetics **41**(4): 393-395.
Supek F, Lehner B. 2015. Differential DNA mismatch repair underlies mutation rate
variation across the human genome. *Nature* **521**(7550): 81-84.
Tanaka Y, Yamashita R, Suzuki Y, Nakai K. 2010. Effects of Alu elements on global
nucleosome positioning in the human genome. *BMC genomics* **11**: 309.
Tempel S. 2012. Using and understanding RepeatMasker. *Methods in molecular*
biology **859**: 29-51.
Terekhanova NV, Seplyarskiy VB, Soldatov RA, Bazykin GA. 2017. Evolution of Local
Mutation Rate and Its Determinants. *Molecular biology and evolution* **34**(5):
1100-1109.

Tolstorukov MY, Choudhary V, Olson WK, Zhurkin VB, Park PJ. 2008. nuScore: a
884   web-interface for nucleosome positioning predictions. *Bioinformatics* **24**(12):
885   1456-1458.
886   Tolstorukov MY, Volfovsky N, Stephens RM, Park PJ. 2011. Impact of chromatin
887   structure on sequence variability in the human genome. *Nature structural &*
888   *molecular biology* **18**(4): 510-515.
889   Tubbs A, Sridharan S, van Wietmarschen N, Maman Y, Callen E, Stanlie A, Wu W,
890   Wu X, Day A, Wong N et al. 2018. Dual Roles of Poly(dA:dT) Tracts in
891   Replication Initiation and Fork Collapse. *Cell* **174**(5): 1127-1142 e1119.
892   Turner TN, Coe BP, Dickel DE, Hoekzema K, Nelson BJ, Zody MC, Kronenberg ZN,
893   Hormozdiari F, Raja A, Pennacchio LA et al. 2017a. Genomic Patterns of De
894   Novo Mutation in Simplex Autism. *Cell* **171**(3): 710-722 e712.
895   Turner TN, Hormozdiari F, Duyzend MH, McClymont SA, Hook PW, Iossifov I, Raja A,
896   Baker C, Hoekzema K, Stessman HA et al. 2016. Genome Sequencing of
897   Autism-Affected Families Reveals Disruption of Putative Noncoding Regulatory
898   DNA. *American journal of human genetics* **98**(1): 58-74.
899   Turner TN, Yi Q, Krumm N, Huddleston J, Hoekzema K, HA FS, Doebley AL, Bernier
900   RA, Nickerson DA, Eichler EE. 2017b. denovo-db: a compendium of human de
901   novo variants. *Nucleic acids research* **45**(D1): D804-D811.
902   Veltman JA, Brunner HG. 2012. De novo mutations in human genetic disease. *Nature*
903   *reviews Genetics* **13**(8): 565-575.
904   Warnecke T, Becker EA, Facciotti MT, Nislow C, Lehner B. 2013. Conserved
905   substitution patterns around nucleosome footprints in eukaryotes and
906   Archaea derive from frequent nucleosome repositioning through evolution.
907   *PLoS computational biology* **9**(11): e1003373.
908   Werling DM, Brand H, An JY, Stone MR, Zhu L, Glessner JT, Collins RL, Dong S, Layer
909   RM, Markenscoff-Papadimitriou E et al. 2018. An analytical framework for
910   whole-genome sequence association studies and its implications for autism
911   spectrum disorder. *Nature genetics* **50**(5): 727-736.
912   West JA, Cook A, Alver BH, Stadtfeld M, Deaton AM, Hochedlinger K, Park PJ,
913   Tolstorukov MY, Kingston RE. 2014. Nucleosomal occupancy changes locally
914   over key regulatory regions during cell differentiation and reprogramming.
915   *Nature communications* **5**: 4719.
916   Winter DR, Song L, Mukherjee S, Furey TS, Crawford GE. 2013. DNase-seq predicts
917   regions of rotational nucleosome stability across diverse human cell types.
918   *Genome research* **23**(7): 1118-1129.
919   Yuen R, Merico D, Bookman M, J LH, Thiruvahindrapuram B, Patel RV, Whitney J,
920   Deflaux N, Bingham J, Wang Z et al. 2017. Whole genome sequencing
921   resource identifies 18 new candidate genes for autism spectrum disorder.
922   *Nature neuroscience* **20**(4): 602-611.
923   Yuen RK, Merico D, Cao H, Pellecchia G, Alipanahi B, Thiruvahindrapuram B, Tong X,
924   Sun Y, Cao D, Zhang T et al. 2016. Genome-wide characteristics of de novo
925   mutations in autism. *NPJ genomic medicine* **1**: 160271-1602710.

Data availability

All the analyses in this study were based on published datasets.

Acknowledgments

We are grateful to Tobias Warnecke, John Diffley, Anob Chakrabarti and Sara Rohban for insightful comments. We thank Peter Van Loo, Jonas Demeulemeester and Maxime Tarabichi for assistance in accessing the PCAWG genomic data. We appreciate obtaining access to the *de novo* mutation data on SFARI Base. This work is supported by the Francis Crick Institute which receives its core funding from Cancer Research UK (FC001110), the UK Medical Research Council (FC001110), and the Wellcome Trust (FC001110) (N.M.L.). N.M.L. is also supported by a Wellcome Trust Investigator Award and core funding from the Okinawa Institute of Science & Technology. C.L. is funded by an EMBO long-term postdoctoral fellowship (ALTF 1499-2016).

Author contributions

C.L. conceived the project, performed the analyses and drafted the manuscript; N.M.L. supervised the project and co-wrote the manuscript.

Competing financial interests

The authors declare no competing financial interests.

Supplementary Tables and Figures

Supplementary Table 1 Coefficients of variables and other parameters/results
from the full regression models for different mutation types (in a separate
Excel file). Note that for each of the categorial variables, the first category was used
by the regression model as reference category (other categories were compared with
the reference category) and thus there is no coefficient for that category. The
statistics (4[th] column) and p-values (5[th] column) in the table were from Wald tests
defaultly produced by 'bayesglm' (shown for reference), which are different from the
LRT-based p-values and were not used in our discussion.

Supplementary Figure 1 Mutations in different nucleosome contexts. (**a**) Information of the *de novo* mutation datasets from seven studies used in analysis. (**b**) Fold enrichment/depletion of gnomAD extremely rare SNVs in different nucleosome contexts. 'Strong', translationally stable positioning; 'Rotational', rotationally but not translationally stable positioning; 'Others', the remaining genomic regions. On the left is the fold enrichment for three subgroups of strong nucleosomes with different stabilities. (**c**) Fold enrichment/depletion of gnomAD INDELs in different nucleosome contexts. When using all INDELs the 'strong.high' group does not have a higher mutation rate than other two groups, but if using the 1-bp INDELs 'strong.high' does have the highest mutation rate among the three groups. We speculated that there may be more false negatives of longer INDELs in the 'strong.high' group.(**d**) Meta-profiles of SNV/INDEL densities (*de novo* or extremely rare variants) around all strong nucleosomes, or in different repeat-associated groups. At the bottom are the G+C content and CpG content profiles.

Supplementary Figure 2 Correlation analysis between nucleosome positioning stability (d_{var}) and other factors. On the top of each panel are the Pearson's correlation coefficients and the corresponding p-values.

977
978
979

Supplementary Figure 3 Results of statistical tests for nine individual SNV mutation types. C/G sites in non-CpG contexts and C/G sites in CpG contexts were tested separately. The red vertical lines represent the significance cut-off (0.05) for the adjusted p values (Bonferroni-Holm correction). 'us', upstream; 'ds', downstream. '#' means adjusted p < 1e-30.

Supplementary Figure 4 Results of statistical tests when considering d_{mean} as a continuous variable. (**a-b**) Statistical significance of each variable when performing likelihood ratio tests for assessing their contribution to mutation rate variation. Panel **a** for *de novo* SNVs and **b** for *de nvo* INDELs. For *de novo* SNVs, we did tests for A/T sites, C/G sites in non-CpG context and C/G sites in CpG context separately. The red vertical lines represent the significance cut-off (0.05) for the adjusted p values (Bonferroni-Holm correction). 'us', upstream; 'ds', downstream. '#' means adjusted p < 1e-30.

1005

Supplementary Figure 5 **Results of statistical tests when considering non-**
additive effects and 7-mer mutability estimates from Carlson et al. (**a**) Adding
the two-way interactions for ±5 nucleotides in the regression models. (**b**) Adding the
7-mer mutability estimates from Carlson et al. as predictors in the regression models.
We tested SNVs at A/T sites, C/G sites in non-CpG context and C/G sites in CpG
context separately. The red vertical lines represent the significance cut-off (0.05) for
the adjusted p values (Bonferroni-Holm correction). 'us', upstream; 'ds', downstream.
'#' means adjusted p < 1e-30.

1014

1015

1016

**Supplementary Figure 6 Analysis of related mutational processes using
bMMRD data.** (**a**) Mutation profiles around strong nucleosomes for bMMRD cancer
genomes and the estimated relative escape ratios of Pol ε or Pol δ, for mutations at
A/T sites and C/G sites respectively. Numbers of cancer genomes used are indicated
in the brackets. (**b**) Comparison of the contribution of COSMIC mutational signatures
predicted by MutationalPatterns in different bMMRD genomes. Highlighted is
Signature 12, which shows a particularly high contribution in POLD1-muated bMMRD
samples. (**c**) the tri-nucleotide mutational profile of Signature 12, obtained from
COSMIC website.

1026

1027

1028

1029

Supplementary Figure 7 Analysis with OK-seq data. (**a**) Schematic illustrating replication strands and Okazaki junctions (OJs). (**b**) Meta-profile of the density of Okazaki junctions inferred from alignments of OK-seq reads around strong nucleosomes (high-mappability). OJ signals for Watson strand and Crick strand were plotted separately. Replication directions of Okazaki fragments are shown by arrows.

1035

1036

1037

Supplementary Figure 8 Analysis related to the DSBs around strong
nucleosomes. (**a**) Density of poly(dA:dT) tracts (based on occurrence of $(dA:dT)_6$
motifs) around strong nucleosomes. (**b-c**) Signal of DSBs based on the END-seq
data around strong nucleosomes associated with different repeat elements. Only the
strong nucleosomes of high 75-mer mappability within ±500bp were considered.
Numbers of usable strong nucleosomes for each group are given in the brackets. HU
(hydroxyurea) is a replicative stress-inducing agent.

1045

Supplementary Figure 9 Densities of strong nucleosome dyads and *de novo*
SNVs along the consensus sequences of 3' ends of different L1 subfamilies. (**a**)
Alignment of the consensus sequences of the 3' end sequences of three L1
subfamilies (from young to old: L1PA5, L1PA10 and L1PA15), indicating the high
similarity between sequences. (**b-d**) Barplots for normalized densities of strong
nucleosome dyads and *de novo* SNVs along the consensus sequences of three L1
subfamilies, using 10-bp bins. Several loci that are enriched for dyads of strong
nucleosomes are shown on the top with ellipses. The red dash lines represent the
average densities for the L1PA5 subfamily. As shown in the **b-d**, the densities of
strong nucleosome dyads and *de novo* SNVs decreased over evolutionary time. (**e**)
Meta-profile for the estimated deformation energy around the strong nucleosomes of
three L1 subfamilies, centered at the dyads.

1058