<u>Region-specific regulation of stem cell-driven regeneration in tapeworms</u>

Tania Rozario[1*], Edward B. Quinn[1], Jianbin Wang[2], Richard A. Davis[2], and Phillip A.

Newmark[1,3,4]*

[1]Morgridge Institute for Research, Madison, WI, USA, [2]Department of Biochemistry and

Molecular Genetics, RNA Bioscience Initiative, University of Colorado School of Medicine,

Aurora, CO, USA, [3] Howard Hughes Medical Institute, [4]Department of Integrative Biology,

University of Wisconsin-Madison, Madison, WI, USA.

* corresponding authors

Email: trozario@morgridge.org

13   pnewmark@morgridge.org

Abstract

Tapeworms grow at rates that rival all metazoan tissues, including during embryonic and

neoplastic growth. For example, the rat intestinal tapeworm, *Hymenolepis diminuta*, produces up

to 2,200 proglottids (segments), increasing in length up to 3,400 fold, and weight up to 1.8 million

fold within the first 15 days of infection[1]. Tapeworms also regenerate: they shed large parts of

their body, releasing their embryos to continue their life cycle, yet are able to continuously

replenish proglottids and maintain an equilibrium length. Such remarkable growth and

regeneration are fueled by adult somatic stem cells, which have yet to be characterized

molecularly. Using *H. diminuta* as a laboratory model, we find that regeneration is limited to the

tapeworm neck, making this tissue a prime source to identify stem cell genes. Using transcriptomic

analyses and RNA interference (RNAi), we characterize and functionally validate the first

molecular regulators of tapeworm growth and regeneration. However, we find no evidence that

stem cells are restricted to the regeneration-competent neck. Instead, we find that lethally irradiated

tapeworms can be rescued from death when cells from both regeneration-competent and

regeneration-incompetent regions are transplanted into the neck. Furthermore, the persistence of

regenerative ability over time requires signal(s) from the head, even though the head itself cannot

regenerate and the head is not necessary for initial regeneration. Together, the head and neck tissue

provide a microenvironment that regulates stem cells to enable region-specific regeneration in this

tapeworm.

Introduction

Tapeworms are parasitic flatworms that infect humans and livestock, causing lost economic output, disease, and in rare cases, death[2]. Tapeworms are well known for their ability to reach enormous lengths. For example, humans infected with the broad or fish tapeworm, *Diphyllobothrium latum*, harbor parasites that average 6 m in length[3]. It is less commonly appreciated that tapeworms regenerate to accommodate their multi-host life cycle. Adult tapeworms in their host intestine develop proglottids that are gravid with embryos. Tapeworms pinch off the posterior and gravid sections of their body, which exit with the host excrement, to be eaten by a suitable intermediate host that supports larval tapeworm development. In spite of losing large body sections, the tapeworm does not progressively shorten; instead, it regenerates proglottids to maintain an equilibrium length. Despite this remarkable biology, tapeworms are an unexplored animal model in the study of regenerative behaviors.

Up until the 1980s, the rat intestinal tapeworm, *H. diminuta*, had been a favorite model organism among parasitologists because these parasites are easily propagated in the laboratory. A great deal of work in cell biology, biochemistry, physiology, and electron microscopy has enriched our understanding of these tapeworms[4]. With the dawn of the molecular age and the preference among scientists to concentrate on a handful of model organisms, *H. diminuta* was largely left behind.

As an obligate endoparasite, adult *H. diminuta* will expire once its host rat dies. However, the lifespan of *H. diminuta* can be uncoupled from that of its host. A single adult tapeworm can be serially amputated and transplanted into a new host intestine, where the fragment can regenerate into a mature tapeworm, even after 13 rounds of amputation over 14 years[5]. These observations

have led to speculation that *H. diminuta* may be inherently immortal, though their life span is

limited by that of their host. This longevity is reminiscent of the free-living cousins of tapeworms:

freshwater planarians like *Schmidtea mediterranea.* Asexual *S. mediterranea* can reproduce

indefinitely by fission, and can regenerate its whole body from tiny fragments. Planarian

immortality and regeneration are enabled by adult somatic stem cells called neoblasts[6-8]. These

stem cells are the only dividing cells within the soma. Like planarians, *H. diminuta* maintains a

population of neoblast-like adult somatic stem cells[9] that are likely responsible for their growth

and regenerative ability. Recently, stem cells of other parasitic flatworms have been defined

molecularly, and their maintenance depends on conserved factors such as argonaute family

members and fibroblast growth factor signaling[10-12]. However, key differences also exist, as a

number of conserved stem cell genes, like *piwi, vasa,* and *tudor,* are absent from the genomes of

parasitic flatworms[13].

Mechanisms of stem cell regulation in parasitic flatworms are still largely unexplored,

though stem cells are crucial for parasite biology and even disease. For example, the tapeworm

Echinococcus multilocularis causes alveolar echinococcosis, in which larvae use their stem cells

to grow as infiltrative cysts that can cause organ failure, and like a malignant tumor, can

metastasize to secondary organs[14]. A better understanding of stem cell regulation in parasitic

flatworms is needed to elucidate parasite biology and potentially, for gaining therapeutic insights.

In this study, we use *H. diminuta* to investigate the molecular basis of tapeworm

regeneration. We established and refined experimental tools such as transcriptomics, *in vitro*

parasite culture, RNA *in situ* hybridization, cycling cell tracing with thymidine analogs, and RNA

interference (RNAi), all reported in this work. We determine that the ability to regenerate is

regionally limited to the neck of adult *H. diminuta*. However, regeneration from the neck is finite

without signals from the tapeworm head. Using RNA sequencing (RNA-seq), we identify and

characterize various markers of the somatic cycling cell population, which includes tapeworm stem

cells. We functionally validate essential regulators of growth and regeneration. Intriguingly, our

analyses failed to uncover a neck-specific stem cell population that explains the regional

regenerative ability displayed by *H. diminuta.* Instead, we demonstrate that tapeworm regeneration

can be rescued by cells that are sourced from outside the neck, regardless of source-tissue

regeneration competence. Thus, extrinsic signals present in the tapeworm neck confer region-

specific regenerative ability in this tapeworm.

Results and Discussion

To examine the regeneration competence of *H. diminuta*, we amputated different anatomical fragments cut from the adult and assayed their ability to regenerate. The body plan of adult *H. diminuta* consists of a head with four suckers, an unsegmented neck (~2.5-3 mm in 6-day old worms), and a body with thousands of proglottids that grow and mature in an anterior-to-posterior direction (Fig. 1a). Previous work showed that amputated *H. diminuta* anterior fragments transplanted into rat intestines were able to regenerate into mature tapeworms[5,15]. However, using transplantation, a complete study of regeneration competence was not possible: only anterior fragments could be tested because the suckers are required to anchor the worms to the intestine. To ascertain which anatomical regions of *H. diminuta* are competent to regenerate, amputated fragments must be grown *in vitro*. We successfully established *H. diminuta in vitro* culture modified from the Schiller method[16], and tested the regeneration competence of several different 1 cm fragments (Fig. 1b-c). The anterior-most fragments (head + neck + body) were competent to regenerate, confirming previous observations using amputation and transplantation[5,15]. Additionally, anterior fragments that were first decapitated (neck + body) were also competent to regenerate proglottids. In contrast, "body only" fragments that did not contain the neck tissue were not competent to regenerate proglottids. All amputated fragments could grow in length (Fig. 1d), differentiate mature reproductive structures, and mate. However, the neck is necessary for regeneration of new proglottids over time. In no case did we observe head regeneration. Also, amputated heads alone could not regenerate proglottids *in vitro* (data not shown) nor *in vivo*[15]. To test if the neck is sufficient for proglottid regeneration, we amputated 2 mm fragments of the neck tissue and grew the fragments *in vitro*. "Neck only" fragments regenerated an average of 383

proglottids (SD=138, n=20) after 12 days *in vitro* (Fig. 1e). Thus, the tapeworm neck is both

necessary and sufficient for proglottid regeneration.

Previous studies have shown that *H. diminuta* can regenerate after serial rounds of

amputation and transplantation for over a decade[5] and perhaps indefinitely. Using *in vitro* culture,

we confirmed that anterior fragments of *H. diminuta* can regenerate proglottids after four rounds

of serial amputation followed by 12-15 days of *in vitro* culture (Fig. 1f-g) without any observable

cessation of regenerative ability. Decapitated fragments were competent to regenerate proglottids

after the first amputation; however, re-amputation abrogated regeneration (Fig. 1f-g). After

decapitation, a definitive neck tissue could not be maintained and eventually, the whole tissue was

comprised of proglottids (Fig. S1). Without the head, proglottid regeneration from the neck is

finite. Thus, the tapeworm head is necessary for persistent proglottid regeneration and to maintain

an unsegmented neck.

If signals from the head regulate regeneration, is regenerative potential asymmetric across

the anterior-posterior (A-P) axis of the neck? We subdivided the neck into three 1 mm fragments

and assayed the number of proglottids regenerated *in vitro* (Fig. 1h-i). We found that the most

anterior neck fragments regenerated more proglottids than the middle or posterior neck fragments

(Fig. 1h-i). Thus, only the neck is competent to regenerate proglottids, but signals from the head

also regulate regeneration, resulting in an anterior bias in regenerative potential.

Why is the neck the only region competent to regenerate? One hypothesis is that pluripotent

stem cells are confined to the neck. There are currently no specific molecular markers for adult

tapeworm stem cells. However, like in other flatworms, the only proliferative somatic cells in *H.*

diminuta are undifferentiated[17,18]. Thus, cycling somatic cells include tapeworm stem cells and

any dividing progeny. To label cycling somatic cells, we used two methods: uptake of the

thymidine analog F-ara-EdU to mark cells in S-phase and fluorescent RNA *in situ* hybridization

(FISH) against conserved cell cycle-regulated transcripts such as the replication licensing factor

minichromosome maintenance complex component 2 (*mcm2*) and *histone h2b* (*h2b*). Both *h2b* and

mcm2 are conserved cycling cell markers expressed in stem cells of free-living and parasitic

flatworms[10,19]. We detected cycling somatic cells throughout the tapeworm body (Fig. S2a-c).

Contrary to previous results[17], we also detected cycling cells in the head, though in small numbers.

The scarcity of these cycling cells may be the reason they were originally missed. Taken together,

cycling cells are present in tissues that are competent or incompetent to regenerate proglottids.

Though cycling cells are widely distributed, it is possible that within this population, there

are stem cells of varying potency and that a pluripotent subpopulation may be restricted to the

neck. To identify potential stem cell markers, we characterized the cycling cell population at the

molecular level. Previously, stem cell genes had been identified in flatworms using irradiation to

deplete cycling cells followed by transcriptome analysis to identify downregulated

transcripts[10,19,20]. In *H. diminuta*, cycling cells are undetectable after exposure to 200 Gy irradiation

(Fig. S2d). Consequently, growth and regeneration were also stunted following irradiation (Fig.

2a-b). We leveraged the sensitivity of *H. diminuta* to irradiation to identify new molecular markers

of cycling somatic cells, including stem cells. Untreated and irradiated worms were cultured *in*

vitro for 3 days, after which the necks were amputated and processed for RNA-seq (Fig. 2c). We

assembled a *de novo* transcriptome for *H. diminuta* consisting of 14,346 transcripts (refer to

methods) to which sequencing reads were mapped. Using differential gene expression analysis,

we identified 683 irradiation-sensitive transcripts (downregulated; FDR $\leq$0.05) (Supplemental

Table 1), cloned 335 genes, and examined their expression patterns using whole-mount

colorimetric *in situ* hybridization (WISH). Using WISH, we observed that expression of

irradiation-sensitive genes was indeed reduced after exposure to irradiation, validating our RNA-

seq approach (Fig. S3).

In tapeworms, the position of cycling cells in the neck is spatially restricted in a pattern

that is conserved in multiple tapeworm species[21] (Fig. 2d-e). Cycling cells are located in the neck

parenchyma, which is bounded by the nerve cords. Cycling cells are not found at the animal edge

where differentiated cells of the muscle and tegument (parasite skin) are located[17]. This pattern

allowed us to use WISH to identify potential cycling cell genes. After excluding genes with

ubiquitous expression or poor signal by WISH, we examined gene expression for 194 irradiation-

sensitive genes. 62% of genes examined were expressed in the neck parenchyma, though in a

variety of patterns (Fig. S4a). 13% of genes examined were predominantly expressed in the neck

parenchyma, similar to *h2b* and *mcm2* (Fig. 2f, S4b). These include the nucleic acid binding factor

Zn finger MYM type 3 (*zmym3)*, the transcription factor *suppressor of hairy wing* (*su(Hw)*)*,* nuclear

lamina component *laminB receptor* (*lbr*), and *leucine-rich WD-containing gene* (*lrwd*). 25% of

irradiation-sensitive genes, were expressed in a minority of cells in the neck parenchyma (Fig.

S4c). 24% were expressed broadly, including within the neck parenchyma and at the animal edge

where non-cycling differentiated cells are located (Fig. S4d). In conclusion, irradiation-sensitive

genes identified by RNA-seq likely represent markers for stem cells, progenitors, and even

differentiated cells that were lost or compromised following irradiation.

To ascertain whether genes that showed expression in the neck parenchyma were indeed

expressed in cycling cells, we performed double fluorescent *in situ* hybridization (dFISH) with

candidate genes and either *h2b* or *mcm2*. We use *h2b* and *mcm2* interchangeably as they

completely co-localize in the neck parenchyma (Fig. S5). Irradiation-sensitive genes from Fig. 2f

were indeed expressed in cycling somatic cells (Fig. 2g-h, S6a-b). One gene, the homeobox factor,

prospero (*pros*), was expressed exclusively in a subset of cycling cells (Fig. 2i). We confirmed

that irradiation-sensitive genes that only partially overlapped in the neck parenchyma, such as the

Zn finger-containing genes *HDt_10981* and *HDt_05316* were expressed in a subset of cycling cells

as well as non-cycling differentiated cells (Fig. S6c-d). We propose that these genes represent

lineage-committed stem cells or progenitors for tissues such as muscle, neurons, tegument, or

protonephridia. In summary, we performed dFISH for 53 candidates and found that 72% of genes

tested were co-expressed in cycling cells (Fig. S6e, Supplemental Table 2). Our analysis revealed

that there is a heterogeneous and complex mixture of cell types or states within the cycling cell

population.

We reasoned that genes expressed in the majority of cycling cells would likely be essential

for tapeworm growth and may regulate stem cell populations that are necessary for regeneration.

Thus, we developed an RNAi protocol that allowed us to functionally investigate how candidate

genes affect tapeworm growth and regeneration (Fig. 3a). As a proof of principle, we used RNAi

to knock down *h2b*. Stem cells are constantly dividing and require new histone synthesis to

maintain the population. Thus, downregulating expression of *h2b* in tapeworms would likely

compromise growth as a result of cycling cell loss, as is the case in other flatworms[19,22]. In our

RNAi paradigm, knockdown of *h2b* induced gross growth defects (Fig. 3b-c). The number of

proglottids regenerated was also reduced; however, this was difficult to quantify as many RNAi

worms were so thin and frail that proglottid definition was lost. We then tested two newly

identified cycling cell genes: *zmym3,* and *su(Hw)* for potential roles in tapeworm growth and

regeneration. Similar to knockdown of *h2b, zmym3(RNAi)* and *su(Hw)(RNAi)* each resulted in

abrogated growth and regeneration. RNAi-mediated knockdown has been demonstrated in other

tapeworm species[23-26], though not in *H. diminuta.* RNAi has not been widely adopted for studying

tapeworm biology due to technical challenges, like poor knockdown efficacy, inefficient

penetrance, and the difficulty of *in vitro* culture. Our RNAi scheme is relatively straightforward

and robust, and shows that gene inhibition can be used to uncover regulators of adult tapeworm

growth and regeneration.

Are the growth defects we observe the consequence of defects in the cycling cell

population? After RNAi, we quantified the number of cycling cells present by detecting F-ara-

EdU[+] cells following 1-hour exposure to the thymidine analog. Knockdown of *h2b, zmym3,* and

su(Hw) severely reduced the number of cycling cells detectable in the neck (Fig. 3d-e). Thus, *h2b,*

zmym3, and *su(Hw)* are necessary for the function of cycling cells.

The mechanism of action by *zmym3* and *su(Hw)* in *H. diminuta* is currently unknown. In

mice and humans, *zmym3* acts as a transcriptional repressor with chromatin-binding properties and

has been shown to regulate cell cycle progression[27] and DNA repair[28]. In *Drosophila*, *su(Hw)* binds

to *gypsy* insulator sequences to regulate chromatin silencing[29]. Similar functions for these genes

may be conserved in tapeworms as cell cycle regulation, DNA repair, and silencing are likely

important for the regulation of tapeworm stem cells.

We have now described molecular heterogeneity within the cycling cells of the neck

parenchyma and identified *h2b, zmym3,* and *su(Hw)* as functionally important in cycling cell

regulation. At this point, we have not yet determined why regeneration competence is restricted to

the neck. Are subsets of cycling cells distributed in a polarized manner? No gene expression pattern

examined was exclusive to cycling cells of the neck. By WISH, all cycling cell genes were

expressed throughout the whole tapeworm body, including *zmym3* and *su(Hw)* (Fig. 4a). Since we

observed an anterior bias in regenerative ability (Fig. 1h-i), we hypothesized that stem cells may

be asymmetrically distributed in the neck and detectable by RNA-seq. We performed RNA-seq of

1 mm anterior, middle, and posterior neck fragments (Fig. 1h) and identified 454 anterior-enriched

and 238 anterior-depleted transcripts (Fold change ≥1.5 or ≤-1.5 respectively; FDR ≤0.05) and

overlaid them with our irradiation-sensitive dataset. The majority of anterior-enriched transcripts

(88%) were not irradiation sensitive (Fig. S7a). Instead, we observed that anterior-enriched and

anterior-depleted genes were often expressed in corresponding gradients (Fig. 4b) but these

patterns were not within the neck parenchyma. Our results suggest that the polarized signals across

the neck tissue are predominantly within the non-cycling compartments.

Since our RNA-seq analysis identified transcripts that were anterior-enriched and

irradiation-sensitive (Fig. S7a), we examined their expression patterns to ascertain if these genes

are expressed within stem cells. We found 15 genes that are expressed in a subset of cells within

the neck parenchyma, including *pros*, *solute carrier family 13* (*slc13*), *hepatocyte nuclear factor*

4 (*hnf4*)*,* and several novel genes (Fig. S7b). However, by dFISH with *mcm2*, 7/8 genes tested

were not expressed in cycling cells (Fig. S7c). Only one gene, *pros,* was co-expressed in cycling

cells (Fig. 2i). Our analyses have not revealed an anteriorly biased subpopulation of stem cells that

confer regenerative ability. This may be due to the coarse resolution of our analysis. A more

thorough characterization of tapeworm cycling cells by single-cell RNA-seq is warranted.

Nonetheless, the variety of cycling cell markers we have described in this work will aid in future

classification of tapeworm stem cells.

So far, we have not found evidence for a unique neck-specific subpopulation of cycling

cells. This led us to hypothesize that stem cells may be distributed throughout the tapeworm, but

that extrinsic signals operating in the neck are necessary to instruct stem cell behavior and/or

proglottid regeneration. To pursue this hypothesis, we designed a functional assay to test

populations of cells for the ability to rescue regeneration. We exposed tapeworms to a lethal dose

of irradiation, injected wild-type cells into the neck region, amputated 5 mm anterior fragments,

and looked for rescue of viability and regeneration after 30 days *in vitro*. Remarkably, bulk-cell

transplants were able to rescue irradiated worms that were destined to die (Fig. 4c; 44% rescue

(n=41)). In no case did we observe rescue of viability or regeneration in irradiated worms that were

not injected with cells or mock-injected (n=34). With this functional assay in hand, we compared

cells from tapeworm anteriors (including the regeneration-competent neck) to cells from tapeworm

posteriors (which are regeneration incompetent) for rescue ability. We found that cells from both

regeneration-competent and regeneration-incompetent regions of *H. diminuta* were able to recue

viability and regeneration in lethally irradiated tapeworms (Fig. 4d; 11% rescue for both A and P

(n=37 and 35, respectively)). These results confirm that the regeneration competence of the neck

tissue is not due exclusively to intrinsic properties of stem cells in this region, but to extrinsic

signals that regulate competence to regenerate.

Our results suggest that in tapeworms, location matters enormously: the head and neck

environment provide cues that regulate the ability of stem cells to mediate proglottid regeneration.

We show that the head plays instructive roles that regulate how proglottids form and regenerate.

Signals from the head may create gradients across the neck that inhibit proglottidization and are

necessary to maintain the neck as an unsegmented tissue. Proglottids can only form once the

inhibitory signals are sufficiently diminished (Fig. 4e). How the head regulates stem cells is less

clear. The head may serve as a niche (directly or indirectly) that is necessary for the maintenance

of stem cells in the neck (Fig. 4e). In this model, stem cells are collectively pluripotent only when

they receive head-derived niche signals, thus limiting regenerative potential to the neck.

Alternatively, stem cells may depend on a local niche that is independent of the head (Fig. 4e). In

this model, stem cells have the capacity to form all cell lineages from any amputated fragment,

however, the extrinsic signals that activate proglottid formation are only found in the neck. The

interplay between intrinsic and extrinsic stem cell regulatory signals has already been shown to

play important roles in regeneration. Head regeneration was induced in three planarian species that

were naturally regeneration deficient by manipulating the gradient of wnt signaling by RNAi[30-32].

These planarians maintained pluripotent stem cells but could not regenerate heads because of

inhibitory cues from the surrounding tissue. In tapeworms, gradients of wnt signaling that delineate

A-P polarity are conserved with planarians[33]. RNA-sequencing of the neck A-P axis in *H. diminuta*

has revealed genes that are expressed in gradients across the neck, which could help us understand

how stem cells and regeneration depend on these extrinsic signals.

Materials and Methods

Animal care and use

Infective *H. diminuta* cysts were obtained from Carolina Biologicals (132232). To obtain adult tapeworms, 100-300 cysts were fed to Sprague-Dawley rats by oral gavage in ~0.5 mL of 0.85% NaCl. Rats were euthanized in a CO_2 chamber typically 6 days post-gavage and tapeworms were flushed out of the small intestine and washed in 1X Hanks Balanced Salt Solution (HBSS; Corning) (140 mg/L $CaCl_2$, 100 mg/L $MgCl_2.6H_2O$, 100 mg/L $MgSO_4.7H_2O$, 400 mg/L KCl, 60 mg/L KH_2PO_4, 350 mg/L $NaHCO_3$, 8 g/L NaCl, 48 mg/L Na_2HPO_4, 1 g/L D-glucose, no phenol red). Rodent care was in accordance with IACUC policies and protocols.

In vitro parasite culture

Biphasic parasite cultures were prepared based on the Schiller method[16]. Briefly, the solid phase was made in 50 mL Erlenmeyer flasks by mixing 30% heat-inactivated defibrinated sheep blood (Hemostat) with 70% agar base for 10 mL blood-agar mixture per flask. Fresh blood was heat-inactivated at 56°C for 30 min then kept at 4°C and used repeatedly for one week by first warming the blood to 37°C. The agar base was prepared from 8 g Difco nutrient agar and 1.75 g NaCl in 350 mL water, autoclaved, and stored at 4°C. Before use, the agar base was microwaved to liquify, and cooled to below 56°C before mixing with warmed blood. After the blood-agar mixture solidified, 10 mL of Working Hanks 4 (WH4; 1X HBSS/4 g/L total glucose/1X antibiotic-antimycotic (Sigma)) was added. Each flask was topped with a gas-permeable stopper (Jaece Identi-plug) and pre-incubated at 37°C in hypoxia (3% CO_2/5% O_2/95% N_2) overnight before use. Before tapeworms were transferred into the flasks, the liquid phase was adjusted to pH7.5 with

200 μL 7.5% NaHCO₃ (Corning). Tapeworms were first washed in WH4 for 10 mins at 37°C in

petri dishes pre-coated with 0.5% BSA to inhibit sticking. Transfers to pre-cultured flasks were

performed by gently lifting the worms with a stainless-steel hook (Moody Tools) and immersing

them in the liquid phase. Tapeworms were transferred to fresh cultures every 3 days.

<u>Fixation and Staining</u>

Tapeworms were heat-killed by swirling in 75-80°C water for a few seconds until the

worms relaxed and elongated, then fixative (4% formaldehyde/10% DMSO/1% NP40 in

Phosphate Buffered Saline with 0.3% TritonX-100 (PBSTx)) was added immediately. Tapeworms

were fixed for 30 min-2hr at room temperature or overnight at 4°C. For DAPI staining, samples

were incubated in 1 μg/mL DAPI (Sigma) in PBSTx overnight at 4°C. Stained tapeworms were

cleared in 80% glycerol/10 mM Tris pH7.5/1 mM EDTA overnight at room temperature before

mounting.

<u>Thymidine analog uptake and staining</u>

For BrdU pulse, tapeworms were incubated in 10 μM BrdU (Sigma) in 1% DMSO at 37°C

in WH4 and processed for staining by an adapted protocol previously used on planarians[34] and

schistosomes[10]. Tapeworms were killed in cold 2% HCl for 2 min with shaking every 30 seconds,

fixed in cold methacarn (6:3:1 methanol:chloroform:glacial acetic acid) for 30 min at 4°C.

Denaturation was performed in 2N HCl for 45 min. Blocking and antibody incubation were

performed in K-block (5% Horse serum/0.45% fish gelatin/0.3% Triton-X/0.05% Tween-20/PBS)

[35], before detection with anti-BrdU antibody (1:500, B35128, Invitrogen) and anti-mouse-HRP

secondary antibody (1:1000; Jackson). Tyramide signal amplification (TSA) reaction[36] was also

used.

For F-ara-EdU pulse, tapeworms were incubated in 0.1 μM F-ara-EdU (Sigma) in 1%

DMSO for 1 hr at 37°C in WH4. Tapeworms were heat-killed, fixed in 4% formaldehyde/10%

DMSO/1% NP40/PBSTx, permeabilized by treatment with 10 μg/mL Proteinase-K/0.1%

SDS/PBSTx for 10-30 min at room temperature, fixed in 4% formaldehyde/PBSTx for 10 min

before samples were cut into small pieces or retained whole in PBSTx. Samples were

permeabilized in PBSTx/10% DMSO/1% NP40 for 20min-1 hr (depending on size) before

performing the click-it reaction[37] with Oregon Green 488 Tyramide (Invitrogen). Signal was

detected using anti-Oregon Green 488-HRP antibody (1:1000; A-21253 Invitrogen) in K-block

followed by TSA reaction. For F-ara-EdU quantification, tiled confocal z-stacks through the

anterior of the worms were taken and cell numbers were counted using background subtraction on

Imaris.

Irradiation

Irradiation was performed using a CellRad irradiator (Faxitron Bioptics) at 200 Gy (150

kV, 5 mA). Tapeworms were irradiated in WH4 in BSA-coated petri dishes. At this dosage,

tapeworms degenerated and were inviable after 30 days in culture.

Transcriptome assembly

To construct a transcriptome for *H. diminuta,* we collected RNA from five regions: 1) head

and neck, 2) immature proglottids, 3) mature reproductive proglottids, 4) gravid proglottids, and

5) mixed larval stages isolated from beetles. The first three regions covered the entirety of 3.5-

week old adult tapeworms. The gravid proglottids were taken from the posteriors of 10-week old

tapeworms. Paired-end libraries were constructed with 2 x 150 bp reads from a HiSeq2500 chip.

2 x ~30 million reads were obtained for each sample. The transcriptome was assembled from three

components: 1) map-based assembly, 2) *de novo* assembly, and 3) Maker predictions from

Wormbase Parasite. The map-based assembly was performed using TopHat2 with the 2014 *H.*

diminuta draft genome courtesy of Matt Berriman at the Wellcome Sanger Institute. 15,859

transcripts were assembled using TopHat. *De novo* assembly was performed using Velvet/Oases

and resulted in 144,682 transcripts. For the Maker predictions, there were 11,275 predicted

transcripts and 73.2% matched (>95% along the length) to the TopHat transcripts. The remaining

predicted transcripts that were not represented in the TopHat dataset were added to make a

combined TopHat/predicted set of 17,651 transcripts. Most of the Oases transcripts matched to the

TopHat/predicted set but 35,300 or 24.4% of the Oases transcripts did not (>75% match cut-off).

These transcripts could be transcripts missed in the genome, transcription noise, non-coding

transcripts, or contamination. We found significant contamination from beetle tissue in the larval

tapeworm sample (more below). Initial filtering for contamination excluded 1,388 transcripts

(from beetle, rat, bacterial, and viral sources). At this point 51,563 transcripts were retained from

the three methodologies described above and were processed for further filtering.

We first filtered out additional contamination from beetle tissues. We expect that this was

a large problem because during dissections, beetle tissues adhered to the tapeworm larvae and were

not easily washed away. These transcripts had best hits to beetle proteins (from *Ixodes scapularis,*

Harpegnathos saltator, Monodelphis domestica, Nasonia vitripennis, Pediculus humanus

corporis, Solenopsis invicta, Tenebrio molitor, and *Tribolium castaneum*). Most of the transcripts

were from the Oases *de novo* assembly and did not match the *H. diminuta* genome. Furthermore,

they were strongly over-represented in the larval sample only. Thus, an additional contamination

filtering step was taken: transcripts from the Oases assembly without matches to the *H. diminuta*

genome that showed >90% expression (by RPKM) in the larval sample were filtered out as

potential contaminants. 11,918 transcripts were filtered out in this step.

To the remaining 39,645 transcripts, we applied additional filters: 1) Remove transcripts if

average RPKM<1 unless the transcript is long (>1000 bp), has a long ORF (>500 bp) or is

annotated. 11,615 transcripts were removed as they met none of these criteria. 2) A stringent

expression cut-off was applied to the remaining Oases transcripts; transcripts were discarded if

average RPKM<5 and maximum RPKM<10 unless the transcripts were long (>1000 bp), had long

ORFs (>500 bp) or were annotated. 8,027 transcripts were removed. 3) 51 transcripts were

removed because they are mitochondrial or rRNAs. 4) An ORF size filter was applied to remove

all transcripts with ORF <300 bp unless they are annotated. 5,331 transcripts were removed. 5)

For the Maker predicted transcripts, expression and size filters were applied to remove transcripts

with expression <1 RPKM and size <500 bp. 275 transcripts were removed.

Our final transcriptome is comprised of 14,346 transcripts (84.9% TopHat, 8.4% Maker

predictions, 6.1% Oases with match to genome, and 0.6% Oases without match to genome). The

total transcriptome size is 34 Mb with average transcript length of 2,354 bp.

<u>RNA-seq for differential gene expression analyses</u>

Tissue was collected and immediately frozen on dry ice in 100 μL Trizol (Life

Technologies) before RNA extraction. Tissue homogenization was performed as the mixture was

in a semi-frozen state using RNase-free pestles and a pestle motor. RNA was purified using the

Direct-zol RNA MiniPrep kit (Zymo). RNA quality was assessed using Bioanalyzer, libraries were

prepared with TruSeq Stranded mRNAseq Sample Prep kit (Illumina), and sequenced on two lanes

on a HiSeq2500 chip. We performed paired-end sequencing and obtained ~20 million reads per

sample. Samples were obtained in triplicate. To identify irradiation-sensitive genes, 2 mm anterior

tapeworm fragments were cut from 10 worms after 3 days *in vitro*. To identify differentially

expressed transcripts across the neck A-P axis, 1 mm fragments were cut from 20 freshly obtained

6-day old tapeworms. Differential gene expression analysis was performed using CLC Genomics

Workbench (Qiagen) by mapping to our assembled transcriptome (above).

In situ hybridization

WISH and FISH protocols were modified from previously published methods for

planarians[36] and the mouse bile-duct tapeworm *Hymenolepis microstoma* (Peter Olson, personal

communication). Tapeworms were heat killed and fixed in 4% formaldehyde/10% DMSO/1%

NP40/PBSTx for 30 min at room temperature before washing and dehydration into methanol.

Dehydrated samples were kept at -30°C for at least 2 days. After rehydration, samples were

permeabilized in 10 µg/mL Proteinase-K/0.1% SDS/PBSTx, washed into x 0.1 M Triethanolamine

pH7-8 (TEA), 2.5 µL/mL acetic anhydride was added for 5 min with vigorous swirling, acetic

anhydride step was repeated, washed in PBSTx and post-fixed in 4% formaldehyde/PBSTx for 10

min. Probe synthesis, hybridization, and staining were performed as previously described[36] using

probe concentrations at ~50 ng/mL for 16-48 hrs at 56°C. All probes were synthesized with either

DIG or DNP haptens and detected using the following antibodies, all at 1:2000: anti-DIG-AP

(Sigma), anti-DIG-POD (Roche), anti-DNP-HRP (Vector Labs). Colorimetric development was

done using NBT (Roche)/BCIP (Sigma) or with Fast-Blue (Sigma)[38]. Fluorescent signal was

visualized after TSA reaction[36]. DAPI staining and mounting were performed as described above.

Microscopy

420   Confocal imaging was performed on a Zeiss LSM 880 with the following objectives: 10X
421 PlanNEOFLUOR air, NA 0.3/ 20X Plan-APOCHROMAT air, NA 0.8/ 40X C-APOCHROMAT
422 oil, NA 1.3/ 63X Plan-APOCHROMAT oil, NA 1.4. WISH samples and whole-mount DAPI-
423 stained worms were imaged using a Zeiss AxioZoom V16. Image processing was performed using
424 ImageJ.

RNAi

Double-stranded RNA (dsRNA) was synthesized as previously described[39] and
resuspended at concentrations ~ 1.5-2 μg/μL. For control injections, 1.5 kb dsRNA derived from
an irrelevant bacterial sequence was used. 6-day old tapeworms were obtained and microinjected
with dsRNA using femtotips II via the Femtojet injection system (Eppendorf) to obtain spreading
across the first ~ 3-4 mm anterior of the tapeworm. The spread of injected fluids could be detected
by a temporary increase in opacity. 500 hPa injection pressure for 0.3-0.8 ms was used per injection
site. Whole tapeworms were cultured *in vitro* for 3 days and then 2 mm anterior fragments were
amputated and cultured for another 3 days. On day 6, the fragments were re-injected with dsRNA
throughout the whole fragment and returned to *in vitro* culture for 9 days before they were
terminated for analysis.

Cell transplantations

For dissociated cell preparations, tapeworms were placed in a drop of calcium-magnesium
free HBSS (CMF HBSS, Gibco), minced into small pieces with a tungsten needle, incubated in

3X Trypsin-EDTA (59418C, Sigma) in CMF HBSS for 20 min at 37°C and dissociated using a

dounce homogenizer (Kontes). Cells were pelleted by centrifugation at 250 g for 5 min. The cell

pellet was washed in CMF HBSS and passed through cell strainers at 100 μm, 40 μm, 20 μm, and

10 μm (Falcon and Sysmex) with one spin and wash in between. Cells were pelleted and

resuspended in ~ 200-400 μL WH4 with 0.05% BSA. Cell injections were performed using the

Cell Tram Oil 4 injection system (Eppendorf) into the necks of irradiated worms. Injected worms

were cultured *in vitro* for 3 days before 5 mm anterior fragments were amputated and grown for

an additional 27 days.

Figure 1. Regeneration competence of *H. diminuta*. a, Schematic of *H. diminuta* adults. The tapeworm head includes suckers that attach the worm to the intestine. The neck is defined as the unsegmented region between the head and body. The proglottids grow in an anterior-to-posterior direction and increase in reproductive maturity as they are displaced posteriorly. The most posterior proglottids are gravid with embryos (dotted). We used 6-day old tapeworms that were not yet gravid. **b**, DAPI-stained 1 cm tapeworm fragments before and after 9 days in culture. **c-d**, Quantification of proglottid number and growth in length from (**b**). Error bars=SD, N=2-5, n=7-21 for each timepoint and condition. **e**, Representative DAPI-stained "neck only" fragments before and after 12 days *in vitro*. **f-g**, DAPI-stained fragments after serial amputation. 2 mm anterior fragments, with or without the head, were grown *in vitro* for 12-15 days and then re-amputated serially. Error bars=SD, one-way ANOVA for +head samples, t-test for -head samples. **h-i**, The neck was subdivided into 1 mm fragments from the anterior, middle, and posterior of the neck and grown *in vitro*. DAPI-stained worms are shown. Error bars=SD, N=3, n=22-29 per condition, one-way ANOVA.

464
Figure 2. Identification of cycling somatic cell genes using RNA-seq. a-b, DAPI staining of 5
mm anterior fragments from control and irradiated worms before and after *in vitro* culture for 6
days. Error bars=SD, one-way ANOVA. **c**, Schematic of RNA-seq strategy to identify irradiation-
sensitive transcripts. **d**, Confocal section of a tapeworm anterior. Cycling cells marked by *mcm2*
(magenta) are confined to the neck parenchyma between the nerve cords marked by *cadherin*
(green). (s: sucker, nc: nerve cord, oc: osmoregulatory canal, t: tegument, m: muscle, and p:
parenchyma). **e**, WISH of known cycling cell markers *h2b* and *mcm2*. **f**, WISH for genes expressed
in the neck parenchyma similar to *h2b* and *mcm2*. **g-i**, Confocal sections of dFISH for irradiation-
sensitive genes (green) with *h2b* or *mcm2* (magenta). Cyan arrowheads indicate cells that are
magnified at the far right.

Figure 3. RNAi to identify genes required for proper growth and regeneration in *H. diminuta*.
a, Schematic of RNAi paradigm. **b**, DAPI-stained worms after RNAi knockdown of *h2b, zmym3*, or *su(Hw)*. **c**, Quantification of worm lengths after RNAi. Error bars=SD, N=3-4, n=26-37, one-way ANOVA. **d**, Maximum-intensity projections from confocal z-stacks at tapeworm necks after RNAi. Panels are oriented anterior facing left. **e**, Quantification of (**d**). Number of F-ara-EdU[+] cells was normalized to total area from maximum-intensity projection of the DAPI stain. Error bars=SD, N=3, n=11-14, one-way ANOVA.

a

A ⟶ P

zmym3

ga   t   o   t

100 µm

su(Hw)

b

anterior-enriched (FC ≥1.5, FDR ≤0.05)   anterior-depleted (FC ≤-1.5, FDR ≤0.05)

frizzled   *delta*

B3ngalt   *gsx*

1 mm

c

cells from whole worms

5 mm

3 days   27 days ⟶ regenerate?

200 Gy   200 Gy + rescue with cells from whole worms

1 mm

d

cells from A or P

5 mm

A   P   3 days   27 days ⟶ regenerate?

200 Gy   200 Gy + rescue with cells from A   200 Gy + rescue with cells from P

1 mm

e

Neck tissue is maintained by signal gradient from head

form proglottid

inhibit proglottidization

Head provides stem cell niche

lineage commitment

niche

pluripotent zone

Stem cell niche is independent of the head and/or neck

local niche

Figure 4. Rescue of lethally irradiated tapeworms by cell transplantation. a, WISH patterns
for cycling cell genes across the tapeworm A-P axis (ga: genital anlagen; t: testis; o: ovary). **b,**
WISH at tapeworm anteriors for genes that were anteriorly enriched or anteriorly depleted by
RNA-seq across the A-P axis of the neck only (refer to Fig. 1h). Panels are oriented anterior facing
left. **c**, DAPI-stained worms after rescue with cell transplants from whole worms. **d**, DAPI-stained
worms after rescue with cell transplants sourced from 5 mm anteriors (A) or 5 mm posteriors (B).
e, Models for how region-specific regeneration is regulated in *H. diminuta.*

Supplemental Figures

Figure S1. Unsegmented neck is depleted after decapitation. a-b, DAPI-stained worms after
decapitation cultured *in vitro* for the indicated number of days. Red arrowheads mark the position
of the first visible proglottid. **b**, Worms were amputated 2 mm posterior to the head to remove
most of the neck tissue before culturing. After 15 days *in vitro* the neck tissue is no longer
identifiable.

Figure S2. Cycling somatic cells are distributed throughout the whole tapeworm body and are irradiation sensitive. a-c, Confocal sections after uptake of F-ara-Edu for 1 hr (**a**) or FISH to detect *mcm2* (**b**) or *h2b* (**c**). Panels are oriented anterior facing left. **d,** Confocal sections of BrdU labeled tapeworms +/- irradiation. Uptake of BrdU for 1 hr, through a section in the worm body after exposure to 200 Gy irradiation.

	0 Gy	200 Gy

zfp (HDt_10981)
FC=10
FDR=6.28e-171

HDp_01159
FC=6
FDR=3.16e-103

lamin (HDt_01618)
FC=1.6
FDR=1.6e-4

100 µm

Figure S3. Validation of RNA-seq results by WISH after irradiation. WISH from two regions of the tapeworm: anterior (left) and body (right), +/- exposure to irradiation followed by 3 days *in vitro.*

Figure S4. WISH patterns of potential cycling cell genes identified using RNA-seq. a, Classification of WISH expression patterns. **b-d**, WISH patterns at tapeworm necks with anterior oriented up. **b**, Examples of genes expressed in the neck parenchyma. **c**, Examples of genes expressed in a subset of cells within the neck. **d**, Examples of genes expressed broadly in the neck including toward the animal edge where differentiated muscle and tegument are located.

Figure S5. Colocalization of *mcm2* and *h2b*. Confocal section of dFISH to detect *mcm2* (green) and *h2b* (magenta) in the neck parenchyma. Panels are oriented anterior up.

Figure S6. The cycling somatic cell population is heterogeneous. a-d, Confocal sections of dFISH to detect irradiation-sensitive genes (green) with *h2b* or *mcm2* (magenta). Cyan arrowheads indicate cells that are magnified at the far right. Yellow arrowheads point to examples of expression in non-cycling cells. **e**, Summary of different co-expression patterns obtained from 53 dFISH experiments.

Figure S7. RNA-seq identifies anterior-enriched genes that are expressed predominantly in non-cycling cells. a, Differential gene expression analysis of anterior, middle, and posterior neck fragments from Fig. 1h was used to identify anterior-enriched and anterior-depleted transcripts. Transcripts were considered anteriorly enriched if they were upregulated in the anterior vs. middle or posterior fragments with fold change ≥1.5 and FDR ≤0.05 and conversely for anterior-depleted transcripts. Overlap with irradiation-sensitive transcripts is shown. **b**, WISH of genes that were anteriorly enriched and irradiation sensitive. All panels are oriented with anterior facing up. **c**, Confocal sections of dFISH to detect genes from (**b**), with *mcm2* (magenta). Cyan arrowheads indicate cells that are magnified at the far right.

References

535   1.   Roberts, L. S. The influence of population density on patterns and physiology of growth in
536   *Hymenolepis diminuta* (Cestoda:Cyclophyllidea) in the definitive host. *Exp. Parasitol.* **11,**
537   332–371 (1961).

538   2.   Del Brutto, O. H. Human cysticercosis (*Taenia solium*). *Trop Parasitol* **3,** 100–103
539   (2013).

540   3.   Craig, P. & Ito, A. Intestinal cestodes. *Curr. Opin. Infect. Dis.* **20,** 524–532 (2007).

541   4.   Arai, H. P. *Biology of the Tapeworm Hymenolepis diminuta.* (Academic Press, Inc., 1980).

542   5.   Read, C. P. Longevity of the tapeworm, *Hymenolepis diminuta. J. Parasitol.* **53,** 1055–
543   1056 (1967).

544   6.   Newmark, P. A. & Sánchez Alvarado, A. Not Your Father's Planarian: A Classic Model
545   Enters The Era Of Functional Genomics. *Nat. Rev. Genet.* **3,** 210–219 (2002).

546   7.   Reddien, P. W. The Cellular and Molecular Basis for Planarian Regeneration. *Cell* **175,**
547   327–345 (2018).

548   8.   Baguñà, J. The planarian neoblast: the rambling history of its origin and some current
549   black boxes. *Int. J. Dev. Biol.* **56,** 19–37 (2012).

550   9.   Roberts, L. S. Development of *Hymenolepis diminuta* in its definitive host in *Biology of*
551   *the Tapeworm Hymenolepis Diminuta,* 357–423 (Academic Press, Inc., 1980).

552   10.   Collins, J., III *et al.* Adult somatic stem cells in the human parasite *Schistosoma mansoni.*
553   *Nature* **494,** 476–479 (2013).

554   11.   Koziol, U., Rauschendorfer, T., Zanon Rodríguez, L., Krohne, G. & Brehm, K. The
555   unique stem cell system of the immortal larva of the human parasite *Echinococcus*
556   *multilocularis. EvoDevo* **5,** 10 (2014).

557   12.   Wang, B., Collins, J., III & Newmark, P. A. Functional genomic characterization of
558   neoblast-like stem cells in larval *Schistosoma mansoni. eLife* **2,** e00768 (2013).

559   13.   Tsai, I. J. *et al.* The genomes of four tapeworm species reveal adaptations to parasitism.
560   *Nature* **496,** 57–63 (2013).

561   14.   Brehm, K. & Koziol, U. On the importance of targeting parasite stem cells in anti-
562   echinococcosis drug development. *Parasite* **21,** 72 (2014).

563   15.   Goodchild, C. G. Transfaunation and repair of damage in the rat tapeworm, *Hymenolepis*
564   *diminuta. J. Parasitol.* **44,** 345–351 (1958).

565   16.   Schiller, E. L. A simplified method for the in vitro cultivation of the rat tapeworm,
566   *Hymenolepis diminuta. J. Parasitol.* **51,** 516–518 (1965).

567   17.   Bolla, R. I. & Roberts, L. S. Developmental physiology of cestodes. IX. Cytological
568   characteristics of the germinative region of *Hymenolepis diminuta. J. Parasitol.* **57,** 267–
569   277 (1971).

570   18.   Sulgostowska, T. The development of organ systems in cestodes. I. A study of histology
571   of *Hymenolepis diminuta* (Rudolphi, 1819) (Hymenolepididae). *Acta Parasitologica*
572   *Polonica* **20,** 449–462 (1972).

573   19.   Solana, J. *et al.* Defining the molecular profile of planarian pluripotent stem cells using a
574   combinatorial RNAseq, RNA interference and irradiation approach. *Genome Biol.* **13,** R19
575   (2012).

576   20.   Eisenhoffer, G. T., Kang, H. & Sánchez Alvarado, A. Molecular analysis of stem cells and
577   their descendants during cell turnover and regeneration in the planarian *Schmidtea*
578   *mediterranea. Cell Stem Cell* **3,** 327–339 (2008).

21. Koziol, U. & Castillo, E. Cell proliferation and differentiation in cestodes. *Research in*
Helminths 121–138 (2011).
22. Collins, J., III, Wendt, G. R., Iyer, H. & Newmark, P. A. Stem cell progeny contribute to
the schistosome host-parasite interface. *eLife* **5,** 243 (2016).
23. Pierson, L. *et al.* RNA interference in a cestode reveals specific silencing of selected
highly expressed gene transcripts. *Int. J. Parasitol.* **40,** 605–615 (2010).
24. Mizukami, C. *et al.* Gene silencing in *Echinococcus multilocularis* protoscoleces using
RNA interference. *Parasitol. Int.* **59,** 647–652 (2010).
25. Spiliotis, M. *et al. Echinococcus multilocularis* primary cells: improved isolation, small-
scale cultivation and RNA interference. *Mol. Biochem. Parasitol.* **174,** 83–87 (2010).
26. Pouchkina-Stantcheva, N. N., Cunningham, L. J., Hrčkova, G. & Olson, P. D. RNA-
mediated gene suppression and in vitro culture in *Hymenolepis microstoma. Int. J.*
Parasitol. **43,** 641–646 (2013).
27. Hu, X. *et al.* Gene knockout of *Zmym3* in mice arrests spermatogenesis at meiotic
metaphase with defects in spindle assembly checkpoint. *Cell Death Dis* **8,** e2910 (2017).
28. Leung, J. W. C. *et al.* ZMYM3 regulates BRCA1 localization at damaged chromatin to
promote DNA repair. *Genes Dev.* **31,** 260–274 (2017).
29. Gdula, D. A., Gerasimova, T. I. & Corces, V. G. Genetic and molecular analysis of the
gypsy chromatin insulator of *Drosophila. Proc. Natl. Acad. Sci. U.S.A.* **93,** 9378–9383
(1996).
30. Sikes, J. M. & Newmark, P. A. Restoration of anterior regeneration in a planarian with
limited regenerative ability. *Nature* **500,** 77–80 (2013).
31. Liu, S. Y. *et al.* Reactivating head regrowth in a regeneration-deficient planarian species.
Nature **500,** 81–84 (2013).
32. Umesono, Y. *et al.* The molecular logic for planarian regeneration along the anterior-
posterior axis. *Nature* **500,** 73–76 (2013).
33. Koziol, U., Jarero, F., Olson, P. D. & Brehm, K. Comparative analysis of Wnt expression
identifies a highly conserved developmental transition in flatworms. *BMC Biol.* **14,** 1
(2016).
34. Forsthoefel, D. J. *et al.* An RNAi screen reveals intestinal regulators of branching
morphogenesis, differentiation, and stem cell proliferation in planarians. *Dev. Cell* **23,**
691–704 (2012).
35. Collins, J., III, King, R. S., Cogswell, A., Williams, D. L. & Newmark, P. A. An atlas for
Schistosoma mansoni organs and life-cycle stages using cell type-specific markers and
confocal microscopy. *PLoS Negl Trop Dis* **5,** e1009 (2011).
36. King, R. S. & Newmark, P. A. In situ hybridization protocol for enhanced detection of
gene expression in the planarian *Schmidtea mediterranea. BMC Dev. Biol.* **13,** 8 (2013).
37. Salic, A. & Mitchison, T. J. A chemical method for fast and sensitive detection of DNA
synthesis in vivo. *Proceedings of the National Academy of Sciences* **105,** 2415–2420
(2008).
38. Currie, K. W. *et al.* HOX gene complement and expression in the planarian *Schmidtea*
mediterranea. EvoDevo **7,** 7 (2016).
39. Rouhana, L. *et al.* RNA interference by feeding in vitro-synthesized double-stranded RNA
to planarians: methodology and dynamics. *Dev. Dyn.* **242,** 718–730 (2013).