1 **A new test of value-modulated attentional bias in mice**

3 **Iasmina Hornoiu, John Gigg & Deborah Talmi**

5 [a]Division of Neuroscience and Experimental Psychology, University of Manchester

Address for correspondence

Deborah Talmi, Division of neuroscience and experimental psychology, School of Biological

Sciences, University of Manchester, Manchester, UK, M139PL.

Telephone: 0161 275 1968

Email: Deborah.Talmi@manchester.ac.uk

Abstract

The cognitive process of attention can be modulated by the emotional value of the stimulus. In order

to understand this association further, we require an animal test of value-modulated attention capture

evoked by ethologically valid stimuli. In mice, female odour triggers arousal and elicits emotional

responses in males. Here, we determined the extent to which objects infused with female odour

captured the attention of male mice. Eight animals were tested in six experiments using a modified

version of the spontaneous Novel Object Recognition task. Attention was operationalised using

differential exploration time of odour-infused objects (O+), identical but odour-neutral objects (O-),

and novel objects (X). We found that when single objects were presented, mice spent comparatively

more time exploring O+ objects both when they were novel (Experiment 1) and when they were

familiar (Experiment 2). When O+ and O- competed with each other for attention, O+ captured

attention preferentially. O+ objects captured attention more than O- objects at a novel (Experiment 3)

or a familiar (Experiment 4) location; they also captured attention more than novel objects at either a

novel (Experiment 5) or a familiar (Experiment 6) location. These results demonstrate that in mice,

female odour can in some circumstances capture attention more than object novelty. These results

pave the way for using odour to modulate memory in tests of novel object recognition.

Keywords: attention, emotional arousal, memory, Novel Object Recognition task, olfactory stimuli

Introduction

Emotional arousal influences cogitation extensively, from early perception, to attention (Golomb, Nguyen-Phuc, Mazer, McCarthy, & Chun, 2010; Pourtois, Schettino, & Vuilleumier, 2013), to higher order cognitive functions (Pessoa, 2009), including memory (Levine & Edelstein, 2009; Talmi, 2013). A central objective of human neuropsychological and neuroimaging research is to trace the neurobiological underpinnings of the link between emotional arousal and attention (Hartikainen, Ogawa, Soltani, & Knight, 2007; Kadohisa, 2013; Lee, Sakaki, Cheng, Velasco, & Mather, 2014; Talmi & McGarry, 2012; Vermeulen, Godefroid, & Mermillod, 2009). In line with this objective, the present study was designed to investigate the impact of an emotionally charged stimulus on attention allocation in mice. Establishing the means to measure such a link would provide a valuable animal task with which to further understand the neurobiology of emotional memory and attention.

Attention is crucial for the effective processing of perceptual information presented by the environment at any given time (Chun, Golomb, & Turk-Browne, 2011). Behaviourally-relevant stimuli are thought to be selected against others and prioritised via two routes: top-down and bottom-up (Corbetta, Patel, & Shulman, 2008; Miller & Cohen, 2001; Pratt & Hommel, 2003). Endogenous attentional control, also referred to as *goal-directed* or *top-down*, is a voluntary process that operates at the level of memory and decision making to modulate information according to the agent's intentions. In contrast, the exogenous, involuntary attentional capture by certain stimuli, known as *bottom-up* or *stimulus-driven attentional process*, depends on the physical characteristics of stimuli and is outside the agent's control. A recently proposed model, called Multiple Attention Gain Control (MAGiC), suggests a third route to attentional selection, through emotion. Emotion is thought to bias perception via distinct gain control mechanisms originating in the amygdala. Neuropsychological, neuroimaging and behavioural studies have provided evidence for the modulatory effects of emotion on sensory processing, whereby emotionally charged stimuli acquire increased representation and access to awareness compared to neutral stimuli (for review, see Pourtois et al., 2013). The majority of studies that informed the MAGiC model utilised threat-related stimuli, due to their obvious behavioural salience (Compton, 2003; Vuilleumier, 2005), as well as their role in various pathological conditions in humans, such as anxiety and phobias (Öhman & Mineka, 2001). However, in order to achieve a more complex understanding of the influence of emotion on the selection of sensory information, more research is needed on the effects of other emotional stimuli on attention, including those with positive value.

Stimuli that predict a valuable outcome – whether positive or negative - capture attention automatically, even when they are task-irrelevant (Anderson, Laurent, & Yantis, 2011; Wang, Duan, Theeuwes, & Zhou, 2014; Wentura, Müller, & Rothermund, 2014). This value-derived influence on

attention is neither top-down, nor bottom-up, since attention in this case is neither voluntarily directed

by contextually relevant goals, nor driven by the sensory significance of the stimuli, respectively.

Rather, attention is directed to stimuli that have acquired the potential to predict valuable outcomes

via associative learning. This mechanism is referred to as *value-modulated attentional capture*

(VMAC) (Le Pelley, Mitchell, Beesley, George, & Wills, 2016). While prioritising unexpected high-

value stimuli might be biologically advantageous in certain situations, in others it could interfere with

goal-directed behaviours. Further research is needed to offer a more in-depth understanding of the

computational control of VMAC (Talmi, Slapkova, & Wieser, 2018).

The neural basis of top-down and bottom-up attention has been studied in detail in humans and in

animal models. In humans, the majority of studies have been conducted on visual attention.

Endogenous (top-down) signals arising in higher-order prefrontal, parietal and limbic cortices interact

with exogenous (bottom-up) signals driven by visual cortical pathways to bias attention in favour of

attended targets, while suppressing representations of unattended information (Desimone & Duncan,

1995; Gitelman, 2003; Katsuki & Constantinidis, 2014). Across the literature, goal-directed

attentional processing has been generally attributed to dorsal fronto-parietal brain areas, while

stimulus-driven attentional control is believed to be mediated by ventral temporal-parietal networks

(Anderson, Laurent, & Yantis, 2014; Corbetta & Shulman, 2002; Yantis et al., 2002). Animal studies

of sustained attention, particularly in task-performing rats, have revealed the role of the cholinergic

system in top-down, as well as bottom-up attentional control. The basal forebrain corticopetal

cholinergic projections, which are activated via direct glutamatergic inputs from the prefrontal cortex

to the basal forebrain, terminate in all cortical areas and are thought to mediate top-down processes in

tasks involving sustained attention (McGaughy & Sarter, 1998; Sarter, Givens, & Bruno, 2001;

Zaborszky, Gaykema, Swanson, & Cullinan, 1997). Lesions of the basal forebrain, affecting inputs of

acetylcholine particularly in fronto-dorsal cortical areas of rat models of attention, result in prolonged

impairments of sustained attention (Sarter et al., 2001). Bottom-up attentional processes have been

shown to largely depend on noradrenergic projections, which originate in the locus coeruleus and

terminate in the thalamus and the basal forebrain. Noradrenergic activation of basal forebrain

corticopetal projections is involved in the processing of threat-related or anxiety-inducing stimuli in a

bottom-up fashion via the recruitment of telencephalic systems (Aston-Jones, Rajkowski, Kubiak,

Valentino, & Shipley, 1996; Sarter et al., 2001). So far, these animal studies have demonstrated that

the basal forebrain cortical cholinergic system represents a core component of the neuronal circuitry

involved in attentional processing, thus contributing to the understanding of the role of the fronto-

parietal cortical regions from human neuropsychological and imaging research. Integrating evidence

from human and animal works is necessary for the development of a reliable model of the neural

mechanisms of attention.

The neural basis of VMAC has been less well understood in animal models. In humans, Anderson et
al. (2014) argue that the neural mechanisms underlying VMAC are mediated by the tail of the caudate
nucleus and the extrastriate visual cortex. Using functional magnetic resonance imaging (fMRI), these
researchers found that task-irrelevant reward-predictive stimuli acting as distractors acquired stronger
representation in the caudate tail and triggered greater activity in the extrastriate visual cortex versus
other non-target stimuli. So far, animal models of attention using stimuli with emotional value have
been derived from the traditional theories of associative learning and conditioning. In one model of
associative learning, proposed by Mackintosh (1975), animals pay more attention to cues that are
reliable predictors of a consequence (high predictiveness) than to non-predictive cues. Selective
attentional bias towards good predictors allows animals to focus on relevant cues, while ignoring
distractors, thereby achieving optimal performance. In contrast, the model by Pearce and Hall (1980)
states that cues with uncertain consequences capture the most attention. The idea behind this model is
that because unreliable cues are surprising, they will be allocated more attentional resources that leads
to rapid learning about their significance. While there is abundant evidence in favour of both
predictability (Duffaud, Killcross, & George, 2007; George & Pearce, 1999) and uncertainty (Kaye &
Pearce, 1984; Wilson, Boumphrey, & Pearce, 1992) models, several hybrid models have emerged in
an attempt to reconcile these principles (for reviews, see Esber & Haselgrove, 2011; Le Pelley et al.,
2016). To date, very little, if any, data exist on the influence of biologically relevant stimuli on
VMAC in rodents. The present study is, to our knowledge, the first research using female odours to
examine VMAC in a rodent model of attention.
149
Unlike humans, who are predominantly influenced by visual stimuli (Shapiro, Egerman, & Klein,
1984), most mammals, including rodents, rely mainly on odours to obtain information about their
environment (Brennan & Kendrick, 2006; Johnston, 2003). Odours from conspecifics of opposite sex
are thought to carry positive reward values, because they elicit an approach response to promote
reproductive behaviours, as opposed to withdrawal or avoidance responses, which supress
reproductive behaviours (Beny & Kimchi, 2014). The motivational significance of odours bearing
reproductive value does not seem to depend on the animal's prior sexual experience; these stimuli
can, therefore, be considered primary reinforcers. This is evident in the laboratory, where sexually
inexperienced male rodents display typical sexual behaviours when exposed to female odours (Beny
& Kimchi, 2014). Given the socio-biological relevance of odours and their known motivational
effects on animals, it is reasonable to assume that odours could be used as stimuli to investigate
VMAC in an animal model.
162
163   Six experiments were designed to quantify the degree of attention allocation in male mice to objects
164   infused with female odour (O+), odour-neutral objects (O-) and novel objects (X) in a modified
165   version of the spontaneous Novel Object Recognition (NOR) task. NOR is one of the most widely

used paradigms in studies of memory functions in rodents. The NOR task can be configured to measure attention (Silvers, Harrod, Mactutus, & Booze, 2007) and has been used in studies focusing on pathological conditions or drug abuse, which result in attentional deficits (Alkam et al., 2011; Piper, Fraiman, & Meyer, 2005). Originally developed by Berlyne (1950) and subsequently adapted for use in mice by Murai *et al.* (2007), this behavioural task relies on the drive of rodents to explore novelty in the absence of any training or external reinforcers. The original NOR task comprises three phases: habituation, familiarisation and test. In the first two phases, the animal is exposed to an open-field arena (habituation) and then to the same arena in the presence of two identical objects (familiarisation). The test phase is similar to the familiarisation phase, with the exception that one of the identical objects is replaced with a novel object. Healthy adult rodents recognise the familiar object at test and pay more attention to the novel (more arousing) object, indicated by a longer exploration time of the novel versus familiar object (Ennaceur, 2010; Gaskin et al., 2010; Mumby, Glenn, Nesbitt, & Kyriazis, 2002). In the modified NOR paradigm used in this study, instead of novel and familiar objects, the attention allocation of male mice was investigated for O+ versus O- or X, which were placed in either novel or familiar locations in a Y-maze.

The first two experiments assessed the attentional capture of mice by an O+ compared with an O-, when both objects were novel (Experiment 1) or familiar (Experiment 2). We hypothesised that because the motivational value of O+ was higher than of O-, the former would attract more attention in both experiments. We then asked whether the same would be true if O+ and O- were placed at different locations in the arena, so that O+ would be found at either a novel (Experiment 3) or a familiar (Experiment 4) location. In these experiments, novel location was used due to its previously demonstrated influence on attention (Ennaceur, Neave, & Aggleton, 1997) and, thus, it was interesting to study how novel location interacted or competed with female odour for attention allocation. We hypothesised that when the location of O+ was novel, O+ would capture more attention than O-, because the former benefited from the advantage of two salient motivational features, namely, motivational value and novel location. However, in Experiment 4 it was less clear whether O+ would still 'win' over O-, due to the fact that the novel location of the latter was now competing with the odour of the former for attention. In the last two experiments, we used another well-known motivational feature, novel identity, and assessed how strong the attention capture by odour was when competing with a novel object (X). Novel objects are expected to recruit more attention than familiar objects in the traditional NOR task and, therefore, we were interested to test how attention to O+ would be influenced by the presence of an X. Experiment 5 tested the attention captured by O+ at a novel location versus an X at a familiar location. Here, O+ had the advantage of odour and location novelty, while X had the advantage of novel identity. In the last experiment, X was placed at a novel location and O+ at a familiar location. Since this time X had the advantage of both

202 novel identity and novel location, the question was how much attention would odour attract when
203 competing with this highly arousing combination assigned to X.

204
205
206 **Methods**

207
Animals
Behavioural experiments were performed using eight adult male C57BL/6J mice (ENVIGO, UK),
which were 14 weeks (3.5 months) old at the start of the experiments and weighed 30±3 g throughout
the project. Mice were weighed prior to each experiment and at the end of the last experiment on a
laboratory scale (Kern FCB, Germany). Mice were maintained by the Biological Services Facility
(BSF), University of Manchester, UK and housed in groups of four individuals in ventilated
Techniplast cages, in standard conditions (20°± 2°C temperature and 55±5% humidity) on a 12:12
light/dark cycle, with *ad libitum* access to food and water. All mice were ear punched for
identification. The experiments were carried out over a period of two months and took place between
9:30am-12:30pm. All procedures were conducted in conformity with the University of Manchester
BSF regulations for animal husbandry and with the Home Office Animals (Scientific Procedures) Act
(1986) and were licenced by the UK Home Office and University of Manchester Ethical Review
Panel.

221
Apparatus
Experiments were performed in custom Y-mazes with three identical white, opaque plastic arms,
(length 160mm, height 280mm) diverging at a 120° angle from each other (designed by Jack Rivers-
Auty and constructed by Plastic Formers Ltd, UK). Each arm became wider at the end to form a small
square arena (length 92mm x width 90mm). The square arenas could be differentiated by the presence
of salient visual cues. Individual mice were randomly assigned to a particular Y-maze throughout
habituation and testing. Mice were always released inside the middle arm (the arm closest to the
nearest room wall), with their backs to the right and left arms, which contained objects (see
Materials). This strategy, following the guidelines by Antunes and Biala (2012), ensured that external
pressure to explore the objects was avoided. During the interval between exposures (habituation) and
the inter-trial interval (ITI, experiments), mice from each cage group were placed in separate holding
cages (standard housing cages, Techniplast), which remained the same throughout the project. Mice
from each cage group were run simultaneously in four Y-mazes placed next to each other. Video
cameras (JVC, 40x optical zoom) placed above the Y-mazes recorded animal performance.

236
Materials

The objects used in experiments were either built from LEGO$^®$ pieces or various other plastic shapes
(Figure 1). Given that the egg halves were identical in shape, the difference in their colour as well as
their position in the maze (face-down or on one side with either convex or concave side facing the
animal) were used to create different object types. All objects were odour-neutral and made of plastic
in order to avoid material preference, minimise odour saturation and facilitate cleaning. The objects
were attached to the floor inside the Y-maze with Blu-Tack$^®$. Objects used in the habituation phase
were plastic letters and were never used in subsequent experimental trials. New object types were
used in each experiment, in order to avoid habituation to any object type.
246
For each experiment, soiled cage bedding from cages containing four female mice (C57BL/6J strain)
was used to label some objects with female odour. The objects to be labelled with female odour were
placed in the bedding the day prior to each trial, around midday (12:00pm). For the experiments that
lasted four days (Experiments 1, 3 and 4), halfway through each experiment (at the end of day 2), the
bedding with female odour was replaced with bedding from a cage containing a second group of
control females of the same strain. For the experiments that lasted only 2 days (Experiments 2, 5 and
6), the bedding remained the same throughout each experiment, but was changed before the start of
the next experiment. The reason for this was that it was observed that the mice had a tendency to
habituate to the smell after day 2 and using new bedding prevented this. Objects with female odour
are here referred to as 'O+'. Copies of the same object but without odour are referred to as 'O-'.
Objects that were never infused with odour are termed 'X'. Y-mazes and objects were cleaned
thoroughly with 70% ethanol and wiped with paper towels between trials with mice from different
group cages, as well as at the end of each daily session.
260

Figure 1. Examples for objects used in the study

Procedure

The tasks used in this study were a modified version of the classical NOR task (Berlyne, 1950). The

procedure of each experiment is depicted in Figure 2. The study began with a familiarisation stage,

during which the animals were handled for one minute on two consecutive days so that they became

accustomed to the experimenter.

Habituation. After familiarisation, the mice were habituated to the testing apparatus and objects over

a five-day period prior to Experiment 1. The first day of habituation consisted of a 10-minute cage

group habituation session to the Y-mazes, in which mice from a given cage were placed together in

one Y-maze, in the absence of objects. On the following day, the mice were exposed individually to

the Y-maze for 10 minutes, again without objects. On the third day, the same steps from day 2 were

repeated, but this time an O- was present in either the left or right choice arm. On the fourth day,

following the same steps as on day 3, each animal was exposed to another O-, but in the opposite arm

to the one on day 3. The last day of habituation consisted of two exposures, this time with an O+ in

either the left or right arm of the maze. The duration of exposure was 10 minutes and the interval

between exposures was 5 minutes. During the 5-minute interval, the mice were placed in their holding

cages and the O+s presented in the first exposure were replaced with different O+s for the second

exposure.

Experimental stage. Following habituation, the mice were tested in six different experiments.

Experiments 1, 3 and 4 were conducted on four consecutive days. Experiments 2, 5 and 6 were
conducted on two consecutive days. There were two experimental trials each day. The interval
between two experimental trials (ITI) was 5 minutes. During the ITI, the animals were placed in
holding cages and the objects were replaced for the next trial. The particular object used in a given
trial (O+, O- or X), the order in which objects were presented and their locations in the maze (left or
right choice arm) were randomised for each animal in each experiment. This randomisation ensured
that each animal was exposed to a given object only in one experimental trial across the entire study.
Each animal experienced the same number of O+, O- and X objects in each experiment. Apart from
Experiment 1, each animal was exposed only to one O+ on each day and whether this was in the first
or second trial of the day was counterbalanced. All of the experiments were performed consecutively
by the same animals. Chronologically, the experiments were carried out in the following order:
Experiment 1, Experiment 3, Experiment 4, Experiment 2, Experiment 5, Experiment 6. Thus, the
mice were 14 weeks (3.5 months) old in Experiment 1, 16 weeks (4 months) old in Experiment 3, 19
weeks (almost 5 months) old in Experiment 4 and 29-30 weeks (around 7.5 months) old in
Experiments 2, 5 and 6.

Experiment 1. This experiment consisted of eight trials in total per animal. Trials included one
exposure to a single object, either O+ or O-. The duration of each trial was one minute (see row 1 of
Fig. 2).

Experiment 2. This experiment was composed of four trials per animal. Each trial included an initial
exposure to O- for one minute (the 'study phase'), either in the left or the right arm of the maze,
followed by an exposure either to O- or O+ for 3 minutes (the 'test phase'; see row 2 of Fig. 2).

Experiment 3. This experiment included eight trials in total per animal. Each trial included a study
phase, where animals were exposed to O- for one minute, either in the left or the right arm of the
maze. The location of O- in the study phase is referred to as the 'familiar' location. The other location
– the arm that was empty during the study phase - is referred to as the 'novel' location. During the test
phase animals were exposed to both O- and O+ for 3 minutes. The O+ was always placed in the
'novel' location (see row 3 of Fig. 2).

Experiment 4. This experiment was identical to experiment 2, except that in the test phase O+ was in
a familiar location (see row 4 of Fig. 2).

Experiment 5. This experiment was composed of four trials in total per animal. Each trial included a
study phase, where animals were exposed to O- for one minute, either in the left or the right arm of
the maze. The location of O- in the study phase is referred to as the 'familiar' location. The other

location – the arm that was empty during the study phase - is referred to as the 'novel' location.

During the test phase animals were exposed to both O+ and X (an entirely novel object) for 3 minutes.

The O+ was always placed in the 'novel' location (see row 5 of Fig. 2).

Experiment 6. This experiment was identical to experiment 5, except that O+ in the test phase always

occupied a familiar location (see row 6 of Fig. 2).

Figure 2. Experimental design.

Trial types in experiments 1-6 are illustrated (one experiment in each row). O+: Odour-infused

object. O-: odour-neutral object. Different copies of objects were used in two-phase experiments

(Experiments 3-6). X: novel objects, never infused with odour.

331
Data analysis
Exploratory behaviour was recorded with video cameras and subsequently, object exploration time for
each object was scored using the Novel Object Timer software (Jack Rivers-Auty; Novel Object
Recognition Task Timer, 2015). The animal was considered to be exploring an object when its nose
was within 2cm of the object and directed at the object. Sitting next to the object or climbing on top of
the object was not regarded as exploratory behaviour. Each trial was scored twice for accuracy and
the average of the two scorings taken as the object exploration time per trial. The mean exploration
time of O+, O- and X was obtained by averaging all trials with these objects for each animal.
340
The displacement index (D2), used to assess object preference from Experiments 3 onwards, was
calculated using the formula: $D2 = (T_{O+} - T_{O-}) / (T_{O+} + T_{O-})$, where T_{O+} is the mean exploration time of
O+ objects and T_{O-} is the mean exploration time of O- objects. The values for this index are bound
within a range of -1 and 1; positive D2 values indicated preference for O+, while negative values
signified preference for O- and a value of 0 signified no preference (Burke, Wallace, Nematollahi,
Uprety, & Barnes, 2010; Oliveira, Hawk, Abel, & Havekes, 2010).
347
348 The differences in total exploration time of O+ and O- were analysed with paired t-tests (two-tailed).
349 D2 values were analysed with one-sample t-tests. In Experiments 3-4, we also investigated the effect
350 of experiment day, using a repeated-measures 2 (Object type: O+, O-) x 4 (day: 1-4) ANOVA.
351
352 <div align="center">**Results**</div>
353
354 **Experiment 1: Novel objects O+ vs. O-**
355 The total time animals spent exploring O+ was significantly higher than the total exploration time for
356 O- ($t(5)=4.11$, $p=0.0046$) as illustrated in Figure 3 (left panel). This finding demonstrates that when
357 both O+ and O- are novel, the former attracts more attention. Two animals were excluded from the
358 analysis of Experiment 1 due to a counterbalancing error.
359
360 **Experiment 2: Familiar objects O+ vs. O-**
361 The total object exploration time in Experiment 2 (Fig. 3 right panel) was significantly higher for O+
362 than O- ($t(7)=3.67$, $p=0.008$). This finding demonstrates that when both O+ and O- are familiar, the
363 former attracts more attention.

Figure 3. Exploration of single objects in Experiments 1 and 2.

Left: *Results of Experiment 1: Novel O+ vs. Novel O-.* ***Right:*** *Results of Experiment 2: Familiar O+ vs. Familiar O-. Bar graphs represent object exploration times (in seconds) averaged across animals. Data are presented as means ± SEM. Statistical significance is expressed with ** (p≤0.01).*

Experiment 3: Competition between O+ in a novel location and O- in a familiar location.

Figure 4 shows that the total exploration time for O+ was significantly higher than that for O- (t(7)=6.99, p=0.0001). Additionally, animals demonstrated a significant preference for O+ over O-, as indicated by D2 value, which was significantly higher than zero (t(7)=9.33, p<0.0001). Object exploration time was analysed across experimental days for any effect of day or interaction between object type and day. A repeated-measures two-way-ANOVA test found that there was a significant effect of object type (F(1,7)=47.84, p<0.001). While numerically this difference diminished across days, neither the effect of day (F(3,21)=2.93, p=0.06), nor the interaction (F(3,21)<1, p= 0.53) were significant.

Experiment 4: Competition between O+ in a familiar location and O- in a novel location.

The total exploration time of O+ was significantly higher than that of O- (*t*(7)=3.48, *p*=0.0051; Figure 4). The D2 value (Figure 4), indicating the preference for O+ over O-, was significantly higher than zero (t(7)=3.67, p=0.008). Object exploration time was analysed across experimental days for any day effect or interaction between object type and day. Replicating the above results, there was a significant effect of object type (F(1,7)=12.02, p=0.01), as determined by a two-way-ANOVA test. As in the previous experiment, here the effect of day (F(3,21)=2.24, p=0.11) and the interaction (F(3,21)=1.13, p=0.36) were not significant.

At the end of the third experiment, it was interesting to determine if the mice had higher preference

for O+ placed in a novel location than for O+ in a familiar location. The overall D2 values from

experiments 3 and 4 were statistically compared with a Student's paired t-test. As illustrated in Figure

4, the D2 value for O+ in a novel location was significantly greater than that for O+ in a familiar

location (t(7)=2.32, p=0.027), indicating that location novelty increased the motivational value of O+.

However, since the D2 values were from two separate experiments, it is important to take into account

possible confounds arising from a counter effect of habituation. Since Experiment 4 followed

Experiment 3, animals might have been less motivated to explore the objects in the former than in the

latter.

Figure 4. Results of Experiments 3 and 4.

Top left: Object exploration time in Experiments 3 and 4. Means of object exploration time (seconds) were calculated by averaging across all animals.

*Top **right**: Effect of location familiarity on odour preference. Mean D2 (discrimination index) value was obtained as the average of the D2 values from all animals.*

*Bottom: Daily object exploration in Experiment 3 (Left) and 4 (Right). Means of object exploration time (seconds) were obtained by averaging across all animals for each experimental day. The bar graphs represent means ±SEM. Statistical significance is expressed as ns (p>0.05), * (p≤0.05), ** (p≤0.01), or *** (p≤0.001).*

Experiment 5: Competition between O+ in a novel location and X in a familiar location.

As shown in Figure 5, the total object exploration time in Experiment 5 was significantly higher for O+ than for X (t(7)=5.63, p=0.0008). A one sample t-test showed that the D2 value was significantly higher than zero, indicating a preference for O+ over X (t(7)=4.92, p=0.0017).

Experiment 6: Competition between O+ in a familiar location and X in a novel location.

The total object exploration time in Experiment 6 was higher for O+ than for X, however, the difference was not statistically significant (t(7)=1.8, p=0.1158, Figure 5). The D2 value in this experiment was also not significantly higher than zero, suggesting no significant preference for O+ over X (t(7)=2.023, p=0.0828).

Additionally, the results from Experiment 5 and Experiment 6 were compared in order to determine the effect of location familiarity and object type (novel identity or odour-infused) on exploration time and object preference. The same caveats hold for this comparison across experiments as for the one across Experiments 3 and 4. A repeated-measures two-way-ANOVA test found that there was a significant effect of location (F(1,7)=19.23, p=0.003) with a preference for the novel object, as well as of object type (F(1,7)=11.68, p=0.011), but no significant interaction (F(1,7)=1.59, p=0.246). It is difficult to interpret these results conclusively as indicative of additive or sub-additive effects, given the lack of counterbalance between experiments and potential effects of habituation.

Figure 5. Results of Experiments 5 and 6.

Left: Effect of object identity and location familiarity on exploration time. Means of object exploration time (seconds) were calculated by averaging across all animals. Bar graphs represent object exploration times (in seconds) averaged across animals.

Right: Effect of object identity and location familiarity on object preference. Mean D2 (discrimination index) value was obtained as the average of the D2 values from all animals. Data are presented as means ± SEM. Statistical significance is expressed as ns (p>0.05), (p≤0.05) or *** (p≤0.001).*

In Experiments 5 and 6, it was also interesting to observe the choice the animal made when it had to

decide which object to explore first (O+ or X); this is referred to as *first object choice*. A binomial test

found that mice were equally likely to choose either object first so they had no first-choice preference

($p=0.1102$ in Experiment 5, $p=0.2153$ in Experiment 6).

Discussion

Evidence for VMAC from sensory domains outside vision is sparse, despite abundant research on

different sensory modalities involved in bottom-up and top-down attention processes (Spence, 2010).

The reason for this is that the initial identification of VMAC was for visual cues and since then, the

scientific focus has been on analysing how learnt value affects visual attention (Chelazzi et al., 2014;

Failing & Theeuwes, 2014; Qi, Zeng, Ding, & Li, 2013). Extending the principles of VMAC to other

sensory domains is essential for a more complex understanding of this attentional process and can

help integrate current knowledge of the modulatory effects of learnt reward on sensory processing

(Pantoja et al., 2007). In the study we report here, we investigated the attentional capture by an

olfactory stimulus and the possibility of any cross-modal interference with visual stimulation by an

object at a novel location or a novel object in the arena.

Previous studies have shown that female odour represents a positive arousing stimulus for laboratory

male mice (Beny & Kimchi, 2014; Connor, 1972; Mackintosh, 1970). According to Beny and Kimchi

(2014), sexually inexperienced (naïve) male mice display sexual behaviours towards female

conspecifics and manifest aggression towards other male mice. Connor (1972) demonstrated that

pheromones found in urine are responsible for these dimorphic behaviours. In his experiments, male

mice displayed milder aggressive behaviours towards male intruders smelling of female urine and

behaved aggressively towards females swabbed with male urine. These behaviours are genetically

determined and, therefore, do not require prior learning. The only experience these laboratory mice

had with female odours was during weaning, in the presence of their mothers, after which they were

isolated from female conspecifics.

In line with these observations, the results from our Experiments 1 and 2 support the conclusion that

male mice pay significantly more attention to O+ than O-. In general, in these experiments, there

seems to be longer exploration of familiar objects compared to novel ones, which might be explained

be a slight neophobia in the mice. Taken together, these experiments confirmed our hypothesis that

female odour captures more attention than an odourless object. These experiments ensured that our

mice displayed behaviours as expected, based on the aforementioned literature, so we were able to

proceed to the next stages of the study.

The following aims were to elucidate how much attention odour would capture when the location of
O+ was either novel or familiar. Previous studies have already established that under normal
conditions, adult rats show preference for an object at a novel location compared to an identical object
at a previously experienced location (Aggleton, Albasser, Aggleton, Poirier, & Pearce, 2010; Barker
& Warburton, 2011; Ennaceur et al., 1997). Therefore, it was not surprising that in Experiment 3, O+
at a novel location attracted more attention than an O- at a familiar location, since both its odour and
location provided O+ with more motivational significance than O-. Interestingly, in Experiment 4,
where O+ at a familiar location competed with O- at a novel location, O+ still attracted more
attention. This represents a notable finding, as it shows that female odour 'wins' over a previously
known arousing factor (location novelty) in the amount of attention captured. In a human study of
auditory attention, Anderson (2016) demonstrated that VMAC by sounds previously associated with a
reward interfere with a visual task and compete with the visual representation of stimuli, reflecting
cross-modal stimulus competition bias by VMAC. Our study shows that, at least in mice performing a
NOR task, a positively stimulating odour outcompetes a visual stimulus (location novelty) for
attention, thus extending the findings from humans to animals and also to a different sensory domain.
467
At first glance, the observation that O+ at a novel location attracts more attention than O+ at a
familiar location might seem intuitive: in the first case, O+ consists of two motivational stimuli –
odour and location novelty, while in the second case O+ only has the odour. However, this also
emphasises an aspect worth taking into consideration – the fact that multiple stimuli can act in
combination to influence attention. Studies focusing on episodic memory have demonstrated that rats
and mice form integrated representations of three distinct object features: its identity, location and the
context in which it was experienced. Being able to associate these separate components allows the
animal to achieve a complex representation and record of environmental experiences, and this has
been termed 'episodic-like memory' in rodent models (Davis, Eacott, Easton, & Gigg, 2013; Eacott,
2004). In light of this work, the interpretation of how O+ at familiar location versus O+ at novel
location affects attention allocation could be further extended. The combination of odour and location
provides the animal with more detailed information about the object and, thus, it attracts more
attention than either odour or location alone. This suggests that odour can act not only on its own, but
also as a component of a stimulating context to modify the degree of attention allocation, thus
influencing other cognitive processes (e.g., episodic memory) in an additive manner.
483
Statistically, there was no day effect on the exploration time of O+ and O- in Experiments 3 and 4.
However, the daily means of exploration time indicate that there was a tendency for the attention
capture by O+ to decline across trials. This might be due to the male mice learning over the first three
trials that if they encounter female olfactory cues, this does not predict the availability of the female
mice, so the female odour starts to lose attentional allocation. This observation should be considered

in future behavioural studies using odours, particularly if such studies involve several trials. In our

study, using odour from different females after four consecutive trials was an attempt to avoid

habituation to odour.

Once we were able to determine that odour elicits more attentional capture than location novelty, the

next logical step was to study attention allocation to odour when competing with another powerfully

arousing stimulus – novel identity. The motivational value of novel identity has been the premise of

many studies using the NOR task, in which rodents are expected to pay more attention to the novel

rather than the familiar object. In our study, odour captured more attention than a novel object when

the location of the former was novel, while the location of the latter was familiar (Experiment 5). If

we assume that location is a visual stimulus, then odour in association with a visually arousing

stimulus captures more attention than another visual stimulus known to be salient to rodents.

However, in Experiment 6, when X was placed at a novel location and O+ at a familiar location,

odour could no longer 'win' over novel identity plus location in the degree of attention allocation.

These findings suggest that, on their own, odour and object identity might have the same motivational

significance to mice and their combination with other arousing stimuli can shift the balance in favour

of one over the other.

Nevertheless, it is not yet clear how visual and olfactory stimuli interact to capture the attention of

mice and whether the two kinds of stimuli are indeed equally motivational. A rather surprising

observation in Experiment 5 was that, in a single case, a particular X type attracted more attention

than its O+ counterpart; this indicates that in some circumstances, visual cues elicit stronger effects on

attention than olfactory cues. Le Pelley et al. (2016) argue that competition between a physically

salient and a less salient cue can alter attention, which can in turn affect the degree of learning. Since

both visual and olfactory cues convey important adaptive information to rodents, the question we

might ask is what determines whether a certain visual or olfactory cue attracts more attention than

other motivational stimuli present in the same environment. This question should motivate further

research into object type preference in mice, as well as more comparisons between the effects of

visual and olfactory cues on attention allocation.

A recent study in monkeys showed that for the visual system, novelty enhanced the motivational

value of stimuli associated with a negative outcome, while for the reward system, the effects of

novelty dissipated for such stimuli (Foley, Jangraw, Peck, & Gottlieb, 2014). This led the researchers

to conclude that novelty acts on attentional mechanisms independent of reward to influence the

processing of information and learning. In light of this finding, our experiments in mice could be

interpreted not just in terms of competition between novelty and a reward-associated stimulus, but

rather as evidence that reward and novelty are dissociable. In the future, it would be interesting to

investigate in humans whether this distinction can be made at both psychological and neurobiological
levels, especially since, contrary to this theory, prior research on animals has suggested that the
effects of novelty can be explained in terms of reward (Horvitz, 2000; Kakade & Dayan, 2002;
Laurent, 2008).
530

Several studies have demonstrated that the value of a particular stimulus feature (e.g., colour) in
predicting reward results in more attention being allocated to the same or similar features (Kalish &
Kruschke, 2000; Lawrence, Sahakian, Rogers, Hodges, & Robbins, 1999; Sutherland & Mackintosh,
1971). In our study, male mice were likely to evaluate female odour as endogenously motivationally
salient and, therefore, as an emotional stimulus. We are still uncertain as to whether naïve males
consider female odour to be reminiscent of their mothers and, thus, have a predictive value acquired
through learning. In the future, it would be useful to compare the degree of attention allocation to
female odours with that to odours or other stimuli that are predictive of an outcome (preferably a
rewarding outcome, since female odour is considered a positively arousing stimulus due to its effects
on reproductive behaviours).
541

542	Our study is currently ongoing and the next step is to compare the mice's attention allocation to
543	female odour-infused objects versus objects infused with a non-emotional odour. Depending on the
544	outcome of this experiment, we would be able to conclude whether mice can form emotionally-
545	charged memories that are different from memories associated with less-emotional odour stimuli.
546

In summary, the present study used object exploration measures to quantify the degree of attention
allocation by male mice to object novelty and/or female odour. Initial experiments demonstrated that
in the absence of other arousing features, objects with female odours capture more attention than an
odour-neutral object. These results agree with previous research showing that under laboratory
conditions, sexually isolated male mice are aroused by the odour of female conspecifics. The study
further demonstrated that odour 'wins' over location novelty in the degree of attention allocation and
that its motivational value is even greater in combination with location novelty. This supports the
conclusion that odour interacts with other arousing stimuli to form arousing contexts. Finally, odour
was able to capture more attention than a novel object, but only when combined with location novelty,
suggesting that the additive effects of visual and olfactory cues on attention exceed those of a single
strongly arousing stimulus. These experiments contribute to the understanding of the effects of female
odour on value-modulated attention capture and provide a reliable protocol to quantify attention
allocation in mice. The findings obtained here should encourage future research to use odour in
investigating the influence of emotional arousal on attention.
561
562

References

Aggleton, J.P., Albasser, M.M., Aggleton, D.J., Poirier, G.L., & Pearce, J.M. (2010). Lesions of the rat perirhinal cortex spare the acquisition of a complex configural visual discrimination yet impair object recognition. *Behavioral Neuroscience, 124*, 55-68. doi:10.1037/a0018320

Alkam, T., Hiramatsu, M., Mamiya, T., Aoyama, Y., Nitta, A., Yamada, K., Kim, H.C., & Nabeshima, T. (2011). Evaluation of object-based attention in mice. *Behavioural Brain Research, 220*, 185-193. doi:10.1016/j.bbr.2011.01.039

Anderson, B.A. (2016). Value-driven attentional capture in the auditory domain. *Attention Perception & Psychophysics, 78*, 242-250. doi:10.3758/s13414-015-1001-7

Anderson, B.A., Laurent, P.A., & Yantis, S. (2011). Value-driven attentional capture. *Proceedings of the National Academy of Sciences of the United States of America, 108*, 10367-10371. doi:10.1073/pnas.1104047108

Anderson, B.A., Laurent, P.A., & Yantis, S. (2014). Value-driven attentional priority signals in human basal ganglia and visual cortex. *Brain Research, 1587*, 88-96. doi:10.1016/j.brainres.2014.08.062

Antunes, M., & Biala, G. (2012) The novel object recognition memory: Neurobiology, test procedure, and its modifications. *Cognitive Processing,* 13, 93–110. doi:10.1007/s10339-011-0430-z

Aston-Jones, G., Rajkowski, J., Kubiak, P., Valentino, R.J., & Shipley, M.T. (1996). Role of the locus coeruleus in emotional activation. *Emotional Motor System, 107*, 379-402. Retrieved from https://doi.org/10.1016/S0079-6123(08)61877-4

Barker, G.R.I., & Warburton, E.C. (2011). When is the hippocampus involved in recognition memory?. *Journal of Neuroscience, 31*, 10721-10731. doi:10.1523/JNEUROSCI.6413-10.2011

Beny, Y., & Kimchi, T. (2014). Innate and learned aspects of pheromone-mediated social behaviours. *Animal Behaviour, 97*, 301-311. doi:10.1016/j.anbehav.2014.09.014

Berlyne, D.E. (1950). Novelty and curiosity as determinants of exploratory behaviour. *British Journal of Psychology, 41*, 68–80. doi:10.1111/j.2044-8295.1950.tb00262.x

Brennan, P.A., & Kendrick, K.M. (2006). Mammalian social odours: Attraction and individual recognition. *Philosophical Transactions of the Royal Society B-Biological Sciences, 361*, 2061-2078. doi:10.1098/rstb.2006.1931

Burke, S.N., Wallace, J.L., Nematollahi, S., Uprety, A.R., & Barnes, C.A. (2010). Pattern Separation Deficits May Contribute to Age-Associated Recognition Impairments. *Behavioral Neuroscience, 124*, 559-573. doi:10.1037/a0020893

Chelazzi, L., Estocinova, J., Calletti, R., Lo Gerfo, E., Sani, I., Della Libera, C., & Santandrea, E. (2014). Altering spatial priority maps via reward-based learning. *Journal of Neuroscience, 34,* 8594-8604. doi:10.1523/JNEUROSCI.0277-14.2014

Chun, M.M., Golomb, J.D., & Turk-Browne, N.B. (2011). A taxonomy of external and internal
attention. *Annual Review of Psychology,* 62, 73-101. doi:
10.1146/annurev.psych.093008.100427

Compton, R.J., 2003. The interface between emotion and attention: A review of evidence from
psychology and neuroscience. *Behavioral and Cognitive Neuroscience Reviews, 2,* 115–129.
doi:10.1177/1534582303255278

Connor, J. (1972). Olfactory control of aggressive and sexual behavior in mouse (Mus-musculus-L.).
Psychonomic Science, 27, 1-3. doi:10.3758/BF03328867

Corbetta, M., Patel, G., & Shulman, G.L. (2008). The reorienting system of the human brain: From
environment to theory of mind. *Neuron, 58*, 306-324. doi:10.1016/j.neuron.2008.04.017

Corbetta, M., & Shulman, G.L. (2002). Control of goal-directed and stimulus-driven attention in the
brain. *Nature Reviews Neuroscience, 3*, 201-215. doi:10.1038/nrn755

Davis, K.E., Eacott, M.J., Easton, A., & Gigg, J. (2013). Episodic-like memory is sensitive to both
Alzheimer's-like pathological accumulation and normal ageing processes in mice. *Behavioural*
Brain Research, 254, 73-82. doi:10.1016/j.bbr.2013.03.009

Desimone, R., & Duncan, J. (1995). Neural mechanisms of selective visual-attention. *Annual Review*
of Neuroscience, 18, 193-222. doi:10.1146/annurev.ne.18.030195.001205

Duffaud, A.M., Killcross, S., & George, D.N. (2007). Optional-shift behaviour in rats: A novel
procedure for assessing attentional processes in discrimination learning. *Quarterly Journal of*
Experimental Psychology, 60, 534-542. doi:10.1080/17470210601154487

Eacott, M.J., & Norman, G. (2004). Integrated memory for object, place, and context in rats: A
possible model of episodic-like memory?. *Journal of Neuroscience, 24*, 1948-1953.
doi:10.1523/jneurosci.2975-03.2004

Ennaceur, A. (2010). One-trial object recognition in rats and mice: Methodological and theoretical
issues. *Behavioural Brain Research, 215*, 244-254. doi:10.1016/j.bbr.2009.12.036

Ennaceur, A., Neave, N., & Aggleton, J.P. (1997). Spontaneous object recognition and object location
memory in rats: The effects of lesions in the cingulate cortices, the medial prefrontal cortex, the
cingulum bundle and the fornix. *Experimental Brain Research, 113*, 509-519.
doi:10.1007/pl00005603

Esber, G.R., & Haselgrove, M. (2011). Reconciling the influence of predictiveness and uncertainty on
stimulus salience: A model of attention in associative learning. *Proceedings of the Royal*
Society B-Biological Sciences, 278, 2553-2561. doi:10.1098/rspb.2011.0836

Failing, M.F., & Theeuwes, J. (2014). Exogenous visual orienting by reward. *Journal of Vision, 14*, 1-
9. doi:10.1167/14.5.6

Foley, N.C., Jangraw, D.C., Peck, C., & Gottlieb, J. (2014). Novelty enhances visual salience
independently of reward in the parietal lobe. *Journal of Neuroscience, 34*, 7947-7957.
doi:10.1523/jneurosci.4171-13.2014

Gaskin, S., Tardif, M., Cole, E., Piterkin, P., Kayello, L., & Mumby, D.G. (2010). Object
familiarization and novel-object preference in rats. *Behavioural Processes, 83*, 61-71.
doi:10.1016/j.beproc.2009.10.003

George, D.N., & Pearce, J.M. (1999). Acquired distinctiveness is controlled by stimulus relevance not
correlation with reward. *Journal of Experimental Psychology-Animal Behavior Processes, 25*,
363-373. doi:10.1037/0097-7403.25.3.363

Gitelman, D.R. (2003). Attention and its disorders. *British Medical Bulletin, 65*, 21-34.
doi:10.1093/bmb/65.1.21

Golomb, J.D., Nguyen-Phuc, A.Y., Mazer, J.A., McCarthy, G., & Chun, M.M. (2010). Attentional
facilitation throughout human visual cortex lingers in retinotopic coordinates after eye
movements. *Journal of Neuroscience, 30*, 10493-10506. doi:10.1523/jneurosci.1546-10.2010

Hartikainen, M.K., Ogawa, H.K., Soltani, T.M., & Knight, T.R. (2007). Emotionally arousing stimuli
compete for attention with left hemispace. *NeuroReport, 18*, 1929-1933.
doi:10.1097/WNR.0b013e3282f1ca18

Horvitz, J.C. (2000). Mesolimbocortical and nigrostriatal dopamine responses to salient non-reward
events. *Neuroscience, 96*, 651–656. doi:10.1016/S0306-4522(00)00019-1

Johnston, R.E. (2003). Chemical communication in rodents: From pheromones to individual
recognition. *Journal of Mammalogy, 84*, 1141-1162. doi:10.1644/BLe-010

Kadohisa, M. (2013). Effects of odor on emotion, with implications. *Frontiers in Systems*
Neuroscience, 7, 66. doi:10.3389/fnsys.2013.00066

Kakade, S., & Dayan, P. (2002). Dopamine: generalization and bonuses. *Neural Networks, 15*, 549 –
559. doi:10.1016/S0893-6080(02)00048-5

Kalish, M. L., & Kruschke, J. K. (2000). The role of attention shifts in the categorization of
continuous dimensioned stimuli. *Psychological Research-Psychologische Forschung, 64,* 105–
116. doi:10.1007/s004260000028

Katsuki, F., & Constantinidis, C. (2014). Bottom-up and top-down attention: Different processes and
overlapping neural systems. *Neuroscientist, 20*, 509-521. doi:10.1177/1073858413514136

Kaye, H., & Pearce, J.M. (1984). The strength of the orienting response during Pavlovian
conditioning. *Journal of Experimental Psychology-Animal Behavior Processes, 10*, 90-109.
doi:10.1037/0097-7403.10.1.90

Laurent, P.A. (2008). The emergence of saliency and novelty responses from reinforcement learning
principles. *Neural Networks, 21*, 1493–1499. doi:10.1016/j.neunet.2008.09.004

Lawrence, A. D., Sahakian, B. J., Rogers, R. D., Hodges, J. R., & Robbins, T. W. (1999).
Discrimination, reversal, and shift learning in Huntington's disease: Mechanisms of impaired
response selection. *Neuropsychologia, 37,* 1359 –1374. doi:10.1016/S0028-3932(99) 00035-4

Le Pelley, M.E., Mitchell, C.J., Beesley, T., George, D.N., & Wills, A.J. (2016). Attention and
associative learning in humans: An integrative review. *Psychological Bulletin, 142*, 1111-1140.
doi:10.1037/bul0000064

Lee, T.-H., Sakaki, M., Cheng, R., Velasco, R., & Mather, M. (2014). Emotional arousal amplifies the
effects of biased competition in the brain. *Social Cognitive and Affective Neuroscience, 9*,
2067-2077. doi:10.1093/scan/nsu015

Levine, L.J., & Edelstein, R.S. (2009). Emotion and memory narrowing: A review and goal-relevance
approach. *Cognition & Emotion, 23*, 833-875. doi:10.1080/02699930902738863

Mackintosh, J.H. (1970). Territory formation by laboratory mice. *Animal Behaviour, 18*, 177-183.
doi:10.1016/0003-3472(70)90088-6

Mackintosh, N. J. (1975). A theory of attention: Variations in the associability of stimuli with
reinforcement. *Psychological Review, 82,* 276– 298. doi:10.1037/h0076778

McGaughy, J., & Sarter, M. (1998). Sustained attention performance in rats with intracortical
infusions of 192 IgG-saporin-induced cortical cholinergic deafferentation: Effects of
physostigmine and FG 7142. *Behavioral Neuroscience, 112*, 1519-1525. doi:10.1037/0735-
7044.112.6.1519

Michael, G.A., Jacquot, L., Millot, J.L., & Brand, G. (2003). Ambient odors modulate visual
attentional capture. *Neuroscience Letters, 352*, 221-225. doi:10.1016/s0304-3940(03)01078-4

Miller, E.K., & Cohen, J.D. (2001). An integrative theory of prefrontal cortex function. *Annual
Review of Neuroscience, 24*, 167-202. doi:10.1146/annurev.neuro.24.1.167

Mumby, D.G., Glenn, M.J., Nesbitt, C., & Kyriazis, D.A. (2002). Dissociation in retrograde memory
for object discriminations and object recognition in rats with perirhinal cortex damage.
Behavioural Brain Research, 132, 215-226. doi:10.1016/s0166-4328(01)00444-2

Murai, T., Okuda, S., Tanaka, T., & Ohta, H. (2007). Characteristics of object location memory in
mice: Behavioral and pharmacological studies. *Physiology & Behavior*, *90*, 116-124.
doi:10.1016/j.physbeh.2006.09.013

Ohman, A., & Mineka, S. (2001). Fears, phobias, and preparedness: Toward an evolved module of
fear and fear learning. *Psychological Review, 108*, 483-522. doi:10.1037//0033-295x.108.3.483

Oliveira, A.M.M., Hawk, J.D., Abel, T., & Havekes, R. (2010). Post-training reversible inactivation
of the hippocampus enhances novel object recognition memory. *Learning & Memory, 17*, 155-
160. doi:10.1101/lm.1625310

Pantoja, J., Ribeiro, S., Wiest, M., Soares, E., Gervasoni, D., Lemos, N.A.M., & Nicolelis, M.A.L.
(2007). Neuronal activity in the primary somatosensory thalamocortical loop is modulated by
reward contingency during tactile discrimination. *Journal of Neuroscience, 27*, 10608-10620.
doi:10.1523/JNEUROSCI.5279-06.2007

Pearce, J.M., & Hall, G. (1980). A model for Pavlovian learning: Variations in the effectiveness of
conditioned but not of unconditioned stimuli. *Psychological Review, 87*, 532-552.
doi:10.1037/0033-295x.87.6.532

Pessoa, L. (2009). How do emotion and motivation direct executive control?. *Trends in Cognitive*
Sciences, 13, 160-166. doi:10.1016/j.tics.2009.01.006

Piper, B.J., Fraiman, J.B., & Meyer, J.S. (2005). Repeated MDMA ("Ecstasy") exposure in adolescent
male rats alters temperature regulation, spontaneous motor activity, attention, and serotonin
transporter binding. *Developmental Psychobiology, 47*, 145-157. doi:10.1002/dev.20085

Pourtois, G., Schettino, A., & Vuilleumier, P. (2013). Brain mechanisms for emotional influences on
perception and attention: What is magic and what is not. *Biological Psychology, 92*, 492-512.
doi: 10.1016/j.biopsycho.2012.02.007

Pratt, J., & Hommel, B. (2003). Symbolic control of visual attention: The role of working memory
and attentional control settings. *Journal of Experimental Psychology-Human Perception and*
Performance, 29, 835-845. doi:10.1037/0096-1523.29.5.835

Qi, S.Q., Zeng, Q.H., Ding, C., & Li, H. (2013). Neural correlates of reward-driven attentional
capture in visual search. *Brain Research, 1532*, 32-43. doi:10.1016/j.brainres.2013.07.044

Sarter, M., Givens, B., & Bruno, J.P. (2001). The cognitive neuroscience of sustained attention:
Where top-down meets bottom-up. *Brain Research Reviews, 35*, 146-160. doi:10.1016/s0165-
0173(01)00044-3

Shapiro, K.L., Egerman, B., & Klein, R.M. (1984). Effects of arousal on human visual dominance.
Perception & Psychophysics, 35, 547-552. doi:10.3758/bf03205951

Silvers, J.M., Harrod, S.B., Mactutus, C.F., & Booze, R.M. (2007). Automation of the novel object
recognition task for use in adolescent rats. *Journal of Neuroscience Methods, 166*, 99-103. doi:
10.1016/j.jneumeth.2007.06.032

Spence, C. (2010). Crossmodal spatial attention. *Annals of The New York Academy of Sciences, 1191*,
182-200. doi:10.1111/j.1749-6632.2010.05440.x

Sutherland, N. S., & Mackintosh, N. J. (1971). *Mechanisms of animal discrimination learning*. New
York, NY: Academic Press

Talmi, D. (2013). Enhanced emotional memory: cognitive and neural mechanisms. *Current*
Directions in Psychological Science, 22, 430-436. doi:10.1177/0963721413498893

Talmi, D., & McGarry, L.M. (2012). Accounting for immediate emotional memory enhancement.
Journal of Memory and Language, 66, 93-108. doi:10.1016/j.jml.2011.07.009

Talmi, D., Slapkova, M., & Wieser, M. (2018). Testing the possibility of model-based Pavlovian
control on attention to threat. *Journal of Cognitive Neuroscience, 31*, 36-48. doi:
10.1162/jocn_a_01329

Vermeulen, N., Godefroid, J., & Mermillod, M. (2009). Emotional modulation of attention: Fear
increases but disgust reduces the attentional blink. *Plos One, 4*, 5.
doi:10.1371/journal.pone.0007924

Vuilleumier, P. (2005). How brains beware: Neural mechanisms of emotional attention. *Trends in*
Cognitive Sciences, 9, 585-594. doi:10.1016/j.tics.2005.10.011

Wang, L.H., Duan, Y.Y., Theeuwes, J., & Zhou, X.L. (2014). Reward breaks through the inhibitory
region around attentional focus. *Journal of Vision, 14*, 7. doi:10.1167/14.12.2

Wilson, P.N., Boumphrey, P., & Pearce, J.M. (1992). Restoration of the orienting response to a light
by a change in its predictive accuracy. *Quarterly Journal of Experimental Psychology Section*
B-Comparative and Physiological Psychology, 44, 17-36. doi:10.1080/02724999208250600

Yantis, S., Schwarzbach, J., Serences, J.T., Carlson, R.L., Steinmetz, M.A., Pekar, J.J., & Courtney,
S.M. (2002). Transient neural activity in human parietal cortex during spatial attention shifts.
Nature Neuroscience, 5, 995-1002. doi:10.1038/nn921

Zaborszky, L., Gaykema, R.P., Swanson, D.J., & Cullinan, W.E. (1997). Cortical input to the basal
forebrain. *Neuroscience, 79*, 1051-1078. doi:10.1016/s0306-4522(97)00049-3

Wentura, D., Muller, P., & Rothermund, K. (2014). Attentional capture by evaluative stimuli: Gain-
and loss-connoting colors boost the additional- singleton effect. *Psychonomic Bulletin &*
Review, 21, 701–707. doi:10.3758/s13423-013-0531-z

759
760