Historical abundance and distributions of *Salpa thompsoni* hot spots in the Southern

Ocean, with projections for further ocean warming

Angelika Wanda Słomska[1]*, Anna Panasiuk[1], Agata Weydmann-Zwolicka[1], Justyna Wawrzynek-Borejko[2], Marta Konik[3], Volker Siegel[4]

[1] University of Gdansk, Department of Marine Plankton Research, Faculty of Oceanography and Geography, 46 M. J. Pilsudskiego Avenue, Gdynia, Poland

[2] University of Gdansk, Department of Marine Ecosystem Functioning, Faculty of Oceanography and Geography, 46 M. J. Pilsudskiego Avenue, Gdynia, Poland

[3] The Institute of Oceanology of the Polish Academy of Sciences, 55 Powstancow Warszawy Street, Sopot, Poland

[4] Sea Fisheries Institute, 9 Palmaille, 22767 Hamburg, Germany

Corresponding Author:

Angelika Słomska[1]

46 M. J. Pilsudskiego Avenue, Gdynia, Pomorskie, 81-378, Poland

Email address: angelika.slomska@ug.edu.pl

Phone: +48 58 523 68 46

Fax: +48 58 523 66 78

ABSTRACT

1. Over the last three decades, a significant variability in *Salpa thompsoni* occurrence has

been observed as a response to the environmental fluctuations of the Southern Ocean ecosystem,

e.g. changes in sea surface temperature as well as shrinking of ice-cover extent around the cold

Antarctic waters.

2. This study presents the historical data of salps abundance from the southwest Atlantic

Sector of the Southern Ocean and covers time span of 20 years. Presented dataset allowed to

track previous fluctuations in Antarctic salp abundance and enabled to combine their distribution

with bottom depth, thermal and ice conditions. The subsequent goal of this work was to reveal

hot spots of salps location and to predict the future range of *S. thompsoni* distribution with

upcoming climate warming in the next 50 years.

3. Results of our study revealed that the highest salp number was located mostly in the

shallow shelf waters with ice-cover and lower temperature. In the studied area, *Salpa thompsoni*

hot spot distributions have been located mostly around Elephant Island but also within islands

around Brensfield and Gerlache Straits, as well as to the south near the cold Bellingshausen Sea.

The inference of future salp distribution demonstrated that the range of *S. thompsoni* would

presumably move southwards enlarging their habitat area by nearly 500 000 km^2.

Key words: climate change, environmental impact assessment, invertebrates, migration, ocean,
zooplantkon

 INTRODUCTION

Over the last 30 years, the rise of atmospheric greenhouse gas has influenced the average

global atmospheric temperature, with an increase of 0.2°C, and the mean global sea surface

temperature (SST) increasing by approximately 0.4°C (Levitus, Antonov & Boyer, 2005;

Levitus, Antonov, Boyer & Stephens, 2000). Ocean heating is particularly pronounced in the

most vulnerable areas which include the Southern Ocean, especially in the Antarctic Peninsula

region (Whitehouse, Meredith, Rothery, Atkinson, Ward & Korb, 2008; Zwally, Comiso,

Parkinson, Cavalieri & Gloersen, 2000). Despite this, study by Turner et al. (2016) based on the

stacked air temperature record since the 1990s, has suggested that there was no evidence

of climate warming in the mentioned area. However, hydrological processes and modifications

recorded in the Southern Ocean are much more complex. It should be recalled that clear proof

of decreasing sea ice extent in the Bellingshausen and Amundsen Seas has been numerously

recorded, and this phenomenon could be tied to continuous ocean heating (Cavalieri

& Parkinson, 2012). In particular, summer water temperature along the Western Antarctic

Peninsula increased by 1.3°C over 50 years (Meredith & King, 2005), and near South Georgia

a 1°C increase has been recorded over the last 80 years (Whitehouse, Meredith, Rothery,

Atkinson, Ward & Korb, 2008). The Weddell Deep Water (WDW) has warmed by ~0.032°C per

decade, which is similar to the temperature changes around South Georgia and in the entire

Antarctic Circumpolar Current (0.03–0.07°C per decade) (Nicol, Croxall, Trathan, Gales

& Murphy, 2007; Whitehouse, Meredith, Rothery, Atkinson, Ward & Korb, 2008).

The current phenomenon of environmental stress caused by climatic and anthropogenic

pressures leads to shifts of frontal hydrological zones, existing habitats, and is also likely

to modify structure, spatial range and seasonal abundance of Antarctic key species (Atkinson,

Siegel, Pakhomov & Rothery, 2004; Loeb & Santora, 2012; Pakhomov, Froneman

& Perissinotto, 2002; Richardson, 2008;; Ross et al., 2014; Steinberg et al., 2015).

The geographical range of crucial Antarctic taxa can move southwards to remain in optimal

thermal conditions, squeezing their distribution range closer to the Antarctic continent (Atkinson,

Siegel, Pakhomov & Rothery, 2004; Loeb & Santora, 2012; Richardson, 2008; Ross et al.,

2014). At the same time, more favorable habitats can be opened for more flexible organisms that

are capable of adaptation like gelatinous salps (Goodall-Copestake, 2016; Jue et al., 2016;

Mackey et al., 2011; Ross et al., 2014; Steinberg et al., 2015).

Antarctic salp *Salpa thompsoni* (Foxton, 1961), together with the krill *Euphausia* superba

(Dana, 1850), are among the most important filter-feeding species of the Southern Ocean, but

only Antarctic krill is considered as a major food source for many top predators, including fish,

penguins, seals and baleen whales in a very short Antarctic trophic chain (Loeb & Santora,

2012). Therefore, regarding the current threat to decreasing krill numbers, and due to a changing

climate conditions, there is a growing concern that salps may locally replace krill. Possible

mechanisms underlying these observations include the southerly movement of the sea ice edge,

water temperature fluctuations, reconstruction of the phytoplankton structure, and changes

in Antarctic Circumpolar Current transport pathways. All these factors might cause shifts

in Antarctic krill populations, and create a free ecological niche available for pelagic tunicates.

Salpa thompsoni is typically oceanic, has an extensive circumpolar distribution (45-55°S)

(Foxton, 1966; Pakhomov, Froneman & Perissinotto, 2002), is not ice-dependent like *Euphausia*

superba and it is usually found in areas with lower food concentrations (Atkinson, Siegel,

Pakhomov & Rothery, 2004; Loeb et al., 1997; Pakhomov, Froneman & Perissinotto,

2002;Siegel & Loeb, 1995; Steinberg et al., 2015). Antarctic salps prefer water masses of higher

temperature and lower productivity across the Circumpolar Current, therefore the highest salp

abundance is observed along with the absence or at least really low number of Antarctic krill

(Atkinson, Siegel, Pakhomov & Rothery, 2004; Siegel & Loeb, 1995). Previous studies provided

some evidence that the greatest changes in abundance and distribution of salps in the Atlantic

Southern Ocean are associated with the sea ice loss, transitional periods between

El Niño-La Niña (ENSO), and with the shifts of the Southern Boundary (SO) of Antarctic

Circumpolar Current (ACC) (Atkinson, Siegel, Pakhomov & Rothery, 2004; Chiba, Ishimaru,

Hosie & Wrigh, 1999; Pakhomov, Froneman & Perissinotto, 2002; Siegel & Loeb, 1995).

The primary focus of these presented studies was to identify the main barriers hindering

the salps occurence and provide missing predictions of the salp population dynamics. Based on

the historical point measurements of *Salpa thompsoni* abundance in the Atlantic Sector of the

Southern Ocean within over twenty years of time span (1975-2001) a spatial distribution of salp

populations was reconstructed. Using the Hot Spot analyses, it revealed the environmental

conditions controlling their propagation in the investigated area with an insight into

the upcoming 50 years in the age of global warming.

METHODS

Study area

The study area covered the region of the Western Antarctic Peninsula and South West

(SW) Atlantic sector of the Southern Ocean (Fig. 1).

The waters flowing around the Antarctic Peninsula are a specific and unique mixture of

various water masses (Martinson et al., 2008). Here we can observe the mixing of warm, nutrient

rich Circumpolar Deep Water (CDW) with cold, Antarctic shelves and coastal water masses,

which have important effects on physical and biological processes (Ducklow et al., 2012).

Antarctic Peninsula (AP) water mass distribution is restricted by bathymetric features such as

continental slope, coastal or shelf regions (Martinson, Stammerjohn, Iannuzzi, Smith & Vernetd,

2008). This region is also more susceptible to the influence of the Antarctic Circumpolar Current

(ACC) than other regions around the Antarctic where the ACC is rather distant from the shelf,

typically separated by a polar gyre (Martinson, Stammerjohn, Iannuzzi, Smith & Vernetd, 2008).

The marine ecosystems of the studied area are characterized by high diatom concentrations and

consequently by a high primary production, krill predominance and a variety of higher vertebrate

consumers (Siegel, Skibowski & Harm, 1992). Macronutrient concentration, such as nitrate and

phosphate, are generally within the AP region compared to more open waters, due to three

factors - high concentrations in deep water, winter deep water mixing resupplying the surface

layer after biological depletion, and micronutrient (iron) limitation (Ducklow et al., 2012).

Sampling

Samples were collected during the German and British research surveys (RV 'Walther

Herwig', 'John Biscoe', 'Polarstern' and 'Meteor' cruises) using a Rectangular Midwater Trawl

(RMT8) (mesh size 4.5 mm, mouth opening of 8 m^2) (Baker, Clarke & Harris, 1973) between

1975 and 2001, during early summer/late autumn. The RMT8 was equipped with a real-time

depth recorder (TDR) and sampled from the upper 200 m of the water column or 10 m above

the bottom, in shallow areas. A double oblique net tow was carried out routinely at all stations

with a standard station grid at a tow speed of 2.5 to 0.5 knots. Calibrated flowmeters, mounted

on the net frame, were used to estimate the volume of filtered water during each haul. Filtered

water volume was calculated using equations from Pommeranz, Hermann & Kühn (1992).

All samples from the surveys were processed at sea. Such approach led to collecting 1872

samples in total, out of which the salps were removed immediately after the tow, counted prior to

other sample processing and were stored in 4% buffered formaldehyde for later measurements.

Water temperature (T) and salinity (S) were measured prior to sampling (upper 200 m), however

this data was available only for a limited number of samples: T for 461 samples and S for 212

samples. Ice coverage was estimated from the bridge of the vessel by visual observation and by

checking the radar, to assess the ice extended for more than one mile around the vessel. The Ice

coverage was expressed in the % coverage to four classifications 0, 1, 2 and 3, where 0 means no

ice, 1 <15%, 2 < 50% and 3 > 50 % of ice cover within one nautical mile.

Reanalysis datasets

Temperature and salinity were not measured during all sampling surveys therefore

we compensated our dataset for years 1970 – 2001 using available complete monthly product

of the Sea Surface Temperature (SST). Due to the fact that in the Antarctic area persistent cloud

cover limits available satellite information we used results of reanalyses HadISST1 published

by the Met Office Hadley Centre for Climate Prediction and Research. The continuity of the SST

maps had been achieved due to use of the Reduced Space Optimal Interpolation (RSOI)

technique. The time series covered ranges from 1871 to date and it was based on the *in situ*

observations. However, data collected in the Met Office Marine Data Bank (MDB) had been

quality controlled e.g. corrected for bias and gridded onto a 1° area grid. Auxiliary data were

included in order to increase quality of the product such as the monthly median SSTs from the

Comprehensive Ocean-Atmosphere Data Set (COADS) which were incorporated in to the

HadISST1 product for years 1871 – 1995. Since 1982 information from the satellite AVHRR

sensor has also been incorporated into the dataset in order to significantly increase spatial data

coverage and thus product quality (Rayner et al., 2003).

Information about the sea ice cover was obtained from the National Snow and Ice Data

Center (NSIDC). The 'Sea Ice Extent' product (ID: G02135) included in this study was

downloaded in the form of complete time-series from the November 1978 to the present. It was

created at the Goddard Space Flight Center (GSFC). The information about sea ice was derived

from passive microwave satellites the Nimbus-7 Scanning Multichannel Microwave Radiometer

(SMMR), the Special Sensor Microwave/Imager (SSM/I) since July 1987, and the Special

Sensor Microwave Imager/Sounder (SSMIS) instruments onboard satellites launched as part

of the Defense Meteorological Satellite Program (DMSP). The data covered the Southern

Hemisphere and were bounded on the north by the parallel of 39.23º southern latitude at spatial

resolution of 25 km x 25 km (Fetter et al., 2017). The sea ice presence was expressed in

a dichotomic (binary) way, where: 1 refers to sea ice cover present and 0 refers to no sea

ice station. The threshold for the ice sheet was adopted as the point, where the average sea ice

concentration in that month drops below 15 percent which is consistent with the operational

definition presented in the IPCC reports (Vaughan et al., 2013; Fetterer et al., 2017).

Data analysis

In order to test possible correlations between significant environmental variables,

the nonparametric Spearman's rank-order correlations were calculated prior analyses. Due to the

log-normal distribution of the data, further analyses were performed on the log-transformed

abundance data [x'=log (x+10)] using STATISTICA 12.0 PL (Statsoft Inc.) software. To reveal

if the sea ice concentration both during sampling and in the preceding winter season (reanalysis

dataset) had any impact on the number of salps during the closest Antarctic summer, the analysis

of variance ANOVA (Kruskal Wallis test) and a probit regression model was used, respectively.

Additionally, the Generalized Additive Models (GAMs, Poisson distribution) implemented

in CANOCO 5 software were used to examine the response of salps abundance by determining

their most probable spatial distribution in relation to sea surface temperature, bottom depth, and

salinity, which were tested separately due to the varying number of available in situ

measurements of bottom depth, T and S (1872, 461 and 212 samples, respectively).

Spatial analyses

The Global Moran's I statistic was used to test whether data are dispersed or there is

a similarity between the collected samples. It was chosen, because The Global Moran's I Index

assesses the spatial autocorrelation based on values as well as features locations. Z-score

statistics and p-value are computed along with the index in order to estimate the significance

of the result. A positive Moran's I Index value proves existence of the clusters, whereas

a negative shows tendency towards dispersion (Moran, 1950). As the statistics pointed to strong

clustering of the data (z-score, representing measures of standard deviation: > 60,

p-value: <0.01) they were divided into three subsets based on the decadal time periods. The

division was based on spatial distribution of the sampling points, as the research stations were

placed over a vast area, and were confirmed as being statistically significant (z-score: >98,

p-value: <0.01). Therefore, both ice cover interpolation and hot spot analysis were applied

separately to each decade '70s, '80s, and '90s (including year 2001, because of its similarity

to the cruises from 94/95 and 96/97). In order to interpolate ice coverage, the Kernel

Interpolation with Barriers method was performed. The coastline of the Antarctic Peninsula from

Antarctic Digital Database (ADD) was used as a barrier in the analysis. The kernel function was

set to first order polynomial with the ridge parameter retaining the default value of 50.

The hot and cold spot analysis was conducted using the Hot Spot Analysis (Getis – Ord

Gi*) tool (Getis and Ord, 1992). The Getis-Ord Gi* statistic was chosen over more global

statistics (e.g. The Global Moran's I Index) to identify more local spatial clusters by finding local

maxima. It was possible to compare of each feature with the surrounding neighbours (Getis

& Ord, 1992). This analysis computes by a default a z-score (a measure of standard deviation)

and p-value. A high z-score and small value of p indicate clustering of high values (hot spots),

whilst low z-score with small p-value inform about spatial clustering of lower values (cold

spots). Point data was exploited to detect the hot spot areas with significantly high salp densities

and cold spots with their lower abundance. In the analysis we used inverse distance band in order

to assure use of every point in the analysis in the decadal subsets.

The future boundary of temperature optimum for the distribution range of salps was

established based on HadISST1 data from the Met Office Hadley Centre's (Rayner et al., 2003).

Mean distribution of surface temperatures during local summer (from December to the beginning

of March) was created by mosaicking SST rasters for the years 1970 to 2001. The first boundary

of salp occurence based on thermal preferences was determined by extracting the periphery

of 0°C isotherm. Temperature range for salp occurrence was based on available literature data

(Foxton, 1966; Ono & Moteki, 2016). The predicted boundary after a temperature increase

of 1°C was extracted analogically from the periphery of 1°C isotherm. According to the previous

studies, water temperature along the Western Antarctic Peninsula has increased by more than

1ºC over the last 50 years (Whitehouse et al., 2008). We adopted a conservative approach and

assumed that water temperature will increase by the next degree over the 50 years ahead. The

maps presenting spatial distribution and abundances of *Salpa thompsoni* were created using

ESRI[®] ArcMap[TM] 10.5.1 software.

RESULTS

The extended collection resulted in 1872 samples, of which *Salpa thompsoni* was present

in 1278. Our results from the long-term data series revealed significant annual variability

in salp abundance. Through the late 1970s and early 1980s mean abundances

of *S. thompsoni* were relatively low in researched areas, with the exceptions of the cruises in

1975/76 and 1983/84. After season 1987/88, with very low salps densities (0.9 ind/1000m^3

±1.26), the extremely high peak abundance occurred in the summer 1989/90, with the mean

density 1408 ind/1000m^3 (±2778.31). Although season 1994/95 was characterized by low salp

abundances, their numbers began to increase over the following years (Fig. 2).

Analysis based on in situ data also confirmed that the high abundance of salps was

influenced by the sea ice concentration in the time of sampling (Kruskal-Wallis test H=33.98,

p<0.0001). Sea ice played a crucial role in the Antarctic salp distribution, however significant

differences were observed only between stations without ice cover (0) and with the presence of

ice (1-3). There was no statistically significant difference between stations with a different

degree of ice cover, which could affect presence of salps. Further analyzes have implied that

S. thompsoni may occur in less favorable ice conditions (ice-cover around 1-3 degree) in smaller

numbers. Interestingly, the probit model for the Sea Ice Index data, confirmed that the ice

concentration during winter months (July and August) had a significant impact on the presence

of salps in the following summer season (N=1311, $\chi2$ =211.36, p<0.0001).

The nonparametric Spearman's rank-order revealed only weak correlations between selected

environmental variables (bottom depth vs T rs=0.2, bottom depth vs sea ice rs=-0.25, and sea ice

vs T rs=-0.16), therefore Generalized Additive Models (GAMs, Fig. 3) were tested

independently and separately for above environmental variables. GAM fitted for sea surface

temperature (F=5.3, p=0.0054) showed a uni-modal response, with the highest number of

S. thompsoni registered in the water layers with temperatures between 1°C and -1°C (Fig. 3).

GAM against salinity (F=5.8, p=0.0039) and bottom depth (F= 15.0, p<0.00001) did not give

such a clear response. The highest numbers of salps seemed to be observed within waters with a

salinity around 33.5-34.25. The same model used for the bottom depth showed the presence of

the major peak, indicating preference for shallow shelf waters with bottom depth around 0-1000

m (Fig. 3).

Hot spot Analysis

Despite the fact that sampling sites in the 1970's were spread along the West Antarctic

Peninsula region, near the South Shetland Islands, around South Orkney Islands in the Atlantic

sector of the Southern Ocean, statistically significant hot spots were located mostly near

Elephant Island, where ice concentration was degree 0 (Z-score > 1.67 and p value < 0.05).

In the 1980's a hot spot was registered further to the south. Regardless of the similarity between

sampling coverages of WAP area, hot spots (Z-score > 1.7 and p value < 0.05) were placed in

regions of the Biscoe Islands, as well as Anvers Island and through the Bransfield Strait (Fig. 4).

Significant high densities of salps, assuming 99% confidence, were still found in the vicinity

of Elephant Island. The stations with high abundance of salps were in the areas with ice coverage

classified as 1. Sampling stations in seasons 1994/95, 1996/97 and 2001 were clustered around

the Elephant Island region. This region showed *S. thompsoni* hot spots, with a high Z-score (up

to 7.89) and p value <0.05.

DISCUSSION

Our results revealed significant annual variability of *Salpa thompsoni* due to differences

in bottom depth and environmental conditions e.g. water temperature or ice condition prevailing

during and before sampling years. Presented data also indicated that the crucial months for salps

were July and August when the lack of sea ice presumably allows these animals to form more

vast and condensed aggregations during the following summer periods. Also, the previous

studies revealed that salps were most frequently present in the ice-free regions (Atkinson, Siegel,

Pakhomov & Rothery, 2004; Chiba, Ishimaru, Hosie & Wrigh, 1999; Lee, Pakhomov, Atkinson

& Siegel, 2010; Mackey et al., 2011; Ross et al., 2014; Steinberg et al., 2015), which our results

initially confirmed. Although, our further analysis showed that *S. thompsoni* may be present even

with less suitable ice conditions, 1-2 ice degree, which means that they might be able to exist

in colder environment. Used GAM model, revealed untypical salp preferences, different than

previously noted, showing that the highest salp number was located in the shallow shelf waters

(bottom depths 0-1000 m), with a low temperature ([-1] – 1°C), and salinity of 33.5 – 34.25

(Fig. 3). Our observation was contrary to the conclusions of Atkinson, Siegel, Pakhomov

& Rothery (2004) who believed that cold shelf water is a place occupied by krill but will not be

preferred by Antarctic salps. This finding may suggest possible shifts of salp densities closer

to the cold continent, or could be partially explained by the dynamics of the ACC water masses.

The ACC pumps warm (>1.5°C), salty (34.65–34.7), and nutrient-rich Circumpolar Deep Water

(CDW) onto the continental shelf below 200 m (Klinck, Hofmann, Beardsley, Salihoglu

& Howard, 2004), and may moves salps towards the Antarctic continent and the surrounding

islands to the Bransfield Strait areas (Tokarczyk, 1987), where they can be found in high

abundances.

The Hot Spots Analysis in three decadal time periods - '70s, '80s, and '90s - were

confirmed that the *Salpa thompsoni* distribution has been located mostly around Elephant Island.

Our dataset from '70s and '90s showed that their greatest numbers were observed mostly within

open ocean as well as around the AP area, however, in the '80s *S. thompsoni* hot spots were

located evenly within Islands around Bransfield and Gerlache Straits, as well as far south near

the Bellingshausen Sea, where sea ice was present and water temperature was lower.

The greatest salp number was recorded between 65-70°S in the presence of ice-cover and water

temperature below 0°C, which can also confirm previous observations by Ono & Moteki (2013),

and may suggest that *S. thompsoni* is able to adapt to less suitable conditions in colder, shelf

waters. S. *thompsoni* generally occurs in warm water masses > 0 °C, between 45-55°S (Foxton,

1966). Results of this study showed that a high number of mature individuals could be observed

even south of the Southern Boundary (SB). Our results, together with the study presented by Ono

& Moteki (2013), has revealed that *S. thompsoni* appeared across the SB of the ACC,

and suggests that presumably it is not a barrier preserving Antarctic waters from *S. thompsoni*

southward migration. A number of genetic studies further proved that salps are highly flexible

(Batta-Lona, Maas, O'Neill, Wiebe & Bucklin, 2017; Goodall-Copestake, 2016; Jue et al., 2016),

capable of adaptation to the changing environmental conditions within different parts of the

Southern Ocean. The comprehensive data about modification of salps/krill population were

presented by Ross et al. (2014). They showed that krill abundance had decreased on the northern

lines and had shifted 200 km to the south. If the trend of shrinking sea ice continues, there

is a certain risk that within two decades, krill may completely disappear from the northern region

of the Southern Ocean. Herein presented result expands foregoing knowledge about the

dynamics of Antarctic salp populations and demonstrates missing predictions of their future

distribution. In our study, we have presented inference about possible future salps distribution

modification with the assumption that the water temperature will increase by 1°C, and results

showed that the *S. thompsoni* range could move southwards by an average of 200 km.

Taking into account climate fluctuations and temperature trends, we tracked the movement of the

temperature boundary for salps for a hypothetical situation of ocean warming by 1°C for the next

50 years. The comparison of the locations of isotherms 0°C and 1°C revealed that massive

blooms of *S. thompsoni* possible may shift their distribution southward e.g. into the Weddell Sea

and closer to the Antarctic continent. Such temperature rise would enlarge their habitat area

by nearly 530 000 km^2, what possibly may exclude Antarctic krill and consequently lead to

changes in the functioning of the Antarctic ecosystem (Fig. 5).

ACKNOWLEDGMENTS

We want to express our gratitude to the crews and scientists who have collected

and analysed these samples over the last 3 decades, and provided this data in a useable format.

We are grateful to Prof. Evgeny Pakhomov from University of British Columbia, M.Sc. Natasha

Waller from Australian Antarctic Division for their willingness and help in improving this

manuscript. We would like to thank Dr. Adrian Zwolicki from the University of Gdansk for his

advice concerning statistical analyses. The authors declare that they have no conflict of interest.

FUNDING SOURCES

This study was supported by the Polish National Science Centre, Project PRELUDIUM no

2016/23/N/NZ8/02801 received by Angelika Słomska. In addition, the work was partly

supported by the external project Miniatura 1 no 2017/01/X/NZ8/01181.

REFERENCES

Atkinson, A., Siegel. V., Pakhomov. E., & Rothery P. (2004). Long-term decline in krill stock

and increase in salps within the Southern Ocean. *Nature,* 432: 100–103.

Baker, A.C., Clarke, M.R., & Harris, M.J. (1973). The N.I.O. combination net (RMT 1+8) and

further developments of rectangular midwater trawls. *Journal of Marine Biology Association of*

the United Kingdom, 53(1): 167–184.

Batta-Lona, P.G., Maas, A.E., O'Neill, R.J., Wiebe, P.H., & Bucklin A. (2017). Transcriptomic

profiles of spring and summer populations of the Southern Ocean salp, *Salpa thompsoni*, in the

Western Antarctic Peninsula region. *Polar Biology,* 40: 1261–1276.

Cai, W., Borlace, S., Lengaigne, M., Rensch, P., Collins, M., Vecchi, G., Timmermann, A.,

Santoso, A., McPhaden, M.J., Wu, L., England, M.H., Wang, G., Guilyardi, E., & Jin, F.-F.

(2014). Increasing frequency of extreme El Niño events due to greenhouse Warming. *Nature*

Climate Change, 4: 111–116.

Cavalieri, D.J., Parkinson, C.L., Gloersen, P., Comiso, J.C., & Zwally, H.J. (1999). Deriving

long-term time series of sea ice cover from satellite passive-microwave multi-sensor data sets.

Journal of Geophysical Research, 104: 15803–15814.

Chiba, S., Ishimaru, T., Hosie, G., & Wrigh, S. (1999). Population structure change of *Salpa*

thompsoni from austral mid–summer to autumn. *Polar Biology,* 22: 341–349.

Dana, J.D. (1850). Synopsis generum crustaceorum ordinis "Schizopoda. *American Journal of*

Science and Arts, 9: 129–133.Ducklow, H., Clarke, A., Dickhut, R., Doney, S.C., Geisz, H.,

Huang, K., Martinson, D.G., Meredith, M.P., Moeller, H.V., Montes-Hugo, M., Schofield, O.,

Stammerjohn, S.E., Steinberg, D., & Fraser, W. (2012). In Rogers, A., Johnston, N., Clarke, A. ,

& Murphy, E. (Eds.), *The marine ecosystem of the West Antarctic Peninsula* (pp. 121–159),

Oxford, UK: Blackwell Scientific..

Fetterer, F., Knowles, K., Meier, W.N., Savoie, M., & Windnagel, A.K. (2017). *Sea Ice Index,*

Version 3. [Th eAntarctic subset]. Boulder, Colorado USA. NSIDC: National Snow and Ice Data

Center. [20 August 2015] https://nsidc.org/data/G02135/versions/3.

Foxton, P. (1961). *Salpa fusiformis* Cuvier and related species. *Discovery Reports,* 32: 3–32.

Foxton, P. (1966). The Distribution and Life-History of *Salpa thompsoni* Foxton with

Observations on a Related Species, *Salpa gerlachei* Foxton. J Grad Res. Cent. *Discovery*

Reports, 34: 1–16.

Getis, A., & Ord, J.K. (1992). The analysis of spatial association by use of distance statistics.

Geographical Analysis, *24*, 189–206.

Goodall-Copestake, WP. (2016). One tunic but more than one barcode: evolutionary insights

from dynamic mitochondrial DNA in *Salpa thompsoni* (Tunicata: Salpida). *Biological Journal of*

the Linnean Society, 120: 637–648.

Henschke, N., Anthony, J.E., Richardson, Suthers J. 2016. Rethinking the Role of Salps in the

Ocean. *Trends in Ecology & Evolution* 31:9: 720-733.

Jue, N.K., Batta-Lona, P.G., Trusiak, S., Obergfell, C., Bucklin, A., & O'Neill, M. (2016). Rapid

evolutionary rates and unique genomic signatures discovered in the first reference genome for

the Southern Ocean salp, *Salpa thompsoni* (Urochordata, Thaliacea). *Genome Biology and*

Evolution, 8: 3171–3186.

Klinck, J.M., Hofmann, E.E., Beardsley, R.C., Salihoglu, B., & Howard, S. (2004). Water-mass

properties and circulation on the west Antarctic peninsula continental shelf in austral fall and

winter 2001. *Deep- Sea Research II,* 51: 1925–1946.

Kokubun, N., Kim, J.H., & Takahashi, A. (2013). Proximity of krill and salps in an Antarctic

coastal ecosystem: evidence from penguin-mounted cameras. *Polar Biology*, *36*(12): 1857–1864.

Lee, Ch., Pakhomov, E., Atkinson, A., & Siegel, V.. (2010). Long-term relationships between

the marine environment, krill, and salps in the Southern Ocean. *Journal of Marine Biology*, DOI:

10.1155/2010/410129.

Levitus, S., Antonov, J.I., Boyer, T.P., & Stephens, C. (2000). Warming of the world

ocean. *Science,* 287(5461): 2225–2229.

Levitus, S., Antonov, J., & Boyer, T. (2005). Warming of the world ocean, 1955–

2003. *Geophysical Research Letters* 39(L10603), DOI: 10.1029/2012GL051106.

Loeb, V., Siegel, V., Holm-Hansen, O., Hewitt, R., Fraser, W., Trivelpiece, W., & Trivelpiece,

S. (1997). Effects of sea-ice extent and krill or salp dominance on the Antarctic food

web. *Nature,* 387: 897–900.

Loeb, V.J, & Santora, J.A. (2012). Population dynamics of *Salpa thompsoni* near the Antarctic

Peninsula: growth rates and interannual variations in reproductive activity (1993–2009).

Progress in Oceanography, 96: 93–97.

Mackey, A.P., Atkinson, A., Hill, S.L., Ward, P., Cunningham, N.J., Johnston, N.M., & Murphy,

E.J. (2011). Antarctic macrozooplankton of the southwest Atlantic sector and Bellingshausen

Sea: Baseline historical distributions (Discovery Investigations, 1928–1935) related to

temperature and food, with projections for subsequent ocean warming. *Deep-Sea Research II,*

59: 130–146.

Martinson, D.G., Stammerjohn, S.E., Iannuzzi, R.A., Smith, R.C., & Vernetd, M. (2008).

Western Antarctic Peninsula physical oceanography and spatio–temporal variability. *Deep-Sea*

Research II Topical Studies in Oceanography, 55: 1964–1987.

Massom, R., & Stammerjohn, S.E. (2010). Antarctic sea ice change and variability – Physical

and ecological implications. *Polar Science,* 4: 149–186.

Meredith, M.P., & King, J.C. (2005). Rapid climate change in the ocean west of the Antarctic

Peninsula during the second half of the 20th century. *Geophysical Research Letters,* 32(L19):

604–609.

Moran, P. (1950). A Test for the Serial Independence of Residuals. *Biometrika,* 37: 178–181.

Nicol, S., Croxall, J., Trathan, P., Gales, N., & Murphy, E. (2007). Paradigm misplaced?

Antarctic marine ecosystems are affected by climate change as well as biological processes and

harvesting*, Antarctic Science,* 19(3): 291–295.

Nicol, S., Foster, J., & Kawaguchi, S. (2011). The fishery for Antarctic krill – recent

developments. *Fish and Fisheries,* DOI: 10.1111/j.1467-2979.2011.00406.x.

Ono, A., & Moteki, M. (2013). Spatial distributions and population dynamics of two salp

species, *Ihlea racovitzai* and *Salpa thompsoni,* in the waters north of Lu☐tzow-Holm Bay (East

Antarctica) during austral summers of 2005 and 2006. *Polar Biology,* 36: 807–907.

Pakhomov, E.A., Froneman, P.W., & Perissinotto, R. (2002). Salp/krill interactions in the

Southern Ocean: spatial segregation and implications for the carbon flux. *Deep−Sea Research II*

Tropical Studies in Oceanography, 49(9-10): 1881–1907.

Pommeranz, T., Hermann, C., & Kühn, A. (1982). Mouth angle of the Rectangular Midwater

Trawl (RMT1+8) during paying out and hauling. *Meeresforschung,* 29: 267–274.Purcell, J.E.,

Sturdevant, M.V., & Galt, C.P. (2005). A review of appendicularians as prey of fish and

invertebrate predators. In Gorsky, G., Youngbluth, M.J. & Deibel, D. (Eds.), *Response of marine*

ecosystems to global change: Ecological impact of appendicularians (pp.359-435), Paris,

France: Contemporary Publishing International.

Rayner, N.A., Parker, D.E., Horton, E.B., Folland, C.K., Alexander, L.V., Rowell, D.P., Kent,

E.C., & Kaplan, A. (2003). Global analyses of sea surface temperature, sea ice, and night marine

air temperature since the late nineteenth century. *Journal of Geophysical Research* 108(D14),

DOI: 10.1029/2002JD002670.

Richardson, P.L. (2008). On the history of meridional overturning circulation schematic

diagrams. *Progress in Oceanography,* 76: 466–486.

Ross, R.M., Questin, L.B., Newberger, T., Shaw, C.T., Jones, J.L., Oakes, S.A., Moore, K.J.

(2014). Trends, cycles, interannual variability for three pelagic species west of the Antarctic

Peninsula 1993-2008. *Marine Ecology Progress Series,* 515: 11–32.

Siegel, V., Skibowski, A., & Harm, U. (1992). Community structure of the epipelagic

zooplankton community under the sea-ice of the northern Weddell Sea. *Polar Biology,* 12(1):

15–24.

Siegel, V., & Loeb, V. (1995). Recruitment of Antarctic krill *Euphausia superba* and possible

causes for its variability. *Marine Ecology Progress Series,* 123: 45–56.

Słomska, A.W., Panasiuk–Chodnicka, A.A., Żmijewska, M.I., & Mańko, M.K. (2010).

Variability of *Salpa thompsoni* population structure in the Drake Passage in summer season

2010. *Polish Polar Research,* 36(4): 391–404.

Steinberg, D.K., Ruck, K.E., Gleiber, M.R., Garzio, L.M., Cope, J.C., Bernard, K.S.,

Stammerjohn, S.E., Schofield, O.M.E., Quetin, L.B., & Ross, R.M. (2015). Long-term (1993-

2013) changes in macrozooplankton of the Western Antarctic Peninsula. *Deep-Sea Research I,*

101: 54–70.

Thiebot, J.B., Ito, K., Raclot, T., Poupart, T., Kato, A., Ropert-Coudert, Y., & Takahashi, A.

(2016). On the significance of Antarctic jellyfish as food for Adélie penguins, as revealed by

video loggers. *Marine Biology 163-108*, DOI: 10.1007/s00227-016-2890-2.

Tokarczyk, R. (1987). Classification of water masses in the Bransfield Strait and southern part of

the Drake Passage using a method of statistical multidimensional analysis. *Polish Polar*

Research, 8(4): 333–366.

Turner, J., Lu, H., White, I., King, J.C., Phillips, T., Hosking, J.S., Bracegirdle, T.J., Marshall,

G.J., Mulvaney, R., & Deb, P. (2016). Absence of 21st century warming on Antarctic Peninsula

consistent with natural variability. *Nature,* 535 (7612): 411–415.

Vaughan, D.G., Comiso, J.C., Allison, I., Carrasco, J., Kaser, G., Kwok, R., Mote, P., Murray,

T., Paul, F., Ren, J., Rignot, E., Solomina, O., Steffen, K., & Zhang, T. (2013) Observations:

Cryosphere. In: *Climate Change* 2013. In Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M.,

Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., & Midgley, P.M. (Eds.), *The Physical*

Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the

Intergovernmental Panel on Climate Change Cambridge, UK: Cambridge University Press.

Whitehouse, M.J., Meredith, M.P., Rothery, P., Atkinson, A., Ward, P., & Korb, R.E. (2008).

Rapid warming of the ocean around South Georgia, Southern Ocean, during the 20th century:

forcings, characteristics, and implications for lower trophic levels. *Deep-Sea Research I,* 55:

1218–28.

Zwally, H.J., Comiso, L.D., Parkinson, C.L., Cavalieri, J.P., & Gloersen, P. (2000). Variability

of Antarctic sea ice 1979–1998. *Journal of* Geophys*ical Research,* 107(C5):

10.1029/2000JC000733.

FIGURES LEGENDS

Fig. 1. The locations of the sampling stations with the year of the cruise depicted in different

color and shape.

Fig. 2. Mean abundance of *Salpa thompsoni* sampled each year (standard deviation values are

given above the line).

Fig. 3. Generalized Additive Models (GAM) fitted for the number of salps against: (A) sea

surface temperature (N= 461), (B) salinity (N=212), and (C) bottom depth (N=1278).

Fig 4. Hot spots of *Salpa thompsoni* distribution in decadal time series: (A) '70s, (B) '80s and

(C) '90s (in the background - the interpolated ice concentration for the respective time series.

Cold spotsmarked: in grey circles - 90% confidence, blue squares - 95% confidence, dark blue

triangles - 99 % confidence; hot spots marked: in orange triangles - 90% confidence, dark orange

sqares - 95% confidence, red circles – 99% confidence; different sea ice degrees marked in:

violet – without sea ice, dark blue- 1, sky blue- 2, light blue- 3).

Fig. 5. Boundary of *Salpa thompsoni* temperature optimum: in the studied year (A - marked in

light gray) and prognosis on its change after 1°C temperature rise (B - marked in dark gray).

▲	1975 - 1976
●	1977 - 1978
■	1979
▲	1980 - 1981
●	1982
■	1983 - 1984
▲	1985
●	1986
■	1987 - 1988
▲	1989 - 1990
●	1994 - 1995
■	1996 - 1997
⊙	1998 - 2001