The most abundant cyst wall proteins of *Acanthamoeba castellanii* are three sets of lectins that

bind cellulose and chitin and localize to distinct structures in cyst walls

Running title: Identification and characterization of *Acanthamoeba* cyst wall proteins

Pamela Magistrado-Coxen[1,#a], Yousuf Aqeel[1], Angelo Lopez[1,2], John R. Haserick[1,2,#b],

Breeanna R. Urbanowicz[3], Catherine E. Costello[2], and John Samuelson[1*]

[1] Department of Molecular and Cell Biology, Boston University Goldman School of Dental

Medicine, Boston, Massachusetts, United States of America

[2] Department of Biochemistry, Boston University School of Medicine, Boston, Massachusetts,

United States of America

[3] Complex Carbohydrate Research Center, University of Georgia, Athens, Georgia, United

States of America

[#a] Current Address: Sarepta Therapeutics, Andover, Massachusetts, United States of America

[#b] Current Address: Glyde Bio, Inc., Cambridge, Massachusetts, United States of America

* Corresponding author

E-mail: jsamuels@bu.edu

## ABSTRACT

*Acanthamoeba castellanii*, cause of keratitis and blindness, is an emerging pathogen because of its association with contact lens use.  The cyst wall contributes to pathogenesis as cysts are resistant to sterilizing reagents in lens solutions and to antibiotics applied to the eye.  We used transmission electron microscopy, as well as structured illumination microscopy and probes for cellulose and chitin, to show that purified cyst walls of *A. castellanii* retain an outer ectocyst layer, an inner endocyst layer, and conical ostioles that connect the layers.  Mass spectrometry showed candidate cyst wall proteins are dominated by three families of lectins (named here Luke, Leo, and Jonah), each of which binds to microcrystalline cellulose and to a lesser degree chitin.  A Jonah lectin, which has one choice-of-anchor A (CAA) domain, localizes to the ectocyst layer of mature cyst walls. Luke lectins, which have two or three carbohydrate-binding modules (CBM49), localize to the endocyst layer and ostioles.  A Leo lectin, which has two domains with eight Cys residues each (8-Cys), also localizes to the endocyst layer and ostioles.  In summary, the most abundant *A. castellanii* cyst wall proteins are three sets of lectins, which have carbohydrate-binding modules that are conserved (CBM49s of Luke), newly characterized (CAA of Jonah), or unique to *Acanthamoebae* (8-Cys of Leo).  Despite their lack of common ancestry, Luke and Leo lectins both localize to the endocyst layer and ostioles, while the Jonah lectin localizes to the ectocyst layer.

## IMPORTANCE

Fifty years ago, investigators identified cellulose in the *Acanthamoeba* cyst wall, which has an outer ectocyst layer, an inner endocyst layer, and conical ostioles that connect the layers.  Here we show cyst walls also contain chitin and three large sets of cellulose- and chitin-binding

lectins, which have distinct localizations.  The *Acanthamoeba* cyst wall, therefore, is more

complicated than cyst walls of *Entamoeba* and *Giardia* (causes of dysentery and diarrhea,

respectively), which have a single layer, a single glycopolymer, and small sets of one or two

lectins.  In contrast, the *Acanthamoeba* cyst wall is far simpler than the walls of fungi and

plants, which have multiple layers, numerous glycopolymers, and hundreds of proteins.  In

addition to providing a better understanding of the cell biology and biochemistry of the

*Acanthamoeba* cyst wall, these studies may lead to diagnostic antibodies that bind to cyst wall

proteins and/or therapeutics that target chitin.

**Introduction**

*Acanthamoebae*, which include the genome project *A. castellanii* Neff strain, are soil protists

named for acanthopods (spikes) on the surface of trophozoites (1).  In immunocompetent

persons, *Acanthamoebae* is a rare but important cause of corneal inflammation (keratitis),

which is difficult to treat and so may lead to scarring and blindness (2-4).  In

immunosuppressed patients, *Acanthamoebae* may cause encephalitis (5).  *Acanthamoeba* is

an emerging pathogen, because 80 to 90% of infections are associated with contact lens use,

which is growing worldwide (6, 7).  Because water for washing hands may be scarce in places

where the free-living protist is frequent, we recently showed that alcohols in concentrations

present in hand sanitizers kill *A. castellanii* trophozoites and cysts (8, 9).

When *A. castellanii* trophozoites are deprived of nutrients in solution or on agar plates,

they form cysts (10-12).  Transmission electron microscopy (TEM) shows cyst walls have two

microfibril-dense layers (outer ectocyst and inner endocyst), which are separated by a

relatively microfibril-free layer (13).  The endocyst and ectocyst layers are connected to each

other by conical ostioles, through which the protist escapes during excystation (14).

The cyst wall of *A. castellanii* protects free-living protists from osmotic shock when

exposed to fresh water, drying when exposed to air, or starvation when deprived of bacteria or

other food sources.  The cyst wall also acts as a barrier, sheltering parasites from killing by

disinfectants used to clean surfaces, sterilizing agents in contact lens solutions, and/or

antibiotics applied directly to the eye (15-17).

We are interested in the cyst wall proteins (CWPs) of *A. castellanii* for three reasons.

First, although monoclonal antibodies to *A. castellanii* have been made, the majority react to

trophozoites, and no CWPs have been molecularly identified (18-21).  Indeed the only cyst-

specific protein identified, which was named for its 21-kDa predicted size (CSP21), is unlikely

to be a CWP, as it lacks a signal peptide (22).  Second, *A. castellanii* and related species are

the only human pathogens that contain cellulose in their wall (23-25).  *Dictyostelium*

*discoideum*, which also has cellulose in its walls, is not pathogenic (26).  Third, the whole

genome of *A. castellanii* predicts a chitin synthase, a chitinase, and two chitin deacetylases,

suggesting the possibility that chitin and chitin-binding proteins are also present in the cyst wall

(1, 27-29).

Our experimental design was simple.  We used transmission electron microscopy

(TEM), as well as structured illumination microscopy (SIM) and probes for glycopolymers, to

judge the intactness and cleanliness of purified *A. castellanii* cyst walls (30).  We used mass

spectrometry to identify candidate CWPs, which were compared to proteins present in walls of

other protists, bacteria, fungi, and plants (31).  We used SIM to localize representative CWPs,

each of which was tagged with a green fluorescent protein (GFP) and expressed under its own

promoter in cyst walls of stably transfected protists (32, 33).  We also determined whether

each CWP, which was made as a maltose-binding protein (MBP) in the periplasm of bacteria,

binds to microcrystalline cellulose and/or chitin beads (34).

## RESULTS

**TEM and SIM showed purified _A. castellanii_ cyst walls contained distinct endocyst and**

**ectocyst layers, as well as ostioles.**  Cyst wall preparations were made by sonicating cysts

and separating walls from cellular contents by density centrifugation and retention on a

membrane with 8 μm-diameter pores.  For TEM, intact cysts and purified cyst walls were

frozen under high pressure, and fixatives were infiltrated at low temperature (35).  Purified cyst

walls had intact ectocyst and endocyst layers, as well as conical ostioles that link them (Fig. 1)

(13).  The purified walls were missing amorphous material that fills the space between the

inner aspect of the cyst wall and the plasma membrane of the trophozoite inside.

For SIM, we used probes that bind chitin (wheat germ agglutinin, WGA) and β-1,3 and

β-1,4 polysaccharides (calcofluor white, CFW) in the walls of fungi and cysts of _Entamoeba_

(36-39).  CFW, a fluorescent brightener, has also been used to diagnose _Acanthamoeba_ cysts

in eye infections (40).  In addition, we made a glutathione-S-transferase (GST) fusion-protein,

which contains the N-terminal CBM49 of a candidate CWP of _A. castellanii_ (Fig. S1 and Excel

file S1) (41).  The GST-AcCBM49 expression construct was designed to replicate that used to

determine the carbohydrate binding properties of SlCBM49, which is a C-terminal

carbohydrate-binding module of the _Solanum lycopersicum_ (tomato) cellulase SlGH9C (42).  In

both intact cysts and purified cyst walls, GST-AcCBM49 predominantly labeled the ectocyst

layer, WGA highlighted the ostioles, and CFW stained the endocyst layer (Fig. 2).  In

summary, TEM and SIM both showed that ectocyst and endocyst layers, as well as ostioles,

were intact in purified cyst walls, which were relatively free of cellular material.

**Mass spectrometry showed candidate CWPs of *A. castellanii* were encoded by**

**multigene families and contain tandem repeats of short domains (CBM49, 8-Cys, and**

**CAA).**  Trypsin treatment of purified *A. castellanii* cyst walls, which was followed by liquid

chromatography mass spectrometry (LC-MS/MS) analysis of the released peptides, gave

similar results in five biological experiments (Table 1 and Excel file S1).  While some proteins

remained in cyst walls after trypsin digestion, their identities were similar to those detected in

the soluble fractions by in gel-digests with trypsin or chymotrypsin.  Candidate CWPs with the

most unique peptides identified by LC-MS/MS belonged to three families, which we named

Luke, Leo, and Jonah lectins, because each bound to cellulose +/- chitin (see below).

Although it was impossible to draw a line that separates actual CWPs from contaminating

proteins, secreted proteins with 18+ unique peptides included six Leo lectins, four Luke lectins,

and three Jonah lectins.  The vast majority of proteins with <18 unique peptides were predicted

to be cytosolic (including CSP21) and so were likely intracellular contaminants of cyst wall

preparations.  The exception to this hypothesis, we think, were additional Luke, Leo, and

Jonah lectins, which are most likely less abundant CWPs.  For readers interested in cytosolic

proteins of *A. castellanii*, we have added Excel file S2, which contains all the mass

spectrometry data including a "dirty" cyst wall preparation that was generated without using the

Percoll gradient or porous filter.

Luke lectins were comprised of an N-terminal signal peptide, followed by two or three

CBM49s that were separated by Ser- and Pro-rich spacers (Figs. 3 and S2) (27, 42-45).  The

N-terminal CBM49 of Luke lectins contained three conserved Trp resides conserved in

SlCBM49 from tomato.  A fourth conserved Trp is present in the CBM49 of *D. discoideum*

cellulose-binding proteins (46).  The other CBM49s (middle and/or C-terminal) of Luke lectins

had two conserved Trp residues.  Luke lectins were acidic (pI 5 to 6) and had formula weights

(FWs) from 27 to 29-kDa (two CBM49s) or 42 to 44-kDa (three CBM49s).  There were no

predicted transmembrane helices or glycosylphosphatidylinositol anchors in the Luke or Leo

lectins (47, 48).  LC-MS/MS of the released cell wall peptides identified at least one unique

peptide corresponding to all 12 genes encoding Luke lectins, although the number of unique

peptides varied from 78 to two (Table 1 and Excel file S1).  In general, Luke lectins with two

CBM49s had more unique peptides than Luke lectins with three CBM49s.  One to four unique

peptides were derived from three CBM49-metalloprotease fusion-proteins, which consisted of

an N-terminal signal peptide followed by a single CBM49 with four conserved Trp residues and

a metalloprotease (ADAM/reprolysin subtype) with a conserved catalytic domain

(HEIGHNLGGNH) (43).  We used a Luke(2) lectin (ACA1_377670) with two CBM49s to

perform RT-PCR, make rabbit anti-peptide antibodies, and make maltose-binding protein

(MBP)- and green fluorescent protein (GFP)-fusions (Figs. 3 and S1) (33, 34).  We also used a

Luke(3) lectin (ACA1_245650) with three CBM49s to make a GFP-fusion (Fig. S2).

Leo lectins had an N-terminal signal peptide followed by two repeats of a unique 8-Cys

domain, some of which were separated by a long Thr-, Lys-, and His-rich spacer (Figs. 3 and

S2).  Leo lectins without a spacer were acidic (pI ~4.8) and had FWs from 19 to 24-kDa, while

Leo lectins with the TKH-rich spacer were basic (pI ~8.3) and had FWs from 36- to 59-kDa.

Leo lectins were encoded by 16 genes, of which 14 proteins were identified by our LC-MS/MS

analysis.  While the number of unique peptides varied from 34 to one, Leo lectins without the

spacer generally had more unique peptides than Leo lectins with the TKH-rich spacer.  We

used a Leo lectin with no spacer (ACA1_074730) to perform RT-PCR, make rabbit anti-peptide

antibodies, and make MBP- and GFP-fusions (Fig. S1).

Jonah lectins had an N-terminal signal peptide followed by one or three choice-of-

anchor A (CAA) domains (Figs 3. and S2) (43).  The binding activity of the CAA domain, which

is adjacent to a collagen-binding domain in a microbial surface component recognizing the

adhesive matrix molecule (MSRAMM) of *Bacillus anthracis*, was not known (49).  Jonah lectins

with a single CAA domain were acidic (pI ~6), had a FW from 44 to 58-kDa and had an N-

terminal Thr-, Lys-, and Cys-rich domain.  Jonah lectins with three CAA domains were basic

(pI ~8.8), had a FW of ~146-kDa, and contained Ser- and Pro-rich spacers between CAA

domains, as well as and hydrophobic regions that may be transmembrane helices (48).  Jonah

lectins were encoded by eight genes, of which five were identified by our LC-MS/MS analysis

based on one to 147 unique peptides.  We used a Jonah lectin (ACA1_164810) with a single

CAA domain to perform RT-PCR, make rabbit anti-peptide antibodies, and make MBP- and

GFP-fusions (Fig. S1).

Other secreted proteins with 18+ unique peptides detected by LC-MS/MS, which are

candidate CWPs, included a laccase with three copper oxidase domains (ACA1_068450), a

protein with a C-terminal ferritin-like domain (ACA1_292810), a Kazal-type serine protease

inhibitor (ACA1_291590), a conserved uncharacterized protein (ACA1_068630), and a protein

unique to *A. castellanii* (ACA1_145900) (43, 44, 50).  Interestingly, a bacterial laccase has

been shown to bind cellulose (51).  There were also three serine proteases, which have been

localized to the secretory system of encysting organisms (52).

These results suggested that the most abundant candidate CWPs of *A. castellanii*

contain tandem repeats of conserved domains (CBM49 in Luke lectins and CAA in Jonah

lectins) or a unique domain (8-Cys in Leo lectins).  Peptides corresponding to nearly all

members of each gene family were detected by mass spectrometry, although the relative

abundances of unique peptides for each CWP varied by more than an order of magnitude,

suggesting marked differences in gene expression.  Because it was not possible to separate

cyst walls into component parts (endocyst and ectocyst layers and ostioles) prior to LC-MS/MS

analysis of tryptic peptides, we used SIM and GFP-tags to localize representative members of

each family of CWPs in cyst walls of transfected *A. castellanii* (see below).

**Origins and diversity of genes that encode Luke, Leo, and Jonah lectins.**  Leo

lectins, which had two domains with 8-Cys each, appeared to be unique to *A. castellanii*, as no

homologs were identified when BLAST analysis were performed using the nonredundant (NR)

database at NCBI (https://www.ncbi.nlm.nih.gov/) (28).  The origin of genes encoding Luke

lectins was difficult to infer, because its CBM49s showed only a 31% identity over a short (77-

amino acid) overlap with a predicted cellulose-binding protein of *D. discoideum* (expect value

of BLASTP was just 7e-05) (28, 46).  In contrast, the CAA domain of Jonah lectins appeared to

derive from bacteria by horizontal gene transfer (HGT), as no other eukaryote contained CAA

domains, and there was a 28% identity over a bigger (263-aa) overlap with a choice-of anchor

A family protein of *Saccharibacillus sp. O16* (5e-12) (1, 28).  The *A. castellanii* laccase (also

known as copper oxidase), whose signals were abundant in the mass spectra, was likely the

product of HGT from bacteria, as there was a 44% identity over a large (526-aa) overlap with a

copper oxidase of *Caldicobacteri oshimai* (6e-135) (50).  The uncertainty was based upon the

presence of similar enzymes in plants, one of which (*Ziziphus jujube*) showed a 39% identity

over a 484-aa overlap (4e-101) with the *A. castellanii* laccase.

No pairs of genes within each lectin family were syntenic, suggesting that individual

genes rather than clusters of genes were duplicated.  With the exception of two Luke lectins

(ACA1_253500 and ACA1_253650) that were 98% identical and two Leo lectins

(ACA1_074770 and ACA1_083920) that were 85% identical, members of each family of CWPs

differed in amino acid sequence by >40%.  Genes that encode CWPs also varied in the

number of introns (zero to two in Luke, two to four in Leo, and zero to 24 in Jonah).  Searches

of genomic sequences of 11 strains of *Acanthamoebae*, deposited in AmoebaDB without

protein predictions by Andrew Jackson of the University of Liverpool, using TBLASTN and

sequences of representative Luke, Leo, and Jonah lectins localized in the next section,

showed four results (28, 29).  First, although stop codons were difficult to identify using this

method, all 11 strains appeared to encode each CWP.  Second, most strains showed 100 to

200-amino acid stretches of each CWP that were 80 to 90% identical to the *A. castellanii* Neff

strain studied here.  These stretches did not include low complexity spacers, which are difficult

to align.  Third, some of the strains showed greater differences from the Neff strain in each

CWP, consistent with previous descriptions of *Acanthamoeba* strain diversity based upon 18S

rDNA sequences (53).  Fourth, while coding sequences and 5' UTRs were well conserved,

introns were very poorly conserved, with the exception of branch-point sequences.

In summary, genes encoding Jonah lectin and laccase likely derived by HGT, while

genes encoding Leo lectins appeared to originate within *Acanthamoeba*.  Although CBM49s of

Luke lectins shared common ancestry with plants and other Amoebazoa, their precise origin

was not clear.  For the most part, gene duplications that expanded each family within the

*Acanthamoeba* genome occurred a long time ago, as shown by big differences in amino acid

sequences of paralogous proteins and variations in the number of introns.  Still, the set of

Luke, Leo, and Jonah lectins identified by mass spectrometry and the sequence of

representative CWPs localized in the next section both appeared to be conserved among 11

sequenced isolates of *Acanthamoebae*.

**SIM showed a representative Jonah lectin was present in the ectocyst layer of**

**mature cyst walls, while representative Luke and Leo lectins were present in the**

**endocyst layer and ostioles.**  To localize candidate CWPs, we expressed a Leo lectin with

no spacer and a Jonah(1) lectin with a single CAA domain, each with a GFP-tag under its own

promoter (446- and 571-bp of the 5' UTR, respectively) in transfected trophozoites of *A.*

*castellanii*, using an episomal vector that was selected with G418 (Fig. S1) (32, 33).  We also

expressed a Luke(2) lectin with two CBM49s and a Luke(3) lectin with three CBM49s, each

with a GFP-tag under its own promoter (486- and 500-bp of the 5' UTR, respectively).  GFP-

tagged candidate CWPs expressed under their own promoter were absent in the vast majority

of log-phase trophozoites, which divided but did not become cysts (data not shown).  GFP-

tagged CWPs were present in small numbers in trophozoites in stationary cultures, where a

few organisms have begun to encyst spontaneously.  In contrast, the vast majority of mature

cysts contained GFP-tagged CWPs in their walls.

Jonah(1)-GFP was present in the ectocyst layer of mature cyst walls, which were

labeled with WGA and CFW (Fig. 4A).  In contrast, Leo-GFP, Luke(2)-GFP, and Luke(3)-GFP

were each present in the endocyst layer and sharply outlined the ostiole (Figs. 4B to 4D).

These results showed that ~500 bp of the 5' UTR was sufficient to cause encystation-specific

expression of GFP-tagged representatives of Luke, Leo, and Jonah lectins.  While the Jonah

lectin localized to the ectocyst layer, Luke and Leo lectins, which do not share common

ancestry, both localized to the endocyst layer and ostioles.  Finally, Luke lectins with two or

three CBM49s localized to the same place.

We think the timing of expression and locations of the GFP-tagged CWPs in cyst walls

were accurate for the following reasons.  First, RT-PCR showed that mRNAs of representative

Luke, Leo, and Jonah lectins, as well as cellulose synthase (ACA1_349650), were absent or

nearly absent from trophozoites but were present during the first three days of encystation

(Fig. S3).  In contrast, glyceraldehyde 3-phosphate dehydrogenase (GAPDH), which catalyzes

the sixth step in glycolysis, was expressed by both trophozoites and encysting *A. castellanii*

(33).  Second, monospecific, polyclonal rabbit antibodies to a 50-amino acid peptide of a

representative Jonah lectin and a 16-amino acid peptide of a representative Leo lectin bound

to Western blots of proteins from cysts but not from trophozoites (Fig. S4).  We were unable to

generate rabbit antibodies to the Luke lectin, using methods that worked to make antibodies to

Jonah and Leo lectins. None of the rabbit anti-peptide antibodies was useful for localizing

CWPs by SIM.  Third, GFP-tagged CSP21 expressed under its own promoter was present in

cytosolic accumulations of mature cysts (Fig. S5) (22, 54).  As CSP21 is homologous to

universal stress proteins and lacks an N-terminal signal peptide, its presence in the cytosol

after nutrient deprivation was expected (43, 55).  Fourth, a GFP-fusion protein (SP-GFP),

which was appended with an N-terminal signal peptide from Luke lectin and expressed under a

GAPDH promoter, localized to secretory vesicles of cysts but not to cyst walls (Fig. S5) (33).

**Luke, Leo, and Jonah lectins all bound to microcrystalline cellulose, while**

**binding of CWPs to chitin beads was variable.**  To test the binding of representative CWPs

to commercially available glycopolymers, we made MBP-CWP fusion-proteins in the periplasm

(secretory compartment) of *E. coli* (34).  The targets were microcrystalline cellulose (used to

characterize binding activities of GST-SlCBM49 from tomato cellulase) and chitin beads (used

to characterize myc-tagged Jacob and Jessie lectins of *Entamoeba histolytica*) (42, 56).

Western blots anti-MBP antibodies showed MBP-Luke(2), which was partially degraded, bound

to microcrystalline cellulose and somewhat less well to chitin beads (Fig. 5).  MBP-Jonah(1)

also bound to microcrystalline cellulose and chitin beads.  MBP-Leo bound less completely to

microcrystalline cellulose and weakly at best to chitin beads.  MBP alone (negative control) did

not bind to microcrystalline cellulose or chitin beads.  These results suggested each CWP is

binding to glycopolymers in the cyst wall, but they did not explain why Jonah localized to the

ectocyst layer, while Luke and Leo localized to the endocyst layer and ostioles.

**Discussion**

**Advantages of studying *Acanthamoeba* cyst walls.**  *A. castellanii* has a number of

properties that make it possible to quickly identify and begin to characterize its CWPs.  The

protist grows axenically in medium without serum or vitamins and synchronously encysts when

placed on non-nutrient agar plates (10, 11).  Cyst walls separate cleanly from cellular contents

without losing essential structures (ectocyst and endocyst layers and ostioles), which we show

are readily visualized with SIM using probes for wall glycopolymers (GST-AcCBM49, WGA,

and CFW) (13, 30, 36-39).  The most abundant CWPs belong to three protein families that

contain tandem repeats of CBM49s (Luke lectins), 8-Cys domains (Leo lectins), or CAA

domains (Jonah lectins).  Stable expression of GFP-tagged proteins under their own promoters

allows localization of candidate CWPs during cyst wall development using SIM (32, 33).

Efficient expression of recombinant proteins in the periplasm (MBP-fusions) of bacteria makes

it possible to assay the ability of CWPs to bind microcrystalline cellulose or chitin beads, by

using techniques to characterize other protist cyst wall lectins and carbohydrate-binding

modules (34, 42, 56-58).

The ease of studying cyst walls of *A. castellanii* is in contrast to studies of cyst walls of

*E. histolytica*, which do not encyst *in vitro* and so are modeled by cysts of the reptilian

pathogen *E. invadens* (38, 56).  While *Giardia* cysts made in mice excyst well, cysts made *in*

*vitro* do not (57).  *Toxoplasma* forms walled oocysts in cats, while *Cryptosporidium* forms

oocysts in cows, mice, or gnotobiotic pigs (58-60).  Oomycetes (water molds that cause potato

blight) also have cellulose and chitin in their walls, but the amount of chitin is very small, and β-

1,3-glucan is also present (61).  The spore coat of *Dictyostelium discoideum*, a cousin of *A.*

*castellanii*, contains cellulose and a heteropolysaccharide of galactose and GalNAc, but it

lacks chitin (26, 62).

Properties of *A. castellanii* that make it challenging to study its cyst wall include the

following.  While cyst walls of *Entamoeba* and *Giardia* have a single layer, a single

glycopolymer (chitin and β-1,3-linked-GalNac, respectively), and small sets of one or two

lectins, *A. castellanii* cyst walls have two layers and ostioles, at least two glycopolymers, and

three large sets of lectins (31, 56, 57, 63, 64).  Although cellulose and chitin synthases are

each single copy, it will be difficult to knock out sets of genes encoding CWPs, even with the

efficient gene editing methods available in other protists (27-29, 65, 66).

**Familiarity and novelty in *A. castellanii* CWPs.**  Although we expected Luke lectins

with two or three CBM49s would be present in the cellulose-rich cyst wall, we could not have

predicted the other abundant CWPs, because the 8-Cys domains of Leo lectins are unique to

*Acanthamoebae* and the CAA domains of Jonah lectins were previously uncharacterized (27,

42, 49).  While Luke lectins have two or three CBM49s, *D. discoideum* has dozens of proteins

with a single CBM49 (Fig. S6) (44).  The Luke lectin binds cellulose and chitin, while the *D.*

*discoideum* proteins with a single CBM49 bind cellulose (46). Chitin-binding by DdCBM49 or

SlCBM49 was not tested, because this glycopolymer is not present in *D. discoideum* and

tomato walls.  Demonstration that CBM49s of the Luke lectin also bind chitin fibrils is new, but

is consistent with recent studies showing CBMs may bind more than one glycopolymer (67).

The metalloprotease fused to an N-terminal CBM49 of *A. castellanii* is absent in *D.*

*discoideum*, while *D. discoideum* adds two CBM49s to a cysteine proteinase, which lacks

these domains in *A. castellanii*.  The CBM49 may act to localize the metalloproteases to the *A.*

*castellanii* cyst wall, as is the case for the chitin-binding domain in *Entamoeba* Jessie lectins or

glucan-binding domain in *Toxoplasma* glucanases (56, 58).  Alternatively, the CBM49 may

suggest the metalloprotease cleaves glycopeptides rather than peptides.  While the GH5

glycoside hydrolases of *A. castellanii* lack CBM49s, CBM49 is present at the C-terminus of

GH9 glycoside hydrolases of *D. discoideum* and *S. lycopersicum* (27, 42, 44).

Even though *A. castellanii* Leo lectins and *E. histolytica* Jacob lectins share no common

ancestry, they have 8-Cys and 6-Cys lectin domains, respectively, often separated by low

complexity sequences (Fig. S7) (63, 64).  *E. histolytica* low complexity sequences vary from

strain to strain, contain cryptic sites for cysteine proteases, and are extensively decorated with

*O*-phosphate-linked glycans (68).  We have not yet identified any Asn-linked or *O*-linked

glycans on Leo lectins or any of the other CWPs, but we expect they will be there.  *A.*

*castellanii* and oomycetes (*Pyromyces* and *Neocallmistic*) each contain proteins with arrays of

CAA domains, but the sequences of the CAAs are so different that it is likely that

concatenation of domains occurred independently (Fig. S8) (43, 44).  Although *A. castellanii* is

exposed to collagen in the extracellular matrix of the cornea, the protist lacks a homolog of the

collagen-binding domain that is adjacent to the CAA domain in the *Bacillus anthracis* collagen-

binding protein (49).

Concatenation of carbohydrate-binding domains in Luke, Leo, and Jonah lectins, which

has previously been shown in WGA, Jacob lectins of *E. histolytica*, and peritrophins of insects,

most likely increases the avidity of the lectins for glycopolymers (31, 63, 64, 69).  The large

number of genes encoding Luke, Leo, and Jonah lectins may be necessary to increase the

quantity of CWPs coating glycopolymers in the cyst wall.  For example, Luke(2) and Luke(3)

lectins with two or three CBM49s, respectively, localize to the same place in mature cyst walls.

Alternatively, there may be differences in timing and localization of CWPs within the same

family, based upon a TKH-rich spacer in Leo lectins or transmembrane helices in Jonah lectins

with three CAA domains (Fig. S2).  Finally, other candidate CWPs, which are abundant but

present at lower copy numbers (e.g. laccase or ferritin-domain protein) and so not tested here,

may have important functions in the cyst wall.

**Possible translation of these results to the clinic.**  While the focus here was on the

biochemistry and cell biology of the cyst wall, Luke, Leo, and Jonah lectins, which are

conserved in all strains of *A. castellanii*, appear to be excellent targets for diagnostic

antibodies, as has been shown for *Giardia* CWP1 and *Entamoeba* Jacob2 lectin (70, 71).

Evidence for chitin in the cyst wall, admittedly not extensively developed here, suggests the

possibility that chitin synthase inhibitors or chitinases might be used as therapeutics (72).

Other investigators have explored the possibility of cellulose synthase inhibitors or cellulases

as therapeutics versus *A. castellanii* cysts (73-76).  The SIM methods, which worked so well

here to study *A. castellanii* cyst walls, might be used to study the fine structure and assembly

of other protist walls.  Insights from studies of the *A. castellanii* cyst wall, which is relatively

simple, may inform studies of fungal and plant walls, which are highly complex.

## MATERIALS AND METHODS

**Culture of trophozoites and preparation of encysting organisms and cysts.**  Culture and

manipulation of *A. castellanii* were approved by the Boston University Institutional Biosafety

Committee.  *A. castellanii* Neff strain trophozoites were purchased from the American Type

Culture collection.  Trophozoites of *A. castellanii* MEEI 0184 strain, which was derived from a

human corneal infection, were obtained from Dr. Noorjahan Panjwani of Tufts University

School of Medicine (9).  Trophozoites were grown in T-75 tissue culture flasks at 30°C in 10 ml

ATCC medium 712 (PYG plus additives) (Sigma-Aldrich Corporation, St. Louis, MO) (11).

Flasks containing log-phase trophozoites (free of cysts that form spontaneously in stationary

cultures) were either chilled or scraped with a cell scraper to release adherent amoebae, which

were concentrated by centrifugation at 500 x g for 5 min and washed twice with phosphate

buffered saline (PBS).  Approximately $10^7$ amoebae obtained from a confluent flask were

induced to encyst by incubation at 30°C on non-nutrient agar plates (10).  After 3 to 6 days

incubation, 15 ml of PBS was added to agar plates, which were incubated on a shaker for 30

min at room temperature (RT).  Encysting organisms were removed using a cell scraper and

concentrated by centrifugation at 1,500 x g for 10 min.

**Preparation of mature cyst walls for SIM, TEM, and mass spectrometry.**  Mature

cysts were washed in PBS and suspended in lysis buffer (10 mM HEPES, 25 mM KCl, 1 mM

$CaCl_2$, 10 mM $MgCl_2$, 2% CHAPS, and 1X Roche protease inhibitor) (Sigma-Aldrich). For SIM,

cysts in 500-µl lysis buffer were broken four times for 2 min each with 200 µl of 0.5 mm glass

beads in a Mini-Beadbeater-16 (BioSpec Products, Bartlesville, OK). For TEM, where glass

beads cannot be used, cysts in 200-µl lysis buffer were broken by sonication four times for 20

seconds each in continuous mode in a Sonicator Cell Disruptor (formerly Heat Systems

Ultrasonic, now Qsonica, Newtown, CT). Broken cysts were added to the top a 15-ml falcon

tube containing 60% sucrose and centrifuged at 4,000 x g for 10 min. The broken cyst wall

pellet was suspended in PBS buffer and washed three times at 10,000 x g in a

microcentrifuge. The cyst wall pellet was used directly for labeling for SIM or fixation for TEM.

For mass spectrometry, the cyst wall pellet broken in the bead beater was overlaid on gradient

containing 2 ml each of 20%, 40%, 60% and 80% Percoll (top to bottom) and centrifuged for

20 min at 3,000 x g. The layer between 60% and 80% Percoll, where the broken cyst walls

were located, was collected and washed in PBS. The cyst wall preparation was suspended in

10 ml of PBS, placed in a syringe, and forced through a 25-mm diameter Whatman Nuclepore

Track-Etched Membrane with 8-µm holes (Sigma-Aldrich). The cellular debris, which passed

through the membranes, was discarded. The membrane was removed from the cassette,

suspended in 5 ml of PBS, and vortexed to release cyst walls. The membrane was removed,

and cyst walls were distributed in microfuge tubes and pelleted at 15,000 x g for 10 min. The

pellet was suspended in 50 µl PBS and stored at -20°C prior to trypsin digestion and mass

spectrometry analysis.

**SIM of glycopolymers of mature cysts and purified cyst walls.** A GST-AcCBM49

fusion-construct, which contains the N-terminal CBM49 of a representative Luke lectin minus

the signal peptide, was prepared by codon optimization (76 to 330-bp coding region of

ACA1_377670) (GenScript, Piscataway, NJ). It was cloned into pGEX-6p-1 (GE Healthcare

Life Sciences, Marlborough, MA) for cytoplasmic expression in BL21(DE3) chemically

competent *E. coli* (Thermo Fisher Scientific, Waltham, MA) (41, 42).  Expression of GST-

AcCBM49 and GST were induced with 1 mM IPTG for 4 hr at RT, and GST-fusions were

purified on glutathione-agarose and conjugated to Alexa Fluor 594 succinimidyl esters (red)

(Molecular Probes, Thermo Fisher Scientific).  Approximately $10^6$ mature cysts or cyst walls

were washed in PBS and fixed in 1% paraformaldehyde buffered with phosphate for 15 min at

RT. Pellets were washed two times with Hank's Buffered Saline Solution (HBSS) and

incubated with HBSS containing 1% bovine serum albumin (BSA) for 1 hour at RT.

Preparations were then incubated for 2 hr at $4^0$C with 2.5 µg GST-CBM49 conjugated to Alexa

Fluor 594 and 12.5 µg of WGA (Vector Laboratories, Burlingame, CA) conjugated to Alexa

Fluor 488 in 100 µl HBSS (37, 38).  Finally, pellets were labeled with 100 µg of calcofluor white

M2R (Sigma-Aldrich) in 100 µl HBSS for 15 min at RT and washed five times with HBSS (39,

40).  Preparations were mounted in Mowiol mounting medium (Sigma-Aldrich) and observed

with widefield and differential interference contrast microscopy, using a 100x objective of a

Zeiss AXIO inverted microscope with a Colibri LED (Carl Zeiss Microcopy LLC, Thornwood,

NY).  Images were collected at 0.2-µm optical sections with a Hamamatsu Orca-R2 camera

and deconvolved using ZEN software (Zeiss).  Alternatively, SIM was performed with a 63-x

objective of a Zeiss ELYRA S.1 microscope at Boston College (Chestnut Hill, MA), and 0.09-

µm optical sections deconvolved using Zen software (30).

**TEM of intact and purified walls.**  High-pressure freezing and freeze substitution were

used to prepare cyst and cyst walls for TEM at the Harvard Medical School Electron

Microscope facility (35).  To make them noninfectious, we fixed mature cysts in 1%

paraformaldehyde for 10 min at RT and washed them 2 times in PBS.  Cyst walls in PBS were

pelleted, placed in 6-mm Cu/Au carriers, and frozen in an EM ICE high-pressure freezer (Leica

Microsystems, Buffalo Grove, Il). Freeze substitution was performed in a Leica EM AFS2

instrument in dry acetone containing 1% $_{dd}H_2O$, 1% $OsO_4$, and 1% glutaraldehyde at -90°C for

48 hr. The temperature was increased 5°C/hour to 20°C, and samples were washed 3 times in

pure acetone and once in propylene oxide for 10 min each. Samples were infiltrated with 1:1

Epon:propylene oxide overnight at 4°C and embedded in TAAB Epon (Marivac Canada Inc. St.

Laurent, Canada). Ultrathin sections (80 to 100 nm thick) were cut on a Leica Reichert

Ultracut S microtome, picked up onto copper grids, stained with lead citrate, and examined in a

JEOL 1200EX transmission electron microscope (JEOL USA, Peabody, MA). Images were

recorded with an AMT 2k CCD camera.

**Mass spectrometry of tryptic and chymotryptic peptides from cyst walls.** Broken

cyst walls, prepared as above, were dissolved into 50 mM $NH_4HCO_3$, pH 8.0, reduced with 50

mM dithiothreithol (DTT) for 20 min at 60°C, alkylated with iodoacetamide (IAA) for 20 min at

RT, and then digested with proteomics grade trypsin (Sigma-Aldrich) overnight at 37°C.

Alternatively broken cyst walls either before or after digestion with trypsin were reconstituted in

1× reducing SDS/PAGE loading buffer and run on a 4–20% precast polyacrylamide TGX gel

(Bio-Rad). Bands stained by colloidal Coomassie blue were excised and washed with 50 mM

$NH_4HCO_3$/acetonitrile (ACN). Reduction, alkylation, and trypsin/chymotrypsin digestion were

performed in-gel. Peptides were dried and desalted using C18 ZipTip concentrators (EMD

Millipore). Peptides from five biological replicates for both *in solution* and *in-situ* hydrolysis

were dissolved in 2% ACN, 0.1% formic acid (FA) and separated using a nanoAcquity-UPLC

system (Waters) equipped with a 5-μm Symmetry C18 trap column (180 μm x 20 mm) and a

1.7-μm BEH130 C18 analytical column (150 μm × 100 mm). Samples were loaded onto the

precolumn and washed for 4 min at a flow rate of 4 μl/min with 100% mobile phase A (99%

Water/1% ACN/0.1% FA).  Samples were eluted to the analytical column with a gradient of 2–

40% mobile phase B (99% ACN/1% Water/0.1% FA) delivered over 40 or 90 min at a flow rate

of 0.5 µl/min.  The analytical column was connected online to a QE or a QE-HF Mass

Spectrometer (Thermo Fisher Scientific) equipped with a Triversa NanoMate (Advion Inc.,

Ithaca, NY) electrospray ionization (ESI) source, which was operated at 1.7 kV.  Data were

acquired in automatic Data Dependent top 10 (QE) or top 20 (QE-HF) mode.  Automated

database searches were performed using the PEAKS software suite version 8.5

(Bioinformatics Solutions Inc., Waterloo, ON, Canada).  The search criteria were set as

follows: trypsin/chymotrypsin as the enzyme with ≤ 3 missed cleavages and ≤ 1 non-specific

cleavage, the error tolerances for the precursor of 5 ppm and 0.05 Da for fragment ions,

carbamidomethyl cysteine as a fixed modification, oxidation of methionine, Pyro-glu from

glutamine, and deamidation of asparagine or glutamine as variable modifications.  The peptide

match threshold (-10 logP) was set to 15, and only proteins with a minimum of two unique

peptides were considered.  The mass spectrometry proteomics data have been deposited to

the ProteomeXchange Consortium (http://proteomecentral.proteomexchange.org) via the

PRIDE partner repository with the dataset identifiers PXD011826 and 10.6019/PXD011826

(77).

**Bioinformatic characterization of candidate CWPs.**  Signal peptides and

transmembrane helices were predicted using SignalP 4.1 and TMHMM, respectively (45, 48).

Glycosylphosphatidylinositol anchors were searched for using big-PI (47).  AmoebaDB, which

contains sequence information from the Neff strain and ten other *Acanthamoeba* strains, was

used to identify genome sequences, predict introns, and identify paralogous proteins (28, 29).

The NR database at the NCBI was used to identify homologs of candidate CWPs in other

species and to identify conserved domains (43). Carbohydrate-binding modules were

searched using CAZy and InterPro databases (27, 44).

**Expression and visualization of GFP-fusions in transfected *A. castellanii*.** We used RT-

PCR from RNA of encysting protists to obtain the coding sequences of a representative

Luke(2) lectin (840-bp CDS of ACA1_377670), Luke(3) lectin (1293-bp CDS of

ACA1_245650), Leo lectin (562-bp CDS of ACA1_074730), and Jonah(1) lectin (1596-bp CDS

of ACA1_164810) (Figs. 3 and S2). Please see Supplemental Table 1 for a list of primers

used to make all the constructs. Using NEBuilder HiFi DNA assembly (New England Biolabs,

Ipswich, MA), we cloned each CDS into the pGAPDH plasmid, which was a kind gift from

Yeonchul Hong of Kyongpook National University School of Medicine, Deagu, Korea (33).

pGAPDH contains a neomycin resistance gene under a TATA-box promoter (for selection with

G418) and a glyceraldehyde 3-phosphate dehydrogenase promoter for constitutive expression

of GFP-fusions (Fig. S1 and Supplemental Table 1). The GFP tag was placed at the C-

terminus of each CWP, and a polyadenylation sequence was added downstream of the GFP-

fusion's stop codon. For expression of CWP genes under their own promoters, we replaced

the GAPDH promoter with 446-bp from the 5 'UTR of the Luke(2) gene, 500-bp from the 5'

UTR of the Luke(3) gene, 486-bp from the 5' UTR of the Leo gene, and 571-bp of the 5'UTR of

the Jonah(1) gene, each cloned from the genomic DNA.

As a control, SP-GFP, which contains a 60-bp sequence encoding an N-terminal signal

peptide of Luke(2) lectin, was expressed behind a GAPDH promoter. As another control, the

470-bp 5' UTR and 525-bp CDS of CSP21 (ACA1_075240) was made with a GFP tag (22).

Transfections in *A. castellanii* were performed as described previously (32, 33) with

some modifications. Briefly, 5 x $10^5$ log-phase trophozoites were allowed to adhere to 6-well

plates in ATCC medium 712 for 30 min at 30°C.  The adherent trophozoites were washed and

replaced with 500 μl of non-nutrient medium (20 mM Tris-HCl (pH 8.8), 100 mM KCl, 8 mM

MgSO$_4$, 0.4 mM CaCl$_2$ and 1 mM NaHCO$_3$).  In an Eppendorf tube, 4 μg of Midiprep (PureLink

HiPure Midiprep Kit, Thermo Fisher Scientific) plasmid DNA was diluted to 100 μl with non-

nutrient medium.  Twenty microliters of SuperFect Transfection Reagent (Qiagen Inc,

Germantown, MD) was added to the DNA suspension, mixed gently by pipetting five times,

and incubated for 10 min at RT.  Six hundred microliters of non-nutrient medium were added to

the DNA-SuperFect mix, and the entire suspension was added to the trophozoites adhering to

the 6-well culture plate.  The culture plate was incubated for 3 hr at 30°C, after which the non-

nutrient medium was replaced with ATCC medium 712 and incubated for another 24 hr at

30°C.  To select for transfectants, we added 12.5 μg/ml of Gibco G418 antibiotic (Thermo

Fisher Scientific) to the culture after 24 hr, and we changed the medium plus antibiotic every 4

days.  After 2 to 4 weeks, the transfectants were growing robustly in the presence of the

antibiotic, and trophozoites and/or cysts expressing GFP were detected by widefield

microscopy.  These organisms were induced to encyst, fixed, labeled with WGA and CFW, and

examined by widefield microscopy and SIM, as described above.

**Binding of MBP-CWP fusions to microcrystalline cellulose and chitin beads.**  MBP-fusion

constructs were prepared by cloning the cDNA of a representative Luke(2) lectin (60 to 843-bp

CDS of ACA1_377670) and a representative Jonah(1) lectin (70 to 1599-bp CDS of

ACA1_164810) without their signal sequences into pMAL-p2x vector (New England Biolabs)

for periplasmic expression in BL21-CodonPlus(DE3)-RIPL (Agilent Technologies, Lexington,

MA) (34).  For the MBP-fusion, the Leo CDS without the signal sequence (67 to 564-bp of

ACA1_074730) was codon optimized (GenScript) and cloned into pMAL-p2x vector.  MBP-

Luke(2) was induced with 250 μM IPTG for 5 hr at RT; MBP-Jonah(1) was induced with 1 mM

IPTG for 5 hr at RT; and MBP-Leo was induced with 250 μM IPTG for 3.5 hr at $37^0$C. MBP-

fusion proteins were purified with amylose resin following the manufacturer's instructions (GE

Healthcare, Pierce, Agilent Technologies and New England Biolabs). MBP-fusions (1 μg each

in 100 μl of 1% NP40 ) were incubated with 0.5 μg Avicel microcrystalline cellulose (Sigma-

Aldrich) or a 50-μl slurry of magnetic chitin beads (New England Biolabs) for 3 hr at $4^0$C with

rocking. Cellulose was centrifuged, and magnetic chitin beads were placed in a magnet

to collect the supernatant (unbound fraction) and pellet (bound fraction). The pellet was

washed three times with 1% NP40. To solubilize proteins, the input material (total), unbound,

and bound fractions were boiled in SDS sample buffer. MBP-proteins were separated on

SDS-PAGE gels, blotted to PVDF membranes, blocked in 5% BSA, and detected using anti-

MBP antibodies (New England Biolabs).

**Western blots of _A. castellanii_ trophozoite and cyst lysates probed with anti-lectin**

**rabbit antibodies.** Log-phase trophozoites and 36-hr-old cysts were harvested, and the total

protein solubilized in SDS sample buffer, run in SDS-PAGE gels, blotted on PVDF

membranes, and blocked in 5% BSA. MBP-CWP fusions and MBP alone were run in adjacent

lanes as positive and negative controls, respectively. The blots were probed with 1:100

dilutions of rabbit polyclonal antibodies (Li International) raised to 16- or 50-amino acid

peptides of representative Luke(2) lectin (residues 230-279 of ACA1_377670), Leo lectin

(residues 124-139 of ACA1_074730) and Jonah(1) lectin (residues 362-411 of ACA1_164810)

(Fig. 3). A 1:1000 dilution of anti-rabbit IgG-HRP (BioRad) was used as secondary antibody

and Super Signal West Pico PLUS (ThermoFisher Scientific) for chemiluminescent detection.

Coomassie stained gels were run in parallel for loading control.

**ACKNOWLEDGMENTS.** We thank Maria Ericsson of Harvard University for help with transmission electron microscopy. We thank Bret Judson and Marc-Jan Gubbels of Boston College for help with structured illumination microscopy, which was supported by NSF grant No. 1626072. This work was also supported by awards from the National Institute of Allergy and Infectious Diseases of the NIH to J.S. (R01 AI110638), from the National Institute of General Medical Science of the NIH to C.E.C. (P41 GM104603), and from the Department of Energy to B.R.U. (DESC0015662). We also thank Dean Jeffrey Hutter of the Boston University Goldman School of Dental Medicine for his support of research in the Samuelson lab.

**REFERENCES**

1. Clarke M, Lohan AJ, Liu B, Lagkouvardos I, Roy S, Zafar N, et al. Genome of *Acanthamoeba castellanii* highlights extensive lateral gene transfer and early evolution of tyrosine kinase signaling. Genome Biol 2013;14(2):R11.

2. Clarke B, Sinha A, Parmar DN, Sykakis E. Advances in the diagnosis and treatment of *Acanthamoeba* keratitis. J Ophthalmol 2012:484892.

3. Carrijo-Carvalho LC, Sant'ana VP, Foronda AS, de Freitas D, de Souza Carvalho FR. Therapeutic agents and biocides for ocular infections by free-living amoebae of *Acanthamoeba* genus. Surv Ophthalmol 2017;62(2):203-218.

4. Carnt N, Robaei D, Minassian DC, Dart JKG. *Acanthamoeba* keratitis in 194 patients: risk factors for bad outcomes and severe inflammatory complications. Br J Ophthalmol. 2018;102(10):1431-35.

5.  Satlin MJ, Graham JK, Visvesvara GS, Mena H, Marks KM, Saal SD, et al. Fulminant and fatal encephalitis caused by *Acanthamoeba* in a kidney transplant recipient: case report and literature review. Transpl Infect Dis 2013;15(6):619-26.

6.  Lorenzo-Morales J, Martín-Navarro CM, López-Arencibia A, Arnalich-Montiel F, Piñero JE, Valladares B. *Acanthamoeba* keratitis: an emerging disease gathering importance worldwide? Trends Parasitol 2013:29(4):181-7.

7.  Carnt N, Hoffman JJ, Verma S, Hau S, Radford CF, Minassian DC, et al. *Acanthamoeba* keratitis: confirmation of the UK outbreak and a prospective case-control study identifying contributing risk factors. Br J Ophthalmol 2018:312544.

8.  Pittet D, Allegranzi B, Boyce J. The World Health Organization Guidelines on Hand Hygiene in Health Care and their consensus recommendations. Infect Control Hosp Epidemiol 2009;30(7):611-22.

9.  Aqeel Y, Rodriguez R, Chatterjee A, Ingalls RR, Samuelson J. Killing of diverse eye pathogens (*Acanthamoeba castellanii*, *Fusarium solani*, and *Chlamydia trachomatis*) with alcohols. PLoS Negl Trop Dis 2017;11(2):e0005382.

10. Neff RJ, Ray SA, Benton WF, Wilborn M. Induction of synchronous encystment (differentiation) in *Acanthamoeba* sp. Methods Cell Biol 1964;1:55-83.

11. Jensen T, Barnes WG, Meyers D.  Axenic cultivation of large populations of *Acanthamoeba castellanii* (JBM). J Parasitol 1970;56(5):904–6.

12. Lloyd D. Encystment in *Acanthamoeba castellanii*: a review. Exp Parasitol 2014;145:Suppl:20-7.

13. Bowers B, Korn ED. The fine structure of *Acanthamoeba castellanii* (Neff strain). II. Encystment. J Cell Biol 1969;41(3): 786–805.

14. Chávez-Munguía B, Omaña-Molina M, González-Lázaro M, González-Robles A, Bonilla

P, Martínez-Palomo A. Ultrastructural study of encystation and excystation in

*Acanthamoeba castellanii*. J Eukaryot Microbiol 2005;52(2):153-8.

15. Coulon C, Collignon A, McDonnell G, Thomas V. Resistance of *Acanthamoeba* cysts to

disinfection treatments used in health care settings. J Clin Microbiol 2010;48(8): 2689-

2697.

16. Dupuy M, Berne F, Herbelin P, Binet M, Berthelot N, Rodier MH, et al. Sensitivity of

free-living amoeba trophozoites and cysts to water disinfectants. Int J Hyg Environ

Health 2014(2-3);217:335-9.

17. Lonnen J, Putt KS, Kernick ER, Lakkis C, May L, Pugh RB. The efficacy of

*Acanthamoeba* cyst kill and effects upon contact lenses of a novel ultraviolet lens

disinfection system. Am J Ophthalmol 2014;158(3):460-8.

18. Flores BM, Garcia CA, Stamm WE, Torian BE. Differentiation of *Naegleria fowleri* from

*Acanthamoeba* species by using monoclonal antibodies and flow cytometry. J Clin

Microbiol 1990;28(9):1999-2005.

19. Hiwatashi E, Tachibanabi H, Kanedab Y, Obazawaa H. Production and characterization

of monoclonal antibodies to *Acanthamoeba castellanii* and their application for detection

of pathogenic *Acanthamoeba spp*. Parasitol Internat 1997;46(3):197-205.

20. Turner ML, Cockerell EJ, Brereton HM, Badenoch PR, Tea M, Coster DJ, et al.

Antigens of selected *Acanthamoeba* species detected with monoclonal antibodies. Int J

Parasitol 2005;35(9):981-90.

21. Kang AY, Park AY, Shin HJ, Khan NA, Maciver SK, Jung SY. Production of a monoclonal antibody against a mannose-binding protein of *Acanthamoeba culbertsoni* and its localization. Exp Parasitol 2018;192:19-24.

22. Hirukawa Y, Nakato H, Izumi S, Tsuruhara T, Tomino S. Structure and expression of a cyst specific protein of *Acanthamoeba castellanii*. Biochim Biophys Acta 1998;1398(1):47-56.

23. Potter JL, Weisman RA. Cellulose synthesis by extracts of *Acanthamoeba castellanii* during encystment. Stimulation of the incorporation of radioactivity from UDP-($^{14}$C)glucose into alkali-soluble and insoluble beta-glucans by glucose 6-phosphate and related compounds. Biochim Biophys Acta 1976;428(1):240-52.

24. Derda M, Winiecka-Krusnell J, Linder MB, Linder E. Labeled *Trichoderma reesei* cellulase as a marker for *Acanthamoeba* cyst wall cellulose in infected tissues. Appl Environ Microbiol 2009;75(21):6827-30.

25. Dudley R, Jarroll EL, Khan NA. Carbohydrate analysis of *Acanthamoeba castellanii*. Exp Parasitol 2009;122(4):338-43.

26. West CM. Comparative analysis of spore coat formation, structure, and function in *Dictyostelium*. Int Rev Cytol 2003;222:237-93

27. Lombard V, Golaconda Ramulu H, Drula E, Coutinho PM, Henrissat B. The Carbohydrate-active enzymes database (CAZy) in 2013. Nucleic Acids Res 2014;42 (Database issue):D490–D495.

28. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, et al. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. Nucleic Acids Res 1997;25:3389-3402.

29. Aurrecoechea C, Barreto A, Brestelli J, Brunk BP, Caler EV, Fischer S, et al. AmoebaDB and MicrosporidiaDB: functional genomic resources for Amoebozoa and *Microsporidia* species. Nucleic Acids Res 2011;39:D612-9.

30. Demmerle J, Innocent C, North AJ, Ball G, Müller M, et al. Strategic and practical guidelines for successful structured illumination microscopy. Nat Protoc 2017;12(5):988-1010.

31. Samuelson J, Bushkin GG, Chatterjee A, Robbins PW. Strategies to discover the structural components of cyst and oocyst walls. Eukaryot Cell 2013;12(2):1578-87.

32. Peng Z, Omaruddin R, Bateman E. Stable transfection of *Acanthamoeba castellanii*. Biochim Biophys Acta 2005;1743(1-2):93-100.

33. Bateman E. Expression plasmids and production of EGFP in stably transfected *Acanthamoeba*. Protein Expr Purif 2010;70(1):95-100.

34. Kapust RB, Waugh DS. *Escherichia coli* maltose-binding protein is uncommonly effective at promoting the solubility of polypeptides to which it is fused. Protein Sci 1999;8(8):1668-74.

35. Vanhecke D, Graber W, Studer D. Close-to-native ultrastructural preservation by high pressure freezing. Methods Cell Biol 2008;88:151-64.

36. Monheit JE, Cowan DF, Moore Dg. Rapid detection of fungi in tissues using calcofluor white and fluorescence microscopy. Arch Pathol Lab Med 1984;108(8);616-8.

37. Shaw JA, Mol PC, Bowers B, Silverman SJ, Valdivieso MH, Durán A, et al. The function of chitin synthases 2 and 3 in the *Saccharomyces cerevisiae* cell cycle. J Cell Biol 1991;114(1):111-23.

38. Frisardi M, Ghosh SK, Field J, Van Dellen K, Rogers R, Robbins P, et al. The most abundant glycoprotein of amebic cyst walls (Jacob) is a lectin with five Cys-rich, chitin-binding domains. Infect Immun 2000;68(7):4217-24.

39. Said-Fernández S, Campos-Góngora E, González-Salazar F, Martínez-Rodríguez HG, Vargas-Villarreal J, Viader-Salvadó JM. Mg2+, Mn2+, and Co2+ stimulate *Entamoeba histolytica* to produce chitin-like material. J Parasitol 2001;87(4):919-23.

40. Wilhelmus KR, Osato MS, Font RL, Robinson NM, Jones DB. Rapid diagnosis of *Acanthamoeba* keratitis using calcofluor white. Arch Ophthalmol 1986;104(9):1309-12.

41. Smith DB. Generating fusions to glutathione S-transferase for protein studies. Methods Enzymol 2000;326:254-70.

42. Urbanowicz BR, Catalá C, Irwin D, Wilson DB, Ripoll DR, Rose JK. A tomato endo-beta-1,4-glucanase, SlCel9C1, represents a distinct subclass with a new family of carbohydrate binding modules (CBM49). J Biol Chem 2007;282(16):12066-74

43. Marchler-Bauer A, Derbyshire MK, Gonzales NR, Lu S, Chitsaz F, Geer LY, et al. CDD: NCBI's conserved domain database. Nucleic Acids Res 2015;43(Database issue):D222-6.

44. Finn RD, Attwood TK, Babbitt PC, Bateman A, Bork P, Bridge AJ, et al. InterPro in 2017 — beyond protein family and domain annotations. Nucleic Acids Res 2017;45(D1):D190-D199.

45. Nielsen H. Predicting Secretory Proteins with SignalP. Methods Mol Biol 2017;1611:59-73.

46. Wang Y, Slade MB, Gooley AA, Atwell BJ, Williams KL. Cellulose-binding modules from

extracellular matrix proteins of *Dictyostelium discoideum* stalk and sheath. Eur J

Biochem 2001;268(15):4334-45.

47. Eisenhaber B, Bork P, Eisenhaber F. Prediction of potential GPI-modification sites in

proprotein sequences. J Mol Biol 1999;292(3):741-58.

48. Krough A, Larsson B, von Heijne G, Sonnhammer EL. Predicting transmembrane

protein topology with a hidden Markov model: application to complete genomes. J Mol

Biol. 2001;305(3):567-80.

49. Xu Y, Liang X, Chen Y, Koehler TM, Hook M. Identification and biochemical

characterization of two novel collagen binding MSCRAMMs of *Bacillus anthracis*. J Biol

Chem 2004;279(50):51760-8.

50. Enguita FJ, Martins LO, Henriques AO, Carrondo MA. Crystal structure of a bacterial

endospore coat component. A laccase with enhanced thermostability properties. J Biol

Chem 2003;278(21):19416-2.

51. Kiiskinen LL, Palonen H, Linder M, Viikari L, Kruus K. Laccase from *Melanocarpus*

*albomyces* binds effectively to cellulose. FEBS Lett 2004;576(1-2):251-5.

52. Moon EK, Chung DI, Hong YC, Kong HH. Characterization of a serine proteinase

mediating encystation of *Acanthamoeba*. Eukaryot Cell 2008 Sep;7(9):1513-7.

53. Corsaro D, Köhsler M, Montalbano Di Filippo M, Venditti D, Monno R, et al. Update on

*Acanthamoeba jacobsi* genotype T15, including full-length 18S rDNA molecular

phylogeny. Parasitol Res 2017 Apr;116(4):1273-1284.

54. Chen L, Orfeo T, Gilmartin G, Bateman E. Mechanism of cyst specific protein 21 mRNA induction during *Acanthamoeba* differentiation. Biochim Biophys Acta 2004;1691(1):23-31.

55. Vollmer AC, Bark SJ. Twenty-Five Years of Investigating the Universal Stress Protein: Function, Structure, and Applications. Adv Appl Microbiol 2018;102:1-36.

56. Chatterjee A, Ghosh SK, Jang K, Bullitt E, Moore L, Robbins PW, Samuelson J. Evidence for a "wattle and daub" model of the cyst wall of entamoeba. PLoS Pathog. 2009;5(7):e1000498.

57. Chatterjee A, Carpentieri A, Ratner DM, Bullitt E, Costello CE, Robbins PW, et al. *Giardia* cyst wall protein 1 is a lectin that binds to curled fibrils of the GalNAc homopolymer. PLoS Pathog 2010;6(8):e1001059 .

58. Bushkin GG, Motari E, Magnelli P, Gubbels MJ, Dubey JP, Miska KB, et al. β-1,3-glucan, which can be targeted by drugs, forms a trabecular scaffold in the oocyst walls of *Toxoplasm*a and *Eimeria*. MBio 2012;3(5): e00258-12.

59. Vinayak S, Pawlowic MC, Sateriale A, Brooks CF, Studstill CJ, Bar-Peled Y, et al. Genetic modification of the diarrhoeal pathogen *Cryptosporidium parvum*. Nature 2015;523(7561):477-80

60. Lee S, Ginese M, Beamer G, Danz HR, Girouard DJ, Chapman-Bonofiglio SP, et al. Therapeutic Efficacy of Bumped Kinase Inhibitor 1369 in a Pig Model of Acute Diarrhea Caused by *Cryptosporidium hominis*. Antimicrob Agents Chemother 2018;62(7): e00147-18.

61. Cherif M, Benhamou N, Blanger RR. Occurrence of cellulose and chitin in the hyphal

walls of *Pythium ultimum:* a comparative study with other plant pathogenic fungi. Can J

Microbiol 1993;39(2):212-223.

62. West CM, Nguyen P, van der Wel H, Metcalf T, Sweeney KR, Blader IJ, et al.

Dependence of stress resistance on a spore coat heteropolysaccharide in

*Dictyostelium.* Eukaryot Cell 2009;8(1):27-36.

63. Van Dellen K, Ghosh SK, Robbins PW, Loftus B, Samuelson J. *Entamoeba histolytica*

lectins contain unique 6-Cys or 8-Cys chitin-binding domains. Infect Immun

2002;70(6):3259-63.

64. Ghosh SK, Van Dellen KL, Chatterjee A, Dey T, Haque R, Robbins PW, et al. The

Jacob2 lectin of the *Entamoeba histolytica* cyst wall binds chitin and is polymorphic.

PLoS Negl Trop Dis 2010;4(7):e750.

65. Gas-Pascual E, Ichikawa HT, Sheikh MO, Serji MI, Deng B, Mandalasi M, et al.

CRISPR/Cas9 and glycomics tools for *Toxoplasma* glycobiology. J Biol Chem 2018:

jbc.RA118.006072.

66. McInally SG, Hagen KD, Nosala C, Williams J, Nguyen K, Booker J, Jones K, Dawson

SC. Robust and stable transcriptional repression in *Giardia* using CRISPRi. Mol Biol

Cell 2018:mbcE18090605.

67. CAZypedia Consortium. Ten years of CAZypedia: a living encyclopedia of

carbohydrate-active enzymes. Glycobiology 2018;28(1):3-8.

68. Van Dellen KL, Chatterjee A, Ratner DM, Magnelli PE, Cipollo JF, Steffen M, et al.

Unique posttranslational modifications of chitin-binding lectins of *Entamoeba invadens*

cyst walls. Eukaryot Cell 2006;5(5):836-48.

69. Shen Z, Jacobs-Lorena M. A type I peritrophic matrix protein from the malaria vector

*Anopheles gambiae* binds to chitin. Cloning, expression, and characterization. J Biol

Chem 1998273(28):17665-70.

70. Stibbs HH. Monoclonal antibody-based enzyme immunoassay for *Giardia lamblia*

antigen in human stool. J Clin Microbiol 1989;27(11):2582-8.

71. Spadafora LJ, Kearney MR, Siddique A, Ali IK, Gilchrist CA, Arju T, et al. Species-

Specific Immunodetection of an *Entamoeba histolytica* Cyst Wall Protein. PLoS Negl

Trop Dis 2016;10(5):e0004697.

72. Chaudhary PM, Tupe SG, Deshpande MV. Chitin synthase inhibitors as antifungal

agents. Mini Rev Med Chem 2013;13(2):222-36.

73. Aqeel Y, Siddiqui R, Khan NA. Silencing of xylose isomerase and cellulose synthase by

siRNA inhibits encystation in *Acanthamoeba castellanii*. Parasitol Res 2013;112:1221-7.

74. Moon EK, Hong Y, Chung DI, Goo YK, Kong HH. Down-regulation of cellulose synthase

inhibits the formation of endocysts in *Acanthamoeba*. Korean J Parasitol 2014;52:131-5.

75. Moon EK, Hong Y, Chung DI, Goo YK, Kong HH. Potential Value of Cellulose Synthesis

Inhibitors Combined With PHMB in the Treatment of *Acanthamoeba* Keratitis. Cornea

2015;34(12):1593-8.

76. Lakhundi S, Siddiqui R, Khan NA. Cellulose degradation: a therapeutic strategy in the

improved treatment of *Acanthamoeba* infections. Parasit Vectors 2015;8:23.

77. Vizcaino JA, Deutsch EW, Wang R, Csordas A, Reisinger F, Rios D, et al.

ProteomeXchange provides globally coordinated proteomics data submission and

dissemination. Nat Biotechnol 2014;32(3):223-6.

 **Figure legends.**

 **FIG 1** TEM showed purified *A. castellanii* cyst walls retained endocyst and ectocyst layers and

 ostioles. A, B. Intact cysts and purified cyst walls had an outer ectocyst layer (red arrows), an

 inner endocyst layer (blue arrows), and ostioles (green arrows) that connect the layers.

 Endocyst and ectocyst layers had the same appearance in intact cysts (C) and purified cyst

 walls (D). Purified cyst walls were missing amorphous material (purple arrow) between the

 wall and the plasma membrane of intact cysts. E. At the edge of the ostiole of an intact cyst,

 the endocyst layer bifurcated, and the outer branch met the ectocyst layer. In the center of the

 ostiole, the ectocyst layer formed a narrow cap over the inner branch of the endocyst layer.

 Scale bars as marked on micrographs.

 **FIG 2** SIM also showed purified *A. castellanii* cyst walls retained distinct ectocyst layer (red

 arrow) and endocyst layer (blue arrow), as well as ostioles (green arrow). The ectocyst layer

 of intact cysts (A) and purified cyst walls (B) labeled red with GST-AcCBM49; the edges of

 ostioles labeled green with WGA; and the endocyst layer labeled blue with CFW. Scale bars

 are 2 µm.

 **FIG 3** Representative proteins of abundant families of CWPs contained two CBM49s (Luke(2)

 lectin), two 8-Cys domains (Leo lectin), or one CAA domain (Jonah(1) lectin). The Luke(2)

 lectin had an N-terminal signal peptide (purple) and two CBM49s separated by short Ser- and

 Pro-rich spacers (light blue). The N-terminal CBM49 contained four Trp residues (red Ws),

 three of which were conserved in a C-terminal CBM49 of a tomato cellulase (larger font) and

 three of which were conserved in a single CBM49 of *Dictyostelium* spore coat proteins

 (underlined). In contrast, the C-terminal CBM49 had two conserved Trp residues. The Leo

 lectin had a signal peptide and two unique domains (dark blue) containing eight Cys residues

each (red Cs).  The Jonah(1) lectin had a signal peptide, a Thr-, Lys-, and Cys-rich domain

(tan), and a single CAA domain (green).  Peptides used to immunize rabbits are underlined.

Representative Luke(3) lectin with three CBM49s, Leo(TRH) lectin with a long Thr-, Lys-, and

His-rich spacer, and Jonah(3) lectin with three CAA domains are shown in Fig. S2.

**FIG 4**  SIM showed a representative Jonah lectin localized to the ectocyst layer of mature cyst

walls, while representative Luke lectins and Leo lectins localized to the endocyst layer and

ostioles.  A. A Jonah(1) lectin with a single CAA domain, which was tagged with GFP and

expressed under its own promoter in transfected *A. castellanii*, localized to the ectocyst layer

of the wall of mature cysts.  WGA labeled both ectocyst and endocyst layers, while CFW

labeled the endocyst layer and ostioles.  B. Leo-GFP with two 8-Cys domains was also present

in the endocyst layer and ostioles.  C. Luke(2)-GFP with two CBM49s was present in the

endocyst layer and sharply outlined conical ostioles of mature cysts.  D. Luke(3) with three

CBM49s was also present in the endocyst layer and ostioles.  A to D. Scale bars are 2 µm.

**FIG 5**  Western blots showed MBP-fusions to Luke, Leo, and Jonah lectins all bound to

microcrystalline cellulose, while binding to chitin beads was variable.  MBP-CWP fusions and

MBP alone, made as recombinant proteins in the periplasm of bacteria, were incubated with

microcrystalline cellulose (A) or chitin beads (B).  Total proteins (T), bound proteins (B), and

unbound proteins (U), as well as molecular weight markers (M), were run on SDS-PAGE,

transferred to PVDF membranes, and detected with an anti-MBP reagent.  Full-length products

in total fractions are underlined in red.  MBP-Leo partially bound to microcrystalline cellulose

and bound weakly, if at all to chitin.  MBP-Luke(2) and MBP-Jonah(1) each bound more

completely to cellulose than to chitin, while MBP alone (negative control) did not bind to

cellulose or chitin.

803 **Table 1. Candidate CWPs identified by mass spectrometry**

| | ID | # Unique peptides | Coverage (%) | Mass (kDa) |
|---|---|---|---|---|
| **Jonah lectins** | | | | |
| three CAAs | ACA1_157320 | 147 | 38 | 146 |
| | | | | |
| one CAA | ACA1_164810 | 83 | 56 | 58 |
| | ACA1_261530 | 18 | 23 | 55 |
| | ACA1_133400 | 9 | 24 | 44 |
| | ACA1_377440 | 6 | 11 | 47 |
| **Luke lectins** | | | | |
| three CBM49s | ACA1_245650 | 72 | 74 | 44 |
| | ACA1_160160 | 8 | 25 | 43 |
| | ACA1_187760 | 7 | 25 | 42 |
| | ACA1_252830 | 6 | 19 | 44 |
| | ACA1_031530 | 6 | 21 | 43 |
| | ACA1_253650 | 5 | 20 | 42 |
| | ACA1_253500 | 5 | 19 | 42 |
| | ACA1_061050 | 3 | 12 | 43 |
| | ACA1_287530 | 2 | 14 | 43 |
| | | | | |
| two CBM49s | ACA1_377670 | 78 | 68 | 29 |
| | ACA1_096300 | 47 | 77 | 28 |
| | ACA1_246110 | 22 | 70 | 27 |
| **Leo lectins** | | | | |
| two 8-Cys domains | ACA1_074730 | 34 | 82 | 20 |
| | ACA1_351320 | 24 | 44 | 20 |
| | ACA1_394030 | 24 | 44 | 20 |
| | ACA1_394280 | 24 | 36 | 24 |
| | ACA1_083920 | 19 | 68 | 20 |
| | ACA1_394560 | 1 | 10 | 19 |
| | | | | |
| two 8-Cys + TKH | ACA1_188350 | 21 | 20 | 59 |
| | ACA1_374130 | 7 | 20 | 52 |
| | ACA1_188550 | 7 | 15 | 46 |
| | ACA1_188370 | 6 | 9 | 68 |
| | ACA1_116240 | 5 | 18 | 56 |
| | ACA1_365840 | 3 | 18 | 44 |
| | ACA1_117050 | 3 | 33 | 36 |
| | ACA1_096640 | 2 | 27 | 37 |

Fig 1

Fig 2

## Luke(2) lectin (ACA1_377670)

MKLLFCLTVLAALIFSCASA

QSCTASLNQTRQAS**W**VDSEQFPRSL<u>W</u>TVEIRNTGQQAVTNVLLSIQGSINQI**W**EVVLVN
DSLYRLPD**W**RLQVGGIPAGQSHVFGFIVNNNSTAAPVSLVSVQC

CBM49

GNVTVPSPSPSASASSVPSPSASASAAPSPSAGASPSSSPRPSPSGG
CSLLASQVARSGAGGS**w**SDNSNRFQIYDITLNNNGASRLTQASLTIAIAADQAIAQF**w**<u>NL</u>
ERVDATNTFNIESFLLPAPGGSQSGLGYVLQTPLNSTSDGSSIGAVNFRC

CBM49

## Leo lectin (ACA1_074730)

MNRFALCFFVVLCAASAAMG QR

**C**PGSDFFPDQRG**CC**PVVVNSVPFYADAQG**C**FPRIISGVAFYADAGG**C**YPSSAGVYRSAGAV**C**PADRV**C**EPE**C**PDMAIVVAEQQGGARL
**C**ANGFFSDVNG**CC**<u>PRVLVDNVPSYRDAQG</u>**C**YPISIASVIFYADVGR**C**YPNSSGVYRPAGAV**C**PAGQQ**C**VVS**C**DRNYA

## Jonah(1) lectin (ACA1_164810)

MKTTYLLLALAALLFCAVVAASA

DDDDRRGSDMNHWDDDDEGHEWDNEWEDEEDGEGRGRGQWDDGREKCRTTTKKCRTTTKKCGTTTKHCPTTTTKRCG
TTTKDCGTTTTKSCGTTTKDCGTTTMKPCTTTSKPVEKCDFAGRDNCLKASEDKFEECVDRKKCAFTVCELEAEEKF
KKCLRGRDGRDGGRENPFEREGENDRQGNERFDQCKKEQLEDKKQCLIQRKASHAGCVAAKECRDFKCLAFIKDCKP
CPTPAATTTTVRVTTH

PPTVGSSCPAEDFFGPAADLNVLVTGSDKLIYDLDMVNGDVEGRVAVNGGFRVKSFGTAQAYSCPDTANYASMFNLI
VNGKMDYSNGQLFC<u>GSSISLSSMNDPSLRQKDFGATNARVAAGKTIKDVTGFDFVATAGYLQGV</u>SNYFSKYAPGTHS
HFRVTEVSPADVNKLNIEGGDYVQVINIKGSGDINFSNFEIPTALNPSQVIYNVVGNNNIKISGFGLKGHLLALNSD
VVLENGHVAGNVYVRSLVGQGSGQVNLAPCP

CAA

Fig 3

Fig 4

## A. Microcrystalline cellulose

anti-MBP antibody

## B. Chitin beads

anti-MBP antibody

Fig 5

**Supplemental files include eight figures, two Excel files, and one table.**

**FIG S1** Constructs made to localize CWPs Ac cyst wall and to determine their binding to microcrystalline cellulose and chitin beads. A. A representative Luke(2) lectin with two CBM49s was used to make GFP-, GST-, and MBP-fusions. A representative Luke(3) lectin with three CBM49s was used to make a GFP-fusion protein. B. A representative Leo lectin was made into GFP- and MBP-fusions. C. A representative Jonah(1) lectin with a single CAA domain was made into GFP- and MBP-fusions. D. Vectors for expressing GFP-fusions in transfected *A. castellanii* under its own promoter contain a neomycin resistance gene under a TATA-binding protein promoter (32, 33). Primers for making constructs are listed in Supplemental Table 1.

**FIG S2** Sequences of candidate CWPs, which differ in at least one essential property from Luke, Leo, and Jonah lectins used for localization and binding studies. A Luke(3) lectin has an N-terminal signal peptide (purple) and three CBM49s separated by short Ser- and Pro-rich spacers (light blue). The CBM49s contain conserved Trp (red Ws) present in the representative Luke lectin (Fig. 3). A Leo(TKH) lectin has a signal peptide, two domains containing eight Cys residues each (red Cs), and a long Thr-, Lys-, and His-rich spacer (brown). A Jonah(3) lectin has three CAA domains (green), hydrophobic regions (tan), and short Ser- and Pro-rich spacers (light blue).

**FIG S3** RT-PCR shows mRNAs of representative Luke, Leo, and Jonah lectins, as well as those of cellulose synthase, are encystation-specific. DNA and total RNA were extracted from trophozoites and organisms encysting for one to three days. RT-PCRs were performed with primers specific for segments of each CWP mRNA, as well as primers specific for segments of mRNAs for GAPDH and cellulose synthase (see Supplemental Table 1). PCR with DNA was used as a positive control, while omission of reverse-transcriptase (-RT) was used as a negative control. Messenger RNAs encoding CWPs and cellulose synthase were absent or nearly absent in trophozoites but were easily detectable in encysting organisms. In contrast, mRNAs for GAPDH were expressed by both trophozoites and encysting organisms (33).

**FIG S4** Western blots with rabbit antibodies to peptides of Jonah and Leo lectins show each CWP is absent in trophozoites but is easily detected in mature cysts. A. Coomassie blue stain of proteins of lysed trophozoites and cysts, as well as molecular weight standards (M). B. Western blotting showed rabbit antibodies to a 50-amino acid peptide of a representative Jonah(1) lectin (underlined in Fig. 3) bound to a cyst protein of the predicted size (red underline) and to an MBP-Jonah(1) fusion-protein made in the periplasm of bacteria. The antibody also bound to degradation products of Jonah(1) lectin. In contrast, the anti-Jonah(1) antibody did not bind to either trophozoites or MBP alone (negative controls). C. Rabbit antibodies to a 16-amino acid peptide of a representative Leo lectin also bound to cyst proteins and to an MBP-Leo fusion but not to trophozoite proteins or to MBP alone. In addition to Leo of the predicted size (red underline), anti-Leo antibodies to a higher molecular weight form, which may be a dimer. These results confirmed encystation-specific expression of Jonah(1) and Leo lectins (Figs. 4 and S3). None of the rabbit antibodies the CWP peptides were useful for labeling cyst walls for widefield microscopy or SIM.

**FIG S5.** Control GFP constructs localize to the cytosol (CSP21-GFP) and secretory vesicles (GFP with an N-terminal signal peptide, SP-GFP) of mature cysts. A. The 21-kDa cyst-specific protein (CSP21) fused to GFP was absent in trophozoites (not shown) but formed punctate structures in the cytosol of cysts (22). B. GFP with an N-terminal signal peptide from Luke lectin and expressed under a GAPDH promoter localized to secretory vesicles of mature cysts (33). These controls make it unlikely that localizations of CWP-GFP constructs to the cyst wall were artifacts (Fig. 4).

**FIG S6**  Contrasting uses of CBM49 by *A. castellanii* and *D. discoideum*.  CBM49, which was first shown to be a cellulose-binding domain at the C-terminus of tomato cellulase, is repeated two or three times in Luke lectins of *A. castellanii* and is also present at the N-terminus of a metalloprotease. In contrast, CBM49 is present in a single copy in the majority of *D. discoideum* proteins and as three copies in rare proteins.  CBM49 is also present in two copies in a *D. discoideum* cysteine protease and as a single copy in a GH5 glycoside hydrolase.

**FIG S7**  *A. castellanii* Leo lectins and *E. histolytica* Jacob lectins have common structures, even though they share no common ancestry (convergent evolution).  Abundant cyst wall proteins of *A. castellanii* (Leo lectins) and *E. histolytica* (Jacob lectins) have unique 8-Cys or 6-Cys domains, respectively, that bind cellulose or chitin.  In each protist, some of the lectins lack spacers, while others have spacers rich in Thr, Lys, and His (*A. castellanii*) or Ser and Thr (*E. histolytica*).

**FIG S8**  Contrasting use of choice of anchor A (CAA) domains in *A. castellanii*, oomycetes, and bacteria.  Jonah lectins, which are abundant in cyst walls of *A. castellanii*, have one or three CAA domains.  The former are preceded by Thr-, Lys-, and Cys-rich sequences (gray), while the latter are separated by Ser- and Pro-rich spacers (blue) and hydrophobic domains (tan).  Predicted proteins of oomycetes (*Pyromyces* or *Neocallimastix*) have three to five CAA domains, while the spore coat protein of *Bacilllus* has a single CAA domain attached to a collagen-binding domain, which is absent in *A. castellanii*.

**Excel file S1**  This file lists the most abundant proteins identified by mass spectrometry of cyst walls purified on the Percoll gradient and retained on a membrane containing 8-μm pores. Proteins with <7 unique peptides have been removed, because they are dominated by cytosolic contaminants.  Luke, Leo, and Jonah lectins have been highlighted in orange.  Other candidate CWPs are marked in yellow.

**Excel file S2**  Complete list of proteins identified by mass spectrometry of cyst walls. This list includes a preparation that was heavily contaminated with cytosolic proteins, because the Percoll gradient and porous membrane were omitted during their purification. Only proteins with at least two unique peptides are included.

**Supplemental Table 1**  Primers used for RT-PCR and construction of GFP- and MBP-fusions.

A. Representative Luke(2) lectin (ACA1_377670)

Representative Luke(3) lectin (ACA1_245650)

B. Representative Leo lectin (ACA1_074730)

C. Representative Jonah lectin (ACA1_164810)

D. Vector for GFP constructs

signal peptide

SP-rich spacer

TKC-rich spacer

GFP — green fluorescent protein

GST — glutathione S-transferase

MBP — maltose-binding protein

GOI — Gene of interest

AcTBP — *A. castellanii* TATA-binding protein promoter

NeoR — Neomycin resistance gene

Fig S1

## Luke(3) lectin (ACA1_245650)

MKQQHQLVLLVFAALFCLTAA

DCAVRIEQSLGSA**W**NGAGGVPMSQ<u>**W**</u>NARIVSSGSETVTALQVAFGAPTTAIDQV**W**GLEPVAGASRVY     CBM49
DLPDYVILNGGLASGQAFN<u>**W**</u>GYIWESSAQAPLSVASVQC

SGGASPTAAPSASSSPVTAPSASSSPVTAPSASP

TIVASPAPCQLSVSQTTRPSSAGGS**w**TEGDFFFQIYDLTLLNSGSRPVSSAVIAIDTTSNQQQVITQF**w**N     CBM49
LERQSATSDLFNVRNPGGNIEVGATLGAGYIVRTPLSAGSQVPPTTRLVSVNC

VGGPSPTIVATPQPSASPTIVASPQPSASPAAG

CNAAVSIVARSAAAGGS**W**TTGPNQFFQIFDITITNTGQRPLNGGVLTFGLPVAGSTITQ**w**WELNRQGNT     CBM49
NVFNVAFNFGPLLVGASQGAGIVVQTSSPSQALPSAVLSNLACAA

## Leo(TKH) lectin (ACA1_374130)

MTPATKLFLVLALGLVLVATAMA DDKKKHDSKGP

**C**KGGYYPDRKG**CC**PAIIKGKAYYRDGSK**C**YPKDVG**C**GVYYADGHGYYPDEKGNYAKDGEE**C**PKERK**C**HER**C**IRPEHHEKS
TTTEECPKETTSTRRETTTTERRTTTPKRTTTTTEECPKETTTTKKHTTTTTEDCPKETTTTKKHTTTTTEECPKETTSTKR
HTTTTEECPKETTSTRKHSTSTKKHTTTTEECPKETTSTKKHYTTSTEECPKETTTTKKHTTTTTEECPKETTTTKKHYTT
STTEECPKETTTTKKHYTTSTKKHHTTSTEECPKETTSTRKETTSTKKHHTTSTTEECPKETTSTRKETTSTRRETTSTK
KHTTTTEECRKTTTEKPCDDDHDKDNK

**C**PGGQWKDGKG**CC**PRKIGHQWYYRDAHG**C**YPVERD**C**GVFYADGKGYYPDENGQYAKAGEY**C**PEYRK**C**KPE**C**DGGYGHY

## Jonah(3) lectin (ACA1_157320)

MSRSRHRVSAPFWGMLLLFSLAALACTCPVAQRSAEPVGLSVGLGVGQSLALALGRGLRFGFGFYSALCFGFRIGQQAAVAFCIR
VGFAGRQQVPFAFGIGLSLALAFCVGFGFGFRFRFGVGVAEQVGLAVSVGFGFAEQVSLALCFGIGVGIGLAIAHGFTVGIGLAI
AHGRNCTTALFNGAQSFDTFLFGSATIQNGQFGGRVAISGNANLTQFGIANALPCTNADARNFNLIVNGNLTSVNGVACGSALVT     CAA
GTVVSAPNFRQEAQGAGVVSGSASMAAVDFDQSFSSLLFTNQQICSLTCKQAAITGSGAVTFDFDNTTNVYCLNAADASRATSFR
FTNFPVDFAGALAINILGSAGETVTLSNAATNLGNVKPTQIVWNVCNGLNVRLSAFTLFGSLLAPASSINITNAQQSLLR
ILLPLGLAFAHPVGLPIGLPLGLGLRFAFGCRLGVTFTLGLCHVITVGFRFALGVSLGGCFALGFGFTLCRRFGFSLGLCHVITV
GIPVRFALGGCVGLCFAFCGRFGFGFGIAGRQRVSLGFTTFCGCLCLTLGLTLCGCLGFSVARRQRLGLALSGRFAIGFCFALCGR
FGFALRCCLGLGLARRHA

SASPAASASASPSASASPSAAASPTASPPIVGSPTPQPPII

GGNCSSPIKALDGAADFDTFVLGQGDALTARNGQFGGRVAVNGNARLSSFGMGAELVCAANDSSAFNLIVQGNLSATNGEIFCGS
ALVNGSIQSAPSLRQINQGANVTQGDLLTLSTLDFTESADNLTRINRQLCEAFADNCTQAEIDAFGIVSLRWDNVTNAGRAQVFC     CAA
LNSSLIANATFFEFAGFPANFSGAIVVNILGSEDVRFANAAISAGNLNATQVLWNVCNATRVDIFAFQLFGSLLAPL
SNVSLINAQQIGTTVALNFTGSSFQKLLAPFIGEICLPDHRRHAHPVGHGLALTRRITLCGRLGIAVALCRGLAIGFALR
VSVGLALCLALGLALCLALGLALCLALGLALCRRLSLGFALRIAFGIALRIAVGVALA

SPSISASPSLSASPTASLSPSISASPSAVSPTLSPSVSPTHPSLSGSPSVSRSPSVSPTLSPSVSPTQSASPSISGSPSS
ASPTQSASPSVSPTASASPSPPSNNVVG

AGNCTNALFGAAQAFDLFVLGTGDQNVTTGLNVVNGQFAGRVAVNSGARLNDFGIGNSLNCTAANATNFNFIVQNDLNATNGELF
CGSALVGGNVTNPLTFRQVNEGANLVSGVDVSISGIDFASATSFLTGVNQAVCALNCTNVTLGQANELRFTGAAGANSTQIFCVR     CAA
SADLSRAASIVFDFAANFTAPIVVNIQGSPNDTTASFRSAAIQRGRLRSEQLLWNVCDSLDVTLSQFQLFGSLLAPNSSLTISNA
EQEGITVVSTLSGQGFQKRLAPFNGFFCQPL

Fig S2

RT-PCRs

encysting organisms

Luke(2) lectin (ACA1_377670)

Leo lectin (ACA1_074730)

Jonah(1) lectin (ACA1_164810)

Cellulose Synthase (ACA1_349650)

GAPDH

Fig S3

A. Coomassie blue

B. rabbit anti-Jonah(1)
peptide antibody

C. rabbit anti-Leo
peptide antibody

Fig S4

Fig S5

Fig S6

## *A. castellanii* Leo lectins

8-Cys  8-Cys

8-Cys  TKH-rich  8-Cys

8-Cys  TKH-rich spacer  8-Cys

8-Cys  TKH-rich spacer  8-Cys

## *E. histolytica* Jacob lectins

6-Cys  6-Cys

6-Cys  6-Cys  ST-rich spacer  6-Cys

signal
peptide

Fig S7

*A. castellanii* Jonah lectins

*Pyromyces spp.* hypothetical protein

*Neocallimastix californiae* hypothetical protein

*Bacillus anthracis* collagen-binding protein

signal peptide

SP-rich spacer

Fig S8

# Supplemental Table 1

| primers | | template | Components for NEBuilder HiFi DNA Assembly (NEB) | construct | referred to in text as: |
|---|---|---|---|---|---|
| forward | CTCTCTCACTAAGACATATGAATGAACCGCTTTGCTCTCTGC | Ac gDNA | N/A* | N/A | RT-PCR primers for Leo lectin (ACA1_074730) |
| reverse | TTGCTCACGGCGTAGTTGCGGTCGC | | | | |
| forward | CTCTCTCACTAAGACATATGAATGAAGCTTCTCTTCTGCTTGACC | Ac cDNA | N/A | N/A | RT-PCR primers for Luke(2) lectin (ACA1_377670) |
| reverse | TTGCTCACGCAGCGGAAGTTGACGGC | | | | |
| forward | CTCTCTCACTAAGACATATGAATGAAGACAACCTACCTGCTTCTG | Ac cDNA | N/A | N/A | RT-PCR primers for Jonah(1) lectin (ACA1_164810) |
| reverse | TTGCTCACAGGGCAGGGGGCC | | | | |
| forward | TGCTCTTCTTCCGCTTCCTC | Ac cDNA | N/A | N/A | RT-PCR primers for cellulose synthase (ACA1_349650) |
| reverse | GAACGGGTAGCTGTCGATGTC | | | | |
| forward | ATGTCGCACATCAAGGTCG | Ac cDNA | N/A | N/A | RT-PCR primers for GAPDH |
| reverse | GCAGGGTGCCTTTGATTATTTACT | | | | |
| NA; linearized by BglII/NdeI digestion | | GAPDH-ACA1_074730cDNA-EGFP-polyAtailRev (unpublished) | vector | own promoter-ACA1_074730cDNA-EGFP-polyAtailRev | Leo-GFP under its own promoter |
| forward | TCACATGGCTCGACACGAAACGGACAGTCTTGATCATCTCTG | Ac gDNA | fragment | | |
| reverse | CAAAGCGGTTCATTCAGTTGGTGTTGTGTTGGTG | | | | |
| forward | CCTCAACCATCATATGAATGAAGCTTCTCTTCTGCTTGACC | GAPDH-ACA1_377670cDNA-EGFP-polyAtailRev (unpublished) | vector | own promoter-ACA1_377670cDNA-EGFP-polyAtailRev | Luke(2)-GFP under its own promoter |
| reverse | GTTAGACCCTAGATCTGTCGAGCCATGTGAGCAAAAG | | | | |
| forward | GACAGATCTAGGGTCTAACACGTGCCAT | Ac gDNA | fragment | | |
| reverse | CATTCATATGATGGTTGAGGGTTGTAGAGGTTG | | | | |
| forward | CGCCGCCATATCAATGGTGAGCAAGGGCG | GAPDH-EGFP-polyAtailRev (unpublished) | vector | own promoter-ACA1_245650gDNA-EGFP-polyAtailRev | Luke(3)-GFP under its own promoter |
| reverse | CAGAGGTAGAAGATCTGTCGAGCCATGTGAG | | | | |
| forward | GACAGATCTTCTACCTCTGATGACGCTTGATCAAG | Ac gDNA | fragment | | |
| reverse | ATTGATATGGCGGCGCAGGC | | | | |
| NA; linearized by BglII/NdeI digestion | | GAPDH-ACA1_164810cDNA-EGFP-polyAtailRev (unpublished) | vector | own promoter-ACA1_164810cDNA-EGFP-polyAtailRev | Jonah(1)-GFP under its own promoter |
| forward | TCACATGGCTCGACACCCTCATTGCAATTTAAATGCCG | Ac gDNA | fragment | | |
| reverse | GGTAGGTTGTCTTCATTCAGATCGCAGATGGGC | | | | |
| forward | GTGCCTCTGCTACTAGTGTGAGCAAGGGCGAG | GAPDH-ACA1_377670cDNA-EGFP-polyAtailRev (unpublished) | vector | GAPDH-ACA1_377670SignalPeptide-EGFP-polyAtailRev | Luke(2) signal peptide-GFP under the GAPDH promoter |
| reverse | GCTCACACTAGTAGCAGAGGCACAGGAGAAGA | | | | |
| forward | GTGGATGGTGAGCAAGGGCGAG | GAPDH-EGFP-polyAtailRev (unpublished) | vector | own promoter-ACA1_075240cDNA-EGFP-polyAtailRev | CSP21-GFP under its own promoter |
| reverse | CAACGTCGTAGATCTGTCGAGCCATGTGAGCAAAAG | | | | |
| forward | ACAGATCTACGACGTTGATGGAGGAGACA | Ac gDNA | fragment 1 | | |
| reverse | GCGGTTCATCTTCAGATAGTTGGTAGT | | | | |
| forward | TATCTGAAGATGAACCGCATCCGTGGC | Ac cDNA | fragment 2 | | |
| reverse | TTGCTCACCATCCACGAGCTCTCGCG | | | | |
| forward | CCTGCCCTTGAGAATTCAATTCGGATCCTCTAGAGTCG | pMAL-p2x-ACA1_074730 (MBP-Leo)** | vector | pMAL-p2x-ACA1_164810 | MBP-Jonah(1) |
| reverse | TCGTCGTCACCCTGAAAATACAAATTCTCGAATTCTGAAA | | | | |
| forward | TCAGGGTGACGACGACGACCGAC | own promoter-ACA1_164810cDNA-EGFP-polyAtailRev | fragment | | |
| reverse | AATTCTCAAGGGCAGGGGGCC | | | | |
| forward | CCGCTGCTGAGAATTCAATTCGGATCCTCTAGAGTCGACC | pMAL-p2x-ACA1_074730 (MBP-Leo) | vector | pMAL-p2x-ACA1_377670 | MBP-Luke(2) |
| reverse | CAACTCTGACCCTGAAAATACAAATTCTCGAATTCTGAAA | | | | |
| forward | TTCAGGGTCAGAGTTGCACGGCATCGC | own promoter-ACA1_377670cDNA-EGFP-polyAtailRev | fragment | | |
| reverse | AATTCTCAGCAGCGGAAGTTGACGG | | | | |
| *N/A = not applicable | | | | | |
| **construct made by restriction enzyme cloning of codon optimized ACA1_074730 (Genscript) into pMAL-p2x vector (NEB) | | | | | |