Empirical examination of the replicability of associations between brain structure and

psychological variables

Shahrzad Kharabian Masouleh[1,2], Simon B. Eickhoff[1,2], Felix Hoffstaedter[1,2] and Sarah

Genon[1,2], for the Alzheimer's Disease Neuroimaging Initiative*

[1]Institute of Neuroscience and Medicine (INM-7: Brain and Behaviour), Research Centre
Ju☐lich, Ju☐lich, Germany
[2]Institute of Systems Neuroscience, Heinrich Heine University Du☐sseldorf, Du☐sseldorf,
Germany;
9

Author's email addresses:

s.kharabian@fz-juelich.de

s.eickhoff@fz-juelich.de

f.hoffstaedter@fz-juelich.de

s.genon@fz-juelich.de

Corresponding authors:
Shahrzad Kharabian Masouleh.
Brain and Behaviour (INM-7)
Institute for Neuroscience and Medicine
Research Centre Jülich
52425 Jülich
E-mail: s.kharabian@fz-juelich.de
22
Dr. Sarah Genon.
Brain and Behaviour (INM-7)
Institute for Neuroscience and Medicine
Research Centre Jülich
52425 Jülich
E-mail: s.genon@fz-juelich.de
29

*Some of the data used in preparation of this article were obtained from the Alzheimer's
Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). As such, the
investigators within the ADNI contributed to the design and implementation of ADNI and/or
provided data but did not participate in analysis or writing of this report. A complete listing of
ADNI investigators can be found at:
http://adni.loni.usc.edu/wp-
content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf
37

Abstract

Linking interindividual differences in psychological phenotype to variations in brain structure is an old dream for psychology and a crucial question for cognitive neurosciences. Yet, replicability of the previously-reported "structural brain behavior" (SBB)-associations has been questioned, recently. Here, we conducted an empirical investigation, assessing replicability of SBB among heathy adults. For a wide range of psychological measures, the replicability of associations with gray matter volume was assessed. Our results revealed that among healthy individuals 1) finding an association between performance at standard psychological tests and brain morphology is relatively unlikely 2) significant associations, found using an exploratory approach, have overestimated effect sizes and 3) can hardly be replicated in an independent sample. After considering factors such as sample size and comparing our findings with more replicable SBB-associations in a clinical cohort and replicable associations between brain structure and non-psychological phenotype, we discuss the potential causes and consequences of these findings.

Introduction:

The early observations of inter-individual variability in human psychological skills and traits have triggered the search for defining their correlating brain characteristics. Studies using in-vivo neuroimaging have provided compelling evidence of a relationship between human skills and traits and brain morphometry that were further influenced by individuals' years of experience, as well as level of expertise. More subtle changes were also shown following new learning/training (Draganski et al., 2004; Taubert et al., 2011), hence further demonstrating dynamic relationships between behavioral performance and brain structural features. Such observations quickly generated a conceptual basis for growing number of studies aiming to map subtle inter-individual differences in observed behavior such as personality traits (Nostro et al., 2017), impulsivity traits (Matsuo et al., 2009) or political orientation (Kanai et al., 2011); to normal variations in brain morphology (for review see (Genon et al., 2018; Kanai and Rees, 2011)). Altogether, these studies created an empirical background supporting the assumption that the morphometry of the brain in humans is related to the wide spectrum of aspects observed in human behavior. Such reports on structural brain behavior (SBB) associations may not only have important implications in psychological sciences and clinical research (Ismaylova et al., 2018; Kim et al., 2015; Luders et al., 2013, 2012; McEwen et al., 2016), but also possibly hold an important key for our understanding of brain functions (Genon et al., 2018) and thus concern many research fields including basic cognitive neuroscience.

Yet, along with the general replication crisis affecting psychological sciences (Button et al., 2013; De Boeck and Jeon, 2018; Open Science Collaboration, 2015), replicability of the previously reported SBB-associations were also questioned recently. In particular, Boekel et al. (2015) in a purely confirmatory replication study, picked on few specific previously

reported SBB-associations. Strikingly, for almost all the findings under scrutiny, they could

not find support for the original results in their replication attempt.

In another study we demonstrated lack of robustness of the pattern of correlations between

cognitive performance and measures of gray matter volume (GMV) in a-priori defined sub-

regions of the dorsal premotor cortex in two samples of healthy adults (Genon et al., 2017). In

particular we found a considerable number of SBB-associations that were counterintuitive in

their directions (i.e., higher performance related to lower gray matter volume). Furthermore,

subsampling revealed that for a given psychological score, negative correlations with GMV

were as likely as positive correlations. Although our study did not primarily aim to address

the scientific qualities of SBB, it revealed, in line with Boekel et al. (2015), that a replication

issue in SBB-associations could seriously be considered. However, ringing the warning bell

of a replication crisis would be premature since these previous studies have approached

replicability questions within very specific contexts and methods and using small sample

sizes (Muhlert and Ridgway, 2016).

In particular, Boekel et al. and Genon et al.'s studies were performed by focusing on a-priori

defined regions-of-interest (ROIs). However, several SBB studies are commonly performed

in groups of dozens of individuals, using an exploratory setting employing a mass-univariate

approach. Thus, the null findings of the two questioning studies could be related to the focus

and averaging of GMV within specific region-of-interests as suggested by (Kanai, 2016) and

discussed in (Genon et al., 2017).

In stark contrast with this argument, in whole-brain mass-univariate exploratory SBB studies,

the multitude of statistical tests that is performed (as the associations are tested for each voxel,

separately) likely yield many false positives. Directly addressing this limitation, several

strategies for multiple comparison correction have been proposed to control the rate of false

positives (Eklund et al., 2016). We could hence assume that the high number of multiple tests

and general low power of neuroimaging studies combined with the flexible analysis choices

(Button et al., 2013; Poldrack et al., 2017; Turner et al., 2018) represent critical factors likely

to lead to the detection of spurious and not replicable associations.

Characterization of spatial consistency of findings across neuroimaging studies is often

performed with meta-analytic approaches, pooling studies investigating similar neuroimaging

markers in relation to a given behavioral function or condition. However, in the case of SBB,

the heterogeneity of the behavioral measures and the large proportion of apriori-ROI analyses

complicate the application of a meta-analytic approach. Illustrating these limitations, previous

meta-analyses have focused on specific brain regions and capitalized on a vast majority of

ROI studies. For example, (Yuan and Raz, 2014) have focused on SBB within the frontal lobe

based on a sample made of approximately 80% of ROI studies. Given these limitations of

meta-analytic approaches for the SBB literature, an empirical evaluation of the replicability of

the findings yielded by an exploratory approach is crucially needed to allow questioning the

replicability of exploratory SBB studies.

Thus in the current study, we empirically examined replicability rates of SBB-association

over a broad range of psychological scores, among heathy adults. In order to avoid the

criticisms raised regarding the low sample size in Boekel et al.'s study, we used an openly

available dataset of a large cohort of healthy participants and assessed replication rate of

SBB-associations using both an exploratory as well as a confirmatory approach. While in the

recent years multivariate methods are frequently recommended to explore the relationship

between brain and behavior (Cremers et al., 2017; Smith and Nichols, 2018), SBB-association

studies using these approaches remain in minority. The mass-univariate approach is still the

main workhorse tool in such studies, not only due to its historical precedence and its wide

integration in common neuroimaging tools, but also possibly owing to more straightforward

interpretability of the detected effects (Smith and Nichols, 2018). The current study, therefore,

focused on the assessment of replicability of SBB-associations using the latter approach.

In particular, we first identified "significant" findings with an exploratory approach based on

mass-univariate analysis, searching for associations of GMV with psychometric variables

across the *whole brain*. Here a linear model was fit between inter-individual variability in the

psychological score and GMV at each voxel. Inference was then made at cluster level, using a

threshold-free cluster enhancement approach (Smith and Nichols, 2009). We then investigated

the reproducibility of these findings, across resampling, by conducting a similar whole-brain

voxel-wise exploratory analysis within 100 randomly generated subsamples of individuals

(discovery samples). Each of these 100 discovery subsamples (of the same size) were

generated by randomly selecting apriori-defined number of individuals (e.g. 70%) from the

original cohort under study. In order to empirically investigate spatial consistency of

significant results from these 100 exploratory analyses, an aggregate map characterizing the

spatial overlap of the significant findings across all discovery samples was generated. This

map denotes the frequency of finding a *significant* association between the behavioral score

and gray matter volume, at each voxel, over 100 analyses and thus provides information about

replicability of "whole brain exploratory SBB-associations" for each behavioral score.

Conceptually, this map gives an estimate of the spatial consistency of the results that one

could expect after re-running 100 times the same SBB study across similar samples.

Additionally, for each of the 100 exploratory analyses, we assessed the replicability of SBB-

associations using a confirmatory approach (i.e. ROI-based approach). For each of the 100

discovery samples, we generated a demographically-matched test pair sample from the

remaining participants of the main cohort. Average GMV within regions showing significant

SBB-association in the initial exploratory analysis, i.e. ROIs, are calculated among the

demographically-matched independent sample and their association with the same

psychological score was compared between the discovery and matched-replication sub-

samples (see Methods for more details).

Confirmatory replication is commonly used in the literature (Boekel et al., 2015; Genon et al.,

2017; Open Science Collaboration, 2015), nevertheless, there is no single standard defined for

evaluating the replication success. Therefore, here, we assessed the replication rate of SBB,

for three different definitions of successful replication in the confirmatory analyses: 1-

Successful replication of the direction of association, only; 2- Detection of significant ($p <$

0.05) association in the same direction as the exploratory results; While the first definition is

arguably too lenient and may result in many very small correlation coefficients defined as

successful replication, it is frequently used as a qualitative measure of replication and may be

used to characterize the possible inconsistency of the direction of associations (that was

observed in our previous study (Genon et al., 2017)). In addition it could be used as a

complement for the possible limitation of the second definition, namely the possibility of

declaring many replications that fell just short of the bright-line of $p< 0.05$ as failed

replication. 3- lastly, in line with previous studies and the reproducibility literature, we

included the Bayes Factors (BF) to quantify evidence that the replication sample provided in

favor of existence or absence of association in the same direction than in the discovery

subsample (Boekel et al., 2015). In other words, when compared to standard p-value

methodology, here hypothesis testing using BF enables quantification of the evidence in favor

of the null hypothesis, i.e. evidence for the absence of a correlation; see Methods for more

details.

If the replication issue of SBB associations can be objectively evidenced, this naturally opens

the questions of the accounting factors. Here, we considered proximal explanatory factors, in

particular at the measurements and analysis level, but also in relation to the object level, that

is, in relation to the nature itself of variations in brain structure and psychometric scores in

healthy individuals. One main proximal factor that is almost systematically blamed is small

sample size. In line with replication studies in other fields (e.g. (Cremers, Wager, & Yarkoni,

2017; Turner, Paul, Miller, & Barbey, 2018)), we thus here investigated the influence of

sample size and replication power on the reproducibility of SBB-associations. More

specifically for every phenotypic score under study we repeated both whole brain exploratory

and ROI-based confirmatory replication analyses using three sample sizes (see Methods for

more details) to assess how sample size influences replication rate of SBB. Furthermore, for

the successfully replicated effects, we also investigated existence of a positive relationship

between the effect size of exploratory and confirmatory analyses.

Finally, in order to promote discussion on the underlying reality which is aimed to be

captured by SBB in the framework of the psychology of individual differences, we included

as benchmarks non-psychological phenotypical measures, i.e. age and body-mass-index

(BMI), and extended our analysis to a clinical sample, where SBB-associations are expected

to enjoy higher biological validity. For this purpose, a subsample of patients drawn from

Alzheimer's Disease Neuroimaging Initiative (ADNI) database were selected, in which

replicability of structural associations of immediate-recall score from Rey auditory verbal

learning task (RAVLT) (Schmidt, 1996) was assessed (see Methods). Due to availability of

the same score within the healthy cohort, this later analysis is used as a "conceptual"

benchmark.

Results:

A total of 10800 exploratory whole brain SBB associations (each with 1000 permutations)

were tested to empirically identify the replicability of the associations of 36 psychological

scores with GMV over 100 splits in independent matched subsamples, at three pre-defined

sample sizes, within the *healthy* cohort; see Supplementary Table 1, for total number of

participants with available score for each of the psychological scores.

Altogether, in contrast to GMV-associations with age and BMI, significant SBB-associations

were highly unlikely. For the majority of the tested psychological variables no significant

association with GMV were found in more than 90% of the whole brain analyses.

SBB-associations among the healthy population:

Replicability of "whole brain exploratory SBB-associations":

Age and BMI structural associations: Voxel-wise associations of age and BMI with GMV, as

suggested by previous studies (Fjell et al., 2014; Kharabian Masouleh et al., 2016; Salat et al.,

2004; Willette and Kapogiannis, 2014), were widespread and strong.

Despite using more stringent thresholds, compared to the threshold used for the psychological

scores (see Methods), for almost all subsamples, we found highly consistent widespread

negative associations of GMV with age. See figure 1A for aggregate maps of spatial overlap

of exploratory findings and density plots, summarizing distribution of "frequency of

significant findings" within each map.

When decreasing the sample size of the discovery cohort, the spatial overlap of significant

findings over 100 splits decreased. More specifically, for the discovery sample of 326

subjects, more than half of the significant voxels were consistently found as being significant

in beyond 90% of the whole-brain exploratory analyses (i.e. high level of spatial consistency

of significant findings). As the size of the subsamples decreased, the shape of the distribution

also changed, and the median of the density plots fell around 50% and even 10% for samples

consisting of 232 and 138 individuals, respectively.

Similar results, though with much lower percent of consistently overlapping voxels, were

seen for negative associations of BMI with GMV. The density plots and the spatial maps of

Figure 1B show that for the larger samples (consisting of 326 and 232 subjects) few voxels

were consistently found in "all" (100%) subsamples as having significant negative

association with BMI. For the smaller samples (with 138 participants) the maximum

replicable association was found in 93% of the splits and 4 out of 100 exploratory analyses

did not result in any significant clusters (Table 1). Additionally, as Figure 2B shows, the

majority of significant voxels had a replicability bellow 50%.

These results highlight the influence of sample size on the replicability (frequency of overlap)

of whole-brain significant associations, even for age and BMI, for which we expected more

stable associations with morphological properties of the brain.

Structural associations of the psychological scores: In contrast, for most of the psychological

scores, only few of the 100 discovery subsamples yielded significant clusters. Table 1 and

supplementary Table 2 show the number of splits for which the exploratory whole-brain SBB-

analysis resulted in *at least one* significant positively or negatively associated cluster for each

score. These results reveal that finding significant SBB-associations using the exploratory

approach in healthy individuals is highly *unlikely* for most of the psychological variables.

Furthermore, the significant findings were spatially very diverse, that is, spatially overlapping

findings were very rare.

We here retained for further analyses the three psychological scores for which the discovery

samples most frequently resulted in at least one significantly associated cluster. These three

scores were the Perceptual reasoning score of WASI (Wechsler, 1999), the number of correct

responses in word-context test and the interference time in the color-word interference task.

For example, for the discovery samples of 326 adults, in 83 out of 100 randomly generated

discovery samples, at least one cluster (not necessarily overlapping) showed a significant

positive association between perceptual reasoning and GMV (Table 1)). Of note, these more

frequently found associations were in the direction linking better task performance with

higher GMV.

Yet again, in line with our observations for BMI associations, the probability of finding at

least one significant cluster tend to decrease in smaller discovery samples (see Table 1).

Likewise, as the discovery sample size decreased, the maximum rate of spatial overlap, as

denoted by the height of the density plots, decreased (see Figure 1C-F). The width of these

plots show that the majority (> 50%) of the significant voxels spatially overlapped only in less

than 10% of the discovery samples. In the same line, the variability depicted by the spatial

maps highlight that many voxels are found as significant only in one out of 100 analyses.

These results highlight that finding a significant association between normal variations on

behavioral scores and voxel-wise measures of GMV among healthy individuals is highly

unlikely, for most of the tested domains. Furthermore, they underscore the extent of spatial

inconsistency and the *poor replicability* of the significant SBB-associations from *exploratory*

analyses.

--------Table 1 ---------

-------figure1---------

Confirmatory ROI-based SBB-replicability:

Age and BMI effects: Irrespective of the size of the test subsamples and definition used to

identify "successful" replication (see Methods), for all ROIs negative age-GMV associations

were "successfully" replicated in the matched test samples. Unlike the perfect replication of

age-associations, replication rate of BMI effects depended highly on the test sample size and

the criteria used to characterize "successful" replication. Over all three tested sample sizes, in

more than 90% of the a-priori defined ROIs, BMI associations were found to be in the same

"direction" in the discovery and test samples (i.e. replicated based on "sign" criteria). The

examination of replicated findings based on "statistical significance" revealed replicated

effects in more than 57% of ROIs. This rate of ROI-based replicability increased from ~57%

to 75%, as the test sample size increased from 140 to 328 individuals (see figure 2).

Furthermore, as the dark blue segments in the outer layers of figure 2 indicates, Bayesian

hypothesis testing revealed moderate-to-strong evidence for H1 in more than 30% of the

ROIs.

286   **--------figure2 ---------**

Psychological variables: Figure 2 also illustrates the replicability rates of structural

associations of the top three psychological measures from the whole brain analyses (the

perceptual reasoning score of WASI, the number of correct responses in word-context test and

the interference time in the color-word interference task).

Despite the structural associations of perceptual reasoning score being in the same direction

(positive SBB-association), for the majority of the ROIs (>85%), less than 31% of all ROIs

showed replicated effects based on "statistical significance" criterion. Finally, less than 4% of

the ROIs were identified as "successfully replicated" based on the Bayes factors. (Figure 2).

For the three tested samples sizes, associations of the word-context task were in the same

direction (positive SBB-association) in the discovery and test pairs in ~75% of ROIs.

Nevertheless, again, the rate of statistically "significantly"-replicated ROIs ranged between 17

to 26%. Furthermore, even less than 8% of all ROIs showed replicated effects based on the

Bayes factors (moderate-to-strong evidence for H1) (Figure 2).

Finally, negative correlations between interference time of the color-word interference task

and average GMV were depicted in ~70 % of the ROIs, but significant-replication was found

in only 11% to 17% of all ROIs, for the three test sample sizes. Along the same line,

replication based on the Bayes factors was below 5% (Figure 2E).

In general, these results show the span of replicability of structural associations from highly

replicable age-effects to very poorly replicable psychological associations. They also

highlight the influence of the sample size, as well as the criteria that is used to define

successful replication on the rate of replicability of SBB-effects in independent samples.

Effect size in the discovery sample and its link with effect size of the test sample and actual

replication:

Figure 3 plots discovery versus replication effect size (i.e. correlation coefficient) for each

ROI and for three test sample sizes. Focusing on by-"sign" replicated ROIs (blue), for the

three psychological scores (perceptual reasoning, word-context and CWI) revealed that the

discovery samples resulted in overall larger effects (magnitude) compared to the test samples.

Indeed, the marginal distributions are centered around smaller correlation coefficients in the

y-dimension (test sample) compared to the x-axis (discovery samples). Furthermore, for these

by-"sign" replicated ROIs, there was no positive relationship between the effect sizes of the

behavioral associations in the discovery and test samples (blue lines in each subplot).

For BMI and age, however, the effect sizes of the discovery and test pairs were generally

positively correlated, suggesting that the ROIs with greater negative structural association

with BMI (or age) in the discovery sample, also tended to show stronger negative associations

within the matched test sample.

To investigate if the replication power, estimated using the correlation coefficient within the

discovery samples, was linked to a higher probability of *actual* replication in the test samples,

the ROIs were grouped into replicated and not-replicated, based on the "statistical

significance" criterion. While the estimations of statistical power were generally higher

among the replicated compared to not-replicated ROIs for BMI associations (p-value of the

Mann-Whitney U tests $< 10^{-5}$), for structural associations of the psychological scores, this was

not the case. Strikingly, for the structural associations of perceptual reasoning, over all sample

sizes, the significantly replicated ROIs tended to have lower estimated power compared to the

ROIs that actually were not-replicated (p-value of the Mann-Whitney U tests $< 10^{-5}$). These

unexpected findings highlight the unreliable aspect of effect size estimations of SBB-

associations within the discovery samples among healthy individuals. They also demonstrate

that these inflated effect sizes result in flawed and thus uninformative estimated statistical

power.

335　　　　　　　　　　　　　　　　**--------figure3 ---------**

Structural associations of total immediate recall score in ADNI cohort:

Replicability of "whole brain exploratory associations":

Within the sample of patients from ADNI-cohort, 84 out of the 100 whole-brain exploratory

analyses resulted in *at least one* significant cluster showing a positive association between the

immediate-recall score and GMV. In the healthy population, however, the same score resulted

in a significant cluster in only less than 10% of exploratory analyses, for any of the three

discovery sample sizes (supplementary Table 2 and supplementary Figure 1).

As could be seen in the spatial maps of Figure 4, significant associations in the ADNI cohort

were found across several brain regions including the bilateral lateral and medial temporal

lobe, the lateral occipital cortex, the precuneus, the superior parietal lobule, the orbitofrontal

cortex and the thalamus. Although most of the significant voxels were found by less than 10%

of the splits, some voxels in the bilateral hippocampus were found to be significantly

associated with the recall score in more than 70% of the subsamples (peak of spatial overlap;

see Figure 4A, B).

Confirmatory ROI-based SBB-replicability:

Figure 4D shows the rates of "successful replication" of associations between the immediate-

recall score and GMV within each ROI in the independent, matched-samples. As the most

inner layer shows, in more than 94% of ROIs, GMV correlated positively with the recall score

in the test subsamples, corroborating the "sign" of the association in the paired-discovery

samples. These correlations were significant in 72% of all ROIs. Furthermore, in more than

50% of all ROIs the correlations in the test sample supported, at least moderately, the link

between higher GMV and higher recall score (using the Bayes factors).

Association between discovery and replication effect size:

The marginal histograms in Figure 4C suggest that overall the **<u>correlations</u>** in the discovery

samples are slightly **stronger** than the **<u>correlation</u>**s in the paired replication samples. When

looking at the ROIs that were successfully replicated (by-sign), there was a positive

association between the discovery and replication effect size (spearman's rho = 0.38, p-value

$< 10^{-11}$) .

Finally, the median replication power was higher among "significantly replicated" ROIs,

compared to not replicated (defined using "statistical significance criterion") ROIs (p-value of

the mann-whiteney U test $< 10^{-3}$). These results showed the superior, yet not perfect,

replicability of SBB-associations within the clinical population (see supplementary Figure 2

for structural associations of immediate recall within healthy cohort). The observed somewhat

robustness of the findings in ADNI suggest that, when the population under study shows clear

variations in both brain structural markers and psychological measurements, such as the

patient group in ADNI cohort, the associations between brain structure and psychological

performance could be relatively reliably characterized. Nevertheless, again, the occurrence of

not-replicated results highlight the importance of confirmatory analyses for a robust

characterization of brain-behavior associations.

377   **--------figure4 ---------**

Discussion:

Our empirical investigation of the replicability of SBB in healthy adults showed that

significant associations between psychological phenotype and GMV are not frequent when

probing a range of psychometric variables with an exploratory approach. Where significant

associations were found, these associations showed a poor replicability.

In the following, we first discussed implications of the very low rate of significant findings

revealed by the exploratory approach. We then discussed the possible causes of the observed

spatial variability of SBB-associations. Those pattern of findings are then compared with the

pattern observed in the clinical cohort. Finally, in line with the replication literature in

psychological sciences and neurosciences (Button et al., 2013; Poldrack et al., 2017; Turner et

al., 2018), we devoted our last section to sample size and power issues in SBB studies in

healthy adults and proposed some recommendations.

Infrequent significant SBB associations in healthy individuals: Importance of reporting null

findings

According to the scientific literature, associations between psychological phenotype

(cognitive performance and psychological trait) and local brain structure are not uncommon

(Kanai and Rees, 2011). However, in our exploratory analyses, when looking at a range of

psychological variables, significant associations with GMV were very rare. It is worth noting

that here by having a-priori fixed analysis design and inference routines, we aimed to avoid

"fishing" for significant findings (Gelman and Loken, 2014). Flexible designs and flexible

analyses routines (Simmons et al., 2011) as well as p-hacking (John et al., 2012) are

considered as inappropriate but frequent research practices (Poldrack et al., 2017). Based on

our findings of infrequent significant SBB-associations, we could assume that flexible

analyses routines, p-hacking and most importantly *publication bias* (Dwan et al., 2013) have

contributed to the high number of significant SBB-reports in the literature.

When considering potential impacts of biased SBB-reports on our confidence of

psychological measures, as well as our conception and apprehension of brain-behavior

relationships and psychological interindividual differences, we would strongly argue for null

findings reports. Such reports would contribute to a more accurate and balanced apprehension

of associations between differences in psychological phenotype and brain morphometric

features, but it would also help to progressively disentangle factors that mediate or modulate

the relationship between brain structure and behavioral outcomes.

Poor spatial overlap of SBB across resampling: possible causes and recommendations

In addition to the low likelihood of finding "any" significant SBB-association using the

exploratory approach, clusters that do survive the significance thresholding did not often

overlap in different subsamples. Furthermore, the probability of spatial overlap further

dropped as the number of participants in the subsamples decreased (Figure 1). Putting this

finding in light of the literature brings two main hypotheses.

First, from the conceptual level, we could hypothesize that the pattern of correlation between

a psychological measure is by nature spatially diffuse at the brain level. Psychological

measures aim to conceptually articulate *behavioral functions and processes*, thus, in most

cases, they have not been developed to identify specific localized *brain functions*. Following

this philosophical segregation between psychological sciences and neurosciences, it is now

widely acknowledged that there is no one-to-one mapping between behavioral functions and

brain regions (Anderson, 2015; Genon et al., 2018; Pessoa, 2014). Instead, mapping a

psychological concept to brain features usually result in a diffuse brain spatial pattern with

small effect sizes (Bressler, 1995; Poldrack, 2010; Tononi et al., 1998). From this axiom, we

can expect that several studies conducted in small samples (specifically after rigorous

corrections for multiple comparisons) are likely to each capture a partial and minor aspect of

the whole true association pattern, resulting in a poor replication rate for each individual study

(i.e. high type II error).

Alternatively, a more parsimonious hypothesis is a methodological one questioning the truth

or validity of the found significant associations hence considering them as spurious (i.e. type I

error). Psychological and MRI measurements are both relatively indirect estimations of

respectively, behavioral features and brain structural features and thus are susceptible to

noise. Correlations in small samples in the presence of noise for both type of variables is

likely to produce spurious significant results (Loken and Gelman, 2017) by fitting a

correlation or regression between random noise in both variables.

Thus, the pattern of poor spatial consistency of SBB findings could result either from factors

at the object of study level, i.e. the relationship between brain and behavior, or, from factors

at the measurement and analysis level. While the latter hypothesis is more parsimonious, one

argument for the former hypothesis comes from the relatively substantial replications by-sign

observed in our confirmatory analyses, of three top behavioral scores (see figure 2). If the

significant SBB findings would be purely driven by noise in the data, we would expect them

to show purely random signs across resampling, which was not the case (but also see

Supplementary figure S2 for example of scores with lower replicability and higher

inconsistent associations across resampling). Therefore, it is actually likely that both

hypotheses hold true and that the spatial variability of significant SBB findings result from

both, factors at the analyses levels and factors at the object level, potentially interacting

together.

It is worth noting that similar complexity and uncertainty have been described for task-based

functional associations studies (Cremers et al., 2017; Turner et al., 2018). In particular,

Cremers et al. (2017) using simulated and empirical data demonstrated that task-based

functional activations have a generally weak and diffuse pattern. Therefore, these authors

concluded that most whole-brain analyses in small samples, specifically when combined with

stringent correction for multiple comparison, to control the false positive rates, would most

likely frequently overlook global meaningful effects and depict results with poor replicability

(type II error). Relatedly, in the present study, higher spatial extent and lower consistency of

significant findings in smaller samples in Figure 1, also suggest higher number of spurious

associations (type I error) in smaller samples (due to winners curse (Button et al., 2013;

Forstmeier and Schielzeth, 2011)) than in larger samples.

These factors, added to the complexity of human behavior, renders the objective of capturing

covariations with psychometric variables in brain structure *locally* particularly challenging.

For that reason, in exploratory studies whose aim is to identify brain structural features

correlating with a specific (set of) psychological variable(s), a multivariate approach could be

advised (Habeck and Stern, 2010; McIntosh and Mišić, 2013). As mentioned earlier, like all

methods, multivariate analyses have their own limitations: in particular, the ensuing difficulty

of interpretability of the revealed pattern. While some authors argue either for one or the other

approach, the use of these approaches are far from being mutually exclusive (Moeller and

Habeck, 2006). Combining both approaches in small datasets indeed revealed that the results

of the univariate approach reflect the "tip of the iceberg" of the behavior's brain correlates,

whose spatial extent are more comprehensively captured with the multivariate analysis, but

interpretability is facilitated by the use of univariate analyses; e.g. (Genon et al., 2016, 2014).

Thus, to partially address the previously described concerns of small and spatially diffuse

effects at the brain level in exploratory whole-brain-behavior study, we here recommend for

the future studies to combine a univariate and a multivariate approach. Although it does not

provide any protection against the influence of noise that may affect both approaches, this

solution may help to reduce the false negatives.

Confirmatory replication of exploratory SBB findings: importance of out of sample

replication

ROI-based analysis further highlighted that significant associations, which have been

discovered when starting with a psychological measure and searching within the whole brain

for a significant association (i.e. "evidenced in an exploratory study"), show poor replicability

(using significance and Bayes factor criteria, but also using a similar sign criterion for most

psychometric scores; For example, see Supplementary Figures S1 and S2.) in a confirmatory

ROI-based study (in line with what was previously shown by Boekel et al. (2015)). These

findings thus call for a general acknowledgment of the uncertainty and fragility of exploratory

findings and the need for *out of sample* confirmatory replications to provide evidence about

the robustness of the reported effects (Ioannidis, 2018; Tukey, 1980).

Further factors influencing replicability of SBB-findings: power of replication and object of

study

Another clear finding of our study is the overestimation of the effect size in the exploratory

approach (Kriegeskorte et al., 2010), specifically in smaller samples (see marginal

distributions of the x- and y-axis in Figure 3). For the majority of the psychological scores, in

the ROI-based approach, we failed to find a clear association between effect size in the

discovery and replication samples. Instead, we observed a rather high estimated statistical

power for replication (due to an inflated effect size estimation (Ioannidis, 2008)), despite very

low actual rate of replicated effects in the independent samples. These findings are

particularly important when considering the current research context, in which power analyses

are encouraged to justify the allocation of financial and human investment in specific future

researches. Prospective studies with power analyses are frequently proposed, where power is

computed based on the findings from previous exploratory analyses in a small sample (Albers

and Lakens, 2018a). An inflated effect size estimation from the exploratory study results in an

unreliable high power, which in turn lead to confidence in prospective studies to find relevant

findings and hence in the allocation and possibly waste of (frequently public) resources

(Albers and Lakens, 2018b; Poldrack et al., 2017). Nevertheless, this provocative conclusion

does not imply that SBB studies should be banished to hell. Our conclusion here mainly

concerns the study of association between variations at *standard psychological measures* and

variations in *measures of gray matter* in "*small*" *samples* of *healthy individuals*. Our results

further show that different types of SBB exploratory studies should not be epistemologically

all put in the same pot.

In support for this argument, in ADNI sample, despite the additional confounding effect of

different scanners and/or scanning parameters due to the multi-site nature of the cohort,

associations between immediate-recall score and GMV were relatively stable. Compared to

associations of the same measure of verbal learning performance within the healthy

population (see supplementary Figure 1), these results highlight the superior reliability of

SBB-associations that are defined in a clinical context. These findings have important

conceptual implications. From an epistemological and conceptual point of view, our

comparative investigation suggests that the object of study matters in the replicability of SBB.

Searching for correlation between variations in cognitive performance and GMV in healthy

adults, on one hand, and in neurodegenerative patients, on the other hand, appear as two

different objects of study, with different replicability rates. While several SBB results in

healthy population are likely to be spurious (see supplementary Table 2), it seems that SBB in

clinical population are more likely to capture true relationships.

Thus, maybe the conceptual objective itself should be questioned: should we expect the

association between normal psychological phenotype, in particular cognitive performance, in

healthy population to be substantially driven by local brain macrostructure morphology?

Brain structure can certainly not be questioned as the primary substrates of behavior and

more than a century of lesion studies recalls this primary principle to our attention (Broca,

1865; Scoville and Milner, 1957), but this does not imply that "normal" variations at standard

psychological tests can be related to variations in markers of local brain macrostructure. Our

results suggest that reliable answer to this important question requires substantially big

samples (bigger than those used here) and independent replications.

Further recommendation: Large sample sizes are important both for exploratory as well as

replication analyses

The sample size and related power issues hold a central position in the current discussions of

the replication crisis in behavioral sciences, as well as in neuroimaging studies (Button et al.,

2013; Ioannidis, 2005; Lilienfeld, 2017; Munafò et al., 2017; Open Science Collaboration,

2015). Higher power is defined as increased probability of finding effects that are genuinely

true. Furthermore, high power experiments have higher positive predictive values (PPV) of

the claimed effects (i.e. probability that the claimed effect reflects a true effect). They also

result in less exaggerated effects sizes when a true effect is discovered (Button et al., 2013).

As such, in the discovery sample, by increasing the sample size, the correlation coefficients

get closer to their real value and their PPV increases. However, in the current study, as the

number of participants in the main sample is limited, the size of the discovery and their

matched replication samples are dependent on each other. Therefore, for each behavioral

measure, larger discovery samples have smaller replication counterparts. These smaller

replication samples have in turn lower power to find the true effects and have lower PPV.

However, in splits with larger replication samples, as the discovery sample gets smaller, apart

from the lower PPV, the estimated correlation coefficients are possibly more exaggerated

(e.g. due to winner's curse) (Cremers et al., 2017) and thus the power of the replication would

be over-estimated. This is a limitation which complicates the interpretation of the relationship

between the calculated replication power and the actual rate of replicability of associations in

the present study. We hoped that the use of a large cohort of healthy individuals as our main

cohort would result in large enough discovery and test cohorts and thus minimize the impact

of above-mentioned limitation. However, large discrepancies between the rate of "significant"

within-split replicability and the a-priori estimated replication power, as we observed in the

ROI-based confirmatory analyses, confirms an exaggerated power estimation in most of our

analyses and thus highlights the insufficiency of the size of the discovery and replication

samples.

Thus overall, these findings suggest that samples consisting of ~200-300 participants have in

reality still low power to identify reliable SBB-associations among healthy participants.

However, the sample size of SBB studies is usually substantially smaller. Figure 5 depicts the

distribution of sample sizes (log-scale) of published studies examining GMV in human

participants with the standard voxel-based morphometry approach across previous years

(BrainMap data (Vanasse et al., 2018)). SBB studies in healthy adults also fall under this

general trend. Based on our current work, we would argue that the probability of finding

spurious or inconclusive results and exaggerated effect size estimations in these studies is thus

quite high (Albers and Lakens, 2018b; Schönbrodt and Perugini, 2013; Yarkoni, 2009).

In addition, to underscore the importance of the sample size, our analyses and results further

show that the size of the replication sample also matters when examining the replicability of a

previous SBB findings. This is an obvious factor that has been frequently neglected in the

discussions about replication crisis. Yet, while many replication studies straightforwardly

blame the sample size of the original studies, it is important to keep in mind that a replication

failure might also come from a too small sample size of the replication study (Muhlert and

Ridgway, 2016).

575   --------figure5 ---------

Limitations:

When interpreting our results, it should be noted that, in order to keep large sample sizes for

the exploratory replication analyses, the discovery subsamples were not necessarily designed

to be independent from each other. Considering this limitation, the poor spatial consistency of

the whole brain exploratory associations that we observed for almost all the behavioral scores

is hence even more alarming. As discussed earlier, another indirect limitation of the limited

size of the selected cohort is the dependence between the size of the discovery and their

matched replication sub-samples. This limitation prevents us to state strong conclusions about

the relationship between the calculated replication power and the actual rate of replicability.

Overall, these acknowledged limitations raise the need for even larger sample sizes for such

investigations. Recent advancements through data collection from much larger number of

participants, such as UK-biobank (Miller et al., 2016) are promising opportunities for

overcoming these limitations in future replication studies.

Moreover, the generalizability of our results are partly limited to our methodological choices

such as the computation of volumetric markers of brain structure (as opposed to surface-based

markers), the size of the smoothing kernel, and the use of a priori-defined ROIs in the

replication sample. Future studies should therefore investigate to which extend our

replicability rates are reproduced with different data preprocessing pipelines and analyses

approaches.

Summary and conclusions

Overall, our work and review of the recent and concomitant replication literature in related

fields demonstrate that several improvements could be recommended to get more accurate

insight on the relationship between psychological phenotype and brain structure and to

progressively answer open questions. Importantly, our recommendations and suggestions

concern different levels of SBB researches: the dataset level, the analyses level, as well as at

the post-publication and replication level.

At the dataset level, our study pointed out the need for big data samples to identify robust

associations between psychological variables and brain structure, with sample size of at least

several hundreds of participants. It should be acknowledged that this conclusion is easier to

achieve than to implement in research practice. Nevertheless, large scale cohort datasets from

healthy adult populations, such as eNKI used in the current study, human connectome project

(HCP) (Van Essen et al., 2013) and UK-biobank (Miller et al., 2016) are now openly

available, hence facilitating endeavor in that direction.

At the analysis level, we recommend the combined use of multivariate analyses, for

comprehensive assessment of the spatial extent of associations and, univariate analyses, to

facilitate interpretability, when studying brain structural correlates of psychological measures.

Furthermore, we emphasis on the importance of *independent* confirmatory replications to

provide evidence about the robustness of the effects.

Finally, *at the post-analysis level*, we concluded from our observations that publication of null

findings should be more encouraged. In addition to directly shaping a more objective picture

of SBB-associations, these null-reports could contribute to new quantitative approaches. In

particular, meta-analyses of published literature (Vanasse et al., 2018) would clearly benefit

from such unbiased reports of null findings.

Sharing raw data would undoubtedly improve the problem of low statistical power, but if not

possible, sharing the unthresholded statistical maps (e.g. through platforms such as

Neurovault (Gorgolewski et al., 2015)) could also be a significant scientific contribution. In

addition to directly contribute to our understanding of brain-behavior relationship, such

efforts would open up new possibilities for estimating the correct size and extent of effects by

integrating unthresholded statistical maps in the estimation of the effects sizes throughout the

brain. Thus, we could hope that sharing initiatives will also contribute indirectly to more valid

and insightful SBB studies in the remote future and hence to a better allocation of resources.

Methods:

Participants:

Healthy adults' data were selected from the enhanced NKI (eNKI) Rockland cohort (Nooner et al., 2012). Data collection received ethics approval through both the Nathan Klein Institute and Montclair State University. Written informed consent was obtained from all participants. We focused only on participants for which good quality T1-weighted scans was available along with timewise-corresponding psychological data. Exclusion criteria consisted of alcohol or substance dependence or abuse (current or past), psychiatric illnesses (eg. Schizophrenia) and current depression (major or bipolar). Furthermore, we excluded participants with missing information on important confounders (age, gender, education) or bad quality of structural scans after pre-processing, resulting in a total sample of 466 healthy participants (age: 48 ± 19, 153 male).

Replicability of SBB-associations within the clinical sample was investigated using a subsample drawn from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database, which was launched in 2003 as a public–private partnership and led by Principal Investigator Michael W. Weiner. The primary goal of ADNI has been to test whether serial magnetic resonance imaging (MRI), positron emission tomography (PET), other biological markers, and clinical and neuropsychological assessment can be combined to measure the progression of mild cognitive impairment (MCI) and early Alzheimer's disease (AD). For up-to-date information, see www.adni-info.org.

We used the baseline measurements from 371 patients (age : 71 ± 7, 200 male ; 47 with significant memory complaint, 177 early MCI, 85 late MCI and 62 AD patients (defined based on ADNI diagnostic criteria, see http://adni.loni.usc.edu/wp-content/themes/freshnews-dev-v2/documents/clinical/ADNI-2_Protocol.pdf), in whom anatomical brain scans had been acquired in a 3Tesla scanner (from 39 different sites).

Phenotypical measurements:

Non-psychological measurements:

Age and body mass index (BMI) are highly reliably assessed and their association with brain

morphology has been frequently examined in previous studies on healthy adults (Fjell et al.,

2014; Kharabian Masouleh et al., 2016; Salat et al., 2004; Willette and Kapogiannis, 2014).

Accordingly, they served here as the initial benchmarks among which SBB framework was

tested in healthy individuals. In order to avoid large clusters that simultaneously cover several

cortical and subcortical regions, we focused on local peaks of associations by increasing the

voxel-level t-threshold of the statistical maps. The modified voxel-level t-threshold was set to

8 and 3, for defining age- and BMI-associated clusters, respectively. These *arbitrary*

thresholds were chosen such that the very large clusters would divide into smaller ones, while

still retaining the general spatial pattern of the significant regions.

Psychological measurements:

The psychological measurements consisted in standard psychometrics and

neuropsychological tests. The testing included: the attention network task (ANT) probing

attention sub-functions (Fan et al., 2002), the Delis-Kaplan testing battery assessing different

aspects of executive functions (Delis et al., 2001) (including trail-making test, color-word

interference task, verbal fluency, 20 questions, proverbs and word-context task) , the Rey

auditory verbal learning task (RAVLT) (Schmidt, 1996) assessing verbal memory

performance, as well as the WASI-II intelligence test (Wechsler, 1999). Psychological

phenotyping also included anxiety (state and trait) (Spielberger et al., 1970) and personality

questionnaires (McCrae and Costa, 2004) in the eNKI cohort. For each test, we used several

commonly derived sub-scores to assess the replicability of their structural associations. For

each psychological measure, participants whose performance deviated more than 3 standard

deviation (SD) from mean of the whole sample were considered as outliers and thus were

excluded from further analysis (See supplementary Table 1).

The list-learning task is a common measure of verbal learning performance and has been

implemented using the same standard tool (RAVLT) in both the eNKI and the ADNI cohort.

Previous studies have shown that the immediate-recall score (sum of recalled items over the

first 5 trials) could be reliably predicted from whole brain MRIs in AD patients (Moradi et al.,

2017). Since this score is a standard measure commonly used in healthy and clinical dataset

and its relations to brain structure in clinical data has been previously suggested, in the current

work we performed SBB with this score in the ADNI cohort as a "conceptual benchmark".

MRI acquisition and preprocessing:

The imaging data of the eNKI cohort were all acquired using a single scanner (Siemens

Magnetom TrioTim, 3.0 T). T1-weighted images were obtained using a MPRAGE sequence

(TR = 1900 ms; TE = 2.52 ms; voxel size = 1 mm isotropic).

ADNI, on the other hand, is a multisite dataset. Here we selected a subset of this data, which

has been acquired in a 3.0 T scanner (baseline measurements from ADNI2 and ADNI GO

cohort) from 39 different sites; see http://adni.loni.usc.edu/methods/documents/ for more

information.

Both datasets were preprocessed using the CAT12 toolbox (Gaser and Dahnke, 2016).

Briefly, each participant's T1-weighted scan was corrected for bias-field inhomogeneities,

then segmented into gray matter (GM), white matter (WM), and cerebrospinal fluid (CSF)

(Ashburner and Friston, 2005). The segmentation process was further extended for accounting

for partial volume effects (Tohka et al., 2004) by applying adaptive maximum a posteriori

estimations (Rajapakse et al., 1997). The gray matter segments were then spatially normalized

into standard (MNI) space using Dartel algorithm (Ashburner, 2007) and further modulated.

The modulation was performed by scaling the normalized gray matter segments for the non-

linear transformations (only) applied at the normalization step. While this procedure ignores

the volume changes due to affine transformation, it allows preserving information about

individual differences in *local* gray matter volume. In other words, it re-introduces individual

differences in local gray matter volume removed in the process of inter-subject registration

and normalization. Finally modulated gray matter images were smoothed with an isotropic

gaussian kernel of 8 mm (full-width-half-maximum).

Statistical analysis:

SBB-associations are commonly derived in an exploratory setting using a mass-univariate

approach, in which a linear model is used to fit interindividual variability in the psychological

score to GMV at each voxel. Inference is then usually made at cluster level, in which groups

of adjacent voxels that support the link between GMV and the tested score are clustered

together.

Replicability of thus-defined associations could be assessed by conducting a similar whole-

brain voxel-wise exploratory analysis in another sample of individuals and comparing the

spatial location of the significant findings that survive multiple comparison correction,

between the two samples. Alternatively, replicability could be assessed, using a confirmatory

approach, in which only regions showing significant SBB-association in the initial

exploratory analysis, i.e. regions of interest (ROIs), are considered for testing the existence of

the association between brain structure and the same psychological score in an independent

sample. The latter procedure commonly focuses on a summary measure of GMV within each

ROI and tests for existence of the SBB-association in the direction suggested by the initial

exploratory analysis. Thus this approach circumvents the need for multiple comparison

correction and therefore increases the power of replication.

Here we assessed replicability of associations between each behavioral measure and gray

mater structure, using both approaches: the whole brain replication approach and the ROI

replication approach, which are explained in details in the following sections.

Replicability of whole brain exploratory SBB-associations:

Whole-brain GLM analyses: 100 random subsamples (of same size) were drawn from the main cohort (eNKI or ADNI). Hereafter, each of these subsamples is called a "discovery sample". In each of these samples, SBB-associations were identified using the voxel-wise exploratory approach after controlling for confounders. This was done by using the general linear model (GLM) as implemented in the "randomise" tool (https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/Randomise), with 1000 permutations. Age, sex and education were modeled as confounders in the eNKI data. As the ADNI dataset is a multi-site study, we further added site and disease category as dummy-coded confounders to GLMs for the analyses in that dataset. Inference was then made using threshold-free cluster enhancement (TFCE) (Smith and Nichols, 2009), which unlike other cluster-based thresholding approaches does not require an arbitrary a-priori cluster forming threshold. Significance was set at $P < 0.05$ (extent threshold of 100 voxels).

Spatial consistency maps and density plots: To quantify the spatial overlap of significant SBB associations over 100 subsamples, spatial consistency maps were generated. To do so, the binarized maps of all clusters that showed significant association in the same direction between each psychological score and GMV were generated (i.e. voxels belonging to a significant cluster get the value "1" and all other voxels were labeled "0") and added over all 100 subsamples. These aggregate maps denote the frequency of finding a *significant* association between the behavioral score and GMV, at each voxel. Accordingly, a voxel with value of 10 in the aggregate map has been found to be significantly associated with the phenotypical score in 10 out of 100 subsamples. Density plots were also generated to represent the distribution of values within each such map, i.e. the distribution of "frequency of significant finding". Hence, the spatial voxel-wise "significance overlap maps" as well as density plots of the distribution of values within each map give indications of the replicability of "whole brain exploratory SBB-associations" for each psychological score.

Replicability of SBB-associations using confirmatory ROI-based approach:

ROI-based confirmatory analyses: The replicability of the SBB associations was also

evaluated with the ROI-based confirmatory approach. For each of the 100 discovery

subsamples, an age- and sex-matched "test sample" was generated from the remaining

participants of the main cohort. In the clinical cohort the discovery and test pairs were

additionally matched for "site". In this analysis, for each psychological variable, the

significant clusters from the above-mentioned exploratory approach from every "discovery

sample" were used as a-priori ROIs. Average GMV over all voxels within the ROI was then

calculated for each participant in the respective "discovery" and "test" pair subsamples.

Within each subsample, association between the average GMV and the psychological variable

was assessed using ranked-partial correlation, controlling for confounding factors. The

correlation coefficient was then compared between each discovery and test pair, providing

means to assess "ROI-based SBB replicability" rates for each psychological score.

Accordingly, each ROI was examined only once, to identify if associations between average

GMV in this ROI and the psychological score from the discovery subsample could be

confirmed in the paired test sample. Replicability rates were quantified according to different

indexes (see below) over all ROIs from 100 discovery samples, yielding a percentage of

"successfully replicated" ROIs based on each index.

Indexes of replicability:

Sign: First, we used a lenient definition of replication, in which we compared only the sign of

correlation coefficients of associations within each ROI between the discovery and the

matched-test sample. Accordingly, any effect that was in the same direction in both samples

(even if very close to zero) was defined as a "successful" replication.

Statistical Significance: Another straightforward method for evaluating replication simply

defines statistically significant effects (e.g. p-value < 0.05) that are in the same direction as

the original effects (from the discovery sample) as "successful" replication. This criteria is

consistent with what is commonly used in the psychological sciences to decide whether a

replication attempt "worked" (Open Science Collaboration, 2015). Yet, a key weakness of

this approach is that it treats the threshold ($p < 0.05$) as a bright-line criterion between

replication success and failure. Furthermore, it does not quantify the decisiveness of the

evidence that the data provides for and against the presence of the correlation (Boekel et al.,

2015; Wagenmakers et al., 2015). However, such an estimation can be provided by using the

"Bayes factors".

Bayes Factor: To compare the evidence that the "test subsample" provided for or against the

presence of an association (H1 and H0, respectively), we additionally quantified SBB-

replication within each ROI, using Bayes factors (Jeffreys, 1961). Similar to Boekel et al.

(2015), here we used the adjusted (one-sided) Jeffry's test (Jeffreys, 1961) based on a uniform

prior distribution for the correlation coefficient. As we intended to confirm the SBB-

associations defined in the discovery subsamples, the alternative hypothesis (H1) in this study

was considered one-sided (in line with Boekel et al. (2015)). We used implementation of the

Bayes   Factors   for   correlations   from   the   R   function   available   at

http://www.josineverhagen.com/?page_id=76.

To facilitate the interpretation, Bayes factors (BF) were summarized into four categories as

illustrated in the bar legend of Figure 2. A BF_{01} lower than 1/3 shows that the data is three

times or more likely to have happened under H1 than H0. Accordingly, this value defines the

"successful" replication.

Investigation on factors influencing replicability of SBB-associations among healthy

individuals:

Sample size: In order to study the influence of sample size on the replicability of SBB-

associations, for each psychological measure, the healthy sample (eNKI) was divided into

discovery and test pairs at three different ratios: 70% discovery and 30% test, 50% discovery

and 50% test and finally 30% discovery and 70% test. As mentioned earlier, in each case, the

discovery and test counterparts were randomly generated 100 times in order to quantify the

replication rates. For example, to assess the replicability of brain structural associations of

age, in the case of "70% discovery and 30% test", the entire NKI sample (n = 466) was

divided into a discovery group of n = 326 participants and an age- and sex-matched test pair

sample of n = 138 and this split procedure was repeated 100 times. Similarly, for generating

equal-sized discovery and test subsamples, 100 randomly generated age and sex matched

split-half samples were generated from the main NKI cohort.

Due to the multi-site structure of the ADNI cohort, when generating unequal sized discovery

and test samples, we did not achieve a good simultaneous matching of age, sex and site, while

trying to maintain samples sizes in each subgroup reasonably large. Thus, in this cohort, we

did not directly study the influence of the sample size and the replicability rates were only

quantified for equal sized discovery and test samples (187 participants matched for age, sex

and site between discovery and test pairs).

Effect size: Furthermore, to study the influence of the effect size on the replication rates, we

focused on the effect sizes within each a-priori ROI in the discovery samples. Here we tested

the following two assumptions:

1) ROIs with larger effect sizes in the discovery sample result in larger effect sizes in the test

sample pairs (i.e. positive association between effect size in the discovery and test samples).

2) ROIs with larger effect sizes in the discovery sample are more likely to result in a

"significant" replication in the independent sample.

To test the first assumption, in the "ROI-based SBB-replicability" the association between

effect size in the discovery and test pairs were calculated for each psychological measure.

These associations were calculated separately for the replicated (defined using "sign"

criterion) and not-replicated ROIs. We expected to find a positive association between

discovery and confirmatory effect sizes, for the "successfully replicated effects".

To test the second assumption, for each ROI, we calculated its replication statistical power

and compared it between replicated and not-replicated ROIs (here replication was defined

using "Statistical Significance" criterion). The statistical power of a test is the probability that

it will correctly reject the null hypothesis when the null is false. In a bias-free case, the power

of the replication is a function of the replication sample size, real size of the effect and the

nominal type I error rate (α). In this study, the replication power was estimated based on the

size of the effects as they were defined in the discovery sample and a significant threshold of

0.05 (one-sided) and was calculated using "pwr" library in R (https://www.r-project.org).

These analyses were performed for each discovery-test split size, separately (i.e. 70%-30%,

50%-50% and 30%-70% discovery-test sample sizes, respectively).

Acknowledgement:

This work was supported by the Deutsche Forschungsgemeinschaft (DFG, GE 2835/1-1, EI

816/4-1), the Helmholtz Portfolio Theme 'Supercomputing and Modelling for the Human

Brain' and the European Union's Horizon 2020 Research and Innovation Programme under

Grant Agreement No. 720270 (HBP SGA1) and Grant Agreement No. 785907 (HBP SGA2).

Clinical data collection and sharing for this project was funded by the Alzheimer's Disease

Neuroimaging Initiative (ADNI) (National Institutes of Health Grant U01 AG024904) and

DOD ADNI (Department of Defense award number W81XWH-12-2-0012). ADNI is funded

by the National Institute on Aging, the National Institute of Biomedical Imaging and

Bioengineering, and through generous contributions from the following: AbbVie,

Alzheimer's

Association; Alzheimer's Drug Discovery Foundation; Araclon Biotech; BioClinica, Inc.;

Biogen; Bristol-Myers Squibb Company; CereSpir, Inc.; Cogstate; Eisai Inc.; Elan

Pharmaceuticals, Inc.; Eli Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd and

its affiliated company Genentech, Inc.; Fujirebio; GE Healthcare; IXICO Ltd.; Janssen

Alzheimer Immunotherapy Research & Development, LLC.; Johnson & Johnson

Pharmaceutical Research & Development LLC.; Lumosity; Lundbeck; Merck & Co., Inc.;

Meso Scale Diagnostics, LLC.; NeuroRx Research; Neurotrack Technologies; Novartis

Pharmaceuticals Corporation; Pfizer Inc.; Piramal Imaging; Servier; Takeda Pharmaceutical

Company; and Transition Therapeutics. The Canadian Institutes of Health Research is

providing funds to support ADNI clinical sites in Canada. Private sector contributions are

facilitated by the Foundation for the National Institutes of Health (www.fnih.org). The

grantee organization is the Northern California Institute for Research and Education, and the

study is coordinated by the Alzheimer's Therapeutic Research Institute at the University of

Southern California. ADNI data are disseminated by the Laboratory for Neuro Imaging at the

University of Southern California.

Competing interests: The authors declare no competing interests.

References:

Albers C, Lakens D. 2018a. When power analyses based on pilot data are biased: Inaccurate
effect size estimators and follow-up bias. *J Exp Soc Psychol* **74**:187–195.
doi:10.1016/j.jesp.2017.09.004
Albers C, Lakens D. 2018b. When power analyses based on pilot data are biased: Inaccurate
effect size estimators and follow-up bias. *J Exp Soc Psychol* **74**:187–195.
doi:10.1016/j.jesp.2017.09.004
Anderson ML. 2015. Précis of after Phrenology: Neural Reuse and the Interactive Brain.
Behav Brain Sci. doi:10.1017/S0140525X15000631
Ashburner J. 2007. A fast diffeomorphic image registration algorithm. *Neuroimage* **38**:95–
113. doi:10.1016/j.neuroimage.2007.07.007
Ashburner J, Friston KJ. 2005. Unified segmentation. *Neuroimage* **26**:839–851.
doi:10.1016/j.neuroimage.2005.02.018
Boekel W, Wagenmakers EJ, Belay L, Verhagen J, Brown S, Forstmann BU. 2015. A purely
confirmatory replication study of structural brain-behavior correlations. *Cortex* **66**:115–
133. doi:10.1016/j.cortex.2014.11.019
Bressler SL. 1995. Large-scale cortical networks and cognition. *Brain Res Rev*.
doi:10.1016/0165-0173(94)00016-I
Broca P. 1865. Sur le siège de la faculté du langage articulé. *Bull la Société d'anthropologie*
Paris **6**:377–393. doi:10.3406/bmsap.1865.9495
Button KS, Ioannidis JPA, Mokrysz C, Nosek BA, Flint J, Robinson ESJ, Munafò MR. 2013.
Power failure: why small sample size undermines the reliability of neuroscience. *Nat Rev*
Neurosci **14**:365–76. doi:10.1038/nrn3475
Cremers HR, Wager TD, Yarkoni T. 2017. The relation between statistical power and
inference in fMRI. *PLoS One* **12**:1–20. doi:10.1371/journal.pone.0184923
De Boeck P, Jeon M. 2018. Perceived Crisis and Reforms: Issues, Explanations, and
Remedies. *Psychol Bull* **144**:757–777. doi:10.1037/bul0000154
Delis DC, Kaplan E, Kramer JH. 2001. Delis-Kaplan Executive Function System (D-KEFS)
examiner's manual. San Antonio, TX: The Psychological Corporation.
Draganski B, Gaser C, Busch V, Schuierer G, Bogdahn U, May A. 2004. Changes in grey
matter induced by training Newly honed juggling skills show up as a transient feature on
a brain-imaging scan . *Nature* **427**:311–312. doi:10.1038/427311a
Dwan K, Gamble C, Williamson PR, Kirkham JJ. 2013. Systematic Review of the Empirical
Evidence of Study Publication Bias and Outcome Reporting Bias - An Updated Review.
PLoS One. doi:10.1371/journal.pone.0066844
Eklund A, Nichols TE, Knutsson H. 2016. Cluster failure: Why fMRI inferences for spatial
extent have inflated false-positive rates. *Proc Natl Acad Sci U S A* **113**:7900–5.
doi:10.1073/pnas.1602413113
Fan J, Mccandliss BD, Sommer T, Raz A, Posner MI. 2002. Testing the Efficiency and
Independence of Attentional Networks. *J Cogn Neurosci* **14**:340–347.
Fjell AM, Westlye LT, Grydeland H, Amlien I, Espeseth T, Reinvang I, Raz N, Dale AM,
Walhovd KB. 2014. Accelerating cortical thinning: unique to dementia or universal in
aging? *Cereb Cortex* **24**:919–34. doi:10.1093/cercor/bhs379
Forstmeier W, Schielzeth H. 2011. Cryptic multiple hypotheses testing in linear models:
Overestimated effect sizes and the winner's curse. *Behav Ecol Sociobiol* **65**:47–55.
doi:10.1007/s00265-010-1038-5
Gaser C, Dahnke R. 2016. CAT - A Computational Anatomy Toolbox for the Analysis of
Structural MRI Data. *HBM Conf 2016* **32**:7743.
Gelman A, Loken E. 2014. The garden of forking paths: Why multiple comparisons can be a
problem, even when there is no "fishing expedition" or "p-hacking" and the research

hypothesis was posited ahead of time. *Psychol Bull* **140**:1272–1280.
doi:dx.doi.org/10.1037/a0037714

Genon S, Bastin C, Angel L, Collette F, Bahri MA, Salmon E. 2014. A partial least squares
analysis of the self reference effect in Alzheimer's disease: A reply to Irish. *Cortex*.
doi:10.1016/j.cortex.2014.02.003

Genon S, Reid A, Langner R, Amunts K, Eickhoff SB. 2018. How to Characterize the
Function of a Brain Region. *Trends Cogn Sci*. doi:10.1016/j.tics.2018.01.010

Genon S, Simon J, Bahri MA, Collette F, Souchay C, Jaspar M, Bastin C, Salmon E. 2016.
Relating pessimistic memory predictions to Alzheimer's disease brain structure. *Cortex*
85:151–164. doi:10.1016/j.cortex.2016.09.014

Genon S, Wensing T, Reid A, Hoffstaedter F, Caspers S, Grefkes C, Nickl-Jockschat T,
Eickhoff SB. 2017. Searching for behavior relating to grey matter volume in a-priori
defined right dorsal premotor regions: Lessons learned. *Neuroimage* **157**:144–156.
doi:10.1016/j.neuroimage.2017.05.053

Gorgolewski KJ, Varoquaux G, Rivera G, Schwarz Y, Ghosh SS, Maumet C, Sochat V V.,
Nichols TE, Poldrack RA, Poline J-B, Yarkoni T, Margulies DS. 2015. NeuroVault.org:
a web-based repository for collecting and sharing unthresholded statistical maps of the
human brain. *Front Neuroinform* **9**:8. doi:10.3389/fninf.2015.00008

Habeck C, Stern Y. 2010. Multivariate data analysis for neuroicaming data: overview and
Application to Alzheimer's disease. *Cell Biochem Biophys* **58**:53–67.
doi:10.1007/s12013-010-9093-0.Multivariate

Ioannidis JPA. 2018. Why replication has more scientific value than original discovery.
Behav Brain Sci **41**:e137. doi:10.1017/S0140525X18000729

Ioannidis JPA. 2008. Why most discovered true associations are inflated. *Epidemiology*
19:640–648. doi:10.1097/EDE.0b013e31818131e7

Ioannidis JPA. 2005. Why most published research findings are false. *PLoS Med*.
doi:10.1371/journal.pmed.0020124

Ismaylova E, Di Sante J, Gouin J-P, Pomares FB, Vitaro F, Tremblay RE, Booij L. 2018.
Associations Between Daily Mood States and Brain Gray Matter Volume, Resting-State
Functional Connectivity and Task-Based Activity in Healthy Adults. *Front Hum*
Neurosci **12**:168. doi:10.3389/fnhum.2018.00168

Jeffreys H. 1961. Theory of probability. Oxford, Uk.: Oxford University Press.

John LK, Loewenstein G, Prelec D. 2012. Measuring the prevalence of questionable research
practices with incentives for truth telling. *Psychol Sci* **23**:524–32.
doi:10.1177/0956797611430953

Kanai R. 2016. Open questions in conducting confirmatory replication studies: Commentary
on Boekel et al., 2015. *Cortex*. doi:10.1016/j.cortex.2015.02.020

Kanai R, Feilden T, Firth C, Rees G. 2011. Political orientations are correlated with brain
structure in young adults. *Curr Biol* **21**:677–680. doi:10.1016/j.cub.2011.03.017

Kanai R, Rees G. 2011. The structural basis of inter-individual differences in human
behaviour and cognition. *Nat Rev Neurosci* **12**:231–242. doi:10.1038/nrn3000

Kharabian Masouleh S, Arélin K, Horstmann A, Lampe L, Kipping JA, Luck T, Riedel-Heller
SG, Schroeter ML, Stumvoll M, Villringer A, Witte AV. 2016. Higher body mass index
in older adults is associated with lower gray matter volume: Implications for memory
performance, Neurobiology of Aging. Elsevier Ltd.
doi:10.1016/j.neurobiolaging.2015.12.020

Kim EJ, Pellman B, Kim JJ. 2015. Stress effects on the hippocampus: A critical review.
Learn Mem. doi:10.1101/lm.037291.114

Kriegeskorte N, Lindquist MA, Nichols TE, Poldrack RA, Vul E. 2010. Everything you never
wanted to know about circular analysis, but were afraid to ask. *J Cereb Blood Flow*
Metab. doi:10.1038/jcbfm.2010.86

Lilienfeld SO. 2017. Psychology's Replication Crisis and the Grant Culture: Righting the
Ship. *Perspect Psychol Sci* **12**:660–664. doi:10.1177/1745691616687745
Loken E, Gelman A. 2017. Measurement error and the replication crisis. *Science (80-)*.
doi:10.1126/science.aal3618
Luders E, Kurth F, Mayer EA, Toga AW, Narr KL, Gaser C. 2012. The Unique Brain
Anatomy of Meditation Practitioners: Alterations in Cortical Gyrification. *Front Hum*
Neurosci **6**:34. doi:10.3389/fnhum.2012.00034
Luders E, Kurth F, Toga AW, Narr KL, Gaser C. 2013. Meditation effects within the
hippocampal complex revealed by voxel-based morphometry and cytoarchitectonic
probabilistic mapping. *Front Psychol* **4**:398. doi:10.3389/fpsyg.2013.00398
Matsuo K, Nicoletti M, Nemoto K, Hatch JP, Peluso MAM, Nery FG, Soares JC. 2009. A
voxel-based morphometry study of frontal gray matter correlates of impulsivity. *Hum*
Brain Mapp **30**:1188–1195. doi:10.1002/hbm.20588
McCrae RR, Costa PT. 2004. A contemplated revision of the NEO Five-Factor Inventory.
Pers Individ Dif **36**:587–596. doi:10.1016/S0191-8869(03)00118-1
McEwen BS, Nasca C, Gray JD. 2016. Stress Effects on Neuronal Structure: Hippocampus,
Amygdala, and Prefrontal Cortex. *Neuropsychopharmacology*.
doi:10.1038/npp.2015.171
McIntosh AR, Mišić B. 2013. Multivariate Statistical Analyses for Neuroimaging Data. *Annu*
Rev Psychol **64**:499–525. doi:10.1146/annurev-psych-113011-143804
Miller KL, Alfaro-Almagro F, Bangerter NK, Thomas DL, Yacoub E, Xu J, Bartsch AJ,
Jbabdi S, Sotiropoulos SN, Andersson JLR, Griffanti L, Douaud G, Okell TW, Weale P,
Dragonu I, Garratt S, Hudson S, Collins R, Jenkinson M, Matthews PM, Smith SM.
2016. Multimodal population brain imaging in the UK Biobank prospective
epidemiological study. *Nat Neurosci* **19**:1523–1536. doi:10.1038/nn.4393
Moeller JR, Habeck CG. 2006. Reciprocal benefits of mass-univariate and multivariate
modeling in brain mapping: Applications to event-related functional MRI, H215O-, and
FDG-PET. *Int J Biomed Imaging* **2006**:1–13. doi:10.1155/IJBI/2006/79862
Moradi E, Hallikainen I, Hänninen T, Tohka J. 2017. Rey's Auditory Verbal Learning Test
scores can be predicted from whole brain MRI in Alzheimer's disease. *NeuroImage Clin*
13:415–427. doi:10.1016/j.nicl.2016.12.011
Muhlert N, Ridgway GR. 2016. Failed replications, contributing factors and careful
interpretations: Commentary on Boekel et al., 2015. *Cortex*.
doi:10.1016/j.cortex.2015.02.019
Munafò MR, Nosek BA, Bishop DVM, Button KS, Chambers CD, Percie Du Sert N,
Simonsohn U, Wagenmakers EJ, Ware JJ, Ioannidis JPA. 2017. A manifesto for
reproducible science. *Nat Hum Behav* **1**:1–9. doi:10.1038/s41562-016-0021
Nooner KB, Colcombe SJ, Tobe RH, Mennes M, Benedict MM, Moreno AL, Panek LJ,
Brown S, Zavitz Stephen TT, Li Q, Sikka S, Gutman D, Bangaru S, Schlachter RT,
Anwar SMK, Hinz CM, Kaplan MS, Rachlin AB, Adelsberg S, Cheung B, Khanuja R,
Yan C, Courtney CCC, King M, Wood D, Cox CL, Kelly AMC, Petkova E, Reiss PT,
Duan N, Thomsen D, Biswal B, Coffey B, Hoptman MJ, Javitt DC, Pomara N, Sidtis JJ,
Koplewicz HS, Castellanos FX, Leventhal BL, Milham MP. 2012. The NKI-Rockland
sample: A model for accelerating the pace of discovery science in psychiatry. *Front*
Neurosci. doi:10.3389/fnins.2012.00152
Nostro AD, Müller VI, Reid AT, Eickhoff SB. 2017. Correlations between Personality and
Brain Structure: A Crucial Role of Gender. *Cereb Cortex* **27**:3698–3712.
doi:10.1093/cercor/bhw191
Open Science Collaboration OS. 2015. Estimating the reproducibility of psychological
science. *Science* **349**:aac4716. doi:10.1126/science.aac4716
Pessoa L. 2014. Understanding brain networks and brain organization. *Phys Life Rev*.

doi:10.1016/j.plrev.2014.03.005

Poldrack RA. 2010. Mapping mental function to brain Structure: How can cognitive
Neuroimaging Succeed? *Perspect Psychol Sci* **5**:753–761.
doi:10.1177/1745691610388777

Poldrack RA, Baker CI, Durnez J, Gorgolewski KJ, Matthews PM, Munafò MR, Nichols TE,
Poline J-B, Vul E, Yarkoni T. 2017. Scanning the horizon: towards transparent and
reproducible neuroimaging research. *Nat Rev Neurosci* **18**:115–126.
doi:10.1038/nrn.2016.167

Rajapakse JC, Giedd JN, Rapoport JL. 1997. Statistical approach to segmentation of single-
channel cerebral mr images. *IEEE Trans Med Imaging* **16**:176–186.
doi:10.1109/42.563663

Salat DH, Buckner RL, Snyder AZ, Greve DN, Desikan RSR, Busa E, Morris JC, Dale AM,
Fischl B. 2004. Thinning of the cerebral cortex in aging. *Cereb Cortex* **14**:721–30.
doi:10.1093/cercor/bhh032

Schmidt M. 1996. RAVLT (Rey Auditory Verbal Learning Test: A Handbook).

Schönbrodt FD, Perugini M. 2013. At what sample size do correlations stabilize? *J Res Pers*
47:609–612. doi:10.1016/j.jrp.2013.05.009

Scoville WB, Milner B. 1957. Loss of recent memory after bilateral hippocampal lesions. *J*
Neurol Neurosurg Psychiatry **20**:11–21. doi:10.1136/jnnp-2015-311092

Simmons JP, Nelson LD, Simonsohn U. 2011. False-positive psychology: Undisclosed
flexibility in data collection and analysis allows presenting anything as significant.
Psychol Sci **22**:1359–1366. doi:10.1177/0956797611417632

Smith SM, Nichols TE. 2018. Statistical Challenges in "Big Data" Human Neuroimaging.
Neuron **97**:263–268. doi:10.1016/j.neuron.2017.12.018

Smith SM, Nichols TE. 2009. Threshold-free cluster enhancement: Addressing problems of
smoothing, threshold dependence and localisation in cluster inference. *Neuroimage*
44:83–98. doi:10.1016/j.neuroimage.2008.03.061

Spielberger CD, Gorsuch RL, Lushene RE. 1970. Manual for the State- Trait Anxiety
Inventory. Palo Alto, CA: Consulting Psychologists Press.

Taubert M, Lohmann G, Margulies DS, Villringer A, Ragert P. 2011. Long-term effects of
motor training on resting-state networks and underlying brain structure. *Neuroimage*
57:1492–1498. doi:10.1016/j.neuroimage.2011.05.078

Tohka J, Zijdenbos A, Evans A. 2004. Fast and robust parameter estimation for statistical
partial volume models in brain MRI. *Neuroimage* **23**:84–97.
doi:10.1016/j.neuroimage.2004.05.007

Tononi G, Edelman GM, Sporns O. 1998. Complexity and coherency: Integrating information
in the brain. *Trends Cogn Sci*. doi:10.1016/S1364-6613(98)01259-5

Tukey JW. 1980. We need both Eploratory and Confirmatory We Need Both Exploratory. *Am*
Stat **34**:23–25. doi:10.2307/2682991

Turner BO, Paul EJ, Miller MB, Barbey AK. 2018. Small sample sizes reduce the
replicability of task-based fMRI studies. *Commun Biol* **1**:62. doi:10.1038/s42003-018-
0073-z

Van Essen DC, Smith SM, Barch DM, Behrens TEJ, Yacoub E, Ugurbil K. 2013. The WU-
Minn Human Connectome Project: An overview. *Neuroimage* **80**:62–79.
doi:10.1016/j.neuroimage.2013.05.041

Vanasse TJ, Fox PM, Barron DS, Robertson M, Eickhoff SB, Lancaster JL, Fox PT. 2018.
BrainMap VBM: An environment for structural meta-analysis. *Hum Brain Mapp* 1–18.
doi:10.1002/hbm.24078

Wagenmakers E-J, Verhagen J, Ly A, Bakker M, Lee MD, Matzke D, Rouder JN, Morey RD.
2015. A power fallacy. *Behav Res Methods* **47**:913–917. doi:10.3758/s13428-014-0517-
4

Wechsler D. 1999. Wechsler Abbreviated Scale of Intelligence. San Antonio, TX: The
Psychologica Corporation.
Willette A a, Kapogiannis D. 2014. Does the brain shrink as the waist expands? *Ageing Res*
Rev 1–12. doi:10.1016/j.arr.2014.03.007
Yarkoni T. 2009. Big Correlations in Little Studies. *Perspect Psychol Sci* **4**:294–298.
doi:10.1111/j.1745-6924.2009.01127.x
Yuan P, Raz N. 2014. Prefrontal cortex and executive functions in healthy adults: A meta-
analysis of structural neuroimaging studies. *Neurosci Biobehav Rev* **42**:180–192.
doi:10.1016/j.neubiorev.2014.02.005
1088

Figure legends:

Figure 1. Replicability of exploratory results within healthy cohort. Frequency of spatial overlap (density plots and aggregate maps) of significant findings from exploratory analysis over 100 random subsamples are depicted for few behavioral score. For each score, columns show the results of three different discovery sample sizes (i.e. when discovery cohorts are generated from 70%, 50% or 30% of the main sample, from left to right respectively (x-axis)). The density plots show the distribution of values within their corresponding aggregate map. The y-axis depicts the frequency of spatial overlap (in %) and the density plots show the distribution of values within their corresponding aggregate maps. In addition to age and BMI (A,B), which are used as benchmarks, the top three behavioral scores with the highest frequency of overlapping findings are depicted (C-E). Within each density plot, the box-plot shows the quartiles and extent of the distribution and the white dot depicts the median of percentage of overlap. On the spatial maps, lighter colors denote higher number of samples with a significant association at the respective voxel. BMI : body mass index; CWI : color-word interference; n = number of participants within the discovery samples.

Figure 2. ROI-based confirmatory replication results within healthy cohort. Donut plots summerising ROI-based replication rates (% of ROI) using three different critera for three different sample sizes among heathy participants. The most inner layers depict replication using "sign" only (blue: replicated, orange: not replciated). The middle layers define replication based on similar "sign" as well as "statistical significance" (i.e. $p < 0.05$) (blue: replicated, orange: not replciate). The most outer layers define replication using "bayes factor" (blue: "moderate-to-string evidece for H1, light blue: anecdotal evidence for H1; light orange: anecdotal evidence for H0, orange: "moderate-to-string evidece for H0);

Figure 3. Discovery versus replication effects sizes: Scatter plots of **correlation coefficients** in the discovery versus replication sample for all ROIs from 100 splits within healthy cohort; each point denotes one ROI, which is color-coded based on its replication

status (by-"sign"). The size of each point is proportional to its estimated statistical power of replication. Regresion lines are drawn for the replicated and unreplicated ROIs, separately.

Figure 4. Replicability of positive association between immediate-recall and GMV within ADNI cohort. A, B: Replicability of exploratory results: Frequency of spatial overlaps (density plot and aggregate maps) over 100 random subsamples. Within the density plot, the box-plot shows the quartiles and extent of the distribution and the white dot depicts the median of percentage of overlap. C, D: ROI-based confirmatory replication results: C: Original versus replication effects sizes (correlation coefficient) for all ROIs from 100 splits; points are color-coded based on their replciation status (by-"sign") and size of each point is proportional to the estimated statistical power of replication. Regresion lines are drawn for the replicated and unreplicated ROIs, separately. D: Donut plots summerising ROI-based replicability rates using three different critera. The most inner layer depicts replicability using "sign" only (blue: replicated, orange: not replciated). The middle layer, defines replication based on similar "sign" as well as "statistical significance" (i.e. $p < 0.05$) (blue: replicated, orange: not replciate). The most outer layer reflects replicability using bayes factor " (blue: "moderate-to-string evidece for H1, light blue: anecdotal evidence for H1; light orange: anecdotal evidence for H0, orange: "moderate-to-string evidece for H0); Discovery and replication samples have equal size (n = 184) and are matched for age, sex and site.

Figure 5. box-plots showing distribution of sample sizes (log-scale) of VBM studies by their publication year (data from the BrainMap database; see (Vanasse et al., 2018)). Each box shows the quantiles (25% and 75%) of the distribution and the gray horizontal line within each box, depicts the median of the distribution.

Table 1. Summary of exploratory findings. For each discovery sample size, the number of clusters in which gray matter volume is positively or negatively associated with the tested phenotypic or psychological score is reported. The number of splits (out of 100) in which the clusters were detected are noted in parentheses (i.e. % of splits with at least one significant cluster [in the respective direction])

Healthy cohort	n_discovery = 70% n_total		n_discovery = 50% n_total		n_discovery = 30% n_total	
	# positively associated clusters (split%)	# negatively associated clusters (split%)	# positively associated clusters (split%)	# negatively associated clusters (split%)	# positively associated clusters (split%)	# negatively associated clusters (split%)
Age (years) n-total = 466	77 (54%)	154 (100%)	5 (4%)	522 (100%)	1 (1%)	1781 (100%)
BMI (kg/m^2) n-total = 466	0	1741 (100%)	0	2276 (100%)	0	1937 (96%)
Perceptual IQ (sum of t-scores) n-total = 466	499 (83%)	0	256 (58%)	0	145 (33%)	0
Word-context (# of consecutively correct) n-total = 262	337 (80%)	0	159 (47%)	0	80 (21%)	0
CWI (interference) (sec) n-total = 449	0	163 (53%)	1 (1%)	122 (39%)	6 (1%)	60 (26%)
Clinical cohort	-		n_discovery = 50% n_total		-	
RAVLT (# total immediate recall)	-	-	309 (84%)	0	-	-

Abbreviations: BMI : body mass index; IQ : intelligence quotient, CWI: color-word interference task; RAVLT : Rey auditory verbal learning task;

Supplementary material:

Supplementary File 1: Including Table S1, Table S2;

Supplementary Tables legends:

Table S1. Distribution of the raw phenotypical and psychological scores in the whole sample.

Table S2. Summary of the exploratory findings. For each discovery sample size, the number of clusters in which gray matter volume is positively or negatively associated with the tested psychological score is reported. Number of splits (out of 100) in which the clusters were detected are noted in parentheses.

Supplementary Figures (Figure S1, Figure S2) legends:

Figure S1. Summary of replication of positive associations between immediate-recall and GMV within healthy cohort. A: Frequency of spatial overlap (density plots and aggregate maps) of significant findings from exploratory analysis over 100 random subsamples. Columns show results of three different discovery sample sizes (i.e. when discovery cohorts are generated from 70%, 50% or 30% of the main sample, from left to right respectively (x-axis)). The density plots show distribution of values within their corresponding aggregate map. The y-axis depicts frequency of spatial overlap (in %) and the density plots show distribution of values within their corresponding aggregate map. On the spatial maps, warmer colors denote higher number of samples with a significant association at the respective voxel. B: ROI-based confirmatory replication results: Top row : Donut plots summerising ROI-based replicability rates (% of ROI) using three different critera for three different sample sizes. The most inner layers depict replicability using "sign" only (blue: replicated, orange: not replciated). The middle layers define replication based on similar "sign" as well as "statistical significance" (i.e. $p < 0.05$) (blue: replicated, orange: not replciate). The most outer layers reflects replicability using bayes factor " (blue: "moderate-to-string evidece for H1, light blue: anecdotal evidence for H1; light orange: anecdotal evidence for H0, orange: "moderate-to-string evidece for H0); Bottom row: Scatter plots of

effect sizes (correlation coefficient) in the discovery versus replication sample for all ROIs from 100 splits within healthy cohort; Points are color-coded based on their replciation status (by-"sign") and size of each point is proportional to the estimated statistical power of replication. Regresion lines are drawn for the replciated and unreplicated ROIs, separately.

Figure S2. ROI-based confirmatory replication results for five personality subscores within healthy cohort. Donut plots summerising ROI-based replication rates (% of ROI) using three different critera for three different sample sizes among heathy participants. The most inner layers depict replication using "sign" only (blue: replicated, orange: not replciated). The middle layers define replication based on similar "sign" as well as "statistical significance" (i.e. $p < 0.05$) (blue: replicated, orange: not replciate). The most outer layers define replication using "bayes factor" (blue: "moderate-to-string evidece for H1, light blue: anecdotal evidence for H1; light orange: anecdotal evidence for H0, orange: "moderate-to-string evidece for H0);