Id proteins promote a cancer stem cell phenotype in triple negative breast cancer via

negative regulation of Robo1

Wee S. Teo[1,2], Holly Holliday[1,2*], Nitheesh Karthikeyan[3*], Aurélie S. Cazet[1,2], Daniel L.

Roden[1,2], Kate Harvey[1], Christina Valbirk Konrad[1], Reshma Murali[3], Binitha Anu Varghese[3],

Archana P. T.[3,4], Chia-Ling Chan[1,2], Andrea McFarland[1], Simon Junankar[1,2], Sunny Ye[1],

Jessica Yang[1], Iva Nikolic[1,2], Jaynish S. Shah[5], Laura A. Baker[1,2], Ewan K.A. Millar[1,6,7,8],

Mathew J. Naylor[1,2,10], Christopher J. Ormandy[1,2], Sunil R. Lakhani[11], Warren Kaplan[1,12],

Albert S. Mellick[13,14], Sandra A. O'Toole[1,9], Alexander Swarbrick[1,2$], Radhika Nair[1,2,3$]

Affiliations

[1]Garvan Institute of Medical Research, Darlinghurst, New South Wales, 2010, Australia

[2]St Vincent's Clinical School, Faculty of Medicine, UNSW Sydney, New South Wales, 2010,

Australia

[3]Cancer Research Program, Rajiv Gandhi Centre for Biotechnology, Kerala, 695014, India

[4]Manipal Academy of Higher Education, Manipal, Karnataka, 576104, India

[5]Centenary Institute, The University of Sydney, New South Wales, 2050, Australia

[6] Department of Anatomical Pathology, NSW Health Pathology, St George Hospital,

Kogarah NSW 2217 Australia

[7] School of Medical Sciences, UNSW Sydney, Kensington NSW 2033, Australia.

[8] School of Medicine, Western Sydney University, Locked Bag 1797, Penrith, NSW 2751,

Australia

[9]Department of Tissue Pathology and Diagnostic Oncology, Royal Prince Alfred Hospital,

Camperdown

[10]Discipline of Physiology & Bosch Institute, School of Medical Sciences, University of

Sydney, NSW 2006

[11]The University of Queensland, UQ Centre for Clinical Research, School of Medicine and

Pathology Queensland, The Royal Brisbane & Women's Hospital, Herston, Brisbane,

Queensland 4029, Australia

[12]Peter Wills Bioinformatics Centre, Garvan Institute of Medical Research, Darlinghurst,

NSW 2010, Australia

[13]UNSW Medicine, University of NSW, 18 High St, Kensington NSW 2052, Australia.

[14]Ingham Institute for Applied Medical Research, 1 Campbell Street, Liverpool, NSW 2170,

Australia.

* These authors contributed equally

$ **Co-corresponding authors**

$ Dr Radhika Nair

Present address

Rajiv Gandhi Centre for Biotechnology

Kerala 695014, India

radhikanair@rgcb.res.in

Phone: +91-471-2781251

Fax: +91-471-2346333

$ A/Prof Alexander Swarbrick

Cancer Research Division, Garvan Institute of Medical Research

370 Victoria St, Darlinghurst, NSW, 2010

Australia

a.swarbrick@garvan.org.au

Phone: +61-2-9355-5780

Fax**:** +61-2-355-5869

Running title: Id-Robo1 axis controls CSC phenotypes in TNBC

Abstract

Background

Breast cancers display phenotypic and functional heterogeneity. Several lines of evidence

support the existence of cancer stem cells (CSCs) in certain breast cancers, a minor

population of cells capable of tumor initiation and metastatic dissemination. Identifying

factors that regulate the CSC phenotype is therefore important for developing strategies to

treat metastatic disease. The Inhibitor of Differentiation Protein 1 (Id1) and its closely related

family member Inhibitor of Differentiation 3 (Id3) (collectively termed Id) are expressed by a

diversity of stem cells and are required for metastatic dissemination in experimental models

of breast cancer. Here, we show that ID1 is expressed in rare neoplastic cells within ER-

negative breast cancers and enriched in brain metastases compared to patient matched

primary tissues. We go on to investigate the molecular mechanism underpinning the role of

Id in CSC biology in the TNBC subtype.

Methods

To address the function of Id1 expressing cells within tumors, we developed three

independent murine models of Triple Negative Breast Cancer (TNBC) in which a genetic

reporter permitted the prospective isolation of Id1$^+$ cells. We also developed an inducible Id

knock down system which allowed us to deplete Id proteins in the 4T1 metastatic murine cell

line.

Results

Using the Id1C3-Tag,p53-/- and 4T1 models, we found that Id1$^+$ cells are enriched for self-

renewal in tumorsphere assays *in vitro* and for tumor initiation *in vivo*. Conversely, depletion

of Id1 and Id3 in the 4T1 murine model of TNBC demonstrates that Id are required for cell

proliferation and self-renewal *in vitro*, as well as primary tumor growth and metastatic

colonization of the lung *in vivo*. Using combined bioinformatic analysis, we have defined a

novel mechanism of Id protein function via negative regulation of the Roundabout Axon

Guidance Receptor Homolog 1 (Robo1) leading to activation of a Myc transcriptional

programme.

Conclusions

Id proteins play a critical role in the proliferative, self renewal and metastatic CSC

phenotypes in TNBC. Our work has shed light on how Id acts on Robo1 to possibly control a

Myc network and could be used to target the CSC population in the TNBC subtype.

Key words: Id proteins, Robo1, Cancer stem cell, Metastasis, Myc signature

Background

Several lines of evidence suggest that rare sub-populations of tumor cells, commonly termed cancer stem cells (CSCs), drive key tumor phenotypes such as self-renewal, drug resistance and metastasis and contribute to disease relapse and associated patient mortality (1-4). Recent evidence points to the hypothesis that CSCs are not static, but they exist in dynamic states, driven by critical transcription factors and are highly dependent on the microenvironmental cues (5-7). Understanding the molecular networks that are critical to the survival and plasticity of CSCs is fundamental to resolving clinical problems associated with chemo-resistance and metastatic residual disease.

The Inhibitor of DNA binding (ID) proteins have previously been recognized as regulators of CSCs and tumor progression (8). These proteins constitute a family of four highly conserved transcriptional regulators (ID1-4) that act as dominant-negative inhibitors of basic helix–loop–helix (bHLH) transcription factors. ID proteins are expressed in a tissue-specific and stage-dependent manner and are required for the maintenance of self-renewal and multipotency of embryonic and many tissue stem cells (9-12) . Previous studies have reported a functional redundancy among the four members of the mammalian Id family, in particular Id1 and Id3 (referred to collectively here as Id), and their overlapping expression patterns during normal development and cancer (13-17). Id2 and Id4 were not investigated in this work as they are found to have independent functions from Id1 and Id3.

A number of studies have implied a significant role for ID1 and ID3 in breast cancer progression and metastasis (14). We have previously demonstrated that Id1 cooperates with activated Ras signalling and promotes mammary tumor initiation and metastasis *in vivo* by supporting long-term self-renewal and proliferative capacity (18). Additional work has clearly implicated ID1 in regulating D- and E-type cyclins and their associated cyclin-

dependant kinases, CDK4 and CDK2 in human breast epithelial cells , p21 (19), the matrix

metalloproteinase MT1-MMP (20), KLF17 (21), Cyclin D1 (22), Bcl-2 (23), and BMI1 (24)

among others.

Even though several Id-dependent targets have been identified, we still lack a comprehensive

picture of the downstream molecular mechanisms controlled by Id and their associated

pathways mediating breast cancer progression and metastasis particularly in the poor

prognostic TNBC subtype. In this study, we demonstrate using four independent mouse

models of TNBC that Id is important for the maintenance of a CSC phenotype. We also

describe a novel mechanism by which Id controls the CSC state by negatively regulating

Robo1 to control proliferation and self-renewal via indirect activation of a Myc

transcriptional programme.

Methods

Lentivirus production

Lentiviral supernatant was produced by transfecting each lentiviral expression vector along

with third-generation lentiviral packaging and pseudotyping plasmids (73) into the packaging

cell line HEK293T. Briefly, 1.4×10^{6} cells were seeded in a 60mm tissue culture dish and

grown to 80% confluence. 3µg of expression plasmid was co-transfected with lentiviral

packaging and pseudotyping plasmids (2.25µg each of pMDLg/pRRE and pRSV-REV and

1.5µg of pMD2.G), using Lipofectamine 2000 (Invitrogen, Mulgrave, Vic, Australia)

according to the manufacturer's protocol. Cell culture medium was replaced after 24hr. The

viral supernatant was collected 48hr post transfection and filtered using a 0.45µm filter. The

filtered lentiviral supernatant was concentrated 20-fold by using Amicon Ultra-4 filter units

(100 kDa NMWL) (Millipore, North Ryde, NSW, Australia).

Lentiviral infection

4T1 cells were plated at a density of 1.0×10^5 cells per well in 6-well tissue culture plates and culture medium was replaced after 24hr with medium containing 8µg/mL of polybrene (Sigma-Aldrich, Lismore, NSW, Australia). The cells were infected overnight with the concentrated virus at 1:5 dilution. Culture medium was changed 24hr post infection and cells were grown until reaching confluence. Cells transduced with both pSLIK-Venus-TmiR-Id1 and pSLIK-Neo-TmiR-Id3 were sorted on FACS using Venus as a marker followed by selection with neomycin at 400µg/mL for 5 days. Cells transduced with pSLIK-Neo-TmiR-EGFP were also selected with neomycin.

Tumorsphere assay

Cells dissociated from modified 4T1 cells and p53-/- Id1/GFP, Id1C3-Tag tumors were put into tumorsphere assay as described previously (28).

Limiting dilution assay

Single-cell suspensions of FACS sorted Id1/GFP+ or unsorted viable tumor cells were prepared as described previously. Tumor cells were transplanted in appropriate numbers into the fourth mammary fat pad of 8- to 12-week-old FVB/N mice and aged till ethical end point. Extreme limiting dilution analysis (75) software was used to calculate the TPF.

Microarray and bioinformatics analysis

Total RNA from the samples were isolated using Qiagen RNeasy minikit (Qiagen, Doncaster, VIC, Australia. cDNA synthesis, probe labelling, hybridization, scanning and data processing were all conducted by the Ramaciotti Centre for Gene Function Analysis (The University of New South Wales). Gene expression profiling was performed using the Affymetrix

GeneChip® Mouse Gene 2.0 ST Array. Normalization and probe-set summarization was

performed using the robust multichip average method (74) implemented in the Affymetrix

Power Tools apt-probeset-summarize software (version 1.15.0) (using the -a rma option).

Differential expression between experimental groups was assessed using Limma (75) via the

limmaGP tool in GenePattern (76). Gene Set Enrichment Analysis (GSEA)

(http://www.broadinstitute.org/gsea) (77) was performed using the GSEA Pre-ranked module

on a ranked list of the limma moderated t-statistics, against gene-sets from v4.0 of the

MSigDB (77) and custom gene-sets derived from the literature. Microarray data are freely

available from GEO: GSE129790

Next generation sequencing

3.5×10^4 4T1 K1 cells were seeded in 6-well plates in 4T1 media and treated with or without

Doxorubicin (1 µg/mL) to induce Id1/3 knockdown. Cells were also transfected with non-

targeting control siRNA (Dharmacon D-001810-10-05) or Robo1 siRNA (Dharmacon M-

046944-01-0010). Cells were harvested after 48 hours and total RNA was extracted using the

automated QiaSymphony magnetic bead extraction system. The Illumina TruSeq Stranded

mRNA Library Prep Kit was used to generate libraries with 1 μg of input RNA following the

manufacturer's instructions. cDNA libraries were sequenced on the NextSeq system

(Illumina), with 75 bp paired-end reads. Quality control was checked using FastQC

(bioinformatics.babraham.ac.uk/projects/fastqc). Reads were then aligned to the mouse

reference genome Mm10 using STAR ultrafast universal RNA-Seq aligner (78).   Gene

feature counting was performed with RSEM (79). Replicate 3 from the Id1 KD group showed

no KD of Id1 by qPCR and was therefore removed prior to down-stream differential

expression analysis. Transcripts with expression counts of 0 across all samples were removed

and then normalised using TMM (80). The normalized counts were then log transformed

using voom (81) and differential expression was performed with limma (75). Differentially

expressed genes were visualized and explored using Degust (http://degust.erc.monash.edu/).

Genes with false discovery rate (FDR)<0.05 were considered significantly differentially

expressed. For GSEA analysis, genes were ranked based on the limma moderated t-statistic

and this was used as input for the GSEA desktop application (77). RNA sequencing data are

freely available from GEO: GSE129858.

Microarray (GSE129790) and RNA-Seq (GSE129858) datasets are available in SuperSeries

GSE129859.

Statistical analysis

Statistical analyses were performed using GraphPad Prism 6. All *in vitro* experiments were

done in 3 biological replicates each with 2 or more technical replicates. 5-10 mice were used

per condition for the *in vivo* experiments. Data represented are means ± standard deviation.

Statistical tests used are Unpaired student t-test and two-way-ANOVA. p-values <0.05 were

considered statistically significant with $*p< 0.05$, $**p< 0.01$, $*** p< 0.001$, $**** p< 0.0001$.

Results

Id marks a subset of cells with stem-like properties in TNBC models

We investigated the role of Id in the context of CSC biology in the TNBC molecular subtype. Immunohistochemistry (IHC) analysis revealed that ID1 is expressed by a small minority of cells (range 0.5-6% of total cancer cells) in ~50 % of ER-negative disease, namely TNBC and Her2+ tumors (Additional File1: Figure S1A, B). No significant difference in the distribution of ID3 expression was observed across different subtypes (data not shown).

To test the hypothesis that $Id1^+$ cells have a unique malignant phenotype, we developed two murine models of TNBC that permit the prospective isolation of $Id1^+$ cells for functional assays. In the first, we used the p53-/- TNBC tumor model where IHC analysis revealed that ~ 5% of neoplastic cells expressed Id1, consistent with the observation in the clinical samples, while Id3 marked a majority of the tumor cells in this model (Figure 1A). To create a genetic reporter cell line, p53-/- mammary tumor cells were transduced with a lentiviral GFP reporter construct under the control of the Id1 promoter (Id1/GFP), as described previously (25) (Additional File1: Figure S1C). FACS sorting for GFP expression followed by immunoblotting confirmed the ability of the Id1/GFP construct to prospectively enrich for Id1+ cells from this model (Additional File1: Figure S1D). We next sought to understand if Id1 marked cells with high self-renewal capacity in this model using tumorsphere assays, a well-established surrogate for cells with high self-renewal capacity (26, 27). We observed an increase in the self-renewal capacity of $Id1/GFP^+$ cells when compared to the unsorted cell population in the p53-/- model (Figure 1B). To establish the *in vivo* relevance of the increased self-renewal capacity of the $Id1/GFP^+$ tumor cells observed *in vitro*, we determined the tumor initiating capacity (TIC) of the $Id1/GFP^+$ cells using the limiting dilution assay (28). $Id1/GFP^+$ cells (1/68) showed a

significant (p value 0.02) 4.6-fold increase in tumor initiating cell frequency over Id1/GFP$^-$

cells (1/314) after serial passage (Figure 1C).

We used the Id1C3-Tag tumor model as a second murine model to assess the phenotype of

Id1$^+$ cells. In the C3-Tag tumor model, the expression of SV40-large T antigen in the

mammary epithelium under the control of the C3 promoter leads to the development of

TNBC in mice (29, 30). These tumors (C3-Tag) closely model the TNBC subtype as assessed

by gene expression profiling (30). To generate a genetic reporter of Id1 promoter activity in

TNBC, the C3-Tag model was crossed to a genetic reporter mouse model in which GFP is

knocked into the intron 1 of the Id1 gene (31). The resulting Id1GFPC3-Tag mice (called

Id1C3-Tag model) developed mammary tumors with similar kinetics as the parental C3-Tag

mice and have a classical basal phenotype characterized by CK14$^+$/CK8$^-$ phenotype

(Additional File1: Figure S1E). 5% and 60% of cells in the Id1C3-Tag tumor were stained

positive for Id1 and Id3 expression, respectively, as observed by IHC (Figure 1D). We were

able to isolate Id1$^+$ tumor cells with a high degree of purity by FACS based on GFP

expression followed by q-RT PCR (Figure 1E). The sorted cells were put into primary

tumorsphere assay and the spheres were serially passaged to secondary and tertiary spheres

which robustly selects for self-renewing cell populations. Similar to the p53-/- Id1/GFP

model, Id1$^+$/GFP$^+$ cells from the Id1C3-Tag model were enriched for sphere-forming capacity

(Figure 1F).

Using the Id1C3-Tag model, we also looked at the association of Id1/GFP expression with

the expression of established CSC markers CD29, CD24 and CD61. CD29$^+$/CD24$^+$ status was

previously reported to mark the tumorigenic subpopulation of cells in murine mammary

tumors (32, 33). The Id1$^+$/GFP$^+$ cells in the Id1C3-Tag model are predominantly of the

CD29$^+$/CD24$^+$ phenotype (Figure 1G), with a 1.6-fold higher proportion of cells expressing

both CD29 and CD24 compared to the Id1$^-$/GFP$^-$ cells which comprise the bulk of the tumor.

Interestingly, Id1$^+$/GFP$^+$ cells are also highly enriched for CD24$^+$/CD61$^+$ expression (more

than 6-fold increase in Id1$^+$/GFP$^+$ cells), which was also reported to mark a murine breast

CSC population (34) (Figure 1G).

We found no correlation between Id1 expression (as indicated by GFP$^+$) and the

CD29$^+$/CD24$^+$ phenotype in the first transplantation round (T1) using the p53-/- model, as the

percentage of CD29$^+$/CD24$^+$ cells was similar across each gating group (Additional File1:

Figure S1F). Interestingly, the Id1$^+$ cells, which are the putative cells that give rise to the

increased TIC as shown in Figure 1C, showed 10 times less CD24$^+$/CD29$^+$ cells in the second

transplantation round (T2) (34). The ability of the markers like CD24, CD29 and CD61 to

identify the CSC population is clearly model-dependent. In addition to CD29 and CD24, the

percentage of GFP$^+$ cells were also analysed and a higher percentage of GFP$^+$ cells was found

in the second transplantation round of the p53-/- tumor compared to the first round tumor

result (Additional File1: Figure S1G), consistent with the increase in TICs reported in Figure

1C.

Id requirement for self-renewal *in vitro* and metastatic competency *in vivo*

We next assessed the requirement for Id1 and Id3 in maintaining the CSC phenotypes.

Numerous studies have shown that there exists a functional redundancy between Id1 and Id3,

so studies typically require depletion of both the factors to reveal a phenotype (35).

Unfortunately we could not generate Id1 and Id3 double out knock mice for the C3Tag and

Id1/3 expressing reporter in the p53-/- tumor models due to technical reasons. Hence we

decided to look at the role of both Id1 and Id3 in the context of a knock down model. We

used the transplantable syngeneic 4T1 TNBC model, which has a high propensity to

spontaneously metastasize to distant sites (including bone, lung, brain and liver), mimicking

the aggressiveness of human breast cancers (36-41). IHC analysis showed that 15% of 4T1

tumor cells express high levels of Id1, and 35% have intermediate levels of Id1 expression,

whereas the expression of Id3 was found in most of the cells (Figure 2A).

In determine whether Id1 marks CSCs in the 4T1 TNBC model, 4T1 cells were transduced

with a lentiviral GFP reporter construct under the control of the Id1 promoter (Id1/GFP)

(Additional File2: Figure S2 A), as used in the $p53^{-/-}$ model and the cells were sorted based

on GFP expression. Id1/GFP+ cells showed more than 2-fold increase in Id1 expression

(Additional File2: Figure S2 B). Even though Id1/GFP+ and Id1/GFP- cells showed similar

proliferation *in vitro* (Additional File2: Figure S2 C), a significant enrichment of *in vitro* self-

renewal capacity was observed in Id1/GFP+ cells when compared to Id1/GFP- cells

(Additional File2: Figure S2 D, E). Finally, we performed the LDT assay wherein we

observed more than 40-fold increase in the metastatic propagating capacity of the Id1/GFP+

cells as compared to the control (p= 0.000161) (Additional File2: Figure S2 F, G).

We next used an inducible lentiviral shRNA system (42) that permits reversible knock down

of Id1 and Id3 in response to doxycycline (Dox) treatment in 4T1 cells. Two clonal 4T1 cell

lines, K1 and K2 were chosen along with a control line (C), based on the efficiency of Id

knock down (Figure 2B, Additional File3: Figure S3A). Id depletion resulted in a significant

decrease in cell proliferation and migration *in vitro* when compared to the control (Additional

File3: Figure S3 B, C, D).

We next interrogated the effect of Id depletion on the self-renewal capacity of the C, K1 and

K2 cell lines. Dox-dependent shRNA induction significantly reduced the ability of the K1

and K2 cells to form primary tumorspheres in the suspension culture (Figure 2C). This effect

was not observed in the control cell line (C; Figure 2C). A significant further decrease in self-

renewal capacity of K1 and K2 lines was observed when primary tumorspheres were

passaged to the secondary stage (Figure 2D, E). The Id depleted tumorspheres were also

markedly smaller in size compared to controls (Figure 2E, Additional File3: Figure S3E).

To assess if the self-renewal phenotype controlled by Id is reversible, we firstly passaged

primary tumorspheres [previously treated with Dox (K+)] to secondary tumorspheres. The

secondary tumorspheres were then cultured in the presence or absence of Dox, to maintain

the Id knockdown status or to allow the re-expression of Id, respectively (Additional File3:

Figure S3F, G). The secondary tumorspheres cultured without Dox (K1+-) re-established

their self-renewal capacity as evidenced by the ability to form new tumorspheres (Figure 2F;

Additional File3: Figure S3H, I), suggesting that Id depletion does not lead to a permanent

loss of self-renewal capacity.

To determine whether Id1 and Id3 are required for primary tumor and metastatic growth *in*

vivo, K1 cells were orthotopically transplanted into the mammary fat pad of BALB/c mice.

Dox-mediated knockdown of Id resulted in modest inhibition of primary tumor growth, with

control tumors growing faster and reaching the ethical endpoint earlier than the Id

knockdown group (Figure 2G). More significantly, mice transplanted with Id depleted K1

cells presented far fewer lung metastatic lesions compared to the control despite growing in

the host for a longer time (p<0.0001; Figure 2H).

To assess the role for Id in metastatic progression *in vivo*, we examined Id expression in lung

metastasis compared to primary tumors in mice injected with K1 cells. An increase in the

expression of Id1 was observed in the lung metastasis in all the samples, while no significant

enrichment of Id3 expression was observed (Additional File4: Figure S4A). To determine

whether altered expression patterns of ID1 are associated with metastasic progression in

patients, ID1 IHC was performed on a cohort of 49 cases with matching primary tumor and

brain metastatic lesions surgically removed from breast cancer patients. Amongst the 13

cases in which ID1 was detected by IHC in the primary tumor, an enrichment of ID1

expression was observed in brain metastases over the patient-matched primary tumor in 11

cases (Additional File4: Figure S4B, Additional File 6: Table S1). Together with data from

the animal model, this result suggests that ID1 promotes metastatic dissemination in a subset

of human breast cancers.

Identification of genes and pathways regulated by Id

The canonical role for Id proteins is to regulate gene expression through association with

transcription factors, yet a comprehensive analysis of Id transcriptional targets in cancer has

not been reported. We performed gene expression profiling of Control (C) and Id depleted K1

cells. The gene expression profiles of four independent replicates (R1, R2, R3 and R4 $\pm$

doxycycline treatment) were compared by microarray analysis (Additional File 5: Figure

S5A). 6081 differentially expressed genes were identified (Q<0.05), with 3310 up-regulated

and 2771 down-regulated genes in Id KD cells (Additional File 7: Table S2) shows the top 25

differentially regulated genes). Network and pathway enrichment analysis was conducted

using the MetaCore^{TM} software. 4301 significant network objects were identified for the Id

knockdown microarray data (adjusted p-value of $\leq$0.05). The top pathways affected by Id

knockdown were mostly associated with the cell cycle (Figure 3A, B) consistent with the loss

of proliferative phenotype described previously (Additional File 3: Figure S3B, C). Similar

results were obtained using Gene Set Enrichment Analysis (GSEA) with significant down

regulation of proliferative signatures (CELL_CYCLE_PROCESS) and mitosis (M_PHASE)

(Additional File 8: Table S3). Genes such as CCNA2, CHEK1 and PLK1 in these gene sets

are down-regulated by Id knockdown. This is consistent with our results (Additional File 3:

Figure S3 B, C) showing Id proteins are necessary for proliferation of 4T1 cells, as well as

previous studies which reported a role of Id in controlling cell cycle progression and

proliferation pathways (17, 43). Enrichment for genes involved in several oncogenic

pathways such as Mek, Vegf, Myc and Bmi1 signalling have also been highlighted

(Additional File 9: Table S4). In order to identify whether Id specifically regulate genes

controlling breast cancer metastasis, GSEA analysis was performed with a collection of

custom "metastasis gene sets". This collection (Table 1) consists of several metastatic

signatures from the C2 collection (MSigDB database; Additional File 10: Table S5),

combined with a list of custom gene sets described in major studies (44-53) as shown in

Figure 3C. Genes differentially expressed in this set included *Robo1* (54, 55), *Il6* (56),

Fermt1 (57), *Foxc2* (58) and Mir30a (59). Three putative Id targets *Robo1, Fermt1* and

Mir30a were then validated using q-RT PCR (Figure 3D) and found to be differentially

regulated in the K1 cell line upon Id KD.

363 Id mediated inhibition of Robo1 controls the proliferative phenotype via activation of

364 Myc transcription

Since Robo1 is known to have a tumor suppressor role in breast cancer biology (54, 60), we

next sought to determine if Robo1 has an epistatic interaction with Id loss of function using

siRNA mediated knockdown of Robo1 followed by proliferation assays. Knockdown of

Robo1 ameliorated the requirement for Id and rescued approximately 55 % of the

proliferative decrease induced by Id KD (Figure 4A).

To understand the mechanisms by which *Robo1* increases the proliferative potential of Id

depleted cells *in vitro*, we performed RNA-Sequencing (RNA-Seq) experiments on K1 cells

with dox-inducible Id KD and/or Robo1 depletion using siRNA. Four replicates per condition

were generated and MDS plots presented in Additional File 5: Figure S5B showed that the

replicates cluster together. Id KD alone in the K1 cells down regulated 4409 genes and up

regulated 5236 genes (FDR<0.05), respectively. The majority of the differentially expressed

genes determined by microarray were found by RNA-Seq analysis (Additional File 5: Figure

S5C). Id depletion led to an increase in *Robo1* expression, as observed in the previous

microarray experiment (Figure 3C, D; Figure 4 B).

Given that Id repressed Robo1 expression, we sought to determine Robo1 target genes in the

absence of Id. Remarkably, under Id depletion conditions, *Robo1* KD restored expression of a

large subset (~45%) of Id target genes to basal levels (Figure 4C). In comparison, knockdown

of Id or Robo1 regulated few targets in the same direction (e.g. both up or both down). This

implies that a large proportion of Id targets may be regulated via suppression of Robo1.

Genes whose expression was repressed by Id KD and rescued by concomitant Robo1 KD

were termed 'Intersect 1' (Figure 4C, Table 2). Genes that were upregulated by Id KD and

downregulated by Robo1 KD (in the absence of Id) were annotated 'intersect 2' (Figure 4C,

Table 3). To investigate the function of these intersect group of genes, we performed GSEA

analysis using the MSigDB hallmark gene set (61). The top signatures in Intersect 1 were

involved in cell proliferation, with enrichment for G2M checkpoint, E2F and Myc targets as

well as mTOR signalling (Table 2). Rank-based analysis revealed strong negative enrichment

for the hallmark Myc targets signature upon Id knockdown alone, and strong positive

enrichment upon Id and Robo1 knockdown (Figure 4D). This suggests that following Id KD,

Robo1 is induced and exerts anti-proliferative effects via suppression of Myc and its target

genes (Additional File 5: Figure S5D, E). Transcription factor motif analysis using EnrichR

revealed that Myc and its binding partner Max, have a high combined score in the Intersect 1

gene list further implicating Myc as downstream effector of Robo1 and Id (Figure 4 E).

We were interested in investigating the possibility that Robo1 may exert its negative effects

on the Myc pathway via regulation of Myc co-factors, which can potently enhance or

suppress Myc transcriptional activity (62). In order to test this hypothesis, we looked at

known Myc co-factors from the literature in our RNA-Seq data to determine if they were

differentially expressed in the Id1 and Robo1 KD conditions. As seen in Additional File 11:

Table S6, we included negative (red) and positive (green) cofactors in the analysis. Scrutiny

of this list suggests that there are numerous negative co-factors (7/10) being induced and

activators being repressed (13/24) by Robo1. For example, putative activation of the gene

Rlim which is an E3 ubiquitin ligase that suppresses the transcriptional activity of MYC (62).

In summary, we have demonstrated that Id depletion leads to a loss in the proliferative and

self-renewal cancer stem cell phenotypes associated with TNBC. Id1 acts by negatively

regulating Robo1 which in turn finally leads to the downstream activation of a Myc

transcriptional program (Figure 5).

Discussion

There is increasing evidence that all cells within a tumor are not equal with some cells having

the plasticity to adapt and subvert cellular and molecular mechanisms to be more tumorigenic

than others. In this study, we demonstrate that Id1 and its closely related family member Id3

are important for the CSC phenotype in the TNBC subtype. Using four independent models

of Id expression and depletion, we demonstrate that the properties of proliferation and self-

renewal are regulated by Id proteins.

Transcription factors like the Id family of proteins can affect a number of key molecular

pathways, allowing switching of phenotypes in response to local cues such as transforming

growth factor-β (TGF-β) (12, 66), receptor tyrosine kinase signalling (67), and steroid

hormones (68) and therefore are able to transduce a multitude of cues into competency for

proliferation and self-renewal. The CSC phenotype as marked by Id is plastic, fitting with the

latest evidence that CSC are not necessarily hierarchically organised, but rather represent a

transient inducible state dependent on the local microenvironment.

We report the first comprehensive analysis of Id transcriptional targets. We go on to identify

a novel epistatic relationship with Robo1, with Robo1 loss sufficient to remove the necessity

for Id in proliferation, suggesting that suppression of Robo1 is an important function for Id in

this setting. Robo1 is a receptor for SLIT1 and SLIT2 that mediates cellular responses to

molecular guidance cues in cellular migration (69). Previous work with mammary stem cells

showed that the extracellular SLIT2 signals via ROBO1 to regulate the asymmetric self-

renewal of basal stem cells through the transcription factor Snail during mammary gland

development (70). Our finding may have significant implications for tumor biology because

SLIT/ROBO signalling is altered in about 40% of basal breast tumors (70). Our work

implicates a novel role for SLIT-ROBO signalling in CSC and shows a new mechanism by

which Id proteins control the self-renewal phenotype by suppressing the Robo1 tumor

suppressor role in TNBC.

The significant decrease in the Myc levels on Id knockdown suggest an Id/ Robo1/ Myc axis

in TNBC (Additional File 5: Figure S5D, E). While the proposed model for regulation of

Myc is not yet clear, we propose two possible modes of regulation of Myc: (1) Robo

independent suppression of Myc expression and (2) Robo dependent regulation of Myc

activity. Though the mechanism still needs to be elaborated, we hypothesise that in the

absence of Id, Robo1 inhibits Myc activity via activation of Myc inhibitors (e.g. Rlim) and/or

inhibition of Myc activators (e.g. Aurka). This is borne out by the analysis of Myc co-factors

in the Id and Id Robo1 KD RNA Seq data (Additional File 11: Table S6). Further work is

needed to determine whether, and which, Myc cofactors are epistatic to Id-Robo1 signalling.

Our data provides further evidence that Robo1 is an important suppressor of proliferation and

self-renewal in TNBC and future work includes extending this work to models of human

TNBC. Prior work showing high Robo1 expression association with good outcome in breast

cancer is consistent with our finding (54). There has been substantial interest in targeting

Myc (60, 71) and Id1, but until now has been very challenging (20, 72). We show that Id1 is

able to reprogram Myc activity possibly via Robo1 and may provide an alternative strategy to

target Myc-dependent transcription.

Conclusion

We have demonstrated that breast cancer cells marked by Id expression have high propensity

for key CSC phenotypes like proliferation and metastasis. We have uncovered a set of genes

that are potential Id targets leading to identification of a mechanism which involves the

negative transcriptional regulation of Robo1 by Id. This suggests an association between Id

and Robo1 that correlates to the activation of a cMyc driven proliferative and self renewal

program. Our observations suggest that we could exploit this pathway to target CSCs in the

difficult to treat TNBC subtype.

Abbreviations

CSC : Cancer stem cell

TNBC: Triple Negative Breast Cancer

Id : Id1 and Id3

References

1.   Chen J, Li Y, Yu TS, McKay RM, Burns DK, Kernie SG, et al. A restricted cell

population propagates glioblastoma growth after chemotherapy. Nature.

2012;488(7412):522-6.

2.   Lawson DA, Bhakta NR, Kessenbrock K, Prummel KD, Yu Y, Takai K, et al. Single-

cell analysis reveals a stem-cell program in human metastatic breast cancer cells. Nature.

2015;526(7571):131-5.

3.   Li X, Lewis MT, Huang J, Gutierrez C, Osborne CK, Wu MF, et al. Intrinsic

resistance of tumorigenic breast cancer cells to chemotherapy. J Natl Cancer Inst.

2008;100(9):672-9.

4.   Malanchi I, Santamaria-Martinez A, Susanto E, Peng H, Lehr HA, Delaloye JF, et al.

Interactions between cancer stem cells and their niche govern metastatic colonization. Nature.

2011;481(7379):85-9.

5.   da Silva-Diz V, Lorenzo-Sanz L, Bernat-Peguera A, Lopez-Cerda M, Munoz P.

Cancer cell plasticity: Impact on tumor progression and therapy response. Seminars in cancer

biology. 2018.

6.   Lee G, Hall RR, 3rd, Ahmed AU. Cancer Stem Cells: Cellular Plasticity, Niche, and

its Clinical Relevance. J Stem Cell Res Ther. 2016;6(10).

7.   Wahl GM, Spike BT. Cell state plasticity, stem cells, EMT, and the generation of

intra-tumoral heterogeneity. NPJ Breast Cancer. 2017;3:14.

8.   Lasorella A, Benezra R, Iavarone A. The ID proteins: master regulators of cancer

stem cells and tumour aggressiveness. Nature reviews Cancer. 2014;14(2):77-91.

9.   Aloia L, Gutierrez A, Caballero JM, Di Croce L. Direct interaction between Id1 and

Zrf1 controls neural differentiation of embryonic stem cells. EMBO Rep. 2015;16(1):63-70.

10.   Hong SH, Lee JH, Lee JB, Ji J, Bhatia M. ID1 and ID3 represent conserved negative

regulators of human embryonic and induced pluripotent stem cell hematopoiesis. J Cell Sci.

2011;124(Pt 9):1445-52.

11.   Liang YY, Brunicardi FC, Lin X. Smad3 mediates immediate early induction of Id1

by TGF-beta. Cell Res. 2009;19(1):140-8.

12.   Stankic M, Pavlovic S, Chin Y, Brogi E, Padua D, Norton L, et al. TGF-beta-Id1

signaling opposes Twist1 and promotes metastatic colonization via a mesenchymal-to-

epithelial transition. Cell reports. 2013;5(5):1228-42.

13.   Anido J, Saez-Borderias A, Gonzalez-Junca A, Rodon L, Folch G, Carmona MA, et

al. TGF-beta Receptor Inhibitors Target the CD44(high)/Id1(high) Glioma-Initiating Cell

Population in Human Glioblastoma. Cancer cell. 2010;18(6):655-68.

14.   Gupta GP, Perk J, Acharyya S, de Candia P, Mittal V, Todorova-Manova K, et al. ID

genes mediate tumor reinitiation during breast cancer lung metastasis. Proceedings of the

National Academy of Sciences of the United States of America. 2007;104(49):19506-11.

15.   Lyden D, Young AZ, Zagzag D, Yan W, Gerald W, O'Reilly R, et al. Id1 and Id3 are

required for neurogenesis, angiogenesis and vascularization of tumour xenografts. Nature.

1999;401(6754):670-7.

16.   Niola F, Zhao X, Singh D, Sullivan R, Castano A, Verrico A, et al. Mesenchymal

high-grade glioma is maintained by the ID-RAP1 axis. The Journal of clinical investigation.

2013;123(1):405-17.

17.   O'Brien CA, Kreso A, Ryan P, Hermans KG, Gibson L, Wang Y, et al. ID1 and ID3

regulate the self-renewal capacity of human colon cancer-initiating cells through p21. Cancer

cell. 2012;21(6):777-92.

18.   Swarbrick A, Roy E, Allen T, Bishop JM. Id1 cooperates with oncogenic Ras to

induce metastatic mammary carcinoma by subversion of the cellular senescence response.

Proceedings of the National Academy of Sciences of the United States of America. 2008;105(14):5402-7.

19.	Swarbrick A, Akerfeldt MC, Lee CS, Sergio CM, Caldon CE, Hunter LJ, et al. Regulation of cyclin expression and cell cycle progression in breast epithelial cells by the helix-loop-helix protein Id1. Oncogene. 2005;24(3):381-9.

20.	Fong S, Itahana Y, Sumida T, Singh J, Coppe JP, Liu Y, et al. Id-1 as a molecular target in therapy for breast cancer cell invasion and metastasis. Proceedings of the National Academy of Sciences of the United States of America. 2003;100(23):13543-8.

21.	Gumireddy K, Li A, Gimotty PA, Klein-Szanto AJ, Showe LC, Katsaros D, et al. KLF17 is a negative regulator of epithelial-mesenchymal transition and metastasis in breast cancer. Nat Cell Biol. 2009;11(11):1297-304.

22.	Tobin NP, Sims AH, Lundgren KL, Lehn S, Landberg G. Cyclin D1, Id1 and EMT in breast cancer. BMC cancer. 2011;11:417.

23.	Kim H, Chung H, Kim HJ, Lee JY, Oh MY, Kim Y, et al. Id-1 regulates Bcl-2 and Bax expression through p53 and NF-kappaB in MCF-7 breast cancer cells. Breast Cancer Res Treat. 2008;112(2):287-96.

24.	Qian T, Lee JY, Park JH, Kim HJ, Kong G. Id1 enhances RING1b E3 ubiquitin ligase activity through the Mel-18/Bmi-1 polycomb group complex. Oncogene. 2010;29(43):5818-27.

25.	Mellick AS, Plummer PN, Nolan DJ, Gao D, Bambino K, Hahn M, et al. Using the transcription factor inhibitor of DNA binding 1 to selectively target endothelial progenitor cells offers novel strategies to inhibit tumor angiogenesis and growth. Cancer research. 2010;70(18):7273-82.

26.	Lee CH, Yu CC, Wang BY, Chang WW. Tumorsphere as an effective in vitro platform for screening anti-cancer stem cell drugs. Oncotarget. 2016;7(2):1215-26.

27.   Pastrana E, Silva-Vargas V, Doetsch F. Eyes wide open: a critical review of sphere-

formation as an assay for stem cells. Cell Stem Cell. 2011;8(5):486-98.

28.   Nair R, Roden DL, Teo WS, McFarland A, Junankar S, Ye S, et al. c-Myc and Her2

cooperate to drive a stem-like phenotype with poor prognosis in breast cancer. Oncogene.

2014;33(30):3992-4002. Epub 2013/09/24.

29.   Green JE, Shibata MA, Yoshidome K, Liu ML, Jorcyk C, Anver MR, et al. The

C3(1)/SV40 T-antigen transgenic mouse model of mammary cancer: ductal epithelial cell

targeting with multistage progression to carcinoma. Oncogene. 2000;19(8):1020-7.

30.   Pfefferle AD, Herschkowitz JI, Usary J, Harrell JC, Spike BT, Adams JR, et al.

Transcriptomic classification of genetically engineered mouse models of breast cancer

identifies human subtype counterparts. Genome biology. 2013;14(11):R125.

31.   Perry SS, Zhao Y, Nie L, Cochrane SW, Huang Z, Sun XH. Id1, but not Id3, directs

long-term repopulating hematopoietic stem-cell maintenance. Blood. 2007;110(7):2351-60.

32.   Herschkowitz JI, Zhao W, Zhang M, Usary J, Murrow G, Edwards D, et al.

Comparative oncogenomics identifies breast tumors enriched in functional tumor-initiating

cells. Proceedings of the National Academy of Sciences of the United States of America.

2012;109(8):2778-83.

33.   Zhang M, Behbod F, Atkinson RL, Landis MD, Kittrell F, Edwards D, et al.

Identification of tumor-initiating cells in a p53-null mouse model of breast cancer. Cancer

research. 2008;68(12):4674-82.

34.   Vaillant F, Asselin-Labat ML, Shackleton M, Forrest NC, Lindeman GJ, Visvader JE.

The mammary progenitor marker CD61/beta3 integrin identifies cancer stem cells in mouse

models of mammary tumorigenesis. Cancer research. 2008;68(19):7711-7.

35.   Konrad CV, Murali R, Varghese BA, Nair R. The role of cancer stem cells in tumor

heterogeneity and resistance to therapy. Can J Physiol Pharmacol. 2017;95(1):1-15.

36.   Yoneda T, Michigami T, Yi B, Williams PJ, Niewolna M, Hiraga T. Actions of

bisphosphonate on bone metastasis in animal models of breast carcinoma. Cancer.

2000;88(12 Suppl):2979-88.

37.   Lelekakis M, Moseley JM, Martin TJ, Hards D, Williams E, Ho P, et al. A novel

orthotopic model of breast cancer metastasis to bone. Clin Exp Metastasis. 1999;17(2):163-

70.

38.   Tao K, Fang M, Alroy J, Sahagian GG. Imagable 4T1 model for the study of late

stage breast cancer. BMC cancer. 2008;8:228.

39.   Aslakson CJ, Miller FR. Selective events in the metastatic process defined by analysis

of the sequential dissemination of subpopulations of a mouse mammary tumor. Cancer

research. 1992;52(6):1399-405.

40.   Pulaski BA, Ostrand-Rosenberg S. Reduction of established spontaneous mammary

carcinoma metastases following immunotherapy with major histocompatibility complex class

II and B7.1 cell-based tumor vaccines. Cancer research. 1998;58(7):1486-93.

41.   Eckhardt BL, Parker BS, van Laar RK, Restall CM, Natoli AL, Tavaria MD, et al.

Genomic analysis of a spontaneous model of breast cancer metastasis to bone reveals a role

for the extracellular matrix. Mol Cancer Res. 2005;3(1):1-13.

42.   Shin KJ, Wall EA, Zavzavadjian JR, Santat LA, Liu J, Hwang JI, et al. A single

lentiviral vector platform for microRNA-based conditional RNA interference and coordinated

transgene expression. Proceedings of the National Academy of Sciences of the United States

of America. 2006;103(37):13759-64.

43.   Nair R, Teo WS, Mittal V, Swarbrick A. ID proteins regulate diverse aspects of

cancer progression and provide novel therapeutic opportunities. Molecular therapy : the

journal of the American Society of Gene Therapy. 2014;22(8):1407-15.

44.   Aceto N, Bardia A, Miyamoto DT, Donaldson MC, Wittner BS, Spencer JA, et al.

Circulating tumor cell clusters are oligoclonal precursors of breast cancer metastasis. Cell.

2014;158(5):1110-22.

45.   Bos PD, Zhang XH, Nadal C, Shu W, Gomis RR, Nguyen DX, et al. Genes that

mediate breast cancer metastasis to the brain. Nature. 2009;459(7249):1005-9.

46.   Charafe-Jauffret E, Ginestier C, Iovino F, Wicinski J, Cervera N, Finetti P, et al.

Breast cancer cell lines contain functional cancer stem cells with metastatic capacity and a

distinct molecular signature. Cancer research. 2009;69(4):1302-13.

47.   Dontu G, Al-Hajj M, Abdallah WM, Clarke MF, Wicha MS. Stem cells in normal

breast development and breast cancer. Cell proliferation. 2003;36 Suppl 1:59-72.

48.   Kang Y, Siegel PM, Shu W, Drobnjak M, Kakonen SM, Cordon-Cardo C, et al. A

multigenic program mediating breast cancer metastasis to bone. Cancer cell. 2003;3(6):537-

49.

49.   Liu H, Patel MR, Prescher JA, Patsialou A, Qian D, Lin J, et al. Cancer stem cells

from human breast tumors are involved in spontaneous metastases in orthotopic mouse

models. Proceedings of the National Academy of Sciences of the United States of America.

2010;107(42):18115-20.

50.   Minn AJ, Gupta GP, Siegel PM, Bos PD, Shu W, Giri DD, et al. Genes that mediate

breast cancer metastasis to lung. Nature. 2005;436(7050):518-24.

51.   Minn AJ, Kang Y, Serganova I, Gupta GP, Giri DD, Doubrovin M, et al. Distinct

organ-specific metastatic potential of individual breast cancer cells and primary tumors. The

Journal of clinical investigation. 2005;115(1):44-55.

52.   Padua D, Zhang XH, Wang Q, Nadal C, Gerald WL, Gomis RR, et al. TGFbeta

primes breast tumors for lung metastasis seeding through angiopoietin-like 4. Cell.

2008;133(1):66-77.

53.   Tang B, Yoo N, Vu M, Mamura M, Nam JS, Ooshima A, et al. Transforming growth

factor-beta can suppress tumorigenesis through effects on the putative cancer stem or early

progenitor cell and committed progeny in a breast cancer xenograft model. Cancer research.

2007;67(18):8643-52.

54.   Chang PH, Hwang-Verslues WW, Chang YC, Chen CC, Hsiao M, Jeng YM, et al.

Activation of Robo1 signaling of breast cancer cells by Slit2 from stromal fibroblast restrains

tumorigenesis   via   blocking   PI3K/Akt/beta-catenin   pathway.   Cancer   research.

2012;72(18):4652-61.

55.   Qin F, Zhang H, Ma L, Liu X, Dai K, Li W, et al. Low Expression of Slit2 and Robo1

is Associated with Poor Prognosis and Brain-specific Metastasis of Breast Cancer Patients.

Scientific reports. 2015;5:14430.

56.   Chang Q, Bournazou E, Sansone P, Berishaj M, Gao SP, Daly L, et al. The IL-

6/JAK/Stat3   feed-forward   loop   drives   tumorigenesis   and   metastasis.   Neoplasia.

2013;15(7):848-62.

57.   Landemaine T, Jackson A, Bellahcene A, Rucci N, Sin S, Abad BM, et al. A six-gene

signature predicting breast cancer lung metastasis. Cancer research. 2008;68(15):6092-9.

58.   Mani SA, Yang J, Brooks M, Schwaninger G, Zhou A, Miura N, et al. Mesenchyme

Forkhead 1 (FOXC2) plays a key role in metastasis and is associated with aggressive basal-

like breast cancers. Proceedings of the National Academy of Sciences of the United States of

America. 2007;104(24):10069-74.

59.   Zhang N, Wang X, Huo Q, Sun M, Cai C, Liu Z, et al. MicroRNA-30a suppresses

breast tumor growth and metastasis by targeting metadherin. Oncogene. 2014;33(24):3119-

28.

60.   Shen L, O'Shea JM, Kaadige MR, Cunha S, Wilde BR, Cohen AL, et al. Metabolic

reprogramming in triple-negative breast cancer through Myc suppression of TXNIP.

Proceedings of the National Academy of Sciences of the United States of America.

2015;112(17):5425-30.

61.   Liberzon A, Birger C, Thorvaldsdottir H, Ghandi M, Mesirov JP, Tamayo P. The

Molecular Signatures Database (MSigDB) hallmark gene set collection. Cell systems.

2015;1(6):417-25.

62.   Gao R, Wang L, Cai H, Zhu J, Yu L. E3 Ubiquitin Ligase RLIM Negatively

Regulates c-Myc Transcriptional Activity and Restrains Cell Proliferation. PloS one.

2016;11(9):e0164086.

63.   Cancer Genome Atlas Research N, Weinstein JN, Collisson EA, Mills GB, Shaw KR,

Ozenberger BA, et al. The Cancer Genome Atlas Pan-Cancer analysis project. Nature

genetics. 2013;45(10):1113-20.

64.   Gao J, Aksoy BA, Dogrusoz U, Dresdner G, Gross B, Sumer SO, et al. Integrative

analysis of complex cancer genomics and clinical profiles using the cBioPortal. Science

signaling. 2013;6(269):pl1.

65.   Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, et al. The cBio

cancer genomics portal: an open platform for exploring multidimensional cancer genomics

data. Cancer discovery. 2012;2(5):401-4.

66.   Kang Y, Chen CR, Massague J. A self-enabling TGFbeta response coupled to stress

signaling: Smad engages stress response factor ATF3 for Id1 repression in epithelial cells.

Molecular cell. 2003;11(4):915-26.

67.   Tam WF, Gu TL, Chen J, Lee BH, Bullinger L, Frohling S, et al. Id1 is a common

downstream target of oncogenic tyrosine kinases in leukemic cells. Blood. 2008;112(5):1981-

92.

68.   Lin CQ, Singh J, Murata K, Itahana Y, Parrinello S, Liang SH, et al. A role for Id-1 in

the aggressive phenotype and steroid hormone response of human breast cancer cells. Cancer

research. 2000;60(5):1332-40.

69.   Huang T, Kang W, Cheng AS, Yu J, To KF. The emerging role of Slit-Robo pathway

in gastric and other gastro intestinal cancers. BMC cancer. 2015;15:950.

70.   Ballard MS, Zhu A, Iwai N, Stensrud M, Mapps A, Postiglione MP, et al. Mammary

Stem Cell Self-Renewal Is Regulated by Slit2/Robo1 Signaling through SNAI1 and mINSC.

Cell reports. 2015;13(2):290-301.

71.   Yang A, Qin S, Schulte BA, Ethier SP, Tew KD, Wang GY. MYC Inhibition

Depletes Cancer Stem-like Cells in Triple-Negative Breast Cancer. Cancer research.

2017;77(23):6641-50.

72.   Dang CV, Reddy EP, Shokat KM, Soucek L. Drugging the 'undruggable' cancer

targets. Nature reviews Cancer. 2017;17(8):502-8.

73.   Dull T, Zufferey R, Kelly M, Mandel RJ, Nguyen M, Trono D, et al. A third-

generation lentivirus vector with a conditional packaging system. Journal of virology.

1998;72(11):8463-71.

74.   Irizarry RA, Bolstad BM, Collin F, Cope LM, Hobbs B, Speed TP. Summaries of

Affymetrix GeneChip probe level data. Nucleic acids research. 2003;31(4):e15.

75.   Smyth GK. Linear models and empirical bayes methods for assessing differential

expression in microarray experiments. Statistical applications in genetics and molecular

biology. 2004;3:Article3.

76.   Reich M, Liefeld T, Gould J, Lerner J, Tamayo P, Mesirov JP. GenePattern 2.0.

Nature genetics. 2006;38(5):500-1.

77.   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al.

Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide

expression profiles. Proceedings of the National Academy of Sciences of the United States of

America. 2005;102(43):15545-50.

78.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR:

ultrafast universal RNA-seq aligner. Bioinformatics. 2013;29(1):15-21.

79.   Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with

or without a reference genome. BMC bioinformatics. 2011;12:323.

80.   Robinson MD, Oshlack A. A scaling normalization method for differential expression

analysis of RNA-seq data. Genome biology. 2010;11(3):R25.

81.   Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers

differential expression analyses for RNA-sequencing and microarray studies. Nucleic acids

research. 2015;43(7):e47.

701 Declarations

702 Ethics approval and consent to participate

703 All experiments involving animal work were performed in accordance with the rules and

704 regulations stated by the Garvan Institute Animal Ethics Committee.

705 Consent for publication

706 All authors have given their consent for publication.

707 Availability of supporting data

708 All microarray and RNA Sequencing data has been deposited into the GEO database.

709 Microarray (GSE129790) and RNA-Seq (GSE129858) datasets are available in SuperSeries

710 GSE129859. Supplementary information is available in the Supplementary files.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by funding from the National Health and Medical

Research Council (NHMRC) of Australia, The National Breast Cancer Foundation and John

and Deborah McMurtrie and in part by Early Career Research (ECR) Award from Science

and Engineering Research Board (SERB), Government of India (ECR/2015/000031). A.S. is

the recipient of a Senior Research Fellowship from the NHMRC. S.O.T. is supported by

NBCF practitioner fellowship and also acknowledges the Sydney Breast Cancer Foundation,

the Tag family, Mr David Paradice, ICAP and the O'Sullivan family and the estate of the late

Kylie Sinclair. RN is the recipient of the Ramanujan Fellowship from the Government of

India (SERB) (SB/S2/RJN/182/2014). WST is funded by International Postgraduate Research

Scholarship and the Beth Yarrow Memorial Award in Medical Science. APT and BAV is

funded through CSIR- Junior Research Fellowship, and DST- INSPIRE Fellowship,

respectively.

Author's contributions

RN and AS contributed to the conceptualizaton. WST, HH ASC, KH and SVK contributed to

the methodology. DLR and CLC to the genomic analysis. RM, BAV, APT, AM, SJ, JY, IN,

JSS and LAB contributed to the investigations. EKAM and SAO carried out the analysis on

patient samples. ASM contributed to the resources. RN and NK wrote the original draft of the

manuscript. MJN, CJO, SRL and WK reviewed, and edited the manuscript. NK contributed

to the visualization. AS and RN contributed to the funding acquisition. All authors read and

approved the final manuscript.

Acknowledgements

We would like to thank the following people for their assistance with this manuscript:

Ms. Nicola Foreman and Ms. Breanna Fitzpatrick for animal handling; Ms. Alice

Boulghourjian and Ms. Anaiis Zaratzian for tissue processing and IHC staining; Mr. Rob

Salomon and Mr. David Snowden for their help with flow sorting.

Author information

Radhika Nair and Alexander Swarbrick are co-senior authors.

Correspondence to radhikanair@rgcb.res.in or a.swarbrick@garvan.org.au

Affiliations

Garvan Institute of Medical Research, Darlinghurst, New South Wales, 2010, Australia

Wee S. Teo, Holly Holliday, Aurélie S. Cazet, Daniel L. Roden, Kate Harvey, Christina

Valbirk Konrad, Chia-Ling Chan , Andrea McFarland, Simon Junankar, Sunny Ye, Jessica

Yang, Iva Nikolic, Laura A. Baker, Ewan K.A. Millar, Mathew J. Naylor, Christopher J.

Ormandy, Warren Kaplan, Sandra A. O'Toole, Alexander Swarbrick, Radhika Nair

St Vincent's Clinical School, Faculty of Medicine, UNSW Sydney, New South Wales, 2010,

Australia

Wee S. Teo, Holly Holliday, Aurélie S. Cazet, Daniel L. Roden, Chia-Ling Chan, Simon

Junankar, Iva Nikolic, Laura A. Baker , Mathew J. Naylor, Christopher J. Ormandy,

Alexander Swarbrick, Radhika Nair

Cancer Research Program, Rajiv Gandhi Centre for Biotechnology, Kerala, 695014, India

Nitheesh Karthikeyan, Reshma Murali, Binitha Anu Varghese, Archana P. T., Radhika Nair

Manipal Academy of Higher Education, Manipal, Karnataka, 576104, India

Archana P. T.

Centenary Institute, The University of Sydney, New South Wales, 2050, Australia

Jaynish S. Shah

Department of Anatomical Pathology, NSW Health Pathology, St George Hospital, Kogarah NSW 2217 Australia

Ewan K.A. Millar

School of Medical Sciences, UNSW Sydney, Kensington NSW 2033, Australia.

Ewan K.A. Millar

School of Medicine, Western Sydney University, Locked Bag 1797, Penrith, NSW 2751, Australia

Ewan K.A. Millar

Department of Tissue Pathology and Diagnostic Oncology, Royal Prince Alfred Hospital, Camperdown

Sandra A. O'Toole

Discipline of Physiology & Bosch Institute, School of Medical Sciences, University of Sydney, NSW 2006

Mathew J. Naylor

The University of Queensland, UQ Centre for Clinical Research, School of Medicine and Pathology Queensland, The Royal Brisbane & Women's Hospital, Herston, Brisbane, Queensland 4029, Australia

Sunil R. Lakhani

Peter Wills Bioinformatics Centre, Garvan Institute of Medical Research, Darlinghurst, NSW

2010, Australia

Warren Kaplan

UNSW Medicine, University of NSW, 18 High St, Kensington NSW 2052, Australia.

Albert S. Mellick

Ingham Institute for Applied Medical Research, 1 Campbell Street, Liverpool, NSW 2170,

Australia.

Albert S. Mellick

Figure legends

Figure 1. Id1 marks tumor cells with high self-renewal in murine models of TNBC.

A. Representative IHC images of Id1 and Id3 expression in p53-/- tumor model. Black arrows

in the inset indicate Id1+ cells. Scale bars = 50μm. **B.** p53-/- tumor cells were transfected

with the Id1/GFP reporter and subsequently sorted for GFP expression. The self-renewal

capacity of Id1/GFP$^+$ p53-/- cells was significantly higher than unsorted Id1/GFP p53-/- cells

upon passage to tertiary tumorspheres. Data are means ± SD (n=3). (*** $p< 0.001$; Two-way

ANOVA). **C.** Id1 expressing cells were sorted from the p53-/- Id1/GFP tumor model and

transplanted into recipient mice by limiting dilution assay. Based on limiting dilution

calculations (ELDA), the Id1$^+$ cells demonstrated a 4.6-fold enrichment in tumor initiating

capacity (TIC) when compared to the Id1$^-$ cells in serial passage. p values for p53-/- Id1GFP+

versus Unsorted Round 1- 0.2920, p53-/- Id1GFP+ versus Id1GFP- Round 2- 0.0221. **D.**

Representative IHC images of the Id1C3-Tag model, confirming its suitability as a model

system. Black arrows in the inset indicate Id1+ cells. Expression of Id1 was less than 5% as

determined by IHC. Bars = 50μm. **E.** Tumor cells from the Id1C3-Tag tumor model were

FACS sorted based on their GFP expression. qRT-PCR analyses on the sorted GFP$^+$ and

GFP$^-$ cell populations showed a significant increase (more than 5-fold) for Id1 expression in

the GFP$^+$ cells compared to cells lacking GFP expression. **F.** *In vitro* self-renewal capacity of

GFP$^+$ cells was measured using the tumorsphere assay. The secondary sphere forming

capacity of Id1$^+$ tumor cells from the Id1C3-Tag model was significantly enriched in

comparison to the Id1$^-$tumor cells. Data are means ± SD (n=3). (**$p< 0.01$, **** $p< 0.0001$;

Two-way ANOVA). **G.** Representative FACS scatterplot and histograms from Id1C3-Tag

tumors showing the expression of the CSC markers CD24, CD29 and CD61 in the Id1$^-$/GFP$^-$

and Id1$^+$/GFP$^+$ cancer cells. Putative CSC populations are highlighted within the red box.

Figure 2. **Depletion of Id1 and Id3 leads to a reduced self-renewal capacity *in vitro* and**

metastatic potential *in vivo*. A. Endogenous levels of Id1 and Id3 expression in 4T1primary

mammary tumors were determined. 4T1 were cells stained for Id1 and Id3 expression

(brown) and counterstained with haematoxylin. Mammary gland tissue from Id1 and Id3 null

(Id1-/- and Id3-/-) mice served as negative controls. Scale bars = 50 μm. Western blot

analysis of protein lysate from 4T1 tumor cells served as positive controls for Id1 and Id3

expression. **B.** Kinetics of conditional Id knockdown in 4T1 cells. Representative Western

blot analysis of Id protein levels in pSLIK K1 cells over time. Cells were cultured in the

presence of 1 μg/ml of Doxycycline (Dox) for 1, 3 and 5 days. β-actin was used as loading

control. **C.** 4T1 Control, pSLIK K1 and K2 clones were assayed for their tumorsphere

forming potential. Dox was added into the culture medium at day 0. Number of primary

tumorspheres formed was quantified by visual examination on day 7. Id knockdown leads to

a decrease in tumorsphere-forming ability of K1 and K2 cell lines. Data are means ± SD

(n=3). (**$p<$ 0.01; Two-way ANOVA). **D.** Primary tumorspheres were passaged and the

number of secondary tumorspheres was quantified on day 14. Knockdown of Id significantly

reduces the ability of the K1 and K2 cells to form secondary tumorspheres in the suspension

culture. Data are means ± SD (n=3). (**$p<$ 0.01; Two-way ANOVA). **E.** Representative

images of primary and secondary tumorsphere formation for the clone K1 ±Dox. **F.**

Quantification and representative images of primary tumorsphere treated with Dox (K1+)

passaged to secondary spheres in Dox free conditions (K1+-) allowing re expression of Id

and restoration of self-renewal capacity. **G.** Knockdown of Id significantly delays tumor

growth in the 4T1 syngeneic model. (*n* = 10 mice; **p value<0.01, Student's *t*-test). **H.** Id

knockdown suppresses spontaneous lung metastasis. Tumors depleted of Id expression

generated fewer spontaneous lung macrometastatic lesions compared to the control despite

growing in the host for a longer time. Inset shows representative images of lungs bearing the

control (K1 - Dox) and Id KD (K1 + Dox) lung metastases at ethical end point. Control; n = 8

mice, Id KD; n=10 mice. Scale bar = 50 um.

Figure 3. Gene expression analysis reveals targets of Id in TNBC

A, B. To characterize the network of genes regulated by Id, functional annotation analyses

were performed on the gene array data from the 4T1 TNBC model. The Id depletion model

attempted to identify downstream targets of Id through a loss of function approach. The gene

expression profile of four independent replicates of the K1 shId clone, with and without

doxycycline treatment, was compared by microarray analysis. This resulted in a list of

differentially expressed genes between control and Id depleted cells, which by further

network and map analysis using Metacore demonstrated was largely driven by genes

controlling cell cycle pathways.

C. Gene expression analysis identified metastasis-related genes that were differentially

expressed in response to Id knockdown. To determine if genes that mediate metastasis were

enriched in the Id signature, gene expression analysis was performed using a manually

curated set of metastasis gene sets. Genes differentially expressed in response to Id

knockdown as well as associated with pathways regulating metastasis were identified based

on reports from the literature which included Robo1.

D. Validation of expression profiling results by quantitative real-time-PCR using the

Taqman® probe based system. Relative mRNA expression of Robo1, Fermt1 and Mir30a, in

the 4T1 pSLIK shId Clonal cell line (K1) and pSLIK control (C), as indicated. Data are

means ± SD (n=3). (**$p< 0.01$, ****$p< 0.0001$; unpaired t-test).

Figure 4. Identification of Myc signature activation by Id via negative regulation of

Robo1

A. Proliferation of K1 cells treated with non-targeting (NT) control siRNA or Robo1 siRNA

in the absence or presence of Doxycycline to induce Id knockdown was measured by the

IncuCyte^{TM} (Essen Instruments) live-cell imaging system. Data shown as mean ±SD (n=3).

(*** p< 0.001; Unpaired two-tailed t-test). **B**. Robo1 expression in Control, Id KD, Robo1

KD and Id Robo1 KD cells was measured by quantitative PCR. Ct values were normalised to

β actin and GAPDH housekeeping genes. Data shown as mean ± SEM (n=4). (** p< 0.01,

**** p < 0.0001; Unpaired two-tailed t-test). **C**. Transcriptional profiling was performed on

Control, Id KD, Robo1 KD and Id Robo1 KD cells. Proportional Venn diagrams (BioVenn)

were generated to visualise the overlapping genes between the different comparisons. **D**.

GSEA Enrichment plots of the hallmark Myc targets version 1 signature from MSigDB. NES

= normalised enrichment score. **E**. Consensus Transcription factor motif analysis using the

Encyclopedia of DNA Elements (ENCODE) and ChIP enrichment analysis (ChEA) data sets

determined using EnrichR. The combined score is a combination of the p-value and z-score.

Figure 5: Model showing the mechanism of Id-Robo1 action in cancer cells.

The proposed model for the regulation of Myc by Id and Robo1. Co-A indicates

representative Myc activator and Co-R indicates representative Myc repressor.

888 **Table 1.** Gene expression signatures of breast cancer metastasis and breast cancer stem cells.

889 This table showed a collection of gene sets which comprised several metastatic signatures

890 that were picked from the C2 collection on the MSigDB database and several other signatures

891 that were manually curated. GSEA analysis was carried out to identify whether any of the

892 Id1/3 targets from the profiling experiment are enriched in these signatures.

Study	Signature	Type	Available on GSEA MSigDB database?
Landemaine et al. A six-gene signature predicting breast cancer lung metastasis. Cancer Res. 2008 Aug 1;68(15):6092-9	Lung metastasis signature of breast cancer	Metastatic tissue tropism	Yes
Bild et al. Oncogenic pathway signatures in human cancers as a guide to targeted therapies. Nature 2006, 439:353-357.	Expression profile of 4 individual genes -- Myc, E2F3, Ras, Src, β-catenin	Signalling pathway	Yes
van 't Veer et al. Gene expression profiling predicts clinical outcome of breast cancer. Nature 2002, 415:530-536.	Poor prognosis signature of breast cancer	Classifier that classifies patients as having good or poor prognosis	Yes
Wang et al. Gene-expression	Poor prognosis	Classifier	Yes

profiles to predict distant metastasis of lymph-node-negative primary breast cancer. Lancet 2005, 365:671-679.	signature of breast cancer		
Ramaswamy et al. A molecular signature of metastasis in primary solid tumors. Nat Genet 2003, 33:49-54.	General metastasis	Classifier	Yes
Finak et al. Stromal gene expression predicts clinical outcome in breast cancer. Nat Med 2008, 14:518-527.	Breast tumor stromal gene expression signature	Classifier	Yes
Farmer et al. A stroma-related gene signature predicts resistance to neoadjuvant chemotherapy in breast cancer. Nat Med 2009, 15:68-74.	Stromal gene expression signature of breast tumor treated with chemotherapy	Classifier	Yes
Kang et al. A multigenic program mediating breast cancer metastasis to bone. Cancer Cell 2003, 3:537-549.	Bone metastasis signature of breast cancer	Metastatic tissue tropism	No
Minn et al. Genes that mediate breast cancer metastasis to lung.	Lung metastasis signature of breast	Metastatic tissue	No

Nature 2005, 436:518-524	cancer	tropism	
Bos et al. Genes that mediate breast cancer metastasis to the brain. Nature 2009, 459:1005-1009.	Bone metastasis signature of breast cancer	Metastatic tissue tropism	No
Padua et al. TGFbeta primes breast tumors for lung metastasis seeding through angiopoietin-like 4. Cell 2008, 133:66-77.	TGF-b signature in lung metastasis of breast cancer	Signalling pathway	No
Aceto et al. Tyrosine phosphatase SHP2 promotes breast cancer progression and maintains tumor-initiating cells via activation of key transcription factors and a positive feedback signaling loop. Nat Med. 2012 Mar 4;18(4):529-37.	Shp2 signature in breast cancer metastasis	Signalling pathway	No
Minn et al. Distinct organ-specific metastatic potential of individual breast cancer cells and primary tumors. J Clin Invest. 2005 Jan;115(1):44-55.	Poor prognosis signature of breast cancer ; Breast cancer metastasis signature; Bone metastasis signature	Metastatic tissue tropism	No

	of breast cancer		
Tang et al. Transforming growth factor-beta can suppress tumorigenesis through effects on the putative cancer stem or early progenitor cell and committed progeny in a breast cancer xenograft model. Cancer Res. 2007 Sep 15;67(18):8643-52.	TGF-b signature in lung metastasis of breast cancer	Signalling pathway	No
Liu et al. The prognostic role of a gene signature from tumorigenic breast-cancer cells. The New England journal of medicine. 2007. 356(3), 217-26.	Gene signatures of CD44+CD24-/low tumorigenic breast-cancer cell-lines and normal breast epithelium	Cancer stem cell	No
Charafe-Jauffret et al. Breast cancer cell lines contain functional cancer stem cells with metastatic capacity and a distinct molecular signature. 2009. Cancer research, 69(4), 1302-13.	Breast cancer stem cell signature	Cancer stem cell	No

Dontu, et al. In vitro propagation and transcriptional profiling of human mammary stem/progenitor cells. 2003. Genes & development, 17(10), 1253-70.	Gene signature of human mammary stem and progenitor cells	Cancer stem cell/ Differentiation	No

896   **Table 2**. GSEA on the Intersect 1 genes from Figure 4C against the MSigDB hallmark gene

897   sets.

INTERSECT 1 GSEA						
Gene Set Name	**# Genes in Gene Set (K)**	**Description**	**#Genes inOverlap (k)**	**k/K**	**p-value**	**FDR (q-value)**
HALLMARK _E2F_TARGETS	200	Genes encoding cell cycle related targets of E2F transcription factors.	158	0.79	5.02E-178	2.51E-176
HALLMARK _G2M_CHE	200	Genes involved in the G2/M checkpoint, as in progression	116	0.58	2.32E-105	5.80E-104

CKPOINT		through the cell division cycle.				
HALLMARK _MYC_TAR GETS_V1	200	A subgroup of genes regulated by MYC - version 1 (v1).	113	0.565	7.79E-101	1.30E-99
HALLMARK _OXIDATIV E_PHOSPHO RYLATION	200	Genes encoding proteins involved in oxidative phosphorylation.	96	0.48	1.04E-76	1.30E-75
HALLMARK _MYC_TAR GETS_V2	58	A subgroup of genes regulated by MYC - version 2 (v2).	42	0.724 1	4.60E-45	4.60E-44
HALLMARK _MTORC1_S IGNALING	200	Genes up-regulated through activation of mTORC1 complex.	66	0.33	1.70E-40	1.42E-39
HALLMARK _MITOTIC_S PINDLE	200	Genes important for mitotic spindle assembly.	60	0.3	2.76E-34	1.97E-33
HALLMARK _DNA_REPA IR	150	Genes involved in DNA repair.	51	0.34	2.54E-32	1.59E-31
HALLMARK	113	Genes up-regulated	31	0.274	3.62E-	2.01E-16

_UNFOLDED_PROTEIN_RESPONSE		during unfolded protein response, a cellular stress response related to the endoplasmic reticulum.		3	17	
HALLMARK_FATTY_ACID_METABOLISM	158	Genes encoding proteins involved in metabolism of fatty acids.	35	0.2215	5.10E-16	2.55E-15
HALLMARK_ADIPOGENESIS	200	Genes up-regulated during adipocyte differentiation (adipogenesis).	39	0.195	1.08E-15	4.89E-15
HALLMARK_CHOLESTEROL_HOMEOSTASIS	74	Genes involved in cholesterol homeostasis.	24	0.3243	1.95E-15	8.11E-15
HALLMARK_ESTROGEN_RESPONSE_LATE	200	Genes defining late response to estrogen.	35	0.175	8.70E-13	3.11E-12

HALLMARK _GLYCOLYSIS	200	Genes encoding proteins involved in glycolysis and gluconeogenesis.	35	0.175	8.70E-13	3.11E-12
HALLMARK _UV_RESPONSE_UP	158	Genes up-regulated in response to ultraviolet (UV) radiation.	28	0.1772	1.18E-10	3.95E-10
HALLMARK _SPERMATOGENESIS	135	Genes up-regulated during production of male gametes (sperm), as in spermatogenesis.	25	0.1852	4.39E-10	1.37E-09
HALLMARK _ANDROGEN_RESPONSE	101	Genes defining response to androgens.	20	0.198	7.15E-09	2.10E-08
HALLMARK _ESTROGEN_RESPONSE_EARLY	200	Genes defining early response to estrogen.	28	0.14	2.76E-08	7.68E-08
HALLMARK _IL2_STAT5	200	Genes up-regulated by STAT5 in	25	0.125	1.31E-	3.27E-06

_SIGNALIN G		response to IL2 stimulation.			06	
HALLMARK _KRAS_SIG NALING_UP	200	Genes up-regulated by KRAS activation.	25	0.125	1.31E-06	3.27E-06

900   **Table 3** .GSEA on the Intersect 2 genes from Figure 4C against the MSigDB hallmark gene

901   sets.

INTERSEC T 2 GSEA						
Gene Set Name	**# Genes in Gene Set (K)**	**Description**	**# Genes in Overlap (k)**	**k/K**	**p-value**	**FDR (q-value)**
HALLMAR K_INTERF ERON_GA MMA_RES PONSE	200	Genes up-regulated in response to IFNG [GeneID=3458].	60	0.3	3.53E-38	1.76E-36
HALLMAR K_INTERF	97	Genes up-regulated in response to alpha	43	0.44 33	4.26E-36	1.06E-34

ERON_ALPHA_RESPONSE		interferon proteins.				
HALLMARK_HYPOXIA	200	Genes up-regulated in response to low oxygen levels (hypoxia).	39	0.195	5.11E-18	8.51E-17
HALLMARK_P53_PATHWAY	200	Genes involved in p53 pathways and networks.	33	0.165	2.66E-13	3.33E-12
HALLMARK_APOPTOSIS	161	Genes mediating programmed cell death (apoptosis) by activation of caspases.	28	0.1739	4.49E-12	4.49E-11
HALLMARK_ESTROGEN_RESPONSE_EARLY	200	Genes defining early response to estrogen.	31	0.155	7.51E-12	4.70E-11
HALLMARK_HEME_METABOL	200	Genes involved in metabolism of heme (a cofactor consisting	31	0.155	7.51E-12	4.70E-11

ISM		of iron and porphyrin) and erythroblast differentiation.				
HALLMARK_MYOGENESIS	200	Genes involved in development of skeletal muscle (myogenesis).	31	0.155	7.51E-12	4.70E-11
HALLMARK_PROTEIN_SECRETION	96	Genes involved in protein secretion pathway.	21	0.2188	2.31E-11	1.28E-10
HALLMARK_EPITHELIAL_MESENCHYMAL_TRANSITION	200	Genes defining epithelial-mesenchymal transition, as in wound healing, fibrosis and metastasis.	30	0.15	3.77E-11	1.89E-10
HALLMARK_ESTROGEN_RESPONSE_LATE	200	Genes defining late response to estrogen.	29	0.145	1.82E-10	8.29E-10

HALLMARK_APICAL_JUNCTION	200	Genes encoding components of apical junction complex.	28	0.14	8.47E-10	3.53E-09
HALLMARK_IL2_STAT5_SIGNALING	200	Genes up-regulated by STAT5 in response to IL2 stimulation.	26	0.13	1.62E-08	5.77E-08
HALLMARK_KRAS_SIGNALING_DN	200	Genes down-regulated by KRAS activation.	26	0.13	1.62E-08	5.77E-08
HALLMARK_ALLOGRAFT_REJECTION	200	Genes up-regulated during transplant rejection.	25	0.125	6.63E-08	2.21E-07
HALLMARK_UNFOLDED_PROTEIN_RESPONSE	113	Genes up-regulated during unfolded protein response, a cellular stress response related to the endoplasmic reticulum.	18	0.1593	1.16E-07	3.62E-07

HALLMARK_ADIPOGENESIS	200	Genes up-regulated during adipocyte differentiation (adipogenesis).	24	0.12	2.60E-07	6.83E-07
HALLMARK_TNFA_SIGNALING_VIA_NFKB	200	Genes regulated by NF-kB in response to TNF [GeneID=7124].	24	0.12	2.60E-07	6.83E-07
HALLMARK_XENOBIOTIC_METABOLISM	200	Genes encoding proteins involved in processing of drugs and other xenobiotics.	24	0.12	2.60E-07	6.83E-07
HALLMARK_IL6_JAK_STAT3_SIGNALING	87	Genes up-regulated by IL6 [GeneID=3569] via STAT3 [GeneID=6774], e.g., during acute phase response.	15	0.1724	4.51E-07	1.13E-06

904   **Supplementary information**

905   Supplementary information is available as Supplementary Files.

Figure 1

Figure 2

A. -log(pValue)

1. Cell cycle_Mitosis
2. Cell cycle_Core
3. Cell cycle_G2-M
4. Cytoskeleton_Spindle microtubules
5. Cell cycle_S phase
6. Inflammation_ Interferon signaling
7. DNA damage_Checkpoint
8. DNA damage_MMR repair
9. Development_Regulation of angiogenesis
10. Development_EMT_Regulation of epithelial-to-mesenchymal transition

Networks

B. -log(pValue)

1. Cell cycle_The metaphase checkpoint
2. Cell cycle_Chromosome condensation in prometaphase
3. DNA damage_ATM/ATR regulation of G2/M checkpoint
4. DNA damage_ATM/ATR regulation of G1/S checkpoint
5. Cell cycle_Spindle assembly, chromosome separation
6. Cell cycle_Cell cycle (generic schema)
7. Cytoskeleton remodeling_TGF,WNT and cytoskeletal remodeling
8. Development_PDGF signaling via STATs and NF-kB
9. Cell cycle_Role of APC in cell cycle regulation
10. Immune response_Oncostatin M signaling via JAK-Stat in human cells

Maps

C. Metastasis associated genes

D. Robo1, Fermt1, Mir30a — Relative mRNA expression; -Dox, +Dox; ns, **, ***

Figure 3

Figure 4

Figure 5

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 3

Supplementary Figure 4

Supplementary Figure 5