1 **The Germline Variants rs61757955 and rs34988193 are Predictive of Survival in**
2 **Low Grade Glioma Patients**
3

4 **Ajay Chatrath[1], Manjari Kiran[1], Pankaj Kumar[1], Aakrosh Ratan[2], and Anindya**
5 **Dutta[1]**
6
7 [1] Department of Biochemistry and Molecular Genetics, University of Virginia,
8 Charlottesville, Virginia
9
10 [2] Center for Public Health Genomics, University of Virginia, Charlottesville, Virginia
11

12 Running Title:
13 Germline Variants are Predictive of Survival in LGG Patients
14
15 Keywords: Low Grade Glioma, Germline Variant, Single Nucleotide Polymorphism,
16 Survival Analysis, *GRB2*, *ANKDD1a*
17

Financial Support:
This work was supported by grants from the NIH R01 CA166054 and CA60499 and a
Cancer Genomics Cloud Collaborative Support grant. The Seven Bridges Cancer
Genomics Cloud has been funded in whole or in part with Federal funds from the
National Cancer Institute, National Institutes of Health, Contract No.
HHSN261201400008C and ID/IQ Agreement No. 17X146 under Contract No.
HHSN261201500003I.

Corresponding Author:
Anindya Dutta
University of Virginia School of Medicine
1240 Jordan Hall, Charlottesville, VA 22908.
Phone: 434-924-1227
Fax: 434-924-5069
E-mail: ad8q@virginia.edu

33
34 Potential Conflicts of Interest:
35 The authors do not have any conflicts of interest to disclose.
36
37 Abstract Word Count: 250
38 Manuscript Word Count: 5,660
39 Number of Figures/Tables: 7
40 Number of References: 50
41 Number of Supplementary Figures/Tables: 8

Abstract

Low grade gliomas are invasive brain tumors that are difficult to completely

resect neurosurgically. They often recur following resection and progress, resulting in

death. Although previous studies have shown that specific germline variants increase the

risk of tumor formation, no previous study has screened many germline variants to

identify variants predictive of survival in glioma patients. In this study, we present an

approach to identify the small fraction of prognostic germline variants from the pool of

over four million variants that we variant called in The Cancer Genome Atlas whole

exome sequencing and RNA sequencing datasets. We identified two germline variants

that are predictive of poor patient outcomes by Cox regression, controlling for eleven

covariates. rs61757955 is a germline variant found in the 3' UTR of *GRB2* associated

with increased *KRAS* signaling, *CIC* mutations, and 1p/19q co-deletion. rs34988193 is a

germline variant found in the tumor suppressor gene *ANKDD1a* that causes an amino

acid change from lysine to glutamate. This variant was found to be predictive of poor

prognosis in two independent low grade glioma datasets and is predicted to be within the

top 0.06% of deleterious mutations across the human genome. The wild type residue is

conserved in all 22 other species with a homologous protein.

Implications: This is the first study presenting an approach to screening many germline

variants to identify variants predictive of survival and our application of this

methodology revealed the germline variants rs61757955 and rs34988193 as being

predictive of survival in low grade glioma patients.

Introduction

Grade II and grade III (low grade) gliomas are primary brain tumors that are

derived from glial cells and include astrocytomas and oligodendrogliomas. They are most

commonly found in the cerebral hemispheres. They are highly invasive and therefore

difficult to completely resect neurosurgically without significant patient morbidity.

Following surgery, patients are typically treated with chemotherapy and radiation, though

these tumors typically recur or progress to grade IV gliomas and are fatal.[1] The median

survival following low grade glioma diagnosis is around 7 years.[2]

While the 2007 World Health Organization's (WHO) classification of central

nervous system neoplasms differentiated between neoplasms primarily based on

histological features, the updated 2016 WHO classification system now utilizes both

molecular and histological parameters.[1] Isocitrate dehydrogenase mutation (*IDH*) status,

1p/19q co-deletion status, telomerase reverse transcriptase (*TERT*) promoter mutation

status, *MGMT* promoter methylation, *TP53* mutation status, and *ATRX* mutation status

may be used to molecularly characterize gliomas.[1,3] The availability of genomic data

from patient glioma samples from groups such as The Cancer Genome Atlas (TCGA), the

Chinese Glioma Genome Atlas (CGGA), and the Ivy Glioblastoma Atlas Project has

substantially contributed to our understanding of these tumors.[4,5]

Many studies have utilized these datasets to identify gene expression signatures,

microRNA expression patterns, somatic mutation status, and imaging characteristics that

are predictive of survival in low grade gliomas.[6–8] While studies have shown that

germline mutations can increase an individual's susceptibility for specific cancers,[9–12]

including a recent study that identified 853 pathogenic or likely pathogenic germline

variants found in 8% of 10,389 cancer patients,[13] no study has comprehensively screened

all of the germline variants in a given cancer type to discover the prognostic variants in

that cancer type. Although germline mutations have been shown to be prognostic in

breast cancer[14] and medulloblastoma[9] in genes that have been well-characterized in the

context of these cancers, these variants were not identified using an unbiased approach

that screened a large number of germline variants. Identifying prognostic germline

variants is challenging due to the limited effect size of germline variants, the large

number of germline variants, and confounding clinical factors that may be associated

with germline variants. Here we present a novel methodology for identifying prognostic

germline variants and report two germline variants that we have found to be associated

with survival in low grade glioma patients.

Methods

Glioma Datasets

491whole exome sequenced normal blood samples (WXS normal), 503 whole

exome sequenced tumor samples (WXS tumor), and 501 RNA sequenced tumor samples

(RNA tumor) from TCGA low grade glioma[4] patients available on the Cancer Genomics

Cloud (CGC)[15] platform were used as part of this analysis. The clinical information was

downloaded directly from the TCGA data portal using the GenomicDataCommons

(https://bioconductor.org/packages/release/bioc/html/GenomicDataCommons.html) R

package available through Bioconductor. Additional molecular characteristics about these

TCGA patients were acquired by downloading the supplement from Ceccarelli et. al.[16]

The raw sequencing data from the Chinese Glioma Genome Atlas patients was

downloaded using accession number SRP027383 from the Sequence Read Archive.

Clinical information for these patients was downloaded directly from the project's

website (http://www.cgga.org.cn/).

Variant Calling

Variant calling was performed on the TCGA low grade glioma whole exome

sequenced normal blood samples (WXS normal), whole-exome sequenced tumor samples

(WXS tumor), and RNA sequenced tumor samples (RNA tumor) using VarDict[17] on

CGC. The VarDict settings were set at default except for requiring mapping quality

greater than 30, base quality greater than 25, a minimum of 3 variant reads, minimum

allele frequency of 5%, and the removal of duplicate reads. We compiled a list of all of

the unique variants and ran 'samtools[18] depth' on all sequencing files requiring a

mapping quality greater than 30. We determined the status of each variant in each patient

from the three datasets (WXS normal sample, WXS tumor sample, and RNA tumor

sample). The variant status at positions with fewer than ten reads for a given patient was

changed to unknown. We used the WXS tumor samples to insert variant calls into the

WXS normal samples at positions at which a variant status was listed as unknown in the

WXS normal samples. If the variant status was still missing in a given patient, we then

used the RNA tumor sample to insert variant calls into the combined WXS variant call

set, allowing us to create the combined set of variant calls.

The same program parameters and approach were used to variant call and process

the CGGA RNA sequencing dataset. All computation on the CGGA dataset was

performed locally and not on CGC.

Quality Control

138   We used annovar[19] to determine the allele frequencies of the variants called by

139   VarDict as listed in gnomAD (http://gnomad.broadinstitute.org/). We calculated the allele

140   frequency of the variants in our study using the following formula:

$$\frac{2 * Number\ of\ Minor\ Allele\ Homozygotes + Number\ of\ Heterozygotes}{2 * Total\ Number\ of\ Patients}$$

The R package GGally (https://cran.r-project.org/web/packages/GGally/index.html) was

used to calculate the correlation between the four variant call sets and to display their

correlations with each other. Only variants with an allele frequency of greater than 5% in

gnomAD and found in 15 or more of the TCGA low grade glioma patients were tested for

an association with survival by Cox regression.

Because we used the WXS tumor and RNA tumor samples to fill in missing

variant calls, we evaluated whether somatic mutations were affecting the validity of our

results. We first determined the percentage of variants called in the WXS tumor sample

that were somatic mutations. To do this, we downloaded the set of somatic mutations

generated by the TCGA Research Network.[20] We then calculated the number of somatic

mutations called in each patient in this variant call set and divided that number by the

total number of variants called in that patient's WXS normal sample. To assess whether

somatic mutations were affecting the integrity of our results, we counted the number of

times that a somatic mutation called by the TCGA Research Network overlapped with the

set of germline variants that we were testing for an association with survival.

Since we used the RNA tumor sample to fill in missing variant calls, we evaluated

whether RNA editing was having a significant impact on our analysis. To do this, we

downloaded the set of over 2.5 million known RNA editing sites from a rigorously

annotated database of RNA editing sites, RADAR.[21] We counted the number of times

that the germline variants that we were testing for an association with survival overlapped

with any of the known 2.5 million RNA editing sites.

Principal Component Analysis

In order to calculate principal components that could separate patients on the basis

of race, we used PLINK[22] to create a pruned set of germline variants to avoid bias from

variants in linkage disequilibrium. Pruning was performed using a window size of 50

variants and a variance inflation factor of 2. These variants were used to calculate

principal components using base R.

Cox Regression and Receiver Operator Characteristic Curves

Lasso in the R package glmnet[23] was run on 17 covariates (**Table 1**). Information

about patient age, gender, tumor location, grade, treatment site, and TP53 mutation status

was acquired from the TCGA data portal, while data for patient somatic mutation count,

percent aneuploidy, TERT expression, IDH mutation status, 1p/19q co-deletion status,

MGMT promoter methylation status, and chromosome 7 gain with chromosome 10 loss

status was acquired from Ceccarelli et. al.[16] The principal components were calculated as

described above. 11 of these 17 covariates were selected for inclusion in the final model

for survival prediction. The R packages survival[24] and survminer[25] were used to run Cox

regression and create Kaplan-Meier curves. For each minor allele, we our model tested

whether the minor allele was associated with a difference in survival outcomes with

respect to the reference allele. False discovery rate correction was performed through

Bonferroni correction.

Receiver operator characteristic (ROC) curves were created and evaluated using

the survivalROC (https://cran.r-

project.org/web/packages/survivalROC/survivalROC.pdf) and pROC (https://cran.r-

project.org/web/packages/pROC/pROC.pdf) R packages. In order to test whether

rs61757955 significantly improves the survival model consisting of the eleven covariates

selected by Lasso, we compared the two ROC curves using the bootstrap method with

1000 iterations. We also used this bootstrapping approach to determine whether

ANKDD1a expression levels, GRB2 expression levels, rs61757955, and rs34988193

together improve the survival model with respect to the eleven covariates selected by

Lasso.

RNA-Sequencing Data Processing

We downloaded the HTSeq FPKM quantification files for each patient from the

Genomic Data Commons data portal. We only used gene quantification files from

primary tumor samples as part of this analysis. Replicate samples from a single patient

were averaged.

Variant Correlation to Covariates and Somatic Mutations

In order to test for associations between the germline variants and genomic and

histological tumor characteristics, we divided patients based on their germline variant

status. We used the Wilcoxon rank-sum test to test for significant differences in each of

the continuous variables between patients with and without a given variant. We used

Fisher's exact test to test for differences in each of the discrete variables using a similar

approach. Somatic mutation calls were downloaded from Ellrott et. al.[20]

Gene Set Enrichment Analysis

Gene set enrichment analysis (GSEA) of mRNA changes associated with

rs61757955 and rs34988193 was performed by dividing the patients into two groups for

each variant based on whether or not they had the reference allele at the position of the

variant. For each germline variant, we calculated the log fold change for all genes

expressed greater than one fragment per kilobase per million mapped reads (FPKM)

between patients with the variant and without the variant. For each gene, fold change was

calculated by dividing the median expression of the gene in patients with the variant by

the median expression of the gene in patients without the variant. We used the log fold

change to rank the genes from greatest log fold change to smallest log fold change. This

file was used as input for GSEA.[26]

Variant Annotation

In order to identify deleterious mutations, we annotated all variants by combined

annotation dependent depletion (CADD) scores and only analyzed the variants predicted

to be within the top 0.1% of all deleterious variants (CADD > 30).[27] This led us to

identify rs34988193 in *ANKDD1a* as a potentially deleterious variant predictive of

survival. Because rs34988193 causes an amino acid change from positively charged

lysine to negatively charged glutamate, we ran a BLASTp

(httpps://blast.ncbi.nlm.nih.gov/Blast.cgi) search so that we could determine how many

species have a protein homologous to ANKDD1a and how consistently the wild type

lysine residue was conserved. We identified homologous sequences in 22 other species.

These sequences were aligned using ClustalW in MEGA.[28] We also annotated this

variant with its PhyloP score.[29] Because the crystal structure for *ANKDD1a* was not

available, we downloaded the predicted model for this protein from Modbase

(https://modbase.compbio.ucsf.edu/modbase-cgi/index.cgi) and calculated the Gribskov

score using prophecy on EMBOSS.[30] We retrieved linked variants from Ensembl using

the population of Utah residents with Northern and Western European ancestry which is

demographically similar to the TCGA low grade glioma patient population.

Results

Identification of High Quality Germline Variants

235   Our variant calling pipeline is shown in **Figure 1**. Briefly, we used the variant

caller VarDict on Cancer Genomics Cloud to identify variants from whole exome

sequencing (WXS) and RNA sequencing samples in about 500 low grade glioma

patients. In total, we found 4,453,701 unique variants. We used 'samtools depth' to

determine the sequencing depth at each of these variants for each patient and changed the

variant status to 'unknown' for patients with sequencing coverage less than 10 reads at a

given position. We created a set of combined variant calls by using the WXS and RNA

tumor samples to fill in unknown values in the whole exome sequenced normal samples

that resulted from having a sequencing coverage of less than 10 reads at a given position.

This approach increased our sample size and enabled us to include many more variants in

our analysis than if we had solely used variant calls from the whole exome sequenced

normal blood samples. Ultimately, this left us with four sets of variants – WXS normal,

WXS tumor, RNA tumor, and a combined set that resulted from merging the other three

variant call sets, giving preference to the WXS normal and then WXS tumor variant calls.

We used the combined variant call set when testing variants for an association with

survival. We only tested variants found in 15 or more low grade glioma TCGA patients

and listed in gnomAD as having an allele frequency of greater than 5%.

Tumor Variant Calls are not Significantly Affected by Somatic Mutations or RNA

Editing After Filtering

Because we used sequencing data from the WXS tumor and RNA tumor samples

to fill in missing calls in the WXS normal samples, we evaluated our variant calls for

contributions from somatic mutations and RNA editing. We first showed that the

majority of variant calls in the tumor sample are germline variant calls. To do this, we

counted the number of somatic mutations called by the TCGA Research Network's

analysis in each patient and divided that number by the number of variants that we called

in the WXS normal sample.[20] The median number of somatic mutations called per patient

was 39. The median number of variants called in the WXS normal sample was 95,794.

We therefore estimated that over 99.9% of variants called in the WXS tumor sample

consisted of germline variants and that the percentage of somatic mutations in the WXS

tumor sample across all patients was quite small (**Figure S1**). Because somatic mutations

rarely occur at the same position, we suspected that the number of somatic mutations

included in our study was extremely small since we limited our analysis to variants found

in 15 or more of the low grade glioma patients and found in gnomAD with an allele

frequency of greater than 5%. Indeed, only one of the 196,022 variants that we tested

overlapped with a somatic mutation. This somatic mutation occurred in only a single

patient (**Table S1**). Ultimately, we did not find any evidence to suggest that somatic

mutations were impacting the quality of our analysis.

We next determined whether RNA editing was affecting our analysis by

downloading the 2.5 million known RNA editing sites from the rigorously annotated

RNA editing database, RADAR.[21] Only 215 of the 196,022 variants that we tested were

located at a position that overlapped with a known RNA editing site. We did not find any

of these variants to be prognostic as part of our analysis. We therefore did not find any

empirical evidence to suggest that somatic mutations or RNA editing impacted our

findings (**Table S1**).

Finally, we established that our four variant call sets (WXS normal, WXS tumor,

RNA tumor, and combined) were concordant with each other by calculating the allele

frequency of each variant called in the four sets and demonstrating a very strong

correlation between all pairs of variants ($r > 0.98$ for all pairs, **Figure S2**). To further

evaluate the quality of our variants calls, we calculated the frequency of each allele and

compared it to the frequency of these alleles as listed in gnomAD. Our alleles frequencies

were well correlated with gnomAD ($r > 0.93$ for all four variant sets, **Table S2**). As

expected, the distribution of allele frequencies is negatively skewed as the majority of the

identified variants are rare (**Figure S2**). We used the variants from the WXS normal

samples to determine the principal components. As expected, these principal components

effectively separate patients on the basis of reported race (**Figure S3**).

Identification of 271 Prognostic Germline Variants that are Independent of Clinical

Covariates

In order to identify clinically relevant germline variants, we restricted our analysis

to variants found in at least 15 patients in the TCGA dataset and found in gnomAD with

an allele frequency of greater than five percent. This restricted our analysis to 196,022

testable variants (**Figure 2A**). In order to reduce the risk of identifying variants that are

prognostic because they are confounded by other covariates known to be associated with

survival, we used the machine learning algorithm Lasso to determine which of 17

covariates should be controlled for in our Cox regression model. Lasso regression was

useful in the screening of these 17 covariates because it penalizes models based on the

number of coefficients, allowing for the elimination of less predictive coefficients from

the model. The algorithm selected 10 covariates known to be associated with differences

in survival in low grade glioma (age, somatic mutation count, percent aneuploidy,

histological subtype of astrocytoma, tumor grade, treatment site, *IDH* mutation status,

1p/19q co-deletion status, *MGMT* promoter methylation status, chromosome 7

gain/chromosome 10 loss status) along with the third principal component that we

calculated (**Table 1**). Although the first two principal components are more effective in

stratifying patients on the basis of race than the third principal component, the selection

of the third principal component over the first two suggests that the third principal

component contributes more information to the survival model than the first two principal

components. This third principal component primarily separates African Americans from

each other, suggesting that a subpopulation of African Americans experienced worse

clinical outcomes in this dataset compared to other groups. We ran Cox regression on all

196,022 variants one at a time, controlling for these 11 covariates, to identify germline

variants predictive of survival.

We identified 271 germline variants that are predictive of survival (p < 0.001)

(**Figure 2A**). As is the case with germline variants in general, the majority of these

germline variants are found in protein-coding genes (**Figure 2B**), are located in introns

(**Figure 2C**), and are single nucleotide polymorphisms (**Figure 2D**). Most single

nucleotide polymorphisms are transitions (**Figure 2E**).

The Germline Variant rs61757955 in *GRB2* is Associated with Poor Prognosis

We identified two germline variants that are highly predictive of survival after

false discovery rate correction (FDR < 0.10) (**Figure 3A, Table 2A**). rs61757955 results

in a mutation in the 3' UTR of Growth Factor Receptor Bound Protein 2 (*GRB2*) and is

associated with a poor prognosis (p=7.08E-10, hazard ratio(HR)=20.4, **Figure 3B, Table**

2A). To determine whether rs61757955 enhances the survival model compared to the

eleven clinical covariates alone, we calculated a risk score for each patient using a Cox

regression model with rs61757955 and the other 11 covariates and a risk score using the

11 covariates alone. Using these risk scores, we determined the rate at which a patient

would be correctly labeled as alive or dead at 7 years with a given false positive rate to

create a receiver operator characteristic curve. The increased area under the curve

suggests that rs61757955 enhances the survival model compared to the eleven clinical

covariates alone (p=0.0489, **Figure 3C**). The allele frequency of rs61757955 is close to

0% according to the 1000 Genomes Project[31] in the Chinese population and, as expected,

did not show up in the Chinese Glioma Genome Atlas. We also found rs28672782, a

germline variant found in the intron of BRSK2, to be associated with a favorable

prognosis, though the testable sample size for this variant was small and the maximum

follow up for patients with this variant was only three years. Therefore, we did not

investigate this variant further (**Figure S4, Table 2A**).

In order to test whether rs61757955 in *GRB2* is associated with an increased risk

of other genomic abnormalities, we separated patients on the basis of this variant to see if

there was a difference in the incidence of the genomic or histological variables (**Table 3**).

We found this variant to be associated with an increased incidence of 1p/19q co-deletions

(p=0.038). Because 1p/19q co-deletions are frequently seen in Capicua transcriptional

repressor (*CIC*) mutated gliomas[32] and *CIC* aberrations are known to be a driver in low

grade glioma tumorigenesis,[33] we tested whether there was a difference in the incidence

of *CIC* mutations in patients with this variant. 38% of patients with this variant had *CIC*

mutated gliomas, whereas only 16% of patients without the variant had a *CIC* mutation

(p=0.0168, **Table 3**). Although the incidence of oligodendrogliomas was elevated in

patients with the variant compared to patients without the variant, consistent with reports

from the literature that 1p/19q co-deletions and *CIC* mutations are enriched in

oligodendrogliomas,[32] this difference was not statistically significant (p=0.475). Since

rs61757955 is in a non-coding region, we also tested whether this variant is associated

with differences in gene expression. We separated patients based on their variant status

and calculated the log fold change of each gene between patients with the variant and

patients without the variant. This data was used as the input for gene set enrichment

analysis (GSEA). We found rs61757955 to be associated with increased *KRAS* signaling

(FDR=0.015) (**Figure 3D**).

Because we only have whole exome sequencing and RNA sequencing data from

The Cancer Genome Atlas, we do not know whether the upregulation of genes in the

KRAS signaling pathway and the increased incidence of *CIC* mutations and 1p/19q

deletions are due to this variant or a linked variant in a regulatory region that we would

be able to analyze with whole genome sequencing data. Therefore, we identified the four

other variants that are genetically linked to rs61757955 in the European population, the

population which is most similar to the TCGA low grade glioma patient population

(**Table S3**). These variants did not pass the criteria to be included within the 196,022

testable variants that we had identified at the beginning of this study but could become

useful in the future.

368   rs34988193 is a Deleterious Germline Variant Present in *ANKDD1a* Associated with

369   Poor Outcomes

In order to identify prognostic variants that are predicted to be deleterious due to

effects on the encoded protein, we repeated our analysis but restricted it to only variants

with a combined annotation dependent depletion (CADD) score greater than 30 and

expression greater than one FPKM on average. 81 variants met this criteria. These

variants correspond to the top 0.1% of deleterious mutations as predicted by this scoring

system. We found the germline variant rs34988193 in the tumor suppressor gene

ANKDD1a to be associated with poor prognosis in the TCGA dataset (p=0.001,

HR=1.73, FDR < 0.10, **Figure 4A-B, Table 2B**). Because this variant is found in both

the European and Asian populations, we were able to test whether this variant is also

predictive of survival in the independent Chinese Glioma Genome Atlas (CGGA) dataset.

We found this variant to be predictive of survival in the CGGA dataset and we found the

hazard ratio that we calculated in CGGA to be very similar to the hazard ratio calculated

in the TCGA dataset (p=0.0743, HR=1.79, **Figure 4C, Table 2B**). rs34988193 is not

linked with any other variant in the European population. We did not find any enriched

pathways after performing gene set enrichment analysis and this variant was not

associated with differences of any of the genomic or histological variables (**Table S4**).

ANKDD1a contains ten ankyrin repeat domains and one death-like domain. This

variant causes a non-synonymous mutation in the last codon of the ninth ankyrin repeat

domain. The AAG to GAG codon change results in the incorporation of negatively

charged glutamate instead of the wild type positively charged lysine residue in the loop

between ankyrin repeats nine and ten (**Figure 4D**). This variant has a CADD score of 32

and is therefore predicted to be in the top 0.06% of deleterious mutations across the

human genome. We performed a BLASTp search using the *ANKDD1a* protein sequence

to identify homologous sequences in 22 other species. We aligned these sequences using

ClustalW and found that this lysine residue is conserved in all 22 of these species (**Figure

4E**). The PhyloP score at this position is 8.42, suggesting that evolution is occurring

much more slowly than expected at this residue assuming no selection pressure. We

determined the position-specific profile Gribskov's score for a lysine to glutamate amino

acid change at this position using the multiple sequencing alignment from 23 species to

be 15 to 3, suggesting that this variant is highly unfavorable.

Combined Model Predicts Survival Better Than Clinical Covariates Alone

As a result of this analysis, we found the germline variants rs61757955 in the 3'

UTR of GRB2 and rs34988193 in the protein-coding region of ANKDD1a to be

predictive of survival in low grade glioma patients. We constructed a survival model

consisting of the eleven clinical covariates, rs61757955, rs34988193, GRB2 expression,

and ANKDD1a expression and generated a receiver operator characteristic curve by

using this model to categorize patients as alive or dead after seven years of follow up.

This combined model is significantly better at predicting survival compared to the eleven

clinical covariates alone (p=0.0279, **Figure 4F**).

Discussion

Up until this point, the identification of prognostic features in gliomas has been

limited to clinical factors, somatic mutations, gene expression changes, and methylation

pattern changes.[6–8] Although many studies have commented on how germline variants

could enable physicians to better individualize patient care by being able to better predict

how a patient might respond to chemotherapeutic treatment,[34–36] most large-scale studies

have focused on identifying germline variants that predispose or protect an individual to a

disease.[13,37] These studies have not focused on understanding how germline variants can

be used to individualize patient care following diagnosis. Identifying prognostic germline

variants is difficult due to the large number of germline variants, the limited effect of any

single germline variant, and clinical factors that may confound the effect of germline

variants. In this study, we have developed a novel method that can be used to identify

prognostic germline variants and we have used that method to identify two variants that

are predictive of survival in the TCGA dataset. The germline variant rs61757955 in

GRB2 is not found in the Asian population and so could not be confirmed in an

independent dataset. In contrast, the germline variant rs34988193 in *ANKDD1a* is found

in both the European and Asian populations, and remarkably, was found to be prognostic

with very similar hazard ratios in both the TCGA and CGGA datasets.

Studies of germline variants using TCGA datasets typically solely utilize the

WXS normal blood samples.[13,38] One major disadvantage to this approach is that it limits

the analysis to genes within the capture regions of the whole exome sequencing kits used

by the study.[4] In this study, we combined the information from both the whole exome

sequencing and RNA sequencing datasets for a given patient to identify germline variants

outside of the whole exome sequencing capture region. Our approach had the added

benefit of providing us with more information for a given variant for variants with low

sequencing depth in the whole exome sequencing datasets. We do not believe that this

approach significantly affected the accuracy of our variant calls because the allele

frequencies calculated from the RNA sequencing dataset were well correlated with the

allele frequencies from gnomAD and with the allele frequencies calculated from the

whole exome sequencing datasets. We showed that somatic mutations and RNA editing

did not affect the integrity of our finding. Only one somatic mutation in a single patient

overlapped with the 196,022 variants that we tested in our analysis and only 215 of the

196,022 variants that we tested overlapped with the 2.5 million known RNA editing sites.

We did not find any of these variants to be predictive of survival. Instead, we feel that the

increased sample size resulting from the additional sequencing coverage greatly

outweighs any effect that somatic mutations or RNA editing had on our results.

We next needed to devise an approach to using these germline variants in a Cox

regression model. We first had to decide how to deal with the absence of a variant in the

variant call file. The variant could be absent because the patient was wild type for that

allele or because the sequencing depth at that position was too low to make the variant

call. We therefore determined the sequencing depth of each variant at each position so

that we could exclude patients with low sequencing depths for the testing of specific

variants. Testing a large number of variants increased the probability of a variant being

significant solely because it was confounded with another significant variable. To avoid

this issue, we tested each variant while controlling for 11 other covariates that we found

to be predictive of survival. In this study, we found rs61757955 to be associated with

differences in 1p/19q co-deletion status. By including the 1p/19q co-deletion as a

covariate in our model, we were able to estimate the effect of rs61757955 independent

from the 1p/19q co-deletion status and the other ten covariates.

GRB2 is a signal transduction adaptor protein that plays an oncogenic role in a

variety of cancers.[39–42] *GRB2* plays an important role in the *RAS/RAF/ERK* pathway. Its

SH2 domain binds the phosphotyrosine of activated growth factor receptor, while its two

SH3 domains bind the guanine nucleotide exchange factor son of sevenless (*SOS*)

protein, resulting in *SOS* recruitment to the plasma membrane and subsequent *RAS*

activation. *RAS* binds and activates the kinase *RAF*, which phosphorylates the kinase

MEK. *MEK* phosphorylates and activates extracellular signal-regulated kinase (*ERK*)

which transmits the signal to transcription factors in the nucleus. This results in cell

proliferation.[43] We found the variant rs61757955 located in the 3' UTR of *GRB2* to be

associated with poor prognosis in glioma patients. Separating patients on the basis of this

variant revealed that the *KRAS* signaling pathway is upregulated in patients with this

variant. As described above, *GRB2* plays a well-characterized role in this pathway.[43] We

also found this variant to be associated with an increased incidence of *CIC* mutations and

1p/19q co-deletions. *CIC* is a known driver of low grade glioma pathogenesis.[33]

Mutations in *CIC* are common in oligodendrogliomas and are associated with poor

prognosis.[4,32] Although patients with rs61757955 variant exhibited an elevation in the

incidence of oligodendrogliomas which we expected given the increased incidence of

CIC mutations and 1p/19q co-deletions,[32] this difference was not statistically significant.

It is possible that this germline variant or the four other germline variants that it is linked

with increase a patient's risk for oligodendrogliomas with the *CIC* mutation and 1p/19q

co-deletion.

In this study, we were only able to study variants in the whole exome or RNA

sequencing data. Although it is possible that the 3' UTR of *GRB2* has regulatory activity

or affects *GRB2* protein translation efficiency, it is also possible that one of the variants

that rs61757955 is linked to regulates the *KRAS* signaling pathway. None of the four

linked variants are in the protein coding sequence of *GRB2* so that if they upregulate

RAS activity, like the rs61757955, they likely do so by regulating the expression of

GRB2. While recent large-scale sequencing studies have published patient whole genome

sequences,[44] this data is not yet available for gliomas. We will be able to apply our

approach to variants in regulatory regions in the future to specifically identify these

prognostic variants when whole genome sequencing data for gliomas is available. Our

inability to further study this variant in the CGGA dataset due to this variant being rare in

Asian populations is a limitation of this study which could be addressed in the future with

the availability of additional glioma sequencing datasets. This result also suggests that the

clinical usefulness of specific germline variants is dependent on the frequency of that

germline variant in the population.

ANKDD1a is a tumor suppressor gene that has been shown to inhibit cell

autophagy and induce apoptosis in glioblastoma multiforme (GBM). It directly interacts

with and upregulates *FIH1*, resulting in inhibition of *HIF1α* activity and decreased

HIF1α half-life. This induces apoptosis in GBM cell lines in hypoxic microenvironments.

Hypermethylation of this gene is common in GBM and leads to decreased *ANKDD1a*

expression and increased cell proliferation.[45] We found the germline variant rs34988193,

located at the end of the ninth of ten ankyrin repeat domains in this protein, to be

associated with a poor prognosis in low grade glioma patients in both the TCGA and

CGGA datasets. The hazard ratio independently calculated using the two datasets is

remarkably similar. The wild type lysine residue is conserved in all 22 species with a

homologue to *ANKDD1a* and this position has a high PhyloP score. This variant is

predicted to be within the top 0.06% of deleterious mutations in the human genome by

CADD score[27] because it causes a change from a positively charged lysine residue to a

negatively charged glutamic acid residue in the loop of this ankyrin repeat. Ankyrin

repeats are common domains known for their involvement with protein-protein

interactions.[46,47] Previous studies have suggested that mutations in the loops of ankyrin

repeats may disrupt protein-protein interactions.[48–50] The change from a positively to

negatively charged amino acid resulting from the germline variant rs34988193 in the loop

of *ANKDD1a* may disrupt *ANKDD1a*'s protein interaction partners and could explain the

poor prognosis associated with this variant seen in two independent datasets. Given the

amino acid change, further studies involving rs34988193 in *ANKDD1a* could be directed

towards experimentally determining whether or not this variant alters *ANKDD1a*'s

protein-protein interactions.

rs61757955 in *GRB2* and rs34988193 in *ANKDD1a* could also be used to enhance

predictions made by survival models clinically, as we found that these variants are

significant predictors of prognosis even after controlling for eleven covariates. The

prognostic effect of rs34988193 in *ANKDD1a* seems to be fairly reliable, as we found

that this variant had a similar hazard ratio in both the TCGA and CGGA datasets. Our

approach could be used in the future to identify sets of germline variants that together

enhance the predictions made by survival models, though the current number of low

grade glioma sequencing samples is small relative to the large number of possible

combinations of germline variants. Focused studies on particular sets of genes or

pathways could potentially get around this low sample size problem by drastically

limiting the number of variants studied. We believe that this study provides researchers

with an effective approach to identifying biologically significant germline variants and

provides clinicians with germline variants that could enhance currently existing survival

models.

Acknowledgements

We thank Dr. Wei Min Chen at the University of Virginia for his statistical input

on our Cox regression models. We thank Dr. Ana Damljanovic and Dr. Liz Williams for

their assistance with computation on the Cancer Genomics Cloud Platform. We thank

dbGAP for providing us with access to The Cancer Genome Atlas data. We thank all of

the patients and their families that participated in The Cancer Genome Atlas and Chinese

Glioma Genome Atlas studies. We thank the Dutta lab members for the valuable

feedback during the drafting of this manuscript.

Figure Legends

Figure 1. A flowchart describing the steps involved in identifying prognostic germline

variants.

Figure 2. Prognostic germline variants in the TCGA dataset.

(A) Of the 4.4 million unique variants called in the TCGA dataset, we ran Cox regression

on the 196,022 germline variants found in gnomAD with an allele frequency greater than

5% and found in 15 or more of the TCGA low grade glioma patients.

(B-E) Similar to the 196,022 germline variants, the 271 prognostic variants are mostly

found in **(B)** protein-coding genes, **(C)** are located in introns, and are **(D)** single

nucleotide polymorphisms (SNP). **(E)** Most single nucleotide polymorphisms cause

transitions.

Figure 3. rs61757955 is a highly prognostic germline variant identified in the TCGA

dataset.

(A) Manhattan plot showing the p-values resulting from testing each germline variant by

Cox regression, controlling for the 11 variables in red in Table 1.Two variants passed the

FDR threshold in the TCGA dataset.

(B) A Kaplan-Meier plot depicting the deleterious outcome associated with rs61757955,

adjusting for the eleven covariates.

(C) Receiver operator characteristic curve at 7 years. rs61757955 increases the area under

the curve compared to the 11 covariates alone, suggesting that it improves the clinical

model.

(D) Separation of patients on the basis of whether or not they have this germline variant

to determine which genes are induced (red) or repressed (blue) in patients with

rs61757955. Subsequent gene set enrichment analysis reveals that patients with this

germline variant exhibit upregulation of the genes involved with *KRAS* signaling.

Figure 4. rs34988193 is a prognostic variant predicted to be highly deleterious.

(A) A Manhattan plot with the p-values resulting from testing each germline variant by

Cox regression, controlling for the eleven covariates in red in Table 1. rs34988193 is

prognostic (FDR<0.10) in the TCGA when restricting the analysis to the top 0.1% most

deleterious variants by combined annotation dependent depletion (CADD).

(B-C) Kaplan-Meier plots depicting the deleterious outcome associated with rs34988193

in the **(C)** TCGA and **(D)** CGGA datasets, adjusted for the eleven covariates.

(D) A schematic showing that this variant is located in the ninth ankyrin repeat of

ANKDD1a. The predicted protein structure of *ANKDD1a* reveals that this variant leads to

an amino acid change from lysine to glutamate on the loop of an ankyrin repeat.

(E) Multiple sequence alignment of *ANKDD1a* in 22 species showing that lysine is

conserved at this position in all of the species with this protein.

(F) Receiver operator characteristic curves comparing the ability of two survival models

to label patients as alive or dead after seven years of follow up. The inclusion of

rs61757955 variant status, rs34988193 variant status, GRB2 expression and ANKDD1a

expression significantly improves the survival prediction compared to the eleven

covariates bolded in red from table 1 alone (p=0.0279).

Table Legends

Table 1. List of variables that are known to be associated with differences in survival in

low grade glioma patients. 11 variables (red) were selected by Lasso for inclusion in the

survival model. We used these 11 variables as covariates in our Cox regression model

when testing each germline variant.

Table 2. Description of the prognostic germline variants identified in this study.

(A) A description of the two prognostic germline variants (FDR<0.10) in the TCGA

dataset identified when testing all 196,022 germline variants.

(B) A description of the prognostic germline variant (FDR<0.10) rs34988193 in

ANKDD1a identified when the analysis was restricted to only germline variants with a

combined annotation dependent depletion (CADD) score greater than 30 in the TCGA

and CGGA datasets.

Table 3. The association between the germline variant rs61757955 and genomic and

histological variables. Patients were divided based on whether or not they had the

germline variant rs61757955. Patients with the germline variant rs61757955 were more

likely to have *CIC* mutated gliomas and the 1p/19q co-deletion.

609
610

References

1. Louis DN, Perry A, Reifenberger G, et al. The 2016 World Health Organization Classification of Tumors of the Central Nervous System: a summary. *Acta Neuropathol*. 2016. doi:10.1007/s00401-016-1545-1

2. Bauman G, Fisher B, Watling C, Cairncross JG, Macdonald D. Adult Supratentorial Low-Grade Glioma: Long-Term Experience at a Single Institution. *Int J Radiat Oncol Biol Phys*. 2009. doi:10.1016/j.ijrobp.2009.01.010

3. Jiao Y, Killela PJ, Reitman ZJ, et al. Frequent ATRX, CIC, FUBP1 and IDH1 mutations refine the classification of malignant gliomas. *Oncotarget*. 2012;3(7):709-722. doi:10.18632/oncotarget.588

4. Network CGAR, Brat DJ, Verhaak RGW, et al. Comprehensive, Integrative Genomic Analysis of Diffuse Lower-Grade Gliomas. *N Engl J Med*. 2015. doi:10.1056/NEJMoa1402121

5. Yan W, Zhang W, You G, et al. Molecular classification of gliomas based on whole genome gene expression: A systematic report of 225 samples from the Chinese Glioma Cooperative Group. *Neuro Oncol*. 2012. doi:10.1093/neuonc/nos263

6. Qian Z, Li Y, Fan X, et al. Prognostic value of a microRNA signature as a novel biomarker in patients with lower-grade gliomas. *J Neurooncol*. 2018. doi:10.1007/s11060-017-2704-5

7. Hu X, Martinez-Ledesma E, Zheng S, et al. Multigene signature for predicting prognosis of patients with 1p19q co-deletion diffuse glioma. *Neuro Oncol*. 2017. doi:10.1093/neuonc/now285

8. Zhou H, Vallières M, Bai HX, et al. MRI features predict survival and molecular markers in diffuse lower-grade gliomas. *Neuro Oncol*. 2017. doi:10.1093/neuonc/now256

9. Guerrini-Rousseau L, Dufour C, Varlet P, et al. Germline SUFU mutation carriers and medulloblastoma: Clinical characteristics, cancer risk, and prognosis. *Neuro Oncol*. 2018. doi:10.1093/neuonc/nox228

10. Dudley B, Karloski E, Monzon FA, et al. Germline mutation prevalence in individuals with pancreatic cancer and a history of previous malignancy. *Cancer*. 2018. doi:10.1002/cncr.31242

11. Zhang J, Walsh MF, Wu G, et al. Germline Mutations in Predisposition Genes in Pediatric Cancer. *N Engl J Med*. 2015. doi:10.1056/NEJMoa1508054

12. Carter H, Marty R, Hofree M, et al. Interaction landscape of inherited polymorphisms with somatic events in cancer. *Cancer Discov*. 2017. doi:10.1158/2159-8290.CD-16-1045

13. Huang K lin, Mashl RJ, Wu Y, et al. Pathogenic Germline Variants in 10,389 Adult Cancers. *Cell*. 2018. doi:10.1016/j.cell.2018.03.039

14. Baretta Z, Mocellin S, Goldin E, Olopade OI, Huo D. Effect of BRCA germline mutations on breast cancer prognosis: A systematic review and meta-analysis. *Medicine (Baltimore)*. 2016. doi:10.1097/MD.0000000000004975

15. Lau JW, Lehnert E, Sethi A, et al. The cancer genomics cloud: Collaborative, reproducible, and democratized - A new paradigm in large-scale computational research. *Cancer Res*. 2017. doi:10.1158/0008-5472.CAN-17-0387

657   16.   Ceccarelli M, Barthel FP, Malta TM, et al. Molecular Profiling Reveals
658   Biologically Discrete Subsets and Pathways of Progression in Diffuse Glioma.
659   *Cell*. 2016. doi:10.1016/j.cell.2015.12.028
660   17.   Lai Z, Markovets A, Ahdesmaki M, et al. VarDict: A novel and versatile variant
661   caller for next-generation sequencing in cancer research. *Nucleic Acids Res*. 2016.
662   doi:10.1093/nar/gkw227
663   18.   Li H, Handsaker B, Wysoker A, et al. The Sequence Alignment/Map format and
664   SAMtools. *Bioinformatics*. 2009. doi:10.1093/bioinformatics/btp352
665   19.   Wang K, Li M, Hakonarson H. ANNOVAR: Functional annotation of genetic
666   variants from high-throughput sequencing data. *Nucleic Acids Res*. 2010.
667   doi:10.1093/nar/gkq603
668   20.   Ellrott K, Bailey MH, Saksena G, et al. Scalable Open Science Approach for
669   Mutation Calling of Tumor Exomes Using Multiple Genomic Pipelines. *Cell Syst*.
670   2018. doi:10.1016/j.cels.2018.03.002
671   21.   Ramaswami G, Li JB. RADAR: A rigorously annotated database of A-to-I RNA
672   editing. *Nucleic Acids Res*. 2014. doi:10.1093/nar/gkt996
673   22.   Purcell S, Neale B, Todd-Brown K, et al. PLINK: A Tool Set for Whole-Genome
674   Association and Population-Based Linkage Analyses. *Am J Hum Genet*. 2007.
675   doi:10.1086/519795
676   23.   Friedman AJ, Hastie T, Simon N, Tibshirani R, Hastie MT. Lasso and Elastic-Net
677   Regularized Generalized Linear Models. *Available online https//cran.r-*
678   *project.org/web/packages/glmnet/glmnet.pdf (Verified 29 July 2015)*. 2015.
679   24.   Therneau TM, T. Lumley. Package ' survival .' *R Top Doc*. 2015.
680   doi:10.1016/j.jhydrol.2011.07.022.
681   25.   *Package "survminer" Type Package Title Drawing Survival Curves Using*
682   *"Ggplot2."*; 2018. https://cran.r-
683   project.org/web/packages/survminer/survminer.pdf. Accessed August 27, 2018.
684   26.   Subramanian A, Tamayo P, Mootha VK, et al. Gene set enrichment analysis: A
685   knowledge-based approach for interpreting genome-wide expression profiles. *Proc*
686   *Natl Acad Sci*. 2005. doi:10.1073/pnas.0506580102
687   27.   Kircher M, Witten DM, Jain P, O'roak BJ, Cooper GM, Shendure J. A general
688   framework for estimating the relative pathogenicity of human genetic variants. *Nat*
689   *Genet*. 2014. doi:10.1038/ng.2892
690   28.   Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics
691   Analysis Version 7.0 for Bigger Datasets. *Mol Biol Evol*. 2016.
692   doi:10.1093/molbev/msw054
693   29.   Pollard KS, Hubisz MJ, Rosenbloom KR, Siepel A. Detection of nonneutral
694   substitution rates on mammalian phylogenies. *Genome Res*. 2010.
695   doi:10.1101/gr.097857.109
696   30.   Rice P, Longden L, Bleasby A. EMBOSS: The European Molecular Biology Open
697   Software Suite. *Trends Genet*. 2000. doi:10.1016/S0168-9525(00)02024-2
698   31.   The 1000 Genomes Project Consortium. A global reference for human genetic
699   variation. *Nature*. 2015. doi:10.1038/nature15393
700   32.   Gleize V, Alentorn A, Connen De Kérillis L, et al. CIC inactivating mutations
701   identify aggressive subset of 1p19q codeleted gliomas. *Ann Neurol*. 2015.
702   doi:10.1002/ana.24443

33. Bailey MH, Tokheim C, Porta-Pardo E, et al. Comprehensive Characterization of
Cancer Driver Genes and Mutations. *Cell*. 2018;173(2):371-385.e18.
doi:10.1016/j.cell.2018.02.060
34. Pesenti C, Gusella M, Sirchia SM, Miozzo M. Germline oncopharmacogenetics, a
promising field in cancer therapy. *Cell Oncol*. 2015. doi:10.1007/s13402-014-
0214-4
35. Pinto N, Cohn SL, Dolan ME. Using germline genomics to individualize pediatric
cancer treatments. *Clin Cancer Res*. 2012. doi:10.1158/1078-0432.CCR-11-1938
36. Schärfe CPI, Tremmel R, Schwab M, Kohlbacher O, Marks DS. Genetic variation
in human drug-related genes. *Genome Med*. 2017. doi:10.1186/s13073-017-0502-5
37. Lu C, Xie M, Wendl MC, et al. Patterns and functional implications of rare
germline variants across 12 cancer types. *Nat Commun*. 2015.
doi:10.1038/ncomms10086
38. Koire A, Katsonis P, Lichtarge O. REPURPOSING GERMLINE EXOMES OF
THE CANCER GENOME ATLAS DEMANDS A CAUTIOUS APPROACH
AND SAMPLE-SPECIFIC VARIANT FILTERING. *Pac Symp Biocomput*. 2016.
39. Gay B, Suarez S, Caravatti G, Furet P, Meyer T, Schoepfer J. Selective Grb2 SH2
inhibitors as anti-Ras therapy. *Int J Cancer*. 1999. doi:10.1002/(SICI)1097-
0215(19991008)83:2<235::AID-IJC15>3.0.CO;2-B
40. Giubellino A, Burke TR, Bottaro DP. Grb2 signaling in cell motility and cancer.
Expert Opin Ther Targets. 2008. doi:10.1517/14728222.12.8.1021
41. Haines E, Saucier C, Claing A. The adaptor proteins p66Shc and Grb2 regulate the
activation of the GTPases ARF1 and ARF6 in invasive breast cancer cells. *J Biol
Chem*. 2014. doi:10.1074/jbc.M113.516047
42. Yu GZ, Chen Y, Wang JJ. Overexpression of Grb2/HER2 signaling in Chinese
gastric cancer: Their relationship with clinicopathological parameters and
prognostic significance. *J Cancer Res Clin Oncol*. 2009. doi:10.1007/s00432-009-
0574-8
43. Lowenstein EJ, Daly RJ, Batzer AG, et al. The SH2 and SH3 domain-containing
protein GRB2 links receptor tyrosine kinases to ras signaling. *Cell*. 1992.
doi:10.1016/0092-8674(92)90167-B
44. Bolouri H, Farrar JE, Triche T, et al. The molecular landscape of pediatric acute
myeloid leukemia reveals recurrent structural alterations and age-specific
mutational interactions. *Nat Med*. 2018. doi:10.1038/nm.4439
45. Feng J, Zhang Y, She X, et al. Hypermethylated gene ANKDD1A is a candidate
tumor suppressor that interacts with FIH1 and decreases HIF1α stability to inhibit
cell autophagy in the glioblastoma multiforme hypoxia microenvironment.
Oncogene. August 2018:1. doi:10.1038/s41388-018-0423-9
46. Sedgwick SG, Smerdon SJ. The ankyrin repeat: A diversity of interactions on a
common structural framework. *Trends Biochem Sci*. 1999. doi:10.1016/S0968-
0004(99)01426-7
47. Shi DJ, Ye S, Cao X, Zhang R, Wang KW. Crystal structure of the N-terminal
ankyrin repeat domain of TRPV3 reveals unique conformation of finger 3 loop
critical for channel function. *Protein Cell*. 2013. doi:10.1007/s13238-013-3091-0
48. Sullivan JM, Zimanyi CM, Aisenberg W, et al. Novel mutations highlight the key
role of the ankyrin repeat domain in TRPV4-mediated neuropathy. *Neurol Genet*.

2015;1(4):e29. doi:10.1212/NXG.0000000000000029
49. Lamandé SR, Yuan Y, Gresshoff IL, et al. Mutations in TRPV4 cause an inherited
arthropathy of hands and feet. *Nat Genet*. 2011. doi:10.1038/ng.945
50. Li J, Mahajan A, Tsai MD. Ankyrin repeat: A unique motif mediating protein-
protein interactions. *Biochemistry*. 2006. doi:10.1021/bi062188q
754

Figure 1

Figure 2

Figure 3

Figure 4

Table 1

Covariate		Median (Min-Max) or Number of Patients
Age		41 (14 - 87)
Gender	Female	250
	Male	200
Somatic Mutation Count		50 (0 - 12255)
Percent Aneuploidy		11% (5.2E-4% - 95%)
log(TERT Expression)		1.0 (0.0 - 9.1) FPKM
Principle Component 1 (PC1)		0.043 (-0.091 - 0.064)
Principle Component 2 (PC2)		-0.017 (-0.23 - 0.17)
Principle Component 3 (PC3)		-0.53 (1.34E-4 - 0.33)
Histological Type	**Astroctyoma**	172
	Oligoastrocytoma	113
	Oligodendroglioma	165
Tumor Location	Frontal Lobe	265
	Temporal Lobe	125
	Parietal Lobe	42
	Other	18
Grade	G2	212
	G3	237
	Cannot Be Assessed	1
Treatment Site	**Henry Fords Hospital**	82
	Case Western St. Joes	90
	Other	278
IDH Mutant	Wild Type	83
	Mutant	367
1p/19q Co-deletion	Absent	303
	Present	147
MGMT Promoter Methylation	Unmethylated	80
	Methylated	370
Chr 7 gain/Chr 10 loss	Absent	399
	Present	51
TP53 Mutant	Absent	232
	Present	218

Table 2

A

Variant	Chrom	Pos	Ref	Alt	Population Frequency	Sample Size	Number of Heterozygotes	Number of Homozygotes	Gene Name	Median Expression (FPKM)	p-value	Hazard Ratio
rs61757955	17	75318086	A	G	5.01%	291	21	0	GRB2	42.2	7.08E-10	20.4
rs28672782	11	1446622	C	T	16.27%	50	15	5	BRSK2	7.29	< 1E-16	1.15E-10

B

Variant	Chrom	Pos	Ref	Alt	Population Frequency	CADD Score	PhyloP Score	Gene Name	Dataset	Sample Size	Number of Heterozygotes	Number of Homozygotes	p-value	Hazard Ratio
rs34988193	15	64943580	A	G	30.90%	32	8.42	ANKDD1a	TCGA	450	199	52	0.00113	1.73
									CGGA	76	18	2	0.0743	1.79

Variable	Mean or Percentage (Wild Type)	Mean or Percentage (Mutant)	p-value
CIC Mutated	**15.9%**	**38.1%**	**0.017**
1p/19q Co-deletion	**25.2%**	**47.6%**	**0.038**
Oligodendroglioma	33.7%	42.9%	0.475
Total Somatic Mutation Count	30.9	30.0	0.766
Percent Aneuploidy	15.1%	11.7%	0.524
Astrocytoma	38.1%	42.9%	0.651
Grade 3	53.0%	42.9%	0.497
IDH Mutated	78.1%	85.7%	0.583
1p/19q Co-deletion	25.2%	47.6%	0.038
MGMT Promoter Methylation	77.8%	81.0%	1.000
Chr 7 Gain/Chr 10 Loss	13.0%	9.5%	1.000
Expression of GRB2 (FPKM)	45.7	44.4	0.636