Rapid evolution by spontaneous mutation increases genetic diversity facilitating plant

population survival

Henning Nottebrock[1], Mao-Lun Weng[1], Matthew T. Rutter[2] & Charles B. Fenster[1]

[1]Department of Biology and Microbiology, South Dakota State University, Brookings, SD USA

[2]Department of Biology, College of Charleston, Charleston, SC USA

Abstract

Using a mechanistic eco-evolutionary trait-based neighborhood-model, we quantify the impact

of mutations on spatial interactions to better understand the potential effect of niche evolution

through mutations on the population dynamics of *Arabidopsis thaliana*. We use 100 twenty-

fifth generation mutation accumulation (MA) lines (genotypes) derived from one founder

genotype to study mutational effects on neighbor responses in a field experiment. We created

individual-based maps (15,000 individuals), including phenotypic variation, to quantify

mutational effects within genotypes versus between genotypes on reproduction and survival.

At small-scale, survival is enhanced but reproduction is decreased when a genotype is

surrounded by different genotypes. At large-scale, seed set is facilitated by different genotypes

while the same genotype has either no effect or negative effects. Mutations may provide a

mechanism for plants to quickly evolve niches and may drive competition, facilitation and

selection with profound consequences for future population and community dynamics.

Introduction

A complete understanding of population and community dynamics requires linking intraspecific

genetic diversity with spatial ecological interactions (Bolnick et al. 2011, Genung et al. 2011).

Evolutionary mechanisms -- mutation, drift, gene flow and selection -- are responsible for

intraspecific genetic variation contributing to both ecological structure and species diversity

(Hart et al. 2016). These community properties impact ecological mechanisms such as

competition and facilitation (Whitlock 2014). Competition and facilitation in turn affect

population and community dynamics that depend on variation in demographic rates (Chase et

al 2002, Solivers et al 2015). Demographic rates are influenced by spatial interactions of genetic

variation and this variation may contribute to species coexistence: genotypes can hinder or

favor the survival of each other by using similar or different resources with negative or positive

consequences for coexisting species (Hart et al. 2016, Hausch et al. 2018). Intraspecific genetic

variation contributing to individual trait variation and environmental adaptability may promote

species coexistence by both increasing habitat heterogeneity and altering competitive

hierarchies (Violle et al. 2012, Ehlers et al. 2016). However, intraspecific genetic variation may

hinder species coexistence when intraspecific genetic variation is diminished by competition

between individuals (Hart et al 2016). To fully understand the causes of coexistence

mechanisms, we need to integrate intraspecific evolutionary mechanisms with intraspecific

ecological mechanisms underlying spatial species interactions (Bolnick et al. 2011, Ehlers et al.

2016).

The number of studies exploring eco-evolutionary dynamics and their potential

feedbacks on population and community dynamics has dramatically increased over the last two

decades (Shefferson and Salguero-Gomez 2015). An important current issue in community and

ecosystem genetics research is determining the relevance of intraspecific genetic variation and

genetic differentiation (divergence) to ecological and evolutionary processes at the community

and ecosystem level (Genung et al. 2011, Pujol et al. 2018). Quantifying intraspecific trait

variation defining the fundamental niches of species is an important link between ecology and

evolution (Violle and Jiang 2009). Spatial structure is another important component of realistic

eco-evolutionary dynamics, such that in spatially structured populations, selection is

determined by the interplay between demographic and genetic structures (Lion et al. 2011).

Demographic structure describes the spatial distribution of individuals through birth, death and

migration resulting in spatial patterns, while genetic structure describes the spatial distribution

of genotypes. The amount and spatial pattern of genetic variation may constrain evolution of

traits influencing competitive ability of individuals (Wilson 2014). Moreover, competitive

abilities depend on species' niches. The strength of competition is determined by how much

individual niches overlap with each other (Hutchinson 1957, Holt 2009) and how long

coexistence has occurred (Connell 1980).

Species niche breadths are reflected by individual trait variability and intraspecific and

interspecific genetic and spatial interactions driven by environmental conditions. Thus,

intraspecific trait variability shaped by intraspecific genetic variation can influence ecological

mechanisms driving variation among individual persistence (Lankau 2009). Only when the

spatial relationship between traits and demographic parameters is explicitly described can

robust hypotheses about the effects of individual variation on competitive outcomes be

accurately formulated (Hart et al. 2016). Especially in controlled common garden environments,

genetic variation in one species can have predictable and heritable effects on associated

communities and ecosystems (Carr and Dudash 1995; Whitham et al. 2003, 2006; Johnson &

Stinchcombe 2007; Bailey et al. 2009; Johnson, Vellend & Stinchcombe 2009). Hence, coupling

evolutionary genetics with community ecology may advance our understanding of species

interactions and population and community dynamics (Baron et al. 2016), especially for

populations and communities that suffer environmental change (Bellard et al. 2012).

An important gap in our understanding of the link between evolutionary and ecological

processes is the nearly complete absence of data quantifying how rapidly genetic variation

governing within species competitive hierarchies evolves (Hausch et al. 2018). An important

source of novel population genetic variation is mutation. Therefore, we aim to investigate the

link between evolutionary genetics and spatial ecological interactions of mutation accumulation

lines (MA lines) in the model plant organism *Arabidopsis thaliana* (Brassicaceae) to advance our

understanding of the genetic origins of population and community dynamics, contributing to a

fuller understanding of the maintenance of biodiversity. We take advantage of 25th generation

A. thaliana mutation accumulation (MA) lines that were planted under field conditions in years

2004 and 2005 with spatial records of each individual each year (e.g., Rutter et al. 2010; 2012,

Rutter et al. 2018) (Fig. 1).

Here we focus on the neighborhood effects of MA lines on focal plants from the same

MA line (within genotypes) or different MA lines (between genotypes) to quantify the evolution

of intraspecific genetic variation through the accumulation of spontaneous mutations. We

implement trait-based neighborhood models (Nottebrock et al. 2017a, Nottebrock et al. 2017b,

Lachmuth et al. 2017) to understand ecological mechanisms of within and between MA lines

genetic variation to examine eco-evolutionary dynamics (Lion 2018). Specifically, we ask: 1) Do

spontaneous mutations rapidly introduce enough genetic variation among MA lines to influence

competition and facilitation of *A. thaliana*? 2) Do spontaneous mutations of MA lines

contribute to competition, facilitation and selection promoting differential MA line survival and

reproduction at different spatial scales? 3) How may competitive and facilitative effects

potentially determine population and community dynamics?

Methods

MA lines are generated from a single nearly homozygous individual founder and cultivated via

limited effective population number. In the case of *A. thaliana*, this occurs through single seed

descent, resulting in Ne= 1. Thus typical MA line cultivation results in an unbiased sample of

mutation effects ranging from deleterious to advantageous, although lethal mutations are

excluded (Lynch and Walsh 1998). Each MA line accumulates independent spontaneous

mutations. After the propagation of a set of MA lines through multiple generations, the genetic

differences among the MA lines and between those lines and the founder reflect the input of

mutation. Significant MA line effects for multiple traits, including performance and trait

measures, were found under both field and greenhouse conditions (Rutter et al. 2010; Roles et

al. 2016, Rutter et al. 2018). Each of the MA lines in our experiment is fixed for an average of 20

different sequence level mutations, single nucleotide mutations (SNMs) and indels combined

(Ossowski et al 2010, Rutter et al. 2012, Weng et al. 2019).

MA lines and field experiments

We used survival and seed set data of *A. thaliana* MA lines and the founder as assessed in field

experiments in 2004 and 2005 from Rutter et al. (2010, 2012 and 2018) planted in a

randomized design (Fig. 1). Rutter et al. (2010, 2012) planted seedlings of 100 MA lines and the

founder at the four-leaf stage, approximately two weeks post germination, into a secondary

successional field at Blandy Experimental Farm (BEF) in Virginia (39°N, 78°W). Each of the 100

MA lines was used to found up to five sublines to minimize biases due to maternal effects

introduced by the specific location within the greenhouse. We founded six sublines from each

of the six lines representing the premutation founder genotype. In 2003, subline plants were

used to generate all seed utilized in all field experiments. In each planting, 7504 individuals

were planted, 7000 individuals of 100 MA lines (70 replicates per MA line, 14 replicates per

subline) and 504 individuals of the founder (14 replicates per subline). The planting

environment corresponds to a spring ephemeral life-history, where plants germinate and

complete the life-cycle in the spring. At the time of planting, vegetation was scant but present.

By harvest, the A. thaliana individuals were dwarfed by naturally occurring vegetation.

The plot was arranged in 14 spatial blocks with each containing 12 subblocks (Fig. 1B)

(total plot area approximately 35 x 25 m). Each block included one seedling from each subline

and in total 7504 individuals. We used the spatial information of each individual within the

described design and created a raster of all plant individuals with R packages (raster, maps,

maptools). We used individual-based maps neighborhood matrices with exact spatial and trait

information of each genotype. If all five sublines did not produce enough seedlings to distribute

in all blocks, seedlings from other sublines of the same MA line were overrepresented in blocks

to maintain the same overall number of plants per MA line. Plants dying within the first 3 days

of transplant (about 50 plants) were considered to have died from transplant shock and were

replaced with another plant from the same MA line. Plants were censused weekly for survival.

Plants were harvested by late May, by which time they had senesced. In 2004 a total of 5915

individuals including 394 founders survived. In 2005, a total of 4506 individuals survived

including 302 founders. Plants were oven dried and biomass was measured. All fruits produced

by each plant were counted and in combination with seed production used as the

measurements of seed set. For our analysis, we used measurements of two response variables

representing an important part of the life history of *A. thaliana* to calculate direct

neighborhood effects on: a) the survival rate of plant individuals and b) individual seed set from

plants that survived and produced fruits.

Statistical Analyses

We use eco-evolutionary trait-based neighborhood models that include intraspecific genetic

variation and phenotypic variation expressed by plant trait biomass to analyze competition and

facilitation between individuals of *A. thaliana*. We analyzed plant survival rate and focal seed

set measurements from years 2004 and 2005 separately. For each year, we considered all

individuals as focal plants in the analysis based on individual-based neighborhood matrices to

analyze 1) rate of plant survival to reproduction and 2) seed set of those plants that survived to

reproduction. We analyzed neighboring plants to focal plants in a radius of 80 cm (small-scale)

or 200 cm (large-scale) of a given focal plant as two spatial scales in the neighborhood analyses

(Fig. 1) to quantify selection, competition or facilitation between plants depending on their

genotypic and phenotypic variation.

159   We used extensions of linear mixed models (package lme4, Bates et al. 2014) in R ver.

3.3.3 (www.r-project.org) to conduct neighborhood analyses of focal seed set and survival. We

assumed binomial errors for the analyses of plant survival and Poisson errors for analyses of

seed set. The mixed models described interactions among plants by including neighborhood

indices as explanatory variables at two spatial scales in separate models. Neighborhood indices

are spatial density effects of surrounding neighborhood plants that affect focal seed set and

survival. For each plant, we used the Euclidian distance between the focal plant and the

neighboring plants to compute response effects of intra- (same genotype) or inter-genotypic

(different genotype) neighbors in a given radius around focal plants (Nottebrock et al. 2017).

Moreover, we used a neighborhood index that accounts for the decline of neighbor effects with

distance from the focal plant (Uriarte et al. 2010) and summed the amount of biomass from all

individuals in a radius of 80 cm or 200 cm respectively by a Gaussian interaction kernel

(Lachmuth et al. 2018, Nottebrock et al. 2017, Damgaard 2004). We used random effects of

block and subblock to correct for environmental variation between local heterogeneous

conditions. Importantly, we correct for between MA line effects by including MA lines as a

random effect. In addition, including a random slope of biomass on each random intercept

corrects for the intraspecifc phenotypic variation depending on local conditions of plant focal

individuals. Moreover, the weighted neighborhood density by plant biomass accounts for

environmental variation between neighboring plants. Our model parameters and a detailed

model description of subblock and block models can be found in S1 supplementary text

(Supplementary Material).

Neighborhood matrices of all individuals (individual-based maps) were used to analyze

the effect of intra- and inter-genotypic neighbors on survival and focal seed set with spatial

interaction kernels of neighborhood (plant biomass) density. By incorporating different

genotypes and phenotypic variation, we can quantify how important genetic variation is for

neighborhood models and if the phenotypic variation explains spatial interactions between

individuals. We assume the consequences of genetic differences to be larger between MA lines

than between any MA line with the founder. This is a valid assumption since each MA line

differs from the other by approximately 20 + 20 = 40 mutations, while any two replicates within

a MA line will differ by one generation, $\leq$ 2 mutations. Thus, we simulated line effects from

parameters derived from MA lines as random effects with the R package 'merTools' and the

function 'plotREsim' in R 2018.

We used the trait values of neighbors to calculate trait-based neighborhood indices

including plant biomass as a trait (Goldberg & Fleetwood, 1987; Goldberg & Landa, 1991; Cahill

et al., 2005). We fitted eco-evolutionary trait-based neighborhood models at two different

spatial scales for response variables (survival and seed set) for each of the two and both years.

To address our objectives, we first analyzed models with differential effect in which intra-

genotypic neighbors (within all MA lines and founder) had a different effect on survival (A1, A2,

Table 1) and seed set (B1, B2, Table1) than inter-genotypic neighbors at small-scale (s) and

large-scale (l). In addition, we analyzed models with neutral effects on survival (A1, A2, Table 1)

and seed set (B1, B2, Table 1) that included total neighbor density without the split between

intra-genotypic and inter-genotypic neighbors at small-scale (s) and large-scale (l). To this end,

all models were fitted with two separate neighborhood indices that were calculated from intra-

and inter-genotypic neighbors. To justify the inclusion of individual plant biomass as trait-values

for interacting plants in the model, we used AICc to compare the models with and without the

trait-proxy (Burnham and Anderson 2002). We found that all models perform better including

biomass as a trait-proxy (ΔAICc > 2). All eco-evolutionary trait-based neighborhood models

contained random effects of subblock nested in block at block scale and subblock scale on the

intercept, MA line identity on the intercept and the focal trait-value (plant biomass) on the

slope. Additionally, because direct environmental variables were not measured during the field

experiments, we included in each model the individual's biomass to correct for environmental

conditions for spatial autocorrelation. All variables are scaled and centered to assure

comparability between predictor variables. Models of differential and neutral effects for 2004

and 2005 (Table 1, A1-A2, B1-B2) are fitted at small-scale (80cm scale) and at large-scale

(200cm scale). Hereafter, the 80 cm scale models are referred to as "small-scale" models and

the 200 cm scale models are referred to as "large scale" models. Neighborhood indices, intra-

and intergenotypic variation and total variation of biomass density are included as inverse

density variables (1/1+density). We compared models of differential and neutral effects

through likelihood ratio tests (LRTs).

Results

Focal plants surrounded by plants of the same genotypes varied from 0 to 6 individuals at small-

scale (sub-block) and 5 to 6 individuals at large-scale (block). After determining that eco-

evolutionary trait-based models perform better when including plant biomass as a trait proxy to

calculate trait-based neighborhood indices we found that all models including plant biomass

perform better than models including only neighborhood density (ΔAIC > 2, Supplementary

Material S2).

Weighted neighbor effects of biomass density on genotypes for plant survival

At small-scale, survival rate is larger when surrounded by inter-genotypic than intra-genotypic

neighbors for both years 2004 and 2005 (model A1 and A2; Table 2 and Fig. 2a). This difference

of survival rate between intra- and inter-genotypic plants indicates that inter-genotypic

neighbors select for genotypic diversity and show stronger competitive effects on the survival

of intra-genotypic neighbors at small-scale. This finding is demonstrated by the superior

performance of the differential model relative to the neutral model (year 2004: LRT: χ^2_{1DF} =

5.21, p < 0.05; year 2005: LRT: χ^2_{1DF} = 14.73, p < 0.001). The differential and neutral models of

survival at large-scale have only non-significant effects (Table 2). In addition, comparing the AIC

we found that models at large-scale perform worse than at small-scale for both 2004 and 2005

(Table 2). We therefore will only discuss small-scale effects on survival. The distance kernel

(alpha) at small-scale shows that neighboring plants of different genotypes reduce plant

survival by 50% at 53 cm in 2004 and 48 cm in 2005.

Weighted neighbor effects of biomass density on genotypes for seed set

At both small and large scales, the differential model performs better than the neutral model:

at small scales (year 2004: LRT: χ^2_{1DF} = 52.48, p < 0.001; year 2005: LRT: χ^2_{1DF} = 17.37, p < 0.001;

respectively) and at large scale (year 2004; LRT: χ^2_{1DF} = 679.44, p < 0.001; year 2005: LRT: χ^2_{1DF} =

21.17, p < 0.001; respectively). The small-scale model of seed set that included weighted

biomass densities demonstrates the reduction of seed set is larger when surrounded by inter-

genotypic than intra-genotypic neighbors for both years 2004 and 2005 (model B1 and B2;

Table 2 and Fig. 2a). The difference between intra- and inter-genotypic plants indicates that

inter-genotypic neighbors have a stronger competitive effect on seed set than intra-genotypic

neighbors at small-scale. In contrast, at large-scale seed set increases when surrounded by

inter-genotypic neighbors but decreases when surrounded by intra-genotypic neighbors for

both years 2004 and 2005 (model B1 and B2; Table 2 and Fig. 2b). Comparing the AIC between

small and large-scales, models at large-scale for 2004 and 2005 perform better (Table 2).

Genotype effects on plant survival and seed set

The distance kernel at small-scale in 2004 indicates a reduction of seed set (competition)

whereas at large-scale are consistent with an increase of seed set (facilitation). Neighboring

plants from all genotypes of *A. thaliana* at small-scale reduce 50% of focal seed set at 23 cm

and at large-scale facilitate 50% of focal seed set at 86 cm. The estimation of genotype effects

(MA lines and founder) simulated as conditional means and expressed as odds ratios show

hierarchical orders of MA line and founder competitive effects at small-scale (Fig. 3). We found

no significance of simulated competitive effects of all MA lines and founder calculated from

survival and seed set models at small-scale (ANOVA, $p > 0.1$, Fig. 4a). In addition, we found no

significant correlation between competitive effects of MA lines and founder at large-scale. In

addition, we did not find selective effects of MA lines and founder on the survival at small-scale

(ANOVA, $p > 0.1$, Fig. 4b). However, we found strong correlation between competitive effects

of MA lines and founder at small-scale and large-scale indicating a genetic trade-off (ANOVA,

$F=460.07$, $P< 0.0001$, Fig. 4c).

Discussion

We are able to quantify individual spatial interactions between and within genotypes (MA lines

and founder) using a mechanistic eco-evolutionary trait-based neighborhood model. We

demonstrate intraspecific genetic variation due to spontaneous mutations can shape

competitive abilities of genetically different individuals *of A. thaliana*. Notably, these

differences arose in just 25 generations of mutation and in the absence of natural selection.

Many of the mutations differentiating the MA lines also appear in nature, contributing to *A.*

thaliana genetic polymorphism (Weng et al., 2019). Plant survival is higher when surrounded by

inter-genotypic neighbors supporting genetic diversity. Effects on focal seed sets were reversed

between small and large scales; inter-genotypic neighbors have stronger negative (competitive)

or positive (facilitative) effects than intra-genotypic neighbors on focal seed sets at small-scale

or large-scale, respectively. Moreover, at small-scale, competitive effects of different MA lines

have similar impacts on survival and seed set. At large-scale, the intra-genotypic effect on

survival turns into less competitive effects on seed set. In contrast, different genotypes show

stronger facilitative effects within the population at large-scale, maybe because plants are more

competitive against other species. Below we discuss how these scale effects provide insight on

community assembly and potential coexistence.

Selection, Competition and Facilitation

Because of stronger statistical signals of effect sizes between genotypes than within genotypes,

niche variation may result in niche expansion or reduction (Ehlers et al. 2016). However, these

competitive effects only occur between plants that survive to reproduction and affect seed-set.

The neighbor effect on plant survival suggests kin recognition despite our randomized field

experiment that likely reduced the spatial autocorrelation of genotypes (Hamilton 1964,

Kubisch et al. 2013). Seed-dispersal is limited in space because the *A. thaliana* fruit do not

dehiscently explode as they do in related species (Hofhuis and Hay 2017), although a seed bank

ensures some dispersal through time (Falahati-Anbaren et al. 2014). The neighbor effect

increasing genetic diversity of *A. thaliana* populations can have a positive effect on the

coexistence of competing species (Vellend 2006). If we assume that the increase of genetic

diversity due to spontaneous mutations of *A. thaliana* affects the extent of individual plant

niches, at large-scale the inclusion of other plant species should result in stronger competitive

effects of *A. thaliana* with other plant species (Hausch et al. 2018). We demonstrate that *A.

thaliana* individuals increase their competitive abilities with an increase of genetic diversity,

perhaps due to the expansion of their individual niches due to more variation in plant traits.

Antagonistic interactions between genotypes on different spatial scales may result in

genetic trade-offs when mutations are advantageous or deleterious, which over many

generations could provide a strong stabilizing force maintaining both species and genetic

diversity in this system and promote coexistence (Lankau 2008). Thus, spontaneous mutation

could also provide an additional evolutionarily stabilizing effect on community dynamics. Our

study demonstrates the potential for a fundamental evolutionary process, mutation, to have

profound consequences for community structure. When *A. thaliana* is rare, selection would

favor genotypes that compete well and enhance the population's survival relative to that of its

interspecific competitor (Lankau 2009). In contrast, when *A. thaliana* is common and the

interspecific competitor is rare, selection would favor *A. thaliana* genotypes that are good

intraspecific competitors. This trade-off may result in a decrease in the interspecific competitive

ability of *A. thaliana*, effectively increasing the competitor's fitness relative to that of *A.*

thaliana (Lankau 2009, Lankau and Strauss 2007). However, the genetic trade-off between

intra-and interspecific competitive abilities due to mutation remains unknown because our

dataset only includes performance of *A. thaliana* individuals without quantifying the

surrounding intraspecific environment (Chesson 2000, Adler et al. 2007).

Competitive Hierarchies of MA lines and Founder lines

Our result show scale dependent competition and facilitation (Nottebrock et al. 2017b)

providing further evidence that in a relative short evolutionary time scale spontaneous

mutations may change the competitive hierarchies between founders and specific MA lines due

to advantageous or deleterious mutations (Rutter et al. 2010). In contrast to Masclaux et al.

2010, we found differential responses to similar genotypes vs different genotypes in *A. thaliana*

depending on the neighbor effect on the plant survival or seed set. However, the interaction

between similar genotypes and different genotypes seems to depend on the strength of the

competitive abilities of the accessions (this study). Comparisons between MA lines and founder

demonstrate that competitive hierarchies follow different orders and competitive effects occur

at different life-history stages at different spatial scales. Eco-evolutionary processes might

reflect spatial selection for diversified genotypes because of niche evolution and individuals of

different genotypes that survived to reproduction have stronger competitive abilities (Ehlers et

al. 2016). Moreover, the hierarchical order of competitive effects between MA lines and

founders shows strong variation.

Eco-evolutionary dynamics

Populations can adapt evolutionarily to their environment on a time scale equivalent to that of

ecological processes and affect present day species interactions, i.e., coexistence (Fussmann et

al. 2007). Yet, sustained and rapid climate change could deplete genetic variance faster than it

can be replenished by mutation (Fournier-Level et al. 2016). In our study, we show that only 25

generations are enough to influence present day plant interactions of *A. thaliana* in the

presence of other plant species. Possible effects of mutations on trait mechanisms might reflect

the increased seed set of *A. thaliana* at large-scale, because of the presence of another plant

species allowing the establishment and growth of *A. thaliana* (Ehlers et al. 2016). Many traits

can influence competition and facilitation between individuals of different genotypes (e.g.

Chapin et al. 1993; Caradus and Woodfield 1998; Hausmann et al. 2005). Intraspecific

competition between *A. thaliana* individuals tends to be higher than between interspecific

competitors (Lankau and Strauss 2007, Van Dam and Baldwin 1998; Tiffin 2002, Strauss and

Irwin 2004). Such spatial mechanisms might be described by the scale difference of competition

at small-scale and facilitation at large-scale between different *A. thaliana* genotypes.

Competition and facilitation are therefore dependent on the extension or reduction of

individual niches, which is based on the ability to advance or distract traits due to spontaneous

mutations. However, we are not able to directly link our results to benefactor or antagonistic

plants, because we focused in our experiment on *A. thaliana* densities and fitness components.

Nevertheless, genotypic-specific interactions are well studied. A number of studies of *A.*

thaliana MA lines demonstrate mutation effects on plant traits such as leaf weight, flowering

time, trichome density, number of leaves at bolting, duration of vegetative and flowering

period (Camara and Pigliucci 1999, Rutter et al. 2010, Stearns and Fenster 2016). It is

conceivable that the increase in variance of these traits through spontaneous mutations could

contribute to the evolution of competitive hierarchies.

Consequences for population and community dynamics

Competition between genetically different individuals of *A. thaliana* at small-scale and

facilitation at large-scale has important implications for population structure (Cahill et al. 2005).

Plants may cooperate by competing less, or act selfishly by competing more (Dudley et al.

2013). However, kin recognition might be the result of kin selection when occurring in a

heterogeneous environment with various genotypes (Hamilton 1964). Especially, at large-scale,

we show that different genotypes facilitate seed set that stabilize the population dynamics by

increasing the performance of neighboring plant individuals (Latzel et al. 2013, Castellanos et al.

2014). Thus, selection for diversified genotypes might reveal a coexistence mechanism, because

mutations at a short time scale (25 generations) are able to alter the competitive hierarchies of

A. thaliana individuals of different genotypes. These genotypes might expand or reduce their

niches and can therefore act more competitive against strangers (Ehlers et al 2016). Moreover,

variation in competitive ability among genotypes due to mutation can lead to intransitive

competitive hierarchies at a small-scale, and allow coexistence of competitors at large-scale

when there is no single dominant competitor (Vellend and Geber 2005, Taylor and Aarssen

1990, Laure et al. 2017). This might also reflect the genetic trade-off between competitive

abilities of MA lines of small-scale competition and of large-scale facilitation (Timan 2004). Low

competition between *A. thaliana* individuals at small-scale transmits into low facilitation

between different species at larger scale. In turn, strong competition within *A. thaliana* at

small-scale results in weaker competition between different species at large-scale. Thus, the

constant input of mutation to genetic variation of competitive hierarchies suggests that genetic

variation is likely never a limiting factor in the evolution of those traits that influence plant

community dynamics.

Implementation of eco-evolutionary trait-based model for population and community ecology

In ecology, two primary contrasting models are crucial to population and community dynamics

for understanding the maintenance of biodiversity: the niche and neutral theories of ecology

(Fisher and Mehta 2015). The niche theory claims that niches are determined by the difference

of abiotic or biotic ecological processes e.g. competition or facilitation (Hutchinson 1957, Bruno

et al. 2003, Colwell and Rangel 2009) whereas the neutral theory predicts that ecological

processes are neutral and are solely modified by stochasticity (Hubbel 2005). Although it is still

under debate which model predicts the dynamic properties of communities and determines the

maintenance of biodiversity (Violle et al. 2017), both models miss the importance of

evolutionary mechanisms and individual variation influencing population and community

dynamics (Violle et al. 2012). The difference of competition and facilitation between or within

genotypes demonstrates that evolutionary mechanisms that shape individual variation could

change hierarchies of intra-specific interactions in natural plant populations over a short

evolutionary time scale. Our results are concordant with the niche theory stressing the

changing of competitive hierarchies among phenotypes in a given environment (Levine and

HilleRisLambers 2009, Nottebrock et al. 2017). Moreover, selection and trait evolution favor

population survival of *A. thaliana* in the wild, which stresses the importance of combining

spatial ecological and evolutionary mechanisms for our understanding of population and

community dynamics.

Conclusion

Rapid evolution of *A. thaliana* due to spontaneous mutation alone has profound consequences

for population and community dynamics. Competition of *A. thaliana* individuals is genotype-

dependent and *A. thaliana* is not a single weak competitor among co-occurring plant species

(Ehlers et al. 2016, Soliveres et al 2017). Moreover, the discovery of the underlying eco-

evolutionary nature of competition in *A. thaliana* supports a shift from species-based to

individual-based community ecology. This would lead to a more predictive ecological theory

(Violle et al 2017). We show that including the intraspecific genetic variation and phenotypic

variation as separate aspects to a trait-based neighborhood model increases the predictive

power to understand population and community dynamics. In particular, non-neutral

intraspecific processes may determine species coexistence, because genetic diversity is

promoted by having stronger competitive abilities at small-scale and stabilizing population

survival at large-scale (Clark et al. 2010). Additionally, our study demonstrates higher genetic

diversity increases population survival due to rapid evolution with implications to forecast the

fate of species and functional diversity in response to environmental changes (Violle et al 2012).

Depending on the time of environmental changes, species may adapt to environmental change

by shifting their fundamental niches (Clark 2010). The combination of intraspecific genotypic

variation and spatial interactions might advance our understanding of community dynamics,

especially of rapid evolution (Koch et al. 2014, Turcotte and Levine 2016). Incorporating genetic

variation and the eco-evolutionary process for determining standing levels of genetic variation

will provide a better understanding of species interactions underlying the maintenance of

biodiversity.

Keywords (10 for indexing purpose): trait-based community ecology, spontaneous mutations,

competition, facilitation, niche evolution, intraspecific genetic variation, intraspecific

phenotypic variation, *Arabidopsis thaliana*, mutation accumulation lines, eco-evolutionary

dynamics.

438
Figure 1. (A) 100 Mutation accumulation lines (here example of four MA-lines) of *Arabidopsis*

thaliana derived from a single founder (Columbia) after 25 generations with a mutation rate of

~0.7 per generation (Weng et al. 2019). Each MA line has a unique genotype that differs from

the founder. Fundamental niche of individual genotypes (different dashed and dotted lines) and

founder genotype (black line) are distributed within the hypothetical fundamental niche of a

population of *A. thaliana* (bold black line). Axes represent niche axis (resource-use,

environmental factors, etc.). (B) Planting design of *A. thaliana* in a natural field experiment

showing spatial biomass variation of individual survivors in 2004.

	Year 2004	Year 2005	Model
Individual survival	A1 (s,l)	A2 (s,l)	Differential
	A1 (s,l)	A2 (s,l)	Neutral
Seed set	B1 (s,l)	B2 (s,l)	Differential
	B1 (s,l)	B2 (s,l)	Neutral

Table 1. Table of all different eco-evolutionary trait-based neighborhood models including

intraspecific variation to test our three objectives of density effects on survival and seed set of

A. thaliana. Differential models split the neighbor responses into within genotypes and

between genotypes. Neutral models describe the overall neighbor response without

differentiating between genotypes. All models include random effects of plant biomass on the

slope, subblock or block and MA-line on the intercept (see Fig. 1 for design), and respectively

depending on small (s) or large-scale (l) (for more details see method section).

Survival model

(A1) 2004		(s) Small scale (80cm)		AIC	(l) Larger scale (200cm)		AIC
		Intra	*Inter*		*Intra*	*Inter*	
Weighted density							
MA Lines	neutral	-0.02 ^{ns}		7551.7	0.004 ^{ns}		7644.5
	differential	-0.02 ^{ns}	-0.37**	7548.6	*0.15* ^{ns}	*-0.30* ^{ns}	7644.2

(A2) 2005		(s) Small scale (80cm)		AIC	(l) Larger scale (200cm)		AIC
		Intra	*Inter*		*Intra*	*Inter*	
Weighted density							
MA lines	neutral	0.03 ^{ns}		9637.3	0.03 ^{ns}		9872.9
	differential	-0.02 ^{ns}	-0.28***	9626.1	0.05 ^{ns}	-0.10 ^{ns}	9873.3

Seed set model

(B1) 2004		(s) Small scale (80cm)		AIC	(l) Larger scale (200cm)		AIC
		Intra	*Inter*		*Intra*	*Inter*	
Weighted density							
MA lines	neutral	-0.02***		52982	0.03		52944
	differential	-0.02**	-0.07***	52948	*0.03****	*0.15****	52939

(B2) 2005		(s) Small scale (80cm)		AIC	(l) Larger scale (200cm)		AIC
		Intra	*Inter*		*Intra*	*Inter*	
Weighted density							
MA	neutral	-0.008*		49103	-0.008*		49049
lines	differential	-0.009**	-0.09***	49095	-0.01*	0.16***	49026

Table 2. The coefficient and AIC from eco-evolutionary trait-based neighborhood models of the

Arabidopsis thaliana field experiment in the years 2004 and 2005. Models include intraspecific

genetic variation defined as intra-genotyptic (within genotypes) and inter-genotypic (between

genotypes) variation. Models are shown at two different spatial scales: weighted subblock

density (80cm radius) and weighted block density (200cm radius). All scales include variables of

neutral effects (absence of intra- and inter-genotypic variation (total sum of all neighboring

genotypes)) and differential effects (presence of intra- and inter-genotypic variation (split of

neighboring genotypes in intra- and inter-genotypic genotypes). Thus, neighbor identity is split

into intra- and inter-genotypes as differential effects or are combined as neutral effects, which

includes all neighbors. P-value levels <0.05*, <0.005**, <0.001*** and >0.05[ns].

Figure 2. The barplots show standardized coefficient effect sizes and associated standard errors

of eco-evolutionary trait-based neighborhood models at small-scale (a, c) and large-scale (b, d)

for all MA-lines for the survival model (a, b) and the seed set model (c, d). Intra-genotypic

effects (yellow) and inter-genotypic effects (grey) are presented for 2004 and 2005. Coefficient

effect sizes indicate reduction or an increase of seed set depending on biomass density of intra-

or inter-genotypic neighbors. In addition, coefficient effect sizes indicate a reduction of survival

of intra or inter-genotypic neighbors.

MA line effect sizes of seed set (a), (b) and survival (c) model

Figure 3. The effect range on seed set (a), (b) and survival (c) computed from eco-evolutionary

trait-based neighborhood models shown as odds ratios for year 2004 (for 2005 see

Supplementary Material). (a) Shows the hierarchical competitive order of MA lines and founder

for the seed set model including range effects of each MA line and founder as random

intercepts and the trait proxy biomass as random slopes at small-scale and (b) at large-scale. (c)

Shows the hierarchical selective order of MA lines and founder for the survival model including

range effects of each MA line and founder as random intercepts at small-scale.

Figure 4. Correlation of the simulated conditional mean represented as odds ratio of same

genotypes (MA lines) reducing or increasing the effect sizes of the seed set model and of the

survival model at small-scale or large-scale for 2004, respectively. For year 2005 see

supplementary material. Each point represents different genotypes of *A. thaliana.* Negative

values means lower effect sizes on the survival or seed set and positive values means increasing

the effect sizes on the survival or seed set. a) Relationship between survival and seed set

model. b) Relationship between survival and seed set model. c) Relationship between small-

scale survival and large-scale seed set model. Since we are plotting effect sizes, the positive

correlation in panel c reflects a trade-off between competition and facilitation (see results).

References

Adler P., B., HilleRisLambers Janneke & Levine J. M. (2007). A niche for neutrality. *Ecology Letters*,

10, 95–104.

Agrawal, A.A., Hastings, A.P., Johnson, M.T.J., Maron, J.L. & Salminen, J.-P. (2012). Insect

herbivores drive real-time ecological and evolutionary change in plant populations. *Science*,

338, 113–116.

Antonovics, J. (1976). The Nature of Limits to Natural Selection. *Annals of the Missouri Botanical*

Garden, 63, 224–247.

Bailey JKet al.2009From genes to ecosystems: a synthesis of the effects of plant genetic factors

across levels of organization. Phil. Trans. R. Soc. B **364,** 1607–1616.

(doi:10.1098/rstb.2008.0336)

Baron E., Richirt J., Villoutreix R., Amsellem L., Roux F. & Bennett A. (2015). The genetics of intra-

and interspecific competitive response and effect in a local population of an annual plant

species. *Functional Ecology*, 29, 1361–1370.

Bataillon, T. (2003). Shaking the 'deleterious mutations' dogma? *Trends in Ecology & Evolution*,

18, 315–317.

Bates, D., Mächler, M., Bolker, B. & Walker, S. (2014). Fitting Linear Mixed-Effects Models using

lme4. *arXiv:1406.5823 [stat]*.

Bellard C., Bertelsmeier C., Leadley P., Thuiller W.& Courchamp F. (2012). Impacts of climate

change on the future of biodiversity. *Ecology Letters*, 15, 365–377.

Bolnick, D.I., Amarasekare, P., Araújo, M.S., Bürger, R., Levine, J.M., Novak, M., *et al.* (2011). Why

intraspecific trait variation matters in community ecology. *Trends in Ecology & Evolution*, 26,

183–192.

Bossdorf Ol., Prati D., Auge H. & Schmid B. (2004). Reduced competitive ability in an invasive

plant. *Ecology Letters*, 7, 346–353.

Burnham and Anderson (2002). Model Selection and Multimodel Inference: A Practical

Information-Theoretical Approach. 2d ed. New York: Springer-Verlag

Bruno, J.F., Stachowicz, J.J. & Bertness, M.D. (2003). Inclusion of facilitation into ecological

theory, 7.

Cahill, F. J., Kembel, S. W. and Gustafson, D. J. (2005), Differential genetic influences on

competitive effect and response in *Arabidopsis thaliana*. Journal of Ecology, 93: 958-967.

doi:10.1111/j.1365-2745.2005.01013.x

Cahill, J.F., Elle, E., Smith, G.R. & Shore, B.H. (2008). Disruption of a belowground mutualism

alters interactions between plants and their floral visitors. *Ecology*, 89, 1791–1801.

Camara, M. d. and M. Pigliucci. (1999). Mutational contributions to genetic variance-covariance

matrices: An experimental approach using induced mutations in *Arabidopsis thaliana*.

Evolution 53 (6), 1692-1703

Caradus, J. & Woodfield, D. Plant and Soil (1998) 200: 63.

https://doi.org/10.1023/A:1004296707631

Carr, D. E. and Dudash, M. R. (1996), Inbreeding depression in two species of *Mimulus*

(Scrophulariaceae) with contrasting mating systems. American Journal of Botany, 83: 586-

593. doi:10.1002/j.1537-2197.1996.tb12743.x

Castellanos Maria Clara, Donat-Caerols Santiago, González-Martínez Santiago C., Verdú Miguel &

Gómez-Aparicio Lorena. (2014). Can facilitation influence the spatial genetics of the

beneficiary plant population? *Journal of Ecology*, 102, 1214–1221.

Chapin, F. S. III, Autumn, K., & Pugnaire, F. (1993). Evolution of suites of traits in response to

environmental stress. *American Naturalist*, 142, S78–S92.

Chase, J. M. et al. 2002. The interaction between predation and competition: a review and

synthesis. − Ecol. Lett. 5: 302–315.

Chave, J., Muller-Landau, H.C. & Levin, S.A. (2002). Comparing Classical Community Models:

Theoretical Consequences for Patterns of Diversity. *The American Naturalist*, 159, 1–23.

Chesson, P. (2000). Mechanisms of Maintenance of Species Diversity. *Annual Review of Ecology*

and Systematics, 31, 343–366.

Clark, J.S. (2010). Individuals and the Variation Needed for High Species Diversity in Forest Trees.

Science, 327, 1129–1132.

Colwell, R.K. & Rangel, T.F. (2009). Hutchinson's duality: The once and future niche. *PNAS*, 106,

19651–19658.

Connell, J. (1980). Diversity and the Coevolution of Competitors, or the Ghost of Competition

Past. *Oikos, 35*(2), 131-138. doi:10.2307/3544421

Damgaard, C. (2004). Inference from plant competition experiments: the effect of spatial

covariance. *Oikos*, 107, 225–230.

Dudley, S. A., Murphy, G. P., File, A. L. and Robinson, D. (2013), Kin recognition and competition

in plants. Funct Ecol, 27: 898-906. doi:10.1111/1365-2435.12121

Ehlers, B.K., Damgaard, C.F. & Laroche, F. (2016). Intraspecific genetic variation and species

coexistence in plant communities. *Biology Letters*, 12, 20150853.

Exposito-Alonso M, Becker C, Schuenemann VJ, Reiter E, Setzer C, Slovak R, et al. (2018) The rate

and potential relevance of new mutations in a colonizing plant lineage. PLoS Genet 14(2):

e1007155. https://doi.org/10.1371/journal.pgen.1007155

Falahati-Anbaran, M., Lundemo, S. and Stenøien, H. K. (2014), Seed dispersal in time can

counteract the effect of gene flow between natural populations of *Arabidopsis thaliana*. New

Phytol, 202: 1043-1054. doi:10.1111/nph.12702

Fisher C.K. and Mehta P.(2014). The transition between the niche and neutral regimes in ecology.

Proc Natl Acad Sci USA 111:13111–13116

Fournier-Level A, et al.(2016) Predicting the evolutionary dynamics of seasonal adaptation to

novel climates in Arabidopsis thaliana. Proc Natl Acad Sci USA 113(20):E2812–E2821.

Fridley, J.D. & Grime, J.P. (2010). Community and ecosystem effects of intraspecific genetic

diversity in grassland microcosms of varying species diversity. *Ecology*, 91, 2272–2283.

Fussmann G. F., Loreau M. & Abrams P. A. (2007). Eco-evolutionary dynamics of communities and

ecosystems. *Functional Ecology*, 21, 465–477.

Genung, M. A., Schweitzer, J. A., Úbeda, F. , Fitzpatrick, B. M., Pregitzer, C. C., Felker-Quinn, E.

and Bailey, J. K. (2011), Genetic variation and community change – selection, evolution, and

feedbacks. Functional Ecology, 25: 408-419. doi:10.1111/j.1365-2435.2010.01797.x

Hamilton, W. D. 1964. The genetical evolution of social behaviour. *I & II. J. Theor. Biol.* 7:1–52.

Hart, S. P., Schreiber, S. J., Levine, J. M. and Coulson, T. (2016), How variation between individuals

affects species coexistence. Ecol Lett, 19: 825-838. doi:<u>10.1111/ele.12618</u>

Hart, S. P., Freckleton, R. P., Levine, J.M. & Kroon, H. (2018). How to quantify competitive ability.

Journal of Ecology, 0.

Hausmann N. J., Juenger T. E., Sen S., Stowe K. A., Dawson T. E., et al., 2005. Quantitative trait loci

affecting δC^{13} and response to differential water availability in *Arabidopsis thaliana*. Evolution

59: 81–96

Hofhuis, H. and Hay, A. (2017), Explosive seed dispersal. New Phytol, 216: 339-342.

doi:10.1111/nph.14541

Holt, R.D. (2009). Bringing the Hutchinsonian niche into the 21st century: Ecological and

evolutionary perspectives. *PNAS*, 106, 19659–19665.

Hubbell S.P. (2005). Neutral theory in community ecology and the hypothesis of functional

equivalence. *Functional Ecology*, 19, 166–172.

Hutchinson G.E. (1957). *Concluding remarks. Cold Spring Harbor Symp Quant Biol 22:415–427.*

Johnson, M.T.J. Stinchcombe, J.R. An emerging synthesis between community ecology and

evolutionary biology Trends Ecol. Evol., 22 (2007), pp. 250-257

Johnson, M.T.J., Vellend, M. & Stinchcombe, J.R. (2009). Evolution in plant populations as a driver

of ecological changes in arthropod communities. *Philosophical Transactions of the Royal*

Society of London B 364:1593-1605.

Kerwin et al. (2015) Natural genetic variation in Arabidopsis thaliana defense metabolism genes

modulates field fitness | eLife Lens. https://lens.elifesciences.org/05604/. Last accessed 4

May 2018.

Koch, H., Frickel, J., Valiadi, M. & Becks, L. (2014). Why rapid, adaptive evolution matters for

community dynamics. *Front. Ecol. Evol.*, 2.

Kubisch A. *et al.* (2013). Kin competition as a major driving force for invasions. *Am. Nat.* 181: 700–

706

Lachmuth S., Henrichmann C., Horn J., Pagel J., Schurr F.M. & Rafferty N. (2018). Neighbourhood

effects on plant reproduction: An experimental–analytical framework and its application to

the invasive Senecio inaequidens. *Journal of Ecology*, 106, 761–773.

Lankau, R.A. (2011). Rapid Evolutionary Change and the Coexistence of Species. *Annual Review of*

Ecology, Evolution, and Systematics, 42, 335–354.

Lankau, R.A. & Strauss, S.Y. (2007). Mutual feedbacks maintain both genetic and species diversity

in a plant community. *Science*, 317, 1561–1563.

Lankau R. A., Wheeler E., Bennett A. E. & Strauss S. Y. (2010). Plant–soil feedbacks contribute to

an intransitive competitive network that promotes both genetic and species diversity. *Journal*

of Ecology, 99, 176–185.

Latzel, V., Allan, E., Bortolini Silveira, A., Colot, V., Fischer, M. & Bossdorf, O. (2013). Epigenetic

diversity increases the productivity and stability of plant populations. *Nat Commun*, 4, 2875.

Laure, G. Zimmermann N. E., Levine J. M., Adler P.r B. & Wootton T (2017). The effects of

intransitive competition on coexistence. *Ecology Letters*, 20, 791–800.

Levine, J.M. & HilleRisLambers, J. (2009). The importance of niches for the maintenance of

species diversity. *Nature*, 461, 254–257.

Lion, S. (2018) Theoretical approaches in evolutionary ecology: Environmental feedback as a

unifying perspective. Am. Nat.191, 21–44.

Lynch, M., and B. Walsh. (1998). Genetics and analysis of quantitative traits. Sinauer Associates,

Inc., Sunderland, MA.

Masclaux, F., Hammond, R. L., Meunier, J. , Gouhier-Darimont, C. , Keller, L. and Reymond, P.

(2010), Competitive ability not kinship affects growth of *Arabidopsis thaliana* accessions. New

Phytologist, 185: 322-331. doi:10.1111/j.1469-8137.2009.03057.x

Mukai, T. (1964). The Genetic Structure of Natural Populations of Drosophila Melanogaster. I.

Spontaneous Mutation Rate of Polygenes Controlling Viability. *Genetics*, 50, 1–19.

Nei, M. (2013). Mutation-driven evolution. Oxford University Press

Nottebrock H., Schmid B., Mayer K., Devaux C., Esler K. J., Böhning-Gaese K., *et al.* (2017a). Sugar

landscapes and pollinator-mediated interactions in plant communities. *Ecography*, 40, 1129–

1138.

Nottebrock H., Schmid B,., Treurnicht M., Pagel J., Esler K. J., Böhning-Gaese K., *et al.* (2017b).

Coexistence of plant species in a biodiversity hotspot is stabilized by competition but not by

seed predation. *Oikos*, 126.

Ossowski, S., K. Schneeberger, J. I. Lucas-Lled´o, N.Warthmann, R. M. Clark,R. G. Shaw, D. Weigel,

and M. Lynch. (2010). The rate and molecularspectrum of spontaneous mutations in

Arabidopsis thaliana. Science 327:92–94.

Roles, A. J., and J. K. Conner. (2008). Fitness effects of mutation accumulation in a natural

outbred population of wild radish (*Raphanus raphanistrum*): comparison of field and

greenhouse environments.. Evolution, 62: 1066-1075. doi:10.1111/j.1558-5646.2008.00354.x

Roles, A. J., M. T. Rutter, C. B. Fenster and J. K. Conner. (2016). Field measurement of genotype by

environment interaction for fitness caused by spontaneous mutations in Arabidopsis thaliana.

Evolution, 70: 1039-1050.

R Development Core Team (2011). R: A language and environment for statistical computing.

Vienna, Austria: R Foundation for Statistical Computing.

Rutter, M. T., F. H. Shaw and C. B. Fenster. (2010). Spontaneous mutation parameters for

Arabidopsis thaliana measured in the wild. Evolution, 64: 1825-1835.

Rutter, M., A. Roles, J. Conner, R. Shaw, F. Shaw, K. Schneeberger, S. Ossowski, D. Weigel and **C.**

B. Fenster. (2012). Brief Communication: Fitness of *Arabidopsis thaliana* mutation

accumulation lines whose spontaneous mutations are known. Evolution, 66: 2335-2339.

Rutter, M. T., A. J. Roles, and C. B. Fenster. (2018). Quantifying natural seasonal variation in

mutation parameters with mutation accumulation lines. *Ecology and Evolution,*

https://doi.org/10.1002/ece3.4085

Schaack, S., Allen, D.E., Latta, L.C., Morgan, K.K. & Lynch, M. (2013). The effect of spontaneous

mutations on competitive ability. *Journal of Evolutionary Biology*, 26, 451–456.

Shaw, R.G., Byers, D.L. & Darmo, E. (2000). Spontaneous Mutational Effects on Reproductive

Traits of Arabidopsis thaliana..Genetics 155:369–378.

Shaw, F. H., C. J. Geyer, and R. G. Shaw. (2002). A comprehensive model of mutations affecting

fitness and inferences for Arabidopsis thaliana. Evolution 56:453–463.

Shefferson Richard P., Salguero-Gómez Roberto & Rees Mark. (2015). Eco-evolutionary dynamics

in plants: interactive processes at overlapping time-scale and their implications. *Journal of*

Ecology, 103, 789–797.

Soliveres, S., Smit, C. & Maestre, F.T. (2015). Moving forward on facilitation research: response to

changing environments and effects on the diversity, functioning and evolution of plant

communities. *Biol Rev Camb Philos Soc*, 90, 297–313.

Soliveres S et al. (2015). Intransitive competition is widespread in plant communities and

maintains their species richness. Ecol. Lett. 18, 790–798. doi:10.1111/ele.12456

Stearns, F. W. and C. B. Fenster. (2016). The effect of induced mutations on quantitative traits in

Arabidopsis thaliana: Natural versus artificial conditions. Ecology and Evolution, doi:

10.1002/ece3.2558

Strauss, S.Y. & Irwin, R.E. (2004). Ecological and evolutionary consequences of multispecies

plant-animal interactions. *Annu. Rev. Ecol. Evol. Syst.*, 35, 435–466.

Taylor, D.R., Aarssen, L.W. & Loehle, C. (1990). On the Relationship between r/K Selection and

Environmental Carrying Capacity: A New Habitat Templet for Plant Life History Strategies.

Oikos, 58, 239–250.

Tiffin, P. (2002). Competition and time of damage affect the pattern of slection acting on plant

defense against herbivores. Ecology, 83: 1981-1990. doi:10.1890/0012-

9658(2002)083[1981:CATODA]2.0.CO;2

Tilman D. (2004). Niche tradeoffs, neutrality, and community structure: a stochastic theory of

resource competition, invasion, and community assembly. Proc. Natl Acad. Sci. USA 101, 10

854–10 861. doi:10.1073/Pnas.0403458101

Turcotte, M.M. & Levine, J.M. (2016). Phenotypic Plasticity and Species Coexistence. *Trends Ecol.*

Evol. (Amst.), 31, 803–813.

Uriarte, M., Swenson, N.G., Chazdon, R.L., Comita, L.S., John Kress, W., Erickson, D., *et al.* (2010).

Trait similarity, shared ancestry and the structure of neighbourhood interactions in a

subtropical wet forest: implications for community assembly. *Ecol. Lett.*, 13, 1503–1514.

Van Dam, N. M. and Baldwin, I. T. (1998), Costs of jasmonate-induced responses in plants

competing for limited resources. Ecology Letters, 1: 30-33. doi:10.1046/j.1461-

0248.1998.00010.x

Vellend Mark & Geber Monica A. (2005). Connections between species diversity and genetic

diversity. *Ecology Letters*, 8, 767–781.

Violle, C., Enquist, B.J., McGill, B.J., Jiang, L., Albert, C.H., Hulshof, C., *et al.* (2012). The return of

the variance: intraspecific variability in community ecology. *Trends in Ecology & Evolution*, 27,

244–252.

Violle, C. & Jiang, L. (2009). Towards a trait-based quantification of species niche. *J Plant Ecol*, 2,

87–93.

Violle, C., Thuiller, W., Mouquet, N., Munoz, F., Kraft, N.J.B., Cadotte, M.W., *et al.* (2017).

Functional Rarity: The Ecology of Outliers. *Trends in Ecology & Evolution*, 32, 356–367.

Weng, M.-L., Becker, C., Hildebrandt, J., Rutter,M.T., Shaw, R.G., Weigel, D. and C. B. Fenster.

Fine-Grained Analysis of Spontaneous Mutation Spectrum and Frequency in *Arabidopsis*

thaliana Genetics. Early online (2018); https://doi.org/10.1534/genetics.118.301721

Whitham, T. G., Young, W. P., Martinsen, G. D., Gehring, C. A., Schweitzer, J. A., Shuster, S. M.,

Wimp, G. M., Fischer, D. G., Bailey, J. K., Lindroth, R. L., Woolbright, S. and Kuske, C. R. (2003),

Community and ecosystem genetics: A consequence of extended phenotype. Ecology, 84:

559-573. doi:10.1890/0012-9658(2003)084[0559:CAEGAC]2.0.CO;2

Whitham, T.G., Bailey, J.K., Schweitzer, J.A., Shuster, S.M., Bangert, R.K., LeRoy, C.J. *et al.* (2006).

A framework for community and ecosystem genetics: from genes to ecosystems. *Nat. Rev.*

Genet., 7, 510–523.

Whitlock R. & Lortie C. (2014). Relationships between adaptive and neutral genetic diversity and

ecological structure and functioning: a meta-analysis. *Journal of Ecology*, 102, 857–872.