Next-generation sequencing based hospital outbreak investigation yields insight

into *Klebsiella aerogenes* population structure and determinants of carbapenem

resistance and virulence

Adel Malek[1], Kelly McGlynn[1], Samantha Taffner[1], Lynn Fine[3], Brenda Tesini[4], Jun

Wang[1], Heba Mostafa[1], Sharon Petry[1], Archibald Perkins[1], Paul Graman[4], Dwight

Hardy[1,2], and Nicole Pecora[1,#]

[1]Department of Pathology and Laboratory Medicine, University of Rochester Medical

Center, New York, USA

[2]Department of Microbiology and Immunology, University of Rochester Medical Center,

New York, USA

[3]Infection Prevention Program, University of Rochester Medical Center, New York

[4]Department of Medicine, Infectious Diseases Division, University of Rochester Medical

Center, New York, USA

Running title: Carbapenem-resistant *Klebsiella aerogenes* outbreak

[#]**Correspondence to:**

Nicole Pecora, MD, PhD

University of Rochester Medical Center

601 Elmwood Avenue, Rochester, NY-14642, USA

Phone: (585) 276-4674, Email: nicole_pecora@urmc.rochester.edu

ABSTRACT

Klebsiella aerogenes is a nosocomial pathogen associated with drug resistance and outbreaks in intensive care units. In a 5-month period in 2017, we experienced an increased incidence of cultures for carbapenem-resistant *K. aerogenes* (CR-KA) from an adult cardiothoracic intensive care unit (CICU) involving 15 patients. Phylogenomic analysis following whole-genome sequencing (WGS) identified the outbreak CR-KA isolates to group together as a tight clonal cluster (<7 SNPs apart), suggestive of a protracted intra-ward transmission event. No clonal relationships were identified between the CICU CR-KA strains and additional hospital CR-KA patient isolates from different wards and/or previous years. Genes encoding carbapenemases or drug-resistant plasmids were absent in the outbreak strains, and carbapenem resistance was attributed to mutations impacting AmpD activity and membrane permeability. The CICU outbreak strains harbored an integrative conjugative element (ICE*Kp*10), which has been associated with pathogenicity in hypervirulent *Klebsiella pneumoniae* lineages. Comparative genomics with global *K. aerogenes* genomes showed our outbreak strains to group closely with global ST4 strains, which along with ST93 likely represent dominant *K. aerogenes* lineages associated with human infections. WGS is a powerful tool that goes beyond high-resolution tracking of transmission events into identifying the genetic basis of drug-resistance and virulence, which are not part of conventional diagnostic workflows. With an increasing availability of sequenced genomes from across the globe, population structure analysis offers opportunities to identify emerging trends and dominant clones associated with specific syndromes and geographical locations for poorly characterized pathogens.

INTRODUCTION

Klebsiella aerogenes (formerly described as *Enterobacter aerogenes*) is a ubiquitous member of the *Enterobacteriaceae* family and a significant nosocomial pathogen associated with drug resistance and a wide variety of infections including pneumonia, bacteremia, urinary tract and surgical site infections (1, 2). In vulnerable patients, *K. aerogenes* infections can arise endogenously (gastrointestinal flora) or be acquired from surroundings in the facility where the patient is admitted (horizontal transmission through colonized healthcare workers, contaminated devices/shared equipment, other patients etc.), with the most critical risk factor for acquiring infection being prolonged broad-spectrum antibiotic administration. Additional risk factors for *K. aerogenes* infections include prolonged stay at healthcare facilities especially ICUs and neonatal wards, complex underlying medical conditions, immunosuppression and mechanical ventilation or the presence of foreign devices. Numerous hospital ward outbreaks in both pediatric and adult populations due to *K. aerogenes* have been described due to a common source or spread via patient-to-patient transmission (1, 2). A particularly high frequency of hospital ICU outbreaks was continually reported from Western Europe in a period between the 1990s and early 2000s, that were largely attributed to the spread and endemic establishment of a clonal *K. aerogenes* strain harboring the extended-spectrum B-lactamase TEM-24 (bla_{TEM-24}) (2).

Within the US and other regions across the globe, *K. aerogenes* has also been reported along with *Klebsiella pneumoniae, Enterobacter cloacae* and *Escherichia coli* to be among the frequently isolated carbapenem resistant *Enterobacteriaceae* (3-5). Clinical CR-KA strains harboring plasmid-borne serine carbapenemases have been

described in the US and worldwide, while metallo-β-lactamases and OXA- 48 have

been reported in Europe, Asia and Brazil (2). However, the primary mechanisms

underlying carbapenem non-susceptibility in *K. aerogenes* are thought to be

carbapenemase independent and mediated by mutations affecting regulation of

chromosomal AmpC β-lactamase expression and membrane permeability (2). The

latter has been well documented in *K. aerogenes* with reports describing mutations

impacting porin function/expression that can arise *in vivo* during antibiotherapy (6), be

reversible (2), and present complex diagnostic and therapeutic management challenges

(7).

Despite the role of *K. aerogenes* as an important opportunistic pathogen and its

epidemic potential, the clinical relevance of intra-species genetic diversity and

significance of specific sequence types (STs) remains unknown. In comparison, in

genomically closely related *K. pneumoniae* and to some extent in *E. cloacae*, clonal

complexes and sequence types (STs) associated with geographical distribution, multi-

drug resistance, hospital outbreaks and disease syndromes have been defined (8, 9).

Recently, a multi-locus sequence-typing (MLST) scheme has been developed for *K.*

aerogenes (10) that can help explore the above-mentioned issues, however, its

performance in evaluating and discriminating clinical/environmental isolates has not yet

been reported.

We pursued this study to investigate an outbreak of CR-KA in a cardiothoracic

intensive care unit (CICU) at our hospital, which persisted for 5 months despite

aggressive infection control measures. The primary goals of our study included whole-

genome sequencing (WGS) based investigation of the clonal relationships among the

CR-KA strains isolated from patients in our hospital and defining putative loci associated

with carbapenem resistance and virulence. In addition, the recently developed publicly

available *K. aerogenes* MLST scheme afforded us the opportunity to delineate the

population structure of CR-KA stains isolated from patients at our hospital. Our initial

findings led us to broadly investigate the origin and significance of specific *K. aerogenes*

sequence types identified in our hospital CR-KA strains by performing comparative

genomics using publicly available global *K. aerogenes* genomes.

RESULTS

Epidemiological and genomic characterization of the CICU CR-KA cluster

In July 2017, five CICU patients had CR-KA isolated from respiratory tract

specimens (room occupancy and patient demographics detailed in Fig. 1 and Table S1).

The first identified case (Patient A) had a past medical history significant for intravenous

drug use and recurrent methicillin-resistant *Staphylococcus aureus* infections. Patient

A's CICU course is described in Fig. S1. The temporal association of subsequent

positive cultures in the CICU prompted an outbreak investigation. Between August to

November 2017, despite extensive infection prevention interventions, 10 additional

CICU patients had positive CR-KA cultures (Table S1). Of the 15 positive cultures from

unique patients, 6 were clinical specimens (5 respiratory, 1 blood) and 9 were rectal

surveillance cultures. Interventions included contact precautions for all patients on the

unit, cohorting of staff, and weekly surveillance cultures of rectal swabs and tracheal

aspirates. Environmental surveillance cultures during this period were negative. An

epidemiological investigation into the possible risk factors among the CICU patients

developing CR-KA infections was non-contributory.

Antibiotic susceptibility testing (AST) of the CICU CR-KA strains revealed some

differences in phenotype (Table 1) among the carbapenems and cephalosporins. Sixty

percent of the CICU cluster strains were resistant to three carbapenems (ertapenem,

imipenem and meropenem), while 93% showed phenotypic resistance to ertapenem.

Susceptibility to cefepime also varied, with a subset of outbreak strains displaying

phenotypic resistance (Table 1).

To address the possibility of an outbreak event, particularly in light of variable

antibiotic susceptibilities, a WGS investigation was undertaken to identify phylogenetic

relationships and transmission links. A total of 26 URMC *K. aerogenes* isolates were

sequenced using Illumina WGS. The primary strains investigated in this study were 15

CR-KA isolates from patients admitted in the CICU, between June and Nov 2017 (Table

S1). For context and comparison, an additional set of 9 CR-KA strains

epidemiologically unlinked to the CICU-cluster strains (patient isolates between years

2015-17), were included in the study (Table S2). Two clinical isolates were also

included as controls, URMC 201 (intermediate susceptible to imipenem) and URMC 223

(susceptible to all carbapenems tested). The AST profiles of the non-CICU CR-KA study

strains are described in Table 1.

All of the 26 sequenced URMC *K. aerogenes* genomes showed high coverage

(>88%) relative to the *K. aerogenes* reference KCTC 2190 strain (ATCC 13048^{T}),

Dataset 1. Single nucleotide polymorphisms (SNPs) were identified across the study

genomes relative to the reference sequence (pair-wise SNP differences ranged from 1-

28,170, see Dataset 2). MLST assignment indicated that all of the CICU clinical and

surveillance isolates belonged to ST4, and SNP-based analyses grouped them in a tight

cluster separately and distantly from non-outbreak isolates (Fig. 2, Dataset 2). Within

the CICU cluster, strains differed from URMC 205 (first case, patient A isolate) by less

than 7 SNPs. In addition, these isolates bore an identical plasmid profile (based on

replicon and plasmid typing) (Table S3). In contrast, all of the 2017 non-CICU isolates

were significantly distant from the CICU outbreak isolates (>20,000 SNPs). The most

closely related non-CICU CR-KA strain was URMC 201 (isolated in 2015). This strain

was also ST4, with 433 SNPs in a pairwise comparison to URMC 205. (Fig. 2, Dataset

2). These relationships, coupled with the epidemiological data, indicated that the 2017

CICU CR-KA strains were of clonal origin.

175
Carbapenem resistance in the URMC CR-KA strains was driven by adaptive

chromosomal gene alterations

A. WGS based identification of acquired antibiotic resistance genes

Despite phenotypic carbapenem resistance, the CICU CR-KA did not harbor

genes for carbapenemases, extended-spectrum β-lactamases, or plasmid-borne AmpC

cephalosporinases. A single non-outbreak CR-KA isolate (URMC 203) harbored a

carbapenemase gene (*bla*$_{nmcA}$). No horizontally acquired genes conferring resistance to

non-β-lactam antibiotics were identified among the study strains, consistent with their

susceptibility profiles (Table 1).

B. Non-synonymous sequence alterations in key chromosomal loci

implicated in carbapenem resistance

In the absence of genes encoding carbapenemases and ESBLs in the CR-KA

outbreak strains, variations in other genetic loci associated with carbapenemase-

independent resistance mechanisms were investigated (11-13). Focusing on the AmpC

cephalosporinase and outer membrane porins, sequence variations in *ampD, ampG,*

ampR, omp35, *omp36* and *ompR* genes in the study strains were assessed relative to

the 'wild-type' allele in the carbapenem-susceptible reference type strain KCTC 2190

(14)*.* The sequences were also compared to alleles in URMC 223 (carbapenem

susceptible) and URMC 201 (intermediate susceptibility). For the *omp* genes, the

upstream DNA sequences were also assessed. The *ampG* and *ompR* genes were wild-

type in all study strains. Variants were identified in all other loci and are described

below:

ampD. Mutations were identified in the *ampD* gene for each of the 24 CR-KA

isolates in this study (Table 2), while the control strains URMC 223 and URMC 201 bore

the 'wild-type' *ampD* allele. The outbreak strains harbored single missense SNPs

(either 284G>T or 482G>A) in *ampD* resulting in Trp95Leu or Arg161His substitutions.

Six non-outbreak CR-KA strains harbored independent non-synonymous single

substitutions while 2 missense substitutions were identified in URMC 221. A single non-

outbreak strain, URMC 202, harbored a nonsense mutation resulting in a truncated

AmpD protein (Table 2). DNA sequence corresponding to *ampD* allele was absent in

isolate URMC 203, (the only study strain to possess a carbapenemase-encoding gene

*bla*NMC-A) due to a large deletion in the genomic region harboring *ampD* and the

neighboring *ampE* gene.

The potential impact of Trp95Leu or Arg161His substitutions on AmpD activity in

the outbreak strains was investigated by homology-based structural modeling of *K.*

aerogenes AmpD using a high-resolution crystal structure of *Citrobacter freundii* AmpD

(15), a close homolog (83.33% amino acid sequence identity). AmpD contains a

hydrophobic surface to accommodate its GlcNAc-anh-MurNAc ligand, which is made up

of tripeptide and glycan moieties. The tripeptide portion of GlcNAc-anh-MurNAc is

coordinated through three salt bridges between carboxyl groups on the tripeptide and

residues Arg71, Arg161 and Arg107, while the peptide backbone is oriented across the

hydrophobic surface. Trp95 forms a planar surface at the end of the ligand-binding

channel to position the diaminopimelate moiety at the distal end of the tripeptide.

Based on the *in silico K. aerogenes* AmpD model, the positively charged

guanidinium group of Arg161 forms two strong electrostatic interactions with the

carboxyl group of D-glutamine on the tripeptide portion of the ligand (Fig. 3). Mutation

of this residue to histidine was predicted to weaken ligand binding, likely affecting the

positioning of the ligand in the active site. In the Trp95Leu mutation, the hydrophobicity

in the region is preserved, but the shorter length of the leucine side chain leaves a gap

at the end of the binding channel likely affecting the positioning of the entire ligand (Fig.

3). These observations lend support that the missense mutations observed in the CICU

outbreak strains would alter ligand docking on AmpD likely reducing/inhibiting its

activity. Four out of the seven CR-KA non-outbreak strains had substitutions within

glycan and peptide interacting regions of AmpD (Table 2), suggesting that these

alterations might also impact activity.

ampR. All of the outbreak strains harbored the reference *ampR* allele. Several

non-outbreak CR-KA strains harbored 2-5 substitutions that likely represent variant

alleles as they were also observed in the control carbapenem-susceptible strain URMC

223.   A single non-outbreak CR-KA strain, URMC 210, bore a nonsense mutation in the

ampR gene resulting in a premature stop codon (Trp117X), likely resulting in a non-

functional truncated AmpR protein.

***omp35* and *omp36*.** Among the 15 outbreak CR-KA strains, three different

omp36 variants were identified relative to the wild-type allele (Table 3). An identical

profile of missense SNPs in these strains and the control strain, URMC 201

(intermediate carbapenem susceptibility) were observed relative to the reference

genome allele. Seven outbreak strains had additional mutations that resulted in

severely truncated proteins. Several non-outbreak CR-KA strains also harbored

missense SNPS of unclear significance. Two strains URMC 202 and URMC 204

harbored distinct frame-shift mutations yielding truncated Omp36 protein variants. A 42

base-pair region of high variation (nucleotides 680-724), corresponding to 15/16 amino

acid substitutions in loop L5 was observed in all the clinical isolates relative to the

reference sequence (Table 3, Fig. S2). The hypervariable region results in a different

charge profile in the region and has been previously reported in a *K. aerogenes* study

describing imipenem resistant clinical isolates (harboring ESBL TEM-24) from patients

in France (16). The predicted Omp36 protein in clinical strain URMC 221 and the

carbapenem susceptible control strain (URMC 223) bore 87% identity relative to the

reference genome Omp36 and were considered significantly distant variants (not

included in the comparative analyses).

All but one of our 24 CR-KA strains had the wild-type allele of *omp35*. A single

strain (URMC 202) bore a deletion in the N-terminal encoding region. The DNA

sequence upstream of the *omp35* gene was investigated to identify mutations in the

promoter sites of the strains, of which one (URMC 204) had a nucleotide difference of

unclear significance at the -22 position.

Virulome analysis identifies a large pathogenicity-associated integrative and

conjugative element (ICE), ICE*Kp*10 in the CICU outbreak CR-KA strains

The prolonged nature of the clonal CR-KA outbreak in our CICU led us to search

for putative virulence genes or pathogenicity loci, which could have promoted

persistence and transmission. A cluster of chromosomally encoded genes encoding

yersiniabactin (*ybt*) metallophore and colibactin (*clb*) genotoxin systems was identified

in all of the outbreak strains (Fig. 4). These loci have been implicated in invasive

infections of *K. pneumoniae* (17).

The *ybt* locus harbored putative genes involved in regulation as well as synthesis

of the siderophore, corresponding transport associated proteins, and a receptor protein

for the uptake of metal-bound siderophore. The *clb* locus included putative homologs

encoding enzymes, transferases and transport proteins involved in production and

secretion of the polyketide colibactin (Fig. 4, Dataset 3). Investigation of the genomic

loci associated with the virulence factor gene cluster identified them to be present on a

mobilizable integrative conjugative element (ICE) inserted in a tRNA-Asn site adjacent

to a gene encoding the glycine cleavage system (Fig. 4). The ICE element bore a

modular arrangement of gene clusters encoding mobile elements, P4 like integrase,

type IV secretion system conjugation machinery and mobilization genes (Fig. 4, Dataset

3). The element was identified to be ICE*Kp*10, using a recently described typing and

interpretation scheme of *ybt* and *clb* sequences (17).

Among the URMC non-outbreak *K. aerogenes* strains, these putative virulence

genes were identified in only 4/10 isolates, which were either ST4 or ST93. The control

strain URMC 223 harbored the yersiniabactin locus exclusively. Detailed descriptions of

these elements in the URMC *K. aerogenes* strains are described in Table S4.

Comparative analyses of URMC CR-KA and publicly available *K. aerogenes*

genomes

To gain insights into the emergence and epidemiology of the CICU outbreak

clones and to place our hospital CR-KA strains in the broader context of global *K.*

aerogenes strains, comparative phylogenomic analyses were performed using Harvest

genomics suite (18). Publicly available *K. aerogenes* genome assemblies (n=110) were

included in the analyses. These included 70 clinical and surveillance strains isolated

from human specimens, 3 'environmental' strains, and 36 strains of unknown origin

(Dataset 4). Based on the newly described MLST scheme, ST4 and ST93 strains were

found to be markedly overrepresented in the available genomes (51.8%, 57/110).

Excellent correlation was observed between Harvest generated tree topologies

(Fig. 5) as well as pairwise SNP differences (Dataset 5) as compared to those

generated by the CFSAN SNP pipeline, for the URMC CR-KA study strains. Based on

Harvest analyses, the CICU outbreak strains clustered closely with each other and to

other global ST4 genomes, as compared to the other URMC CR-KA strains (outbreak-

unrelated), which were distantly dispersed throughout the phylogenomic distribution

(Fig. 5). The MLST based sequence types of URMC CR-KA and the global *K.*

aerogenes genomes also correlated tightly with HARVEST generated core-genome

based topologies (Fig. 5).

Six publicly available assembled genomes grouped closely with the CICU

outbreak strain genomes (< 200 SNPs apart). These included ST4 strains, UCI 27, UCI

28, UCI 45, which have been described in a carbapenem resistance surveillance study

by Cerqueria et al. (19), and GN04794, GN05662 and GN02525; strains derived from

varied US patients' clinical specimens (blood, sputum, wound drainage) Fig. 5, Dataset

4). The above-mentioned strains had fewer SNP differences in relation to the CICU

outbreak strains as compared to URMC 201, the closest and only non-outbreak ST4

strain isolated in our hospital (Fig. 5, Dataset 5).

Carbapenemase encoding genes were identified in a total 13/111 global *K.*

aerogenes genome assemblies (11.8%) (Fig. 5). These included genes encoding KPC-

2 (n=7), KPC-3 (n=1), OXA-48 (n=4) and NDM-6 (n=1). The chromosomal serine

carbapenemase, *bla*$_{NMC-A,}$ was solely present in URMC 203, a CR-KA strain isolated

from a patient in our hospital in 2015. Among the thirty ST4 genomes, 4 strains (13%)

harbored genes encoding carbapenemases (4/4, KPC-2), and these were clinical

strains isolated from non-USA patients. The relative contribution of carbapenemase

mediated versus non-carbapenemase mediated mechanisms of resistance to

carbapenems in the global *K. aerogenes* could not be assessed due to the absence of

antibiotic susceptibility metadata for most strains.

A characteristic genomic feature of the URMC CICU outbreak strains and a

subset of non-outbreak associated strains was the presence of the yersiniabactin

siderophore and colibactin systems. Using Kleborate (17), we investigated the

distribution and organization of these systems in the global *K. aerogenes* genomes to

identify associations, if any, with specific ST types and geographical regions (Fig. 5,

Datasets 6). The prevalence of yersiniabactin and colibactin encoding systems in the

global *K. aerogenes* genomes was found to be 53.64% (59/110) and 52.72% (58/111)

respectively. A 100 % association was found between the presence of colibactin in the

genomes and concurrent presence of yersiniabactin. Higher prevalence of the virulence

cluster was observed in ST4 and ST93 types; 85% (12/14) and 95% (42/44)

respectively, although these two ST types were also the most abundantly represented in

the available set of genomes. The other ST types were less well represented in the

study set (n<3), so the prevalence of these systems in them could not be accurately

established. Interestingly, the strains exclusively designated 'environmental' isolates

(B3, FGI35, and B) did not harbor genes encoding the above-mentioned virulence

systems (Fig. 5, Dataset 6). These trends correlated with analyses of our 26 hospital

strains, where 75% of the ST 93 strains (3/4) were positive for yersiniabactin and

colibactin systems, while none of the strains with novel/unassigned ST types harbored

genes encoding the same (0%, 5/5).

DISCUSSION

Whole genome sequencing presents a powerful resource that can be deployed

for prospective and comprehensive outbreak investigations offering tremendous

resolution in tracking transmission events and delineating genomic determinants

associated with drug resistance and virulence (20). This WGS study was initiated in

order to establish the molecular epidemiology of CR-KA strains isolated from patients in

our hospital. Beyond the outbreak investigation, whole genome data enabled us to

address additional fundamental questions associated with carbapenem resistance,

virulence attributes and population structure of *K. aerogenes* that are poorly understood.

Fifteen patients were associated with the CICU outbreak event in the period

between July-November 2017. Based on the differences in AST patterns in a subset of

study strains, it was initially questioned whether the strains were clonal. WGS

investigations demonstrated that the 15 CICU CR-KA isolates differed from each other

by less than 7 SNPs and grouped distantly from the other hospital CR-KA isolates,

which were used a baseline for context and comparison (Fig. 2). Based on these

findings, the CICU CR-KA strains were concluded to be part of a single clonal cluster

indicating protracted intra-ward transmission. Environmental sampling within the CICU

remained negative and a common source of the outbreak or contributory risk factors

could not be identified. After the last positive patient surveillance culture (Nov 2017), no

CR-KA strains were isolated in a 2-month period of continued surveillance and the

outbreak was deemed to have subsided. Phylogenomic analysis (albeit limited) of the

non-CICU CR-KA isolates did not identify specific dominant clones circulating in the

hospital (Fig. 2, Dataset 2).

Infections due to carbapenem-resistant organisms present complex diagnostic

and therapeutic management challenges and a better understanding of how adaptive or

acquired resistance emerges in the healthcare environment is needed (21). In clinical

CR-KA strains, carbapenem resistance has been associated with carbapenemase

production or adaptive mutations following antibiotic exposure (2). Among our 24 study

CR-KA isolates, a single non-CICU strain harbored a carbapenemase-encoding gene,

bla_{NMC-A}. NmcA has been reported in *E. cloacae* (22), but to our knowledge this is the

first report of this carbapenemase being identified in *K. aerogenes*. All but one of the 24

CR-KA strains were found to harbor mutations in genes involved in the synthesis or

regulation of the inducible AmpC cephalosporinase (*ampD, ampR*) and outer membrane

porins (*omp 35, omp36*). A majority of mutations were found to be within the open-

reading frames of *ampD* and *omp36* (catalogued in Tables 2 and 3).

There are limited reports regarding the role of AmpD in adaptive carbapenem

resistance in *K. aerogenes* strains (23, 24), although the role of AmpD in AmpC

expression in *E. cloacae* has been well established (25). *In silico* modeling of *K.*

aerogenes AmpD using the *C. freundii* AmpD crystal structure (15) predicted that

Arg161His and Trp95Leu substitutions due to SNPs in the outbreak strains would likely

impact enzymatic activity (Fig. 3). Other *ampD* mutations in several non-CICU CR-KA

isolates were found to result in substitutions that could affect substrate-enzyme

interactions (Table 2).

Mutations in *omp36* have been described in clinical CR-KA strains and functional

studies investigating their impact have been reported (6, 26). A diverse array of

mutations was identified in the *omp36* gene among the CR-KA study strains, including

non-synonymous mutations resulting in frame-shifts or premature stop codons resulting

in truncated and likely non-functional Omp36 variants (Table 3). Additional SNPs

relative to the carbapenem-susceptible reference genome that resulted in substitutions

of unknown significance in predicted β-sheets and extracellular loop regions (Table 3,

Fig. S2).

Even within the CICU CR-KA clonal cluster, heterogeneity was identified in *ampD*

and *omp36* genes (Table 2, Table 3). These genetic loci likely represent mutational

hotspots associated with adaptive carbapenem resistance in *K. aerogenes.* It is likely

that the CICU CR-KA cluster represents a population of clonal origin, albeit with micro-

heterogeneity in the above regions. The testing and archiving of single isolated

carbapenem resistant colonies instead of multiples during individual patient specimen

workup likely represents a limitation that did not allow us to capture the full complement

of CR-KA strain microdiversity associated with individual patients during the outbreak

event.

The detailed significance of the alleles and mutations described above in

development of carbapenem resistance in *K. aerogenes* needs to be verified by

additional genetic (allelic exchange and complementation) and biochemical approaches.

Moreover, complex regulatory networks including transcriptional activators, sensor

kinases and two-component systems have been implicated in adaptive drug resistance

in *Enterobacteriaceae spp*. (13)*,* and their roles in contributing towards carbapenem

resistance in our study strains cannot be ruled out.

Despite their lack of horizontally acquired drug resistance elements, the outbreak

strains in this study managed to persist for several months in the face of an active

infection prevention effort prompting us to assess their virulence determinants. The

outbreak strains and a subset of non-outbreak CR-KA strains harbored an ICE

encoding the metallophore yersiniabactin (Ybt) and genotoxin colibactin (Clb)

systems (Fig. 4, Table S4). A Ybt encoding pathogenicity island has been described in

association with a prolonged nation-wide outbreak in the Netherlands involving multiple

hospitals and >100 patients due to a multi-drug resistant *Enterobacter hormaechei*

clone (27). A recent study by Lam *et al.* described the prevalence of *ybt* to be

particularly high (>80%) in certain hypervirulent *K. pneumoniae* clonal-groups (CG23)

(17). The study also reported a significant association of Ybt with an increased risk of

invasive infections (bacteremia, liver abscesses etc.). Ybt was first described in

pathogenic *Yersinia spp.*, encoded by a chromosomal gene cluster termed the High

Pathogenicity Island (HPI), with a critical role in iron scavenging during infection (28).

Subsequent studies reported acquisition of the HPI in other clinical *Enterobacteriaceae*

strains (17, 29) and additional functions ascribed to Ybt include evasion of host

lipocalin-2 (30) and sequestration/import of heavy metals (31). The polyketide colibactin

is frequently associated with yersiniabactin and has been shown to induce

chromosomal instability and DNA damage in eukaryotic cells (32). Our findings present

new avenues for research investigating the role of ICE elements encoding Ybt and Clb

in the virulence of clinical *K. aerogenes* strains. While horizontal transmission of

carbapenemases via mobile elements is increasingly recognized as a major public

health issue (33), the transmission of mobilizable virulence factors present an

underappreciated threat in the healthcare environment that warrants more surveillance.

In order to set our hospital strains (outbreak and non-outbreak) into a broader

context, core-genome comparisons and MLST were used to examine the population

structure of our strains relative to global *K. aerogenes* strains pulled from public

databases (Fig. 5, Datasets 5, 6). This analysis is the first evaluation of the nascent *K.*

aerogenes MLST scheme in discriminating clinical *K. aerogenes* isolates. The scheme

was found to be robust with distribution of STs correlating closely with topologies based

on *K. aerogenes* strains core-genomes. Our preliminary analyses suggest that ST4 and

ST93 might be dominant global clones associated with *K. aerogenes* infections.

Outbreak CR-KA strains clustered closest to other clinical US ST4 strains, suggesting a

clonal expansion of this ST (Fig. 5). It is noteworthy that the ST4 group also included

carbapenemase-producing *K. aerogenes* isolates from international sites. These

included CR-KA strains associated with drug-resistant intra-abdominal and urinary tract

infections in patient samples from Brazil, Canada, Colombia and China that had been

sequenced as a part of large surveillance study, SMART (Study for Monitoring

Antimicrobial Resistance Trends) (34). ST93 was the most prevalent sequence type in

the global *K. aerogenes* assembled genomes (43/110, 39%), with a wide geographical

distribution including the US (Fig. 5).   Four of the eleven non-CICU CR-KA strains from

our hospital belonged to this group. Two closely related ST93 isolates, *K. aerogenes*

1509E and G7, have been described as representatives of clonal strains associated

with multiple multi-drug resistant *K. aerogenes* outbreaks in France (35, 36).

Incidentally, global ST4 and ST93 isolates also had a higher prevalence of HPI

encoding *ybt* and *clb* (85% and 95% respectively). The success of these potentially

"high-risk" clones needs to be examined more closely by undertaking large-scale

studies with strains from diverse global sites and patient populations. These studies will

help examine niche adaptation, emergence of antibiotic resistance and evolution of

pathogenicity leading to a better understanding of *K. aerogenes.*

In summary, genomic approaches for surveillance and outbreak investigations

are emerging as critical functions for infection prevention and diagnostic microbiology

laboratories. Apart from evaluating the effectiveness of infection measures and

dimension of transmission events, WGS applied across sets of isolates is a powerful

tool for assessing virulence factors and identifying clinically relevant and emerging

sequence types.

MATERIALS AND METHODS

Setting, study design, *K. aerogenes* strains and metadata.

The University of Rochester Medical Center (URMC) is an 830-bed tertiary-care medical

center, with a 14-bed cardiothoracic intensive care unit, serving the Greater Rochester

Area, New York. Following approval by the University of Rochester Institutional Review

Board (RSRB00068143), a total of 26 *K. aerogenes* strains isolated from patients at

URMC in the course of regular clinical care and/or surveillance efforts were selected for

the study. Each isolate corresponded to a single first CR-KA strain isolated during the

course of hospitalization. For context and comparison, an additional set of *K.*

aerogenes strains epidemiologically unlinked to the CICU outbreak was included in the

study. Ward occupancy and pertinent clinical and epidemiological information were

obtained through review of patient medical records and the laboratory information

system, and are described in Tables S1 and S2.

Antibiotic susceptibility testing (AST) of the study *K. aerogenes* strains

AST of the study strains was performed as part of routine diagnostics using the

VITEK®2 (BioMérieux, France) system and/or Kirby Bauer disk-diffusion methods. AST

interpretations were based on interpretive criteria defined by the Clinical and Laboratory

Standards Institute (M100 document, 27th Edition).

Sequencing library preparation and raw data acquisition

The isolates were cultured on standard laboratory media from archived frozen stocks,

examined for purity, and re-identified by Vitek MALDI-TOF MS (BioMérieux, France).

Single colony genomic DNA extractions were performed using MagNA Pure Compact

instrument (Roche Diagnostics, Indianapolis, IN); the DNA quality was analyzed using

QuantiFluor dsDNA system (Promega). Nextera XT kit (Illumina, San Diego, CA) was

used for preparing dual-indexed WGS libraries. For QC dsDNA quality of individual

samples was analyzed using the 4200 TapeStation system (Agilent). Following library

cleanup, individual libraries were pooled at equimolar ratios and denatured; DNA

concentration was determined using Qubit ssDNA kit (Thermo fisher Scientific). The

libraries were combined with20ρM PhiX control and sequenced on the Illumina Miseq^{TM}

benchtop sequencer (Illumina, San Diego, CA) at the URMC Genomic Core Facilities,

using V3 kit and 2X300 bp paired-end protocol.

Genomic analyses

Analyses were performed using an in-house bioinformatics pipeline 'URMC Bacterial

Genomic Analysis Pipeline' (v2.0.6), run on a high-performance computer cluster at the

Center for Integrated Research Computing at University of Rochester. **A) Quality**

control: For each sample, the read quality scores across all bases were assessed

using FastQC v0.11.5 (37) and low quality reads were trimmed using Trimmomatic

v0.36 (38). Genomic coverage of the sequencing reads relative to the reference *K.*

aerogenes KCTC 2190 genome (ATCC 13048^{T}; Refseq accession

number: NC_015663.1) were determined using Bowtie 2 v2.2.9 (39). Across the study

isolates, alignments showed average genomic coverage of 90.84% (minimum 89%) with

an average depth of coverage: 100X (excluding regions below 12X). Quality metrics

are detailed in Dataset 1. **B) Core-genome SNP calling workflow:** A read mapping

approach was used to assess SNPs in the genomes of the study CR-KA strains relative

to the reference genome *K. aerogenes* KCTC 2190. Mapping, variant calling and

phylogenetic analysis were performed by locally installed CFSAN SNP Pipeline v1.0.0

(40), an analysis workflow developed by the U.S Food and Drug Administration (FDA).

CFSAN pipeline employs a 2-phase variant calling workflow and the 'optimized' version

of the pipeline with criteria as applied by Saltykova et al (higher stringency in allele

frequency thresholds and coverage) (41) was applied. In the first phase, variants were

called based on *mpileup function* of SAMTools and *mpileup2snp tool* from VarScan

(minimum average base quality=20, minimum read depth of coverage at site=12,

minimum allele frequency=90%). Densely clustered SNPs that could arise due to

recombination were excluded (>/=3 SNPs in a 50 bp window). High-confidence SNP

variants meeting the criteria were composed to a list. In the 2[nd] phase, nucleotide sites

at the listed positions were determined for all sites (minimum allele frequency threshold

for SNP filtering=90%). SNPs located in mobile elements as annotated by RAST were

excluded. The final nucleotide sites post filtering corresponding to the listed positions

were used to build a SNP matrix. A multi FASTA file with concatenated SNP matrix

entries were used for inferring phylogeny by FastTree (42), which were visualized using

ITOL (43). **C) Sequence assembly, scaffolding, annotation and analyses:** De novo

genome assemblies (average N_{50}: 344943 across all isolates) were generated by

SPAdes genome Assembler v3.11.1 (44), and the assembly quality was assessed by

QUAST v4.5 (45). Ordering and orientation of contigs was performed using Medusa .

Draft genomes were annotated with RAST (46) and Prokka (47). MLST on the *K.*

aerogenes genomes was performed using a newly developed publicly available scheme

(10). Alleles and genetic markers corresponding to acquired antibiotic resistance,

virulence and plasmid replicons were identified using PlasmidFinder (48), ResFinder

(49), and Virulence Factor Database (50). Typing of genetic loci associated with

mobilizable yersiniabactin siderophore and genotoxin colibactin systems was performed

using Kleborate (17), and the loci were visualized with MacVector software (MacVector,

Cary, NC). Sequence variations in chromosomal genes associated with carbapenem

resistance in the study strains were assessed relative to alleles in control strains

(URMC 201, URMC 223) and 'wild-type' alleles in carbapenem susceptible (14)

reference genome KCTC 2190 [Genbank ID-*ampD*: 10792472 (EAE_11350), *ampG*:

10792757 (EAE_12735), *ampR*: 10792632 (EAE_12115), *omp35*: 10793271

(EAE_15245), *omp36*: 10795060 (EAE_24205) and *ompR*: 10791222 (EAE_05245)].

Multiple sequence alignments of gene alleles and corresponding proteins was

performed using Vector NTI software (Invitrogen, Carlsbad, CA) and the alignments

were manually inspected to identify substitutions that would likely impact function. **D)**

Comparative genomics of publicly available global *K. aerogenes* assembled

genomes with URMC CR-KA assemblies: Using a custom shell script, publicly

available *K. aerogenes* genomes available as of July 2018 in the NCBI genome

database were downloaded from the NCBI FTP site. Harvest genomics suite was used

to perform intraspecific core-genome alignments as described before (18); phylogenies

were visualized using ITOL (43). **E) Accession number(s):** WGS and metadata

corresponding to the study URMC *K. aerogenes* isolates were deposited at NCBI under

BioProject accession number PRJNA504784. Accession numbers for individual isolates

are listed in Table S1.

***In silico* protein analyses**

For homology modeling, SWISS-MODEL (51) was used to thread *K. aerogenes*

AmpD (acc WP_015704411.1, aa1-187) through the structure of *C. freundii* AmpD (15)

(PDB 2y2c, aa1-187). The overall quaternary structure of *K. aerogenes* AmpD was

predicted with high precision (95% confidence, 99% coverage). Comparative analyses

and imaging of protein structures were performed with PyMOL (52). JalView was used

to create alignments (53).

ACKNOWLEDGEMENTS

We are grateful to the URMC Clinical Microbiology Laboratories and Infection

Prevention staff in specimen processing, data collection and epidemiological

investigations. We wish to acknowledge the URMC Genomics Research Center for

support with WGS. We also thank Steve Gill (URMC, Genomics Research Center) for

reviewing the manuscript draft. Internal funding from University of Rochester

Department of Pathology and Laboratory Medicine supported this study. We report no

conflicts of interest relevant to this article.

REFERENCES

1. **Sanders WE, Jr., Sanders CC.** 1997. *Enterobacter spp*.: pathogens poised to flourish at the turn of the century. Clin Microbiol Rev **10:**220-241.

2. **Davin-Regli A, Pages JM.** 2015. *Enterobacter aerogenes* and *Enterobacter cloacae*; versatile bacterial pathogens confronting antibiotic treatment. Front Microbiol **6:**392.

3. **Guh AY, Bulens SN, Mu Y, Jacob JT, Reno J, Scott J, Wilson LE, Vaeth E, Lynfield R, Shaw KM, Vagnone PM, Bamberg WM, Janelle SJ, Dumyati G, Concannon C, Beldavs Z, Cunningham M, Cassidy PM, Phipps EC, Kenslow N, Travis T, Lonsway D, Rasheed JK, Limbago BM, Kallen AJ.** 2015. Epidemiology of Carbapenem-Resistant *Enterobacteriaceae* in 7 US Communities, 2012-2013. JAMA **314:**1479-1487.

4. **Lee HJ, Choi JK, Cho SY, Kim SH, Park SH, Choi SM, Lee DG, Choi JH, Yoo JH.** 2016. Carbapenem-resistant *Enterobacteriaceae:* Prevalence and Risk Factors in a Single Community-Based Hospital in Korea. Infect Chemother **48:**166-173.

5. **Robert J, Pantel A, Merens A, Lavigne JP, Nicolas-Chanoine MH, Group ONsCRS.** 2014. Incidence rates of carbapenemase-producing *Enterobacteriaceae* clinical isolates in France: a prospective nationwide study in 2011-12. J Antimicrob Chemother **69:**2706-2712.

6. **Philippe N, Maigre L, Santini S, Pinet E, Claverie JM, Davin-Regli AV, Pages JM, Masi M.** 2015. In Vivo Evolution of Bacterial Resistance in Two Cases of

Enterobacter aerogenes Infections during Treatment with Imipenem. PLOS One

10:e0138828.

7. **Boucher HW, Talbot GH, Bradley JS, Edwards JE, Gilbert D, Rice LB,**

Scheld M, Spellberg B, Bartlett J. 2009. Bad bugs, no drugs: no ESKAPE! An

update from the Infectious Diseases Society of America. Clin Infect Dis **48:**1-12.

8. **Wyres KL, Holt KE.** 2016. *Klebsiella pneumoniae* Population Genomics and

Antimicrobial-Resistant Clones. Trends Microbiol **24:**944-956.

9. **Gomez-Simmonds A, Annavajhala MK, Wang Z, Macesic N, Hu Y, Giddins**

MJ, O'Malley A, Toussaint NC, Whittier S, Torres VJ, Uhlemann AC. 2018.

Genomic and Geographic Context for the Evolution of High-Risk Carbapenem-

Resistant *Enterobacter cloacae* Complex Clones ST171 and ST78. MBio **9**.

10. https://pubmlst.org/kaerogenes/

11. **Jacoby GA.** 2009. AmpC beta-lactamases. Clin Microbiol Rev **22:**161-182.

12. **Dupont H, Choinier P, Roche D, Adiba S, Sookdeb M, Branger C, Denamur**

E, Mammeri H. 2017. Structural Alteration of OmpR as a Source of Ertapenem

Resistance in a CTX-M-15-Producing *Escherichia coli* O25B:H4 Sequence Type

131 Clinical Isolate. Antimicrob Agents Chemother **61**.

13. **Pages JM, James CE, Winterhalter M.** 2008. The porin and the permeating

antibiotic: a selective diffusion barrier in Gram-negative bacteria. Nat Rev

Microbiol **6:**893-903.

14. **Lavigne JP, Sotto A, Nicolas-Chanoine MH, Bouziges N, Bourg G, Davin-**

Regli A, Pages JM. 2012. Membrane permeability, a pivotal function involved in

antibiotic resistance and virulence in *Enterobacter aerogenes* clinical isolates.

Clin Microbiol Infect **18:**539-545.

15. **Carrasco-Lopez C, Rojas-Altuve A, Zhang W, Hesek D, Lee M, Barbe S,**

Andre I, Ferrer P, Silva-Martin N, Castro GR, Martinez-Ripoll M, Mobashery

S, Hermoso JA. 2011. Crystal structures of bacterial peptidoglycan amidase

AmpD and an unprecedented activation mechanism. J Biol Chem **286:**31714-

31722.

16. **Thiolas A, Bornet C, Davin-Regli A, Pages JM, Bollet C.** 2004. Resistance to

imipenem, cefepime, and cefpirome associated with mutation in Omp36

osmoporin of *Enterobacter aerogenes*. Biochem Biophys Res Commun **317:**851-

856.

17. **Lam MMC, Wick RR, Wyres KL, Gorrie CL, Judd LM, Jenney AWJ, Brisse S,**

Holt KE. 2018. Genetic diversity, mobilisation and spread of the yersiniabactin-

encoding mobile element ICE*Kp* in *Klebsiella pneumoniae* populations. Microb

Genom.

18. **Treangen TJ, Ondov BD, Koren S, Phillippy AM.** 2014. The Harvest suite for

rapid core-genome alignment and visualization of thousands of intraspecific

microbial genomes. Genome Biol **15:**524.

19. **Cerqueira GC, Earl AM, Ernst CM, Grad YH, Dekker JP, Feldgarden M,**

Chapman SB, Reis-Cunha JL, Shea TP, Young S, Zeng Q, Delaney ML, Kim

D, Peterson EM, O'Brien TF, Ferraro MJ, Hooper DC, Huang SS, Kirby JE,

Onderdonk AB, Birren BW, Hung DT, Cosimi LA, Wortman JR, Murphy CI,

Hanage WP. 2017. Multi-institute analysis of carbapenem resistance reveals

remarkable diversity, unexplained mechanisms, and limited clonal outbreaks.

Proc Natl Acad Sci U S A **114:**1135-1140.

20. **Quainoo S, Coolen JPM, van Hijum S, Huynen MA, Melchers WJG, van**

Schaik W, Wertheim HFL. 2017. Whole-Genome Sequencing of Bacterial

Pathogens: the Future of Nosocomial Outbreak Analysis. Clin Microbiol Rev

30:1015-1063.

21. **Goodman KE, Simner PJ, Tamma PD, Milstone AM.** 2016. Infection control

implications of heterogeneous resistance mechanisms in carbapenem-resistant

Enterobacteriaceae (CRE). Expert Rev Anti Infect Ther **14:**95-108.

22. **Pottumarthy S, Moland ES, Juretschko S, Swanzy SR, Thomson KS,**

Fritsche TR. 2003. NmcA carbapenem-hydrolyzing enzyme in *Enterobacter*

cloacae in North America. Emerging infectious diseases **9:**999-1002.

23. **Tzouvelekis LS, Tzelepi E, Kaufmann ME, Mentis AF.** 1994. Consecutive

mutations leading to the emergence in vivo of imipenem resistance in a clinical

strain of *Enterobacter aerogenes*. J Med Microbiol **40:**403-407.

24. **Babouee Flury B, Ellington MJ, Hopkins KL, Turton JF, Doumith M,**

Woodford N. 2016. The differential importance of mutations within AmpD in

cephalosporin resistance of *Enterobacter aerogenes* and *Enterobacter cloacae*.

Int J Antimicrob Agents **48:**555-558.

25. **Babouee Flury B, Ellington MJ, Hopkins KL, Turton JF, Doumith M, Loy R,**

Staves P, Hinic V, Frei R, Woodford N. 2016. Association of Novel

Nonsynonymous Single Nucleotide Polymorphisms in *ampD* with Cephalosporin

Resistance and Phylogenetic Variations in *ampC*, *ampR*, *ompF*, and *ompC* in

Enterobacter cloacae Isolates That Are Highly Resistant to Carbapenems.

Antimicrob Agents Chemother **60:**2383-2390.

26.   **De Gheldre Y, Maes N, Rost F, De Ryck R, Clevenbergh P, Vincent JL,**

Struelens MJ. 1997. Molecular epidemiology of an outbreak of multidrug-

resistant *Enterobacter aerogenes* infections and in vivo emergence of imipenem

resistance. J Clin Microbiol **35:**152-160.

27.   **Paauw A, Caspers MP, Leverstein-van Hall MA, Schuren FH, Montijn RC,**

Verhoef J, Fluit AC. 2009. Identification of resistance and virulence factors in an

epidemic *Enterobacter hormaechei* outbreak strain. Microbiology **155:**1478-1488.

28.   **Heesemann J, Hantke K, Vocke T, Saken E, Rakin A, Stojiljkovic I, Berner R.**

1993. Virulence of *Yersinia enterocolitica* is closely associated with siderophore

production, expression of an iron-repressible outer membrane polypeptide of

65,000 Da and pesticin sensitivity. Molecular Microbiology **8:**397-408.

29.   **Putze J, Hennequin C, Nougayrede JP, Zhang W, Homburg S, Karch H,**

Bringer MA, Fayolle C, Carniel E, Rabsch W, Oelschlaeger TA, Oswald E,

Forestier C, Hacker J, Dobrindt U. 2009. Genetic structure and distribution of

the colibactin genomic island among members of the family *Enterobacteriaceae*.

Infect Immun **77:**4696-4703.

30.   **Bachman MA, Lenio S, Schmidt L, Oyler JE, Weiser JN.** 2012. Interaction of

lipocalin 2, transferrin, and siderophores determines the replicative niche of

Klebsiella pneumoniae during pneumonia. MBio **3**.

31.   **Robinson AE, Lowe JE, Koh EI, Henderson JP.** 2018. Uropathogenic

enterobacteria use the yersiniabactin metallophore system to acquire nickel. J

Biol Chem.

32.   **Fais T, Delmas J, Barnich N, Bonnet R, Dalmasso G.** 2018. Colibactin: More

Than a New Bacterial Toxin. Toxins (Basel) **10**.

33.   **Gupta N, Limbago BM, Patel JB, Kallen AJ.** 2011. Carbapenem-resistant

Enterobacteriaceae: epidemiology and prevention. Clin Infect Dis **53:**60-67.

34.   **Morrissey I, Hackel M, Badal R, Bouchillon S, Hawser S, Biedenbach D.**

2013. A Review of Ten Years of the Study for Monitoring Antimicrobial

Resistance Trends (SMART) from 2002 to 2011. Pharmaceuticals (Basel)

6:1335-1346.

35.   **Diene SM, Merhej V, Henry M, El Filali A, Roux V, Robert C, Azza S, Gavory**

F, Barbe V, La Scola B, Raoult D, Rolain JM. 2013. The rhizome of the

multidrug-resistant *Enterobacter aerogenes* genome reveals how new "killer

bugs" are created because of a sympatric lifestyle. Mol Biol Evol **30:**369-383.

36.   **Thiolas A, Bollet C, La Scola B, Raoult D, Pages JM.** 2005. Successive

emergence of *Enterobacter aerogenes* strains resistant to imipenem and colistin

in a patient. Antimicrob Agents Chemother **49:**1354-1358.

37.   http://www.bioinformatics.babraham.ac.uk/projects/fastqc/ .

38.   **Bolger AM, Lohse M, Usadel B.** 2014. Trimmomatic: a flexible trimmer for

Illumina sequence data. Bioinformatics **30:**2114-2120.

39.   **Langmead B, Salzberg SL.** 2012. Fast gapped-read alignment with Bowtie 2.

Nat Methods **9:**357-359.

40.   **Davis S, Pettengill JB, Luo Y, Payne J, Shpuntoff A, Rand H, Strain E.** 2015.

CFSAN SNP Pipeline: an automated method for constructing SNP matrices from

next-generation sequence data. PeerJ Computer Science **1:**e20.

41.   **Saltykova A, Wuyts V, Mattheus W, Bertrand S, Roosens NHC, Marchal K,**

De Keersmaecker SCJ. 2018. Comparison of SNP-based subtyping workflows

for bacterial isolates using WGS data, applied to *Salmonella enterica* serotype

Typhimurium and serotype 1,4,[5],12:i. PLOS One **13:**e0192504.

42.   **Price MN, Dehal PS, Arkin AP.** 2010. FastTree 2-approximately maximum-

likelihood trees for large alignments. PLOS One **5:**e9490.

43.   **Letunic I, Bork P.** 2016. Interactive tree of life (iTOL) v3: an online tool for the

display and annotation of phylogenetic and other trees. Nucleic Acids Res

44:W242-245.

44.   **Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS,**

Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV,

Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new

genome assembly algorithm and its applications to single-cell sequencing. J

Comput Biol **19:**455-477.

45.   **Gurevich A, Saveliev V, Vyahhi N, Tesler G.** 2013. QUAST: quality

assessment tool for genome assemblies. Bioinformatics **29:**1072-1075.

46.   **Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K,**

Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL,

Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD,

Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The

RAST Server: rapid annotations using subsystems technology. BMC genomics

9:75.

47.   **Seemann T.** 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics

30:2068-2069.

48.   **Carattoli A, Zankari E, Garcia-Fernandez A, Voldby Larsen M, Lund O, Villa**

L, Moller Aarestrup F, Hasman H. 2014. In silico detection and typing of

plasmids using PlasmidFinder and plasmid multilocus sequence typing.

Antimicrob Agents Chemother **58:**3895-3903.

49.   **Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O,**

Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial

resistance genes. J Antimicrob Chemother **67:**2640-2644.

50.   **Chen L, Zheng D, Liu B, Yang J, Jin Q.** 2016. VFDB 2016: hierarchical and

refined dataset for big data analysis--10 years on. Nucleic Acids Res **44:**D694-

697.

51.   **Waterhouse A, Bertoni M, Bienert S, Studer G, Tauriello G, Gumienny R,**

Heer FT, de Beer TAP, Rempfer C, Bordoli L, Lepore R, Schwede T. 2018.

SWISS-MODEL: homology modelling of protein structures and complexes.

Nucleic Acids Res **46:**W296-W303.

52.   **Seeliger D, de Groot BL.** 2010. Ligand docking and binding site analysis with

PyMOL and Autodock/Vina. J Comput Aided Mol Des **24:**417-422.

53.   **Waterhouse AM, Procter JB, Martin DM, Clamp M, Barton GJ.** 2009. Jalview

Version 2--a multiple sequence alignment editor and analysis workbench.

Bioinformatics **25:**1189-1191.

FIGURE LEGENDS

Fig. 1. Patient occupancy and overlap in CICU ward during the *K. aerogenes* outbreak. Black bars-first positive CR-KA clinical cultures, blue bars-first positive CR-KA surveillance cultures

Fig. 2. Dendogram showing pairwise SNP differences based phylogenetic relatedness of URMC *K. aerogenes* strains. Whole genome sequence of *K. aerogenes* KCTC 2190 (ATCC 13048) was used for reference mapping. A total of 28,170 discriminatory high-quality SNPs in the core genomes obtained by the CFSAN SNP pipeline were used to plot the tree (excluding mobile elements and putative recombination sites). CICU outbreak strains: IDs highlighted in red, clinical isolates in bold, patient A isolate (1[st] case) with yellow background. Year strain isolated: orange-2015, yellow-2016, blue-2017. SNP differences relative to patient A shown. Scale bar indicates nucleotide substitutions per site.

Fig. 3. Computational modeling of the impact of *ampD* mutations on AmpD in URMC outbreak CR-KA strains. A) *K. aerogenes* AmpD modeled on *C. freundii* AmpD structure. Key residues interacting with glycan and peptide portions of ligand are shown. B, C) Surface models of *K. aerogenes* AmpD depicting the wild-type binding surface for the diaminopimelate moiety (W95, top) versus the binding surface of AmpD containing the W95L mutation (blue surfaces indicate amino acid positions from panel A; bottom, altered surface highlighted in red).

Fig. 4. Yersiniabactin and colibactin encoding gene loci on integrative conjugative element ICE*Kp*10 in the CICU outbreak CR-KA strains. Blue arrows-integrase encoding genes, brown arrows-Zn^{2+}/Mn^{2+} modules grey arrows-genes encoding mobilization proteins, pink arrows: vir-T4SS system.

Fig. 5. Harvest based phylogenomic comparisons of URMC *K. aerogenes* genomes with global *K. aerogenes* genomes. Discriminatory SNPs based on core genome comparisons were used to plot the tree. URMC study *K. aerogenes* strains in purple (outbreak strains in bold, index patient strain highlighted yellow). Presence of genes encoding yersiniabactin siderophore system (green squares), colibactin synthesis cluster (blue triangles), and carbapenamases (red circles) in assembled genomes shown. Scale bar indicates nucleotide substitutions per site.

Table 1. Phenotypic antibiotic susceptibility profiles of *K. aerogenes* strains in this study

	Study Isolates	Amikacin	Gentamicin	Tobramycin	Ciprofloxacin	Moxifloxacin	Trimethoprim-Sulfa	Piperacillin-Tazobactam	Ceftriaxone	Cefepime	Ertapenem	Imipenem	Meropenem
2017 URMC CICU associated CR-KA strains	**URMC 205***	S	S	S	S	S	S	R	R	S	R	R	R
	URMC 206	S	S	S	S	S	S	R	R	S	R	R	R
	URMC 207	S	S	S	S	S	S	R	R	S	R	R	R
	URMC 208	S	S	S	S	S	S	R	R	S	S	S	R
	URMC 209	S	S	S	S	S	S	R	R	S	R	R	R
	URMC 211	S	S	S	S	S	S	R	R	D	R	R	R
	URMC 212	S	S	S	S	S	S	R	R	R	R	I	R
	URMC 213	S	S	S	S	S	S	R	R	R	R	I	R
	URMC 215	S	S	S	S	S	S	R	R	S	R	R	R
	URMC 216	S	S	S	S	S	S	R	R	S	R	S	S
	URMC 218	S	S	S	S	S	S	R	R	S	R	R	R
	URMC 219	S	S	S	S	S	S	R	R	S	R	R	R
	URMC 224	nd	S	S	S	S	S	R	R	R	R	nd	R
	URMC 225	S	S	S	S	S	S	R	R	R	R	R	R
	URMC 226	S	S	S	S	S	S	R	R	R	R	I	I
Other URMC CR-KA strains	URMC 200	S	S	S	S	nd	S	R	R	R	R	nd	R
	URMC 202	S	S	S	S	S	S	R	R	S	R	R	R
	URMC 203	S	S	S	S	S	S	R	R	S	R	R	R
	URMC 204	S	S	S	S	nd	S	R	R	S	R	nd	R
	URMC 210	S	S	S	S	nd	S	R	R	S	R	nd	S
	URMC 214	S	S	S	S	nd	S	I	R	S	R	I	S
	URMC 217	S	S	S	R	nd	S	R	R	S	R	I	S
	URMC 221	S	S	S	S	nd	S	S	S	S	R	nd	S
	URMC 222	S	S	S	S	S	S	R	R	R	R	I	S
Con	URMC 201	S	S	S	S	S	S	S	S	S	S	I	S
	URMC 223	S	S	S	S	nd	S	S	S	S	S	nd	S

R=Resistant, S= Sensitive, I =Intermediate, D= Dose Dependent, nd=not determined; *Patient A (1[st] case)-URMC 205; Con: control strains: URMC 223, carbapenem susceptible and URMC 201, intermediate carbapenem resistance

Table 2. Non-synonymous SNPs in *ampD* gene associated with carbapenem resistance in the URMC *K. aerogenes* isolates

Study Isolates		Differences in *ampD* gene relative to wild-type allele		Effect on encoded AmpD sequence	
		MS SNPs	NS SNPs	Single AA substitutions	PS
CICU outbreak associated CR-KA strains	URMC 205*, URMC 207, URMC 209, URMC 211, URMC 212, URMC 213, URMC 215, URMC 218, URMC 219, URMC 224, URMC 224, URMC 226	482G>A		Arg161His	
	URMC 206, URMC 208, URMC 216	284G>T		Trp95Leu	
Outbreak unrelated CR-KA strains	URMC 200	338T>G		Ile113Ser	
	URMC 202		412C>T		Gln138X
	URMC 204	117C>T		Pro39Ser	
	URMC 210	335C>T		Ser112Leu	
	URMC 214	280G>A		Ala94Thr	
	URMC 217	496G>C		Gly166Arg	
	URMC 221	478A>G, 501C>A		Ile160Val, Ala168Asp	
	URMC 222	492T>G		Glu164Asp	
Control strains	URMC 201, URMC 223				

No sequencing reads mapped to the *ampD* gene in URMC 203.

 *Index patient isolate.

Abbr: SNP: single nucleotide polymorphism, MS: missense, NS: nonsense, AA: amino acids, PS: premature stop codon

Table 3. Non-synonymous genetic lesions in *omp36* genes and resulting alterations in Omp36 sequence in the URMC *K. aerogenes* isolates

Study Isolates		Differences in *omp36* gene sequence relative to wild-type allele				Effect on encoded Omp36 amino acid sequence relative to wild-type sequence				
		FS	HV region (Nt 680-724)	NS SNPs	MS SNPs	FS	HV region (AA 226-241)	PS	Truncated	Single AA substitions in protein synthesized
CICU outbreak associated CR-KA strains	URMC 205*, URMC 207	+	+		175A>G, 564T>C, 566A>G	pAsp91ThrfsX12		102X	+	Ile59Val
	URMC 206, URMC 208, URMC 212, URMC 213, URMC 216, URMC 224, URMC 225, URMC 226		+		175A>G, 564T>C, 566A>G		+			Ile59Val, Asp189Gly
	URMC 209, URMC 211, URMC 215, URMC 218, URMC 219		+	184A>T	175A>G, 564T>C, 566A>G			62X	+	Ile59Val
Outbreak unrelated CR-KA strains	URMC 200, URMC 203, URMC 210, URMC 214		+		564T>C, 566A>G, 615 T>G, 834T>C, 835A>G		+			Asp189Gly, Asp205Glu, Asn279Asp
	URMC 202	+	+		564T>C, 566A>G, 615 T>G, 834T>C, 835A>G	pTrp77LysfsX2		78X	+	Trp77Lys
	URMC 204	+	+		564T>C, 566A>G, 615 T>G, 834T>C, 835A>G	pSer304ProfsX21	+	324X	+	Asp189Gly, Asp205Glu, Asn279Asp
	URMC 217		+		564T>C, 566A>G, 569T>A		+			Asp189Gly, Phe190Tyr
	URMC 222		+		175A>G,564T>C, 566A>G, 615 T>G		+			Ile59Val, Asp189Gly, Asp205Glu
Control strain	URMC 201 (intermediate carbapenem R)		+		175A>G,564T>C, 566A>G		+			Ile59Val, Asp189Gly

*Index patient isolate.

Abbr: SNP: single nucleotide polymorphism, NS: nonsense, MS; missense, FS: frame-shift, PS:

premature stop codon, HV: hypervariable region, Nt: nucleotides, AA: amino acids

Fig. 1. Patient occupancy and overlap in CICU ward during the *K. aerogenes* outbreak. Black bars-first positive CR-KA clinical cultures, blue bars-first positive CR-KA surveillance cultures

Fig. 2. Dendogram showing pairwise SNP differences based phylogenetic relatedness of URMC *K. aerogenes* strains. Whole genome sequence of *K. aerogenes* KCTC 2190 (ATCC 13048) was used for reference mapping. A total of 28,170 discriminatory high-quality SNPs in the core genomes obtained by the adapted CFSAN pipeline were used to plot the tree (excluding mobile elements and putative recombination sites). CICU outbreak strains: IDs highlighted in red, clinical isolates in bold, patient A isolate (1st case) with yellow background. Year strain isolated: orange-2015, yellow-2016, blue-2017. SNP differences relative to patient A shown. Scale bar indicates nucleotide substitutions per site.

Fig. 3. Computational modeling of the impact of *ampD* mutations on AmpD in URMC outbreak CR-KA strains. A) *K. aerogenes* AmpD modeled on *C. freundii* AmpD structure. Key residues interacting with glycan and peptide portions of ligand are shown. B, C) Surface models of *K. aerogenes* AmpD depicting the wild-type binding surface for the diaminopimelate moiety (W95, top) versus the binding surface of AmpD containing the W95L mutation (blue surfaces indicate amino acid positions from panel A; bottom, altered surface highlighted in red).

Fig. 4. Yersiniabactin and colibactin encoding gene loci on integrative conjugative element ICE*Kp*10 in the CICU outbreak CR-KA strains. Blue arrows-integrase encoding genes, brown arrows-Zn^{2+}/Mn^{2+}modules grey arrows-genes encoding mobilization proteins, pink arrows: vir-T4SS system.

Fig. 5. Harvest based phylogenomic comparisons of URMC *K. aerogenes* genomes with global *K. aerogenes* genomes. Discriminatory SNPs based on core genome comparisons were used to plot the tree. URMC study *K. aerogenes* strains in purple (outbreak strains in bold, index patient strain highlighted yellow). Presence of genes encoding yersiniabactin siderophore system (green squares), colibactin synthesis cluster (blue triangles), and carbapenamases (red circles) in assembled genomes shown. Scale bar indicates nucleotide substitutions per site.

Supplementary Data

Fig. S1. Patient A timeline following CICU admit highlighting diagnostic workup, microbiological findings and antibiotic administration prior to development of CR-KA respiratory infection.

Fig. S2. Comparison of *K. aerogenes* Omp36 with representative porins of the OmpC and OmpF families. Predicted β-sheets, loops and helixes are shown. Non-synonymous SNP mutations in CICU outbreak CR-KA strains resulting in substitutions relative to reference highlighted in blue.

Table S1: Attributes of patients associated with the 2017 URMC CICU CR-KA cluster

Patient ID	Isolate ID	Underlying condition/disease and indications for admission	Age range	Culture site and Type
A	URMC 205*	Cardiogenic shock, endocarditis, sepsis, respiratory failure, history of ECMO	20-30	Tracheal aspirate
B	URMC 206	Cardiac tamponade, cardiogenic shock, history of ECMO, biventricular failure, LVAD present	70-80	Sputum
C	URMC 207	Ischemic mitral valve dysfunction, rapid atrial fibrillation, COPD, acute respiratory failure, mitral stenosis	70-80	Sputum
D	URMC 208	Heart failure, LVAD present, hemorrhagic stroke and recurrent infections	60-70	Sputum
E	URMC 209	Congestive heart failure, elevated mycocardial infraction, sarcoidosis	50-60	Sputum
F	URMC 211	History of fibromyalgia, cardiogenic shock, history of ECMO	40-50	Sputum
G	URMC 212	Cardiac tamponade, critical aaortic stenosis	60-70	Rectal Surveillance
H	URMC 213	Chronic systolic heart failure, ischemic cardiomyopathy	60-70	Rectal Surveillance
I	URMC 215	Ascending aortic aneurysm repair	60-70	Rectal Surveillance
J	URMC 216	Coronary artery disease, aortic valve insufficiency	70-80	Rectal Surveillance
K	URMC 218	Congestive heart failure, history of ECMO, acute heart transplant rejection	70-80	Rectal Surveillance
L	URMC 219	Paroxysmal atrial firbrillation, ventricular dysfunction	60-70	Rectal Surveillance
M	URMC 224	Endocarditis, end-stage renal disease	40-50	Blood
N	URMC 225	Chronix systolic heart failure, LVAD complication, cardiac tamponade	50-60	Rectal Surveillance
O	URMC 226	Coronary artery disease, end stage renal disease	70-80	Rectal Surveillance

*URMC 205, first patient diagnosed with clinical CR-KA infection in the URMC CICU in 2017

Abbr: ECMO: Extracorporeal membrane oxygenation, LVAD: left ventricular assist device present, COPD: chronic obstructive pulmonary disease

Table S2: Summary of additional hospital *K. aerogenes* strains in this study
(isolates not associated with CICU CR-KA cluster)

Isolate ID	Year of positive culture	Gender	Age range	Culture site and type	Patient's unit/ward
URMC 200	2015	M	20-30	Catheter urine	General Medicine
URMC 201	2015	M	30-40	Blood	Acute Care Medicine
URMC 202	2016	M	60-70	Broncholaveolar lavage	Burn/Trauma
URMC 203	2016	M	70-80	Biliary drain fluid	Solid Organ Transplant
URMC 204	2016	M	70-80	Catheter urine	Cardiovascular Progressive Care
URMC 210	2017	F	80-90	Urine	Outpatient
URMC 214	2017	F	90-100	Urine	Outpatient
URMC 217	2017	F	70-80	Urine, cytoscopy	Acute Medicine
URMC 221	2017	F	90-100	Urine	Outpatient
URMC 222	2017	F	40-50	Abdominal ascites fluid	Emergency Department
URMC 223*	2017	F	70-80	Urine	Cardiovascular Step Down

* All strains except URMC 223 and URMC 201 were deemed carbapenem resistant by CDC and CLSI definitions

URMC 223 was a pan-sensitive *K. aerogenes* isolated from the urine of a patient in the CICU in Oct 2017 (within the outbreak period)

Table S3: Acquired antibiotic resistance markers, plasmid replicon types and plasmid multilocus sequence typing of the study *K. aerogenes* strains

Study Isolates		Antimicrobial resistance marker	Replicon types	pMLST summary
2017 URMC CICU associated CR-KA strains	URMC 205		IncFIB(pENTAS01),IncFII(Y),ColRNAI	IncF[Y2*:A-:B-]
	URMC 206		IncFIB(pENTAS01),IncFII(Y),ColRNAI	IncF[Y2*:A-:B-]
	URMC 207		IncFIB(pENTAS01),IncFII(Y),ColRNAI	IncF[Y2*:A-:B-]
	URMC 208		IncFIB(pENTAS01),IncFII(Y),ColRNAI	IncF[Y2*:A-:B-]
	URMC 209		IncFIB(pENTAS01),IncFII(Y),ColRNAI	IncF[Y2*:A-:B-]
	URMC 211		IncFIB(pENTAS01),IncFII(Y),ColRNAI	IncF[Y2*:A-:B-]
	URMC 212		IncFIB(pENTAS01),IncFII(Y),ColRNAI	IncF[Y2*:A-:B-]
	URMC 213		IncFIB(pENTAS01),IncFII(Y),ColRNAI	IncF[Y2*:A-:B-]
	URMC 215		IncFIB(pENTAS01),IncFII(Y),ColRNAI	IncF[Y2*:A-:B-]
	URMC 216		IncFIB(pENTAS01),IncFII(Y),ColRNAI	IncF[Y2*:A-:B-]
	URMC 218		IncFIB(pENTAS01),IncFII(Y),ColRNAI	IncF[Y2*:A-:B-]
	URMC 219		IncFIB(pENTAS01),IncFII(Y),ColRNAI	IncF[Y2*:A-:B-]
	URMC 224		IncFIB(pENTAS01),IncFII(Y),ColRNAI	IncF[Y2*:A-:B-]
	URMC 225		IncFIB(pENTAS01),IncFII(Y),ColRNAI	IncF[Y2*:A-:B-]
	URMC 226		IncFIB(pENTAS01),IncFII(Y),ColRNAI	IncF[Y2*:A-:B-]
Other URMC KA strains	URMC 200		ColRNAI	NA
	URMC 201		No replicons found	NA
	URMC 202		ColRNAI	NA
	URMC 203	bla_{NMC-A}	IncFIB(pENTAS01),ColRNAI	IncF(unknown ST)
	URMC 204		No replicons found	NA
	URMC 210		IncFIB(pKPHS1),IncFIB(K),IncFII(Yp)	IncF[S3*:A-:B-]
	URMC 214		ColRNAI	NA
	URMC 217		IncFIB(pENTAS01),ColRNAI	IncF(unknown ST)
	URMC 221		No replicons found	NA
	URMC 222		IncFII(SARC14),IncFII(29),IncFIB(K)	IncF[S3*:A-:B-]
	URMC 223		IncFIB(K),IncHI1B	IncHI1(unknown ST); IncF(unknown ST)

Table S4: Typing yersiniabactin and colibactin encoding gene clusters in URMC *K. aerogenes* strains using Kleborate

Study Isolates		Yersiniabactin	YbST	Colibactin	CbST
2017 URMC CICU associated CR-KA strains	URMC 205	ybt 17; ICEKp10	289-1LV	clb 3	17-3LV
	URMC 206	ybt 17; ICEKp10	289-1LV	clb 3	17-3LV
	URMC 207	ybt 17; ICEKp10	289-1LV	clb 3	17-3LV
	URMC 208	ybt 17; ICEKp10	289-1LV	clb 3	17-3LV
	URMC 209	ybt 17; ICEKp10	289-1LV	clb 3	17-3LV
	URMC 211	ybt 17; ICEKp10	289-1LV	clb 3	17-3LV
	URMC 212	ybt 17; ICEKp10	289-1LV	clb 3	17-3LV
	URMC 213	ybt 17; ICEKp10	289-1LV	clb 3	17-3LV
	URMC 215	ybt 17; ICEKp10	289-1LV	clb 3	17-3LV
	URMC 216	ybt 17; ICEKp10	289-1LV	clb 3	17-3LV
	URMC 218	ybt 17; ICEKp10	289-1LV	clb 3	17-3LV
	URMC 219	ybt 17; ICEKp10	289-1LV	clb 3	17-3LV
	URMC 224	ybt 17; ICEKp10	289-1LV	clb 3	17-3LV
	URMC 225	ybt 17; ICEKp10	289-1LV	clb 3	17-3LV
	URMC 226	ybt 17; ICEKp10	289-1LV	clb 3	17-3LV
Other URMC KA strains	URMC 200	ybt 0; ICEKp10	220-2LV	clb 3	13-2LV
	URMC 201	ybt 17; ICEKp10	289-1LV	clb 3	17-1LV
	URMC 202	ybt 17; ICEKp10	267	clb 3	19-1LV
	URMC 203	-	0	-	0
	URMC 204	ybt 16; ICEKp12	277-3LV	clb 3	17-1LV
	URMC 210	-	0	-	0
	URMC 214	-	0	-	0
	URMC 217	-	0	-	0
	URMC 221	-	0	-	0
	URMC 222	-	0	-	0
	URMC 223	ybt 4; plasmid	221-3LV	-	0

Abbr: YbST: Yersiniabactin locus sequence type, CbST: Colibactin locus sequence type