1

Caterpillars on a phytochemical landscape

Forister, M.L.[1,2], Yoon, S.[1,2], Philbin, C.S.[2,3], Dodson, C.D.[2,3], Hart, B.[4], Harrison, J.G.[5], Shelef, O.[6],

Fordyce, J.A.[7], Marion, Z.H.[8], Nice, C. C.[9], Richards, L.A.[1,2], Buerkle, C.A.[5], Lebeis, S.[10], and Z.

Gompert[11]

[1] Department of Biology, Program in Ecology, Evolution and Conservation Biology, University of

  Nevada, Reno, NV 89557, USA; Email: forister@gmail.com

[2] Hitchcock Center for Chemical Ecology, University of Nevada, Reno, NV 89557, USA

[3] Department of Chemistry, University of Nevada, Reno, NV 89557, USA

[4] Department of Biochemistry, University of Nevada, Reno, NV 89557, USA

[5] Department of Botany and Program in Ecology, University of Wyoming, Laramie, WY 82071, USA

[6] Department of Natural Resources, Institute of Plant Sciences, Volcani Center, Agricultural Research

Organization, Rishon LeZion, Israel

[7] Department of Ecology and Evolutionary Biology, University of Tennessee, Knoxville, TN 37996, USA

[8] Department of Ecology, University of Canterbury, Christchurch, New Zealand

[9] Department of Biology, Population and Conservation Biology Program, Texas State University, San

  Marcos, TX 78666, USA

[10] Department of Microbiology, University of Tennessee, Knoxville, TN 37996, USA

[11] Department of Biology, Utah State University, Logan, UT 84322, USA

Abstract

Foundational theories in plant-animal interactions are being updated with input from modern metabolomic approaches that offer more comprehensive phytochemical profiles than were previously available. Here we use a recently-formed plant-insect interaction, the colonization of alfalfa (*Medicago sativa*) by the Melissa blue butterfly (*Lycaeides melissa*), to describe the landscape of primary and secondary plant metabolites and the performance of caterpillars as affected by both individual compounds and suites of covarying phytochemicals. We find that survival, development time and adult weight are all affected by a large number of compounds, including biomolecules associated with plant cell function and putative anti-herbivore action. The dimensionality of the plant-insect interface is high, with clusters of covarying compounds in many cases encompassing divergent effects on different aspects of caterpillar performance. The sapogenic glycosides are represented by more than 20 individual compounds with some of the strongest beneficial and detrimental effects on caterpillars, which highlights the value of metabolomic data as opposed to previous approaches that relied on total concentrations within defensive classes. Considering positive and negative effects of both secondary compounds and primary metabolites (possibly associated with nutritional imbalance), theories of the evolution of plant defense based on a simple dichotomy between investment in defense or primary metabolism appear to be overly simplistic. Results are also discussed in light of previous work on local adaptation to alfalfa by the focal herbivore.

Keywords

Lycaeides melissa, Medicago sativa, metabolomics, plant defense, specialization

1. Introduction

One of the conceptual pillars of trophic ecology is the idea that herbivores must overcome the barrier of plant secondary chemistry before extracting the nutrients necessary for growth and reproduction [1]. The success of this idea is reflected in areas of research that include coevolution [2], ecological specialization [3], and nutrient flow in ecosystems [4]. In most cases, progress has been made by chemical ecologists focusing on tiny subsets of the secondary metabolites produced by plants and consumed by herbivores. The focus on a few charismatic molecules or classes of compounds, such as furanocoumarins [5] or cardiac glycosides [6], was at least in part necessitated by early methods in natural products chemistry that were targeted and not easily optimized for the discovery of large suites of co-occurring primary and secondary metabolites [7,8]. As technological limitations have dissipated, the opportunity now exists for a deeper understanding of the challenges faced by herbivores, with the possibility of discovering, among other things, novel compounds and synergistic interactions among compounds [9–11]. More generally, an important task is to quantify the phytochemical dimensionality of the antagonistic interaction between plants and herbivores, with an eye towards understanding constraints on the evolution of both players [12,13] and predicting the formation of new plant-herbivore interactions [14]. Here we use the example of a specialized herbivore and a novel host plant to investigate the phytochemical landscape from the perspective of developing caterpillars.

The Melissa blue butterfly, *Lycaeides melissa*, specializes on larval host plants in the pea family (Fabaceae), primarily in the genera *Astragalus* and *Lupinus*. Within the last 200 years, *L. melissa* has colonized introduced alfalfa, *Medicago sativa* (Fabaceae), at least twice and probably multiple times [15], forming a heterogeneous patchwork of association throughout the range of the butterfly in western North America. *M. sativa* is a suboptimal host, relative to native hosts that have been examined, and populations of the butterfly that persist on *M. sativa* show evidence of loss of preference for native hosts [16], reduced caterpillar performance on native hosts, and a slight increase in ability to develop on the suboptimal novel host [17]. The genetic architecture of host use in this system is known to be polygenic and characterized by loci with conditionally neutral (host-specific) effects [17], but what is needed next is

an understanding of how many independent or covarying phytochemical compounds have consequential effects on caterpillars eating the novel host. For example, will the trajectory of further local adaptation by *L. melissa* to *M. sativa* be a matter of evolving the ability to detoxify one or a large number of compounds? We would also like to know how key compounds covary among individual plants, which should help us understand the puzzle faced by ovipositing females facing an array of co-occurring *M. sativa* chemotypes. Here we use a common garden approach (to minimize non-genetic phenotypic variation among plants) and caterpillars individually reared in a controlled environment to address these questions while describing the effects of metabolomic variation in *M. sativa* on *L. melissa*.

2. Methods

(a) Plants and caterpillars

Plants used in this project were grown at the University of Nevada, Reno, Main Station experimental farm. The common garden was planted in 2016 with seeds collected the previous year from 45 plants (previously studied by Harrison et al. [18]) growing in a fallow field in north-western Nevada on the western edge of the Great Basin Desert. The focal butterfly, *L. melissa*, was present in the source field but has not yet colonized the university farm where experimental plants were grown. The 45 maternal plants each contributed 15 offspring to a randomized grid design in the common garden, irrigated with broadcast sprayers in 2016 and drip in 2017, without supplemental fertilization. Out of each maternal family, a single plant was randomly selected for use in the rearing experiment reported here as a way to capture as much genetic and phenotypic variation as possible.

On 17 and 18 July 2017, a total of 45 *L. melissa* females were collected from an alfalfa-associated population near Verdi, NV, and confined to oviposition arenas (500 mL plastic cups) in groups of three with host plant leaves and supplied with Gatorade on mesh lids. After three days, eggs were removed from leaves, pooled, and kept at room temperature until hatching, at which time caterpillars were placed individually in Petri dishes (100 x 25 mm) with leaves of a particular *M. sativa* individual (which became the only plant from which they were fed throughout the experiment). Ten caterpillars were assigned to

each of the 45 experimental *M. sativa* plants (for a total of 450 independently-reared caterpillars) and kept

in a growth chamber set to 25 C and a 12 hour light / 12 hour dark cycle. From each caterpillar we

recorded survival to adult, date of eclosion (if successful) and adult weight to the nearest 0.01 mg on a

Mettler Toledo XP26 microbalance.

(b) Phytochemistry and plant traits

Metabolomic variation among individual plants was characterized with liquid chromatography–mass

spectrometry (LC-MS, [19]) using leaves collected on a single day at the start of the rearing experiment.

Dried, ground leaves (10 mg) were extracted in 2 mL of 70% aqueous ethanol and injected into an

Agilent 1200 analytical high performance liquid chromatograph paired with an Agilent 6230 Time-of-

Flight mass spectrometer via an electrospray ionization source. Resulting chromatograms were analyzed

using MassHunter Quantitative Analysis (v.B.06.00, Agilent, Santa Clara, CA); see electronic

supplementary material for additional phytochemical protocols. Major classes of compounds were

identified using characteristic relative mass defects [20]. Leaf protein content was quantified with three

replicates (~2 mg each) per plant using the Bicinchoninic acid assay (Pierce Biotechnology, Waltham,

MA). Before grinding, five dried leaflets from each sample were weighed to the nearest 0.1 mg, scanned,

and area was measured using ImageJ (v.1.52a); specific leaf area (SLA) was calculated as leaf area

divided by dry mass. Finally, leaf toughness was measured on fresh material in the common garden, at

the start of the experiment (mid-July, when leaves were also sampled for chemistry and protein) and at the

end of the experiment (mid-August), from three leaves per plant at each date, with a penetrometer

(Chatillon 516 Series) through the center of the middle leaflet (as in [18]); the three leaves were selected

haphazardly, avoiding the oldest and youngest leaves. The six leaf toughness measurements per plant

were averaged for a single toughness measure used in analyses.

(c) Analyses of plant traits and caterpillar performance

(i) Overview

Our analytical strategy to understand the association between phytochemical variation and caterpillar

performance followed two complementary paths, one focusing on dimension reduction and feature

selection to produce relatively simple models, and the other on the estimation of effects of all individual

compounds. For the first path, involving dimension reduction and feature selection, we utilized an

approach developed for gene transcription studies that identifies groups or modules of correlated variables

with hierarchical clustering [21]; after clustering, we reduced the number of independent variables by

selecting among modules and other plant traits (specific leaf area, protein and leaf toughness) using lasso

regression [22]. Selected modules (and other plant traits) were analyzed in Bayesian linear models that

are useful in this context because they allowed us to quantify our confidence in the sign of effects

(positive or negative) as continuous probabilities (as opposed to relying on arbitrary significance cutoffs).

For the second analytical path, we utilized ridge regression [22] to estimate effects for all compounds

simultaneously, which allowed us to investigate the distribution of effects among compounds and classes

of compounds. Both analytical paths incorporated cross-validation during the lasso and ridge regressions,

and as a means of evaluating the predictive success of the Bayesian models. We also used randomization

tests to compare the performance of modules and individual compounds with randomly-chosen suites of

compounds.

(ii) Dimension reduction and feature selection

We chose an approach that reduces the number of independent variables while allowing us to learn

something about the correlational structure of the data, specifically unsupervised hierarchical clustering as

implemented in the blockwiseModules function of the WGCNA package [21] in R [23]. Among the

options in the pipeline, we used positive correlations among variables ("signed" network type), merge cut

height at 0.25, and correlations raised to the power of five (which is where the scale free topology index

reached a plateau). Through experimentation, we found that our results with LC-MS data were robust to

variation in these choices, including the choice of signed or unsigned networks. After an initial round of

clustering, we took a remaining 19 unassigned compounds and put them through a second round of

clustering (although the majority of consequential compounds were identified in the first round). One

output of the WGCNA procedure is the first eigenvector from each cluster of compounds, which reduced

our number of predictor variables by a factor of ten. The resulting eigenvectors plus protein, SLA

(specific leaf area) and leaf toughness were then put through the feature reduction step of lasso regression

[22], a penalized regression that allows beta coefficients to be constrained to zero (thus excluding

variables). We used the cv.glmnet function of the glmnet package [24] with cross-validation during error

reduction set to leave out one plant (and associated caterpillars) at each iteration. The variables selected

by the lasso were then put into a Bayesian linear model to estimate coefficients and associated credible

intervals using JAGS (version 3.2.0) run in R with the rjags package [25]. Two Markov chains were run

for 10,000 steps for each analysis (no burn in was required) and chain performance was assessed by

plotting chain histories, and calculating the Gelman and Rubin convergence diagnostic and effective

sample sizes [26,27]. For all models, uninformative priors for the regression coefficients were modeled

as a Normal distribution with a mean of zero and variance of 0.01. We quantified our confidence in the

sign of coefficients (positive or negative) as the fraction of the posterior samples that were less than zero

(for coefficients with a median negative value) or greater than zero (for coefficients with a median

positive value).

167   All analyses were done using the R statistical language [23] on scaled (z-transformed) predictor

variables, and both lasso and Bayesian models used binomial (for survival), Poisson (for development

time) and Gaussian (for adult weight) errors. The latter two analyses (development time and adult

weight) included sex as a factor. The analysis of development time also included adult weight as a

covariate; while (reciprocally) the analysis of adult weight included development time as a predictor.

These variables are negatively correlated (at -0.52), and they function as useful covariates of each other,

allowing us to investigate the possibility of unique plant effects on weight gain and development time,

which could not be discovered if, for example, these variables were combined into a single performance

index.

The success of models developed with the dimension reduction and feature selection pipeline was judged in two ways. We used a cross-validation procedure in which we left out one plant (and associated caterpillars) in each iteration of the Bayesian model and then used the estimated coefficients (for phytochemical variables and other plant traits) to predict the performance of the unobserved caterpillars. After 45 iterations (one for each plant), we calculated a simple correlation coefficient between the observed and predicted performance of caterpillars across plants. In addition, we repeatedly resampled the original LC-MS data to match the structure of the reduced set of predictor variables to ask to what extent randomly assembled modules could outperform the empirically-derived modules.

(iii) Individual compound effects

The second path of our two-part analytical strategy involved simultaneous estimation of the effects of all individual chemical compounds on caterpillar survival, development time and adult weight. For this approach, we again used penalized regression (in the glmnet package [24]), but this time with ridge regression (instead of lasso) which constrains beta coefficients to avoid variance inflation but does not eliminate variables. As with the analyses above, ridge regression was done using error structures appropriate to the specific response variables, and included additional covariates where possible (in models of development time and adult weight). The resulting coefficients associated with all individual compounds were examined as a second perspective on the modules examined in the first set of analyses, and were used to ask to what extent individual compound effects could be predicted by the degree to which they vary among individual plants as quantified with the simple coefficient of variation. To assess confidence in the results of ridge regressions, we used a bootstrap approach, repeatedly resampling the data and estimating coefficients 1000 times, noting the compounds whose bootstrap confidence intervals did or did not overlap zero [28]. We also allowed for the discovery of interactions among compounds using penalized regression on all individual compounds and all pairwise interactions between compounds. For ease of interpretation, this final analysis of potential interactions used lasso (not ridge) regression, letting the coefficients for many of the individual compounds and pairwise interactions go to zero.

3. Results

Of the 450 caterpillars that started the experiment, 261 were reared to eclosion as adults (a mortality rate similar to previous work with this system [17]) on leaves from 45 individual alfalfa plants that were characterized for protein, leaf toughness, specific leaf area and 163 individual metabolomic features (see figure 1 for variation among plants in caterpillar performance and a subset of plant traits, and electronic supplementary material table S1 for a list of compounds). Hierarchical clustering identified 14 subsets (or modules) of compounds with generally low correlations among modules and high correlations within modules (see electronic supplementary material figures S1 and S2 for correlations within and among modules, and figure S3 for module variation among plants). The correlational structure of the phytochemical data is illustrated as an adjacency network in figure 2, where it can be seen that some modules (e.g., modules 1, 2, and 3) contain a great diversity of compound types, while other modules are made up of more narrow classes (e.g. modules 7 and 8 which are mostly saponins, a class of defensive secondary metabolites [29]). From the 14 eigenvectors summarizing variation in the modules, as well as the other plant traits, lasso regression produced a reduced set of potential predictors which were then used in Bayesian multiple regression models that included between seven and nine independent variables (table 1). The models had reasonably high performance in leave-one-out cross-validation (correlations between observed and predicted values were between 0.50 and 0.59, table 1), and also in resampling analyses (electronic supplementary material figure S4), where a small fraction (never more than 4%) of randomly-generated models exceeded the variance explained of the models reported in table 1.

Variation among plants in the suites of covarying compounds had large effects on caterpillar performance: for example, the beta coefficient of -2.33 (on the log-odds scale) associated with module 3 corresponds to a reduction in mean survival from 0.58 to 0.12 associated with a one unit change in variation associated with that phytochemical module (table 1). The phytochemical predictor variables are eigenvectors from clustering analysis, and thus are not entirely straightforward to interpret, especially when the clustering analysis was itself based on z-transformed data. It is useful to note that our LC-MS

data consists of peak areas divided by the peak of an internal standard, and again divided by the dry

weight of the sample (thus, in total, referred to as "relative abundance per dry weight"). Variation in

these numbers reflects variation in concentrations within compounds (among plants), but care should be

used in comparing among compounds because of different ionization responses relative to the standard

(thus the use of z-transformation for among-compound analyses). Nevertheless, the effects reported in

table 1 reflect real variation in suites of compounds, as can be seen in correlations between eigenvectors

and individual compounds in electronic supplementary material figure S2, and in variation among plants

in average z-scores in figure S3.

Modules included in multiple regression models frequently had common effects across response

variables (e.g., the positive association of module 10 with both survival and adult weight), with the

exceptions of module 11 that had a solitary effect on survival and module 6 with an effect only on

development time (although the probability of the latter having a negative effect was only 0.75). Specific

leaf area had a negative effect on survival and adult weight, and the coefficients for specific leaf area (-

0.31 for survival and -0.42 for weight) were of smaller magnitude than most phytochemical effects.

Module-based analyses (as in table 1) focused on feature reduction with lasso regression; as a

complementary analytical approach, we used ridge regression on all of the compounds (which estimates

effects of individual compounds without excluding variables as in lasso regression). Analyses of

individual compounds by ridge regression (figure 3) were broadly consistent with the strongest module-

specific effects, as can be seen, for example, with module 10 having positive effects on survival and adult

weight in module analyses (table 1) and in compound-specific analyses (figure 3). Similarly, the

individual compounds in module 3 had negative compound-specific effects on survival (figure 3), and

that module had the strongest negative effect on survival in the eigenvector-based analyses in table 1. Not

surprisingly, the larger modules (with a greater number of covarying compounds, including many primary

metabolites) tended to have a more complex mix of positive and negative effects (for examples, modules

1 and 2, figure 3). For ease of interpretation, the coefficients from compound-specific regressions of

survival and development time (in figures 3 and 4) have been back-transformed to be on the scales of

probability and days (respectively), and displayed as changes relative to intercepts. For example, a

compound with a relatively large effect on survival in figure 3 could be associated with a one half percent

(0.005) reduction in the probability of survival relative to average survival and while holding other

compounds constant.

We also considered potential pairwise interactions among individual compounds, and found few

interactions that passed the filter of the penalized regression (electronic supplementary material table S2),

at least relative to the large number of potential interactions. We did not find evidence that more or less

variable compounds had differential effects on caterpillars, although there was a trend towards both

greater positive and greater negative effects being associated with less variable compounds (figure S5).

We did, however, see variation among classes of compounds in their effects on caterpillars (figure 4). All

classes included positive and negative effects, although phenolic effects on survival were more often

negative. The widest breadth of effects (including the most extreme positive and negative compound-

specific coefficients) tended to be found among peptides, saponins and phospholipids, with lipids and

terpenoids having a more narrow range (figure 4).

4. Discussion

The results reported here represent a dissection of the phytochemical landscape facing a specialized insect

herbivore attacking a novel host plant [30]. The phytochemical landscape is both physical, referring to

variation in compounds among individual plants in a common garden, and hypothetical to the extent that

effects of individual compounds on caterpillars are estimated, although compounds are of course not

encountered in isolation. Our exploration of the phytochemical landscape facing *L. melissa* on *M. sativa*

is necessarily a first draft based on a snapshot in time. Nevertheless, models including suites of covarying

compounds and other plant traits had predictive success and suggested different natural products affecting

survival, development time and adult weight (performance measured as adult weight is a proxy for

fecundity [31]). Previous work with *M. sativa* and insect herbivores has focused on sapogenic glycosides

[29], and a simple outcome from our study could have been that one or a small number of saponins have

anti-herbivore properties that reduce fitness of our focal insect. Instead, we find large numbers of

compounds with potentially consequential effects on caterpillars (figure 3).

The precise identification of specific compounds is not as important as the more general result

that the prominent class of defensive secondary chemistry, saponins, includes compounds with both

positive and negative effects. Moreover, saponin effects tend slightly towards the positive with improved

survival, faster development and increased adult size (figure 4). Positive effects are potentially associated

with feeding stimulation, as has been observed (along with other positive effects) for other specialist

herbivores and plant toxins [32,33]. Negative effects of saponins on insects potentially include disruption

of hormone production [34], although exact modes of action on *L. melissa* will await further study.

Perhaps more intriguing than the range of effects associated with saponins are the negative effects

associated with variation in certain primary metabolites (figure 4). For example, some of the largest

negative effects in module 3 (figure 2) are phospholipids, peptides and even sucrose. These could be

direct effects if a compound is suboptimal for development, or they could be associated with nutritional

imbalance [35], such that too much of one nutrient makes it difficult for caterpillars to consume a

balanced diet. It has been suggested that the presentation of unbalanced nutrition can be a kind of anti-

herbivore strategy for a plant that does not depend on secondary metabolites [36]. Although this

possibility has not been thoroughly investigated in many systems with full metabolomic profiling, the

idea that nutritional imbalance could be as important as direct toxicity suggests that we might update

theories of the evolution of plant defense that were built on differential investment into simple categories

of plant growth versus defense [37].

The finding that our specialist herbivore is affected by a wide range of metabolites, primary and

secondary, that vary greatly even within a single host population has implications for our understanding

of heterogeneity in the system and for the course of local adaptation of the herbivore to the novel host.

Lycaeides melissa typically colonizes weedy or feral patches of *M. sativa* on roadsides or integrated into

natural communities, and previous work has documented dramatic variation among individual alfalfa

locations (often in close proximity) in the extent to which they can support caterpillar development [38].

Previous phytochemical data with a lower resolution was less successful in explaining that variation [38],

and the results reported here suggest that among patch variation could be explained by future studies

using metabolomic data as used here. The within-population complexity described in the current study

combined with previous evidence for dramatic among-population variation in *M. sativa* suitability for the

focal herbivore also raises the possibility that the novel host presents a multi-faceted and potentially ever-

shifting target from the perspective of evolving butterfly populations. In particular, it is possible that *M.*

sativa defense against a specialist herbivore might be realized through different combinations (within and

among populations) of individually-acting compounds, thus making it unlikely for butterflies in any one

population to possess an effective suite of alleles that improve fitness on the novel host. In this context, it

is interesting to note that a molecular genetic dissection of caterpillar performance in this system found a

large number (potentially hundreds) of individual loci associated with performance on *M. sativa*, yet

evolution in populations associated with the novel host is primarily associated with a loss (through

genetic drift) of the ability to eat a native host and only slight improvement in the ability to eat the novel

host [17].

The correlational structure of the phytochemical variation that we observed has implications for

the evolution of plant defense and the accumulation of insect herbivores on *M. sativa*. Specifically,

correlations among modules should make it possible to hypothesize directions of least resistance for

defense evolution. Compounds in module 9 had a negative effect on survival (table 1), and module 9

negatively covaried with module 10 (electronic supplementary material figure S1), which itself had a

positive association with caterpillar survival. Thus an increase in module 9 and an associated decrease in

10 would be beneficial for the plant, at least with respect to herbivory by our focal herbivore. Of course,

most plants do not have the luxury of optimizing defense against a single herbivore, and it is easy to

imagine that improvements in defense against one enemy could lead to increased attraction to another,

especially given the diversity of effects even within major classes studied here, including saponins and

phenols. Compounds in the latter class (phenolics) were found to have strong positive and negative

effects on assemblages of arthropods associated with the maternal plants from which seeds were collected

to start the common garden used in the present study [18]. We have not attempted to separate constitutive and induced defenses [39] as the plants in our common garden were exposed to natural and continuous levels of herbivory. We have also focused on simple effects rather than interactions among compounds, although some were detected (electronic supplementary material table S2), and interactions could certainly add to the complexity of effects on different herbivores. Future studies in this system involving greater numbers of plants will have greater power to test for robust interactive effects. In the meantime, it is clear that metabolomic data such as that analyzed here has the potential to both open up new avenues of conceptual development in plant-insect interactions and to link micro-evolutionary trajectories across hosts and herbivores.

Acknowledgements

This work was supported by National Science Foundation grant DEB-1638793 to MLF and CDD, DEB-1638768 to ZG, DEB-1638773 to CCN, DEB-1638922 to SL and JAF, and DEB-1638602 to CAB; MLF was additionally supported by a Trevor James McMinn professorship. Thanks to Ian Wallace, the Hitchcock Center for Chemical Ecology and the P.I.G. group for discussion and expertise.

References

1. Feeny P, Rosenthal GA, Berenbaum MR. 1992 The evolution of chemical ecology: contributions from the study of herbivorous insects. *Herbiv. their Interact. with Second. plant Metab.* **2**, 1–44.

2. Agrawal AA, Petschenka G, Bingham RA, Weber MG, Rasmann S. 2012 Toxic cardenolides: chemical ecology and coevolution of specialized plant--herbivore interactions. *New Phytol.* **194**, 28–45.

3. Dyer LA. 1995 Tasty generalists and nasty specialists - antipredator mechanisms in tropical lepidopteran larvae. *Ecology* **76**, 1483–1496.

4. Olson JS. 1963 Energy storage and the balance of producers and decomposers in ecological systems. *Ecology* **44**, 322–331.

5.   Berenbaum M. 1983 Coumarins and caterpillars: a case for coevolution. *Evolution (N. Y)*. **37**, 163–179.

6.   Zalucki MP, Brower LP, Alonso A. 2001 Detrimental effects of latex and cardiac glycosides on survival and growth of first-instar monarch butterfly larvae *Danaus plexippus* feeding on the sandhill milkweed Asclepias humistrata. *Ecol. Entomol.* **26**, 212–224.

7.   Dyer LA *et al.* 2018 Modern approaches to study plant–insect interactions in chemical ecology. *Nat. Rev. Chem.* **1**. (doi:10.1038/s41570-018-0009-7)

8.   Maag D, Erb M, Glauser G. 2015 Metabolomics in plant-herbivore interactions: challenges and applications. *Entomol. Exp. Appl.* **157**, 18–29.

9.   Richards LA, Dyer LA, Smilanich AM, Dodson CD. 2010 Synergistic effects of amides from two Piper species on generalist and specialist herbivores. *J. Chem. Ecol.* **36**, 1105–1113.

10. Sardans J, Penuelas J, Rivas-Ubach A. 2011 Ecological metabolomics: overview of current developments and future challenges. *Chemoecology* **21**, 191–225.

11. Prince EK, Pohnert G. 2010 Searching for signals in the noise: metabolomics in chemical ecology. *Anal. Bioanal. Chem.* **396**, 193–197.

12. Fordyce JA, Nice CC. 2008 Antagonistic, stage-specific selection on defensive chemical sequestration in a toxic butterfly. *Evolution (N. Y)*. **62**, 1610–1617. (doi:10.1111/j.1558-5646.2008.00388.x)

13. Macel M, van Dam NM, Keurentjes JJB. 2010 Metabolomics: the chemistry between ecology and genetics. *Mol. Ecol. Resour.* **10**, 583–593.

14. Erbilgin N, Ma C, Whitehouse C, Shan B, Najar A, Evenden M. 2014 Chemical similarity between historical and novel host plants promotes range and host expansion of the mountain pine beetle in a native host ecosystem. *New Phytol.* **201**, 940–950.

15. Chaturvedi S, Lucas LK, Nice CC, Fordyce JA, Forister ML, Gompert Z. 2018 The predictability of genomic changes underlying a recent host shift in Melissa blue butterflies. *Mol. Ecol.*

16. Forister ML, Scholl CF, Jahner JP, Wilson JS, Fordyce JA, Gompert Z, Narala D, Buerkle CA,

Nice CC. 2012 Specificity, rank preference and the colonization of a non-native host plant by the

Melissa blue butterfly. *Oecologia* **DOI: 10.10**.

17. Gompert Z *et al.* 2015 The evolution of novel host use is unlikely to be constrained by trade-offs

or a lack of genetic variation. *Mol. Ecol.* **24**, 2777–2793.

18. Harrison JG *et al.* 2018 Deconstruction of a plant-arthropod community reveals influential plant

traits with nonlinear effects on arthropod assemblages. *Funct. Ecol.* **32**, 1317–1328.

19. Jorge TF, Mata AT, António C. 2016 Mass spectrometry as a quantitative tool in plant

metabolomics. *Phil. Trans. R. Soc. A* **374**, 20150370.

20. Ekanayaka EAP, Celiz MD, Jones AD. 2015 Relative mass defect filtering of mass spectra: a path

to discovery of plant specialized metabolites. *Plant Physiol.* **167**, 1221–1232.

(doi:10.1104/pp.114.251165)

21. Langfelder P, Horvath S. 2008 WGCNA: an R package for weighted correlation network analysis.

BMC Bioinformatics **9**, 559.

22. Ogutu JO, Schulz-Streeck T, Piepho H-P. 2012 Genomic selection using regularized linear

regression models: ridge regression, lasso, elastic net and their extensions. In *BMC proceedings*, p.

S10.

23. Team RDC, R Development Core Team R. 2016 R: A Language and Environment for Statistical

Computing. *R Found. Stat. Comput.* (doi:10.1007/978-3-540-74686-7)

24. Friedman J, Hastie T, Simon N, Tibshirani R. 2016 Lasso and Elastic-Net Regularized

Generalized Linear Models. R-package version 2.0-5. 2016.

25. Plummer M, others. 2003 JAGS: A program for analysis of Bayesian graphical models using

Gibbs sampling. In *Proceedings of the 3rd international workshop on distributed statistical*

computing,

26. Gelman A, Rubin DB, others. 1992 Inference from iterative simulation using multiple sequences.

Stat. Sci. **7**, 457–472.

27. Brooks SP, Gelman A. 1998 General methods for monitoring convergence of iterative simulations.

J. Comput. Graph. Stat. **7**, 434–455.

28. Delaney NJ, Chatterjee S. 1986 Use of the bootstrap and cross-validation in ridge regression. *J. Bus. Econ. Stat.* **4**, 255–262.

29. Levin DA. 1976 The chemical defenses of plants to pathogens and herbivores. *Annu. Rev. Ecol. Syst.* **7**, 121–159.

30. Hunter MD. 2016 *The phytochemical landscape: linking trophic interactions and nutrient dynamics*. Princeton University Press.

31. Forister ML, Nice CC, Fordyce JA, Gompert Z. 2009 Host range evolution is not driven by the optimization of larval performance: the case of Lycaeides melissa (Lepidoptera: Lycaenidae) and the colonization of alfalfa. *Oecologia* **160**, 551–561. (doi:10.1007/s00442-009-1310-4)

32. Seigler D, Price PW. 1976 Secondary compounds in plants: primary functions. *Am. Nat.* **110**, 101–105.

33. Smilanich AM, Fincher RM, Dyer LA. 2016 Does plant apparency matter? Thirty years of data provide limited support but reveal clear patterns of the effects of plant chemistry on herbivores. *New Phytol.* **210**, 1044–1057.

34. Chaieb I. 2010 Saponins as insecticides: a review. *Tunis. J. Plant Prot.* **5**, 39–50.

35. Behmer ST. 2009 Insect herbivore nutrient regulation. *Annu. Rev. Entomol.* **54**.

36. Berenbaum MR. 1995 Turnabout is fair play - secondary roles for primary compounds. *J. Chem. Ecol.* **21**, 925–940.

37. Stamp N. 2003 Out of the quagmire of plant defense hypotheses. *Q. Rev. Biol.* **78**, 23–55.

38. Harrison JG *et al.* 2016 The many dimensions of diet breadth: phytochemical, genetic, behavioral, and physiological perspectives on the interaction between a native herbivore and an exotic host. *PLoS One* **11**, e0147971.

39. Jansen JJ, Allwood JW, Marsden-Edwards E, van der Putten WH, Goodacre R, van Dam NM. 2009 Metabolomic analysis of the interaction between plants and herbivores. *Metabolomics* **5**, 150.

Author contributions:

MLF: designed experiment, conducted analyses, wrote first draft.

SY: conducted experiment and contributed to experimental design.

CSP, CDD, BH: generated and interpreted phytochemistry and protein data.

MLF, JGH, OS: developed and maintained common garden.

JAF, ZHM, CCN, LAR: contributed to analyses and experimental design.

CAB, JAF, ZG, SL, CCN: contributed to experimental design.

All authors: contributed to writing.

Figure 1. Variation among plants in caterpillar survival (a), development time (b) adult weight (c), three individual compounds (d-e) and two external plant traits, specific leaf area (g) and leaf toughness (h). The three example compounds shown here (out of the 163 assayed) were among the top five most influential compounds for survival, development time and adult weight: cpd. 9 is a peptide with a negative association with survival, cpd. 94 (another peptide) has a negative association with development time, and cpd. 160 is a phospholipid with a negative association with adult weight. Individual plants in all panels are organized from left to right by decreasing caterpillar survival in the top panel (a). Standard errors are shown for panels b, c, g and h. The units for d-e are compound relative abundance per dry weight of sample; the units for specific leaf area are cm^2/mg, and grams/newton for leaf toughness.

Figure 2. Illustration of correlational structure among compounds: each node in the network is a compound, and compounds are linked by a line if they are correlated among individual plants at 0.5 or above (links among compounds in modules 12-14 represent weaker correlations, greater than 0.1; see main text for details). Two letter codes within nodes indicate compound classes, as explained in the legend. Colors of nodes correspond to membership in modules as determined by hierarchical cluster analysis; the color key to the 14 modules is shown in the lower left. Not shown are a small number of compounds with weak connections to all other compounds, including two compounds that were not included in any module (shown as module zero in figure 3).

Figure 3. Effects of individual compounds on survival, development time and adult weight, as estimated by ridge regression (using binomial, Poisson and Gaussian models, respectively). The strength of effect for each compound is indicated by the horizontal extent of each bar, and compounds are grouped by modules (m1, m2, etc.); the order of compounds along the vertical axis is arbitrary within modules and fixed across columns. Orange colors indicate negative effects on survival, development and weight, while blue colors are positive effects. The darker shades of orange and blue mark coefficients whose 95% confidence intervals did not overlap zero in 1,000 bootstrap samples. Values for survival and development time have been back-transformed from units on the log-odds and log scales to units of probability and days to pupation, and are shown as changes from the mean or intercept values. For example, a negative (orange) survival coefficient of 0.005 means a one-half percent reduction in average probability of survival associated with variation in a particular compound. The ten compounds with the largest coefficients (by absolute value) and bootstrap intervals not overlapping zero are labelled by compound classes (see figure 2 for abbreviations) in each panel.

Figure 4. Violin plots of compound-specific effects (coefficients from ridge regressions) summarized by chemical classes. Plots show median (black dot), interquartile range (box) and 95% confidence intervals (whiskers) surrounded by kernel density envelopes. Sample sizes for each category as follows: l5 lipids, 17 terpenoids, 24 saponins, 48 peptides, 43 phospholipids, 7 phenolics and 9 other. Categories are as shown in electronic supplementary material table S1, with the exception of terpenoids (which is shown here as a pooled category of 5 sterols, 5 vitamins and 7 carotenoids) and "other" (which is 3 alkaloids, 2 amino acids, 1 halogenated compound and 3 sugars). Categories are arranged from left to right based on the gradient of median positive to negative effects on survival. Coefficients for survival (a) and development time (b) have been back-transformed from the units of log-odds and log to probability and days to pupation, respectively, and shown as deviations from the mean or intercept value (as in figure 3).

Table 1. Results from Bayesian regressions of module eigenvectors and covariates predicting caterpillar survival, development time and adult weight (as binomial, Poisson, and Gaussian regressions, respectively, with corresponding units in log-odds, log number of days, and milligrams). For each regression coefficient, numbers in parentheses are 95% credible intervals (the first two numbers) and the probability that the coefficient has the estimated sign (e.g., 0.63 for the m2 survival coefficient of 0.37 indicates a 63% probability that the m2 module has a positive effect on survival). Note that negative coefficients for development time indicate faster caterpillar development (fewer days) associated with variation in a particular compound. Modules (listed in the left column) are only shown if they were included in one of the three regressions following feature selection using lasso regression (see main text for additional details). Empty spaces in the table appear if a particular module was selected through lasso regression for one or two analyses but not all three (m3, for example, was not selected by lasso regression for development time). Slash marks (/) indicate variables not considered for a particular analysis (e.g., sex, adult weight [mg] and development time [days] were not possible for the survival analysis because they are not observed on dead individuals). Values for "validation" shown in the last row are the correlation between observed and predicted values in a cross-validation analysis (electronic supplementary material figure S4).

	Survival coefficient (CI; prob.)	Development time coefficient (CI, prob.)	Weight coefficient (CI, prob.)
m2	0.37 (-1.87, 2.67; 0.63)	-0.07 (-0.26, 0.12; 0.78)	
m3	-2.33 (-3.94, -0.72; >0.99)		-2.33 (-4.87, 0.24; 0.96)
m6		-0.05 (-0.19, 0.09; 0.75)	
m9	-2.31 (-4.49, -0.15; 0.98)	0.064 (-0.11, 0.24, 0.78)	
m10	2.54 (0.81, 4.23; >0.99)		3.56 (1.22, 5.84; >0.99)
m11	2.01 (0.64, 3.42; >0.99)		
SLA	-0.31 (-0.55, -0.08; >0.99)		-0.42 (-0.78, -0.06; 0.99)
Protein	0.061 (-0.16, 0.28; 0.71)		0.08 (-0.26, 0.42; 0.68)
Tough.	0.036 (-0.17, 0.24; 0.36)	-0.002 (-0.02, 0.02; 0.57)	
Sex	/	0.06 (0.02, 0.10; >0.99)	1.10 (0.38, 1.83; >0.99)
mg	/	-0.03 (-0.05, -0.01; >0.99)	/
Days	/	/	1.10 (0.38, 1.83; >0.99)
Intercept	0.34 (0.14, 0.53; >0.99)	3.48 (3.45, 3.52; >0.99)	10.36 (9.81, 10.92; >0.99)
Validation	0.53	0.59	0.50

Figure 1. Variation among plants in caterpillar survival (a), development time (b) adult weight (c), three individual compounds (d-e) and two external plant traits, specific leaf area (g) and leaf toughness (h). The three example compounds shown here (out of the 163 assayed) were among the top five most influential compounds for survival, development time and adult weight: cpd. 9 is a peptide with a negative association with survival, cpd. 94 (another peptide) has a negative association with development time, and cpd. 160 is a phospholipid with a negative association with adult weight. Individual plants in all panels are organized from left to right by decreasing caterpillar survival in the top panel (a). Standard errors are shown for panels b, c, g and h. The units for d-e are compound relative abundance per dry weight of sample; the units for specific leaf area are cm^2/mg, and grams/newton for leaf toughness.

Figure 2. Illustration of correlational structure among compounds: each node in the network is a compound, and compounds are linked by a line if they are correlated among individual plants at 0.5 or above (links among compounds in modules 12-14 represent weaker correlations, greater than 0.1; see main text for details). Two letter codes within nodes indicate compound classes, as explained in the legend. Colors of nodes correspond to membership in modules as determined by hierarchical cluster analysis; the color key to the 14 modules is shown in the lower left. Not shown are a small number of compounds with weak connections to all other compounds, including two compounds that were not included in any module (shown as module zero in figure 3).

Figure 3. Effects of individual compounds on survival, development time and adult weight, as estimated by ridge regression (using binomial, Poisson and Gaussian models, respectively). The strength of effect for each compound is indicated by the horizontal extent of each bar, and compounds are grouped by modules (m1, m2, etc.); the order of compounds along the vertical axis is arbitrary within modules and fixed across columns. Orange colors indicate negative effects on survival, development and weight, while blue colors are positive effects. The darker shades of orange and blue mark coefficients whose 95% confidence intervals did not overlap zero in 1,000 bootstrap samples. Values for survival and development time have been back-transformed from units on the log-odds and log scales to units of probability and days to pupation, and are shown as changes from the mean or intercept values. For example, a negative (orange) survival coefficient of 0.005 means a one-half percent reduction in average probability of survival associated with variation in a particular compound. The ten compounds with the largest coefficients (by absolute value) and bootstrap intervals not overlapping zero are labelled by compound classes (see figure 2 for abbreviations) in each panel.

Figure 4. Violin plots of compound-specific effects (coefficients from ridge regressions) summarized by chemical classes. Plots show median (black dot), interquartile range (box) and 95% confidence intervals (whiskers) surrounded by kernel density envelopes. Sample sizes for each category as follows: l5 lipids, 17 terpenoids, 24 saponins, 48 peptides, 43 phospholipids, 7 phenolics and 9 other. Categories are as shown in electronic supplementary material table S1, with the exception of terpenoids (which is shown here as a pooled category of 5 sterols, 5 vitamins and 7 carotenoids) and "other" (which is 3 alkaloids, 2 amino acids, 1 halogenated compound and 3 sugars). Categories are arranged from left to right based on the gradient of median positive to negative effects on survival. Coefficients for survival (a) and development time (b) have been back-transformed from the units of log-odds and log to probability and days to pupation, respectively, and shown as deviations from the mean or intercept value (as in figure 3).