1

Caterpillars on a phytochemical landscape

Matthew L. Forister[1,2], Su'ad Yoon[1,2], Casey S. Philbin[2,3], Craig D. Dodson[2,3], Bret Hart[4], Joshua G. Harrison[5], Oren Shelef[6], James A. Fordyce[7], Zachary H. Marion[8], Christopher C. Nice[9], Lora A. Richards[1,2], C. Alex Buerkle[5] , and Zach Gompert[10]

[1] Department of Biology, Program in Ecology, Evolution and Conservation Biology, University of Nevada, Reno, NV 89557, USA

[2] Hitchcock Center for Chemical Ecology, University of Nevada, Reno, NV 89557, USA

[3] Department of Chemistry, University of Nevada, Reno, NV 89557, USA

[4] Department of Biochemistry, University of Nevada, Reno, NV 89557, USA

[5] Department of Botany and Program in Ecology, University of Wyoming, Laramie, WY 82071, USA

[6] Department of Natural Resources, Institute of Plant Sciences, Volcani Center, Agricultural Research Organization, Rishon LeZion, Israel

[7] Department of Ecology and Evolutionary Biology, University of Tennessee, Knoxville, TN 37996, USA

[8] Department of Ecology, University of Canterbury, Christchurch, New Zealand

[9] Department of Biology, Population and Conservation Biology Program, Texas State University, San Marcos, TX 78666, USA

[10] Department of Biology, Utah State University, Logan, UT 84322, USA

Author for correspondence: Matthew L. Forister, +1 (775) 784-6770, email: forister@gmail.com

Summary

• Modern metabolomic approaches that generate more comprehensive phytochemical profiles than were previously available are providing new opportunities for understanding plant-animal interactions. Specifically, we can characterize the phytochemical landscape by asking how many individual compounds affect herbivores and how compounds covary among plants.

• Here we use the recent colonization of alfalfa (*Medicago sativa*) by the Melissa blue butterfly (*Lycaeides melissa*) to quantify primary and secondary plant metabolites and the performance of caterpillars as affected by both individual compounds and suites of covarying phytochemicals.

• We find that survival, development time and adult weight are all associated with variation in a large number of compounds, including biomolecules associated with plant cell function as well as putative anti-herbivore action. The dimensionality of the plant-insect interface is high, with clusters of covarying compounds in many cases encompassing divergent effects on different aspects of caterpillar performance.

• Individual compounds with the strongest associations tend to be secondary metabolites, including alkaloids, phenolic glycosides and saponins. The saponins are represented in our data by more than 25 individual compounds with beneficial and detrimental effects on caterpillars, which highlights the value of metabolomic data as opposed to approaches that rely on total concentrations within defensive classes.

Keywords: *Lycaeides melissa, Medicago sativa*, metabolomics, plant defense, specialization

Introduction

One of the conceptual pillars of trophic ecology is the idea that herbivores must overcome the barrier of plant secondary chemistry before extracting the nutrients necessary for growth and reproduction (Feeny *et al.*, 1992). The success of this idea is reflected in areas of research that include coevolution (Agrawal *et al.*, 2012), ecological specialization (Dyer, 1995), and nutrient flow in ecosystems (Olson, 1963). In most cases, progress has been made by chemical ecologists focusing on small subsets of the secondary metabolites produced by plants and consumed by herbivores. The focus on a few charismatic molecules or classes of compounds, such as furanocoumarins (Berenbaum, 1983) or cardiac glycosides (Zalucki *et al.*, 2001), was at least in part necessitated by early methods in natural products chemistry that were targeted and not easily optimized for the discovery of large suites of co-occurring primary and secondary metabolites (Maag *et al.*, 2015; Dyer *et al.*, 2018). As technological limitations have dissipated, the opportunity now exists for a deeper understanding of the challenges faced by herbivores, with the possibility of discovering, among other things, novel compounds and synergistic interactions among compounds (Prince & Pohnert, 2010; Richards *et al.*, 2010; Sardans *et al.*, 2011). More generally, an important task is to quantify the phytochemical dimensionality of the antagonistic interaction between plants and herbivores, with an eye towards understanding constraints on the evolution of both players (Fordyce & Nice, 2008; Macel *et al.*, 2010) and predicting the formation of new plant-herbivore interactions (Erbilgin, 2018). Here we use the example of a specialized herbivore and a novel host plant to investigate the phytochemical landscape from the perspective of developing caterpillars.

The Melissa blue butterfly, *Lycaeides melissa*, specializes on larval host plants in the pea family (Fabaceae), primarily in the genera *Astragalus* and *Lupinus*. Within the last 200 years, *L. melissa* has colonized introduced alfalfa, *Medicago sativa* (Fabaceae), at least twice and probably multiple times (Chaturvedi *et al.*, 2018), forming a heterogeneous patchwork of association throughout the range of the butterfly in western North America. *M. sativa* is a suboptimal host, relative to native hosts that have been examined, and populations of the butterfly that persist on *M. sativa* show evidence of loss of preference for native hosts (Forister *et al.*, 2012), reduced caterpillar performance on native hosts, and a slight increase in ability to develop on the suboptimal novel host (Gompert *et al.*, 2015). The genetic architecture of host

use in this system is known to be polygenic and characterized by loci with conditionally neutral (host-specific) effects (Gompert *et al.*, 2015). What is needed next is an understanding of how many independent or covarying phytochemical compounds have consequential effects on caterpillars eating the novel host. For example, will the trajectory of further local adaptation by *L. melissa* to *M. sativa* be a matter of evolving the ability to detoxify one or a large number of compounds? We would also like to know how key compounds covary among individual plants, which should help us understand the puzzle faced by ovipositing females encountering an array of co-occurring *M. sativa* chemotypes. Here we use a common garden approach (to minimize non-genetic phenotypic variation among plants) and caterpillars individually reared in a controlled environment to address these questions while describing the effects of metabolomic variation in *M. sativa* on *L. melissa*.

Materials and Methods

Plants and caterpillars

Plants used in this project were grown at the University of Nevada, Reno, Main Station experimental farm. The common garden was planted in 2016 with seeds collected the previous year from 45 plants (previously studied by Harrison et al. (Harrison *et al.*, 2018)) growing in a fallow field in north-western Nevada on the western edge of the Great Basin Desert. The focal butterfly, *L. melissa*, was present in the source field but has not yet colonized the university farm where experimental plants were grown. The 45 maternal plants each contributed 15 offspring to a randomized grid design in the common garden, irrigated with broadcast sprayers in 2016 and drip in 2017, without supplemental fertilization. Out of each maternal family, a single plant was randomly selected for use in the rearing experiment reported here as a way to capture as much genetic and phenotypic variation as possible.

On 17 and 18 July 2017, a total of 45 *L. melissa* females were collected from an alfalfa-associated population near Verdi, NV, and confined to oviposition arenas (500 mL plastic cups) in groups of three with host plant leaves and supplied with Gatorade on mesh lids. After three days, eggs were removed from leaves, pooled, and kept at room temperature until hatching, at which time caterpillars were placed individually in Petri dishes (100 x 25 mm) with leaves of a particular *M. sativa* individual (which became the only plant from which they were fed

throughout the experiment). Ten caterpillars were assigned to each of the 45 experimental *M. sativa* plants (for a total of 450 independently-reared caterpillars) and kept in a growth chamber set to 25 C and a 12 hour light / 12 hour dark cycle. From each caterpillar we recorded survival to adult, date of eclosion (if successful) and adult weight to the nearest 0.01 mg on a Mettler Toledo XP26 microbalance.

Phytochemistry and plant traits

Metabolomic variation among individual plants was characterized with liquid chromatography–mass spectrometry (LC-MS, (Jorge *et al.*, 2016)) using leaves collected on a single day at the start of the rearing experiment. Dried, ground leaves (10 mg) were extracted in 2 mL of 70% aqueous ethanol and injected into an Agilent 1200 analytical high performance liquid chromatograph paired with an Agilent 6230 Time-of-Flight mass spectrometer via an electrospray ionization source. Resulting chromatograms were analyzed using MassHunter Quantitative Analysis (v.B.06.00, Agilent, Santa Clara, CA); see electronic supporting information for additional phytochemical protocols. Major classes of compounds were identified using characteristic relative mass defects (Ekanayaka *et al.*, 2015). Leaf protein content was quantified with three replicates (~2 mg each) per plant using the Bicinchoninic acid assay (Pierce Biotechnology, Waltham, MA). Before grinding, five dried leaflets from each sample were weighed to the nearest 0.1 mg, scanned, and area was measured using ImageJ (v.1.52a); specific leaf area (SLA) was calculated as leaf area divided by dry mass. Finally, leaf toughness was measured on fresh material in the common garden, at the start of the experiment (mid-July, when leaves were also sampled for chemistry and protein) and at the end of the experiment (mid-August), from three leaves per plant at each date, with a penetrometer (Chatillon 516 Series) through the center of the middle leaflet, as in (Harrison *et al.*, 2018); the three leaves were selected haphazardly, avoiding the oldest and youngest leaves. The six leaf toughness measurements per plant were averaged for a single toughness measure used in analyses.

Analyses of plant traits and caterpillar performance

Overview

Our analytical strategy to understand the association between phytochemical variation and caterpillar performance followed two complementary paths, one focusing on dimension reduction and feature selection to produce relatively simple models, and the other on the estimation of effects of all individual compounds. For the first path, involving dimension reduction and feature selection, we utilized an approach developed for gene transcription studies that identifies groups or modules of correlated variables with hierarchical clustering (Langfelder & Horvath, 2008); after clustering, we reduced the number of independent variables by selecting among modules and other plant traits (specific leaf area, protein and leaf toughness) using lasso regression (Ogutu *et al.*, 2012). Selected modules (and other plant traits) were analyzed in Bayesian linear models that are useful in this context because they allowed us to quantify our confidence in the sign of effects (positive or negative) as continuous probabilities (as opposed to relying on arbitrary significance cutoffs). For the second analytical path, we utilized ridge regression (Ogutu *et al.*, 2012) to estimate effects for all compounds simultaneously, which allowed us to investigate the distribution of effects among compounds and classes of compounds. Both analytical paths incorporated cross-validation during the lasso and ridge regressions, and as a means of evaluating the predictive success of the Bayesian models. We also used randomization tests to compare the performance of modules and individual compounds with randomly-chosen suites of compounds.

Dimension reduction and feature selection

We chose an approach that reduces the number of independent variables while allowing us to learn something about the correlational structure of the data, specifically unsupervised hierarchical clustering as implemented in the blockwiseModules function of the WGCNA package (Langfelder & Horvath, 2008) in R (Team & R Development Core Team, 2016). Among the options in the pipeline, we used positive correlations among variables ("signed" network type), merge cut height at 0.25, and correlations raised to the power of five (which is where the scale free topology index reached a plateau). Through experimentation, we found that our results with LC-MS data were robust to variation in these choices, including the choice of signed or unsigned networks. After an initial round of clustering, we took a remaining 19 unassigned compounds and put them through a second round of clustering (although the majority

of consequential compounds were identified in the first round). One output of the WGCNA procedure is the first eigenvector from each cluster of compounds, which reduced our number of predictor variables by a factor of ten. The resulting eigenvectors plus protein, SLA (specific leaf area) and leaf toughness were then put through the feature reduction step of lasso regression (Ogutu *et al.*, 2012), a penalized regression that allows beta coefficients to be constrained to zero (thus excluding variables). We used the cv.glmnet function of the glmnet package (Friedman *et al.*, 2016) with cross-validation during error reduction set to leave out one plant (and associated caterpillars) at each iteration. The variables selected by the lasso were then put into a Bayesian linear model to estimate coefficients and associated credible intervals using JAGS (version 3.2.0) run in R with the rjags package (Plummer & others, 2003). Two Markov chains were run for 10,000 steps for each analysis (no burn in was required) and chain performance was assessed by plotting chain histories, and calculating the Gelman and Rubin convergence diagnostic and effective sample sizes (Gelman *et al.*, 1992; Brooks & Gelman, 1998). For all models, uninformative priors for the regression coefficients were modeled as a normal distribution with a mean of zero and variance of 100 (variance = 1/precision). We quantified our confidence in the sign of coefficients (positive or negative) as the fraction of the posterior samples that were less than zero (for coefficients with a median negative value) or greater than zero (for coefficients with a median positive value).

All analyses were done using the R statistical language (Team & R Development Core Team, 2016) on scaled (z-transformed) predictor variables, and both lasso and Bayesian models used binomial (for survival), Poisson (for development time) and Gaussian (for adult weight) errors. The latter two analyses (development time and adult weight) included sex as a factor. The analysis of development time also included adult weight as a covariate; while (reciprocally) the analysis of adult weight included development time as a predictor. These variables are negatively correlated (at -0.52), and they function as useful covariates of each other, allowing us to investigate the possibility of unique plant effects on weight gain and development time, which could not be discovered if, for example, these variables were combined into a single performance index.

The success of models developed with the dimension reduction and feature selection pipeline was judged in two ways. We used a cross-validation procedure in which we left out one plant (and associated caterpillars) in each iteration of the Bayesian model and then used the

estimated coefficients (for phytochemical variables and other plant traits) to predict the performance of the unobserved caterpillars. After 45 iterations (one for each plant), we calculated a simple correlation coefficient between the observed and predicted performance of caterpillars across plants. In addition, we repeatedly resampled the original LC-MS data to match the structure of the reduced set of predictor variables to ask to what extent randomly assembled modules could outperform the empirically-derived modules.

Individual compound effects

The second path of our two-part analytical strategy involved simultaneous estimation of the effects of all individual chemical compounds on caterpillar survival, development time and adult weight. For this approach, we again used penalized regression (in the glmnet package (Friedman *et al.*, 2016)), but this time with ridge regression (instead of lasso) which constrains beta coefficients to avoid variance inflation but does not eliminate variables. As with the analyses above, ridge regression was done using error structures appropriate to the specific response variables, and included additional covariates where possible (in models of development time and adult weight). The resulting coefficients associated with all individual compounds were examined as a second perspective on the modules examined in the first set of analyses, and were used to ask to what extent individual compound effects could be predicted by the degree to which they vary among individual plants as quantified with the simple coefficient of variation. To assess confidence in the results of ridge regressions, we used a bootstrap approach, repeatedly resampling the data and estimating coefficients 1000 times, noting the compounds whose bootstrap confidence intervals did or did not overlap zero (Delaney & Chatterjee, 1986). We also allowed for the discovery of interactions among compounds using penalized regression on all individual compounds and all pairwise interactions between compounds. For ease of interpretation, this final analysis of potential interactions used lasso (not ridge) regression, letting the coefficients for many of the individual compounds and pairwise interactions go to zero.

Results

Of the 450 caterpillars that started the experiment, 261 were reared to eclosion as adults (a

mortality rate similar to previous work with this system (Gompert *et al.*, 2015)) on leaves from

45 individual alfalfa plants that were characterized for protein, leaf toughness, specific leaf area

and 163 individual metabolomic features (see Fig. 1 for variation among plants in caterpillar

performance and a subset of plant traits, and Supporting Information Table S1 for a list of

compounds). Hierarchical clustering identified 14 subsets (or modules) of compounds with

generally low correlations among modules and high correlations within modules (see Supporting

Information Figs. S1 and S2 for correlations within and among modules, and Fig. S3 for module

variation among plants). The correlational structure of the phytochemical data is illustrated as an

adjacency network in Fig. 2, where it can be seen that some modules (e.g., modules 1 and 2)

contain a great diversity of compound types, while other modules are made up of more narrow

classes (e.g. modules 7 and 8 are mostly saponins, a class of defensive secondary metabolites

(Levin, 1976)). From the 14 eigenvectors summarizing variation in the modules, as well as the

other plant traits, lasso regression produced a reduced set of potential predictors which were then

used in Bayesian multiple regression models that included between seven and nine independent

variables (Table 1). The models had reasonably high performance in leave-one-out cross-

validation (correlations between observed and predicted values were between 0.50 and 0.59,

Table 1), and also in resampling analyses (Supporting Information Fig. S4), where a small

fraction (never more than 4%) of randomly-generated models exceeded the variance explained of

the models reported in Table 1.

Variation among plants in the suites of covarying compounds had large effects on

caterpillar performance: for example, the beta coefficient of -2.33 (on the log-odds scale)

associated with module 3 corresponds to a reduction in mean survival from 0.58 to 0.12

associated with a one unit change in variation associated with that phytochemical module (Table

1; note that in Table 1 and elsewhere negative coefficients for development time are associated

with fewer days, and thus can be thought of as potentially beneficial effects, in contrast to

negative coefficients for survival and weight that are detrimental to caterpillars). The

phytochemical predictor variables are eigenvectors from clustering analysis, and thus are not

entirely straightforward to interpret, especially when the clustering analysis was itself based on

z-transformed data. It is useful to note that our LC-MS data consists of peak areas divided by the

peak of an internal standard, and again divided by the dry weight of the sample (thus, in total,

referred to as "relative abundance per dry weight"). Variation in these numbers reflects variation

in concentrations within compounds (among plants), but care should be used in comparing

among compounds because of different ionization responses relative to the standard (thus the use

of z-transformation for among-compound analyses). Nevertheless, the effects reported in Table

1 reflect real variation in suites of compounds, as can be seen in correlations between

eigenvectors and individual compounds in Supporting Information Fig. S2, and in variation

among plants in average z-scores in Fig. S3.

Modules included in multiple regression models frequently had common effects across

response variables (e.g., the positive association of module 10 with both survival and adult

weight), with the exceptions of module 11 that had a solitary effect on survival and module 6

with an effect only on development time (although the probability of the latter having a negative

effect was only 0.75). Specific leaf area had a negative effect on survival and adult weight, and

the coefficients for specific leaf area (-0.31 for survival and -0.42 for weight) were of smaller

magnitude than most phytochemical effects.

Module-based analyses (as in Table 1) focused on feature reduction with lasso regression;

as a complementary analytical approach, we used ridge regression on all of the compounds

(which estimates effects of individual compounds without excluding variables as in lasso

regression). Analyses of individual compounds by ridge regression (Fig. 3) were broadly

consistent with the strongest module-specific effects, as can be seen, for example, with module

10 having positive effects on survival and adult weight in module analyses (Table 1) and in

compound-specific analyses (Fig. 3). Similarly, the individual compounds in module 3 had

negative compound-specific effects on survival (Fig. 3), and that module had the strongest

negative effect on survival in the eigenvector-based analyses in Table 1. Not surprisingly, the

larger modules (with a greater number of covarying compounds, including many primary

metabolites) tended to have a more complex mix of positive and negative effects (for example,

modules 1 and 2, Fig. 3). For ease of interpretation, the coefficients from compound-specific

regressions of survival and development time (in Figs. 3 and 4) have been back-transformed to

be on the scales of probability and days (respectively), and displayed as changes relative to

intercepts. For example, a compound with a relatively large effect on survival in Fig. 3 could be

associated with a one half percent (0.005) reduction in the probability of survival relative to

average survival and while holding other compounds constant.

We also considered potential pairwise interactions among individual compounds, and

found few interactions that passed the filter of the penalized regression (Supporting Information

Table S2), at least relative to the large number of potential interactions. Saponins and alkaloids

tended to be overrepresented in the interactions that were detected, and phenolic glycosides were

involved in stronger negative interactions relative to other compounds (Fig. S5). We did not find

evidence that more or less variable compounds had differential effects on caterpillars, although

there was a trend towards both greater positive and greater negative effects being associated with

less variable compounds (Fig. S6). We saw some variation among classes of compounds in their

effects on caterpillars (Fig. 4). All classes included positive and negative effects, with saponins,

alkaloids and phenolic glycosides including some of the stronger negative effects of individual

compounds, while lipids and sterols tended towards positive associations with survival and

development (Fig. 4).

Discussion

The results reported here represent a dissection of the phytochemical landscape facing a

specialized insect herbivore attacking a novel host plant (Hunter, 2016). The phytochemical

landscape is both physical, referring to variation in compounds among individual plants in a

common garden, and hypothetical to the extent that effects of individual compounds on

caterpillars are estimated, although compounds are of course not encountered in isolation. Our

exploration of the phytochemical landscape facing *L. melissa* on *M. sativa* is necessarily a first

draft based on a snapshot in time, and the implications of this are discussed further (below) in the

context of within and among-population heterogeneity in the system. Nevertheless, models

including suites of covarying compounds and other plant traits had predictive success and

suggested different natural products affecting survival, development time and adult weight

(performance measured as adult weight is a proxy for fecundity (Forister *et al.*, 2009)). Previous

work with *M. sativa* and insect herbivores has focused on sapogenic glycosides (Levin, 1976),

and a simple outcome from our study could have been that one or a small number of saponins

have anti-herbivore properties that reduce fitness of our focal insect. Instead, we find large

numbers of compounds with potentially consequential effects on caterpillars (Fig. 3), and the

effects of those compounds were greater than the effects of leaf toughness and specific leaf area

(Carmona *et al.*, 2011).

The precise identification of specific compounds is perhaps not as important as the result

that prominent classes of secondary metabolites, such as saponins and peptides, include

compounds with both positive and negative effects. Positive effects of these compounds are

potentially associated with feeding stimulation, as has been observed (along with other positive

effects) for other specialist herbivores and plant toxins (Seigler & Price, 1976; Smilanich *et al.*,

2016). Negative effects of saponins on insects potentially include disruption of hormone

production (Chaieb, 2010), although exact modes of action on *L. melissa* will await further study.

Although most of the individual compounds with strong effects appear to be secondary

metabolites (including alkaloids and phenolic glycosides, as well as saponins and peptides), we

also find both positive and negative effects associated with variation in certain primary

metabolites (Fig. 4). For example, the larger detrimental effects of individual compounds on

caterpillars include phospholipids (Fig. 2). These could be direct effects if a compound is

suboptimal for development, or they could be associated with nutritional imbalance (Behmer,

2009), such that too much of one nutrient makes it difficult for caterpillars to consume a

balanced diet. It has been suggested that the presentation of unbalanced nutrition can be a kind

of anti-herbivore strategy (Berenbaum, 1995). Although this possibility has not been thoroughly

investigated in many systems with full metabolomic profiling, the idea that nutritional imbalance

could be as important as direct toxicity suggests that we might update theories of the evolution of

plant defense that were built on differential investment into simple categories of plant growth

versus defense (Stamp, 2003).

The finding that our specialist herbivore is affected by a wide range of metabolites,

primary and secondary, that vary greatly even within a single host population has implications

for our understanding of heterogeneity in the system and for the course of local adaptation of the

herbivore to the novel host. *Lycaeides melissa* typically colonizes weedy or feral patches of *M.*

sativa on roadsides or integrated into natural communities, and previous work has documented

dramatic variation among individual alfalfa locations (often in close proximity) in the extent to

which they can support caterpillar development (Harrison *et al.*, 2016). Previous phytochemical

data with a lower resolution was less successful in explaining that variation (Harrison *et al.*,

2016), and the results reported here suggest that among patch variation could be explained by

future studies using metabolomic data as used here. The within-population complexity described in the current study combined with previous evidence for dramatic among-population variation in *M. sativa* suitability for the focal herbivore also raises the possibility that the novel host presents a multi-faceted and potentially ever-shifting target from the perspective of evolving butterfly populations. In particular, it is possible that *M. sativa* defense against a specialist herbivore might be realized through different combinations (within and among populations) of individually-acting compounds, thus making it unlikely for butterflies in any one population to possess an effective suite of alleles that improve fitness on the novel host. In this context, it is interesting to note that a molecular genetic dissection of caterpillar performance in this system found a large number (potentially hundreds) of individual loci associated with performance on *M. sativa*, yet evolution in populations associated with the novel host is primarily associated with a loss (through genetic drift) of the ability to eat a native host and only slight improvement in the ability to eat the novel host (Gompert *et al.*, 2015).

The correlational structure of the phytochemical variation that we observed has implications for the evolution of plant defense and the accumulation of insect herbivores on *M. sativa*. Specifically, correlations among modules should make it possible to hypothesize directions of least resistance for defense evolution. Compounds in module 9 had a negative effect on survival (Table 1), and module 9 negatively covaried with module 10 (Supporting Information Fig. S1), which itself had a positive association with caterpillar survival. Thus an increase in module 9 and an associated decrease in 10 would be beneficial for the plant, at least with respect to herbivory by our focal herbivore. Of course, most plants do not have the luxury of optimizing defense against a single herbivore, and it is easy to imagine that improvements in defense against one enemy could lead to increased attraction to another, especially given the diversity of effects even within major classes studied here, including saponins and phenolic glycosides. Compounds in the latter class (phenolics) were found to have strong positive and negative effects on assemblages of arthropods associated with the maternal plants from which seeds were collected to start the common garden used in the present study (Harrison *et al.*, 2018).

The results reported here raise a number of avenues for future exploration, including the apparent overrepresentation of both saponins and alkaloids in interactions with other compounds (Fig. S5). Relative mass defect (RMD) is a useful tool for the categorization of compounds, but it has limitations in complex mixtures; we are developing methods that use other data from high

resolution mass spectrometry to further refine categorization of *Medicago* metabolites. Also, in the present study, we have not attempted to separate constitutive and induced defenses (Jansen *et al.*, 2009) as the plants in the common garden were exposed to natural and continuous levels of herbivory. We also acknowledge that feeding under laboratory conditions is of course not natural, although we found in a previous study that genetic variants (in caterpillars) associated with success in laboratory feeding trials were at least partially predictive of genetic variation associated with alfalfa host use in the wild (Chaturvedi *et al.*, 2018). Thus it is clear that metabolomic data such as that analyzed here has the potential to both open up new avenues of conceptual development in plant-insect interactions and to link micro-evolutionary trajectories across hosts and herbivores.

Acknowledgements

This work was supported by National Science Foundation grant DEB-1638793 to MLF and CDD, DEB-1638768 to ZG, DEB-1638773 to CCN, DEB-1638922 to SL and JAF, and DEB-1638602 to CAB; MLF was additionally supported by a Trevor James McMinn professorship. Thanks to Ian Wallace, the Hitchcock Center for Chemical Ecology and the P.I.G. group for discussion and expertise.

Author contributions

MLF: designed experiment, conducted analyses, wrote first draft. SY: conducted experiment and contributed to experimental design. CSP, CDD, BH: generated and interpreted phytochemistry and protein data. MLF, JGH, OS: developed and maintained common garden. JAF, ZHM, CCN, LAR: contributed to analyses and experimental design. CAB, JAF, ZG, CCN: contributed to experimental design. All authors: contributed to writing.

References

Agrawal AA, Petschenka G, Bingham RA, Weber MG, Rasmann S. **2012**. Toxic cardenolides: chemical ecology and coevolution of specialized plant--herbivore interactions.

New Phytologist **194**: 28–45.

Behmer ST. **2009**. Insect herbivore nutrient regulation. *Annual review of entomology* **54**.

Berenbaum M. **1983**. Coumarins and caterpillars: a case for coevolution. *Evolution* **37**: 163–
179.

Berenbaum MR. **1995**. Turnabout is fair play - secondary roles for primary compounds. *Journal*
of Chemical Ecology **21**: 925–940.

Brooks SP, Gelman A. **1998**. General methods for monitoring convergence of iterative
simulations. *Journal of computational and graphical statistics* **7**: 434–455.

Carmona D, Lajeunesse MJ, Johnson MTJ. **2011**. Plant traits that predict resistance to
herbivores. *Functional Ecology* **25**: 358–367.

Chaieb I. **2010**. Saponins as insecticides: a review. *Tunisian Journal of Plant Protection* **5**: 39–
50.

Chaturvedi S, Lucas LK, Nice CC, Fordyce JA, Forister ML, Gompert Z. **2018**. The
predictability of genomic changes underlying a recent host shift in Melissa blue butterflies.
Molecular ecology.

Delaney NJ, Chatterjee S. **1986**. Use of the bootstrap and cross-validation in ridge regression.
Journal of Business & Economic Statistics **4**: 255–262.

Dyer LA. **1995**. Tasty generalists and nasty specialists - antipredator mechanisms in tropical
lepidopteran larvae. *Ecology* **76**: 1483–1496.

Dyer LA, Philbin CS, Ochsenrider KM, Richards LA, Massad TJ, Smilanich AM, Forister
ML, Parchman TL, Galland LM, Hurtado PJ, *et al.* 2018. Modern approaches to study
plant–insect interactions in chemical ecology. *Nature Reviews Chemistry* **1**.

Ekanayaka EAP, Celiz MD, Jones AD. **2015**. Relative mass defect filtering of mass spectra: a
path to discovery of plant specialized metabolites. *Plant Physiology* **167**: 1221–1232.

Erbilgin N. **2018**. Phytochemicals as mediators for host range expansion of a native invasive
forest insect herbivore. *New Phytologist*.

Feeny P, Rosenthal GA, Berenbaum MR. **1992**. The evolution of chemical ecology:
contributions from the study of herbivorous insects. *Herbivores: their interactions with*
secondary plant metabolites, **2**: 1–44.

Fordyce JA, Nice CC. **2008**. Antagonistic, stage-specific selection on defensive chemical
sequestration in a toxic butterfly. *Evolution* **62**: 1610–1617.

Forister ML, Nice CC, Fordyce JA, Gompert Z. **2009**. Host range evolution is not driven by the optimization of larval performance: the case of Lycaeides melissa (Lepidoptera: Lycaenidae) and the colonization of alfalfa. *Oecologia* **160**: 551–561.

Forister ML, Scholl CF, Jahner JP, Wilson JS, Fordyce JA, Gompert Z, Narala D, Buerkle CA, Nice CC. **2012**. Specificity, rank preference and the colonization of a non-native host plant by the Melissa blue butterfly. *Oecologia* **DOI: 10.10**.

Friedman J, Hastie T, Simon N, Tibshirani R. **2016**. Lasso and Elastic-Net Regularized Generalized Linear Models. R-package version 2.0-5. 2016.

Gelman A, Rubin DB, others. **1992**. Inference from iterative simulation using multiple sequences. *Statistical science* **7**: 457–472.

Gompert Z, Jahner JP, Scholl CF, Wilson JS, Lucas LK, Soria-Carrasco V, Fordyce JA, Nice CC, Buerkle CA, Forister ML. **2015**. The evolution of novel host use is unlikely to be constrained by trade-offs or a lack of genetic variation. *Molecular Ecology* **24**: 2777–2793.

Harrison JG, Gompert Z, Fordyce JA, Buerkle CA, Grinstead R, Jahner JP, Mikel S, Nice CC, Santamaria A, Forister ML. **2016**. The many dimensions of diet breadth: phytochemical, genetic, behavioral, and physiological perspectives on the interaction between a native herbivore and an exotic host. *PloS One* **11**: e0147971.

Harrison JG, Philbin CS, Gompert Z, Forister GW, Hernandez-Espinoza L, Sullivan BW, Wallace IS, Beltran L, Dodson CD, Francis JS, *et al.* 2018. Deconstruction of a plant-arthropod community reveals influential plant traits with nonlinear effects on arthropod assemblages. *Functional Ecology* **32**: 1317–1328.

Hunter MD. **2016**. *The phytochemical landscape: linking trophic interactions and nutrient dynamics*. Princeton University Press.

Jansen JJ, Allwood JW, Marsden-Edwards E, van der Putten WH, Goodacre R, van Dam NM. **2009**. Metabolomic analysis of the interaction between plants and herbivores. *Metabolomics* **5**: 150.

Jorge TF, Mata AT, António C. **2016**. Mass spectrometry as a quantitative tool in plant metabolomics. *Phil. Trans. R. Soc. A* **374**: 20150370.

Langfelder P, Horvath S. **2008**. WGCNA: an R package for weighted correlation network analysis. *BMC bioinformatics* **9**: 559.

Levin DA. **1976**. The chemical defenses of plants to pathogens and herbivores. *Annual review of*

Ecology and Systematics **7**: 121–159.

Maag D, Erb M, Glauser G. **2015**. Metabolomics in plant-herbivore interactions: challenges and applications. *Entomologia Experimentalis et Applicata* **157**: 18–29.

Macel M, van Dam NM, Keurentjes JJB. **2010**. Metabolomics: the chemistry between ecology and genetics. *Molecular ecology resources* **10**: 583–593.

Ogutu JO, Schulz-Streeck T, Piepho H-P. **2012**. Genomic selection using regularized linear regression models: ridge regression, lasso, elastic net and their extensions. In: BMC proceedings. S10.

Olson JS. **1963**. Energy storage and the balance of producers and decomposers in ecological systems. *Ecology* **44**: 322–331.

Plummer M, others. **2003**. JAGS: A program for analysis of Bayesian graphical models using Gibbs sampling. In: Proceedings of the 3rd international workshop on distributed statistical computing.

Prince EK, Pohnert G. **2010**. Searching for signals in the noise: metabolomics in chemical ecology. *Analytical and bioanalytical chemistry* **396**: 193–197.

Richards LA, Dyer LA, Smilanich AM, Dodson CD. **2010**. Synergistic effects of amides from two Piper species on generalist and specialist herbivores. *Journal of chemical ecology* **36**: 1105–1113.

Sardans J, Penuelas J, Rivas-Ubach A. **2011**. Ecological metabolomics: overview of current developments and future challenges. *Chemoecology* **21**: 191–225.

Seigler D, Price PW. **1976**. Secondary compounds in plants: primary functions. *The American Naturalist* **110**: 101–105.

Smilanich AM, Fincher RM, Dyer LA. **2016**. Does plant apparency matter? Thirty years of data provide limited support but reveal clear patterns of the effects of plant chemistry on herbivores. *New Phytologist* **210**: 1044–1057.

Stamp N. **2003**. Out of the quagmire of plant defense hypotheses. *The Quarterly Review of Biology* **78**: 23–55.

Team RDC, R Development Core Team R. **2016**. R: A Language and Environment for Statistical Computing. *R Foundation for Statistical Computing*.

Zalucki MP, Brower LP, Alonso-M A. **2001**. Detrimental effects of latex and cardiac glycosides on survival and growth of first-instar monarch butterfly larvae Danaus plexippus

feeding on the sandhill milkweed Asclepias humistrata. *Ecological Entomology* **26**: 212–224.

Figure legends (also reproduced below individual figures)

Fig. 1. Variation among plants in caterpillar survival (a), development time (b) adult weight (c), three individual compounds (d-e) and two external plant traits, specific leaf area (g) and leaf toughness (h). The three example compounds shown here (out of the 163 assayed) were among the top five most influential compounds for survival, development time and adult weight: cpd. 9 is an alkaloid with a negative association with survival, cpd. 94 (a peptide) has a negative association with development time, and cpd. 160 is a phospholipid with a negative association with adult weight. Individual plants in all panels are organized from left to right by decreasing caterpillar survival in the top panel (a). Standard errors are shown for panels b, c, g and h. The units for d-e are compound relative abundance per dry weight of sample; the units for specific leaf area are cm^2/mg, and grams/newton for leaf toughness.

Fig. 2. Illustration of correlational structure among compounds: each node in the network is a compound, and compounds are linked by a line if they are correlated among individual plants at 0.5 or above (links among compounds in modules 12-14 represent weaker correlations, greater than 0.1; see main text for details). Two letter codes within nodes indicate compound classes, as explained in the legend. Colors of nodes correspond to membership in modules as determined by hierarchical cluster analysis; the color key to the 14 modules is shown in the lower left. Not shown are a small number of compounds with weak connections to all other compounds, including two compounds that were not included in any module (shown as module zero in Fig. 3).

Fig. 3. Effects of individual compounds on survival, development time and adult weight, as estimated by ridge regression (using binomial, Poisson and Gaussian models, respectively). The strength of effect for each compound is indicated by the horizontal extent of each bar, and compounds are grouped by modules (m1, m2, etc.); the order of compounds along the vertical axis is arbitrary within modules and fixed across columns. Orange colors indicate negative effects on survival, development and weight, while blue colors are positive effects (note that negative effects for development time correspond to fewer days, or more rapid development). The darker shades of orange and blue mark coefficients whose 95% confidence intervals did not overlap zero in 1,000 bootstrap samples. Values for survival and development time have

been back-transformed from units on the log-odds and log scales to units of probability and days to pupation, and are shown as changes from the mean or intercept values. For example, a negative (orange) survival coefficient of 0.005 means a one-half percent reduction in average probability of survival associated with variation in a particular compound. The fifteen compounds with the largest coefficients (by absolute value) and bootstrap intervals not overlapping zero are labelled by compound classes (see Fig. 2 for abbreviations) in each panel. Structural annotations are shown to the right for six compounds based on matches from the METLIN metabolomics database, as follows by compound number: 154 (unidentified sterol); 9 (unidentified alkaloid); 60 (soyasaponin A3); 40 (unidentified saponin); 46 (medicagenic acid 3-O-triglucoside); 45 (medinoside E). Those same compounds are identified in parentheses in the main panels next to the bars corresponding to their individual effects.

Fig. 4. Violin plots of compound-specific effects (coefficients from ridge regressions) summarized by chemical classes. Plots show median (black dot), interquartile range (box) and 95% confidence intervals (whiskers) surrounded by kernel density envelopes. Sample sizes for each category are shown above the top panel ("Other" includes 1 sugar, 2 pigments and 2 halogenated compounds). Categories are arranged from left to right based on the gradient of median positive to negative effects on survival. Coefficients for survival (a) and development time (b) have been back-transformed from the units of log-odds and log to probability and days to pupation, respectively, and shown as deviations from the mean or intercept value (as in Fig. 3). Note that negative effects for development time correspond to fewer days (more rapid development).

Table 1 Results from Bayesian regressions of module eigenvectors and covariates predicting caterpillar survival, development time and adult weight (as binomial, Poisson, and Gaussian regressions, respectively, with corresponding units in log-odds, log number of days, and milligrams).

	Survival coefficient (CI; prob.)	Development time coefficient (CI, prob.)	Weight coefficient (CI, prob.)
m2	0.37 (-1.87, 2.67; 0.63)	-0.07 (-0.26, 0.12; 0.78)	
m3	-2.33 (-3.94, -0.72; >0.99)		-2.33 (-4.87, 0.24; 0.96)
m6		-0.05 (-0.19, 0.09; 0.75)	
m9	-2.31 (-4.49, -0.15; 0.98)	0.064 (-0.11, 0.24, 0.78)	
m10	2.54 (0.81, 4.23; >0.99)		3.56 (1.22, 5.84; >0.99)
m11	2.01 (0.64, 3.42; >0.99)		
SLA	-0.31 (-0.55, -0.08; >0.99)		-0.42 (-0.78, -0.06; 0.99)
Protein	0.061 (-0.16, 0.28; 0.71)		0.08 (-0.26, 0.42; 0.68)
Tough.	0.036 (-0.17, 0.24; 0.36)	-0.002 (-0.02, 0.02; 0.57)	
Sex	/	0.06 (0.02, 0.10; >0.99)	1.10 (0.38, 1.83; >0.99)
mg	/	-0.03 (-0.05, -0.01; >0.99)	/
Days	/	/	1.10 (0.38, 1.83; >0.99)
Intercept	0.34 (0.14, 0.53; >0.99)	3.48 (3.45, 3.52; >0.99)	10.36 (9.81, 10.92; >0.99)
Validation	0.53	0.59	0.50

For each regression coefficient, numbers in parentheses are 95% credible intervals (the first two numbers) and the probability that the coefficient has the estimated sign (e.g., 0.63 for the m2 survival coefficient of 0.37 indicates a 63% probability that the m2 module has a positive effect on survival). Note that negative coefficients for development time indicate faster caterpillar development (fewer days) associated with variation in a particular compound. Modules (listed in the left column) are only shown if they were included in one of the three regressions following feature selection using lasso regression (see main text for additional details). Empty spaces in the Table appear if a particular module was selected through lasso regression for one or two analyses but not all three (m3, for example, was not selected by lasso regression for development time). Slash marks (/) indicate variables not considered for a particular analysis (e.g., sex, adult weight [mg] and development time [days] were not possible for the survival analysis because they are not observed on dead individuals). Values for "validation" shown in the last row are the correlation between observed and predicted values in cross-validation (Supporting Information Fig. S4).

Fig. 1. Variation among plants in caterpillar survival (a), development time (b) adult weight (c), three individual compounds (d-e) and two external plant traits, specific leaf area (g) and leaf toughness (h). The three example compounds shown here (out of the 163 assayed) were among the top five most influential compounds for survival, development time and adult weight: cpd. 9 is a peptide with a negative association with survival, cpd. 94 (another peptide) has a negative association with development time, and cpd. 160 is a phospholipid with a negative association with adult weight. Individual plants in all panels are organized from left to right by decreasing caterpillar survival in the top panel (a). Standard errors are shown for panels b, c, g and h. The units for d-e are compound relative abundance per dry weight of sample; the units for specific leaf area are cm^2/mg, and grams/newton for leaf toughness.

Fig. 2. Illustration of correlational structure among compounds: each node in the network is a compound, and compounds are linked by a line if they are correlated among individual plants at 0.5 or above (links among compounds in modules 12-14 represent weaker correlations, greater than 0.1; see main text for details). Two letter codes within nodes indicate compound classes, as explained in the legend. Colors of nodes correspond to membership in modules as determined by hierarchical cluster analysis; the color key to the 14 modules is shown in the lower left. Not shown are a small number of compounds with weak connections to all other compounds, including two compounds that were not included in any module (shown as module zero in Fig. 3).

Fig. 3. Effects of individual compounds on survival, development time and adult weight, as estimated by ridge regression (using binomial, Poisson and Gaussian models, respectively). The strength of effect for each compound is indicated by the horizontal extent of each bar, and compounds are grouped by modules (m1, m2, etc.); the order of compounds along the vertical axis is arbitrary within modules and fixed across columns. Orange colors indicate negative effects on survival, development and weight, while blue colors are positive effects (note that negative effects for development time correspond to fewer days, or more rapid development). The darker shades of orange and blue mark coefficients whose 95% confidence intervals did not overlap zero in 1,000 bootstrap samples. Values for survival and development time have been back-transformed from units on the log-odds and log scales to units of probability and days to pupation, and are shown as changes from the mean or intercept values. For example, a negative (orange) survival coefficient of 0.005 means a one-half percent reduction in average probability of survival associated with variation in a particular compound. The fifteen compounds with the largest coefficients (by absolute value) and bootstrap intervals not overlapping zero are labelled by compound classes (see Fig. 2 for abbreviations) in each panel. Structural annotations are shown to the right for six compounds based on matches from the METLIN metabolomics database, as follows by compound number: 154 (unidentified sterol); 9 (unidentified alkaloid); 60 (soyasaponin A3); 40 (unidentified saponin); 46 (medicagenic acid 3-O-triglucoside); 45 (medinoside E). Those same compounds are identified in parentheses in the main panels next to the bars corresponding to their individual effects.

Fig. 4. Violin plots of compound-specific effects (coefficients from ridge regressions) summarized by chemical classes. Plots show median (black dot), interquartile range (box) and 95% confidence intervals (whiskers) surrounded by kernel density envelopes. Sample sizes for each category are shown above the top panel ("Other" includes 1 sugar, 2 pigments and 2 halogenated compounds). Categories are arranged from left to right based on the gradient of median positive to negative effects on survival. Coefficients for survival (a) and development time (b) have been back-transformed from the units of log-odds and log to probability and days to pupation, respectively, and shown as deviations from the mean or intercept value (as in Fig. 3). Note that negative effects for development time correspond to fewer days (more rapid development).