Non-random sister chromatid segregation mediates rDNA copy number maintenance in

Drosophila

George J. Watase[1, 2] and Yukiko M. Yamashita[1, 2, *]

[1] Whitehead Institute for Biomedical Research, Department of Biology, Massachusetts Institute

of Technology, [2] Howard Hughes Medical Institute, 455 Main Street, Cambridge, MA 02142

*Corresponding author: yukikomy@wi.mit.edu

Abstract (125 words)

Although considered to be exact copies of each other, sister chromatids can segregate non-randomly in some cases. For example, sister chromatids of the X and Y chromosomes segregate non-randomly during asymmetric division of male germline stem cells (GSCs) in *D. melanogaster*. Here we identify that the ribosomal DNA (rDNA) loci, which are located on the X and Y chromosomes, and an rDNA-binding protein, Indra, are required for non-random sister chromatid segregation (NRSS). We provide the evidence that NRSS is a mechanism by which GSCs recover rDNA copy number, which occurs through unequal sister chromatid exchange, counteracting the spontaneous copy number loss that occurs during aging. Our study reveals an unexpected role for NRSS in maintaining germline immortality through maintenance of a vulnerable genomic element, rDNA.

One Sentence Summary (125 characters)

rDNA copy number maintenance by non-random sister chromatid segregation contributes to germline immortality in *Drosophila*

Sister chromatids, generated through the precise process of DNA replication, are considered identical. Nevertheless, it has been proposed that sister chromatids might carry distinct information or mutation loads, and their non-random segregation may underlie asymmetric cell division (*1-3*). However, the underlying mechanism remains elusive, preventing investigation of the physiological relevance of non-random sister chromatid segregation (NRSS). Using *D. melanogaster* male germline stem cells (GSCs) as a model system, where asymmetric stem cell division can be observed at single cell resolution, we previously showed that the X and Y chromosomes exhibit strikingly biased sister chromatid segregation (*4*). By using chromosome-orientation fluorescence *in situ* hybridization (CO-FISH) with chromosome-specific probes (Fig. 1A), (+)-strand templated vs. (-)-strand templated sister chromatids of each chromosome can be differentiated (Fig. 1A and B). If sister chromatids are equivalent, (+)- vs. (-)-strand templated sister chromatids would segregate to the GSC or GB (gonialblast, the differentiating daughter of a GSC) at random (50:50). Although we observed random sister

chromatid segregation for autosomes (chromosome 2 and 3), the X and Y chromosomes

segregated their sister chromatids non-randomly, with a specific strand segregating to GSC in

~80% of observed divisions (Fig. 1C, 'red strand')(*4*). This demonstrated that sister chromatids,

which supposedly carry the same genetic information, can be distinguished and segregated non-

randomly during asymmetric stem cell division.

To elucidate the underlying molecular mechanism of NRSS, we sought to identify the

chromosomal loci that mediate NRSS and found that ribosomal DNA (rDNA) is required. An

X chromosome without rDNA, *Df(1)bb^{158}* (*bb^{158}* hereafter), as well as a Y chromosome

without rDNA (*Ybb^-*), exhibited randomized sister chromatid segregation (Fig. 1C, table S1).

Importantly, the intact Y chromosome in the *bb^{158}* strain, as well as the intact X chromosome

in the *Ybb^-* strain, still exhibited NRSS (Fig. 1C, table S1), suggesting that the rDNA loci

likely act as cis-elements to mediate NRSS. A chromosome 2 containing an rDNA locus

translocated from the Y chromosome also exhibited NRSS ('2^Y with rDNA' in T(Y;2)A77

translocation, Fig. 1D, table S2), suggesting that rDNA is sufficient to induce NRSS. As a

critical control, a chromosome 2 carrying a similar translocation from the Y chromosome that

does not include the rDNA did not exhibit NRSS ('2^Y without rDNA' in T(Y;2)P8

translocation, Fig. 1D, table S2). This is the first demonstration that a specific region of a

chromosome is responsible for NRSS, opposing the widely-held speculation that NRSS

depends on chromosome-wide information such as epigenetic information and replication-

induced mutations (*5*).

To understand how rDNA mediates NRSS, we isolated rDNA binding proteins from the

GSC extract. Each rDNA locus consists of 150-225 repeated rDNA units in order to support the

high demand of ribosome biogenesis (*6*). Each rDNA unit contains the 18S, 5.8S/2S, 28S rRNA

genes and three spacer sequences [the external transcribed spacer (ETS), internal transcribed

spacer (ITS) and intergenic spacer (IGS)] (Fig. 2A). Interestingly, the Y chromosome of *D.*

simulans, a species closely related to *D. melanogaster,* has IGS repeats but no rRNA genes, ETS

or ITS (*7*), yet exhibited NRSS (fig. S1, table S1). We hypothesized that IGS may be responsible

for NRSS. Thus, we isolated IGS-binding proteins by mass spectrometry followed by a

secondary screen based on subcellular localization (Fig. 2B, table S3). In this study, we focus on

a previously-uncharacterized zinc finger protein, CG2199, which we named Indra after the

Hindu god who lost immortality due to a curse from Durvasa. Using a specific anti-Indra

antibody (fig. S2A) and an Indra-GFP line, we found that Indra localizes to the nucleolus (the

site of rDNA transcription) in interphase (Fig. 2C, fig. S3A) and rDNA loci during metaphase

(Fig. 2D, fig. S3B). ChIP-qPCR further demonstrated that Indra preferentially binds to IGS (Fig.

2E). Strikingly, RNAi-mediated knockdown of *indra* in the germline (fig. S2A) compromised

NRSS for both the X and Y chromosomes (Fig. 2F, table S4). Taken together, these results show

that IGS and its binding protein Indra mediate NRSS.

We found that *indra* is required for rDNA copy number maintenance. RNAi-mediated

depletion of *indra* (*nos-gal4>UAS-indra$^{TRiP.HMJ30228}$*) resulted in drastically fewer progeny

compared to control (Fig. 3A, P_0). Some of the offspring from *indra$^{TriP.HMJ30228}$* males exhibited a

bobbed phenotype, a hallmark of rDNA copy number insufficiency characterized by abnormal

cuticle patterns on the abdomen (Fig. 3B; (*8*)). The frequency of *bobbed* flies increased when the

Y chromosome from *indra$^{TriP.HMJ30228}$* fathers was placed in the background of *bb^{158}*, the X

chromosome that lacks rDNA (Fig. 3B). Quantitative droplet digital PCR (ddPCR) confirmed

that rDNA copy number was reduced in *indra$^{TriP.HMJ30228}$* animals (Fig. 3C, P_0). Depletion of

indra over successive generations resulted in a progressive loss of fecundity (Fig. 3A, F_1-F_2)

associated with a reduction in rDNA copy number (Fig. 3C, F_1-F_2). Moreover, *indra* is required

for 'rDNA magnification', a phenomenon by which an X chromosome with insufficient rDNA

copy number is induced to recover copy number, when the fly lacks rDNA on the Y

chromosome (*Ybb$^-$*) (Fig. 3, D and E, fig. S4; (*9*)). These results suggest that *indra* is required for

rDNA copy number maintenance over generations.

Although the repetitiveness of rDNA loci is critical to support ribosome biogenesis, it

also makes rDNA loci susceptible to intrachromatid recombination, which leads to spontaneous

copy number loss (Fig. 4A). To maintain the integrity of rDNA loci, copy number loss must be

counteracted by copy number recovery. In yeast, rDNA copy number recovery is mediated by

unequal sister chromatid recombination (*10*). Similarly, rDNA magnification, which we

postulated to mediate rDNA copy number recovery in the *Drosophila* male germline (*11*), is

proposed to utilize unequal sister chromatid exchange (USCE) (*9*). USCE allows for copy

number recovery on one of the sister chromatids at the expense of the other (Fig. 4B; (9)),

generating asymmetry between two sister chromatids (Fig. 4B). We hypothesized that this

asymmetry may underlie NRSS.

Strikingly, asymmetry in rDNA amount was detected during anaphase in GSCs under

'magnifying conditions'(bb^{z9}/Ybb^-) with the GSCs preferentially inheriting the stronger signal

(Fig. 4, D and E). As an important control, asymmetry in rDNA amount was not observed in flies

with sufficient rDNA copy number (Fig. 4, C and E). Interestingly, there were rare cases where

rDNA asymmetry was created, but the stronger signal was inherited by the GBs (Fig. 4E, top

panel). Plotting the ratio of stronger over weaker rDNA signal (Fig. 4E, bottom panel) suggests

that the magnifying condition strongly induced USCE. rDNA copy number and segregation

asymmetries were absent under magnifying conditions following depletion of *indra* (Fig. 4E),

suggesting that *indra* may be involved not only in NRSS but also in generating copy number

asymmetry through USCE.

These results are consistent with a model in which rDNA magnification is mediated by

USCE, and the sister chromatid with increased rDNA copy number is selectively retained by

GSCs by NRSS. We further tested this idea by using additional CO-FISH probes. The probe for

the 359-bp repeat, located proximal to the rDNA (Fig. 4F), was used to detect NRSS in the

experiments described above. $(TAGA)_n$, which is located distal to the rDNA, also exhibited

NRSS in control (non-magnifying) conditions (Fig. 4G, table S5). Strikingly, under magnifying

conditions, 359-bp maintained NRSS but $(TAGA)_n$ exhibited random segregation, suggesting

that sister chromatid exchange occurred between 359-bp and $(TAGA)_n$, most likely within the

rDNA locus (Fig. 4F). Taken together, these data suggest that GSCs undergo rDNA

magnification through USCE followed by NRSS (see Supplementary text).

Our study reveals the unexpected molecular mechanisms and biological significance of

NRSS. We propose that NRSS is a key process to recover and maintain inherently unstable

rDNA copy numbers such that the integrity of the germline genome is upheld over generations,

supporting germline immortality. Future work is required to understand how rDNA copy number

134 differences between sister chromatids are recognized and faithfully segregated to the GSCs to

135 achieve rDNA copy number recovery.

137 **Competing interests**

138 The authors declare no competing interests.

140 **Acknowledgements**

141 We thank the Bloomington, Kyoto, and National Drosophila Species Stock Centers, the Vienna

142 Drosophila Resource Center and the Developmental Studies Hybridoma Bank for reagents. We

143 thank the Yamashita lab members, Drs. Swathi Yadlapalli and Sue Hammoud, and the Life

144 Science Editors for comments on the manuscript. We thank MS Bioworks for mass-spectrometry

145 analysis and the laboratories of Drs. Stephen Weiss and Jiandie Lin for sharing equipment. This

146 research was supported by Howard Hughes Medical Institute.

148 **Author contributions**

149 G.W. and Y. Y. designed and conducted experiments, interpreted results and wrote and edited

150 the manuscript.

152 **References**

153 1. P. M. Lansdorp, Immortal strands? Give me a break. *Cell* **129**, 1244-1247 (2007).

154 2. T. A. Rando, The immortal strand hypothesis: segregation and reconstruction. *Cell* **129**,
155   1239-1243 (2007).

156 3. S. Tajbakhsh, C. Gonzalez, Biased segregation of DNA and centrosomes: moving
157   together or drifting apart? *Nat Rev Mol Cell Biol* **10**, 804-810 (2009).

158 4. S. Yadlapalli, Y. M. Yamashita, Chromosome-specific nonrandom sister chromatid
159   segregation during stem-cell division. *Nature* **498**, 251-254 (2013).

160 5. S. Tajbakhsh, Stem cell identity and template DNA strand segregation. *Curr Opin Cell*
161   *Biol* **20**, 716-722 (2008).

162 6. R. S. Hawley, C. H. Marcus, Recombinational controls of rDNA redundancy in
163   Drosophila. *Annu Rev Genet* **23**, 87-120 (1989).

164 7. A. R. Lohe, P. A. Roberts, An unusual Y chromosome of Drosophila simulans carrying
165   amplified rDNA spacer without rRNA genes. *Genetics* **125**, 399-406 (1990).

166 8. F. M. Ritossa, K. C. Atwood, S. Spiegelman, A molecular explanation of the bobbed
167   mutants of Drosophila as partial deficiencies of "ribosomal" DNA. *Genetics* **54**, 819-834
168   (1966).

169 9. K. D. Tartof, Unequal mitotic sister chromatin exchange as the mechanism of ribosomal
170   RNA gene magnification. *Proc Natl Acad Sci U S A* **71**, 1272-1276 (1974).

171   10.   T. Kobayashi, Ribosomal RNA gene repeats, their stability and cellular senescence. *Proc*
172   *Jpn Acad Ser B Phys Biol Sci* **90**, 119-129 (2014).
173   11.   K. L. Lu, J. O. Nelson, G. J. Watase, N. Warsinger-Pepe, Y. M. Yamashita,
174   Transgenerational dynamics of rDNA copy number in Drosophila male germline stem
175   cells. *Elife* **7**, (2018).
176

Figures

Fig. 1. rDNA loci are required for NRSS of the X and Y chromosomes in *D. melanogaster*

(A) Chromosome-orientation *in situ* hybridization (CO-FISH) to assess NRSS. Plus (+) vs.
minus (-) templated strands are indicated by red and blue lines, newly-synthesized strands
by black dotted lines. Following removal of BrdU-containing, newly-synthesized strands,
strand-specific probes were applied to distinguish red vs. blue templated strands.

(B) Representative images of Y chromosome CO-FISH results where a GSC inherits the 'red'
strand $((AATAC)_n)$, whereas a GB inherits the 'blue' strand $((GTATT)_n)$. The hub, the
stem cell niche to which GSCs are attached, is indicated by an asterisk, GSC-GB pairs
are outlined by dotted lines and the CO-FISH signals by arrowheads. Vasa: germ cells.
Arm: hub. Add: the connection between GSC and GB. Bar: 10μm.

(C, D) Schematics of *D. melanogaster* X and Y chromosomes (C) and Y-2 translocation
chromosomes (D). Summary of sister chromatid segregation patterns in indicated genotypes
is shown (see table S1 and S2). Data shown as mean ± s.d. from three independent
experiments. n, number of GSC-GB pairs scored. *P*-values: Fisher's exact test.

Fig. 2. Indra is a novel zinc-finger protein that binds to rDNA and mediates NRSS.

(A) Schematic of rDNA loci in *D. melanogaster*.

(B) Experimental scheme to isolate IGS-binding proteins (see Materials and Methods).

(C) Localization of Indra at the apical tip of the testis. The hub is indicated by an asterisk. An

enlarged image of a GSC is shown in the inset. Fibrillarin: nucleolus. Vasa: germ cells.

Bar: 10µm.

(D) Localization of Indra on a metaphase chromosome spread from germ cells. The X rDNA

locus (arrow) and Y rDNA locus (arrowhead) can be identified by their relative location

to the centromere (Cid). An additional example of the Y chromosome is shown in the

inset. Bar: 5 µm.

(E) Indra ChIP-qPCR showing enrichment of Indra on rDNA/IGS. The 5S rDNA sequence

on chromosome 2 outside of the rDNA loci was used as a negative control. Mean and s.d.

from three technical replicates of qPCR are shown. Similar results were obtained from

two biological replicates.

(F) Summary of sister chromatid segregation patterns upon knockdown of *indra* (see table
S4). Data shown as mean ± s.d. from three independent experiments. n, number of GSC-
GB pairs scored. *P*-values: Fisher's exact test.
212

Fig. 3. *indra* is required for rDNA copy number maintenance

(A) Fertility of control and *nos-gal4>UAS-indra$^{TRiP.HMJ30228}$* males across generations (P$_0$-F$_2$).

Total number of progeny from 0-5 day old males in each generation and 5-10 day old

males in P$_0$ were scored. Data shown as median with 95% confidence interval and

individual data points. n, number of individual crosses scored. *P*-value: two-tailed Mann-

Whitney test.

(B) Frequency of *bobbed* animals in progeny of 0-5 day-old control and *nos-gal4>UAS-*

indra$^{TRiP.HMJ30228}$ males. Mean and s.d. from three independent experiments with

individual data points are shown. n, total number of progeny scored. *P*-values: two-tailed

chi-squared test. Examples of normal and '*bobbed*' cuticle phenotypes are shown on the

left.

(C) 28S rRNA gene copy number in the testes from 0-5 day old control and *nos-gal4>UAS-*

indra$^{TRiP.HMJ30228}$ males in successive generations (P$_0$-F$_2$) assessed by ddPCR. Data shown

as median with 95% confidence interval and individual data points. n, number of

individual crosses scored. *P*-value: two-tailed Mann-Whitney test.

(D) Schematic diagram of rDNA magnification assay (see fig. S4 for the details).

Magnification was detected by normal cuticle phenotype in the offspring.

(E) Frequency of rDNA magnification in the indicated genotypes/conditions. Data shown as

mean and s.d. from four (*nos-gal4>UAS-indra*^{GD9748}, *UAS-Dcr-2*) or three (the rest)

independent experiments with individual data points. n, total number of progeny scored.

P-values: Fisher's exact test.

Fig. 4. rDNA loci undergoe sister chromatid exchange and segregate asymmetrically in GSCs during rDNA magnification

(A) Diagram of spontaneous rDNA copy number loss by intrachromatid recombination.

(B) A proposed model for rDNA copy number recovery by unequal sister chromatid exchange.

(C, D) DNA-FISH for IGS during anaphase in GSCs under normal (*yw*) (C) and magnifying (*bb^{z9}/Ybb$^-$*) (D) conditions. The hub is indicated by an asterisk. An enlarged image of the IGS signal from GSC and GB sides is shown in the inset. Bar: 5 μm.

(E) Quantification of IGS signal intensity presented as GSC side/GB side (top panel) or stronger/weaker (bottom panel) during anaphase in GSCs in control and *nos-gal4>UAS-*

indra^{GD9748}, *UAS-Dcr-2* males. The median and individual data point are shown. 95%

confidence interval is also shown in the bottom panel. n, number of anaphase cells

scored. *P*-value: two-tailed Mann-Whitney test. Note that due to rare cases where the GB

side exhibited stronger IGS signal than GSC side, the data did not reach statistical

significance between control and *indra*^{GD9748} under magnifying conditions in the top

panel.

(F) Diagram of the X chromosome showing the location of the rDNA and the 359-bp and

(TAGA)$_n$ repeats. Sister chromatid exchange at rDNA loci would flip (TAGA)$_n$

segregation pattern relative to 359-bp.

(G) Summary of sister chromatid segregation patterns assessed by 359-bp and (TAGA)$_n$

probes in the indicated genotypes/conditions (see table S5). Data shown as mean ± s.d.

from three independent experiments. n, number of GSC-GB pairs scored. *P*-values:

Fisher's exact test.

Supplementary Materials

Materials and Methods

Supplementary Text

Figs. S1 to S5

Tables S1 to S7

References (1-18)

Materials and method

<u>Fly husbandry and strains</u>

All fly stocks were raised on standard Bloomington medium at 25°C containing 0.15% of tegosept

as anti-fungal (no propionic acid was added). The following fly stocks were used: *Df(1)bb^158,*

y^1/Dp(1;Y)y^+/C(1)*; ca^1 awd^K (BDSC3143), *FM6/C(1)DX, y* f^1/Y* (BDSC784), *UAS-*

indra^{TRiP.HMJ30228} (BDSC63661), *UAS-Dcr-2* (BDSC24650), *indra-GFP* (BDSC67660;

http://flybase.org/reports/FBti0186577) were obtained from the Bloomington Drosophila Stock

Center. *y^1 eq^1/Df(YS)bb^-* (DGRC101260), *T(Y;2)A77, B^S, y^+/SM1; C(1)RM, y^1/C(1;Y)1, y^1*

(DGRC130079), *T(Y;2)P8, B^S, y^+/SM1; C(1)RM, y^1/C(1;Y)1, y^1* (DGRC130170) were obtained

from the Kyoto Stock Center. *D. simulans W^{501}*(DSSC14021-0251.195) was obtained from the

National Drosophila Species Stock Center. *UAS-indra^{GD9748}* (v20839) was obtained from the

Vienna Drosophila Resource Center. *nos-gal4* (*1*), *UAS-Upd* (*2*), *tub-gal80^{ts}* (*3*), *nos-gal4* without

VP16 (*4*) have been previously described.

To examine the sister chromatid segregation patterns of the 2^Y and Y^2 chromosomes,

T(Y;2)A77/SM1; C(1)RM/O or *T(Y;2)P8/SM1; C(1)RM/O* females were crossed to *yw* males and

the resulting *T(Y;2)A77/+; X/O* and *T(Y;2)P8/+; X/O* male flies were examined. The details of the

translocation are shown in Fig. 1D.

Two independent RNAi lines, *UAS-indra^{TRiP.HMJ30228}* and *UAS-indra^{GD9748}*, were used to

knockdown *indra* specifically in early germ cells using *nos-gal4* as the driver. *UAS-indra^{GD9748}*

was combined with *UAS-Dcr-2* to increase RNAi efficiency. The knockdown efficiency of these

RNAi lines was validated by immunostaining using an anti-Indra antibody (fig. S2A). Since *nos-*

gal4>UAS-indra^{TRiP.HMJ30228} results in severe germ cell loss due to high RNAi efficiency (Fig. 3A,

fig. S2A), a temperature-sensitive GAL4 inhibition system (*tub-gal80^{ts}*; *nos-gal4ΔVP16>UAS-*

indra^{TRiP.HMJ30228}) was used as necessary (e.g. Fig. 2F). Upon shifting from the permissive

temperature (18˚C) to the non-permissive temperature (29˚C), GSCs were lost gradually over 2-4

days (fig. S2, B and C), and the CO-FISH assay (Fig. 2F) was conducted 3 days after temperature

shift. In assays that required a sustained germline (e.g. magnification assays, Fig. 3E and Fig. 4E),

we used *nos-gal4>UAS-indra^{GD9748}, UAS-Dcr-2*.

<u>Immunofluorescence staining and confocal microscopy</u>

Drosophila adult testes were dissected in phosphate-buffered saline (PBS), transferred to 4%

formaldehyde in PBS and fixed for 30 min. The testes were then washed in PBST (PBS containing

0.1% Triton X-100) for at least 30 min, followed by incubation with primary antibody in 3%

bovine serum albumin (BSA) in PBST at 4°C overnight. Samples were washed for 60 min (3 x 20

min washes) in PBST, incubated with secondary antibody in 3% BSA in PBST at 4°C overnight,

washed as above, and mounted in VECTASHIELD with 4',6-diamidino-2-phenylindole (DAPI;

Vector Labs, Burlingame, CA). To examine Indra localization on mitotic chromosome spreads,

Drosophila 3rd instar larval testes were dissected in PBS, transferred to 0.5% sodium citrate and

incubated for 10 min, fixed in 4% formaldehyde in PBS for 4 min, then squashed between the

cover slip and slide glass. The sample was frozen in liquid nitrogen, the cover slip was removed,

and immediately washed in PBS, followed by immunofluorescence staining as described above,

except that the incubation was performed on the slide glass in a humid chamber with the sample

covered with a small piece of parafilm.

The primary antibodies used were as follows: rabbit anti-Vasa (1:200; d-26; Santa Cruz

Biotechnology, Santa Cruz, CA), mouse anti-Adducin-like [1:20; 1B1; developed by H.D. Lipshitz,

obtained from Developmental Studies Hybridoma Bank (DSHB)](*5*), mouse anti-Armadillo

(1:100; N2 7A1; developed by E. Wieschaus, obtained from DSHB)(*6*), rat-anti Vasa (1:20;

developed by A.C. Spradling and D. Williams, obtained from DSHB), mouse anti-Fibrillarin

(1:200; 38F3; Abcam), chicken anti-Cid (1:500)(*7*). The anti-Indra antibody was generated by

injecting a peptide (RKITDVLETITHRSIPSSLPIKIC) into guinea pig (Covance, Denver, PA) and

used at a dilution of 1:500. Specificity of the antibody was validated by the lack of signal in

indra^{RNAi} testis (fig. S2A). Alexa Fluor-conjugated secondary antibodies (Life Technologies) were

used at a dilution of 1:200. Images were taken on a Leica TCS SP8 confocal microscope with a

63x oil immersion objective (NA = 1.4) and processed using Adobe Photoshop software.

For DNA FISH combined with immunofluorescent staining, whole mount *Drosophila* testes were

prepared as described above, and the immunofluorescence staining protocol was carried out first.

Upon completion of the wash post incubation with the secondary antibody, samples were fixed

with 4% formaldehyde for 10 min and washed in PBST for 30 min. Fixed samples were incubated

with 2 mg/ml RNase A solution at 37°C for 10 min, then washed with PBST. Samples were washed

in 2x SSC with increasing formamide concentrations (20% and 50%) for 10 min each.

Hybridization buffer (50% formamide, 10% dextran sulfate, 2x SSC, 1 mM EDTA, 1 mM probe)

was added to washed samples. Samples were denatured at 91°C for 2 min, then incubated overnight

at 37°C. Following the hybridization, testes were washed once in 50% formamide/2x SSC, once

in 20% formamide/2x SSC, and 3 times in 2x SSC. All reagents contained 1 mM EDTA except

for the washes prior to RNase A treatment. Fluorescence quantification was done on merged z-

stacks using Image J 'Sum of pixel intensity (RawIntDen)' to compare signal intensity between

sister chromatids. To avoid the effect of signal intensity changes along the Z plane, we scored

anaphase GSCs only when two IGS FISH signals were found within the same Z plane for Fig. 4,

C-E.

<u>Chromosome orientation fluorescence *in situ* hybridization (CO-FISH)</u>

CO-FISH in whole mount *Drosophila* testes was performed as previously described (*8*). Briefly,

young adult flies (day 1-3) were fed with 5-bromodeoxyuridine (BrdU)-containing food (950 µl

of 100% apple juice, 7 mg of agar, and 50 µl of 100 mg/ml BrdU solution in a 1:1 mixture of

acetone and DMSO) for 12 hours. After the feeding period, flies were transferred to regular fly

food for 13.5 hours. Because the average GSC cell cycle length is ~12 hour, most GSCs undergo

a single S phase in the presence of BrdU followed by mitosis during this feeding procedure. GSCs

that have undergone more or less than one S phase or mitosis were excluded from our analysis by

limiting the scoring to GSC-GB pairs that have complementary CO-FISH signals in the GSC and

GB (red signal in one cell, blue signal in the other). Note that GSC and GB stay connected by the

fusome until mid-S phase, which allowed identification of the GSC-GB sister pairs. Testes were

dissected, fixed and immunostained as described above. Then, testes were fixed for 10 min with

4% formaldehyde in PBS, followed by 3 washes with PBST. Following the washes, the testes were

rinsed once with PBST and treated with RNase A (Roche; 2 mg/ml in PBS) for 10 min at 37°C,

washed with PBST for 5 min, and stained with 100 µl of 2 µg/ml Hoechst 33258 (Invitrogen) in

2x SSC for 15 min at room temperature. The testes were then rinsed 3 times with 2x SSC,

transferred to a tray, and irradiated with ultraviolet light in a CL-1000 Ultraviolet Crosslinker

(UVP; wavelength: 365 nm; calculated dose: 5400 J/m^2). Nicked BrdU positive strands were

digested with exonuclease III (New England BioLabs) at 3 U/µl in 1x NEB1 buffer or 1x NEB

cutsmart buffer for 10 min at 37°C. The testes were washed once with PBST for 5 min and then

fixed with 4% formaldehyde in PBS for 2 min. Subsequently, the fixed testes were washed 3 times

with PBST. Testes were incubated sequentially for a minimum of 10 min each in 20%

formamide/2x SSC and 50% formamide/2x SSC. The testes were incubated with hybridization

buffer (50% formamide, 2x SSC, 10% dextran sulfate) containing 1 µM of each probe for 16 hours

at 37°C. Following hybridization, testes were washed once in 50% formamide/2x SSC, once in

20% formamide/2x SSC and 3 times in 2x SSC. Images were taken on either a Leica TCS SP5 or

STELLARIS 8 confocal microscope with a 63x oil immersion objective (NA = 1.4) and processed

using Adobe Photoshop software. For CO-FISH in GSCs from *tub-gal80^{ts}*; *nos>indra$^{TRiP.HMJ30228}$*,

the BrdU pulse was conducted 3 days after temperature shift (fig. S2C). BrdU was fed at 29°C for

9 hours, followed by an 11-hour chase at 29°C in which the flies were fed regular fly food. The

probes are described in table S6. All reagents contained 1 mM EDTA except for the washes

immediately preceding an enzymatic reaction (RNase A and exonuclease III).

<u>IGS DNA pull down and mass-spectrometry</u>

200 pairs of *upd*-expressing testes (*nos-gal4>UAS-upd*) were dissected in Schneider's *Drosophila*

Medium (Gibco) and washed 3 times with ice-cold PBS. *upd* expression causes overproliferation

of GSC-like cells. The testes were homogenized in lysis buffer [20 mM Tris-HCl pH8.0, 1 mM

EDTA, 10% Glycerol, 0.2% NP-40, 1 mM DTT, 1x solution of PhoSTOP cocktail (Roche), 1x

solution of c0mplete EDTA-free protease inhibitor cocktail (Roche)], and the homogenate was

incubated on ice for 20 min. Following this incubation, the lysate was centrifuged at 3,000 rpm for

10 min at 4°C, and the supernatant was saved as whole cell extract. The pellet, which contains the

nuclear fraction, was resuspended in lysis buffer containing 100 mM NaCl and incubated on ice

for 1 hour. During incubation, the sample was vortexed at highest setting for 15 sec every 10 min.

The nuclear fraction was isolated by centrifugation at 14,000 rpm for 30 min at 4°C and mixed

with the whole cell extract prepared above. Protein concentration was measured by absorbance at

562 nm using Pierce^TM BCA Protein Assay Kit (Thermo Scientific) to adjust the concentration

between samples.

The 240-bp IGS sequence (repeated 4 times, 4xIGS), or the *Kr* gene promoter sequence (control),

was cloned into pBluescript SK⁻. Biotin end-labeling at the 5' of one strand of the 4xIGS or *Kr*

gene promoter was performed by PCR using a T7 primer with Biotin-TEG and a T3 primer.

Biotinylated 4xIGS and *Kr* gene promoter DNA were then purified by QIAGEN's PCR

purification kit. 2 µg of each biotinylated DNA was immobilized to 100 µl of streptavidin-bound

M-280 Dynabeads^TM (invirogen). The beads were washed 3 times with 1x Binding and Washing

buffer (5 mM Tris-HCl pH8.0, 0.5 mM EDTA, 500 mM NaCl) and then blocked with 0.5% BSA

in TGEDN buffer (120 mM Tris-HCl pH8.0, 1 mM EDTA, 100 mM NaCl, 1 mM DTT, 0.1%

Triton X-100, 10% Glycerol). 20% volume of each of the biotinylated DNA-conjugated

Dynabeads^TM was incubated with 20 µg of herring sperm DNA (Sigma-Aldrich) and the cell

extract prepared above (containing 500 µg of protein at 3.8-4.8 µg/µl concentration, matched

between control vs. IGS beads). After incubating for 2 hours at 4°C, the beads were washed 5

times with TGEDN buffer. The proteins bound to either the 4xIGS or *Kr* gene promoter DNA were

eluted in LDS sample loading buffer (1.5x) at 100°C for 15 min. 50% volume of each DNA bound

proteins was separated on a 10% Bis-Tris Novex mini-gel (invitrogen) using the MES buffer

system. The gel was stained with coomassie and excised into ten equally sized segments. These

segments were analyzed by LC/MS/MS (MS Bioworks, Ann Arbor, MI). The gel digests were

analyzed by nano LC/MS/MS with a Waters NanoAcquity HPLC system interfaced to a Thermo

Fisher Q Exactive. Peptides were loaded on a trapping column and eluted over a 75 µm analytical

column at 350 nL/min; both columns were packed with Luna C18 resin (phenomenex). The mass

spectrometer was operated in data-dependent mode, with MS and MS/MS performed in the

Orbitrap at 70,000 FWHM resolution and 17,500 FWHM resolution, respectively. The fifteen most

abundant ions were selected for MS/MS.

ChIP-qPCR

200 pairs of *upd*-expressing testes (*nos-gal4>UAS-upd*) were dissected in ice-cold PBS containing

protease inhibitor [1x solution of c0mplete protease inhibitor cocktail (Roche) and 1 mM PMSF].

The testes were crosslinked by incubating with 1% formaldehyde for 15 min at 37°C and rinsed

twice in ice-cold PBS containing protease inhibitor to stop the crosslink reaction. The testes were

homogenized in 200 µl of ice-cold ChIP Sonication Buffer [1% triton X-100, 0.1% sodium

deoxycholate, 50 mM Tris-HCl (pH 8.0), 150 mM NaCl, 5 mM EDTA], and the homogenate was

incubated on ice for 15 min. Following the incubation, the homogenate was aliquoted into 0.5 ml

PCR tubes, placed in a Biorupter® Plus sonication system (DIAGENODE) and sonicated in 4°C

water bath for 10 cycles of 30 sec 'ON' and 30 sec 'OFF' at 'HIGH' setting. The sonicated lysate

was centrifuged at 14,000 rpm for 10 min at 4°C to pellet cell debris. The volume of supernatant

was brought up to 1 ml with ChIP sonication buffer, and 40 µl of Dynabeads^{TM} Protein A

(invitrogen) was added to the supernatant. After a 1-hour preabsorption with Dynabeads^{TM} Protein

A at 4°C, 30 µl of supernatant (3%) was kept as 'input'. The rest was split into two and incubated

overnight with 10 µl of anti-Indra antibody (1:10 dilution from the original serum; generated as

described above) or 10 µl of pre-immune guinea pig serum (1:10 dilution from the original serum),

respectively. After incubating for 16 hours, 40 µl Dynabeads^{TM} Protein A was added to each

reaction and incubated for an additional 4 hours at 4°C with rotation. The beads were then washed

for 5 min at 4°C with 1 ml of the following buffers: 2 washes with ChIP sonication buffer, followed

by 3 washes with High Salt Wash buffer [1% triton X-100, 0.1% sodium deoxycholate, 50 mM

Tris-HCl (pH 8.0), 500 mM NaCl, 5 mM EDTA]; 2 washes with LiCl Immune Complex Wash

buffer [250 mM LiCl, 0.5% NP-40, 0.5% deoxycholate, 1 mM EDTA, 10 mM Tris-HCl (pH8.0)];

1 wash with TE buffer [10 mM Tris-HCl (pH8.0), 1 mM EDTA). For elution, each ChIP sample

was incubated with 250 µl of Elution Buffer (1% SDS, 100 mM NaHCO$_3$) for 30 min at 65°C,

vortexing gently every 10 min. After repeating the elution process once more, the supernatants

were combined. 500 µl of elution buffer was added to the 'input' sample. 20 µl of 5 M NaCl and

10 µl of RNase A [Roche; 2 mg/ml in 10 mM Tris-HCl (pH7.5) and 15 mM NaCl] were added to

each sample and incubated for overnight at 65°C. After incubating for 16 hours, 2 µl of Proteinase

K (New England BioLabs), 10 µl of 500 mM EDTA, and 20 µl of 1 M Tris-HCl (pH8.0) were

added to each sample and incubated at 45°C for 2 hours. The precipitated DNA was purified using

QIAGEN's PCR purification kit. Real-Time PCR was conducted to quantify precipitated DNA

using the Standard Curve method. Power SYBR Green PCR Master Mix (appliedbiosystems) was

used as the PCR reaction buffer. The QuantStudio^{TM} 6 Flex System (appliedbiosystems) was used

for Real-Time PCR reaction and analyzing the data. Primers used for Real-Time PCR are listed in
table S7.

<u>Fertility assay</u>

Newly eclosed single males (control (*nos-gal4*) or *nos-gal4>UAS-indra*^{TRiP.HMJ30228}) were
individually crossed to three *yw* virgin females. After 5 days, each male was transferred to a new
vial with three new virgin females. The number of adult flies that eclosed in each vial was scored.
nos-gal4>UAS-indra^{TRiP.HMJ30228} females are completely sterile. Therefore, to examine the fertility
of *nos>indra*^{TRiP.HMJ30228} across generations, newly eclosed *nos-gal4>UAS-indra*^{TRiP.HMJ30228}
males for each generation were crossed to *nos-gal4* females to deplete *indra* in germline in the
subsequent generation. Then, newly eclosed single males (control (*nos-gal4*) or *nos-gal4>UAS-*
indra^{TRiP.HMJ30228}) at each generation were individually crossed to three *yw* virgin females and the
number of adult flies that eclosed in each vial was scored.

<u>Droplet Digital PCR (ddPCR)</u>

20 pairs of testes/sample were dissected from 0-3 day-old control (*nos-gal4*) or *nos-gal4>UAS-*
indra^{TRiP.HMJ30228} males. Genomic DNA isolation was performed as previously described (*9*).
Briefly, the testes were homogenized in 200 µl of buffer A (100 mM Tris-HCl pH8.0, 100 mM
EDTA pH8.0, 100 mM NaCl, 0.5% SDS), and then an additional 200 µl of buffer A were added
to the homogenate. The homogenate was incubated at 65°C for 30 min. Then 800 µl of LiCl/KAc
(2.5:1 mixture of 6 M LiCl and 5 M KAc) was added to the homogenate, and the sample was left
on ice for 15 min. Subsequently, the sample was centrifuged at 14,000 rpm for 15 min, and 1 ml
of supernatant was transferred to a new tube. The supernatant was mixed with 600 µl of
isopropanol and centrifuged at 14,000 rpm for 15 min. The pellet (containing genomic DNA) was
washed once in 1 ml of 70% ethanol, air dried for 30min and dissolved in 35 µl of TE buffer. The
quality and concentration of genomic DNA were measured on a NANODROP ONE (Thermo
Scientific).

30 ng of genomic DNA was used per 20 µL ddPCR reaction for control gene reactions (RpL and
Upf1), and 0.3 ng of genomic DNA was used per 20 µL ddPCR reaction for 28S rRNA gene
reactions. The primers and probes are listed in table S7. ddPCR reactions were carried out

according to the manufacturer's protocol (Bio-Rad). In short, master mixes containing ddPCR

Supermix for Probes (No dUTP) (Bio-Rad), genomic DNA, primer/probe mixes, and *Hind*III-HF

restriction enzyme (New England Biolabs) for 28S rRNA gene reactions (no restriction enzyme is

needed for the control gene reactions) were prepared in 0.2 mL PCR tubes, and incubated at room

temperature for 15 min to allow for restriction enzyme digestion. ddPCR droplets were generated

from samples using a QX200 Droplet Generator (Bio-Rad), and droplets then underwent complete

PCR cycling on a C100 deep-well thermocycler (Bio-Rad). Droplet fluorescence was read using a

QX200 Droplet Reader (Bio-Rad). Sample copy number was determined using Quantasoft

software (Bio-Rad). rDNA copy number per genome was determined by 28S sample copy number

multiplied by 100 (due to the 100x dilution of genomic DNA in the 28S reaction compared to

control reaction) divided by control gene copy number multiplied by the expected number of

control gene copies per genome (2 for RpL samples; 1 for Upf1 samples). The 28S rRNA gene

copy number values normalized by each control were then averaged to determine 28S copy number

for each sample.

<u>Magnification assay</u>

The experimental design to assay rDNA magnification is shown in fig. S4. The bb^{z9} allele carries

an insufficient rDNA copy number on the X chromosome (*10*), which exhibits a '*bobbed*' cuticle

phenotype when combined with the bb^{158} allele (no rDNA on X chromosome) in females. To

induce magnification, the bb^{z9} allele was combined with a Y chromosome lacking rDNA

(bb^{z9}/Ybb^-)('magnifying condition'). These bb^{z9}/Ybb^- males were crossed to $bb^{158}/FM6$ females,

and the resulting bb^{z9}/bb^{158} females were examined for the *bobbed* cuticle phenotype. If

magnification occurred, a magnified allele ($bb^{z9\text{-}mag}$) combined with bb^{158} would produce a wild

type cuticle, whereas a non-magnified allele combined with bb^{158} would show a *bobbed* cuticle

phenotype. The frequency of wild type cuticle among total female progeny without FM6 (i.e. bb^{z9}

and $bb^{z9\text{-}mag}$ / bb^{158}) was scored as 'magnification frequency'.

<u>Statistical analysis</u>

For comparison of sister chromatid segregation patterns in Fig. 1, C and D, Fig. 2F, Fig. 4G, and

fig. S1, significance was determined by two-sided Fisher's exact tests. For comparison of

frequencies of *bobbed* animals in Fig. 3B and wild-type cuticle animals in Fig. 3E, significance

was determined by two-tailed chi-squared tests using a 2 x 2 contingency table (normal; *bobbed*).

Other than these, significance was determined by two-tailed Mann-Whitney tests.

Supplementary Text

The results shown in Fig. 4 have a few critical implications. Under non-magnifying conditions,

USCE appears to be rare, based on the equal amount of IGS FISH signal in anaphase GSCs (Fig.

4, C and E) and based on the fact that both 359-bp repeats and $(TAGA)_n$ repeats segregate non-

randomly (Fig. 4G). It is interesting to note that without asymmetry in rDNA copy number (under

non-magnifying conditions), GSCs still faithfully retain a specific strand ('red strand') (Fig. 4, F

and G). This suggests that NRSS is not mediated by actual copy number differences, but rather

implies that sister chromatids (of rDNA loci) may have additional inherent asymmetries. We

speculate that such asymmetries may correlate with the propensity of a specific sister chromatid

to gain rDNA copy number, should USCE occur. An attractive candidate for such an asymmetry

is the molecular asymmetry during DNA replication that is specific to rDNA loci. It is well

established that DNA replication occurs unidirectionally in rDNA loci (*11, 12*) due to the presence

of a replication fork block on one side of the replication origin (fig. S5A). Accordingly, one sister

chromatid is mostly replicated as the leading strand, whereas the other is mostly replicated as the

lagging strand. In yeast, the DNA break that induces rDNA copy number recovery is known to

occur on the leading strand (the strand mostly replicated as the lagging strand) at the replication

fork block (fig. S5B; (*13*)). If this is universal, the broken end of the leading strand has limited

choices as to where to recombine with the sister chromatid to repair the DNA break. The broken

end would not recombine with a region that is not yet replicated. The recently replicated region of

the lagging strand, where Okazaki fragments have not been processed, may not be a good substrate

for sister chromatid recombination, either. The remaining possible region would be the sister

chromatid that was replicated as the leading strand (fig. S5B). If this happens, the strand mostly

replicated as the lagging strand is likely to gain the copy number. Thus, we speculate that the

mechanism that mediates NRSS may have the ability to distinguish leading vs. lagging strands and

specifically connects the lagging strand to the GSC side.

Although the mechanism that ensures the retention of a specific strand to the GSC remains elusive,

the CO-FISH results shown in Fig. 4, F and G provide a critical hint. Under magnifying conditions,

where USCE occurs, it is the $(TAGA)_n$ repeats whose segregation pattern is randomized. This

suggests that the 359-bp side of the rDNA is responsible for the retention in GSCs. This side of

the chromosome contains the centromere, whose asymmetry has been suggested to mediate non-

random segregation of chromosomes (*14, 15*). However, we have no evidence thus far to suggest

that the centromere is responsible for NRSS of the X and Y chromosomes. Additionally, the loss

of rDNA with retention of most of 359-bp and the entire centromere was sufficient to compromise

NRSS (Fig. 1C). Therefore, it is highly unlikely that 359-bp or the centromere contains sufficient

information to mediate NRSS. Future investigation is required to address these key molecular

mechanisms.

A

rDNA IGS only

D. simulans Y chr.

$(AATAAAC)_n$

B

Fig. S1. The *D. simulans* Y chromosome segregates sister chromatids non-randomly

    (A) Schematic of the *D. simulans* Y chromosome

    (B) Summary of the sister chromatid segregation pattern assessed by CO-FISH in the *D. simulans* control strain (w^{501}) (see table S1 for detailed data). Data shown as mean ± s.d. from three independent experiments. n, number of GSC-GB pairs scored. #, *P*-value of Fisher's exact test by comparing to hypothetical random sister chromatid segregation is shown.

Fig. S2. Validation of *indra^RNAi* efficiency and antibody specificity

(A) Examples of testis apical tips after *indra* knockdown by indicated *indra^RNAi* lines. Anti-Indra antibody staining was lost from germ cells upon *indra* knockdown by *nos-gal4>indra^TRiP.HMJ30228* or *nos-gal4>indra^GD9748 dcr-2*. (UAS-dcr-2 was added to enhance the efficiency of *indra^GD9748*). This experiment also demonstrates the specificity of the anti-Indra antibody. The hub is indicated by an asterisk. Germ cells are indicated by dotted lines and somatic cells are indicated by solid lines. Bar: 25 µm.

(B) Examples of testis apical tips before and after induction of *nos-gal4>indra^TRiP.HMJ30228*. GSCs are indicated by dotted lines and the hub is indicated by an asterisk. Bar: 10 µm.

(C) GSC number after induction of *indra^TRiP.HMJ30228*. n, number of testes scored. *P*-values: two-tailed Mann-Whitney test.

315
316 **Fig. S3. Localization of Indra-GFP to the nucleolus and rDNA loci**

(A) Localization of Indra-GFP at the apical tip of the testis. Indra localizes to nucleolus visualized by Fibrillarin. The hub is indicated by an asterisk. Bar: 25 µm.

(B) Localization of Indra-GFP on a metaphase chromosome spread. The X and Y chromosomes are indicated by dotted lines. Cid: centromere. Bar: 5 µm.

Fig. S4. Diagram of phenotypic assessment to detect rDNA magnification of the bb^{z9} allele

The bb^{z9} allele carries an insufficient rDNA copy number on the X chromosome, which causes flies to exhibit a '*bobbed*' cuticle phenotype when combined with the bb^{158} allele (no rDNA on X chromosome) in females (Fig. 3B). To induce magnification, the bb^{z9} allele was combined with a Y chromosome without rDNA (bb^{z9}/Ybb^-)('magnifying condition'). To assess whether the bb^{z9} allele magnified, these bb^{z9}/Ybb^- males were crossed to $bb^{158}/FM6$ female, and cuticle phenotype of the resulting bb^{z9}/bb^{158} females was examined. If magnification occurred, the magnified allele (bb^{z9-mag}) combined with bb^{158} would have a wild type cuticle, whereas the non-magnified allele combined with bb^{158} would have the *bobbed* phenotype. The frequency of wild type cuticle among total female progeny without FM6 (i.e. bb^{z9} and bb^{z9-mag}/bb^{158}) was scored as 'magnification frequency'.

337
Fig. S5. Diagram of DNA replication at rDNA loci

    (A) Replication fork block on one side of the replication origin leads to mostly unidirectional DNA replication at the rDNA loci. This causes one sister chromatid to be synthesized primarily as leading strand and the other as lagging strand.

    (B) In yeast, double strand DNA breaks primarily occur on the leading strand when fork progression is prevented at the replication fork block (top). An appropriate donor for DNA repair may be found in the region of the sister chromatid replicated as leading strand. If such recombination happens, the sister chromatid mostly replicated as lagging strand (bottom strand) will gain copy number.

Table S1. CO-FISH results in *D. melanogaster* rDNA deficient stocks and *D. simulans*

		Outcome	
		Y chromosome	**X chromosome**
D. melanogaster	wild type (*yw*)	76.9%:23.1% (±2.2%) (n=299)	77.4%:22.6% (±1.0%) (n=124)
	Df(1)bb158/Y	76.2%:23.8% (±3.5%) (n=63)	42.4%:57.6% (±11.9%) (n=33)
	X/Df(YS)bb-	45.7%:54.3% (±8.7%) (n=46)	75.5%:24.5% (±5.7%) (n=53)
D. simulans	wild type (*w501*)	80.0%:20.0% (±1.8%) (n=45)	N.D.

Probes used:

D. mel Y chromosome: Cy3-(AATAC)$_6$, Cy5-(GTATT)$_6$

D. mel X chromosome: Cy3-359 forward, Cy5-359 reverse

D. sim Y chromosome: Cy5-(GTTTATT)$_6$, Cy3-(AATAAAC)$_6$

Table S2. CO-FISH results of Y-2 translocation chromosomes

		Outcome
		T(Y;2) chromosome
T(Y;2)A77/+; XO	2^Y (with rDNA)	79.3%:20.7% (±0.5%) (n=58)
	Y^2	N.D.
T(Y;2)P8/+; XO	2^Y	46.6%:53.4% (±5.8%) (n=73)
	Y^2 (with rDNA)	78.1%:21.9% (±9.3%) (n=32)

Probes used:

T(Y;2) chromosome 2^Y: Cy5-(GTTTATT)$_6$, Cy3-(AATAAAC)$_6$

T(Y;2) chromosome Y^2: Cy3-(AATAC)$_6$, Cy5-(GTATT)$_6$

365 **Table S3. List of proteins that were enriched in IGS-beads pull-down**

Gene Name	Experiment 1	Experiment 2	Localized to nucleolus?	Localized to rDNA in mitosis?
Rrp1	19/0	7/0	N.D.	N.D.
CG2199/Indra	8/0	2/0	**YES**	**YES**
Iswi	8/0	0/0	**YES** (*16*)	N.D.
D1	60/3	37/0	**NO**	**NO**
apt	9/0	13/0	N.D.	N.D.
IleRS	26/0	11/0	**NO** (*17*)	N.D.
Dp1	21/0	9/0	N.D.	N.D.
dre4	21/0	9/0	**(YES)***	**(NO)***
Dsp1	14/0	8/0	**YES**	**NO**
clu	15/0	7/0	**NO** (*18*)	N.D.
Hrb27C	14/0	6/0	**NO** (*19*)	N.D.
TppII	20/0	6/0	**NO** (*20*)	N.D.
Prosα3	6/0	5/0	N.D.	N.D.
l(2)37Cc	14/2	5/0	**NO** (*21*)	N.D.
Cyt-c-p	9/0	5/0	**NO** (*22*)	N.D.
RpL8	9/2	5/0	**YES** (*23*)	N.D.
CG3995	5/0	5/0	**NO**	**NO**
TFAM	25/4	24/4	**NO** (*24*)	N.D.

367 * As no reagents to visualize the localization of dre4, which is a component of the FACT
368 complex, were available, the localization of SSRP1 (another component of the FACT complex)
369 was used when deciding whether or not to follow up dre4 in this study.

371 Data are shown as peptide counts in IGS beads/control beads.

373 **Table S4. CO-FISH results upon knockdown of *indra***

	Outcome	
	Y chromosome	**X chromosome**
indra^{TRiP.HMJ30228} control*	75.0%:25.0% (±2.6%) (n=40)	80.0%:20.0% (±1.7%) (n=30)
tub-gal80^{ts}, nos-gal4ΔVP16>UAS-indra^{HMJ30228}	40.0%:60.0% (±0.0%) (n=30)	43.6%:56.4% (±9.6%) (n=39)
indra^{GD9748} control (*nos-gal4>UAS-Dcr-2*)	76.9%:23.1% (±7.4%) (n=26)	76.9%:23.1% (±7.9%) (n=39)
nos-gal4>UAS-indra^{GD9748}, UAS-Dcr-2	42.9%:57.1% (±2.1%) (n=28)	43.2%:56.8% (±7.5%) (n=44)

374 Probes used:
375 Y chromosome: Cy3-(AATAC)$_6$, Cy5-(GTATT)$_6$
376 X chromosome: Cy3-359 forward, Cy5-359 reverse
377 *: Cross siblings of *tub-gal80^{ts}, nos-gal4ΔVP16>UAS-indra^{TRiP.HMJ30228}* that do not express
378 *indra^{HMJ30228}* (either *nos-gal4ΔVP16* only or *UAS-indra^{TRiP.HMJ30228}* only) were used as control
379
380
381 **Table S5. CO-FISH results of X chromosome in magnifying condition**

		Outcome
		X chromosome
Non-magnifying (*yw*)	TAGA	87.5%:12.5% (±11.3%) (n=32)
	359	81.1%:18.9% (±0.9%) (n=37)
Magnifying (bb^{z9}/Ybb^-)	TAGA	53.3%:46.7% (±4.7%) (n=45)
	359	74.2%:25.8% (±7.7%) (n=31)

382 Probes used:
383 (TAGA)$_n$: Cy3-(TAGA)$_8$, Cy5-(TCTA)$_8$
384 (359)$_n$: Cy3-359 forward, Cy5-359 reverse

391 **Table S6. Probe sequences for CO-FISH and DNA FISH**

Probe target	5'-sequence-3'	Source or reference	Related figure
(AATAC)$_n$ (forward)	Cy3-(AATAC)$_6$	(8)	Fig. 1, B-D, Fig. 2F
(AATAC)$_n$ (reverse)	Cy5-(GTATT)$_6$		Fig. 1, B-D, Fig. 2F
359-bp (forward)	Cy3-CCACATTTTGCAAATTTTGATGACCCCCCTCCTTACAAAAAATGCG		Fig. 1C, Fig. 2F, Fig. 4G
359-bp (reverse)	Cy5-AGGATTTAGGGAAATTAATTTTTGGATCAATTTTCGCATTTTTTGTAAG		Fig. 1C, Fig. 2F, Fig. 4G
(AATAAAC)$_n$ (forward)	Cy3-(AATAAAC)$_6$	This study	Fig. 1D, fig. S1B
(AATAAAC)$_n$ (reverse)	Cy5-(GTTTATT)$_6$	(25)	Fig. 1D, fig. S1B
(TAGA)$_n$ (forward)	Cy3-(TAGA)$_8$	(25)	Fig. 4G
(TAGA)$_n$ (reverse)	Cy5-(TCTA)$_8$	This study	Fig. 4G
240-bp IGS	Cy5-TCCATTCACTAAAATGGCTTTTCTCTATAATACTTAGAGAATATGGGAATATTTCAACATTTTTCACT	(26)	Fig. 4, C-E

397 **Table S7: Primer and probe sequences for Real-Time PCR and Droplet-Digital PC**

Primer name	5'-sequence-3'	Source or reference
rt-5S rDNA (forward)	AAGTTGTGGACGAGGCCAAC	(27)
rt-5S rDNA (reverse)	CGGTTCTCGTCCGATCACCGA	
rt-IGS #1 (forward)	GCTGTTCTACGACAGAGGGTTC	
rt-IGS #1 (reverse)	CAATATGAGAGGTCGGCAACCAC	
rt-IGS #2 (forward)	GGTAGGCAGTGGTTGCCG	
rt-IGS #2 (reverse)	GGAGCCAAGTCCCGTGTTC	
rt-ETS (forward)	ATTACCTGCCTGTAAAGTTGG	
rt-ETS (reverse)	CCGAGCGCACATGATAATTCTTCC	
rt-18S rDNA (forward)	TTCTGGTTGATCCTGCCAGTAG	
rt-18S rDNA (reverse)	CGTGTGTACTTAGACATGCATGGC	
rt-28S rDNA (forward)	CCTCAACTCATATGGGACTACC	This study
rt-28S rDNA (reverse)	CACTGCATCTCACATTTGCC	
dd-RpL32 (forward)	GCTTCAAGGGACAGTATCTG	(10)
dd-RpL32 (reverse)	AACGCGGTTCTGCATGAG	

dd-RpL32 (probe)	**HEX**-ATGCCCAACATCGGTTAC-**Iowa Black FQ**
dd-28S (forward)	GAGCTGCCATTGGTACAG
dd-28S (reverse)	GCTTTCGCCTTGAACTTAG
dd-28S (probe)	**HEX**-TGGTGGATAGTAGCAAATAATCG-**Iowa Black FQ**
dd-Upf1 (forward)	CACACTTTATGTCCACCATTATTG
dd-Upf1 (reverse)	GAGTTTCCGTAGGGACCAC
dd-Upf1 (probe)	**HEX**-CCGTAACCGCCACTGCGGT-**Iowa Black FQ**

399 **References**
400

1. M. Van Doren, A. L. Williamson, R. Lehmann, Regulation of zygotic gene expression in
Drosophila primordial germ cells. *Curr Biol* **8**, 243-246 (1998).
2. M. P. Zeidler, N. Perrimon, D. I. Strutt, Polarity determination in the Drosophila eye: a
novel role for unpaired and JAK/STAT signaling. *Genes Dev* **13**, 1342-1353 (1999).
3. S. E. McGuire, P. T. Le, A. J. Osborn, K. Matsumoto, R. L. Davis, Spatiotemporal rescue
of memory dysfunction in Drosophila. *Science* **302**, 1765-1768 (2003).
4. M. Inaba, M. Buszczak, Y. M. Yamashita, Nanotubes mediate niche-stem-cell signalling
in the Drosophila testis. *Nature* **523**, 329-332 (2015).
5. M. Zaccai, H. D. Lipshitz, Differential distributions of two adducin-like protein isoforms
in the Drosophila ovary and early embryo. *Zygote* **4**, 159-166 (1996).
6. B. Riggleman, P. Schedl, E. Wieschaus, Spatial expression of the Drosophila segment
polarity gene armadillo is posttranscriptionally regulated by wingless. *Cell* **63**, 549-560
(1990).
7. M. Jagannathan, R. Cummings, Y. M. Yamashita, A conserved function for
pericentromeric satellite DNA. *Elife* **7**, (2018).
8. S. Yadlapalli, Y. M. Yamashita, Chromosome-specific nonrandom sister chromatid
segregation during stem-cell division. *Nature* **498**, 251-254 (2013).
9. A. M. Huang, E. J. Rehm, G. M. Rubin, Quick preparation of genomic DNA from
Drosophila. *Cold Spring Harb Protoc* **2009**, pdb prot5198 (2009).
10. J. O. Nelson, A. Slicko, Y. M. Yamashita, The retrotransposon R2 maintains Drosophila
ribosomal DNA repeats. *bioRxiv*, 2021.2007.2012.451825 (2021).
11. Y. Akamatsu, T. Kobayashi, The Human RNA Polymerase I Transcription Terminator
Complex Acts as a Replication Fork Barrier That Coordinates the Progress of Replication
with rRNA Transcription Activity. *Mol Cell Biol* **35**, 1871-1881 (2015).
12. T. Kobayashi, Ribosomal RNA gene repeats, their stability and cellular senescence. *Proc
Jpn Acad Ser B Phys Biol Sci* **90**, 119-129 (2014).
13. M. D. Burkhalter, J. M. Sogo, rDNA enhancer affects replication initiation and mitotic
recombination: Fob1 mediates nucleolytic processing independently of replication. *Mol
Cell* **15**, 409-421 (2004).
14. T. Akera, E. Trimm, M. A. Lampson, Molecular Strategies of Meiotic Cheating by
Selfish Centromeres. *Cell* **178**, 1132-1144 e1110 (2019).
15. R. Ranjan, J. Snedeker, X. Chen, Asymmetric Centromeres Differentially Coordinate
with Mitotic Machinery to Ensure Biased Sister Chromatid Segregation in Germline
Stem Cells. *Cell Stem Cell* **25**, 666-681 e665 (2019).
16. A. V. Emelyanov *et al.*, Identification and characterization of ToRC, a novel ISWI-
containing ATP-dependent chromatin assembly complex. *Genes Dev* **26**, 603-614 (2012).
17. J. Lu, S. J. Marygold, W. H. Gharib, B. Suter, The aminoacyl-tRNA synthetases of
Drosophila melanogaster. *Fly (Austin)* **9**, 53-61 (2015).
18. R. T. Cox, A. C. Spradling, Clueless, a conserved Drosophila gene required for
mitochondrial subcellular localization, interacts genetically with parkin. *Dis Model Mech*
2, 490-499 (2009).
19. M. J. Matunis, E. L. Matunis, G. Dreyfuss, Isolation of hnRNP complexes from
Drosophila melanogaster. *J Cell Biol* **116**, 245-255 (1992).

20. S. C. Renn, B. Tomkinson, P. H. Taghert, Characterization and cloning of tripeptidyl
peptidase II from the fruit fly, Drosophila melanogaster. *J Biol Chem* **273**, 19173-19182
(1998).
21. S. J. Lee, R. Feldman, P. H. O'Farrell, An RNA interference screen identifies a novel
regulator of target of rapamycin that mediates hypoxia suppression of translation in
Drosophila S2 cells. *Mol Biol Cell* **19**, 4051-4061 (2008).
22. L. Dorstyn, K. Mills, Y. Lazebnik, S. Kumar, The two cytochrome c species, DC3 and
DC4, are not required for caspase activation and apoptosis in Drosophila cells. *J Cell Biol*
167, 405-410 (2004).
23. K. N. Rugjee *et al.*, Fluorescent protein tagging confirms the presence of ribosomal
proteins at Drosophila polytene chromosomes. *PeerJ* **1**, e15 (2013).
24. K. Takata *et al.*, Drosophila mitochondrial transcription factor A: characterization of its
cDNA and expression pattern during development. *Biochem Biophys Res Commun* **287**,
474-483 (2001).
25. M. Jagannathan, N. Warsinger-Pepe, G. J. Watase, Y. M. Yamashita, Comparative
Analysis of Satellite DNA in the Drosophila melanogaster Species Complex. *G3*
(Bethesda) **7**, 693-704 (2017).
26. K. L. Lu, J. O. Nelson, G. J. Watase, N. Warsinger-Pepe, Y. M. Yamashita,
Transgenerational dynamics of rDNA copy number in Drosophila male germline stem
cells. *Elife* **7**, (2018).
27. Q. Zhang, N. A. Shalaby, M. Buszczak, Changes in rRNA transcription influence
proliferation and cell fate within a stem cell lineage. *Science* **343**, 298-301 (2014).
466