Decoupling of rates of protein synthesis from cell expansion leads to supergrowth

Benjamin D. Knapp[1,2], Pascal Odermatt[1,3], Enrique R. Rojas[3,4], Wenpeng Cheng[5], Xiangwei He[5], Kerwyn Casey Huang[2,3,4,6,7,*], and Fred Chang[1,*]

[1]Department of Cell and Tissue Biology, University of California, San Francisco, San Francisco, CA 94143

[2]Biophysics Program, Stanford University, Stanford, CA 94305, USA

[3]Department of Bioengineering, Stanford University, Stanford, CA 94305, USA

[4]Department of Biochemistry, Stanford University School of Medicine, Stanford, CA 94305, USA

[5]Life Sciences Institute, Zhejiang University, Hangzhou, Zhejiang, China, 310027

[6]Department of Microbiology and Immunology, Stanford University School of Medicine, Stanford, CA 94305, USA

[7]Chan Zuckerberg Biohub, San Francisco, CA 941586

[*]To whom correspondence should be addressed:

Fred Chang

513 Parnassus Avenue, HSW616

San Francisco, CA 94143 USA

Phone: +1 (415) 476-8922

E-mail: fred.chang@ucsf.edu

Kerwyn Casey Huang

Shriram Center, Room 007 MC: 4245

443 Via Ortega

Stanford, CA 94305-4125 USA

Phone: +1 (650) 721-2483

E-mail: kchuang@stanford.edu

ABSTRACT

Cell growth is a complex process in which the biosynthesis of cellular components must be coordinated with increases in cell size in order to maintain their concentrations. In many cell types, the steady-state growth rate is proportional to cell size such that larger cells grow proportionally faster than smaller cells. Little is known about the mechanisms coordinating biosynthesis, cell volume, and cytoplasmic density. Here, we reveal a global mechanism for proteome homeostasis by transiently decoupling protein synthesis from volume growth rate in the fission yeast *Schizosaccharomyces pombe*. Rapid osmotic oscillations strongly inhibited growth, but when oscillations were terminated cells subsequently underwent a period of "supergrowth" in which growth was significantly faster than steady-state growth for multiple generations. Protein concentrations and cytoplasmic density increased steadily during oscillations, and gradually decreased to steady-state values during supergrowth. Transient growth inhibition by disabling the secretory pathway produced similar behaviors to osmotic oscillations. The accumulation of biomass was responsible for driving subsequent rapid growth, yielding a homeostatic mechanism for maintaining global protein concentration.

INTRODUCTION

Proliferating cells generally increase their biomass and volume during the cell cycle before dividing. Although much is understood about how duplication of certain cellular components such as the chromosomes is accomplished, much less is known about how the proteome itself is duplicated. The concentrations of many proteins are thought to be maintained during cell growth [1-3]. While global mechanisms of proteome homeostasis have been identified [4], it is unknown the extent to which the concentrations of individual proteins are coordinated with cell volume [5], whether this coordination can be altered, and the effects of such a perturbation.

The rate of cellular growth, which we define here as the increase of cellular volume, is determined by numerous factors. The biosynthesis of cellular components has been speculated to set growth rate, as decreasing protein translation, for instance, can slow or halt growth. The rate of cellular growth is also affected by cell size; many organisms, including bacteria, fungi, and mammalian cells, exhibit exponential growth [6,7] in which the absolute growth rate at steady-state is proportional to cell size. Although the mechanism(s) for achieving exponential growth remains to be determined, it may arise from the scaling of the biosynthetic machinery with cell size: if protein synthesis is coupled to cell size, and cell size is dictated by protein concentration, then exponential growth will result.

The fission yeast *Schizosaccharomyces pombe* is an established model for cell-cycle regulation and growth. The simple, rod-shaped morphology and regular growth and division patterns of these cells make them highly amenable to quantitative studies. During their cell cycle, *S. pombe* cells exhibit polarized tip growth at one or both cell tips during interphase [8,9], and growth halts during mitosis and cytokinesis [10,11]. Like other tip-growing cells, the growth of the cell surface is directed by polarity machinery that ultimately mediates remodeling and insertion of new cell wall at the cell tips. Growth of the surface is further impacted by mechanical factors, such as the turgor pressure due to osmolyte concentration imbalances across the membrane that expands the elastic cell wall [8,10]. While exponential volume growth at the single-cell level has been observed in many cell types, whether individual fission yeast cells exhibit such behavior has been a source of controversy for decades; the current consensus is a bilinear growth behavior with an increased slope later in the cell cycle [11-16].

Biomass synthesis (largely driven by protein synthesis) and volume increase (driven by

membrane and cell-wall synthesis) must be coordinated during growth, but it is unknown how

this fundamental coupling is achieved. Here, we present multiple ways of uncoupling the

coordination between biomass and volume in *S. pombe* cells, and thus provide new insights into

the role of proteome homeostasis in cell growth. We first establish that fission yeast cells exhibit

cell size-dependent, exponential growth for a large fraction of the cell cycle. Through

manipulation of turgor pressure or secretion, we decouple the rate of global protein synthesis

from growth rate. These conditions produce a global excess of proteins within the cytoplasm,

which in turn results in an extended period of extremely rapid growth. These findings thus

provide new insights into the role of cytoplasmic density in cell growth and proteome

homeostasis, whereby the rapid expansion of dense cells accelerates the re-equilibration of

protein concentrations.

RESULTS

Fission yeast cells grow exponentially during a large fraction of the cell cycle

To determine the growth behavior of individual *S. pombe* cells, we imaged cells in time lapse in microfluidic chambers and implemented automated image analysis methods to quantify cellular dimensions at subpixel resolution (Figure 1A, S1A; Methods). In agreement with previous studies, these rod-shaped cells elongated from ~8 μm to 14 μm before entering mitosis (Figure 1B, S1B), and maintained a constant width of ~4 μm (Figure 1B, S1C). Instantaneous growth rates (defined as dL/dt, where L is cell length) of wild-type cells ranged from 2-5 μm/h at 30 °C (Figure 1C), with cell length highly correlated with elongation rate, indicating that growth rate accelerates during the cell cycle. In purely exponential growth, absolute growth rate scales linearly with cell size, hence $\dot{e} = \frac{1}{L}\frac{dL}{dt}$ (size-normalized growth rate) is a constant. Quantification of $\dot{e}$ revealed that fission yeast growth during the cell cycle is organized into at least three phases (Figure 1D): (I) growth acceleration for the first ~10-20% of the cell cycle (immediately after cell separation), (II) exponential growth for the majority of interphase, in which $\dot{e} = 0.35 \pm 0.06$ h^{-1} (corresponding to doubling biomass during interphase in 2.0±0.3 h), and (III) deceleration to zero growth in mitosis and cytokinesis. These data are inconsistent with simple linear or bilinear growth models (Figure S2). We further tested the effect of cell size on growth rate by analyzing abnormally large cells (*cdc25-22* cells at the semi-permissive temperature 30 °C, which divide at 30±4 μm in length). These large cells grew much faster than wild-type cells and followed a similar scaling of single-cell elongation rate with size, with a size-normalized growth rate $\dot{e}$ of 0.30 ± 0.05 h^{-1} (Figure 1C, S1D,E). These findings establish exponential growth behavior in fission yeast, with absolute growth rates that scale with cell size over a wide range of volumes.

Cells grow slowly during oscillatory osmotic shocks, and exhibit super-fast growth upon exit from oscillations

In terms of the mechanics of cell growth, internal turgor pressure provides physical force for expansion of the cell wall to increase cellular volume. Fission yeast cells have a thick (~200 nm), elastic cell wall inflated by high turgor pressures of ~1.5 MPa (15 atm) [10,17]. Examining the effects of adding the osmotic agent D-sorbitol (hereafter sorbitol) to the media has proven useful

for probing the mechanical effects of reducing turgor pressure [10,18-20]. We investigated the effect of mechanical perturbations on cell growth using osmotic oscillations in repeated cycles of hyper- and hypoosmotic shocks (Methods) [21]. We exposed *S. pombe* cells in a microfluidic device to repeated switches between rich medium (YE5S) and YE5S+0.5 M D-sorbitol (Figure 2A, S3A). In these initial experiments, we applied 24 oscillatory cycles of 0.5 M sorbitol shocks with a 10-min period for 4 hours, then subsequently followed cells in YE5S media without sorbitol. Each acute addition of 0.5 M sorbitol caused rapid water efflux, loss of turgor pressure, and shrinkage in volume by ~20%, with mean longitudinal and radial contractions in cell size of 3.5% and 7%, respectively (Figure 2B) [10]. During each 5-min hyperosmotic period, cells partially re-inflated, dependent on osmotic adaption mechanisms (Figure 2B, S3B). Upon hypoosmotic shifts back to YE5S medium without sorbitol, cells rapidly returned to their normal width in the absence of shocks (Figure 2B, S3B), suggesting that *S. pombe* cells can rapidly downregulate turgor pressure to a preferred value when envelope stresses increase above a certain point.

Over the course of 4 hours of these osmotic cycles, cells remained viable and continued to grow and divide (Movie S1, Figure S3C). However, the mean growth rate was lower than that of control cells: growth initially slowed (Figure 2C) for the first ~2 h (to <0.06 h^{-1}, Figure 2D), followed by adaptation to the oscillations after ~3 h that resulted in a size-normalized growth rate increase to ~0.2 h^{-1} (Figure 2D). When cells were returned after the cycles of osmotic oscillations to YE5S without any sorbitol, they grew unusually fast, with tip growth rates >8 μm/h (population mean 6.5±0.9 μm/h), almost two-fold higher than mean control tip growth rates (3.8 μm/hr, Figure 2C); we refer to this period of unprecedented rapid growth as "supergrowth." Size-normalized growth rates indicated that cells were growing abnormally rapidly for their size ($\dot{e} = 0.6$ h^{-1} vs. 0.35 h^{-1} for control cells, Figure 2D). These growth behaviors were highly stereotypical throughout the population (>96% cells, 49/51), and were substantially faster than any previously reported growth rates for fission yeast. Acceleration began by 15 minutes after exit from oscillations (Figure 2C), and growth rate reached a peak just before cells entered mitosis (Figure 2C, S4A). Elevated growth rates persisted for 2-3 generations (~4 hours, Figure 2C,D, S4B), but in each cell cycle, the rate decreased in a step-wise manner from the previous generation even though cells still exhibited exponential growth (Figure S4B,C). Growth still decelerated normally during mitosis and cytokinesis. Interestingly, the initial acceleration

phase of the cell cycle (Figure 1D) was largely absent during supergrowth (Figure S4B),

suggesting that cells were primed for exponential growth immediately after division.

Despite the large changes in growth rate during and after the oscillations, cell morphology and

size were remarkably normal. Cells exhibited tip growth during interphase without large

changes in cell width or tip shape (Figure S5). Importantly, cells entered mitosis at the normal

cell length of 14 μm during oscillations and during supergrowth despite a >10-fold range in

growth rates, indicating that changes in growth rate did not affect cell-size control (Figure 2E).

During supergrowth, cell-cycle periods were shorter because due to a decrease in the duration of

interphase, but the durations of mitosis and cytokinesis were normal, indicating that not all

cellular processes were sped up (Figure S4C,D). These data suggest that cell size and the periods

of cell division are controlled independently of growth rate.

To determine the requirements for supergrowth, we examined its dependence on oscillation

parameters (amplitude, period, number of periods). We systematically varied each parameter

around our baseline values (0.5 M, 10-min period, 24 periods), keeping the other two

parameters fixed. In general, the maximal supergrowth rate increased gradually with increasing

amplitude, period, or number of periods (Figure 2F). Single shocks were generally not sufficient

(Figure 2Fii). Oscillations with 1 M sorbitol, which caused ~50% volume loss, resulted in a near-

cessation of growth during oscillations, but then subsequently led to rapid supergrowth (Figure

2Fiii). These data suggest that growth and cell-cycle progression during oscillations are not

required for supergrowth. The graded effects suggest that the rapid growth state is not regulated

by an all-or-nothing switch, but rather accumulates over time during the osmotic shifts.

Supergrowth is independent of established osmotic stress and growth pathways

To probe the mechanism underlying supergrowth, we considered two non-exclusive models. The

first model involves the osmotic stress-response pathway, which could signal to growth

pathways to regulate growth rate. The second model is based on material storage, in which

components important for growth accumulate during periods of slow volume expansion because

biosynthesis does not slow commensurately with growth rate. The excess of these materials

upon exit from oscillations could then drive the subsequent supergrowth.

As an initial test of the signaling model, we assayed various mutants for their ability to undergo

supergrowth. We assessed strains lacking Sty1 and Pmk1 (MAP kinases at the hub of the

pathways that response to osmotic, cell-wall, and other environmental stresses) [22,23]; Gpd1,

which regulates glycerol synthesis during turgor adaptation to osmotic stresses [19,22]; and Cch1, a

calcium channel involved in response to cell-wall stresses [24]. We also queried regulators in the

TOR pathway, a central regulator of growth: Ssp2 (an AMPK-like protein kinase that regulates

TORC1 activity) [25,26] and Gad8 (an AGC protein kinase effector of TORC2) [27]. All these mutants

exhibited supergrowth responses similar to that of wildtype after oscillatory osmotic shock

(Figure S6). Ribosomal mutants with slower basal growth rates showed a proportional

supergrowth response (Figure S6). Thus, these signaling pathways regulating cellular responses

to stress, turgor, and growth are not required for supergrowth.

The proteome globally increases in concentration during osmotic oscillations

The material storage model predicts that components important for growth rise in concentration

during oscillations. As an initial test, we monitored a fluorescent protein marker, E2-mCrimson

expressed from the *ACT1* (actin gene) promoter. E2-mCrimson is stable and folds relatively

rapidly [28]. Under normal growth, mCrimson intensity remained approximately constant

throughout the cell cycle (Figure S7A), indicating that this marker is produced at a rate

proportional to volume growth rate (Figure S7B), the expected behavior for many native

proteins. In contrast, during osmotic oscillations, mCrimson fluorescence concentration

increased, rising linearly to ~50% above normal levels after 4 h of 10-min, 500-mM oscillations

(Figure 3A-C). When cells entered supergrowth, mCrimson intensity progressively decreased

back to wild-type levels over 2-3 generations (Figure 3A-C), similar to the time scale for

restoration of normal growth rates.

To investigate why mCrimson accumulates during osmotic oscillations, we computed the rates of

mCrimson production and compared to growth rate. During oscillations, the rate of mCrimson

production remained close to normal levels even though cells grew in volume at a much slower

rate than normal (Figure S7D). Thus, mCrimson increased in concentration not because of a large increase in protein production, but because of a failure to decrease protein production as much the reduction in volume growth rate.

We also examined how mCrimson concentration relaxed to equilibrium levels during supergrowth. In this phase, the cells grew in volume much faster than mCrimson production (Figure S7D). Much of this decrease in concentration can thus be explained by a dilution mechanism, which is enhanced by the rapid growth. The rate of "production" was also even lower than normal levels (Figure S7D), which may reflect decreased protein synthesis or increased degradation rates. Calculations using our growth-rate measurements to predict the decay constant of mCrimson fluorescence, assuming steady-state production after oscillations, indicated that at least 75% of the fall in mCrimson levels relative to controls can be explained by dilution during growth. These findings highlight dilution by rapid growth as a mechanism contributing to protein homeostasis.

Next, we determined whether native cellular proteins also increase in concentration during oscillations. We imaged fluorescently tagged proteins representing a variety of cellular processes, such as metabolism, chromosomal organization, cell-size control, and cell growth (Figure 3D, Table S1). These proteins all maintained a constant concentration during normal growth. Much like E2-mCrimson, the concentrations of many of these cellular proteins increased during 4 hours of 10-min, 0.5 M osmotic oscillations, and then returned gradually to normal during supergrowth (Figure 3D). No aggregates of these tagged proteins were observed. Similar changes in intensities were observed for fusions to various fluorescent proteins (Figure 3D), suggesting that the behavior was not an artifact of any particular fluorescent protein. Among the different protein fusions, there was a large range of behavioral variation, with different rates of increase and decrease during oscillations and supergrowth. Impressively, some proteins accumulated by ~100% during oscillations. In contrast, a subset of proteins displayed modest or no increases in concentration during oscillations. For example, the concentration of histone H2A was relatively steady throughout the experiment (Figure 3D), perhaps reflecting the fact that the amount of DNA is not increasing abnormally in these cells. The cell-size sensor Cdr2 also increased only modestly during oscillations (Figure 3D), consistent with the maintenance of size

control. Ribosomal proteins also showed only modest increases (Figure 3E; see section below).

These findings suggest that osmotic oscillations caused large changes in the concentrations of a

significant subset of cellular proteins.

We next employed two complementary methods to measure the global state of cellular

components. Staining of individual cells for total protein with fluorescein isothiocyanate (FITC)

showed a similar rise and fall, with a mean 20% increase during 2 hours of osmotic oscillations

(Figure 3F), while DNA concentration remained constant (Figure S8). Quantitative-phase

imaging (Methods) showed that intracellular density, as measured from the diffractive index of

the cell, also exhibited a ~10% and 20% increase during 0.5-M and 1-M oscillations, respectively

(Figure 3G). Both measurements were in approximate agreement with the average increases in

protein concentrations (Figure 3D). Thus, osmotic oscillations clearly produce global changes in

protein concentration and cytoplasmic density.

Other oscillation regimes induced even larger increases in protein concentration. For instance,

over the course of 48 oscillation cycles with 1 M sorbitol, cells halted growth and accumulated

150% higher mCrimson concentrations (Figure S9). These additional oscillatory cycles of 1M

sorbitol displayed no sign of saturation in linear concentration accumulation (Figure S9B),

suggesting that even higher increases are likely possible.

We sought to determine if a simple model of growth kinetics could quantitatively predict such a

linear increase during growth inhibition. Experimental measurements have shown that

translational efficiency $\mu = \lambda_1 \frac{1}{V}\frac{dV}{dt} + \lambda_o$ contains a growth-dependent and growth-independent

component λ_0 [29], which leads to a prediction of steady-state growth rate that is within 30% of

experimental measurements (see Supplementary Information for more details). In our

experiments with 1-M oscillatory shocks, growth halts completely and ribosome concentration is

constant. In this scenario, our model predicts that the baseline translational efficiency λ_0 [29]

results in accumulation of biomass M at a constant rate $dM/dt = \lambda_0 R$, where R is the ribosome

abundance. Hence, M is predicted to increase linearly during osmotic shock oscillations, at a

reduced rate relative to unperturbed growth that yields a ~12% increase over 4 hours, similar to

our experimental measurements (Figure 3G). Taken together, our experimental measurements

and theoretical model indicate that oscillations and supergrowth represent periods of

decoupling of protein production and cell expansion, and lead to profound accumulation of

intracellular proteins.

Increased protein concentrations drive supergrowth

Next, we asked whether the increased concentration of proteins is responsible for supergrowth.

First, further analysis of our extensive exploration of oscillation parameters revealed that the

maximal supergrowth rate was highly correlated with the observed increase in mCrimson

concentration during oscillations (Figure 4A, r = 0.85). Second, we tested whether the increase of

protein concentration is necessary for supergrowth. To globally inhibit protein synthesis, we

treated cells with 100 µg/mL cycloheximide [30] during oscillations, and then washed out the

inhibitor upon exit from oscillations. These cycloheximide-treated cells recovered to near-

normal growth rates but never achieved supergrowth (Figure 4B). Thus, efficient protein

translation during oscillations is required for supergrowth.

Finally, we tested whether the accumulation of growth materials during oscillations is sufficient

to drive growth in the absence of new protein synthesis. For control cells that were not exposed

to oscillatory shocks, cycloheximide treatment caused a rapid cessation in cell growth. In

contrast, cells that had been exposed to oscillations still grew substantially, at ~3-fold faster

growth rates than control cells for 2 h (Figure 4C,D). Cells after oscillations also reestablished

steady-state growth rates upon removal of cycloheximide more quickly (Figure 4C). Collectively,

these results support the storage model in which cells with accumulated components are able to

undergo some growth even in the absence of new protein synthesis.

Effect of ribosomal concentration, cell wall synthases, and turgor pressure on supergrowth

To probe whether the up-regulation of particular cellular processes drives supergrowth, we

assessed three candidates involved in growth regulation: ribosomes, the cell-wall biosynthetic

machinery, and turgor pressure. We monitored ribosomal protein abundances in single cells

using ribosomal protein-GFP fusions expressed from their endogenous promoters (Methods) and

found that their steady-state concentrations were relatively constant (Figure 3E). The

concentrations of these ribosomal proteins increased only modestly during oscillations (from 0-

20%) (Figure 3E) and then quickly reverted to baseline levels during supergrowth. Some

ribosomal proteins displayed nearly constant concentration throughout the experiment (Figure

3E, S10), indicating that their production rates decreased and increased concurrently with the

growth rate. Thus, rapid supergrowth is not caused by a large increase in ribosome

concentration.

As cell-wall assembly likely limits the rate of cell expansion, a candidate factor for controlling

cell-wall growth is the major cell-wall glucan synthase Bgs4. Bgs4 is a trans-membrane protein

that localizes to growing cell tips during interphase and to the developing septum during

cytokinesis [31]. During the oscillations, as cells grew slowly (Figure S11A), Bgs4 puncta were

delocalized across the cell membrane (Figure S11B), consistent with the general depolarization

of actin and Cdc42 seen in the osmotic-shock response [31-33]. Bgs4 intensities at cell tips

decreased by 50%, although total Bgs4 concentration gradually increased (by ~50%) during this

period (Figure 3D). Upon exit from oscillations, Bgs4 rapidly repolarized to the cell tip, and tip

intensities rose to more than 50% more than in control cells within 30-45 min (Figure S11B).

The amount of Bgs4 at the cell tips strongly correlated with the growth rate in these experiments

(r = 0.95) (Figure S11C). It is likely that such increases in the levels of Bgs4 and additional

polarity factors contributing to cell wall growth collectively contribute to rapid growth. These

results reveal mechanistic insight into why cell growth slows down during osmotic shock

oscillations and how polarized growth is rapidly established after oscillations.

Another potential contributor to supergrowth could be an increase in turgor pressure that may

arise from increased cytoplasmic density. We examined turgor pressure by measuring cell

width, as changes in turgor pressure are expected to lead to concurrent reversible swelling in

cell width independent of tip elongation. Although cell widths increased slightly during osmotic

oscillations, they did not decrease during supergrowth as cytoplasmic density decreases (Figure

S5A, S12). These data indicate that the small width increases during oscillations are not

reversible and thus may arise from delocalized cell wall synthesis (Figure S11B,C) [32], rather than

from increases in turgor pressure. In sum, turgor pressure is unlikely to increase substantially

during supergrowth, and is not the driver of the increase in growth rate during supergrowth.

Transient inhibition of protein secretion also drives subsequent supergrowth

Our model predicted that other perturbations that transiently inhibit cell volume growth would

also lead to proteome accumulation and subsequent supergrowth. One way of stopping growth

and increasing buoyant density is through inhibition of secretion, as shown initially in budding

yeast [34]. To halt secretion, we treated mCrimson-expressing cells for 4 hours with 100 μg/mL

brefeldin A (BFA), which inhibits protein transport from the endoplasmic reticulum to the Golgi

apparatus [35,36]. Cells exhibited a rapid decrease in growth rate to essentially zero throughout

BFA treatment, during which time mCrimson concentration increased approximately linearly

(Figure 4E). Many cells eventually died, but the 45% of cells that survived the treatment

exhibited supergrowth upon BFA wash out (Figure 4E) with growth-rate trajectories similar to

those observed after 0.5 M sorbitol oscillatory osmotic shocks (Figure 2D). Thus, rather than just

a response to osmotic stress, supergrowth and proteome accumulation can be produced by other

treatments that limit volume expansion without proportional inhibition of protein synthesis.

DISCUSSION

Here, our interrogation of the growth response of *S. pombe* cells to osmotic perturbations has revealed unexpected insights into the relationships between cell size, growth, and biosynthesis. In investigating the basis for supergrowth, we find that these unprecedented rapid growth rates arise from a decoupling of biosynthesis and volume expansion. During the osmotic oscillations, volume expansion is impaired far more than biosynthesis, which leads to global accumulation of cellular components inside the cell. Release from these conditions then results in abnormally rapid growth that drives an accelerated return to normal proteome concentrations relative to dilution. Our results thus support a model in which rapid growth is driven by the accumulation of cellular materials. This effect of material storage on growth rate may be connected with the mechanisms underlying the near-universal scaling of growth rate with cell size (exponential growth) observed under steady-state environmental conditions across the kingdoms of life. In the simplest interpretation, cells in supergrowth grow rapidly because they have the internal resources of a larger cell. Whether the cellular materials that promote growth are some subset of the cytoplasmic constituents, or the entire collection, remains to be seen. Our data suggests that growth rate is set by a combination of cellular processes, including the biosynthetic machinery, metabolism, and cell-wall assembly, rather than by a single component such as ribosomes.

This work reveals that relatively simple manipulations can decouple cell growth and protein synthesis, yielding mechanistic insights into how these fundamental processes are normally coordinated. As concentrations of many proteins do not vary greatly during steady-state growth, it has been speculated that extensive feedback mechanisms exist for protein homeostasis; for instance, increased concentration of a certain protein may inhibit its own translation or transcription through a signaling pathway that senses its concentration, leading to a decrease back to normal levels; such a protein-specific mechanism has been suggested for tubulin homeostasis for instance [37]. In this scenario, we speculate that osmotic oscillations lead to breakdown of the putative feedback mechanisms, possibly because of the rapid changes in cell size and concentrations. However, our findings also raise a second, simpler possibility in which signaling feedback mechanisms may not exist for many proteins [38-40]. We speculate that volume expansion itself could act as a homeostatic mechanism, wherein biomass increases largely independent of volume and small variations in cytoplasmic protein concentrations are brought

back into balance by a correction in short-term volume growth rate. Abnormal increases in protein concentrations would drive faster growth than synthesis, which would decrease their concentration in part through dilution, similar to what we observe during supergrowth. A decrease in protein concentration would lead to slowed growth, allowing protein concentration to build back up. Our experiments were able to reveal such a mechanism for the coordination of growth and volume, because they allowed us to uncouple this coordination and examine the recovery process.

Such a mechanism however may apply to a subset of proteins, but not to all. During osmotic oscillations, certain groups of proteins (such as metabolic enzymes) maintained relatively high synthesis rates and hence rose in concentration. By contrast, other proteins – including ribosomal proteins, histones, and cell size regulators – are likely regulated so that their expression scales with growth to maintain their concentrations; the level of accumulation of these proteins during osmotic oscillations would then reflect the robustness of this regulation. Tight control of ribosomal protein concentration is predicted to be another factor to keep the proteome in check [41], preventing accumulation at more rapid rates when expansion is perturbed.

The remarkable cellular phenotype we have presented in which the concentrations of much of the proteome are altered, accompanied by substantial increases in cytoplasmic density that likely involve a sizeable replacement of water within the cytoplasm by protein biomass. Nevertheless, *S. pombe* cells retain viability, grow with normal morphologies, and exhibit normal periods of mitosis and cytokinesis, underscoring the robust control of critical biochemical processes. In particular, this work provides a striking demonstration of how cell size can be maintained even over a 10-fold range of growth rates, a finding inconsistent with models of cell-size control based directly on growth rate [42]. Rather, these results support the existence of cell-size control mechanisms in fission yeast cells that involve direct assessment of cell size through "sizers" [43,44]. In general, our findings indicate that cellular density can be regulated or perturbed in various settings such as stress. Procedures such as osmotic oscillations that increase protein expression per unit volume may have practical applications, for instance in protein production. It will be valuable to explore in future studies the myriad ways in which such global changes in cytoplasmic density affect cellular processes.

METHODS

Strains and growth conditions

All strains used in this study are listed in Table S1.

Single colonies were inoculated from plates into liquid rich medium (YE5S, Sunrise Science Products YES-225) overnight with shaking at 30 °C. Once cells reached mid-log phase ($OD_{600}\sim0.5$), they were diluted 1:10 directly into ONIX Y04C-02 microfluidic flow cells (EMD Millipore) that had been primed with fresh medium for 15 min. Cells were grown for 15-30 min prior to imaging to allow them to equilibrate to the flow-cell chamber. YE5S was supplemented with D-sorbitol (Sigma) for hypertonic conditions, and fresh medium was introduced into the culture chambers at 5 psi. Control measurements were performed by exchanging YE5S between wells in adjacent culture chambers. By adding a tracer dye (0.5 μg/mL Alexa Fluor 647 carboxylic acid; Life Technologies) to the sorbitol-supplemented medium, we determined that media exchange occurred within 10 s (Figure 2A, S3A). Where applicable, drug (cycloheximide or brefeldin A; Sigma) was added to the medium.

Microscopy

Cells were imaged in phase-contrast using a Ti-Eclipse stand (Nikon Instruments) with a 40X (NA: 0.95) or 100X (NA: 1.45) objective, and a Zyla 4.2 sCMOS camera (Andor Technology). Temperature was maintained with a stage-top incubator (OkoLab), which was warmed for at least 1 h prior to imaging. Images were acquired at 1-s or 5-s frame intervals for measuring rapid volume changes during osmotic shocks (Figure S3B), and 1-min or 5-min frame intervals for osmotic shock and growth measurements. For strains expressing fluorescent proteins, cells were imaged in phase-contrast and with laser illumination in spinning-disk confocal mode (Yokogawa CSU-10), and images were acquired with an EM-CCD camera (Hamamatsu) every 10 min at low power to avoid bleaching. The microscope systems were integrated using μManager v. 1.41 [45].

Cell segmentation and size/fluorescence quantification

We first utilized the deep neural network-based machine learning framework *DeepCell* (Van Valen et al., 2017) to segment cells. For each imaging technique (wide-field, confocal) and

441 objective, approximately 200 cells were manually outlined to produce a training dataset. Trained
442 networks were used to generate binary images for feature (extracellular/cell
443 perimeter/cytoplasm) identification. These images were used as the input for gradient
444 segmentation in *Morphometrics* v. 1.1 [46] to define cell contours at sub-pixel resolution.
445
The centerline between the two poles was calculated through an iterative method in which
symmetric bisections were created starting from the contour's centroid toward the poles. Cell
length was defined as the total length of the centerline, and width as the median length of lines
running perpendicular to the centerline and stretching between the two sides of the cell. Cell
surface area and volume were calculated by integrating disks of revolution at each point on the
centerline.
452
453 For fluorescence quantification, the total signal within each cell contour was summed.
454
455 ***Estimation of supergrowth contribution to mCrimson dilution***
456 Estimates for growth-based dilution during supergrowth were calculated using the equation
457

$$c_{\text{dilution}}(t) = c_0 + (c_{\max} - c_0)e^{-\int_0^{t_1} \frac{1}{V}\frac{dV}{dt}dt},$$

459
460 where the rate of protein production during supergrowth is assumed to follow the growth rate
461 and $\frac{1}{V}\frac{dV}{dt}$ is the population-averaged growth rate. The difference between the actual and
462 estimated concentrations $(c(t) - c_{\text{dilution}}(t))$ was used to estimate the contribution to dilution
463 from growth at >75%.
464
Bgs4 tip localization and intensity quantification
Cell tips were identified based on peaks in the contour curvature, which is defined at each
contour point as the second derivative of the local vector normal. The fluorescence intensity was
integrated along a line scan perpendicular to the contour at each contour point whose curvature
was greater than 0.12 μm^{-1}, a threshold that was determined visually as an optimal identifier of
cell tip regions.

Osmotic oscillations on bulk cultures

Using a tabletop upright glass vacuum filter holder (Fisherbrand) with 0.45-µm HA membrane

filters (Millipore, HAWP04700), the medium was rapidly exchanged (<30 s) for cultures of up to

200 mL. All media were maintained at 30 °C, and cells were resuspended and shaken at 30 °C

during each half-period. One milliliter samples were taken approximately each hour during the

experiment and immediately fixed in 70% cold ethanol. Cells from the culture were imaged

directly after these oscillations to measure growth rate and mCrimson fluorescence.

Fluorescein isothiocyanate (FITC) and propidium iodide (PI) staining

Staining protocols were adapted from [47]. One-milliliter samples were collected at various time

points during the bulk oscillations protocol at $OD_{600} \sim 0.5$, and cells were fixed with 70% cold

ethanol for at least 24 h at 4 °C. Approximately 10^6 cells (300 µL) were washed with 50 mM

sodium citrate buffer (pH 7.4), and then treated with 0.1 mg/mL RNAse A (ThermoFisher,

EN0531) for at least 2 h. For total protein staining, cells were washed and then stained with 5

ng/mL FITC (Sigma, F7250) for at least 30 min, then washed three times. To determine total DNA

content, cells were stained with PI at 0.1 mg/mL for at least 30 min and then washed.

To quantify FITC/PI ratios, cells were imaged on 1% agarose + EMM pads (Figure S8). *Z*-stacks

were acquired to obtain sum-projections after background subtraction, with cell outlines

obtained manually due to the poor phase contrast of fixed cells.

Quantitative Phase Imaging

Images were acquired with a Ti-Eclipse Ti inverted microscope (Nikon) equipped with a 680-nm

bandpass filter (D680/30, Chroma Technology) in the illumination path with a 60X (NA: 1.4) DIC

oil objective (Nikon). Before imaging, Koehler illumination was configured and the peak

illumination intensity with 10-ms exposure time was set to the middle of the dynamic range of

the Zyla sCMOS 4.2 camera (Andor). µManager v. 1.41 [45] was used to automate acquisition of

brightfield *z*-stacks with a step size of 250 nm from ±3 µm around the focal plane (total of 25

imaging planes) at 10-min intervals.

For analysis, all imaging planes between maximal distances of +1.25 μm and -1.25 μm around the
focal plane were selected. Based on these images, the phase information was calculated using a
custom Matlab script implementing a previously published algorithm [48]. In brief, this method
relates the phase information of the cell to intensity changes along the *z*-direction. Equidistant,
out-of-focus images above and below the focal plane are used to estimate intensity changes at
various defocus distances. A phase-shift map is reconstructed in a non-linear, iterative fashion to
solve the transport-of-intensity equation.
509
A Gaussian peak was fitted to the background of each image and then corrected to be at zero
phase shift using an image-wide subtraction of the mean of the peak. These corrected images
were segmented using *DeepCell* and *Morphometrics* v. 1.1. From the cell outlines, the median
intensity of each cell was obtained with ImageJ and used to calculate the cytoplasmic density as
follows. The difference of the reconstructed phase map median intensity of cells in YE5S medium
compared to cells in YE5S medium supplemented with 100 mg/mL BSA was used to define the
phase shift contribution equivalent to 100 mg/mL biomass. The phase shift corresponding to
100 mg/mL BSA was then calculated by first averaging the intensity of all cells in YE5S medium
at timepoints 10 min before and after BSA addition. Then, the difference relative to the median
intensity of all cells during the measurement in YE5S supplemented with 100 mg/mL BSA was
defined as the signal contributed by 100 mg/mL biomass.
521
522 Throughout the experiment, multiple measurements in YE5S supplemented with 100 mg/mL
523 BSA were performed and calibration values for each imaging timepoint in between two
524 consecutive measurements in YE5S + BSA were obtained by linear interpolation. These values
525 were then used for quantification of cell density.

AUTHOR CONTRIBUTIONS

B.D.K., K.C.H., and F.C. conceptualized the study. B.D.K., P.O., E.R.R., X.H., K.C.H., and F.C. designed the experiments. B.D.K. and P.O. performed oscillatory osmotic shock experiments. P.O. performed quantitative phase imaging. X.X. and W.C. constructed ribosomal-GFP strains. B.D.K., P.O., E.R.R., K.C.H., and F.C. analyzed data. B.K., K.C.H., and F.C. wrote the manuscript. All authors reviewed the manuscript before submission.

ACKNOWLEDGMENTS

We thank members of the Huang, Chang, J. Skotheim, and S. Dumont labs for helpful discussions. This work was funded by NIH CAREER Award MCB-1149328 (to K.C.H.), the Allen Discovery Center at Stanford University on Systems Modeling of Infection (to K.C.H.), the National Natural Science Foundation of China (NSFC; 3167080548) (to X.H.), and NIH R01-GM56836 and Bilateral BBSRC-NSF/BIO Award 1638195 (to F.C.). K.C.H. is a Chan Zuckerberg Biohub Investigator.

    FIGURE LEGENDS

Figure 1: *S. pombe* cells grow exponentially.

A) Wild-type (WT) fission yeast cells imaged in phase contrast. Green outlines were obtained via automated segmentation (Methods). Manual inspection revealed that segmentation was accurate for virtually all cells, regardless of cell crowding.

B) Population-averaged length and width of growing fission yeast cells, measured as a function of time after the mechanical separation of daughter cells after septation normalized by cell-cycle time (n = 1129 cells). Solid lines and shaded areas represent the mean ± 1 standard deviation (S.D.). Cell width was essentially invariant, while

length increased over the cell cycle until the cell reaches the division length of ~14

μm.

C) Instantaneous growth rate dL/dt increased with cell length. The trend for wild-type

(WT) cells (blue) continued for *cdc25-22* cells (red), which are substantially longer.

Black lines represent best linear fits for each strain (wildtype, *n*=1129 cells; *cdc25-22*,

n=246 cells).

D) Instantaneous size-normalized growth rate $\left(\frac{1}{L}\,dL/dT\right)$ of wild-type cells, as a function

of normalized cell-cycle time as in (B). The size-normalized growth rate was constant

for a large fraction of the cell cycle, signifying growth proportional to size (exponential

growth). Cells had three phases of growth: (I) super-exponential acceleration, (II)

steady-state exponential growth, and (III) deceleration as cells approach mitosis.

Figure 2: Exit from osmotic shock oscillations induces robust supergrowth.

A) Cells were equilibrated to achieve steady-state growth in YE5S, and then exposed to cycles of hyperosmotic and hypoosmotic shocks with 500 mM sorbitol, with 5 min between shocks.

B) Length of a representative wild-type (WT) cell increased more rapidly during periods in YE5S+500 mM sorbitol, but returned to the value expected of growth in YE5S after

570   hypoosmotic shocks. Width increased in YE5S+500 mM sorbitol, and was maintained at

571   an approximately constant level in YE5S.

C)   Length (top) and instantaneous growth rates (bottom) of wild-type cells (n = 45) before,
during (yellow shading), and after 24 cycles of 0.5-M sorbitol osmotic shock oscillations
with 10-min period. Cells are the same color in both plots. Cells were imaged every 5 min.
After the oscillations, cells exhibited faster growth (supergrowth) for multiple
generations. Traces that begin at later times points are newly formed daughter cells.

D)   Size growth rates of wild-type cells throughout an oscillatory osmotic shock experiment
(red, n = 973 cells). Growth rate during oscillations (yellow, dashed box) was lower than
the control that was kept in YE5S throughout (blue, n = 1242). Growth rate increased
after cells exited the oscillations, decreasing back to the control growth rate with a time
constant of 1.3 h.

E)   Cell-size control was unaffected by oscillations or supergrowth, with constant length (top)
and width (bottom) at the time of division throughout the experiment in (C).

F)   Maximal size-normalized growth rates during supergrowth as a function of (i) oscillatory
shock period length, (ii) number of periods, and (iii) sorbitol concentration (n = 46-536
cells per data point). Each parameter was varied while the other two parameters were
held fixed. Red circles represent our standard conditions of 24 cycles of 500-mM shocks
with 10-min period. Error bars represent one standard deviation.

Figure 3: Osmotic-shock oscillations disrupt coupling between protein synthesis and cell size.

A) Representative time-lapse images of mCrimson-expressing cells before, during, and after 24 10-min periods of 0.5-M sorbitol osmotic shocks. Fluorescence intensity

increased during oscillations, and then gradually reverted to pre-shock levels during
supergrowth.
B) Representative trajectories of integrated cellular fluorescence before, during, and
after oscillations (n =54 cells). Synthesis of mCrimson continued relatively unabated
during oscillations. The yellow box in this and subsequent panels denotes the 24 10-
min periods of 0.5-M sorbitol osmotic shocks. (n = 54 cells)
C) mCrimson concentration increases by ~60% during oscillations (red), despite large
decreases in volume growth rate (blue) (n = 843 cells). During supergrowth, the
increased growth rate led to dilution of the mCrimson concentration. Dark centerline
is the population-averaged mean, and the shading represents one standard deviation.
D) Concentrations of various proteins labeled with fluorescent proteins before, during,
and after oscillations. All protein concentrations increased during oscillations, but to
varying degrees.
E) Ribosomal proteins also increased during oscillations, but to a lesser degree; RPS802
concentration remained approximately constant throughout the experiment.
F) The ratio of FITC (proteins) to PI (DNA) fluorescence, a proxy for global protein
density, increased during oscillations and then decreased back to steady-state levels
during supergrowth. Cells were subjected to 12 cycles of 500-mM sorbitol shocks with
10-min period (Methods). Error bars are one standard deviation (n = 200 cells each
datapoint).
G) Cytoplasmic density, as measured by quantitative phase imaging (Methods), increased
during osmotic shocks and decreased during supergrowth. Dark centerline is the
population-averaged mean cell density, and the shading represents one standard
deviation. Green: normal growth (n = 88 cells); black, 500-mM oscillations (n = 187
cells); purple, 1-M oscillations (n = 106 cells).
620

Figure 4: Excess protein concentration is necessary for supergrowth.

 A) Maximum size-normalized growth rates were highly correlated with the mean

   mCrimson concentration at exit from oscillations, across all shock magnitudes,

   periods, and number of shock cycles in Figure 2F. Error bars represent one standard

   deviation.

 B) Cells treated with 100 μg/mL of the translation inhibitor cycloheximide (CHX; black

   shading, n = 228 cells) during oscillations (24 10-min periods, 0.5-M sorbitol shocks;

   yellow shading) rapidly halted growth during oscillations and exhibited no

   supergrowth afterward, unlike untreated cells (blue, n = 545 cells). Shading

   represents one standard deviation.

 C) Cells exposed to osmotic-shock oscillations as in (B) (black, n = 175 cells) or

   maintained in constant YE5S growth conditions (blue, n = 260 cells) were treated

   (directly after the exit from oscillations) for 3 h with 100 μg/mL cycloheximide. Cells

that underwent oscillations grew more during CHX treatment and resumed normal

growth more quickly after CHX washout. Shading represents one standard deviation.

D) During cycloheximide treatment, the ratio of growth rates of cells that did and did not

undergo oscillations was always ≥1 and reached a plateau within 30 min. At the

plateau, cells that underwent oscillations grew approximately three-fold faster than

control cells. The growth-rate ratio remained >1 for 2-3 h after cycloheximide

treatment.

E) Brefeldin A (green shading) treatment completely inhibited growth, while mCrimson

continued to accumulate throughout treatment. After the drug was washed out, cells

initiated supergrowth with dynamics similar to those achieved after osmotic shocks.

REFERENCES

Newman, J. R. *et al.* Single-cell proteomic analysis of S. cerevisiae reveals the architecture of biological noise. *Nature* **441**, 840-846, doi:10.1038/nature04785 (2006).

Schmoller, K. M., Turner, J. J., Koivomagi, M. & Skotheim, J. M. Dilution of the cell cycle inhibitor Whi5 controls budding-yeast cell size. *Nature* **526**, 268-272, doi:10.1038/nature14908 (2015).

Zhou, X. *et al.* Bacterial division. Mechanical crack propagation drives millisecond daughter cell separation in Staphylococcus aureus. *Science* **348**, 574-578, doi:10.1126/science.aaa1511 (2015).

You, C. *et al.* Coordination of bacterial proteome with metabolism by cyclic AMP signalling. *Nature* **500**, 301-306, doi:10.1038/nature12446 (2013).

Padovan-Merhar, O. *et al.* Single mammalian cells compensate for differences in cellular volume and DNA copy number through independent global transcriptional mechanisms. *Mol Cell* **58**, 339-352, doi:10.1016/j.molcel.2015.03.005 (2015).

Willis, L. & Huang, K. C. Sizing up the bacterial cell cycle. *Nat Rev Microbiol* **15**, 606-620, doi:10.1038/nrmicro.2017.79 (2017).

Tzur, A., Kafri, R., LeBleu, V. S., Lahav, G. & Kirschner, M. W. Cell growth and size homeostasis in proliferating animal cells. *Science* **325**, 167-171, doi:10.1126/science.1174294 (2009).

Abenza, J. F. *et al.* Wall mechanics and exocytosis define the shape of growth domains in fission yeast. *Nat Commun* **6**, 8400, doi:10.1038/ncomms9400 (2015).

Chang, F. & Martin, S. G. Shaping fission yeast with microtubules. *Cold Spring Harb Perspect Biol* **1**, a001347, doi:10.1101/cshperspect.a001347 (2009).

Atilgan, E., Magidson, V., Khodjakov, A. & Chang, F. Morphogenesis of the Fission Yeast Cell through Cell Wall Expansion. *Curr Biol* **25**, 2150-2157, doi:10.1016/j.cub.2015.06.059 (2015).

Mitchison, J. M. & Nurse, P. Growth in cell length in the fission yeast Schizosaccharomyces pombe. *J Cell Sci* **75**, 357-376 (1985).

12 Baumgartner, S. & Tolic-Norrelykke, I. M. Growth pattern of single fission yeast cells is
bilinear and depends on temperature and DNA synthesis. *Biophys J* **96**, 4336-4347,
doi:10.1016/j.bpj.2009.02.051 (2009).

13 Sveiczer, A., Novak, B. & Mitchison, J. M. The size control of fission yeast revisited. *J Cell Sci*
109 (Pt 12), 2947-2957 (1996).

14 Cooper, S. Distinguishing between linear and exponential cell growth during the division
cycle: single-cell studies, cell-culture studies, and the object of cell-cycle research. *Theor*
Biol Med Model **3**, 10, doi:10.1186/1742-4682-3-10 (2006).

15 Horvath, A., Racz-Monus, A., Buchwald, P. & Sveiczer, A. Cell length growth patterns in
fission yeast reveal a novel size control mechanism operating in late G2 phase. *Biol Cell*
108, 259-277, doi:10.1111/boc.201500066 (2016).

16 Horvath, A., Racz-Monus, A., Buchwald, P. & Sveiczer, A. Cell length growth in fission yeast:
an analysis of its bilinear character and the nature of its rate change transition. *FEMS*
Yeast Res **13**, 635-649, doi:10.1111/1567-1364.12064 (2013).

17 Chang, F., Atilgan, E., Burgess, D. & Minc, N. Manipulating cell shape by placing cells into
micro-fabricated chambers. *Methods Mol Biol* **1136**, 281-290, doi:10.1007/978-1-4939-
0329-0_13 (2014).

18 Basu, R., Munteanu, E. L. & Chang, F. Role of turgor pressure in endocytosis in fission
yeast. *Mol Biol Cell* **25**, 679-687, doi:10.1091/mbc.E13-10-0618 (2014).

19 Minc, N., Boudaoud, A. & Chang, F. Mechanical forces of fission yeast growth. *Curr Biol* **19**,
1096-1101, doi:10.1016/j.cub.2009.05.031 (2009).

20 Proctor, S. A., Minc, N., Boudaoud, A. & Chang, F. Contributions of turgor pressure, the
contractile ring, and septum assembly to forces in cytokinesis in fission yeast. *Curr Biol*
22, 1601-1608, doi:10.1016/j.cub.2012.06.042 (2012).

21 Rojas, E., Theriot, J. A. & Huang, K. C. Response of Escherichia coli growth rate to osmotic
shock. *Proc Natl Acad Sci U S A* **111**, 7807-7812, doi:10.1073/pnas.1402591111 (2014).

22 Chen, D. *et al.* Global transcriptional responses of fission yeast to environmental stress.
Mol Biol Cell **14**, 214-229, doi:10.1091/mbc.E02-08-0499 (2003).

23 Sanchez-Mir, L. *et al.* Rho1 GTPase and PKC Ortholog Pck1 Are Upstream Activators of the
Cell Integrity MAPK Pathway in Fission Yeast. *Plos One* **9**, doi:ARTN e88020
10.1371/journal.pone.0088020 (2014).

24   Ma, Y. *et al.* Transient receptor potential (TRP) and Cch1-Yam8 channels play key roles in
the regulation of cytoplasmic Ca2+ in fission yeast. *PLoS One* **6**, e22421,
doi:10.1371/journal.pone.0022421 (2011).

25   Davie, E., Forte, G. M. & Petersen, J. Nitrogen regulates AMPK to control TORC1 signaling.
Curr Biol **25**, 445-454, doi:10.1016/j.cub.2014.12.034 (2015).

26   Schutt, K. L. & Moseley, J. B. Transient activation of fission yeast AMPK is required for cell
proliferation during osmotic stress. *Molecular Biology of the Cell* **28**, 1804-1814,
doi:10.1091/mbc.E17-04-0235 (2017).

27   Ikeda, K., Morigasaki, S., Tatebe, H., Tamanoi, F. & Shiozaki, K. Fission yeast TOR complex 2
activates the AGC-family Gad8 kinase essential for stress resistance and cell cycle control.
Cell Cycle **7**, 358-364, doi:10.4161/cc.7.3.5245 (2008).

28   Al-Sady, B., Greenstein, R. A., El-Samad, H. J., Braun, S. & Madhani, H. D. Sensitive and
Quantitative Three-Color Protein Imaging in Fission Yeast Using Spectrally Diverse,
Recoded Fluorescent Proteins with Experimentally-Characterized In Vivo Maturation
Kinetics. *Plos One* **11**, doi:ARTN e0159292
10.1371/journal.pone.0159292 (2016).

29   Waldron, C. & Lacroute, F. Effect of Growth-Rate on Amounts of Ribosomal and Transfer
Ribonucleic-Acids in Yeast. *J Bacteriol* **122**, 855-865 (1975).

30   Polanshek, M. M. Effects of heat shock and cycloheximide on growth and division of the
fission yeast, Schizosaccharomyces pombe. With an Appendix. Estimation of division
delay for S. pombe from cell plate index curves. *J Cell Sci* **23**, 1-23 (1977).

31   Cortes, J. C. *et al.* The novel fission yeast (1,3)beta-D-glucan synthase catalytic subunit
Bgs4p is essential during both cytokinesis and polarized growth. *J Cell Sci* **118**, 157-174,
doi:10.1242/jcs.01585 (2005).

32   Mutavchiev, D. R., Leda, M. & Sawin, K. E. Remodeling of the Fission Yeast Cdc42 Cell-
Polarity Module via the Sty1 p38 Stress-Activated Protein Kinase Pathway. *Curr Biol* **26**,
2921-2928, doi:10.1016/j.cub.2016.08.048 (2016).

33   Haupt, A., Ershov, D. & Minc, N. A Positive Feedback between Growth and Polarity
Provides Directional Persistency and Flexibility to the Process of Tip Growth. *Current*
Biology **28**, 3342-+, doi:10.1016/j.cub.2018.09.022 (2018).

34 Novick, P., Field, C. & Schekman, R. Identification of 23 complementation groups required
for post-translational events in the yeast secretory pathway. *Cell* **21**, 205-215 (1980).
35 Brazer, S. C., Williams, H. P., Chappell, T. G. & Cande, W. Z. A fission yeast kinesin affects
Golgi membrane recycling. *Yeast* **16**, 149-166, doi:10.1002/(SICI)1097-
0061(20000130)16:2<149::AID-YEA514>3.0.CO;2-C (2000).
36 Lippincottschwartz, J., Yuan, L. C., Bonifacino, J. S. & Klausner, R. D. Rapid Redistribution of
Golgi Proteins into the Er in Cells Treated with Brefeldin-a - Evidence for Membrane
Cycling from Golgi to Er. *Cell* **56**, 801-813, doi:Doi 10.1016/0092-8674(89)90685-5
(1989).
37 Cleveland, D. & Kirschner, M. Autoregulatory control of the expression of alpha- and beta-
tubulins: implications for microtubule assembly. *Cold Spring Harb Symp Quant Biol* **46 Pt
1**, 171-183 (1982).
38 Springer, M., Weissman, J. S. & Kirschner, M. W. A general lack of compensation for gene
dosage in yeast. *Mol Syst Biol* **6**, 368, doi:10.1038/msb.2010.19 (2010).
39 Torres, E. M., Springer, M. & Amon, A. No current evidence for widespread dosage
compensation in S. cerevisiae. *Elife* **5**, e10996, doi:10.7554/eLife.10996 (2016).
40 Torres, E. M. *et al.* Effects of aneuploidy on cellular physiology and cell division in haploid
yeast. *Science* **317**, 916-924, doi:10.1126/science.1142210 (2007).
41 Zhao, Y., Sohn, J. H. & Warner, J. R. Autoregulation in the biosynthesis of ribosomes. *Mol
Cell Biol* **23**, 699-707 (2003).
42 Lucena, R. *et al.* Cell Size and Growth Rate Are Modulated by TORC2-Dependent Signals.
Curr Biol **28**, 196-210 e194, doi:10.1016/j.cub.2017.11.069 (2018).
43 Pan, K. Z., Saunders, T. E., Flor-Parra, I., Howard, M. & Chang, F. Cortical regulation of cell
size by a sizer cdr2p. *Elife* **3**, e02040, doi:10.7554/eLife.02040 (2014).
44 Facchetti, G., Chang, F. & Howard, M. Controlling cell size through sizer mechanisms.
Current Opinion in Systems Biology **5**, 86-92,
doi:https://doi.org/10.1016/j.coisb.2017.08.010 (2017).
45 Edelstein, A. D. *et al.* Advanced methods of microscope control using µManager software.
Journal of biological methods **1**, e10 (2014).

46   Ursell, T. *et al.* Rapid, precise quantification of bacterial cellular dimensions across a

genomic-scale knockout library. *BMC Biol* **15**, 17, doi:10.1186/s12915-017-0348-8

(2017).

47   Porro, D., Brambilla, L. & Alberghina, L. Glucose metabolism and cell size in continuous

cultures of Saccharomyces cerevisiae. *FEMS Microbiol Lett* **229**, 165-171 (2003).

48   Bostan, E., Froustey, E., Nilchian, M., Sage, D. & Unser, M. Variational Phase Imaging Using

the Transport-of-Intensity Equation. *IEEE Trans Image Process* **25**, 807-817,

doi:10.1109/TIP.2015.2509249 (2016).