Long title: Host evolutionary history predicts virus prevalence across bumblebee

species

Short title: Evolutionary signals in bumblebee-virus interaction networks

David J. Pascall[1,6], Matthew C. Tinsley[2], Darren J. Obbard[3,4] and Lena Wilfert[1,5]

[1] College of Life and Environmental Science, University of Exeter Cornwall Campus,

Treliever Road, Penryn, UK

[2]

[3] Institute of Evolutionary Biology, University of Edinburgh, Charlotte Auerbach

Road, Edinburgh, UK

[4] Centre for Infection, Evolution and Immunity, University of Edinburgh, Charlotte

Auerbach Road, Edinburgh, UK

[5] Institute of Evolutionary Ecology and Conservation Genomics, University of Ulm,

Albert-Einstein-Allee 11, 89069 Ulm, Germany

[6] Institute of Biodiversity, Animal Health and Comparative Medicine, Graham Kerr

Building, University of Glasgow, Glasgow, UK

Author Contributions

DJP – Conceptualization, Data Curation, Formal Analysis, Investigation,

Methodology, Project Administration, Software, Visualization, Writing – Original

Draft Preparation, Writing – Review & Editing

MCT – Investigation, Resources, Writing – Review & Editing

DJO – Conceptualization, Data Curation, Formal Analysis, Funding Acquisition,

Methodology, Project Administration, Software, Visualisation, Writing – Review &

Editing

LW – Conceptualization, Funding Acquisition, Investigation, Methodology, Project

Administration, Resources, Supervision, Writing – Review & Editing

Author's Summary

Despite the importance of disease in the regulation of animal populations, our

understanding of the distribution of pathogen burden across wild communities

remains in its infancy. In this study, we investigated the distribution of viruses across

natural populations of 13 different bumblebee species in Scotland. In order to

accurately assess this distribution, we first searched for viruses using a transcriptomic

approach, finding at least 30 new viruses of bumblebees, and assayed a subset of them

for their presence and absence in different species. Then, in the first application of

these methods to an animal-virus system, we used co-phylogenetic mixed models to

investigate the factors that lead to species being infected to different degrees by a

subset of these viruses. We found that, while much of the variation in the prevalence

of the viruses can be explained by the specifics of individual bumblebee-virus

pairings, related bumblebee species being infected to similar degrees with the same

sets of viruses has an important contribution to the distribution of viruses across hosts.

Consistent with previous work, our study indicates that, while in general the

interaction between a host and a virus may be unpredictable, closely related species

are more likely to exhibit similar patterns.

Abstract

Why a pathogen associates with one host but not another is one of the most important

questions in disease ecology. Here we use transcriptome sequencing of wild-caught

bumblebees from 13 species to describe their natural viruses, and to quantify the

impact of evolutionary history on the realised associations between viruses and their

pollinator hosts. We present 37 novel virus sequences representing at least 30

different viruses associated with bumblebees. We verified 17 of them by PCR and

estimate their prevalence across species in the wild. Through small RNA sequencing,

we demonstrate that at least 10 of these viruses form active infections in wild

individuals. Using a phylogenetic mixed model approach, we show that the

evolutionary history of the host shapes the current distribution of virus/bumblebee

associations. Specifically, we find that related hosts share viral assemblages, viruses

differ in their prevalence averaged across hosts and the prevalence of infection in

individual virus-host pairings depends on precise characteristics of that pairing.

Introduction

Pathogens that naturally infect more than one host species have a particularly high

risk of disease emergence (Woolhouse & Gowtage-Sequeria 2005). One especially

important group of pathogens are the viruses, whose ubiquity leads them to have a

disproportionate role in the regulation of natural populations (Suttle 2007). Viruses

are relevant in populations that humans manage for economic and conservation

reasons, such as bumblebees, which are both in decline (Williams & Osborne 2009)

and important providers of ecosystem services (Garibaldi et al. 2013).

Bumblebees, genus *Bombus*, are a primitively eusocial group of important wild

pollinators; many bumblebee species have experienced population declines, linked to

biotic and abiotic stressors such as habitat degradation, pesticide use and shared

infectious diseases for example caused by viral pathogens (Vanbergen & the Insect

Pollinators Initiative 2013). While honeybee viruses have been intensively studied,

and have in many cases been found to represent multihost pathogens (see Manley et

al. (2015) and the references within), bumblebee-specific viruses are comparatively

poorly studied, and how widely they are shared between species is unknown.

In order for a species to be a multihost pathogen, some degree of opportunity for

cross-species transmission must exist. Our definition of multihost pathogens follows

that of Fenton et al. (2015). As such, multihost pathogens are defined to include two

conceptually distinct groups: 'facultative multihost pathogens' that are able to

maintain transmission chains in multiple host species (i.e. $R_0 >= 1$ in multiple host

species) and 'obligate multihost pathogens', which rely on sufficiently high rates of

cross-species transmission to offset unsustainable transmission within individual host

species (i.e. $0 < R_0 < 1$ within host species, $R_0 >= 1$ overall). In addition, pathogens that

maintain transmission in a single host ($R_0 >= 1$) but experience regular spillover (with

or without the expectation of onward transmission: $0 <= R_0 < 1$) are included as being

effectively multihost pathogens within our definition. R_0 is defined as the expected

number of secondary infections caused by a single typical infected individual in an

entirely naïve host population (Heesterbeek 2002). We define cross-species

transmission as the movement of a multihost pathogen between host species within its

host range. This contrasts with host shifting, which we define as a transmission event

to a new host species, leading to a change in host range; however there is necessarily

some unavoidable ambiguity between cross-species transmission and host shifting in

the case of pathogens that exhibit rare spillover events.

The opportunity for cross-species transmission, which explains the large number of

viruses originally detected in honeybees present in bumblebees, may be created by

niche overlap in foraging (Salathé & Schmid-Hempel 2011). Bumblebee nests are

provisioned by foraging workers who gather pollen and nectar from flowers in the

surrounding area. Considerable interspecific differences in plant species utilization by

foragers of different species are commonly observed (e.g. Arbulo, Santos, Salvarrey

& Invernizzi 2011; Goulson & Darvill 2004; Goulson et al. 2008; Harder 1985), but

this is not a universal phenomenon (Lye et al. 2010), and the degree of overlap may

depend on the diversity of flowers currently in bloom. Flower choice of foragers is

correlated with species tongue length (Goulson et al. 2008; Harder 1985), which

implicitly incorporates shared behavioural characteristics between closely related

bumblebee species as there is phylogenetic correlation between tongue length and

relatedness (Harmon-Threatt & Ackerly 2013). Different species of bumblebee also

exhibit incomplete temporal separation throughout the year, causing some degree of

partitioning in niche space even when they are spatially sympatric (Goodwin 1995).

This ecology leads to a complex interaction network between bumblebee species as

well as sympatric honeybees, which may structure cross-species transmission.

The prevalence of pathogens, including viruses, across host species, such as

bumblebees, is structured on two levels. First, a virus may be present or entirely

absent in a potential host species. Second, other factors may then influence how

prevalent a pathogen is within that species. At the presence/absence level, a complete

lack of infection in nature can occur in three ways: 1) a host and virus may exist in

allopatry or in completely non-interacting ecological niches, preventing transmission

irrespective of the host's susceptibility; 2) a physiological or molecular mismatch

(including immunity) between a host and virus can prevent infection; and 3)

environmental conditions may be such that transmission cannot occur between two

sympatric species. None of these mechanisms represent an immutable barrier, and all

represent ends of a continuum, where lesser forms simply reduce transmission.

Spatially or ecologically separated hosts and parasites may come into contact through

migrations or human facilitated invasions, allowing new associations to emerge. For

example, the arrival of *Plasmodium relictum* to the Hawaiian islands led to avian

population declines and contributed to extinctions in the naturally susceptible but

naïve populations (van Riper et al. 1986). Incompatibility can break down if evolution

in the pathogen or host removes the physiological or molecular barriers to infection,

as shown when Canine parvovirus type 2 emerged from Feline panleukopenia virus

after gaining the ability to bind to canine transferrin receptors (Hueffer et al. 2003).

For virus-host associations where infection can and does occur, quantitative

differences in infection risk between species can be driven by ecological variation in

transmission rates. These differences can be driven by, for example, the propensity for

group living (Johnson et al. 2011), population densities (Arneberg et al. 1998), the

biodiversity of the community (Civitello et al. 2015) and host avoidance behaviours

(Curtis 2014). Variation in infection risk among host species can also be driven by

physiological and molecular factors, with hosts having varying suitability for the

replication of a given parasite. In the extreme case, a host species may exhibit

condition-dependent susceptibility; where infection can only occur when the immune

system is suppressed, either directly, through an immunosuppressant disease or

chemical agent, or indirectly, through trade-offs in resource allocation brought about

by malnutrition (Chandra 1983). Both behavioral and ecological factors, leading to

differences in contact rate, and physiological factors, leading to differences in

infection probability on contact, may be phylogenetically correlated (Harmon-Threatt

& Ackerly 2013; Longdon et al. 2011).

Box 1 – Definition of Terms

Co-phylogenetic generalized linear mixed models that incorporate phylogenetic

variance from multiple clades (Hadfield et al. 2014; Rafferty & Ives 2013) have been

used relatively rarely, and a biological interpretation of the model terms may not be

immediately familiar. In the host-parasite context, this approach can be used to

model how the probability of infection is predicted by both host and parasite species,

allowing for covariance induced by the relationships within each group, and the

interactions between these model terms. This can be considered either at the species-

wide level (i.e. the probability that infection will occur at all in a given host/parasite

pairing), or at the level of individuals within species (i.e. infection prevalence). Here

we provide verbal descriptions of how the terms can be interpreted, as well as

references to a figure in Hadfield et al. (2014) where each of these effects is

illustrated graphically:

Phylogenetic Effect: Variation in the mean value of a trait among species that is

explained by phylogenetic divergence. For example, more closely-related hosts might

be more similar in susceptibility to viral infection (display higher viral prevalence),

irrespective of virus species. Equivalently, more closely-related viruses might be

more similar in infectiousness, irrespective of host species (Figures 1a and b in

Hadfield et al. (2014)).

Species Effect: Variation in the mean value of a trait among species that is

not explained by a *Phylogenetic Effect*. For example, much of the variation in

prevalence among viral species (irrespective of host), may not be explained by the

virus phylogeny but instead depend on lineage-specific viral traits (Figures 1g and f in

Hadfield et al. (2014)).

Non-phylogenetic Interaction: The interaction term between host and parasite *Species*

Effects, such that variation in the mean value of a trait within particular host/parasite

pairings depends the specifics of the host and parasite involved in a way not affected

by their evolutionary divergence. For example, variation in prevalence between

particular pairings that is caused by the interaction between lineage-specific host and

viral traits (Figure 1h in Hadfield et al. (2014)).

Coevolutionary Interaction: The interaction term between host and parasite

Phylogenetic Effects, such that variation in the mean value of a trait within particular

host/parasite pairings depends on the evolutionary divergence among species in both

host and parasite clades. For example, if the prevalence of infection is more similar

among pairings of closely-related hosts and closely-related parasites than would be

expected from the host and parasite phylogenies and species-means alone. (Figure 1e

in Hadfield et al. (2014)).

Evolutionary Interaction: The interaction term between the host (or parasite)

Phylogenetic Effect and the partners' *Species Effect*, such that the variation in the

mean value of a trait within particular host/parasite pairings depends on the

evolutionary divergence among hosts (or parasites) and on the identity of particular

partner species, but is not predicted by the evolutionary divergence between partners

species. For example, if the similarity in viral prevalence for one virus species is

strongly predicted by the evolutionary divergence among hosts, but a completely

different relationship (unrelated to the evolutionary divergence among viruses) is seen

for other virus species. (Figure 1c and d in Hadfield et al. (2014)).

A new multihost virus can arise in two ways, either through a virus species gaining a

new host (Longdon et al. 2014) (Figure 1a), or through a speciation event in a

multihost virus (Figure 1b). When novel multihost viruses are generated through host

shifting (Figure 1a), a 'host evolutionary interaction' effect (see Box1) can result, as

the consistent switching of viruses (V) to hosts (H) closely related to their ancestral

host will lead to related hosts having correlated viral assemblages. When novel

multihost viruses arise through speciation, i.e. if the ability to infect multiple hosts is

an ancestral trait (Figure 1b), a 'virus evolutionary interaction' effect can result (see

Box 1) through the inheritance of the ancestral host range, leading to the daughter

virus species having correlated host assemblages. These effects can also be generated

in ecological time through mechanisms that lead to biased cross-species transmission.

Figure 1 Mechanisms for the generation of novel multihost viruses. The generation of novel multihost

viruses through host shifting (1a) leads to a 'host evolutionary interaction' effect (Box1), as the

consistent switching of viruses (V) to hosts (H) closely related to their ancestral host will lead to

related hosts having correlated viral assemblages. The generation of novel multihost viruses through

speciation (1b) can lead to a 'virus evolutionary interaction' effect (Box 1) through the inheritance of

the ancestral host range, leading to the daughter virus species having correlated host assemblages.

We tested for a role of evolutionary history in shaping the current host/virus

assemblage using species from an ecologically and economically important group, the

bumblebees. We cataloged the virome of wild-caught bumblebees from across

Scotland by RNAseq, finding at least 30 new viruses. We then tested multiple

bumblebee species for a subset of these novel viruses and three previously reported

honeybee viruses: Slow bee paralysis virus (Bailey & Woods 1974), Acute bee

paralysis virus (Bailey et al. 1963) and Hubei partiti-like virus 34 (Cornman et al.

2012; Shi et al. 2016). We analysed virus prevalence using co-phylogenetic models to

determine the presence or absence and relative strengths of the evolutionary signals

that are expected to shape the host/virus assemblage in this system, and performed

tests to attempt to determine the mechanisms driving this.

Methods

Sampling strategy

A total of 926 individual bumblebees of 13 species were collected on the wing from

nine sites across Scotland in July and August of 2009 and 2011, and frozen in liquid

nitrogen or at -80°C. In 2009, we sampled the Ochil Hills, Glenmore, Dalwhinnie,

Stirling, Iona, Staffa, and the Pentlands, and in 2011 we sampled Edinburgh and

Gorebridge (Supplementary Table 1). The cryptic species complex of *Bombus*

terrestris, *Bombus lucorum, Bombus cryptarum* and *Bombus magnus* was resolved

using RFLP analysis following Murray et al. (2008). All individuals were bisected

longitudinally prior to RNA extraction. One half of each bumblebee was used in

pooled RNA extractions of 2-11 individuals per species (median 10; Supplementary

Table 2). Two of these pools ('DIV' and 'P11') were included in the RNAseq, but

excluded from prevalence testing. The groups of bumblebees were ground in liquid

nitrogen and added to TRIzol reagent (Life Technologies) for RNA extractions,

following the manufacturer's standard protocol. The RNA concentrations in the

pooled samples were equalized to approximately 200 ng/ul/individual based on

Nanodrop measurements.

RNA Sequencing and Bioinformatics

The RNA was combined by species for *B. terrestris* (239 individuals), *Bombus*

pascuorum (212 individuals)*, B. lucorum* (182 individuals) and other *Bombus* (293

individuals) into four large RNA pools. These large pools were sequenced using the

Illumina HiSeq platform with 100bp paired end reads (Beijing Genomics Institute)

after poly-A selection. This excludes ribosomal and bacterial RNA, and will enrich

for mRNAs and those RNA viruses that have polyadenylated genomes or products.

The single-species bumblebee pools were subsequently re-sequenced following

duplex specific nuclease normalization, to reduce rRNA representation, and enrich for

rare transcripts while retaining non-polyadenylated viruses and products. The small

RNAs of the same RNA pools of *B. terrestris*, *B. lucorum* and *B. pascuorum* were

also sequenced to test for the replication of viruses identified via the transcriptome

sequencing.

For each pool, paired end RNAseq data were initially mapped to the published

Bombus terrestris and *B. impatiens* genomes using bowtie2 (Langmead & Salzberg

2012) to reduce the representation of conserved bumblebee sequences. Read pairs that

did not map concordantly, including divergent bumblebee sequences and other

associated microbiota, were assembled *de novo* using Trinity 2.2.0 (Grabherr et al.

2011) as paired end libraries, following automated trimming ('--trimmomatic') and

digital read normalisation ('--normalize_reads'). Where two RNAseq libraries (Poly-

A and DSN) had been sequenced, these were combined for assembly.

To identify putative viruses, all long open reading frames from each contig were

identified and concatenated to provide a 'bait' sequence for similarity searches using

Diamond (Buchfink et al. 2015) and BLASTp (Altschul et al. 1990). Contigs shorter

than 500 base pairs were discarded. These contig translations were used to search

against a Diamond database comprising all of the virus protein sequences available in

NCBI database 'nr', and all of the Dipteran, Hymenopteran, Nematode, Fungal,

Protist, and prokaryotic proteins available in NCBI database 'refseq_protein' (mode

'blastp'; e-value 0.001; maximum of one match). Matches to phage and short matches

to large DNA viruses were excluded. Remaining contigs were manually curated to

identify and annotate high-confidence virus-like sequences. To quantify approximate

fold-coverage, and to assess viRNA properties, the raw RNAseq and trimmed small

RNA reads were mapped against the putative viral contigs using bowtie2's '--very-

sensitive' setting and retaining only the top map (Langmead & Salzberg 2012), from

this we recorded the number of mapped reads per kilobase of transcript per million

mapped reads. We considered viruses to show strong evidence of replication in the

host if they had at least 50 mapping siRNA reads with a size distribution sharply

peaked at 22nt (viRNAs are generated from replicating viruses by Dcr2). Following

Fauquet and Stanley (Fauquet & Stanley 2005), we defined contigs exhibiting less

than 90% nucleotide identity as separate viruses and those exhibiting greater than or

equal to 90% identity as strains of known viruses.

PCR Validation and Testing

A subset of contigs were chosen for manual validation. All chosen contigs met both of

the following conditions: the presence of mapping reads in the bumblebee small

RNAs (for the *B. terrestris*, *B. pascuorum* and *B. lucorum* pools; not a condition for

the mixed *Bombus* pool) or the transcriptomic RNAs (for the other *Bombus* pool

where small RNAs were not generated), and the closest blast match being viral RNA-

dependent RNA-polymerase. Internal primers for these contigs were generated using

primer3 (Untergasser et al. 2012) and amplification of the target was verified via

Sanger sequencing. See Supplementary Table 3 for PCR conditions and primer

sequences. Mayfield virus 1 and 2 were Sanger sequence validated over the entirety

of the contig. The Loch Morlich and River Liunaeg virus sequences were generated

by the connection of several disjoint contigs by Sanger sequencing. Black Hill virus

was excluded from further analysis as it was found that that the PCR reaction

amplified a host sequence that could not be visually differentiated from the virus

product.

Phylogenetic Inference

Following Cameron et al. (2007), we inferred the bumblebee phylogeny using

cytochrome oxidase I, elongation factor 1-alpha, opsin, phosphoenolpyruvate

carboxykinase, 16S and arginine kinase genes. To break up long branches and allow

dating, additional species not sampled in the field were added (see Supplementary

Table 4 for genbank accession numbers and species included). The DNA sequences

were aligned with MAFFT using the L-INS-i setting (Katoh et al. 2017; Katoh et al.

2005). The 6 gene alignments were then used to generate the phylogeny in BEAST

v2.4.5 (Bouckaert et al. 2014), treating each file as a separate partition, using

bModelTest (Bouckaert & Drummond 2015) with the 'transitionTransversion split'

setting and a calibrated Yule tree prior (Heled & Drummond 2012). An uncorrelated

lognormal relaxed clock was fitted to each partition, with exponential(λ=1) priors

placed over the mean rate and the default gamma(α=0.5396, β=0.3819) priors being

placed over the standard deviation (Drummond et al. 2006). The bumblebees were

constrained to be monophyletic, with the honeybee, *Apis mellifera*, as an outgroup. A

gamma (α=74.85889, β=0.4366812) distributed divergence time prior was placed

over the tMRCA of the *Bombus* clade, with parameters optimised to match the 2.5[th]

and 97.5[th] percentiles of the posterior distribution of ages previously estimated by

Hines (2008). Four separate runs of the MCMC were performed for 100,000,000 steps

from random starting trees, with the first 50,000,000 steps being discarded as burn in.

Convergence of the posterior among runs was assessed in Tracer v1.7 (Rambaut et al.

2017). The posterior distribution was thinned to 1000 trees.

For the virus phylogeny, amino acid sequences were inferred based on the translated

ORFs for regions predicted to contain RdRp motifs using the GenomeNet MOTIF

search function (Kanehisa et al. 2002) against the Pfam database (Finn et al. 2014),

with an expectation cut-off of 0.00001. If a virus had no annotated motifs, the

canonical GDD RdRp amino acid motif (Kamer & Argos 1984) was identified

manually. Additional virus species (Supplementary Table 5) were added to the

phylogeny to anchor species with short generated contigs, and to break up long

branches. Given the long evolutionary distance between the viruses, PROMALS3D

(Pei et al. 2008) was used to align viral sequences. The alignments were trimmed to

the first conserved secondary structural element at both ends as predicted by

PROMALS3D with the 0.95 conservation metric. Two of the novel viruses (Agassiz

Rock virus and Cnoc Mor virus) were not included in this phylogeny because the

section of the RdRp gene required fell outside the available contig. Given that it is

unclear whether there was a universal common ancestor of all RNA viruses (Koonin

et al. 2015), we aligned the sequences and generated the phylogeny twice, with and

without the negative sense RNA viruses (Supplementary Table 5). The trees serve

purely to quantify expected variance (under a Brownian motion model of evolution)

between closely related viruses. The deep splits in the phylogeny are poorly resolved

with RdRp data (Zanotto et al. 1996), due to the fast evolutionary rates of RNA

viruses, the considerable time since divergence and permutations in the RdRp

sequence (Gorbalenya et al. 2002). However, this should not overly bias the

conclusions as beyond a certain evolutionary distance, the viruses would be expected

to become essentially uncorrelated when averaged across the posterior

(Supplementary Table 6 for realised correlations).

Phylogenetic models used the BLOSUM62 rate matrix (Henikoff & Henikoff 1992)

with gamma distributed rate variation using 4 gamma categories, an uncorrelated

lognormal relaxed clock (Drummond et al. 2006) and a Yule tree prior. A CTMC rate

reference prior (Ferreira & Suchard 2008) was placed over the clock mean and an

exponential($\lambda=1$) prior was placed over the standard deviation. The alpha parameter

of the gamma distributed rate variation was given an exponential($\lambda=1$) prior.

Absolute dating of viral trees is difficult due to the inconsistency in estimated ages

provided by estimated clock rates and known orthologous insertions between sister

host species (Holmes 2003), but is not essential for our analysis, which depends only

on relative branch lengths. Nevertheless, we chose to use orthologous insertions to

provide approximate dates for our tree. To account for the estimated ages of RNA

viral families (Katzourakis & Gifford 2010), we set a uniform lognormal prior with

an offset of 97 Mya, a mean of 500 Mya and a logged standard deviation of 0.5 on the

age of the root of the tree including the negative sense RNA viruses and a lognormal

prior with an offset of 76 Mya, a mean of 500 Mya and a logged standard deviation of

0.5 on the age of the tree excluding them. Two partitiviruses (Rosellinia necatrix

partitivirus 2 and Raphanus sativus cryptic virus 1) known to have a common ancestor

older than 10 Mya (Chiba et al. 2011) were included for dating purposes. We placed a

diffuse lognormal prior with an offset of 10 Mya, a mean of 30 Mya and a logged

standard deviation of 0.5, on the age of the MRCA of these species. Both models

were run over 10 separate chains for 50,000,000 generations on a cluster in BEAST

v1.8.4 (Drummond et al. 2012), with 25,000,000 generations being discarded as burn-

in. Convergence of the posterior was assessed in Tracer v1.7 (Rambaut et al. 2017).

The posterior distributions were combined and thinned to 1000 trees.

Prevalence Estimation

Maximum likelihood prevalence and 2-log-likelihood confidence intervals were

estimated for each host/virus combination with more than one pool using the code

from Webster et al. (2015). As the samples were small pooled groups of individuals,

such that a PCR 'positive' represents one or more infections, we modelled the

prevalence using a "pooled binomial" likelihood (Ebert et al. 2010; Gibbs & Gower

1960; Thompson 1962). This approach requires that the underlying prevalence of a

virus is the same in all pools, which is unlikely for bumblebees sampled from

different locations. Estimates should therefore be treated with caution.

Co-phylogenetic Mixed Model Analysis

To test for the evolutionary effects on association, the presence/absence data and the

phylogenetic trees were analysed using a co-phylogenetic mixed model (Hadfield et

al. 2014) implemented in Stan (Carpenter et al. 2017). Our model is explicitly focused

at the individual-level, and the model's predictions represent the predicted probability

of infection within an individual of the species. This is in contrast with Hadfield et

al.'s original implementation where the focus was at the species- or population- level

and the model was estimating the probability that the parasite would be found in the

species or population at all. In all cases, the presented models showed no divergences,

acceptable Rhat and E-BFMI values and effective sample sizes of over 200.

We fitted host and virus phylogenetic effects, which measure the extent to which

variation in prevalence is clustered on the host and viral phylogenies respectively. We

also fitted host and viral evolutionary interaction effects, which measure the extent to

which related species have similar probabilities of infection in the sets of their

interaction partners. The final phylogenetic term fitted was a coevolutionary

interaction, which measures the extent to which related hosts are infected to similar

degrees by related viruses.

In addition to the phylogenetic terms, non-phylogenetic host and virus terms, an

interaction between these terms and a pool ID term were fitted. The non-phylogenetic

host and virus terms measure variation that can be partitioned between host species

and virus species in average infection risk that is not consistent with trait evolution by

Brownian motion. The interaction term measures variation that can be partitioned

between pairwise interactions between individual hosts and viruses that is not

consistent with the linear sum of their individual means from the non-phylogenetic

host and virus terms. The pool ID effect measures variation between pools in

infection risk averaged over all the viruses tested. As the pools combined hosts by

species rather than by location, so that some had individuals from multiple locations,

we treated each location and each realised combination of locations as levels of a

random effect, terming this the "spatial composition effect". This describes the

variation in average infection level between realised combinations of locations

averaged across viruses. Model 1 included all the viruses, Model 2 excluded the

negative sense RNA viruses and Model 3 fitted a pseudo-taxonomic model. In Model

3, the relationship among the viruses was represented by a polytomic viral tree with

arbitrary branch lengths (with a root-to tip distance of 1 unit, and equal length

between each taxonomic level) with the viruses being split first by their genomic type

(+ve sense RNA, -ve sense RNA and dsRNA) implying a covariance of 0 between

genome structures, followed by splitting by the putative viral clades identified by Shi

et al. (2016). This was done to test for potential bias caused the by the possibility of

systematic misidentification of the correct relationship between families in the

estimated viral trees.

The form of the models is shown below, where i indexes the data points, $group_i$

represents the level of a categorical variable that the ith pool belongs to, y_i represents

the 1/0 indicator for the presence or absence of infection in the ith pool, k_i represents

the number of individuals in the ith pool, p_i is the unmeasured probability of infection

of a single individual in the ith pool, y'_i is the estimated value of $\log_e(p_i/(1 - p_i))$, μ is

the global mean of the latent variable, ε is a normally distributed error term. All terms

were fitted as random effects (i.e. estimated by partial pooling). As above, a "pooled

binomial" likelihood was used (Ebert et al. 2010; Gibbs & Gower 1960; Thompson

1962).

$$y_i \sim \text{Bernoulli}(1 - (1 - p_i)^{k_i})$$

$p_i = \exp(y'_i)/\exp(1 + y'_i)$

$y'_i = \mu + \text{host}_i + \text{virus}_i + \text{interaction}_i + \text{host phylogenetic effect}_i + \text{virus}$

$\text{phylogenetic effect}_i + \text{host evolutionary interaction effect}_i + \text{virus evolutionary}$

$\text{interaction}_i + \text{coevolutionary interaction}_i + \text{pool ID}_i + \text{species composition}_i + \varepsilon$

All variance-covariance matrices were generated as described in Hadfield et al.

(2014), with the variance-covariance matrices scaled to correlation matrices. A

standard logistic prior was placed over the global intercept on the latent scale, μ,

representing a flat prior on the probability scale. An exponential(λ=1) prior was

placed on each variance term in the model. In the full model with all variances being

estimated, this is equivalent to a gamma(α=11, β=1) prior over the total variance,

which gives a prior mean variance of 11, and an appropriate prior on the standard

deviation of a variable on the logit scale. Intraclass correlations, which represent the

proportion of the variance explained by each effect, were calculated on the link scale

(with an addition of $\pi^2/3$ to the denominator to account for the variance of the logistic

distribution of the latent variable) from the model outputs and reported. Highest

posterior density intervals were calculated by the SPIn method (Liu et al. 2015) and

90% credible intervals are reported as these are more robust to sampling in the tails of

the posterior distribution (Stan Development Team 2017).

The total phylogenetic variance was calculated as:

$(\sigma^2_{\text{host phylogenetic}} + \sigma^2_{\text{host interaction}} + \sigma^2_{\text{virus phylogenetic}} + \sigma^2_{\text{virus interaction}} + \sigma^2_{\text{coevolutionary}}$

$_{\text{interaction}})/(\sigma^2_{\text{total}} + \pi^2/3)$

The total non-phylogenetic variance was calculated as:

$(\sigma^2_{\text{host}} + \sigma^2_{\text{virus}} + \sigma^2_{\text{interaction}} + \sigma^2_{\text{poolID}} + \sigma^2_{\text{spatial composition}})/(\sigma_{\text{total}} + \pi^2/3)$

Uncertainty in the inferred phylogenies was accounted for by direct marginalisation.

This dramatically increased the runtime of the model, as, given the input of H host

phylogenies and V viral phylogenies from their posterior distributions, the likelihood

of each datapoint has to be calculated *HV* times. As such, we included only 10 trees

from each posterior, as a trade-off between runtime and accounting for uncertainty in

the tree hypotheses. The marginalisation is show below, with **y** being the total vector

of presences and absences, *H* being the number of host phylogenies used, *V* being the

number viral phylogenies used, θ being all the non-variance-covariance parameters in

the model, Ω_j being the set of variance-covariance matrices generated by the *j*th

combination of host and virus phylogenies, Ω_{HV} representing the set of all variance-

covariance matrices being marginalised over and $\mathcal{L}$ representing a likelihood.

$\mathcal{L}(\boldsymbol{y} \mid \theta, \Omega_{HV}) = \sum_{j=1}^{HV} \frac{1}{HV} \mathcal{L}(\boldsymbol{y} \mid \theta, \Omega_j)$

Tongue Length Analysis

After finding that the posterior for the host evolutionary interaction was well resolved

from zero, we designed a post-hoc test to attempt to detect signal for one of the

obvious mechanistic explanations for this; structured transmission networks driven by

evolutionarily conserved anatomical factors. We tested for an association between the

tongue length differences between bumblebee species and the differences in their viral

community structures, as a proxy for signal of differential transmission at flowers

driven by evolutionarily conserved flower choice. Average tongue lengths for each

bumblebee species except *Bombus bohemicus* and *Bombus cryptarum* were taken

from Goulson, Hanley and Darvill (2005). No published tongue length could be found

for *Bombus cryptarum*, so we assumed that it was identical to that of *Bombus*

lucorum, a species of which it is near indistinguishable in the field. *Bombus*

bohemicus was excluded from this analysis, because it is an inquiline parasite, and

therefore its ecology differs from the other species in such a way that tongue length

would not expected be expected to be correlated with the viral community distance.

In order to test for a correlation between tongue length and virus similarity, estimates

of the distance in viral communities between host species are required. These were

generated as follows: For each host-virus combination, the package 'prevalence' was

used to generate posterior draws of the underlying prevalences under a Beta (1,1)

prior. Then 1000 draws per species were taken from these sets of MCMC draws to

generate 1000 matrices of host-virus prevalences consistent with the raw data. For

each of these matrices, the distance between each species' viral community was

calculated by taking the vector of estimated prevalences for the 16 viruses of a given

species as a coordinate in a 16-dimensional space then calculating the Euclidean

distance between these points. The rank correlation (Kendall's τ-b) between each pair

of species' viral community distances and their tongue length distances was then

calculated, using the mantel function in the R package vegan. The point estimate

presented is the median of the 1000 initial correlations accounting for the uncertainty

in the underlying prevalences. The 95% confidence interval is the 2.5^{th} and 97.5^{th}

percentiles distribution of estimated correlations.

Results

RNA was extracted from 13 species of bumblebee from nine sites, to identify new

viruses, assay their prevalence and their pattern of distribution across host species and

to test whether the evolutionary histories of the viruses and hosts have impacted the

current distribution.

Read and Assembly Statistics

A total of 134,026,056 sequencing read pairs were generated for *Bombus lucorum*,

135,590,922 for *Bombus terrestris*, 128,670,194 for *Bombus pascuorum* and

26,838,390 for the other *Bombus* species with 0.37, 0.38, 3.36 and 15.12 percent of

reads mapping to the known viruses or the novel bee viruses found in the study. The

poly-A and DSN normalized datasets were unexpectedly highly correlated, given their

expected biases (1.000 for *Bombus terrestris*, 0.999 for *Bombus pascuorum* and 0.998

for *Bombus lucorum*) implying that the sequences results were highly consistent

irrespective of the selection method used.

Previously Described Viruses Present in the Metagenomic Pools

RNAseq reads mapped to three previously described bee viruses. The majority of

these reads mapped either to the Acute bee paralysis virus/Kashmir bee virus complex

(henceforth ABPV) (Bailey et al. 1963) or to Slow bee paralysis virus (SBPV) (Bailey

& Woods 1974). Additionally, in the mixed *Bombus* pool, reads were found mapping

to Hubei partiti-like virus 34 (HPLV34) a virus initially detected, though not named,

in honeybees by Cornman et al. (2012), then subsequently also reported in a sample

from Chinese landsnails by Shi et al (2016).

No RNAseq reads were mapped to Deformed wing virus – type A (Bailey & Ball

1991), Chronic bee paralysis virus (Bailey et al. 1963), Bee macula-like virus (de

Miranda et al. 2015), Ganda bee virus (Schoonvaere et al. 2016), Scaldis River bee

virus (Schoonvaere et al. 2016), Black queen cell virus (Bailey & Woods 1977), Apis

rhabdovirus 1 (Remnant et al. 2017), Apis rhabdovirus 2 (Remnant et al. 2017), Apis

bunyavirus 1 (Remnant et al. 2017), Apis bunyavirus 2 (Remnant et al. 2017), Apis

flavivirus (Remnant et al. 2017), Apis dicistrovirus (Remnant et al. 2017), Apis Nora

virus (Remnant et al. 2017) and members of the Lake Sinai virus complex (Runckel et

al. 2011). A small number of small RNA reads did map to these viruses, however, this

likely represents cross-mapping, given the lower stringency of 22nt reads. Two of the

viral contigs generated by the *de novo* assembly had high similarity to previously

described plant viruses; both RNAs of White clover cryptic virus 2 (Boccardo et al.

1985) (96% identity), both RNAs of strain of Arabis mosaic virus

(MH614320/MH614321) (Smith & Markham 1944) distant to previously sequenced

strains (91% identity) and a strain of Red Clover nepovirus A (MH614312) (Koloniuk

et al. 2018) distant to previously sequenced strains (90% identity).

Putative Novel Viral-like Sequences

We identified 37 putative novel viral contigs, four mapping to DNA viruses (4

densovirus-like contigs) and 33 to RNA viruses (4 Reo group contigs, 2 Toti-Chryso

group contigs, 4 Bunya-Arena group contigs, 1 Orthomyxoviridae-like contig, 8

Hepe-Virga group contigs, 12 Picorna-Calici group contigs and 2 Tombus-Noda

group contigs). Based on the supposition that a contig represents a separate virus if it

maps to a different viral grouping than the other contigs, or if it can be aligned to all

other contigs within its assigned viral grouping, this represents 30 new viruses with

seven remaining contigs that may represent other genomic regions of these 30 viruses

or separate viruses that cannot be confirmed as such. See Table 1 for information on

the viruses tested for prevalence using PCR and Supplementary Table 7 for detailed

information on all of the identified contigs. The numbers of reads mapping these

contigs were variable and are shown in Table 2.

Table 1 The names, genome structures and groupings (following Shi et al. (2016)) of the newly

discovered viruses for which prevalence was assessed.

Putative viral contig	Abbreviations	Clade	Genome structure
Agassiz Rock virus	ARV	Reo	dsRNA
Elf Loch virus	ELV	Reo	dsRNA
Dumyat virus	DV	Toti-Chryso	dsRNA
Sheriffmuir virus	SV	Toti-Chryso	dsRNA
Clamshell Cave virus	CCV	Bunya-Arena	- ssRNA
Allermuir Hill virus 1	AHV1	Hepe-Virga	+ssRNA
Allermuir Hill virus 2	AHV2	Hepe-Virga	+ssRNA
Allermuir Hill virus 3	AHV3	Hepe-Virga	+ssRNA
Mill Lade virus	MLV	Hepe-Virga	+ssRNA
Boghill Burn virus	BBV	Picorna-Calici	+ssRNA
Gorebridge virus	GV	Picorna-Calici	+ssRNA
Loch Morlich virus	LMV	Picorna-Calici	+ssRNA
Mayfield virus 1	MV1	Picorna-Calici	+ssRNA
Mayfield virus 2	MV2	Picorna-Calici	+ssRNA
River Liunaeg virus	RLV	Picorna-Calici	+ssRNA
Castleton Burn virus	CBV	Tombus-Noda	+ssRNA

Table 2 The RNAseq reads per kilobase per mapped million reads in the *Bombus terrestris*, *Bombus*

lucorum, Bombus pascuorum and mixed *Bombus* pools. Structural zeros are indicated by dashes, zeros

in the table indicate below 0.005. Contigs with names in bold meet the criterion of having at least 50

mapping small RNA reads with a sharp peak in the size distribution at 22nt in *Bombus terrestris,*

Bombus lucorum and *Bombus pascuorum* providing evidence of replication (see main text).

Putative viral contig	Accession number	*Bombus terrestris*	*Bombus lucorum*	*Bombus pascuorum*	mixed *Bombus*
Bombus-associated Densovirus-like Contig 1	MH614322	0.09	-	-	12.34
Bombus-associated Densovirus-like Contig 2	MH614323	0.39	2.66	-	14.23
Agassiz Rock virus	MH614287	3.77	0.49	-	-
Cnoc Mor virus	MH614297	0.97	-	-	20.38
Bombus-associated Reoviridae-like Contig 1	MH614298	0.42	0.03	-	1.63
Elf Loch virus	MH614300	-	0.00	1.00	0.05
Dumyat virus	MH614299	-	-	-	10.30
Sheriffmuir virus	MH614317	-	-	-	2.55

Clamshell Cave virus	MH614294	0.07	-	-	2.10
Bombus-associated Bunyaviridae-like Contig 1	MH614295	0.70	-	-	5.61
Bombus-associated Bunyaviridae-like Contig 2	MH614296	-	-	-	12.13
Bombus-associated Phlebovirus-like Contig 1	MH614315	3.07	2.77	0.95	1.63
Bombus-associated Orthomyxovirus-like Contig 1	MH614314	-	-	0.44	-
Allermuir Hill virus 1	MH614288	15.27	0.64	0.02	1.50
Allermuir Hill virus 2	MH614289	0.01	0.03	12.71	0.13
Allermuir Hill virus 3	MH614290	0.40	2.62	0.50	3.35
Mill Lade virus	MH614306	0.40	0.48	0.03	7.28
Bombus-associated Virga-like Contig 1	MH614308	-	0.65	-	-
Bombus-associated Virga-like Contig 2	MH614309	-	0.33	-	-
Bombus-associated Virga-like Contig 3	MH614318	16.03	4.83	0.52	90.37
Bombus-associated Virga-like Contig 4	MH614319	1.66	0.11	-	0.54
Black Hill virus	MH614291	-	-	-	2.96
Boghill Burn virus	MH614292	0.00	10.92	0.00	0.11
Gorebridge virus	MH614301	2.97	0.02	-	0.15
Bombus-associated Picornavirus-like Contig 1	MH614302	6.27	0.02	-	0.29
Loch Morlich virus	MH614303	0.00	-	-	7.83
Mayfield virus 1	MH614304	391.31	232.93	0.59	7.67
Mayfield virus 2	MH614305	4.87	0.79	336.97	558.82
Bombus-associated Nepovirus-like Contig 1	MH614310	1.72	1.02	0.09	1.70
Bombus-associated Nepovirus-like Contig 2	MH614311	0.33	0.39	0.11	0.21
Bombus-associated Picornavirus-like Contig 2	MH614316	0.11	0.29	1.02	0.04
River Liunaeg virus	MH614307	0.24	0.47	0.01	9.58
Castleton Burn virus	MH614293	2.39	12.30	8.10	122.99
Bombus-associated Nodavirus-like Contig 1	MH614313	-	-	-	1.50

siRNA-based Evidence for Infection

RNA interference is an important component of antiviral defence in arthropods

(Bronkhorst et al. 2012). As part of this defence mechanism, homologs of *Drosophila*

Dicer-2 cleave dsRNA, usually in the form of replication intermediates, giving rise to

a characteristically narrow and sharply peaked distribution of virus-derived small

RNAs. Thus the presence of such small RNAs from both strands of an ssRNA virus

provide compelling evidence that the virus was replicating. In bumblebees the

characteristic Dicer-mediated viral siRNAs peak sharply at 22nt (Remnant et al.

2017), and the viruses that displayed at least 50 characteristic viral siRNAs are

marked in Table 2. The distribution of the mapped small RNA reads is shown in

Figure 3 for all viruses where the siRNAs are described in the main text, with full data

in Supplementary Figure 1.

In the three bumblebee species with siRNA data, a sequence similar to a phlebovirus

glycoprotein (AEL29653.1) displayed >50 siRNA reads. However, the size spectra of

these reads is centered on 24nt with a strong bias for a 5' terminal uracil, with the

antisense mapping orientation being more prevalent. This 5' U-bias is consistent with

insect piRNAs (Brennecke et al. 2007), and the predominant antisense orientation is

consistent with the piRNA mapping pattern to endogenous viral elements (EVE) in

mosquitoes (Suzuki et al. 2017). However, the size of piRNAs in bumblebees is

generally larger than this (Lewis et al. 2018). This sequence is therefore potentially an

EVE that has either been gained multiple times or has been maintained in the

bumblebee genome since at least the *B. pascuorum-B. terrestris/B. lucorum* split.

It is notable that the size distribution of viral siRNAs is less sharply peaked in

Mayfield virus 1, Mayfield virus 2 and Slow bee paralysis virus (excepting Mayfield

virus 1 in *B. lucorum*, which is sharply peaked), with broad 'shoulders'. This is

reminiscent of the pattern seen for Drosophila C virus and Drosophila Nora virus in

wild-collected D. *melanogaster* (Webster et al. 2015), both of which contain a viral

suppressor of RNAi (van Rij et al. 2006; van Mierlo et al. 2012).

B. pascuorum also had siRNA reads mapping to a sequence with 19% identity to

Tomato black ring virus (CAA56792.1). However, the read length spectra were

sharply peaked at 21nt, rather than the 22nt of bumblebee viRNAs. This is consistent

with siRNA's produced from DLC4, the key antiviral dicer in *Arabidopsis thaliana*

(Deleris et al. 2006) implying acquisition of the small RNAs through nectar or pollen

contamination.

Figure 2 The mapping of small RNA reads to Mayfield virus 1, Mayfield virus 2, a contig similar to

Tomato black ring virus, a contig similar to a phlebovirus glycoprotein and Slow bee paralysis virus.

Blue lines represent reads mapping to the positive-sense strand at that genomic position, red lines

represent reads mapping to the negative sense strand. The histogram of read size spectra shows the

count of reads of each length mapping in the positive (above) and negative (below) directions. The

colouring of each bar shows the counts of the reads beginning with each 5' base (red-U, blue-C, green-

A, yellow-G).

Prevalence

Species level prevalences differed dramatically among the different viruses (Figure

3). Prevalences were generally low to intermediate, with modal viral prevalences for

most host-virus combinations being below 15%. Slow bee paralysis virus was by far

the most common virus in the sample, with estimated prevalences of greater than 25%

in multiple species. Our ability to estimate the prevalence of common viruses is

limited by the pooling, leading us to only be able to assign lower bounds to

prevalences in these cases, but in 7 of 11 species, all pools were positive for SBPV.

Acute bee paralysis virus, Hubei partiti-like virus 34, Castleton Burn virus,

Gorebridge virus, Mayfield virus 1 and Mayfield virus 2 all reached 15-25%

prevalences in multiple species. Several viruses showed strong signals of species

specificity, having very low to zero prevalences in multiple host species but high

prevalences in others. Examples of this pattern include Allermuir Hill virus 1 in *B.*

terrestris, Allermuir Hill virus 2 in *B. pascuorum*, Allermuir Hill virus 3 in *B. magnus*

and *B. monticola*, as well as Loch Morlich virus and River Luinaeg virus in *B.*

jonellus.

Figure 3 A heatmap of maximum likelihood estimates for prevalence. Hosts and viruses are ordered by

phylogenetic relatedness, the trees represent the maximum clade credibility topology. Squares in red

with maximum likelihood estimates of the prevalence of 1 correspond to cases where all pools were

positive. The maximum likelihood estimate is likely extremely upwardly biased in this case.

Host-Pathogen Co-phylogenetic Models

All models that included a virus phylogeny term gave qualitatively similar results

(Figure 4, Table 3). This suggests that the results are robust to both phylogenetic

uncertainty and the assumption of a common ancestor of all RNA viruses. For this

reason, for the rest of this section, estimates will be given from the model containing

the estimated phylogeny with all the RNA viruses included. All estimates represent

the percentage of the total variance in the model (the sum of all estimated variance

components adjusted for the variance of the link function by the addition of $\pi^2/3$)

explained by a term. In all cases, the presented point estimate is the posterior mean,

and 90% shortest posterior density intervals (Liu et al. 2015) are presented following

in square brackets. Shortest posterior density intervals are a variant of highest

posterior density intervals and describe the shortest possible interval containing (in

this case) 90% of the probability density for the parameter. We present 90% intervals

rather than the standard 95% intervals, as 95% intervals calculated from simulation

draws are less computationally stable (Stan Development Team 2017). In all cases but

the virus phylogenetic effect, the posterior estimates for the proportion of variance

explained by each effect differed strongly from their induced priors (Supplementary

Figure 2).

Summary of Model Results

We find evidence that which viral species infects a host, the specific interaction

between individual hosts and individual viruses and related hosts having similar

prevalences with the same sets of viruses all explain variation in infection prevalence.

Total Evolutionarily-associated Variation

In the models containing the virus phylogeny, approximately a quarter (25.9% [11.6-

40.4]) of the total variation in prevalence was explained by terms accounting for the

evolutionary histories of hosts and viruses (host phylogenetic effect, virus

phylogenetic effect, host evolutionary interaction effect, virus evolutionary interaction

effect and coevolutionary interaction).

Host and Virus Level Effects

The host and virus phylogenetic effects measure the extent to which related hosts

have similar average prevalences of virus infection and related viruses have similar

average prevalences across hosts. The host and virus non-phylogenetic terms measure

the extent to which hosts and viruses differ in their average infection levels in manner

not consistent with evolution by Brownian motion along a phylogeny. The host

species and phylogenetic effects explained a small proportion of the total variance in

infection probability (species: 1.9%, [0.0-4.6]; phylogenetic: 2.9%, [0.0-6.5]). The

shape of the posterior distributions for the two parameters visualised in Figure 6,

makes it clear that the most credible values for both of these parameters are 0. While

it is unlikely that there is no variation in average prevalence between host species, it is

clear that the amount of prevalence explained by hosts differing in their infection

levels averaged across viruses is small relative to the other effects.

The virus species and phylogenetic effects explained a larger proportion of the total

variance in infection probability, with non-phylogenetic variation dominating, but

were imprecisely estimated (species: 13.8%, [3.8-23.7]; phylogenetic: 8.5%, [0.0-

17.8]). The posterior density for the phylogenetic effect is concentrated at 0. So, while

it is clear there is a virus species effect, the data does not appear informative for the

presence or absence of a viral phylogenetic effect in this system. The posterior draws

for the viral species and phylogenetic effect were negatively correlated within

iterations, indicating that the model had difficulty partitioning the two. This partial

non-identifiablility explains the broad posteriors on both.

Interaction Effects

A host evolutionary interaction effect measures the extent to which more closely

related hosts have more similar prevalences with the same sets of viruses, and a virus

evolutionary interaction effect measures the extent to which more closely related

viruses infect the same sets of hosts to more similar degrees. A coevolutionary

interaction term measures the extent to which more related hosts are infected to

similar degrees with viruses that are themselves related. The non-phylogenetic

interaction term measures the extent to which there is variance in the mean prevalence

of specific host-virus pairings, that are is not consistent with the other interaction

effects.

There was little evidence for a virus evolutionary interaction effect or coevolutionary

interaction having a large effect on the observed prevalences (virus: 2.9% [0.0-6.1];

coevolutionary: 2.0% [0.0-4.6]). In both cases, the marginal posterior distributions

were peaked at 0.

There was evidence for a host evolutionary interaction explaining some of the total

variance in prevalence (9.5% [0.0-18.3]). This is the only parameter in the model

where the estimated size of the effect depended strongly on the specific treatment of

the virus phylogeny (see Figure 4), and indeed whether the lower 90% bound of the

credible interval rounded to 0.0 or 0.1 depended on the phylogenetic matrix (or set of

phylogenetic matrices) inputted. The marginal posterior distribution of the parameter

was concentrated at lower values when the estimated phylogeny including the

negative-sense RNA viruses was used, and at higher values in the other two cases.

However, irrespective of the choice of virus phylogeny, the mode of the distribution

and majority of the density was distant from zero, implying that the effect is likely to

be biologically relevant. As the virus phylogenies themselves are not actually directly

involved in this term, this must be due to the partitioning of variance across other

terms being cryptically different depending on the assumptions about the virus

phylogeny.

There was also a clear non-phylogenetic interaction (15.8% [5.8-25.4]), implying that

much of the variation in prevalence is due to the specifics of individual host-virus

combinations.

As with the virus species effect and the virus phylogenetic effect, the MCMC draws

for the proportion of variance explained by the host evolutionary interaction and non-

phylogenetic interaction were negatively correlated within an iteration, implying that

separating these parameters was proving difficult. While this lead to a diffuse

posterior with wide credible intervals for both, they remain individually interpretable,

and both effects appear present simultaneously.

764
Figure 4 Comparison of estimated proportion of variance in prevalence explained by different

parameters between models. As each factor explains a proportion of the attributed variance in the

model total over all factors must sum to 1. For each parameter, the circle represents the modal estimate,

the thick bars represent the 50% shortest posterior density interval and the thin bars represent the 90%

shortest posterior density interval. "Pool ID" is the proportion of the total variation in prevalence

explained by pools within species differing in the degree to which they were infected by viruses.

"Spatial Composition" is the proportion of the total variation in prevalence explained by the

combination of locations from which the bees in the pool originate. "Host Species Effect" is the

proportion of variation in prevalence explained by hosts having different average viral prevalences.

"Virus Species Effect" is the proportion of variation in prevalence explained by viruses differing in

their average prevalences. "Non-phylogenetic Interaction" is the proportion of variation in prevalence

explained by host-virus combinations differing in the their average prevalences beyond that which

would be expected by their host and virus species effects alone. "Host Phylogenetic Effect" is the

proportion of variation in prevalence explained by hosts having average viral prevalences correlated

across the host phylogeny. "Virus Phylogenetic Effect" is the proportion of variation in prevalence

explained by viruses having average prevalences correlated across the viral phylogeny. "Host

Evolutionary Interaction" is the proportion of variation in prevalence explained by related hosts having

correlated viral assemblages. "Virus Evolutionary Interaction" is the proportion of the variation

explained by related viruses having correlated host assemblages. "Coevolutionary Interaction" is the

proportion of the variation explained by related hosts having similar prevalences of related viruses.

"Total Non-phylogenetic" is the proportion of the variation that can be explained by terms not

involving the host and virus phylogeny and excluding the residual ("Host Species Effect", "Virus

Species Effect", "Pool ID", "Spatial Composition Effect", "Non-phylogenetic Interaction"). "Total

phylogenetic" is the proportion of the variation that can be explained by terms involving a host or virus

phylogeny ("Host Phylogenetic Effect", "Virus Phylogenetic Effect", "Host Evolutionary Interaction",

"Virus Evolutionary Interaction", "Coevolutionary Interaction"). "Residual + Link" is the proportion of

the total variance that is explained by the residual variance and variance of the logistic distribution

$(\pi^2/3)$.

Table 3 Mean estimates for the intra-class correlations of each variance component. The point estimate

is the posterior mean, the numbers in brackets represent the 90% shortest posterior density interval.

	All Viruses	No Negative Sense Viruses	Pseudo-taxonomic
Virus Species Effect	13.8 (3.8, 23.7)	13.6 (2.1, 23.2)	14.4 (4.1, 24.2)
Virus Phylogenetic Effect	8.5 (0, 17.8)	8.9 (0, 19.4)	7.9 (0, 17.1)
Host Species Effect	1.9 (0, 4.6)	2.1 (0, 4.8)	1.9 (0, 4.6)
Host Phylogenetic Effect	2.9 (0, 6.5)	3.0 (0, 6.9)	2.9 (0, 6.6)
Virus Evolutionary Interaction	2.9 (0, 6.1)	2.3 (0, 5.1)	2.9 (0, 6.6)
Host Evolutionary Interaction	9.5 (0, 18.3)	10.4 (0.2, 18.8)	10.5 (0, 19.0)
Non-phylogenetic Interaction	15.8 (5.8, 25.4)	15.8 (6.5, 25.2)	14.9 (4.6, 24.0)
Coevolutionary Interaction	2.0 (0, 4.6)	2.2 (0, 5.0)	2.0 (0, 4.5)
Pool ID	3.0 (1.1, 4.8)	3.3 (1.4, 5.3)	3.0 (1.2, 4.8)
Pool Spatial Composition	1.4 (0, 2.9)	1.3 (0, 2.8)	1.3 (0, 2.9)
Residual + Link	38.3 (31.5, 45.1)	37.2 (30.3, 43.9)	38 (31, 44.7)
Total Non-phylogenetic	35.9 (22.4, 50.3)	36 (21.2, 49.8)	35.7 (22.7, 50.1)
Total Phylogenetic	25.9 (11.6, 40.4)	26.8 (11.6, 41.5)	26.3 (11.7, 40.4)

Tongue Length-Viral Community Correlation

Given that the co-phylogenetic model found that related hosts share viral

communities, one potential mechanism for this is phylogenetically-biased exposure,

driven by phylogenetically correlated floral preferences. If bumblebee species with

similar flower preferences had similar viral communities, it would be expected that

there would be a positive correlation between tongue length similarity (as this is an

important factor in floral preference) and viral community similarity between pairs of

species. The point estimate of the correlation between the two distances was small

and negative (-0.06), but the 95% confidence intervals for that point estimate

overlapped zero (-0.13, 0.00), so given the uncertainty in the data, a correlation of

zero cannot be rejected. None-the-less, given this data, a strong positive relationship

between tongue length and viral community similarity seems unlikely, a result

inconsistent with phylogenetically-biased exposure driven by tongue length-mediated

floral choice.

Discussion

Using wild bumblebee species that share transmission opportunities, we have shown

that variation in the prevalence of infection in the wild is explained by related hosts

being infected with the same viruses to similar degrees, viruses differing in their

average prevalence and individual virus-host pairings having greater or lesser

prevalence than would be expected by the interaction of the host and virus species

effects alone.

Virus Discovery

There is now an extensive diversity of viruses known in bees, with most new studies

finding novel viruses (Cornman et al. 2012; Mordecai et al. 2015; Remnant et al.

2017; Runckel et al. 2011; Schoonvaere et al. 2016; Schoonvaere et al. 2018; Roberts

et al. 2018). We have found 37 novel putative viral contigs in the transcriptomes of

wild-caught bumblebees from across Scotland, suggesting that virus discovery in this

taxonomic group is far from saturation. As with any metagenomic study, it is hard to

be confident that the virus-like contigs represent real infections of the sampled host,

rather than surface or gut contaminants. However, the presence of 22nt virus siRNAs,

generated from double-stranded viruses by Dicer as part of an antiviral response in the

host, provides compelling evidence that at least 10 of these contigs (Densovirus 2 and

3, Elf Loch virus, Allermuir Hill virus 1, 2 and 3, Mill Lade virus, Mayfield virus 1

and 2, and Castleton Burn virus) represent active viral infections in bumblebees.

Mites and nematodes both parasitise bumblebees and therefore could potentially be an

alternative source of the small RNAs. Mite viRNAs are reported to be centered at

24nt (Remnant et al. 2017), and could therefore not produce the small RNA patterns

observed. Nematode viRNAs are centered at 22nt, like bumblebee viRNAs, (Félix et

al. 2011) and thus could potentially produce this pattern. While, outside of queens

infected with *Sphaerularia bombi*, nematode infection of wild bumblebees appears to

be very rare (Rao et al. 2017), nematodes cannot be categorically ruled out as a source

of the observed small RNAs. One contig's (MH614312) mapped small RNAs were

centered at 21nt, and the closest known virus was a nepovirus of plants. As DCL4, a

major plant Dicer, produces viRNAs of this size (Wang et al. 2011), this is consistent

with that particular virus being a plant virus, which was transferred in collected nectar

or pollen.

Phylogenetic Effects

We found no evidence for a large host phylogenetic effect, where related hosts have

correlated average viral prevalences. To the best of our knowledge no studies have

previously applied these methods to viruses sampled from wild animals. However,

other traits relating to viral disease in a series of studies in *Drosophila* species under

experimental conditions have consistently detected host phylogenetic effects in

factors that would be expected to be correlated with prevalence in the wild, such as

infection probability (Longdon et al. 2011), virulence and viral load (Longdon et al.

2015) and viral load alone (Roberts et al. 2018). However, two of these studies

focused on a single isolate of Drosophila C virus. Therefore, the variation that they

attribute to a phylogenetic effect may be partitioned into the host evolutionary

interaction in our study, as a host evolutionary interaction is equivalent to separate

inconsistent host phylogenetic effects for each interaction partner. Our were not

particularly informative for the presence or absence of a virus phylogenetic effect,

with the posterior being very diffuse with a majority of the density near zero. This

appears partly due to difficulties partitioning the variation between the virus species

effect and virus phylogenetic effect. Irrespective of the cause, we can make no strong

statements about whether related viruses exhibit similar prevalences across hosts from

this dataset.

Host Species Effect

There was little evidence for an important host species effect, implying that hosts do

not strongly vary in the average degree to which they are infected with viruses. The

previous studies using these methods have universally found host species effects

(Hadfield et al. 2014; Waxman et al. 2014). However, as both of these studies have

used mammal-eukaryotic parasite datasets, the degree of relevance for them as a

comparison in unclear. Experimental evidence from virus studies across drosophilid

flies have found weak to zero host species effects on the titre of sigma viruses

(Longdon et al. 2011) and Drosophila C virus virulence (Longdon et al. 2015) but

considerably larger host species effects on Drosophila C virus load (Longdon et al.

2015). This between-study variation potentially indicates a reason we did not detect a

host species effect. With a single pathogen, the average and particular degrees of

variation in infection between hosts are identical. As soon as multiple pathogens are

involved, they diverge, such that it is possible for there to be no variation in the

average prevalence between hosts, but still considerable variation in the prevalence of

particular viruses between hosts, which is consistent with the presence of a host

species effect in correlates of prevalence in some viruses but not others, as noted

above.

Virus Species Effect

A clear virus species effect was detected in the dataset, despite the uncertainty added

by the difficulty partitioning the virus species and phylogenetic effects. Therefore,

viruses differed in their prevalences averaged across hosts. This is not a surprising

result as viruses differ in host range (Bandín & Dopazo 2011), virulence (Langsjoen

et al. 2018; Baker & Antonovics 2012) and infectious period (Baker & Antonovics

2012) at both the species and the strain level. Variation in host range changes the size

of the host pool available for infection, and variation in virulence and infectious

period both change the length of time any infected host is available for sampling. All

these factors would be expected to drive consistent differences in long-run

prevalences between viruses. Additionally, as our sites were only sampled once,

short-term effects will also drive between virus variation. Any virus that was

experiencing an epizootic at the time of sampling will be overrepresented relative to

its long-run prevalence, further increasing the between virus variation.

Host Evolutionary Interaction Effect

A host assemblage effect was found, where phylogenetically related hosts share viral

assemblages, showing that more closely related hosts are more similar in virus

prevalence for groups of viruses. The statistical machinery required for estimating this

effect is quite new and, as such in the disease ecology field, has predominantly been

applied to mammal-parasite and plant-parasite systems where there are good datasets

already existing. None-the-less, host evolutionary interactions have always been

found when searched for using these methods (Waxman et al. 2014; Hadfield et al.

2014) and analogous effects are commonly found using different methods (Davies &

Pedersen 2008; Huang et al. 2014; Cooper et al. 2012). In a system where these

viruses were host limited, this pattern could be explained by preferential host shifting,

where parasites more frequently gain the ability to infect hosts closely related to ones

that they are already capable of infecting. Preferential host shifting is known to be a

general phenomenon, and has been observed in macroparasites, viruses and

protozoans (see Longdon et al. 2014 and the references within).

While some of the viruses in this study were not detected in a subset of host species,

most of the viruses found here appear to genuinely be multihost viruses, with the

majority being detected in over half the sampled species. Given this, a combination of

biased cross-species transmission and preferential host shifting appears a better

explanation in this system. Biased cross-species transmission occurs when

transmission occurs more frequently between some species that a pathogen is already

capable of infecting than others. This biased cross-species transmission could be

driven by two non-exclusive mechanisms: phylogenetically-biased transmission

probabilities and phylogenetically-biased exposure.

Phylogenetically-biased transmission probabilities occurs when cross-species

transmission is more frequent among close relatives, due to the probability of

infection after contact with the virus being similar between related species. Related

hosts present correlated environments from the perspective of the virus at the

molecular and anatomical level, therefore adaptation to one should provide

corresponding fitness increases on the other. Experimental results have shown that

correlated mutations occur on viral entry to related hosts (Longdon et al. 2018),

implying that this cross-adaptation does occur. However, this is probabilistic, and

different routes the mutations fixed on entry can differ between replicate entries

(Longdon et al. 2018; Streicker et al. 2012). Therefore, if there is antagonistic

pleiotropy between mutations that are adaptive in two different groups of hosts and

cross-adaptation predicts the probability of successful infection on contact, then a

phylogenetically-biased transmission network will result.

Phylogenetically-biased exposure represents an evolutionarily-driven ecological

phenomenon that biases cross-species transmission rates, mediated by niche overlap.

Contaminated flowers are likely to be an important source of intra- and inter- specific

pathogen transmission in bumblebees and pollinators more generally (Durrer &

Schmid-Hempel 1994; Graystock et al. 2015; McArt et al. 2014). The flower

visitation network has been shown to be associated with the partitioning of genetic

diversity of *Crithidia bombi* between bumblebee hosts (Salathé & Schmid-Hempel

2011), and the network itself is highly structured, though temporally variable (Ruiz-

González et al. 2012). Different bumblebee species show tongue length differences,

which are phylogenetically associated (Harmon-Threatt & Ackerly 2013), and the

differences in tongue length correlate with differential flower usage between

bumblebee species (Goulson et al. 2008; Inouye 1978). If infection occurs at

contaminated flowers, the structuring of the flower usage network could cause

different flowers to build up different surface viral communities. This could drive

consistent phylogenetically-correlated differences in viral infection rates through

differential exposure.

Post-hoc testing did not find a positive relationship between tongue length

dissimilarity (a rough proxy for species-level flower choice dissimilarity) and viral

community composition dissimilarity, which provides some evidence against

phylogenetically-biased exposure as the causative mechanism. However, the study

design in this case is not optimal for disentangling biased transmission probabilities

and biased exposure, as species were sampled from different locations at a single

timepoint and prevalence of the viruses varied spatially. Given this, drawing strong

conclusions as to the relative impact of the two mechanisms outlined above based on

this data would be premature. Similarly, the subgenus *Psithyrus* contains socially

parasitic species that are coevolved to parasitise particular social bumblebee species,

which could also lead to phylogenetically-correlated differences in viral infection

rates. We were unable to test whether socially parasitic cuckoo bumblebee species

have similar viral communities to their hosts, as our study included only a single

parasitic bumblebee species, *B. bohemicus*, but the possibility of brood parasitism

being an important driver of between-colony disease transmission is worth further

study.

Virus Evolutionary Interaction Effect and Coevolutionary Interaction

We found no evidence of a large virus evolutionary interaction or coevolutionary

interaction. This is largely unsurprising as it would appear implausible that the host

assemblages have been conserved over evolutionary time, as the deep splits in the

viral families predate the most recent common ancestor of bumblebees by many

millions of years (Koonin et al. 2008).

Non-phylogenetic Interaction

A non-phylogenetic interaction was detected. This interaction represents variation in

prevalence caused by specific host-virus pairings having prevalences beyond that

which would be expected by the simple addition of the individual host and virus

means. A non-phylogenetic interaction could be caused by a large range of factors,

some biological and some due to the specifics of the model, many of which would be

likely to be acting simultaneously to generate this signal. One possibility is

coevolution between the host and virus that occurred after both diverged from their

common ancestor with the closest related species in the study. Another is the

complete absence of coevolution, where spillover from a primary or group of primary

hosts causes either a constant very low prevalence of dead-end infections, which are

none-the-less detectable by PCR. Related to this is a statistical issue involving cases

where not every species in the study is within a virus' host range, and the species that

are within the host range are not closely related. In this case, the variation is not

absorbed by the host evolutionary interaction and almost no host has a prevalence

close to the mean across hosts, as in many species the prevalence is zero, which

causes the mean to be considerably lower than the average prevalence in the species

the virus does infect. This effect would be magnified if the sampling occurs during an

epizootic. More broadly, anything that changes the epidemiological parameters of a

virus in a specific host will lead to a non-phylogenetic interaction. Considering the

variation in the natural history of viruses and the lesser, but still significant, variation

in the natural history of bumblebees, a large non-phylogenetic interaction is to be

expected.

1007

Conclusion

While it is clear that viruses are abundant in pollinators, the factors that determine the

distribution of pollinator viruses have remained uncertain, outside of a few well-

studied cases (Fürst et al. 2014; McMahon et al. 2015). With the novel viruses

discovered in this study, we have investigated predictors of these virus/host

associations and found that both the host evolutionary history and the identity of the

virus contributes to this distribution. This supports both theory and prior empirical

evidence that related species are more at risk of infection from each other's diseases

than the diseases of distantly related species. However, the importance of the viral

identity and unique interactions between host-virus pairs suggests that the

introduction of a novel virus into a community is likely to have unpredictable effects

even when no close relatives of currently known hosts are present. This highlights the

risk posed by disease spillover for the conservation not only of wild pollinator

communities, but also to communities consisting of related animal or plant species in

general.

1023

1024 **Data Availability**

1025 The data and code for running the analyses is available on github under a GPLv3

1026 licence, as code uses code taken from other GPLv3 licenced works

1027 (https://github.com/dpascall/bumblebee-virus-cophylo).

1028

Acknowledgements

We thank Jarrod Hadfield for extensive statistical advice, modifications to

MCMCglmm and for helpful comments on this manuscript, Dave Goulson for

assistance identifying some specimens and to Ben Longdon and Bethany Clark for

further comments. Rowan Doff assisted in the lab with RFLP analyses and Claire

Webster with DNAse treatments. Claire Webster, Jarrod Hadfield and Florian Bayer

helped with fieldwork.

1036

1037   **Funding**

1038   This work was funded by a BBSRC SWBIO DTP PhD stipend to DP, a Royal Society

1039   Dorothy Hodgkin Fellowship to LW, Wellcome Trust Research Career Development

1040   Fellowship WT085064 to DJO.

1041

References

Altschul, S.F. et al., 1990. Basic local alignment search tool. *Journal of Molecular*

Biology, 215(3), pp.403–410.

Arbulo, N, Santos, E, Salvarrey, S, & Invernizzi, C., 2011. Proboscis length and

resource utilization in two ruguayan bumblebees: Bombus atratus Franklin and

Bombus bellicosus Smith (Hymenoptera: Apidae). *Neotropical Entomology*,

40(4), pp.483–488.

Arneberg, P. et al., 1998. Host densities as determinants of abundance in parasite

communities. *Proceedings of the Royal Society B: Biological Sciences*,

265(1403), pp.1283–1289.

Bailey, L. & Ball, B. V., 1991. *Honey Bee Pathology* 2nd Editio., London: Academic

Press Inc.

Bailey, L., Gibbs, A.J. & Woods, R.D., 1963. Two viruses from adult honey bees

(Apis mellifera Linnaeus). *Virology*, 21(3), pp.390–395.

Bailey, L. & Woods, R.D., 1974. Three previously undescribed viruses from the

honey bee. *Journal of General Virology*, 25(2), pp.175–186.

Bailey, L. & Woods, R.D., 1977. Two more small RNA viruses from honey bees and

1059  further observations on sacbrood and acute bee-paralysis viruses. *Journal of*

1060  *General Virology*, 37(1), pp.175–182.

1061 Baker, C. & Antonovics, J., 2012. Evolutionary determinants of genetic variation in

1062  susceptibility to infectious diseases in humans. *PLoS ONE*.

1063 Bandín, I. & Dopazo, C.P., 2011. Host range, host specificity and hypothesized host

1064  shift events among viruses of lower vertebrates. *Veterinary Research*, 42(67).

1065 Boccardo, G. et al., 1985. Three seedborne cryptic viruses containing double-stranded

1066  RNA isolated from white clover. *Virology*, 147(1), pp.29–40.

1067 Bouckaert, R. et al., 2014. BEAST 2: A Software Platform for Bayesian Evolutionary

1068  Analysis A. Prlic, ed. *PLoS Computational Biology*, 10(4), p.e1003537.

1069 Bouckaert, R. & Drummond, A., 2015. bModelTest: Bayesian phylogenetic site

1070  model averaging and model comparison. *bioRxiv*, p.020792.

1071 Brennecke, J. et al., 2007. Discrete Small RNA-Generating Loci as Master Regulators

1072  of Transposon Activity in Drosophila. *Cell*, 128(6), pp.1089–1103.

1073 Bronkhorst, A.W. et al., 2012. The DNA virus Invertebrate iridescent virus 6 is a

1074  target of the Drosophila RNAi machinery. *Proceedings of the National Academy*

1075  *of Sciences*, 109(51), pp.E3604–E3613.

1076 Buchfink, B., Xie, C. & Huson, D.H., 2015. Fast and sensitive protein alignment

1077  using DIAMOND. *Nature methods*, 12(1), pp.59–60.

1078 Cameron, S.A., Hines, H.M. & Williams, P.H., 2007. A comprehensive phylogeny of

1079  the bumble bees (Bombus). *Biological Journal of the Linnean Society*, 91(1),

1080  pp.161–188.

1081 Carpenter, B. et al., 2017. Stan: A probabilistic programming language. *Journal of*

1082  *Statistical Software*, 76(1), pp.1–32.

1083 Chandra, R.K., 1983. Nutrition, immunity, and infection: Present knowledge and

future directions. *The Lancet*, 321(8326), pp.688–691.

Chiba, S. et al., 2011. Widespread endogenization of genome sequences of non-

retroviral RNA viruses into plant genomes. *PLoS Pathogens*, 7(7).

Civitello, D.J. et al., 2015. Biodiversity inhibits parasites: Broad evidence for the

dilution effect. *Proceedings of the National Academy of Sciences*, 112(28),

pp.8667–8671.

Cooper, N. et al., 2012. Phylogenetic host specificity and understanding parasite

sharing in primates. *Ecology Letters*, 15(12), pp.1370–1377.

Cornman, R.S. et al., 2012. Pathogen webs in collapsing honey bee colonies. *PLoS

ONE*, 7(8), pp.1–21.

Curtis, V.A., 2014. Infection-avoidance behaviour in humans and other animals.

Trends in Immunology, 35(10), pp.457–464.

Davies, T.J. & Pedersen, A.B., 2008. Phylogeny and geography predict pathogen

community similarity in wild primates and humans. *Proceedings of the Royal

Society B: Biological Sciences*, 275(1643), pp.1695–1701.

Deleris, A. et al., 2006. Hierarchical action and inhibition of plant dicer-like proteins

in antiviral defense. *Science*, 313(5783), pp.68–71.

Drummond, A.J. et al., 2012. Bayesian phylogenetics with BEAUti and the BEAST

1.7. *Molecular Biology and Evolution*, 29(8), pp.1969–1973.

Drummond, A.J. et al., 2006. Relaxed phylogenetics and dating with confidence D.

Penny, ed. *PLoS Biology*, 4(5), pp.699–710.

Durrer, S. & Schmid-Hempel, P., 1994. Shared Use of Flowers Leads to Horizontal

Pathogen Transmission. *Proceedings of the Royal Society B: Biological

Sciences*, 258(1353), pp.299–302.

Ebert, T.A., Brlansky, R. & Rogers, M., 2010. Reexamining the Pooled Sampling

Approach for Estimating Prevalence of Infected Insect Vectors. *Annals of the Entomological Society of America*, 103(6), pp.827–837.

Fauquet, C.M. & Stanley, J., 2005. Revising the way we conceive and name viruses below the species level: A review of geminivirus taxonomy calls for new standardized isolate descriptors. *Arch Virol*, 150, pp.2151–2179.

Félix, M.A. et al., 2011. Natural and experimental infection of Caenorhabditis nematodes by novel viruses related to nodaviruses J. Hodgkin, ed. *PLoS Biology*, 9(1), p.e1000586.

Fenton, A. et al., 2015. Are All Hosts Created Equal? Partitioning Host Species Contributions to Parasite Persistence in Multihost Communities. *The American Naturalist*, 186(5), pp.610–622.

Ferreira, M.A.R. & Suchard, M.A., 2008. Bayesian analysis of elapsed times in continuous-time Markov chains. *Canadian Journal of Statistics*, 36(3), pp.355–368.

Finn, R.D. et al., 2014. Pfam: The protein families database. *Nucleic Acids Research*, 42(D1), pp.D222–D230.

Fürst, M. a. et al., 2014. Disease associations between honeybees and bumblebees as a threat to wild pollinators. *Nature*, 506(7488), pp.364–366.

Garibaldi, L.A. et al., 2013. Wild pollinators enhance fruit set of crops regardless of honey bee abundance. *Science*, 340(6127), pp.1608–1611.

Gibbs, A.J. & Gower, J.C., 1960. The use of a multiple-transfer method in plant virus transmission studies - Some statistical points arising in the analysis of results. *Annals of Applied Biology*, 48(1), pp.75–83.

Goodwin, S., 1995. Seasonal phenology and abundance of early-, mid-and long-season bumble bees in southern England. *Journal of Apicultural Research*, 34(2),

pp.79–87.

Gorbalenya, A.E. et al., 2002. The palm subdomain-based active site is internally

permuted in viral RNA-dependent RNA polymerases of an ancient lineage.

Journal of Molecular Biology, 324(1), pp.47–62.

Goulson, D. et al., 2005. Causes of rarity in bumblebees. *Biological Conservation*,

122(1), pp.1–8.

Goulson, D. & Darvill, B., 2004. Niche overlap and diet breadth in bumblebees; are

rare species more specialized in their choice of flowers? *Apidologie*, 35(1),

pp.55–63.

Goulson, D., Lye, G.C. & Darvill, B., 2008. Diet breadth, coexistence and rarity in

bumblebees. *Biodiversity and Conservation*, 17(13), pp.3269–3288.

Grabherr, M.G. et al., 2011. Full-length transcriptome assembly from RNA-Seq data

without a reference genome. *Nature Biotechnology*, 29(7), pp.644–652.

Graystock, P., Goulson, D. & Hughes, W.O.H., 2015. Parasites in bloom: flowers aid

dispersal and transmission of pollinator parasites within and between bee

species. *Proceedings of the Royal Society B-Biological Sciences*, 282(1813),

p.20151371.

Hadfield, J.D. et al., 2014. A tale of two phylogenies: comparative analyses of

ecological interactions. *The American Naturalist*, 183(2), pp.174–87.

Harder, L.D., 1985. Morphology as a predictor of flower choice by bumble bees.

Ecology, 66(1), pp.198–210.

Harmon-Threatt, A.N. & Ackerly, D.D., 2013. Filtering across Spatial Scales:

Phylogeny, Biogeography and Community Structure in Bumble Bees. *PLoS*

ONE, 8(3), p.e60446.

Heesterbeek, J.A.P., 2002. A brief history of R0 and a recipe for its calculation. *Acta*

Biotheoretica, 50, pp.189–204.

Heled, J. & Drummond, A.J., 2012. Calibrated tree priors for relaxed phylogenetics

and divergence time estimation. *Systematic Biology*, 61(1), pp.138–149.

Henikoff, S. & Henikoff, J.G., 1992. Amino acid substitution matrices from protein

blocks. *Proceedings of the National Academy of Sciences of the United States of*

America, 89(22), pp.10915–10919.

Hines, H.M., 2008. Historical Biogeography, Divergence Times, and Diversification

Patterns of Bumble Bees (Hymenoptera: Apidae: Bombus). *Systematic Biology*,

57(1), pp.58–75.

Holmes, E.C., 2003. Molecular Clocks and the Puzzle of RNA Virus Origins. *Journal*

of Virology, 77(7), pp.3893–3897.

Huang, S. et al., 2014. Phylogenetically related and ecologically similar carnivores

harbour similar parasite assemblages. *Journal of Animal Ecology*, 83(3), pp.671–

680.

Hueffer, K. et al., 2003. The Natural Host Range Shift and Subsequent Evolution of

Canine Parvovirus Resulted from Virus-Specific Binding to the Canine

Transferrin Receptor The Natural Host Range Shift and Subsequent Evolution of

Canine Parvovirus Resulted from Virus-Specific Bind. *Journal of Virology*,

77(3), pp.1718–1726.

Inouye, D.W., 1978. Resource Partitioning in Bumblebees: Experimental Studies of

Foraging Behavior. *Ecology*, 59(4), pp.672–678.

Johnson, M.B. et al., 2011. Parasite transmission in social interacting hosts:

Monogenean epidemics in guppies. *PLoS ONE*, 6(8).

Kamer, G. & Argos, P., 1984. Primary structural comparison of RNA-dependent

polymerases from plant, animal and bacterial viruses. *Nucleic Acids Research*,

12(18), pp.7269–7282.

Kanehisa, M. et al., 2002. The KEGG databases at GenomeNet. *Nucleic Acids*

Research, 30(1), pp.42–6.

Katoh, K. et al., 2005. MAFFT version 5: improvement in accuracy of multiple

sequence alignment. *Nucleic Acids Research*, 33(2), pp.511–518.

Katoh, K., Rozewicki, J. & Yamada, K.D., 2017. MAFFT online service: multiple

sequence alignment, interactive sequence choice and visualization. *Briefings in*

Bioinformatics.

Katzourakis, A. & Gifford, R.J., 2010. Endogenous viral elements in animal genomes.

PLoS Genetics, 6(11), p.e1001191.

Koloniuk, I., Přibylová, J. & Fránová, J., 2018. Molecular characterization and

complete genome of a novel nepovirus from red clover. *Archives of Virology*,

163(5), pp.1387–1389.

Koonin, E. V., Dolja, V. V. & Krupovic, M., 2015. Origins and evolution of viruses

of eukaryotes: The ultimate modularity. *Virology*, 479–480, pp.2–25.

Koonin, E. V et al., 2008. The Big Bang of picorna-like virus evolution antedates the

radiation of eukaryotic supergroups. *Nature Reviews Microbiology*, 6(12),

pp.925–939.

Langmead, B. & Salzberg, S.L., 2012. Fast gapped-read alignment with Bowtie 2.

Nature Methods, 9(4), pp.357–359.

Langsjoen, R.M. et al., 2018. Chikungunya virus strains show lineage-specific

variations in virulence and cross-protective ability in murine and nonhuman

primate models. *mBio*.

Lewis, S.H. et al., 2018. Pan-arthropod analysis reveals somatic piRNAs as an

ancestral defence against transposable elements. *Nature Ecology and Evolution*,

2(1), pp.174–181.

Liu, Y., Gelman, A. & Zheng, T., 2015. Simulation-efficient shortest probability

intervals. *Statistics and Computing*, 25(4), pp.809–819.

Longdon, B. et al., 2011. Host phylogeny determines viral persistence and replication

in novel hosts. *PLoS Pathogens*, 7(9), p.e1002260.

Longdon, B. et al., 2018. Host shifts result in parallel genetic changes when viruses

evolve in closely related species. *PLoS Pathogens*.

Longdon, B. et al., 2015. The causes and consequences of changes in virulence

following pathogen host shifts. *PLoS Pathogens*, 11(3), p.e1004728.

Longdon, B. et al., 2014. The Evolution and Genetics of Virus Host Shifts. *PLoS*

Pathogens, 10(11), p.e1004395.

Lye, G.C. et al., 2010. Forage use and niche partitioning by non-native bumblebees in

New Zealand: Implications for the conservation of their populations of origin.

Journal of Insect Conservation, 14(6), pp.607–615.

Manley, R., Boots, M. & Wilfert, L., 2015. Emerging viral disease risk to pollinating

insects: Ecological, evolutionary and anthropogenic factors. *Journal of Applied*

Ecology, 52(2), pp.331–340.

McArt, S.H. et al., 2014. Arranging the bouquet of disease: Floral traits and the

transmission of plant and animal pathogens. *Ecology Letters*, 17(5), pp.624–636.

McMahon, D.P. et al., 2015. A sting in the spit: Widespread cross-infection of

multiple RNA viruses across wild and managed bees. *Journal of Animal*

Ecology, 84(3), pp.615–624.

van Mierlo, J.T. et al., 2012. Convergent Evolution of Argonaute-2 Slicer Antagonism

in Two Distinct Insect RNA Viruses. *PLoS Pathogens*, 8(8), p.e1002872.

de Miranda, J.R. et al., 2015. Genome characterization, prevalence and distribution of

a macula-like virus from Apis mellifera and Varroa destructor. *Viruses*, 7(7),

pp.3586–3602.

Mordecai, G.J. et al., 2015. Diversity in a honey bee pathogen: first report of a third

master variant of the Deformed Wing Virus quasispecies. *The ISME Journal*,

10(5), pp.1–10.

Murray, T.E. et al., 2008. Cryptic species diversity in a widespread bumble bee

complex revealed using mitochondrial DNA RFLPs. *Conservation Genetics*,

9(3), pp.653–666.

Pei, J., Kim, B.H. & Grishin, N. V., 2008. PROMALS3D: A tool for multiple protein

sequence and structure alignments. *Nucleic Acids Research*, 36(7), pp.2295–

2300.

Rambaut, A. et al., 2017. Tracer v1.7. Available at: http://beast.bio.ed.ac.uk/Tracer.

Rao, S., Poinar, G. & Henley, D., 2017. A scientific note on rare parasitism of the

bumble bee pollinator, Bombus impatiens, by a mermithid nematode,

Pheromermis sp. (Nematoda: Mermithidae). *Apidologie*, 48(1), pp.75–77.

Remnant, E.J. et al., 2017. A Diverse Range of Novel RNA Viruses in Geographically

Distinct Honey Bee Populations. *Journal of Virology*, (May), p.JVI.00158-17.

van Rij, R.P. et al., 2006. The RNA silencing endonuclease Argonaute 2 mediates

specific antiviral immunity in Drosophila melanogaster. *Genes and*

Development, 20(21), pp.2985–2995.

van Riper, C. et al., 1986. The Epizootiology and ecological significance of malaria in

Hawaiian land birds. *Ecological Monographs*, 56(4), pp.327–344.

Roberts, J.M.K., Anderson, D.L. & Durr, P.A., 2018. Metagenomic analysis of

Varroa-free Australian honey bees (Apis mellifera) shows a diverse

Picornavirales virome. *Journal of General Virology*, 99, pp.818–826.

Roberts, K. et al., 2018. Changes in temperature alter susceptibility to a virus
    following a host shift. *bioRxiv*.

Ruiz-González, M.X. et al., 2012. Dynamic transmission, host quality, and population
    structure in a multihost parasite of bumblebees. *Evolution*, 66(10), pp.3053–
    3066.

Runckel, C. et al., 2011. Temporal analysis of the honey bee microbiome reveals four
    novel viruses and seasonal prevalence of known viruses, Nosema, and Crithidia.
    PLoS ONE, 6(6), p.e20656.

Salathé, R.M. & Schmid-Hempel, P., 2011. The Genotypic Structure of a Multi-Host
    Bumblebee Parasite Suggests a Role for Ecological Niche Overlap. *PLOS ONE*,
    6(8), p.e22054.

Schoonvaere, K. et al., 2018. Study of the Metatranscriptome of Eight Social and
    Solitary Wild Bee Species Reveals Novel Viruses and Bee Parasites. *Frontiers in
    Microbiology*, 9, p.177.

Schoonvaere, K. et al., 2016. Unbiased RNA Shotgun Metagenomics in Social and
    Solitary Wild Bees Detects Associations with Eukaryote Parasites and New
    Viruses R. Lu, ed. *PLOS ONE*, 11(12), p.e0168456.

Shi, M. et al., 2016. Redefining the invertebrate RNA virosphere. *Nature*, 540, pp.1–
    12.

Smith, K.M. & Markham, R., 1944. Two new viruses affecting tobacco and other
    plants. *Phytopathology*, 34, pp.324–329.

Stan Development Team, 2017. *Stan Modeling Language: User's Guide and
    Reference Manual* 2.17.0.

Streicker, D.G. et al., 2012. Variable evolutionary routes to host establishment across
    repeated rabies virus host shifts among bats. *Proceedings of the National

Academy of Sciences.

Suttle, C.A., 2007. Marine viruses--major players in the global ecosystem. *Nature*

Reviews Microbiology, 5(10), pp.801–812.

Suzuki, Y. et al., 2017. Uncovering the repertoire of endogenous flaviviral elements

in Aedes. *Journal of Virology*.

Thompson, K.H., 1962. Estimation of the Proportion of Vectors in a Natural

Population of Insects. *Biometrics*, 18(4), pp.568–578.

Untergasser, A. et al., 2012. Primer3-new capabilities and interfaces. *Nucleic Acids*

Research, 40(15), p.e115.

Vanbergen, A.J. & the Insect Pollinators Initiative, 2013. Threats to an ecosystem

service: Pressures on pollinators. *Frontiers in Ecology and the Environment*,

11(5), pp.251–259.

Wang, X.-B. et al., 2011. The 21-Nucleotide, but Not 22-Nucleotide, Viral Secondary

Small Interfering RNAs Direct Potent Antiviral Defense by Two Cooperative

Argonautes in Arabidopsis thaliana. *The Plant Cell*, 23(4), pp.1625–1638.

Waxman, D. et al., 2014. Inferring Host Range Dynamics from Comparative Data:

The Protozoan Parasites of New World Monkeys. *The American Naturalist*,

184(1), pp.65–74.

Webster, C.L. et al., 2015. The discovery, distribution, and evolution of viruses

associated with drosophila melanogaster. *PLoS Biology*, 13(7), p.e1002210.

Williams, P.H.P.H.P.H. & Osborne, J.L., 2009. Bumblebee vulnerability and

conservation world-wide. *Apidologie*, 40(3), pp.367–387.

Woolhouse, M.E.J. & Gowtage-Sequeria, S., 2005. Host range and emerging and

reemerging pathogens. *Emerging Infectious Diseases*, 11(12), pp.1842–1847.

Zanotto, P.M. et al., 1996. A reevaluation of the higher taxonomy of viruses based on

RNA polymerases. *Journal of Virology*, 70(9), pp.6083–6096.

1310