Analysis of the structural variability of topologically associated domains as revealed by Hi-C

Natalie Sauerwald[1], Akshat Singhal[2], and Carl Kingsford[1,*]

[1] Computational Biology Department, Carnegie Mellon University, Pittsburgh, PA 15213, USA

[2] Department of Computer Science, Stony Brook University, Stony Brook, NY 11790, USA

[*] to whom correspondence should be addressed: carlk@cs.cmu.edu

Abstract

Three-dimensional chromosome structure plays an integral role in gene expression and regulation, replication timing, and other cellular processes. Topologically associating domains (TADs), one of the building blocks of chromosome structure, are genomic regions with higher contact frequencies within the region than outside the region. A central question is the degree to which TADs are conserved or vary between conditions. We analyze a set of 137 Hi-C samples from 9 different studies under 3 measures in order to quantify the effects of various sources of biological and experimental variation. We observe significant variation in TAD sets between both non-replicate and replicate samples, and show that this variability does not seem to come from genetic sequence differences. The effects of experimental protocol differences are also measured, demonstrating that samples can have protocol-specific structural changes, but that TADs are generally robust to lab-specific differences. This study represents a systematic quantification of the key factors influencing comparisons of chromosome structure.

While it is recognized that the three-dimensional structure of the chromosome is an integral part of many key genomic functions, we lack a full understanding of the variability of this structure across biological sources or experimental conditions. Changes in chromosome structure at specific genomic regions and under certain conditions have been implicated in a variety of human diseases

and disabilities, including many cancers [26, 28, 18, 19], deformation or malformation of limbs during development [25], and severe brain anomalies [41], but we do not have a sense of the expected variation of TADs genome-wide. In healthy cells, genome shape is heavily linked to key processes such as gene regulation and expression [8, 5, 22, 14, 36], replication timing [37, 29, 33, 2], and DNA accessibility and nuclear organization [44, 34, 6].

Features of genome-wide three-dimensional chromosome structure can be measured by Hi-C [24], a high-throughput variant of the chromosome conformation capture protocol [9]. The experiment involves cross-linking and ligating spatially close genomic segments, then aligning them back to the genome to find their genomic positions. The output of this experiment is a matrix in which the rows and columns represent segments of the genome along a chromosome, and each matrix entry records the pairwise interaction frequency of the genome fragments of the associated row and column. These values reflect 3D proximity, quantifying the frequency of contact between every pair of genomic segments.

A hierarchical architecture has emerged from these Hi-C matrices, in which chromosome structure is composed of several different scales of components, from multi-megabase compartments to sub-megabase topologically-associating domains (TADs) and sub-TADs [15, 4]. TADs represent chromosomal regions with significantly higher interaction frequency among segments within the TAD than with those outside of it [11]. TADs are considered to be a primary structural building block of chromosome architecture [13], and several methods have been developed to computationally identify them [11, 35, 7, 43, 15, 31].

One challenge in the interpretation of TADs is that we have little understanding of the variability of TAD structures under different conditions. As more Hi-C data has become available recently, it is now possible to perform a substantial analysis of the relative consistency or variability of TADs across a variety of human cell conditions. Previous work has suggested that TADs are largely conserved across human cell types and possibly even species, however the degree of this conservation is unclear and has been tested in only a small set of samples [11, 35].

We quantify the influence of both technical and biological variation on TAD structures across several experimental and biological conditions. We observe that approximately 10–70% of combined TAD boundaries differ between replicates, regardless of sequencing depth or contact coverage, pointing to a potentially dynamic or disordered arrangement. Across 69 samples of different cell

lines and tissue types, we observe ~20–80% unshared TAD boundaries, suggesting that there can be fairly large differences in TAD sets across biological conditions. We find that samples of the same cell or tissue type have elevated structural similarities, suggesting that biological function is a key driver of structural similarity. By analyzing the structural similarity of sets of parents and their children, as well as tissue samples taken from different individuals, we observe that the genetic sequence differences between individuals and the genetic sequence similarities between parents and their children have little impact on TAD structural similarity. Of the possible sources of technical variation considered in this work, the choice of *in situ* (in nucleus) ligation versus dilution (in solution) ligation protocols has the greatest influence on Hi-C and TAD structures. In contrast, we demonstrate that Hi-C measurements and corresponding TAD calls are robust to other technical differences such as the choice of restriction enzyme and lab-specific variations.

Results

Data and comparison measures

We analyzed 69 publicly available human Hi-C samples (137 including replicates) from nine different studies [11, 35, 42, 10, 21, 12, 46, 17, 39] covering 52 unique biological sources and a variety of experimental conditions (Table 1). All samples were processed from raw reads (.fastq files) to normalized Hi-C matrices at 100kb resolution through the HiC-Pro pipeline [40]. TADs were computed on all resulting Hi-C matrices using Armatus [15].

Pairwise similarity of these 69 samples (2346 pairs plus 83 replicate pairs) was quantified with three different similarity measures. HiCRep [45] computes a statistical similarity score on Hi-C matrices directly. The Jaccard Index (JI) quantifies the similarity between two sets, in this case defined as sets of TAD boundaries, returning a value that represents the fraction of TAD boundaries shared between the two samples. TADsim is a measure adapted from a method that identifies structurally similar regions between two TAD sets [38]; it quantifies the fraction of the genome covered by similar TAD structures using the variation of information (VI) metric [27]. See the Online Methods section for details on each comparison measure.

Distributions of similarity values are compared using the Mann-Whitney U test, which is a non-parametric statistical test quantifying the probability that two independent samples were drawn

from populations with the same underlying distribution. More specifically, the alternative hypothesis for this test is that one distribution is greater than the other. A low p-value suggests that one of the two distributions is likely drawn from a distribution with a higher median than the other; we will use this to test whether various conditions reflect elevated structural similarity, relative to a background or other comparison distribution.

Structural similarity of replicate samples

By quantifying the similarity of all 83 replicate pairs in our dataset, we find that the TAD sets of replicate pairs are significantly more similar than those of non-replicate pairs (Figure 1**a**–**c**, $p < 10^{-20}$ for all comparison measures), despite some previous work that suggested much lower TAD reproducibility between replicates [16]. Forcato *et al.* [16] reported an average replicate JI value of below 0.3 for TAD sets generated by Armatus, which is much lower than the average JI value of our data at 0.6277. Among the samples studied here, replicates share an average of 62.77% of their combined TAD boundaries, which is consistent with previous studies on different data using different methods (Dixon *et al.* [11]: 62.28%, 73.73% and Rao *et al.* [35]: 61.88%). This leaves almost 40% of TAD boundaries that vary across replicates. Between non-replicate pairs, almost 60% TAD boundaries are not shared on average. This variability could not be explained by experimental limitations, as we found that reproducibility is only weakly correlated with sequencing coverage (see Figure S5). This therefore points to a dynamic or disordered structure, as suggested by a recent imaging study [32].

Variability across tissues and individuals

The chromosome structure of tissue samples has not been as widely studied as that of cell lines, but these structures may provide valuable insight into tissue-specific genomics. Among our set of 69 Hi-C experiments, 13 different human tissues are represented, and there are 16 pairs of the same tissue type taken from different donor individuals. The similarity values of the chromosome structures of these pairs are statistically indistinguishable from those of replicate samples (Figure 2**a**–**c**; HiCRep: $p = 0.4792$, JI: $p = 0.1300$, TADsim: $p = 0.09559$). There is much less variation across individuals than across tissue types (Figure 2**a**–**c**; HiCRep: $p = 1.5577 \times 10^{-6}$, JI: $p = 3.876 \times 10^{-6}$, TADsim: $p = 1.017 \times 10^{-7}$), suggesting that individual genetic differences have less influence on chromosome

Figure 1: **Hi-C and TAD reproducibility.** The violin plots show distributions of **a:** HiCRep, **b:** Jaccard Index, and **c:** TADsim values on pairs of either replicates or non-replicates. All of these plots show a statistically significant ($p < 10^{-20}$) elevation of similarity among replicate pairs, demonstrating that both Hi-C matrices and TADs are reproducible.

structure than the biological function of the sample.

We observe some conservation of chromosome structure across different tissue types, as reported by Schmitt *et al.* [39]. In order to determine whether TAD structure is more similar across tissues than across cell lines, we compared the similarities among tissue types to the background distribution consisting of all non-replicate pairs with at least one cell line. Two of our three measures suggest that there is elevated conservation among tissues, but the two TADsim distributions are statistically similar (Figure 2**a**–**c**, HiCRep: $p = 2.120 \times 10^{-19}$, JI: $p = 0.004806$, TADsim: $p = 0.4235$). The average JI value across tissue samples of 41.6% implies that while there is a significant level of similarity among chromosome structures of different tissue types, close to 60% of TAD boundaries vary across different tissue samples. This level of variability across tissue types may indicate the existence of tissue-specific structural features.

TAD structure is largely invariant across family relationships

Hi-C measurements from blood lymphocyte cells of three parent-parent-child triplets (trios) permit a glimpse into the heritability of chromosome structure. We find that unrelated individuals (parents) share just as much chromosomal structural similarity as each parent and their child (Figures 2**d**–**f**). We therefore see no evidence that chromosome structure is an especially heritable trait, at

Figure 2: **Biological sources of TAD variation. a,b,c:** Comparisons between and within tissue samples using **a** HiCRep, **b** JI and **c** TADsim. Each figure shows four violin plots representing distributions of similarity values of the background (non-replicate pairs), replicates, pairs from the same tissue type but different donor individuals, and pairs from different tissue types. **d,e,f:** Comparisons with three trio samples of blood lymphocyte cells using **d** HiCRep, **e** JI and **f** TADsim. The background distribution consists of all non-replicate pairs, and the blood lymphocyte pair distribution shows all similarity values of two blood lymphocyte samples outside of the trios. The trio replicates refer to the similarity values of the replicate pairs from within each individual of the trio samples. The scattered points on the right side of each figure represent all within-trio comparisons, colored by family relationship.

least in blood lymphocyte cells. The similarity values within trios are generally much higher than the background of non-replicate comparisons, however they are similar to the distribution of pairs of blood lymphocytes, so this is likely a result of the shared cell type rather than genetic similarity. As with the tissue data, the biological source (cell or tissue type) seems to be a much stronger driver of structural similarity than genetic similarity.

Variations across Hi-C protocols

In order to investigate technical sources of variation, we compare several common variations in the Hi-C protocol, and test whether they affect the similarity of the TADs that are identified. There are two main protocol variants that differ in the cross-linkage step. *In situ* Hi-C [35] (also termed "in nucleus Hi-C" [30]) involves cross-linking the DNA within the nucleus, while dilution Hi-C (or "in solution Hi-C") performs cross-linking in a dilute solution. Each protocol also requires the choice of a restriction enzyme, which could be any of four common options: HindIII, MboI, NcoI, and DpnII. While there has been some study of the differences in Hi-C data resulting from *in situ* and dilution protocols [35, 30], the influence on TAD sets and the effect of the restriction enzyme had not been systematically studied previously.

In situ and dilution Hi-C reproducibility

Across all replicate pairs (12 in situ, 71 dilution), the intra-chromosomal Hi-C matrices of *in situ* replicates are statistically significantly more similar than dilution replicates (Figure **3a**, $p = 1.180 \times 10^{-5}$). However, the TAD sets of *in situ* replicates only show statistically significantly higher similarity than those of dilution replicates under the JI measure (Figures **3b,c**; JI: $p = 0.02703$, TADsim: $p = 0.1547$). TADs capture only relatively short-range interactions, and it therefore appears that the difference between *in situ* and dilution Hi-C is not as significant a factor in TAD reproducibility as in overall Hi-C matrix reproducibility. It has been previously shown that *in situ* Hi-C matrices are more reproducible than dilution Hi-C matrices [35, 30], specifically with respect to long-range and inter-chromosomal contacts.

Figure 3: **Comparing Hi-C samples generated from the *in situ* and dilution protocols. a:** HiCRep shows that *in situ* Hi-C matrices are more reproducible than dilution matrices ($p < 0.0005$). **b,c:** TAD set reproducibility according to JI ($p = 0.02703$) and TADsim ($p = 0.1547$) shows that protocol choice has less of an impact on reproducibility of TAD sets than full Hi-C matrices. **d,e,f:** Comparisons of same cell type pairs generated by the same and different protocols. The background distribution is all comparisons of different cell types. Under all measures, there is a clear and statistically significant ($p < 0.05$) drop in similarity values of samples generated by different protocols compared to samples generated by the same protocol.

Comparing *in situ* and dilution samples

In order to study whether both *in situ* and dilution protocols result in the same structures, we compared samples across protocols. Among pairs of the same cell type, mixed protocol pairs, where one sample came from *in situ* and one from dilution, were consistently statistically significantly less similar than the single protocol pairs, in which both samples came from the same protocol (Figures 3d,e,f, HiCRep: $p = 0.0003423$, JI: $p = 0.03967$, TADsim: $p = 0.02002$). The chromosomal structures identified from these two protocol variants are therefore not entirely consistent, although pairs from the same cell type still showed more similarity than pairs of different cell types, even among mixed protocol pairs (HiCRep: $p = 3.436 \times 10^{-15}$, JI: $p = 1.2645 \times 10^{-9}$, TADsim: $p = 0.006010$). We observe a similar trend across all non-replicate pairs as well (Figure S6). Overall, we observe some structural differences induced by the protocol variations, but not enough to obscure the general similarities expected from samples of the same cell type.

Restriction enzyme choice has minimal impact on TAD sets

By comparing samples from the same lab of the same cell type, generated with a different restriction enzyme, we see no significant variation in similarity measures induced by restriction enzymes, as shown in Figure 4. As expected, the pairs of the same cell type with a different restriction enzyme tend to be more structurally similar than the background distribution, which includes all 2333 other pairwise comparisons. The choice of restriction enzyme does not appear to be a significant source of technical variation in measurements of chromosome structure, as both Hi-C matrices and TAD sets appear robust to this experimental variable.

TAD variation induced by lab-specific differences

Variation induced by differences in labs or experimenters, rather than true biological differences, can be an obstacle to meaningful and accurate analysis of biological data. Across all of our data, we see no pattern of elevated structural similarity among samples from the same lab (Figure 5a, JI and TADsim heat maps can be seen in Figures S7 and S8). A comparison of pairs of the same cell type from different labs shows that these pairs are generally more similar than non-replicate pairs, with similarity values near those of replicate pairs (Figures 5b,c,d). Consistent with the protocol-

Figure 4: **Measurements of structural similarity across the use of different restriction enzymes.** The scattered points represent the similarity of a pair of samples of the same cell type (grey is hESC, red is HFF-hTERT), generated by using different restriction enzymes. The violin plot shows the distribution of all other non-replicate cmoparisons. As expected, the points that differ only in restriction enzyme are largely more similar than the background, suggesting that this choice is not a significant source of technical variability.

driven variation described above, the three lowest pairwise scores for IMR90 in both JI and TADsim are the three mixed protocol comparisons; all other points represent pairs generated by the same protocol. Chromosome structure seems to be robust to the variability across experimental labs.

Discussion

We have demonstrated that cell or tissue type, rather than individual or genetic difference, appears to be the greatest driver of biological variation in TAD structures and Hi-C matrices. Between replicates, TAD structures are shown to share only 60% of their boundaries, suggesting that chromosome structure is not a static feature, but remains variable even in identical cell populations. We observe elevated similarities between samples of the same cell type, suggesting that TAD structures are likely correlated with cellular function, rather than individual genetics. The largest differences due to technical variations appeared in comparing structures generated through *in situ* or dilution protocols, while lab-specific differences and restriction enzyme choices had a smaller impact on the resulting similarities of Hi-C measurements.

In order to maximize the number of samples analyzed in this work, all comparisons were per-

Figure 5: **Quantifying variation across samples from different labs. a** Summary of all 2346 pairwise sample comparisons as a heatmap of HiCRep scores. The dotted lines outline groups of samples from the same study. **b,c,d**: The effects of lab-specific variation on chromosome structure measurements. The points represent similarity scores of the same cell type (red for IMR90, grey for hESC) from different studies. These can be compared to the distribution of non-replicate pairs and that of replicate pairs, showing that samples from different labs achieve similarity values near those of replicate pairs.

formed at a fairly low resolution of 100kb, so structural features that would be clearer at higher resolution may have been overlooked. A few observations noted in this work are consistent with previous smaller-scale studies of higher-resolution matrices. In particular, the similarities between replicates that we observed were consistent with those found in Dixon *et al.* [11] and Rao *et al.* [35] and the similarity across tissue types was also found by Schmitt *et al.* [39].

There are still relatively few available Hi-C data sets compared with other genomic analyses, and many of the observations made here would be strengthened with more samples or with confirmation through single-cell Hi-C. In particular, more trios from other cell types would help to confirm whether there is truly no elevated similarity in genetically related individuals, or whether this conclusion was specific to the blood lymphocyte samples studied in this work. As more single-cell Hi-C data becomes available and analysis methods improve, cell-to-cell variations in chromosome structure will be easier to assess, and we will be able to determine whether these population trends hold within individual cells.

All samples studied here were processed from sequencing reads to Hi-C matrices through the same pre-processing pipeline, and all TADs were computed using Armatus [15]. These choices may have influenced the trends we observed in this work, because different pre-processors, Hi-C normalization methods, or TAD callers could result in different patterns in the resulting structural measurements. The consistencies with previous work using different methods for each of these steps suggests that they did not have a major effect, but more study is needed to assess the overall robustness of Hi-C measurements to these processing choices. Additionally, there may be other possible comparison methods for Hi-C matrices and TAD sets, which may or may not agree with the three measures used here.

Further study of the structural differences across cell types may lead to insights into the mechanisms of chromosome structure. These comparison techniques could also be used to determine the differences between chromosome structures in healthy and diseased cells and could point to the locations of structural changes that are present across diseased cells. There is already significant evidence of structural abnormalities in many diseases (review, [25]). Additional systematic, genome-wide analyses of TAD structures could increase our understanding of a range of human diseases.

This work compares Hi-C data and TAD structures from nine studies using three different

measures, in order to identify trends in the variables controlling chromosomal structural similarity. We observe that even replicates display a certain amount of variability in chromosome structure. Chromosome structure appears most conserved within cell types and tissue types and not influenced more strongly by genetic similarity or differences across individuals. Differences in the cross-linkage step of the Hi-C protocol can induce variation in the resulting Hi-C and TAD measurements, but they seem robust to both lab-specific differences and choice of restriction enzyme.

The scripts to reproduce the analysis performed here are available at `https://github.com/Kingsford-Group/localtadsim/tree/master/analysis`.

Online Methods

Data

A total of 76 human Hi-C samples were processed from sequencing reads (.fastq files) downloaded from various publicly available sources (Sequence Read Archive (SRA) [23], ENCODE [42], Gene Expression Omnibus (GEO) [3], or 4DN portal [10]). Normalized Hi-C matrices were computed from the reads through the HiC-Pro pipeline [40], and each sample was tested for quality at 100kb resolution. Using the criteria suggested by Ay and Noble [1] and Rao *et al.* [35] (at least 80% of all bins must contain more than 1000 contacts), we found 7 experiments which could not be analyzed at 100kb resolution or less (Table S1), leaving 69 human Hi-C data sets (137 including all replicate samples) representing 52 unique cell types or biological sources from 9 studies. The details of these experiments, including accession numbers, are found in Table 1. All samples were normalized using iterative correction and eigenvector decomposition (ICE) [20], and all analyses presented here were performed at 100kb resolution. For analyses that do not explicitly compare replicates, all aligned reads from each replicate of a sample were merged and processed into a single combined Hi-C matrix for optimal data quality.

From the Hi-C matrices, TADs were computed using Armatus [15], a commonly used method for identifying TADs efficiently. Armatus takes one parameter γ, which controls the expected TAD size. For every sample and chromosome, Armatus was run with γ values ranging from 0 to 1 at intervals of 0.1, and the γ value was chosen to ensure TADs as close as possible to the expected median TAD size of 880kb [4] on each sample and chromosome.

Comparison measures

In order to comprehensively compare chromosome structures, we use three different measures: HiCRep [45], Jaccard Index (JI), and TADsim [38]. HiCRep measures similarity between Hi-C matrices directly, and both JI and TADsim compare similarity of predicted TADs. All three measures were computed on all 2346 pairs of non-replicate samples, in addition to all 83 replicate pairs.

HiCRep was designed to assess the reproducibility of replicates or the similarity of two Hi-C matrices. This measure uses a stratum-adjusted correlation coefficient to reliably compute a statistical similarity score between two Hi-C matrices, explicitly accounting for both the strong distance dependence found in Hi-C and the known domain structure [45]. This method returns a value that represents the overall similarity of the full Hi-C matrix, and distinguishes between replicate and non-replicate samples significantly better than simple correlation coefficients. We ran HiCRep on all intra-chromosomal matrices of our samples and averaged over all chromosomes to get a single value per cell type pair. HiCRep requires a smoothing parameter h, which was selected for each comparison according to the heuristic optimization procedure provided by the software, which chooses the minimum h value at which the score begins to converge. We allow a range of 0 to 3, which is expanded from the 0 to 2 range shown in the documentation example, to allow more options while maintaining computational efficiency.

The Jaccard Index (JI), a simple set similarity metric, is defined as the size of the intersection of two sets A and B divided by the size of the union of the sets: $JI(A, B) = |A \cap B|/|A \cup B|$. When comparing TADs, the two sets A and B represent the two lists of TAD boundaries, as used in Forcato *et al.* [16]. The resulting JI value is an easily interpretable number representing the fraction of shared boundaries between the two TAD sets.

While JI is an effective way to compare boundary locations, it does not take into account the total overlap between TAD interiors. We therefore also adopted a measure from Sauerwald and Kingsford [38], which presented a method to identify structurally similar regions between two TAD sets. We updated the original method to address some methodological artifacts and improve efficiency, as detailed in the Supplementary materials. The measure used here, which we will call "TADsim", is simply the fraction of the genome covered by structurally similar regions identified

by the method described in Sauerwald and Kingsford [38].

Statistical comparisons

Distributions of similarity values under all three measures were checked for statistical significance through the Mann-Whitney U test, also called the Mann-Whitney-Wilcoxon (MWW) test. This nonparametric statistical test assesses the null hypothesis that a randomly selected value from one sample is equally likely to be less than or greater than a randomly selected value from the other sample. The alternative hypothesis can then be formulated as a randomly selected value from one distribution being likely to be greater than (or less than) a randomly selected value from the other distribution.

Without knowing the underlying distribution of structural similarity values, a nonparametric statistical test is required for all of our comparisons. The Kolmogorov-Smirnov two-sample test (KS test) is another commonly used nonparametric test, but it does not include any assessment of which distribution is greater than the other. The KS test is additionally sensitive not only to differences in the median or mean between two distributions, but any differences in their shapes, dispersion, or skewness as well. We therefore chose the MWW test for these analyses, given that we specifically are testing for the difference in relative magnitudes of the values in the distributions, rather than differences their overall shapes.

Funding

This research is funded in part by the Gordon and Betty Moore Foundation's Data-Driven Discovery Initiative through Grant GBMF4554 to C.K., and by the US National Institutes of Health (R01HG007104 and R01GM122935). Research reported in this publication was supported by the NIGMS of the NIH under award number P41GM103712 and by the Richard K. Mellon Presidential Fellowship in Life Sciences to N.S. This work was partially funded by The Shurl and Kay Curci Foundation.

Cell type	Description	Replicates	Res frag	Protocol	Accession(s)	Citation
IMR90	lung fibroblast	2	MboI	*in situ*	SRR1658672, SRR1658673, SRR1658674, SRR1658675, SRR1658676, SRR1658677, SRR1658678	[35]
GM12878	blood lymphocyte	2	MboI	*in situ*	SRR1658570, SRR1658571, SRR1658572, SRR1658573, SRR1658574, SRR1658575, SRR1658576, SRR1658577, SRR1658578, SRR1658579, SRR1658580, SRR1658581, SRR1658582, SRR1658583, SRR1658584, SRR1658585, SRR1658586, SRR1658587, SRR1658588, SRR1658589, SRR1658590, SRR1658591, SRR1658592, SRR1658593, SRR1658594, SRR1658595, SRR1658596, SRR1658597, SRR1658598, SRR1658599, SRR1658600, SRR1658601, SRR1658602, SRR1658603	[35]
HMEC	mammary epithelial	2	MboI	*in situ*	SRR1658680, SRR1658681, SRR1658682, SRR1658683, SRR1658684, SRR1658685	[35]
HUVEC	umbilical vein endothelial	1	MboI	*in situ*	SRR1658709, SRR1658710, SRR1658711, SRR1658712, SRR1658713, SRR1658714	[35]
K562	chronic myeloid leukemia	2	MboI	*in situ*	SRR1658693, SRR1658694, SRR1658695, SRR1658696, SRR1658697, SRR1658698, SRR1658699, SRR1658700, SRR1658701, SRR1658702	[35]
KBM7	chronic myeloid leukemia	2	MboI	*in situ*	SRR1658703, SRR1658704, SRR1658705, SRR1658706, SRR1658707, SRR1658708	[35]
NHEK	epidermal keratinocyte	1	MboI	*in situ*	SRR1658689, SRR1658690, SRR1658691	[35]
A549	adenocarcinomic alveolar basal epithelial	2	HindIII	dilution	ENCLB571HTP, ENCLB222WYT	[42]
Caki2	clear cell renal cell carcinoma (epithelial)	2	HindIII	dilution	ENCLB555CZE, ENCLB858SVS	[42]
G401	rhabdoid tumor kidney epithelial	2	HindIII	dilution	ENCLB506SDM, ENCLB589RBY	[42]
LNCaP-FGC	prostate carcinoma epithelial-like	2	HindIII	dilution	ENCLB191OGC, ENCLB473XWD	[42]
NCI-H460	large cell lung cancer	2	HindIII	dilution	ENCLB118KAE, ENCLB104ZTM	[42]
Panc1	pancreas ductal adenocarcinoma	2	HindIII	dilution	ENCLB951HSJ, ENCLB134IVX	[42]
RPMI-7951	malignant melanoma	2	HindIII	dilution	ENCLB210AAY, ENCLB016TGU	[42]
SKMEL5	malignant melanoma	2	HindIII	dilution	ENCLB296ZFT, ENCLB462TWE	[42]
SKNDZ	neuroblastoma	2	HindIII	dilution	ENCLB524GGK, ENCLB952BSP	[42]
SKNMC	neuroepithelioma	2	HindIII	dilution	ENCLB215KZO, ENCLB914GYK	[42]
T47D	ductal carcinoma	2	HindIII	dilution	ENCLB758KFU, ENCLB183QHG	[42]
IMR90	lung fibroblast	2	HindIII	dilution	SRX116345, SRX128222	[11]
hESC	human embryonic stem cell	2	HindIII	dilution	SRX116344, SRX128221	[11]
H1-hESC	human embryonic stem cell	1	NcoI	*in situ*	4DNES4DGHDMX	[10]
H1-hESC	human embryonic stem cell	3	DpnII	*in situ*	4DNESRJ8KV4Q	[10]
H1-hESC	human embryonic stem cell	1	HindIII	dilution	4DNES78Y8Y5K	[10]
H1-hESC	human embryonic stem cell	2	DpnII	*in situ*	4DNES2M5JIGV	[10]
HFF-hTERT	foreskin fibroblast	4	HindIII	dilution	4DNES9L4AK6Q	[10]
HFF-hTERT	foreskin fibroblast	2	DpnII	*in situ*	4DNESVUMGLG2	[10]
HFF-hTERT	foreskin fibroblast	1	NcoI	*in situ*	4DNESY859VLG	[10]
HFF-hTERT	foreskin fibroblast	2	HindIII	*in situ*	4DNESB6MNCFE	[10]
HFF-hTERT	foreskin fibroblast	1	HindIII	*in situ*	4DNES8J78WV2	[10]
HFF-hTERT	foreskin fibroblast	1	MboI	*in situ*	4DNESAPF27TG	[10]
HFFc6	subclone of HFF-hTERT	2	DpnII	*in situ*	4DNES2R6PUEK	[10]
HG00733	blood lymphocyte	2	HindIII	dilution	4DNESTAPSPUC	[10]
HG00732	blood lymphocyte	2	HindIII	dilution	4DNESI2UKI7P	[10]
HG00731	blood lymphocyte	2	HindIII	dilution	4DNESJ1VX52C	[10]
HG00514	blood lymphocyte	2	HindIII	dilution	4DNESE3ICNE1	[10]
HG00513	blood lymphocyte	2	HindIII	dilution	4DNESJIYRA44	[10]
HG00512	blood lymphocyte	2	HindIII	dilution	4DNES4GSP9S4	[10]
GM19238	blood lymphocyte	2	HindIII	dilution	4DNESYUYFD6H	[10]
GM19239	blood lymphocyte	2	HindIII	dilution	4DNESVKLYDOH	[10]
GM19240	blood lymphocyte	2	HindIII	dilution	4DNESHGL976U	[10]
hESC	human embryonic stem cell	1	HindIII	dilution	SRR639047, SRR639048, SRR639049	[21]
IMR90	lung fibroblast	6	HindIII	dilution	SRX212172, SRX212173, SRX294948, SRX294949, SRX294950, SRX294951	[21]
IMR90	lung fibroblast	6	HindIII	dilution	SRX212174, SRX212175, SRX294952, SRX294953, SRX294954, SRX294955	[21]
IMR90	lung fibroblast	1	HindIII	dilution	SRR639045, SRR639046	[21]
hESC	human embryonic stem cell	2	HindIII	dilution	SRX378271, SRX378272	[12]
GM20431	blood lymphocyte	3	HindIII	dilution	ENCLB097VEW, ENCLB167NGL, ENCLB938LSX	[42]
skeletal muscle tissue	gastrocnemius medialis, 4 donors	4	MboI	*in situ*	ENCLB925XYW, ENCLB361HQM, ENCLB966EDS, ENCLB645GUM	[42]
transverse colon	from 4 donors	4	MboI	*in situ*	ENCLB584CUK, ENCLB920LTI, ENCLB724QSQ, ENCLB527HSP	[42]
brain microvascular	endotheilial	2	HindIII	dilution	SRX3322341, SRX3322340	[42]
astrocyte	cerebellum	2	HindIII	dilution	ENCLB672PAB, ENCLB174TEA	[42]
astrocyte	spinal cord	2	HindIII	dilution	SRX3322978, SRX3322979	[42]
DLD1	colon adenocarcinoma epithelial	2	HindIII	dilution	SRX3321987, SRX3321988	[42]
pericyte	brain	2	HindIII	dilution	SRX3322286, SRX3322287	[42]
HEMEC	endometrial microvascular endothelial	2	HindIII	dilution	SRX3322599, SRX3322600	[42]
hepatic sinusoid	endothelial	2	HindIII	dilution	ENCLB284TIY, ENCLB618NVM	[42]
ACHN	renal cell adenocarcinoma epithelial	2	HindIII	dilution	SRX3322373, SRX3322374	[42]
IMR90	lung fibroblast	2	HindIII	dilution	GSM2595584, GSM2595585	[46]
hESC (H9)	human embryonic stem cell	1	HindIII	dilution	GSM2309023	[17]
adrenal gland	tissue	1	HindIII	dilution	SRX2179246	[39]
bladder	tissue	2	HindIII	dilution	SRX2179247, SRX2179248	[39]
DPC	dorsolateral prefrontal cortex tissue	1	HindIII	dilution	SRX2179249	[39]
hippocampus	tissue	1	HindIII	dilution	SRX2179250	[39]
lung	tissue from 2 donors	2	HindIII	dilution	SRX2179252, SRX2179251	[39]
ovary	tissue	1	HindIII	dilution	SRX2179253	[39]
pancreas	tissue from 2 donors	2	HindIII	dilution	SRX2179254, SRX2179255, SRX2179256, SRX2179257	[39]
psoas muscle	tissue from 2 donors	2	HindIII	dilution	SRX2179260, SRX2179258, SRX2179259	[39]
right ventricle	tissue	1	HindIII	dilution	SRX2179261	[39]
small bowel	tissue	1	HindIII	dilution	SRX2179262	[39]
spleen	tissue from 2 donors	2	HindIII	dilution	SRX2179264, SRX2179263	[39]

Table 1: All Hi-C data used in this study

Author contributions

N.S. collected, processed, and analyzed all of the data. C.K. and N.S. designed the experiments and wrote the manuscript. A.S. conceived of and implemented all improvements to the original TADsim method. All authors contributed to the final manuscript.

Financial disclosure

C.K. is a co-founder of Ocean Genomics, Inc.

References

[1] Ay, F. and Noble, W. S. (2015). Analysis methods for studying the 3D architecture of the genome. *Genome Biology*, **16**(1), 183.

[2] Ay, F., Bailey, T. L., and Noble, W. S. (2014). Statistical confidence estimation for Hi-C data reveals regulatory chromatin contacts. *Genome Research*, **24**(6), 999–1011.

[3] Barrett, T., Wilhite, S. E., Ledoux, P., Evangelista, C., Kim, I. F., Tomashevsky, M., Marshall, K. A., Phillippy, K. H., Sherman, P. M., Holko, M., *et al.* (2012). NCBI GEO: archive for functional genomics data sets – update. *Nucleic Acids Research*, **41**(D1), D991–D995.

[4] Bonev, B. and Cavalli, G. (2016). Organization and function of the 3D genome. *Nature Reviews Genetics*, **17**(11), 661–678.

[5] Cavalli, G. and Misteli, T. (2013). Functional implications of genome topology. *Nature Structural and Molecular Biology*, **20**(3), 290–299.

[6] Chen, Y., Zhang, Y., Wang, Y., Zhang, L., Brinkman, E. K., Adam, S. A., Goldman, R., van Steensel, B., Ma, J., and Belmont, A. S. (2018). Mapping 3D genome organization relative to nuclear compartments using TSA-Seq as a cytological ruler. *The Journal of Cell Biology*, **217**(11), 4025–4048.

[7] Crane, E., Bian, Q., McCord, R. P., Lajoie, B. R., Wheeler, B. S., Ralston, E. J., Uzawa, S.,

Dekker, J., and Meyer, B. J. (2015). Condensin-driven remodelling of X chromosome topology during dosage compensation. *Nature*, **523**(7559), 240–244.

[8] Cremer, T. and Cremer, C. (2001). Chromosome territories, nuclear architecture and gene regulation in mammalian cells. *Nature Reviews Genetics*, **2**(4), 292–301.

[9] Dekker, J., Rippe, K., Dekker, M., and Kleckner, N. (2002). Capturing chromosome conformation. *Science*, **295**(5558), 1306–1311.

[10] Dekker, J., Belmont, A. S., Guttman, M., Leshyk, V. O., Lis, J. T., Lomvardas, S., Mirny, L. A., O'Shea, C. C., Park, P. J., Ren, B., *et al.* (2017). The 4D nucleome project. *Nature*, **549**(7671), 219–226.

[11] Dixon, J. R., Selvaraj, S., Yue, F., Kim, A., Li, Y., Shen, Y., Hu, M., Liu, J. S., and Ren, B. (2012). Topological domains in mammalian genomes identified by analysis of chromatin interactions. *Nature*, **485**(7398), 376–380.

[12] Dixon, J. R., Jung, I., Selvaraj, S., Shen, Y., Antosiewicz-Bourget, J. E., Lee, A. Y., Ye, Z., Kim, A., Rajagopal, N., Xie, W., *et al.* (2015). Chromatin architecture reorganization during stem cell differentiation. *Nature*, **518**(7539), 331–336.

[13] Dixon, J. R., Gorkin, D. U., and Ren, B. (2016). Chromatin domains: the unit of chromosome organization. *Molecular Cell*, **62**(5), 668–680.

[14] Duggal, G., Wang, H., and Kingsford, C. (2014). Higher-order chromatin domains link eQTLs with the expression of far-away genes. *Nucleic Acids Research*, **42**(1), 87–96.

[15] Fillipova, D., Patro, R., Duggal, G., and Kingsford, C. (2014). Identification of alternative topological domains in chromatin. *Algorithms for Molecular Biology*, **9**, 14.

[16] Forcato, M., Nicoletti, C., Pal, K., Livi, C. M., Ferrari, F., and Bicciato, S. (2017). Comparison of computational methods for Hi-C data analysis. *Nature Methods*, **14**(7), 679–685.

[17] Freire-Pritchett, P., Schoenfelder, S., Várnai, C., Wingett, S. W., Cairns, J., Collier, A. J., García-Vílchez, R., Furlan-Magaril, M., Osborne, C. S., Fraser, P., *et al.* (2017). Global reorgan-

isation of cis-regulatory units upon lineage commitment of human embryonic stem cells. *eLife*, **6**, e21926.

[18] Fudenberg, G., Getz, G., Meyerson, M., and Mirny, L. (2011). High order chromatin architecture shapes the landscape of chromosomal alterations in cancer. *Nature Biotechnology*, **29**(12), 1109–1113.

[19] Hnisz, D., Weintraub, A. S., Day, D. S., Valton, A.-L., Bak, R. O., Li, C. H., Goldmann, J., Lajoie, B. R., Fan, Z. P., Sigova, A. A., *et al.* (2016). Activation of proto-oncogenes by disruption of chromosome neighborhoods. *Science*, **351**(6280), 1454–1458.

[20] Imakaev, M., Fudenberg, G., McCord, R. P., Naumova, N., Goloborodko, A., Lajoie, B. R., Dekker, J., and Mirny, L. A. (2012). Iterative correction of Hi-C data reveals hallmarks of chromosome organization. *Nature Methods*, **9**, 999–1003.

[21] Jin, F., Li, Y., Dixon, J. R., Selvaraj, S., Ye, Z., Lee, A. Y., Yen, C.-A., Schmitt, A. D., Espinoza, C. A., and Ren, B. (2013). A high-resolution map of the three-dimensional chromatin interactome in human cells. *Nature*, **503**(7475), 290–294.

[22] Le Dily, F., Baù, D., Pohl, A., Vicent, G. P., Serra, F., Soronellas, D., Castellano, G., Wright, R. H., Ballare, C., Filion, G., *et al.* (2014). Distinct structural transitions of chromatin topological domains correlate with coordinated hormone-induced gene regulation. *Genes & Development*, **28**(19), 2151–2162.

[23] Leinonen, R., Sugawara, H., Shumway, M., and Collaboration, I. N. S. D. (2010). The sequence read archive. *Nucleic Acids Research*, **39**(suppl_1), D19–D21.

[24] Lieberman-Aiden, E., van Berkum, N. L., Williams, L., Imakaev, M., Ragoczy, T., Telling, A., Amit, I., Lajoie, B. R., Sabo, P. J., Dorschner, M. O., Sandstrom, R., Bernstein, B., Bender, M. A., Groudine, M., Gnirke, A., Stamatoyannopoulos, J., Mirny, L. A., Lander, E. S., and Dekker, J. (2009). Comprehensive mapping of long-range interactions reveals folding principles of the human genome. *Science*, **326**(5950), 289–293.

[25] Lupiáñez, D. G., Spielmann, M., and Mundlos, S. (2016). Breaking TADs: how alterations of chromatin domains result in disease. *Trends in Genetics*, **32**(4), 225–237.

[26] Meaburn, K. J., Gudla, P. R., Khan, S., Lockett, S. J., and Misteli, T. (2009). Disease-specific gene repositioning in breast cancer. *The Journal of Cell Biology*, **187**(6), 801–812.

[27] Meilă, M. (2003). Comparing clusterings by the variation of information. In *Learning Theory and Kernel Machines*, pages 173–187. Springer.

[28] Misteli, T. (2010). Higher-order genome organization in human disease. *Cold Spring Harbor Perspectives in Biology*, **2**(8), a000794.

[29] Moindrot, B., Audit, B., Klous, P., Baker, A., Thermes, C., de Laat, W., Bouvet, P., Mongelard, F., and Arneodo, A. (2012). 3D chromatin conformation correlates with replication timing and is conserved in resting cells. *Nucleic Acids Research*, **40**(19), 9470–9481.

[30] Nagano, T., Várnai, C., Schoenfelder, S., Javierre, B.-M., Wingett, S. W., and Fraser, P. (2015). Comparison of Hi-C results using in-solution versus in-nucleus ligation. *Genome Biology*, **16**(1), 175.

[31] Norton, H. K., Emerson, D. J., Huang, H., Kim, J., Titus, K. R., Gu, S., Bassett, D. S., and Phillips-Cremins, J. E. (2018). Detecting hierarchical genome folding with network modularity. *Nature Methods*, **15**(2), 119–122.

[32] Ou, H. D., Phan, S., Deerinck, T. J., Thor, A., Ellisman, M. H., and O'Shea, C. C. (2017). ChromEMT: Visualizing 3D chromatin structure and compaction in interphase and mitotic cells. *Science*, **357**(6349), eaag0025.

[33] Pope, B. D., Ryba, T., Dileep, V., Yue, F., Wu, W., Denas, O., Vera, D. L., Wang, Y., Hansen, R. S., Canfield, T. K., *et al.* (2014). Topologically associating domains are stable units of replication-timing regulation. *Nature*, **515**(7527), 402–405.

[34] Ramani, V., Cusanovich, D. A., Hause, R. J., Ma, W., Qiu, R., Deng, X., Blau, C. A., Disteche, C. M., Noble, W. S., Shendure, J., *et al.* (2016). Mapping 3D genome architecture through in situ DNase Hi-C. *Nature Protocols*, **11**(11), 2104–2121.

[35] Rao, S. S. P., Huntley, M. H., Durand, N. C., Stamenova, E. K., Bochkov, I. D., Robinson, J. T., Sanborn, A. L., Machol, I., Omer, A. D., Lander, E. S., and Aiden, E. L. (2014). A 3D

map of the human genome at kilobase resolution reveals principles of chromatin looping. *Cell*, **159**(7), 1665–1880.

[36] Rennie, S., Dalby, M., van Duin, L., and Andersson, R. (2018). Transcriptional decomposition reveals active chromatin architectures and cell specific regulatory interactions. *Nature Communications*, **9**(1), 487.

[37] Ryba, T., Hiratani, I., Lu, J., Itoh, M., Kulik, M., Zhang, J., Schulz, T. C., Robins, A. J., Dalton, S., and Gilbert, D. M. (2010). Evolutionarily conserved replication timing profiles predict long-range chromatin interactions and distinguish closely related cell types. *Genome Research*, **20**(6), 761–770.

[38] Sauerwald, N. and Kingsford, C. (2018). Quantifying the similarity of topological domains across normal and cancer human cell types. *Bioinformatics*, **34**(13), i475–i483.

[39] Schmitt, A. D., Hu, M., Jung, I., Xu, Z., Qiu, Y., Tan, C. L., Li, Y., Lin, S., Lin, Y., Barr, C. L., *et al.* (2016). A compendium of chromatin contact maps reveals spatially active regions in the human genome. *Cell Reports*, **17**(8), 2042–2059.

[40] Servant, N., Varoquaux, N., Lajoie, B. R., Viara, E., Chen, C.-J., Vert, J.-P., Heard, E., Dekker, J., and Barillot, E. (2015). HiC-Pro: an optimized and flexible pipeline for Hi-C data processing. *Genome Biology*, **16**(1), 259.

[41] Spielmann, M., Lupiáñez, D. G., and Mundlos, S. (2018). Structural variation in the 3D genome. *Nature Reviews Genetics*, **19**, 453–467.

[42] The ENCODE Project Consortium (2012). An integrated encyclopedia of DNA elements in the human genome. *Nature*, **489**(7414), 57–74.

[43] Weinreb, C. and Raphael, B. J. (2015). Identification of hierarchical chromatin domains. *Bioinformatics*, **32**(11), 1601–1609.

[44] Yaffe, E. and Tanay, A. (2011). Probabilistic modeling of Hi-C contact maps eliminates systematic biases to characterize global chromosomal architecture. *Nature Genetics*, **43**(11), 1059–1065.

[45] Yang, T., Zhang, F., Yardımcı, G. G., Song, F., Hardison, R. C., Noble, W. S., Yue, F., and Li, Q. (2017). HiCRep: assessing the reproducibility of Hi-C data using a stratum-adjusted correlation coefficient. *Genome Research*, **27**(11), 1939–1949.

[46] Zirkel, A., Nikolic, M., Sofiadis, K., Mallm, J.-P., Brackley, C. A., Gothe, H., Drechsel, O., Becker, C., Altmüller, J., Josipovic, N., *et al.* (2018). HMGB2 loss upon senescence entry disrupts genomic organization and induces CTCF clustering across cell types. *Molecular Cell*, **70**(4), 730–744.