Multiple species drive flexible lake food webs with warming

Timothy J. Bartley[1,2]* (timothy.bartley@gmail.com), Tyler D. Tunney[3,4] (tyler.tunney@dfo-mpo.gc.ca), Nigel P. Lester[5,6] (nigel.lester@ontario.ca), Brian J. Shuter[5,6] (brian.shuter@utoronto.ca), Robert H. Hanner[2,7] (rhanner@uoguelph.ca), and Kevin S. McCann[2] (ksmccann@uoguelph.ca)

[1]Department of Biology, University of Toronto Mississauga, 3359 Mississauga Road, Mississauga, Ontario, Canada, L5L 1C6

[2]Department of Integrative Biology, University of Guelph, 50 Stone Road East, Guelph, Ontario, Canada N1G 2W1

[3]Gulf Fisheries Centre, Fisheries and Oceans Canada, Moncton, New Brunswick E1C 9B6, Canada

[4]Center for Limnology, University of Wisconsin–Madison, Madison, WI, USA 53706

[5]Science and Research Branch, Ontario Ministry of Natural Resources, Peterborough, Ontario, Canada K9J 7B8

[6]Department of Ecology and Evolution, University of Toronto, Toronto, Ontario, Canada M5S 3G5

[7]Biodiversity Institute of Ontario, University of Guelph, 50 Stone Road East, Guelph, Ontario, Canada N1G 2W1

22   *correspondence to Timothy J. Bartley, Department of Biology, University of Toronto

23   Mississauga, 3359 Mississauga Road, Mississauga, Ontario, Canada, L5L 1C6 e-mail:

24   timothy.bartley@gmail.com; phone: 1 (519) 546-4652

26   Statement of Authorship

31   Data Accessibility Statement

32   Upon acceptance, we will archive our data on Zenodo, https://zenodo.org/

34   Running Title

35   Lake food webs flex with warming

37   Keywords

Abstract

1. Climate change is rewiring the food webs that determine the fate of diverse ecosystems. Mobile generalist consumers are responding to climate change by rapidly shifting their behaviour and foraging, driving food webs to flex. Although these responsive generalists form a key stabilizing module in food web structure, the extent to which they are present throughout whole food webs is largely unknown.

2. Our goal was to examine if the thermal preferences of multiple species drive predictable behavioural and foraging responses to changes in temperature that drive flexes in food web structure.

3. We used stable isotope and catch-per-unit-effort data to investigate the habitat use and diet of four widespread and abundant species of fishes that represent different key trophic roles in food webs of boreal lakes that span a natural 7°C air temperature gradient.

4. We found significant reductions in nearshore derived carbon and nearshore habitat use with increased temperature in three of four fish species, indicating that multiple species drive flexible lake food webs with warming. We also found evidence that the response of lake trout to increased temperatures may reduce their biomass and cascade to release their preferred prey, the pelagic forage fish cisco.

5. Our results suggest that climate warming will shift lake food webs toward increased reliance on offshore habitats and resources. We argue that species across trophic levels broadly couple lake macrohabitats, suggesting that potentially stabilizing responsive consumers are present throughout food webs. However, climate change appears to limit their ability to responsively forage, critically undermining a repeated stabilizing mechanism in food webs.

Introduction

Ongoing rapid climate change is altering species interactions and reorganizing the food webs that

determine the fate of diverse ecosystems (Barton, Beckerman, & Schmitz, 2009; Kortsch,

Primicerio, Fossheim, Dolgov, & Aschan, 2015; T. D. Tunney, McCann, Lester, & Shuter,

2014). As environmental conditions change, food web structure changes—or 'flexes'—with it

(McMeans et al., 2016). Identifying these changes in food web structure is central to

understanding the consequences of climate change on the major energy pathways of carbon and

nutrient flow in ecosystems but is challenging given that food web structure is inherently

complex (Polis, 1991). Importantly, climate change is expected to 'rewire' food webs (Bartley et

al., n.d.) by adding or removing whole trophic links (e.g., Kortsch *et al.* 2015) and by modifying

the strengths of existing interactions (e.g., Barton & Schmitz 2009). These changes in the flow of

energy among species are likely to have myriad consequences in ecosystems, from species to

whole ecosystem function (Guzzo, Blanchfield, & Rennie, 2017; Kondoh, 2010; Křivan &

Schmitz, 2003). If so, future warming may alter food webs in ways that lead to ecosystems that

differ substantially from past and present conditions (Kortsch et al., 2012). Despite the potential

for these drastic changes, our understanding of food webs responses to climate change and the

consequences of these response remain limited.

Food webs are flexible because of the foraging responses of mobile generalist consumers that

can rapidly shift their foraging behaviour in response to environmental change. Many generalist

consumers move across the landscape to forage in various spatially-distinct macrohabitats

(Rooney, McCann, & Moore, 2008) and can feed omnivorously across trophic levels (Thompson

et al. 2007). As a result, these generalist consumers can respond in two key ways: by changing

the degree to which they couple different macrohabitats (i.e., 'horizontal' shifts between energy

pathways, Schindler & Scheuerell 2002; Vander Zanden & Vadeboncoeur 2002; Rooney *et al.*

2008) and by shifting their degree of omnivory (i.e., 'vertical' foraging shifts along an energy

pathway, Barton & Schmitz 2009; Sentis *et al.* 2014; Ruiz-Cooley *et al.* 2017). Generalists that

exhibit both horizontal and vertical foraging flexibility form the generalist module (*sensu*

McMeans *et al.* 2016), a widely documented patterning of food web interactions (Eloranta et al.,

2015; Nakano & Murakami, 2001; Polis, Anderson, Anderson, & Holt, 1997; Rooney et al.,

2008) that can expand and contract food webs in response to changing environmental conditions

(T. D. Tunney, McCann, Lester, & Shuter, 2012). This generalist module may be a powerful

stabilizing structure, allowing mobile generalists to respond to changing densities in different

habitats in a manner that buffers population dynamics (Kevin Shear McCann & Rooney, 2009;

McMeans et al., 2016). Despite this potential, few studies have empirically documented the

presence of fundamental module throughout food webs or how human impacts like climate

change may interrupt this module.

One well-documented example of a responsive generalist consumer that drives food web

flexibility is lake trout (*Salvelinus namaycush*), a common generalist top predator in lakes of

boreal North America. Lake trout forage on fishes and invertebrates in both of the two thermally

distinct lake macrohabitats: the nearshore (i.e., littoral) habitat and the offshore (i.e., pelagic)

habitat (Vander Zanden & Vadeboncoeur, 2002). Because lake trout is cold-adapted and

thermally sensitive, the foraging habits of lake trout are mediated by thermal accessibility of the

relatively warm nearshore habitat (Guzzo et al., 2017; Plumb & Blanchfield, 2009). Lake trout

show reduced nearshore habitat use with increasing nearshore temperature (Guzzo et al., 2017;

T. D. Tunney et al., 2014). When lake trout avoid entering the physiologically taxing nearshore

habitat, they forage less on nearshore resources and more on fish (T. D. Tunney et al., 2012,

2014). This reduced access to abundant nearshore prey may undermine lake trout's ability to

responsively forage to resource densities in different macrohabitats (Guzzo et al., 2017; T. D.

Tunney et al., 2014). Importantly, this phenomenon may not be limited to lake trout; many fishes

are generally highly mobile and flexible foragers (Dill, 1983), and boreal lakes in North America

contain numerous cold-adapted fish species that may exhibit similar behavioural and foraging

responses due to thermal limitations (Hasnain, Shuter, & Minns, 2013; Vander Zanden &

Vadeboncoeur, 2002). In contrast, the species that prefer warmer waters may not be thermally

limited and thus may not exhibit behavioural or foraging responses to warming. However, the

roles that species other than lake trout play in determining how lake food webs flex are largely

unexplored. To understand how whole food webs are being altered by climate change requires

that we examine how many species across trophic levels and with various thermal preferences

respond to warming.

Here, we use a natural climate gradient of approximately 7°C to examine how multiple

species behavioural and foraging responses drive flexes in food web structure. We investigate the

habitat use and diet of four widespread and abundant species of boreal shield fishes that

represent different key roles and trophic levels in the food web: the cold-adapted generalist top

predator lake trout, the cold-adapted pelagic planktivorous cisco (*Coregonus artedi*), the cool-

adapted piscivorous top predator walleye (*Sander vitreus*), and the cool-adapted mesopredator

yellow perch (*Perca flavescens*) (Coker, Portt, & Minns, 2001; Hasnain et al., 2013). We use

stable-isotope-based food-web indices of nearshore feeding and trophic position along with

catch-per-unit-effort based metrics of behaviour to show that several species across trophic levels

and from multiple thermal guilds respond predictably to changes in temperature. We also use

catch-per-unit-effort data to show that lake trout show reduced biomass index under increased

temperatures, consistent with a reduction of the nearshore resource availability. Our results

suggestion that multiple species comprising key trophic roles drive flexible lake food webs with

warming. We argue that by studying the foraging and behavioral responses of this set of key

players, we can predict how food webs will flex with climate change. We end by discussing how

these changes suggest climate change will impact the function and stability of lake ecosystems,

and how many generalist consumers may be a fundamental feature of food web architecture.

Materials and Methods

Lake Selection

The Canadian boreal shield includes hundreds of thousands of lakes that span various

natural environmental gradients, including climate (Gunn, Steedman, & Ryder, 2004; Keller,

2007). We used data for 66 lakes in the province of Ontario, all of which have been used

previously to study food web structure through stable isotope analysis (Dolson, McCann,

Rooney, & Ridgway, 2009; T. D. Tunney et al., 2014; T. Tunney, McCann, Jarvis, Lester, &

Shuter, 2018). We use lakes that have both nearshore and offshore intermediate consumers to

ensure that changes in food web structure or predator behaviour are not driven by differences in

the presence or absence of whole trophic groups. For 59 of our 66 lakes, we used catch-per-unit-

effort (CUE) data from the Ontario Ministry of Natural Resources and Forestry (OMNRF)

Broad-scale Fisheries Monitoring (BSM) Program (Sandstrom, Rawson, & Lester, 2013). In

summary, for each lake, standardized fish community surveys took place one time between May

and September from 2008 to 2012 using overnight sets of two types of multipaneled mesh

gillnets: North American standard multipaneled gill nets (with 8 mesh sizes varying from 38 to

127 mm) and small mesh gillnets (with 5 mesh sizes from 13 to 38 mm). Netting occurred

randomly throughout the lake and depending on the maximum depth of each lake, in each of up

to eight depth strata: 1-3m, 3-6m, 6-12m, 12-20m, 20-35m, 35-50m, 50-75m, and >75m.

Complete details can be found in Sandstrom *et al.* 2013. Catch data were not available for the

seven lakes from Dolson *et al.* (2009).

Lake Attributes

We used two measures of temperature for our analysis: average recent air temperature in

°C and growing degree days > 5 °C (Table S1). Our primary temperature variable was average

recent air temperature, which was calculated as the average hourly air temperature in °C for the

30 days prior to field sampling because it corresponds to the time period reflected by the isotopic

signature of fish muscle tissue (Peterson & Fry, 1987). This temperature variable was derived

from FetchClimate (Grechka et al., 2016) using the latitude and longitude of each lake. In

addition to the average recent air temperature, we corroborated our results with and growing

degree days above 5°C from 1981 to 2010 (provided by the Ontario Ministry of Natural

Resources and Forestry), which measures the accumulation of heat over time. We also account

for several physical and chemical factors that have been previously identified as important

drivers of habitat coupling or trophic position in boreal shield lakes (Table S1, all provided by

the Ontario Ministry of Natural Resources, Dolson *et al.* 2009; Tunney *et al.* 2014, 2018): lake

surface area in hectares, mean lake depth in metres, Secchi depth in metres, and total phosphorus

in $\mu g \cdot L^{-1}$, and lake shape (calculated the shoreline development index): SDI = (Shoreline

Distance) $/2 \sqrt{\pi \times Surface\ Area}$ (equation 1).

Species Selection

We used four species that are typical of their respective trophic groups. Stable isotope

data for these species was available for 40 or more lakes and the trophic level, thermal

classifications, and habitat preferences for these species are known (Coker et al., 2001; Hasnain

et al., 2013). Lake trout (*Salvelinus namaycush*) is the most common top predator of offshore

habitats in boreal shield lakes, and much research on boreal shield lake food webs has focused on

this coldwater species (Dolson et al., 2009; Guzzo et al., 2017; Plumb & Blanchfield, 2009; T. D.

Tunney et al., 2014). Cisco (or lake herring, *Coregonus artedi*) is one of the most common cold-

adapted planktivores and is a common prey item for lake trout and walleye. Walleye (*Sander*

vitreus) is a common cool-adapted piscivore and popular sport fish that is present in lakes across

the boreal shield. Yellow perch (*Perca flavescens*) is a widespread and abundant cool-adapted

mesopredator that is consumed by a wide variety of predatory fishes throughout its range,

including both lake trout and walleye. These four species comprise a large portion of the average

catch in the 59 lakes that we use here from the OMNRF's BsM surveys (see Supplementary

Information). The number of individual for each species sampled for stable isotope analysis in

each lake varied from two to 21 (mean 13.5) for lake trout, three to 32 (mean 17.0) for walleye,

one to 20 (mean 10.8) for cisco, and one to 18 (mean 9.1) for yellow perch. For both lake trout

and walleye, only individuals greater than 250 mm were used for stable isotope analysis because

these species are known to show ontogenic diet shifts (Galarowicz, Adams, & Wahl, 2006;

Mittelbach & Persson, 1998; Sherwood, Pazzia, Moeser, Hontela, & Rasmussen, 2002).

Food Web Metric Calculations using Stable Isotopes

As in many previous studies, we used stable isotopic signatures from our four fish species

and baseline invertebrates to calculate both the nearshore carbon index (based on the proportion

nearshore carbon, e.g., Tunney *et al.* 2014) and trophic position (David M Post, 2002; Vander

Zanden, Casselman, & Rasmussen, 1999; Vander Zanden, Shuter, Lester, & Rasmussen, 1999).

Collection and processing methods for stable isotope data can be found in Dolson *et al.* 2009;

Tunney *et al.* 2014, 2018). We used baseline invertebrates from the nearshore and offshore zones

to account for variability in isotopic signatures across lakes. We incorporated data for multiple

trophic groups into both our nearshore and offshore baseline isotopic signatures to reduce the

number of estimated baseline values required for our analysis and to increase the sample size for

our baseline isotopic signatures. We corrected all $\delta^{13}C$ signatures using C:N ratios as $\delta^{13}C_{corr} =$

$\delta^{13}C_{raw} + (-3.32 + 0.99 * CN)$ (equation 2), where $\delta^{13}C_{corr}$ is the corrected $\delta^{13}C$ signature,

$\delta^{13}C_{raw}$ is the raw $\delta^{13}C$ signature, and CN is the C:N ratio of that tissue sample (David M. Post

et al., 2007). For lakes that were missing either nearshore or offshore baseline isotopic

signatures, we estimated the $\delta^{13}C$ and $\delta^{15}N$ signatures of the missing baseline using the available

baseline and simple linear regression between the baselines across lakes (see Supplementary

Information).

We used two source mixing models to estimate the nearshore carbon index and the

trophic position of each species in each lake based on their relative isotopic signatures (David M

Post, 2002). We calculated the nearshore carbon index for each fish species as $NCI_{fish} =$

$(\delta^{13}C_{fish} - \delta^{13}C_{osb})/(\delta^{13}C_{nsb} - \delta^{13}C_{osb})$ (equation 3), where NCI_{fish} is the nearshore carbon

index in the diet of a fish species, $\delta^{13}C_{fish}$ is the average $\delta^{13}C$ signature for that fish species,

$\delta^{13}C_{osb}$ is the average or estimated $\delta^{13}C$ signature for all offshore baselines (i.e., mussels and/or

zooplankton), and $\delta^{13}C_{nsb}$ is the average or estimated $\delta^{13}C$ signature for all nearshore baselines

(i.e., snails and/or aquatic insect larvae). The nearshore carbon index is similar to the proportion

nearshore carbon used by others (e.g., Tunney *et al.* 2014) but is not constrained between zero

and one. This approach minimizes data transformation requirements of the stable isotopic

signatures and increases the number of lakes in our analyses while still allowing us to

qualitatively understand changes in the relative contribution of nearshore resources to fish diets.

Based on the nearshore feeding index, we estimated the trophic position of each species

as $\text{TP}_{\text{fish}} = 2 + \left(\delta^{15}\text{N}_{\text{fish}} - \left(\delta^{15}\text{N}_{\text{nsb}} \times \text{NCI}_{\text{fish}} + \delta^{15}\text{N}_{\text{osb}} \times (1 - \text{NCI}_{\text{fish}})\right)\right)/3.4$ (equation 4),

where TP_{fish} is the trophic position of a fish species, NFI_{fish} is the nearshore carbon index for that

fish species, $\delta^{15}\text{N}_{\text{fish}}$ is the average $\delta^{15}\text{N}$ signature for that fish species, $\delta^{15}\text{N}_{\text{nsb}}$ is the average or

estimated $\delta^{15}\text{N}$ signature for all nearshore baselines (i.e., snails and aquatic insect larvae),

$\delta^{15}\text{N}_{\text{osb}}$ is the average or estimated $\delta^{15}\text{N}$ signature for all offshore baselines (i.e., mussels and

zooplankton), 3.4 is the assumed increase in $\delta^{15}\text{N}$ due to fractionation (David M Post, 2002), for

each trophic level, and 2 is the assumed trophic position of the baseline invertebrates.

Behaviour Metrics and Biomass Index Using Catch-per-unit-effort

We used catch-per-unit-effort data for each depth stratum (Sandstrom *et al.* 2013) to

calculate a weighted average mean depth of capture for each species as $\text{SMD} = (\sum_{i=1}^{4} d_i \times$

$CUE_i \times p_i)/(\sum_{i=1}^{4} CUE_i)$ (equation 5), where SMD is the mean depth of capture of a fish species,

CUE_i is the CUE of that species for depth stratum i, p_i is the proportion of the lake in depth

stratum i, and d_i is the depth of at the center of the depth range for stratum i (2m for stratum 1,

4.5m for stratum 2, 9m for stratum 3, and 16m for stratum 4). We only used depth strata 1

through 4 to calculate mean depth of capture because these are the only depth strata that are

sampled with both small-mesh and large-mesh gear (Sandstrom et al., 2013). The CUE data that

we use here is the number of fish per 100 metre nights. For CUE biomass index, we used the kg

of fish per gang per 100 metre nights, calculated by summing the area-weighted CUE (number of

fish per 100 metre nights) for each depth stratum and then multiplying by the mean weight of

that fish species.

Statistical Analyses

To test for the effect of temperature on each response variable (nearshore carbon index,

trophic position, mean depth of capture, nearshore/offshore presence, and the log-transformed

CUE biomass index of each species), we used a two-step approach. Firstly, to account for

morphometric or chemical factors known or suspected to influence the foraging and behaviour of

these fish species, we first constructed multiple regression models for each response variable and

lake size in hectares, lake shape (SDI), Secchi depth in m, mean lake depth in m, and total

phosphorous in $\mu g \bullet L^{-1}$. We then compared all subsets of each full multivariate regression model

using AIC. We selected the best model as the model with the smallest number of explanatory

variables from the set of models within 2 AIC points of the lowest AIC value (Aho *et al.* 2014,

see Supplementary Information for best models). Secondly, to test the effect of each temperature

variable, we took the best model for each response variable and ran a multiple regression that

included the temperature variable (Figs 2, 3, and 4, Table S3). To meet statistical requirements,

we used $\log_{10}$ transformed lake surface area and natural log transformed mean lake depth and

total phosphorous. We removed all data points with a Cook's distance greater than 1 from their

respective regression models, and we used variance inflation factor to check for inflated

coefficient estimates in multiple linear regressions. All regression analyses were performed in

the R statistical language (v3.2.3), and model selection was conducted using the R package

"glmulti" (Calcagno, 2013).

Results

Food Web Metric Calculations using Stable Isotopes

We found strong evidence that lake food web structure varied across an approximately 7° C

climate gradient. Three of the four species showed evidence of changes in nearshore feeding

with increased temperature. Both top predators (lake trout and walleye) showed a significant

decrease in the nearshore carbon index with increasing average recent air temperature, and the

cold-adapted cisco showed a similar marginally significant decrease (Fig. 2a, c, and e, Table S3).

The cool-adapted yellow perch showed no significant relationship between average recent air

temperature and the nearshore carbon index (Fig. 2g, Table S3). Lake trout showed a significant

increase in trophic position with increased average recent air temperature (Fig. 2b, Table S3), but

cisco, walleye and yellow perch showed no significant relationship between trophic position with

increasing average recent air temperature (Fig. 2d, Table S3).

Behaviour Metrics and Biomass Index Using Catch-per-unit-effort

In agreement with changes in nearshore carbon index, the same three species (lake trout,

walleye, and cisco) showed strong evidence of reduced nearshore habitat use. All three species

showed a significant increase in mean depth of capture with increasing average recent air

temperature (Fig. 3a, c, and e, Table S3). Consistent with these results, lake trout and cisco

showed a significant decrease in probability of nearshore presence and walleye showed a

marginally significant increase in probability of offshore presence with increasing average recent

air temperature (see Supplementary Information). In contrast, yellow perch showed no

relationship between either mean depth of capture or probability of offshore presence and

average recent air temperature (Fig. 3g and inset, Table S3).

We found evidence of a reduction in biomass with increasing temperature for one

species, lake trout, which showed a significant decrease in CUE biomass index with increasing

average recent air temperature (Fig. 3b, Table S3). In contrast, cisco, walleye, and yellow perch

showed no significant relationship between average recent air temperature CUE biomass index

(Fig. 3d, f and h, Table S3).

Discussion

Understanding how climate change is altering whole food webs is a challenging but critical

endeavor for understanding the responses of whole ecosystems. Previous research showed that

lake trout (*Salvelinus namaycush*), a generalist top predator in lake ecosystems, responds to

increased temperature by reducing its nearshore habitat and resource use, and its degree of

omnivory (Guzzo et al., 2017; T. D. Tunney et al., 2014), thus forming a flexible 'generalist

module' (McMeans et al., 2016). Here, we extend that research to examine the impacts of

climate warming on four key species that represent different key roles and trophic levels in

boreal lake food webs: lake trout, walleye (*Sander vitreus*), cisco (*Coregonus artedi*), and yellow

perch (*Perca flavescens*). Our stable-isotope-based food-web metrics and our catch-per-unit-

effort-based behavioural metrics show a consistent response of species across multiple trophic

levels to warming to nearshore-derived carbon and offshore habitat use. In fact, temperature was

the most significant variable for predicting food web structure in this study (see Supplementary

Information). By harnessing the variation in food web structure across natural temperature

gradients, we have detected consistent food web responses to warming in boreal shield lake

ecosystems.

We found strong evidence of foraging and behavioural responses to increased temperature in

three of the four key food web members that we examined. As expected, our cold-adapted

species, the predatory lake trout and the planktivorous cisco, showed decreased nearshore

feeding and decreased near-shore habitat use with increasing temperatures. Our results are

consistent with the thermally accessibility hypothesis that has been previously shown for lake

trout (Guzzo et al., 2017; T. D. Tunney et al., 2014). Because offshore fish communities are

largely comprised of species that prefer cold water (Coker et al., 2001; Hasnain et al., 2013),

these thermal constraints likely operate on whole offshore fish communities, causing them to

respond uniformly to warming. We found that the cool-adapted piscivorous walleye showed

behavioural and foraging responses through decreased coupling and decreased nearshore habitat

use with higher temperature, but in contrast the cool-adapted mesopredator yellow perch showed

no such response. Since walleye and yellow perch have similar thermal tolerances (Hasnain *et al*.

2013), the reason for the different responses of these species is unclear. Despite the lack of

response in yellow perch, the foraging responses of these fish species strongly indicate that

boreal shield lake food webs will collectively flex towards offshore resource and habitat use in

response to warming.

We found evidence that a reduction in nearshore coupling with warming may impact the

biomass of lake trout. The biomass of lake trout decreased significantly with increased

temperature, but none of the other species showed this pattern (Fig. 3). Such reductions in

biomass with increasing temperature may be a consequence of a reduction in overall energy

availability due to decreased nearshore coupling. Mobile top predators that garner energy from

multiple spatially distinct macrohabitats are expected to have elevated biomass as a result of a

higher overall resource availability, producing relatively top-heavy food webs (i.e., non-Eltonian,

Mccauley *et al.* 2018; Woodson *et al.* 2018). Therefore, reduced access to the nearshore energy

pathway in warm lakes could therefore reduce overall energy availability for lake trout and so

lower lake trout biomass and population density (Guzzo et al., 2017; T. Tunney et al., 2018). A

reduction in lake trout biomass across could subsequently reduce the top-down control of lake

trout on cisco, their preferred prey. Reduced top-down control by lake trout in warmer lakes in

conjunction with a bottom-up reduction in their biomass from reduced access to nearshore

resources may result in no discernable trend in standing biomass of cisco with temperature,

consistent with our results here. Changes in trophic biomass structure and concomitant changes

in top-down control are one possible consequence of the flexes in lake food webs towards the

offshore with warming. If climate change broadly decouples predators from multiple resource

pools, changes in biomass structure may be a widespread signatures of climate change, possibly

altering top-down effects within ecosystems (K. S. McCann, Rasmussen, & Umbanhowar, 2005;

McCauley et al., 2018).

Our results demonstrate that many mobile generalist consumers at trophic levels and

throughout compartments drive flexibility throughout food webs in boreal shield lakes. Our work

builds on previous studies that have started to expose the role that single generalist top predators

species play in determining food web responses to climate change in many ecosystems (Barton

& Schmitz, 2009; Eloranta et al., 2016; Woodward, Perkins, & Brown, 2010). The presence of

flexible foragers at various trophic levels is consistent with both recent ecological theory that

predicts large, mobile generalists ought to rapidly respond to environmental change (Rooney &

McCann, 2012), and is consistent with empirical evidence of flexible foraging from various

ecosystems (Edmunds, Laberge, & McCann, 2016; Gonçalves, Azeiteiro, Pardal, & De Troch,

2012; Held, Mendl, Devereux, & Byrne, 2002; Rooney et al., 2008; Schmitz, Beckerman, &

O'Brien, 1997; Shochat, Lerman, Katti, & Lewis, 2004; Taipale, Kankaala, Tiirola, & Jones,

2008; Vander Zanden & Vadeboncoeur, 2002). The presence of responsive foraging throughout

diverse food webs rather than just at the top of food chains would make the generalist module the

type of remarkably scale-invariant feature of food web architecture that has long interested

ecologists (Sugihara, Schoenly, & Trombla, 1989).

The presence of responsive consumers across trophic levels and throughout food web

compartments has strong implications for food web stability. Responsive foraging by generalists

can be potently stabilizing in food webs (McMeans et al., 2016; D M Post, Conners, & Goldberg,

2000; Rip, McCann, Lynn, & Fawcett, 2010; Rooney, McCann, Gellner, & Moore, 2006).

Generalist consumers can average variability in their resources (akin to the portfolio effect of

Tilman *et al.* (1998)) by rapidly responding to resource variability in space and time by shifting

away from low-density resources and towards high density resources, impeding that variability

from emanating throughout food webs (Valdovinos, Ramos-Jiliberto, Garay-Narváez, Urbani, &

Dunne, 2010). This switching ability also increases trophic redundancy, which can prevent

secondary extinctions (Borrvall, Ebenman, & Jonsson, 2000; Sanders, Thébault, Kehoe, & Frank

van Veen, 2018). If a broad suite of organisms can make rapid, smart foraging responses to

environmental change, they may accentuate the stabilizing effect of a single generalist module.

The presence of many responsive consumers increases stability in complex food web models

(Kondoh, 2010; Valdovinos et al., 2010). This reveals a potential powerful, repeated stabilizing

mechanism throughout food webs that Levin (1998) argues is fundamental to persistence of

ecosystems. However, when generalist consumers decouple macrohabitats to avoid entering

physiologically taxing habitats, their ability to respond to consumer densities is limited,

undermining their stabilizing potential (Guzzo et al., 2017; Schmitz & Barton, 2014; T. D.

Tunney et al., 2014). Thus, the decoupling of habitats with warming may be widespread across

ecosystems, forcing the food webs of many ecosystems to flex with climate change in ways that

alter ecosystem stability worldwide.

Our work here adds a key piece to the mounting evidence that generalist consumers likely to

play a vital role in determining how climate change rewires food webs (Bartley et al., n.d.).

Generalist consumers positioned throughout food webs may ultimately reorganize whole

ecosystems in response to climate change (Blanchard, 2015; McMeans et al., 2016). Determining

both the prevalence and positions of these responsive generalist consumers and the factors that

constrain these species responses are critical tasks for predicting the myriad ways that food webs

will adapt with a changing climate. Importantly, our results suggest that these species throughout

food webs contribute to adaptive capacity—the ability of an ecosystem to change in response to

changing conditions—a key part of the resilience of ecosystems in the face of environmental

variation (McMeans et al., 2016). Although we have showed that lake food webs flex at many

axes, it is unclear when this adaptive capacity will be exceeded and entire ecosystems collapse.

This uncertainty highlights the clear need to forecast major changes in food web structure.

Monitoring the behaviour and foraging of responsive consumers is a promising way to track

changes in food web structure and give prescient warnings of future drastic changes in ecosystem

dynamics (Bartley et al., n.d.; Velarde, Ezcurra, & Anderson, 2013). This makes responsive

generalist consumers critical to understanding of how ecosystems will fare through climate

change.

Acknowledgements

This work was supported by the National Science and Engineering Research Council of Canada through a Discovery Grant to K.S.M., a Strategic Grant to K.S.M., B.J.S. and N.P.L., and a Canada Graduate Scholarship to T.J.B. We thank K. Armstrong, J. Wright and many other biologists and technical staff at the Ontario Ministry of Natural Resources and Forestry and the Broad-scale Fisheries Monitoring Program. We also thank M. Berger, I. Farahbakhsh, B. Graham, B. Heavenrich for assistance with gathering and organizing climate data. D. Schindler, M. Granados, M. Guzzo, and B. Robinson provided helpful comments on an early draft of this manuscript.

Author Contributions

T.J.B. and K.S.M. conceived and designed this study. T.J.B. and T.D.T. collected field data and conducted lab work. T.J.B. analyzed data and wrote the initial draft of the manuscript. T.J.B., T.D.T., N.P.L., B.J.S., R.H.H., and K.S.M. all contributed substantially to writing and revisions.

Data Accessibility

Upon acceptance, our data will be archived on Zenodo, https://zenodo.org/

References

Aho, K., Derryberry, D., & Peterson, T. (2014). Model selection for ecologists: The worldviews of AIC and BIC. *Ecology*. doi:10.1890/13-1452.1

Bartley, T. J., McCann, K. S., Bieg, C., Cazelles, K., Granados, M., Guzzo, M. M., … McMeans, B. C. (n.d.). Food Web Rewiring in a Changing World. *Nature Ecology & Evolution*.

Barton, B. T., Beckerman, A. P., & Schmitz, O. J. (2009). Climate warming strengthens indirect interactions in an old-field food web. *Ecology*, *90*(9), 2346–2351.

Barton, B. T., & Schmitz, O. J. (2009). Experimental warming transforms multiple predator effects in a grassland food web. *Ecology Letters*, *12*(12), 1317–1325. doi:10.1111/j.1461-0248.2009.01386.x

Blanchard, J. L. A. (2015). A rewired food web. *Nature*, *527*(7577), 7–8. doi:10.1038/nature16311

Borrvall, C., Ebenman, B., & Jonsson, T. (2000). Biodiversity lessens the risk of cascading extinctions in model food webs. *Ecology Letters*, *3*, 131–136. doi:10.1046/j.1461-0248.2000.00130.x

Calcagno, V. (2013). glmulti: Model selection and multimodel inference made easy. Retrieved from https://cran.r-project.org/web/packages/glmulti/

Coker, G., Portt, C., & Minns, C. K. (2001). *Morphological and ecological characteristics of Canadian freshwater fishes*. Burlington, Ontario.

Dill, L. M. (1983). Adaptive Flexibility in the Foraging Behavior of Fishes. *Canadian Journal of Fisheries and Aquatic Sciences*, *40*(4), 398–408. doi:10.1139/f83-058

Dolson, R., McCann, K., Rooney, N., & Ridgway, M. (2009). Lake morphometry predicts the degree of habitat coupling by a mobile predator. *Oikos*, *118*(8), 1230–1238. doi:10.1111/j.1600-0706.2009.17351.x

Edmunds, N. B., Laberge, F., & McCann, K. S. (2016). A role for brain size and cognition in food webs. *Ecology Letters*, *19*(8), 948–955.

Eloranta, A. P., Helland, I. P., Sandlund, O. T., Hesthagen, T., Ugedal, O., & Finstad, A. G. (2016). Community structure influences species' abundance along environmental gradients. *Journal of Animal Ecology*, *85*(1), 273–282. doi:10.1111/1365-2656.12461

Eloranta, A. P., Kahilainen, K. K., Amundsen, P. A., Knudsen, R., Harrod, C., & Jones, R. I. (2015). Lake size and fish diversity determine resource use and trophic position of a top predator in high-latitude lakes. *Ecology and Evolution*, *5*(8), 1664–1675. doi:10.1002/ece3.1464

Galarowicz, T. L., Adams, J. A., & Wahl, D. H. (2006). The influence of prey availability on ontogenetic diet shifts of a juvenile piscivore. *Canadian Journal of Fisheries and Aquatic Sciences*, *63*(8), 1722–1733. doi:10.1139/f06-073

Gonçalves, A. M. M., Azeiteiro, U. M., Pardal, M. A., & De Troch, M. (2012). Fatty acid
profiling reveals seasonal and spatial shifts in zooplankton diet in a temperate estuary.
Estuarine, Coastal and Shelf Science, *109*, 70–80. doi:10.1016/j.ecss.2012.05.020

Grechka, D. A., Berezin, S. B., Emmott, S., Lyutsarev, V., Smith, M. J., & Purves, D. W. (2016).
Universal, easy access to geotemporal information: FetchClimate. *Ecography*. Ecography.
doi:10.1111/ecog.02321

Gunn, J. M., Steedman, R. J., & Ryder, R. A. (2004). *Boreal shield watersheds: lake trout
ecosystems in a changing environment*. CRC Press.

Guzzo, M. M., Blanchfield, P. J., & Rennie, M. D. (2017). Behavioral responses to annual
temperature variation alter the dominant energy pathway, growth, and condition of a cold-
water predator. *Proceedings of the National Academy of Sciences*, 201702584.
doi:10.1073/pnas.1702584114

Hasnain, S. S., Shuter, B. J., & Minns, C. K. (2013). Phylogeny influences the relationships
linking key ecological thermal metrics for North American freshwater fish species.
Canadian Journal of …, *972*(April), 964–972. doi:70(7): 964-972, 10.1139/cjfas-2012-0217

Held, S., Mendl, M., Devereux, C., & Byrne, R. W. (2002). Foraging pigs alter their behaviour in
response to exploitation. *Animal Behaviour*, *64*, 157–165. doi:10.1006/anbe.2002.3044

Keller, W. (2007). Implications of climate warming for Boreal Shield lakes: a review and
synthesis. *Environmental Reviews*, *15*(NA), 99–112. doi:10.1139/A07-002

Kondoh, M. (2010). Foraging Adaptation and the Relationship between Food-Web Complexity
and Stability. *Science*, *299*(5611), 1388–1391. doi:10.1126/science.1079154

Kortsch, S., Primicerio, R., Beuchel, F., Renaud, P. E., Rodrigues, J., Lønne, O. J., & Gulliksen,
B. (2012). Climate-driven regime shifts in Arctic marine benthos. *Proceedings of the
National Academy of Sciences of the United States of America*, *109*(35), 14052–7.
doi:10.1073/pnas.1207509109

Kortsch, S., Primicerio, R., Fossheim, M., Dolgov, A. V., & Aschan, M. (2015). Climate change
alters the structure of arctic marine food webs due to poleward shifts of boreal generalists.
Proceedings of the Royal Society B: Biological Sciences, *282*(1814), 20151546.
doi:10.1098/rspb.2015.1546

Křivan, V., & Schmitz, O. J. (2003). Adaptive foraging and flexible food web topology.
Evolutionary Ecology Research, *5*(5), 623–652.

Levin, S. A. (1998). Ecosystems and the biosphere as complex adaptive systems. *Ecosystems*,
1(5), 431–436. doi:10.1007/s100219900037

McCann, K. S., Rasmussen, J. B., & Umbanhowar, J. (2005). The dynamics of spatially coupled
food webs. *Ecology Letters*, *8*(5), 513–523. doi:10.1111/j.1461-0248.2005.00742.x

McCann, K. S., & Rooney, N. (2009). The more food webs change, the more they stay the same.
Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences,

364(1524), 1789–1801. doi:10.1098/rstb.2008.0273

McCauley, D. J., Gellner, G., Martinez, N. D., Williams, R. J., Sandin, S. A., Micheli, F., …
McCann, K. S. (2018). On the prevalence and dynamics of inverted trophic pyramids and
otherwise top-heavy communities. *Ecology Letters*, *21*(3), 439–454. doi:10.1111/ele.12900

McMeans, B. C., McCann, K. S., Tunney, T. D., Fisk, A. T., Muir, A. M., Lester, N. P., …
Rooney, N. (2016). The adaptive capacity of lake food webs: from individuals to
ecosystems. *Ecological Monographs*, *86*(1), 4–19.

Mittelbach, G. G., & Persson, L. (1998). The ontogeny of piscivory and its ecological
consequences. *Canadian Journal of Fisheries and Aquatic Sciences*, *55*(6), 1454–1465.
doi:10.1139/f98-041

Nakano, S., & Murakami, M. (2001). Reciprocal subsidies: Dynamic interdependence between
terrestrial and aquatic food webs. *Proceedings of the National Academy of Sciences of the*
United States of America, *98*(1), 166–170. doi:10.1073/pnas.98.1.166

Peterson, B. J., & Fry, B. (1987). Stable isotopes in ecosystem studies. *Annual Reviews in*
Ecology and Systems, *18*, 293–320.

Plumb, J. M., & Blanchfield, P. J. (2009). Performance of temperature and dissolved oxygen
criteria to predict habitat use by lake trout (Salvelinus namaycush). *Canadian Journal of*
Fisheries and Aquatic Sciences, *66*(11), 2011–2023. doi:10.1139/F09-129

Polis, G. A. (1991). Complex trophic interactions in deserts□: an empirical critique of food-web
theory. *The American Naturalist*. doi:10.1086/285208

Polis, G. A., Anderson, W. B., Anderson, R., & Holt, R. D. (1997). Toward an Integration of
Landscape and Food Web Ecology□: The Dynamics of Spatially Subsidized Food Webs.
Annual Review of Ecology and Systematics, *28*(1997), 289–316.
doi:10.1146/annurev.ecolsys.28.1.289

Post, D. M. (2002). Using stable isotopes to estimate trophic position: models, methos, and
assumptions. . *Ecology*, *83*(3), 703–718. doi:Doi 10.2307/3071875

Post, D. M., Conners, M. E., & Goldberg, D. S. (2000). Prey preference by a top preator and the
stability of linked food chains. *Ecology*, *81*(1), 8–14. doi:10.1890/0012-
9658(2000)081[0008:PPBATP]2.0.CO;2

Post, D. M., Layman, C. A., Arrington, D. A., Takimoto, G., Quattrochi, J., & Montaña, C. G.
(2007). Getting to the fat of the matter: Models, methods and assumptions for dealing with
lipids in stable isotope analyses. *Oecologia*, *152*(1), 179–189. doi:10.1007/s00442-006-
0630-x

Rip, J. M. K., McCann, K. S., Lynn, D. H., & Fawcett, S. (2010). An experimental test of a
fundamental food web motif. *Proceedings. Biological Sciences / The Royal Society*,
277(February), 1743–1749. doi:10.1098/rspb.2009.2191

Rooney, N., McCann, K., Gellner, G., & Moore, J. C. (2006). Structural asymmetry and the

stability of diverse food webs. *Nature*, *442*(July), 265–269. doi:10.1038/nature04887

Rooney, N., & McCann, K. S. (2012). Integrating food web diversity, structure and stability. *Trends in Ecology and Evolution*, *27*(1), 40–45. doi:10.1016/j.tree.2011.09.001

Rooney, N., McCann, K. S., & Moore, J. C. (2008). A landscape theory for food web architecture. *Ecology Letters*, *11*(8), 867–881. doi:10.1111/j.1461-0248.2008.01193.x

Ruiz-Cooley, R. I., Gerrodette, T., Fiedler, P. C., Chivers, S. J., Danil, K., & Ballance, L. T. (2017). Temporal variation in pelagic food chain length in response to environmental change. *Science Advances*, *3*(10), e1701140. doi:10.1126/sciadv.1701140

Sanders, D., Thébault, E., Kehoe, R., & Frank van Veen, F. J. (2018). Trophic redundancy reduces vulnerability to extinction cascades. *Proceedings of the National Academy of Sciences*, *0*, 201716825. doi:10.1073/pnas.1716825115

Sandstrom, S., Rawson, M., & Lester, N. (2013). *Manual of instructions for broad-scale fish community monitoring using North American (NA1) and Ontario small mesh (ON2) gillnets*.

Schindler, D. E., & Scheuerell, M. D. (2002). Habitat coupling in lake ecosystems. *Oikos*, *98*, 177–189. doi:10.1034/j.1600-0706.2002.980201.x

Schmitz, O. J., & Barton, B. T. (2014). Climate change effects on behavioral and physiological ecology of predator-prey interactions: Implications for conservation biological control. *Biological Control*, *75*, 87–96. doi:10.1016/j.biocontrol.2013.10.001

Schmitz, O. J., Beckerman, A. P., & O'Brien, K. M. (1997). Behaviorally mediated trophic cascades: Effects of predation risk on food web interactions. *Ecology*, *78*(5), 1388–1399. doi:10.1890/0012-9658(1997)078[1388:BMTCEO]2.0.CO;2

Sentis, A., Hemptinne, J. L., & Brodeur, J. (2014). Towards a mechanistic understanding of temperature and enrichment effects on species interaction strength, omnivory and food-web structure. *Ecology Letters*, *17*(7), 785–793. doi:10.1111/ele.12281

Sherwood, G. D., Pazzia, I., Moeser, A., Hontela, A., & Rasmussen, J. B. (2002). Shifting gears: enzymatic evidence for the energetic advantage of switching diet in wild-living fish. *Canadian Journal of Fisheries and Aquatic Sciences*, *59*(2), 229–241. doi:10.1139/f02-001

Shochat, E., Lerman, S. B., Katti, M., & Lewis, D. B. (2004). Linking optimal foraging behavior to bird community structure in an urban☐desert landscape: field experiments with artificial food patches. *The American Naturalist*, *164*(2), 232–243. doi:10.1086/422222

Sugihara, G., Schoenly, K., & Trombla, A. (1989). Scale invariance in food web properties. *Science*, *245*(4913), 48–52. doi:10.1126/science.2740915

Taipale, S., Kankaala, P., Tiirola, M., & Jones, R. I. (2008). Whole-Lake Dissolved Inorganic 13C Additions Reveal Seasonal Shifts in Zooplankton Diet. *Ecology*, *89*(2), 463–474. doi:10.1890/07-0702.1

Thompson, R. M., Hemberg, M., Starzomski, B. M., Shurin, J. B., Hompson, R. M. T., Emberg, M. H., & Tarzomski, B. R. M. S. (2007). Trophic Levels and Trophic Tangles□: the Prevalence of Omnivory in Real Food Webs. *Ecology*, *88*(3), 612–617.

Tilman, D., Lehman, C. L., Bristow, C. E., Tilman, D., Lehman, C. L., & Bristow, C. E. (1998). Notes and Comments Diversity-Stability Relationships□: Statistical Inevitability or Ecological Consequence□? *American Naturalist*, *151*(3), 277–282.

Tunney, T. D., McCann, K. S., Lester, N. P., & Shuter, B. J. (2012). Food web expansion and contraction in response to changing environmental conditions. *Nature Communications*, *3*(May), 1105. doi:10.1038/ncomms2098

Tunney, T. D., McCann, K. S., Lester, N. P., & Shuter, B. J. (2014). Effects of differential habitat warming on complex communities. *Proceedings of the National Academy of Sciences of the United States of America*, *111*(22), 8077–82. doi:10.1073/pnas.1319618111

Tunney, T., McCann, K., Jarvis, L., Lester, N., & Shuter, B. (2018). Blinded by the light? Nearshore energy pathway coupling and relative predator biomass increased with reduced water transparency across lakes. *Oecologia*. doi:10.1007/s00442-017-4049-3

Valdovinos, F. S., Ramos-Jiliberto, R., Garay-Narváez, L., Urbani, P., & Dunne, J. A. (2010). Consequences of adaptive behaviour for the structure and dynamics of food webs. *Ecology Letters*, *13*(12), 1546–1559. doi:10.1111/j.1461-0248.2010.01535.x

Vander Zanden, M. J., Casselman, J. M., & Rasmussen, J. B. (1999). Stable isotope evidence for the food web consequences of species invasions in lakes. *Nature*, *401*(6752), 464–467. doi:10.1038/46762

Vander Zanden, M. J., Shuter, B. J., Lester, N., & Rasmussen, J. B. (1999). Patterns of food chain length in lakes: A stable isotope study. *American Naturalist*, *154*(154), 406–416. doi:10.1086/303250

Vander Zanden, M. J., & Vadeboncoeur, Y. (2002). Fishes As Integrators of Benthic and Pelagic Food Webs in Lakes. *Ecology*, *83*(8), 2152–2161. doi:10.1890/0012-9658(2002)083[2152:FAIOBA]2.0.CO;2

Velarde, E., Ezcurra, E., & Anderson, D. W. (2013). Seabird diets provide early warning of sardine fishery declines in the Gulf of California. *Scientific Reports*, *3*(FEBRUARY), 1332. doi:10.1038/srep01332

Woodson, C. B., Schramski, J. R., & Joye, S. B. (2018). A unifying theory for top-heavy ecosystem structure in the ocean. *Nature Communications*, *9*(1), 23. doi:10.1038/s41467-017-02450-y

Woodward, G., Perkins, D. M., & Brown, L. E. (2010). Climate change and freshwater ecosystems: impacts across multiple levels of organization. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, *365*(1549), 2093–2106. doi:10.1098/rstb.2010.0055

Figure with eight panels (a–h) showing nearshore foraging index and mean log(depth) of capture versus average recent air temperature for lake trout, cisco, walleye, and yellow perch.

Figure 1. Nearshore foraging index and the mean log(depth) of capture of lake trout (a and b), cisco (c and d), walleye (e and f), and yellow perch (g and h) in lakes across a gradient of average recent air temperature. A solid line indicates a p-value less than 0.05, and a dashed line indicates a p-value less than 0.1. Regression model details can be found in Table S3. Fish silhouettes were taken from PhyloPic (http://www.phylopic.org).

618
619
620 **Figure 2.** Trophic position of lake trout (a), walleye (b), cisco (c), and yellow perch (d) in lakes
621 across a gradient of average recent air temperature. A solid line indicates a p-value less than
622 0.05, and a dashed line indicates a p-value less than 0.1. Regression model details can be found
623 in Table S3. Fish silhouettes were taken from PhyloPic (http://www.phylopic.org).
624
625

627
628
629 **Figure 3.** The log-transformed CUE biomass index for lake trout (a), walleye (b), cisco (c), and
630 yellow perch (d) in lakes across a gradient of average recent air temperature. A solid line
631 indicates a p-value less than 0.05, and a dashed line indicates a p-value less than 0.1. Regression
632 model details can be found in Table S3. Fish silhouettes were taken from PhyloPic
633 (http://www.phylopic.org).
634

a) typical cold lake

i) thermal conditions

ii) species' responses

b) warmed lake after climate change

iii) thermal conditions

iv) species' responses

Figure 4: Schematic of the effects of warming on four widespread and abundant fish species of
boreal shield lake food web: lake trout, cisco, walleye, and yellow perch. The x axis indicates
shifts in the degree of nearshore foraging based on $\delta^{13}C$ signatures. The y axis indicates shifts in
habitat use in terms of the mean depth of capture. Lake trout, cisco, walleye all show shifts
towards offshore foraging and offshore habitat use with increased temperature. Fish silhouettes
were taken from PhyloPic (http://www.phylopic.org). Adapted from Tunney *et al.* 2014.