SGK phosphorylates Cdc25 and Myt1 to trigger cyclin B-Cdk1 activation at the

meiotic G2/M transition

Daisaku Hiraoka[1*], Enako Hosoda[2], Kazuyoshi Chiba[2], and Takeo Kishimoto[1*]

[1]Science & Education Center, Ochanomizu University, Tokyo 112-8610, Japan.

[2]Department of Biological Sciences, Ochanomizu University, Tokyo 112-8610, Japan.

[*]Correspondence to:

Daisaku Hiraoka: E-mail: hiraoka.daisaku@ocha.ac.jp, Tel: +81-3-5978-5370

Takeo Kishimoto: E-mail: kishimoto.takeo@ocha.ac.jp, Tel: +81-3-5978-5370

Abstract

The kinase cyclin B-Cdk1 complex is a master regulator of M-phase in both mitosis and

meiosis. At the G2/M transition, cyclin B-Cdk1 activation is initiated by a trigger that

reverses the balance of activities between Cdc25 and Wee1/Myt1, and is further

accelerated by autoregulatory loops. In somatic cell mitosis, this trigger was recently

proposed to be the cyclin A-Cdk1/Plk1 axis. However, in the oocyte meiotic G2/M

transition, in which hormonal stimuli induce cyclin B-Cdk1 activation, cyclin A-Cdk1

is non-essential and hence the trigger remains elusive. Here, we show that SGK directly

phosphorylates Cdc25 and Myt1 to trigger cyclin B-Cdk1 activation in starfish oocytes.

After hormonal stimulation of the meiotic G2/M transition, SGK is activated by

cooperation between the Gβγ-PI3K pathway and an unidentified pathway downstream

of Gβγ, called the atypical Gβγ pathway. These findings identify the trigger in oocyte

meiosis and provide insights into the role and activation of SGK.

Introduction

Activation of cyclin-dependent kinase 1 complexed with cyclin B (cyclin B-Cdk1)

induces entry into M-phase during somatic cell mitosis and germ cell meiosis[1-3]. Cyclin

B-Cdk1 is regulated by synthesis and degradation of cyclin B, and by inhibitory

phosphorylation of Cdk1 at Thr14 and Tyr15. These residues are phosphorylated by

kinases belonging to the Wee1/membrane-associated tyrosine/threonine 1 (Myt1)

family, but dephosphorylated by cell division cycle 25 (Cdc25)[4]. Cyclin B accumulates

before and/or during M-phase entry. However, Wee1/Myt1 activity is dominant over

Cdc25 activity before M-phase; therefore, cyclin B-Cdk1 remains inactive due to

inhibitory phosphorylation. At M-phase entry, a small population of cyclin B-Cdk1 is

first activated by a trigger that reverses the balance between Cdc25 and Wee1/Myt1

activities, thereby making Cdc25 activity predominant. Thereafter, cyclin B-Cdk1 itself

further accelerates dephosphorylation of Cdk1 via feedback loops, leading to maximal

activation[4-8]. Although the molecular identity of the trigger of cyclin B-Cdk1 activation

has received great attention, it remains elusive.

In mitotic M-phase entry (G2/M transition), the trigger may be affected by various

factors, such as checkpoints[5,9,10], and may involve redundant or stochastic processes[5,9].

At least in normal mitotic cell cycles, polo-like kinase 1 (Plk1) is activated by Aurora A

in a cyclin A-Cdk1-dependent manner, and in turn phosphorylates Cdc25 to trigger

activation of cyclin B-Cdk1[11-13]. Consistently, the cyclin A-Cdk1/Plk1 axis functions as

the trigger for cyclin B-Cdk1 activation in the first embryonic cell division cycle[14].

In contrast with mitotic cell cycles in proliferating somatic cells, meiotic cell

cycles in animal oocytes mostly arrest at prophase of meiosis I (prophase-I), which

corresponds to late mitotic G2 phase[15]. Release from this arrest is also induced by cyclin

B-Cdk1, which is activated downstream of extracellular hormonal stimuli[16]; hence this

is known as the meiotic G2/M transition. However, this transition has not been reported

to require cyclin A[16-21]. Furthermore, Plk1[14,22,23] and Aurora[24-26] are non-essential in

some oocytes. Thus, mechanisms other than the cyclin A-Cdk1/Plk1 axis likely trigger

cyclin B-Cdk1 activation at the meiotic G2/M transition in oocytes.

In vertebrate oocytes, meiotic G2 arrest requires cAMP-dependent protein kinase

A (PKA), and downregulation of this kinase leads to meiotic G2/M transition[16,20,27,28].

In mice, PKA appears to directly upregulate Wee1B and downregulate Cdc25; however,

it remains unclear how downregulation of PKA reverses the balance between Wee1B

and Cdc25 activities to trigger cyclin B-Cdk1 activation[29]. In *Xenopus*, the hormone

progesterone stimulates protein synthesis of Moloney sarcoma oncogene (Mos) and

cyclin B, which redundantly trigger cyclin B-Cdk1 activation[30]. This redundant process

is possibly mediated in two manners: activating phosphorylation of Cdc25 by

Mos/mitogen activated protein kinase (MAPK) cascade, and inhibitory phosphorylation

of Myt1 by newly synthesized cyclin B-associated Cdk1[23]. However, the molecular link

between downregulation of PKA and *de novo* protein synthesis of Mos and cyclin B

remains elusive[31,32].

In contrast with vertebrate oocytes, the mechanisms by which activation of cyclin

B-Cdk1 is triggered at the meiotic G2/M transition have been well studied in

invertebrate starfish oocytes since the existence of a trigger kinase was first

reported[16,21,33]. In starfish, cAMP and PKA are likely not involved in meiotic G2

arrest[34]. The physiological maturation-inducing hormone 1-methyladenine (1-MeAde)

induces the meiotic G2/M transition[35] without a requirement for new protein synthesis[36].

Cyclin A, Wee1, and Mos are not present, and Aurora and Plk are not required for

cyclin B-Cdk1 activation, in the meiotic G2/M transition[14,19,25,37]. In unstimulated

oocytes, cyclin B is already accumulated[38], but Myt1 inactivates cyclin B-Cdk1[39].

Stimulation by 1-MeAde is mediated by an unidentified cell surface G-protein-coupled

receptor (GPCR), which induces dissociation of Gβγ from Gαi on the plasma

membrane[40-43]. Gβγ binds to and activates phosphoinositide-3-kinase (PI3K), resulting

in production of phosphatidylinositol 3,4,5-triphosphate (PI-345P_3)[44,45]. Dependent on

PI-345P_3, phosphoinositide-dependent kinase 1 (PDK1) and target of rapamycin

complex 2 (TORC2) phosphorylate Akt in its activation loop (A-loop) and hydrophobic

motif (HM), respectively, leading to its activation[46,47]. Thereafter, Akt inactivates Myt1

by directly phosphorylating Ser75 (ref. 39) and activates Cdc25 by directly

phosphorylating multiple residues including Ser188 (ref. 48), resulting in reversal of the

balance of Cdc25 and Myt1 activities, and consequently initial activation of cyclin

B-Cdk1 (refs 33,39). These observations suggest that Akt likely represents the trigger

kinase at the meiotic G2/M transition in starfish oocytes[39].

Nevertheless, our recent observations suggested that the PI3K-Akt pathway is

insufficient[48]. Expression of a constitutively active form of Akt (CA-Akt) induces the

meiotic G2/M transition, but the required expression level of CA-Akt is 40-fold higher

than that of endogenous Akt. Furthermore, expression of a constitutively active form of

PI3K (CA-PI3K) induces Akt activation, but results in low levels of Cdc25 and Myt1

phosphorylation and therefore fails to induce the meiotic G2/M transition. We reported

that another pathway, called "the atypical pathway", which is activated by Gβγ in

parallel with PI3K activation, compensates for this insufficiency[48]. Although the

molecular details of this pathway remain unclear, we proposed that it cooperates with

the PI3K-Akt pathway to trigger cyclin B-Cdk1 activation[48]. One possibility is that

another kinase, which can redundantly phosphorylate Akt substrates, functions in these

pathways to trigger cyclin B-Cdk1 activation.

One such candidate kinase is serum- and glucocorticoid-regulated kinase

(SGK)[49,50]. SGK is a member of the AGC kinase family, which includes Akt[51], and has

the same substrate consensus motif (RXRXXS/T, where X represents any amino acid)

as Akt[52]. In mammalian cells, similar to Akt, SGK family members are activated in a

PI3K-dependent manner, dependent on phosphorylation of the A-loop and HM by

PDK1 and TORC2, respectively[52,53]. However, the contribution of SGK to the meiotic

G2/M transition has not been investigated.

Here, we carefully evaluated whether Akt and SGK play redundant roles in the

meiotic G2/M transition in starfish oocytes. We showed that SGK is necessary and

sufficient for the initial phosphorylation of Cdc25 and Myt1 to trigger cyclin B-Cdk1

activation upon 1-MeAde stimulation. By contrast, although contradictory to our

previous report[39], inhibition of Akt did not affect the meiotic G2/M transition. Several

experiments supported the importance of SGK, rather than of Akt. This discrepancy is

discussed. Furthermore, as upstream pathways, we found that PI3K and the atypical

Gβγ pathway cooperatively activate SGK. All these observations clearly explain the

insufficiency of the PI3K-Akt pathway. Thus, our findings indicate that SGK is the

trigger kinase at the meiotic G2/M transition in starfish oocytes, and provide new

insights into SGK activation.

Results

A newly generated phospho-specific antibody detects A-loop phosphorylation of endogenous starfish Akt (sfAkt)

In previous studies[47,48], we monitored phosphorylation of the HM by immunoblotting as a marker of starfish Akt (sfAkt) activity because attempts to generate an antibody that detects the phosphorylated A-loop were unsuccessful. Nevertheless, it is important to analyze A-loop phosphorylation to precisely evaluate Akt activation. Thus, we again attempted to generate a phospho-specific antibody against a 17-amino acid phospho-peptide derived from the A-loop of sfAkt that includes phospho-Thr at the site of phosphorylation by PDK1 (Fig. 1a). In immunoblot analysis of whole-oocyte samples using this antibody, phosphorylation of sfAkt was detectable following 1-MeAde stimulation only when Akt was overexpressed (Fig. 1b, closed arrowhead). When endogenous Akt was concentrated by immunoprecipitation using an antibody raised against a C-terminal fragment of sfAkt (anti-sfAkt-C antibody), the anti-phospho-A-loop antibody detected a band at the same size as Akt after 1-MeAde stimulation (Fig. 1c for input and flow-through, d for precipitate, closed arrowhead). The band disappeared upon treatment with the PDK1 inhibitor BX795 (Fig. 1d, closed arrowhead), indicating that the anti-phospho-A-loop antibody successfully detected A-loop phosphorylation of endogenous Akt by PDK1.

SGK is activated after 1-MeAde stimulation in a PDK1- and TORC2-dependent manner

The phospho-A-loop antibody detected another band that migrated slightly slower (~60 kDa) than Akt (Fig. 1b, c, open arrowhead). This band was still observed in

flow-through samples in which Akt had been removed by immunoprecipitation (Fig. 1c,

open arrowhead) and disappeared in the presence of BX795 (Fig. 1c, open arrowhead).

Thus, we speculated that the band corresponds to PDK1-dependent A-loop

phosphorylation of a kinase other than Akt.

In parallel with our research, Hosoda et al. found that starfish SGK (sfSGK) is

activated after 1-MeAde stimulation in starfish oocytes in a study of intracellular pH

regulation as shown in the accompanying paper[54]. They isolated a cDNA encoding

sfSGK and collaborated with us to generate an anti-sfSGK neutralizing antibody raised

against a C-terminal peptide of sfSGK including the HM (anti-sfSGK-C antibody). A

predicted molecular weight of sfSGK is 56 kDa, and eleven of the seventeen residues in

the antigen peptide used to raise the anti-phospho-A-loop antibody against sfAkt are

conserved in the A-loop of SGK (Fig. 1a). Thus, the other band detected by the

anti-phospho-A-loop antibody may correspond to A-loop phosphorylation of SGK. The

mobility of this band was the same as that of the top band of SGK detected by the

anti-sfSGK-C antibody in 1-MeAde-treated oocytes (Fig. 1e, open arrowhead). In

addition, the band disappeared after depletion of SGK from oocyte extracts (Fig. 1f),

whereas it was detected in a sample of immunoprecipitated SGK (Fig. 1f), indicating

that the anti-phospho-A-loop antibody also detects A-loop phosphorylation of sfSGK.

Taken together, these results confirmed that SGK is activated after 1-MeAde

stimulation.

SGK and Akt phosphorylate a Ser or Thr residue in the same consensus motif[52]

and may therefore play a redundant role in 1-MeAde signaling. We next analyzed the

dynamics of SGK activation in starfish oocytes in detail. As previously reported[48],

phosphorylation of Akt in the HM and phosphorylation of Cdc25 at Ser188 were

detected within 2 min after 1-MeAde addition (Fig. 1g). SGK was activated at a similar

time (Fig. 1g, pA-loop). Thereafter, Tyr15 of Cdk1 was dephosphorylated, which is

indicative of cyclin B-Cdk1 activation (Fig. 1g, 12 min and later). In mammalian cells,

SGK is activated by phosphorylation of its A-loop and HM, which is catalyzed by

PDK1 and mTORC2, respectively[52,53]. In addition, A-loop phosphorylation depends on

HM phosphorylation[52]. In starfish oocytes, a mobility shift of SGK was still observed

even when A-loop phosphorylation was abolished by BX795 (Fig. 1c). Using chemical

inhibitors, Hosoda et al. suggested that this mobility shift represents TORC2-dependent

phosphorylation[54]. Consistently, inhibition of TOR by microinjection of an anti-TOR

neutralizing antibody[47] completely abolished the mobility shift of SGK after 1-MeAde

stimulation (Fig. 1h). In these oocytes, A-loop phosphorylation of SGK was also

abolished (Fig. 1h). Therefore, A-loop phosphorylation of SGK likely depends on

TORC2-dependent HM phosphorylation in starfish oocytes, as reported in mammalian

cells.

SGK, but not Akt, is required for the meiotic G2/M transition

Next, we examined the involvement of Akt and SGK in the meiotic G2/M transition. To

this end, we specifically inhibited each kinase using antibodies. Non-specific bands

were detected by immunoblotting of starfish oocytes with the anti-sfAkt-C antibody

(Fig. 2a), but were not immunoprecipitated by this antibody (Fig. 2a). This indicates

that the anti-sfAkt-C antibody does not interact with these non-specific proteins in

non-denaturing conditions. In addition, although immunoprecipitates obtained using the

anti-sfAkt-C antibody contained a trace amount of SGK (Fig. 1d, SGK with long

exposure), this was not due to cross-reaction of SGK with this antibody because a

similar amount of SGK was detected in a control precipitation obtained using a control

IgG (Fig. 1d, SGK with long exposure). Furthermore, the amount of SGK was not

reduced in extracts after immunoprecipitation of Akt (Fig. 1c). These observations

confirm the specificity of the anti-sfAkt-C antibody. The anti-sfSGK-C antibody

specifically detected SGK by immunoblotting (Fig. 1e). In addition, the protein level of

Akt remained constant before and after depletion of SGK using this antibody (Fig. 1f),

confirming the specificity of the anti-sfSGK-C antibody.

The antigens of these antibodies include the phosphorylation site in the HM of

each protein. Thus, the antibodies were expected to bind to the HM and block its

phosphorylation. As expected, microinjection of the anti-sfAkt-C antibody abolished

HM phosphorylation of Akt after 1-MeAde-stimulation (Fig. 2b). We previously

suggested that A-loop phosphorylation of Akt depends on its HM phosphorylation in

starfish oocytes based on analysis using exogenously expressed human Akt[47].

Consistently, A-loop phosphorylation was also disrupted by injection of the

anti-sfAkt-C antibody (Fig. 2c, closed arrowhead). SGK was activated as normal in

these oocytes (Fig. 2b, open arrowhead). Thus, the anti-sfAkt-C antibody specifically

inhibits Akt. On the other hand, microinjection of the anti-sfSGK-C antibody disrupted

both A-loop phosphorylation and the mobility shift of SGK after 1-MeAde stimulation

(Fig. 2d) without affecting phosphorylation of Akt (Fig. 2d for HM, e for A-loop, closed

arrowhead), indicating that this antibody specifically inhibits SGK.

We next examined the effect of microinjection of these antibodies on the meiotic

G2/M transition. Oocytes were injected with the anti-sfAkt-C antibody and then treated

with 1-MeAde. As a marker of the meiotic G2/M transition, we monitored germinal

vesicle breakdown (GVBD), which is equivalent to nuclear envelope breakdown in

somatic cells. Surprisingly, GVBD occurred as normal in these Akt-inhibited oocytes

(Fig. 2f). By contrast, microinjection of the anti-sfSGK-C antibody suppressed GVBD

in a dose-dependent manner (Fig. 2g), suggesting that SGK, but not Akt, is required for

the meiotic G2/M transition.

We previously reported that microinjection of an anti-sfAkt antibody inhibits the

meiotic G2/M transition[39]. The antibody used in that previous study was raised against

an 88-amino acid C-terminal fragment of sfAkt. In the present study, there were no

stocks of the antiserum left and therefore we used an antibody purified from an

antiserum raised against the same fragment of sfAkt in another rabbit. Thus, the

specificity of the previously used antibody may differ from that of the presently used

antibody. In addition, 33 of the 88 residues in the antigen peptide are conserved in

sfSGK and therefore the previously used antibody may have cross-reacted with sfSGK

and consequently inhibited SGK as well as Akt. Check of the specificity of this

antibody was limited during our previous study because we lacked an anti-sfSGK

antibody and phospho-specific antibodies against both sfAkt and sfSGK at that time.

SGK directly phosphorylates Cdc25 and Myt1 to trigger cyclin B-Cdk1 activation

To trigger activation of cyclin B-Cdk1, we previously demonstrated that Cdc25 and

Myt1 are activated and inactivated, respectively, by phosphorylation of Akt/SGK

consensus motifs, which correspond to multiple sites including Ser188 in Cdc25 (ref.

48), and Ser75 in Myt1 (ref. 39). Consistent with these previous findings and with the

requirement of SGK for the meiotic G2/M transition, phosphorylation of Cdc25 at

Ser188 and Myt1 at Ser75 was inhibited by injection of the anti-sfSGK-C antibody (Fig.

2d), but not by injection of the anti-sfAkt-C antibody (Fig. 2b). Furthermore, in

Phos-tag SDS-polyacrylamide gel electrophoresis (PAGE) analysis, which emphasizes

the mobility shifts of phosphorylated proteins, the upward shifts of Cdc25 and Myt1

after 1-MeAde stimulation were completely inhibited by the anti-sfSGK-C antibody,

but not by the anti-sfAkt-C antibody (Fig. 3a). This suggests that the regulatory

phosphorylation of Cdc25 and Myt1 to trigger activation of cyclin B-Cdk1 depends on

SGK, but not on Akt.

To investigate whether SGK directly phosphorylates Cdc25 and Myt1, *in vitro*

phosphorylation experiments were performed using recombinant proteins. Attempts to

prepare active recombinant sfSGK were unsuccessful; therefore, we used commercially

available recombinant human SGK3 (hSGK3), which is the best related to sfSGK

among the three isoforms of human SGK. Ser188 of starfish Cdc25 (sfCdc25) was

phosphorylated upon incubation with hSGK3 (Fig. 3b). In addition, the mobility shift

was still observed when Ser188 of sfCdc25 was substituted with Ala (Fig. 3b),

suggesting that hSGK3 directly phosphorylates sfCdc25 at multiple residues including

Ser188. This is consistent with the fact that sfCdc25 has five Akt/SGK consensus motifs.

hSGK3 also phosphorylated Ser75 of starfish Myt1 (sfMyt1), leading to a slight

mobility shift (Fig. 3c). Consistent with the fact that sfMyt1 has only one Akt/SGK

consensus motif, the mobility shift of sfMyt1 was not observed when Ser75 was

substituted with Ala (Fig. 3c), suggesting that Ser75 is the only site phosphorylated by

hSGK3 in sfMyt1. These results suggest that Cdc25 and Myt1 are direct substrates for

SGK in starfish oocytes.

The importance of SGK in 1-MeAde signaling was further supported by

experiments using oocyte extracts. A GST-conjugated Akt/SGK substrate peptide

(GST-AS peptide), which contained an Akt/SGK consensus motif, was incubated in

261 oocyte extracts. Thereafter, phosphorylation of the peptide was evaluated by

262 immunoblotting with an anti-pan-phospho-Akt/SGK substrate antibody. The GST-AS

263 peptide was not phosphorylated in the extract of unstimulated oocytes, but was

264 phosphorylated in the extract of 1-MeAde-treated oocytes (Fig. 4a, b). This

265 phosphorylation of the GST-AS peptide was not affected by immunodepletion of Akt

266 (Fig. 4c for confirmation of Akt depletion and d, e for activity), but was abolished upon

267 depletion of SGK (Fig. 4f for confirmation of SGK depletion and g, h for activity).

268 These results suggest that SGK is a major kinase activated after 1-MeAde stimulation

269 and phosphorylates the Akt/SGK consensus motif in starfish oocytes.

270  Taken together, these observations support the conclusion that SGK, but not Akt,

271 is required for the regulatory phosphorylation that makes Cdc25 activity dominant over

272 Myt1 activity to trigger cyclin B-Cdk1 activation.

274 **SGK is sufficient for the regulatory phosphorylation of Cdc25 and Myt1 to trigger**

275 **cyclin B-Cdk1 activation at the meiotic G2/M transition**

276 Next, we investigated whether sfSGK is sufficient to trigger cyclin B-Cdk1 activation.

277 Mammalian SGK3 and Akt have a regulatory domain N-terminal to their catalytic

278 domain, which is named the PX domain and PH domain, respectively[51]. Binding of

279 these domains to phosphoinositides on the cellular membrane contributes to activation

280 of these kinases[55,56]. Membrane targeting of Akt via substitution of the PH domain with

281 a myristoylation sequence results in its activation without extracellular stimulation[39,47,57].

282 To generate constitutively active sfSGK (CA-SGK), we substituted the PX domain of

283 sfSGK with a myristoylation sequence. Furthermore, Thr479, which is the conserved

284 phosphorylation site by TORC2, was substituted with Glu to mimic phosphorylation of

the HM. Injection of mRNA encoding CA-SGK into unstimulated oocytes induced

GVBD (Fig. 5a). In these oocytes, the level of A-loop phosphorylation of CA-SGK at

GVBD was comparable to that of endogenous SGK after 1-MeAde stimulation (Fig. 5b,

c), suggesting that CA-SGK is activated in the absence of 1-MeAde stimulation and that

activation of SGK is sufficient to induce the meiotic G2/M transition.

We then examined whether CA-SGK expression induces the regulatory

phosphorylation of Cdc25 and Myt1 to trigger cyclin B-Cdk1 activation. To analyze

CA-SGK-induced signaling without any effect of Cdk-dependent feedback pathways,

CA-SGK was expressed in the presence of the Cdk inhibitor roscovitine. These oocytes

were collected at the time point at which 50% of CA-SGK-expressing oocytes exhibited

GVBD in the absence of roscovitine (5 h and 45 min in Fig. 5a). Although expression

and A-loop phosphorylation levels of CA-SGK were lower than those of endogenous

SGK (Fig. 5d, pA-loop), phosphorylation of Cdc25 at Ser188 and of Myt1 at Ser75

were detectable in these oocytes (Fig. 5d). Taken together, these findings suggest that

activation of SGK is sufficient for the regulatory phosphorylation of Cdc25 and Myt1 to

trigger cyclin B-Cdk1 activation.

SGK is activated by cooperation of the Gβγ-PI3K and atypical Gβγ pathways in

starfish oocytes

In a previous report[48], we proposed the existence of an atypical Gβγ pathway that is

activated by Gβγ and cooperates with the Gβγ-PI3K pathway to induce the meiotic

G2/M transition. Therefore, we examined whether SGK is activated by these pathways.

First, we investigated whether SGK is activated by signaling pathways

downstream of Gβγ. Expression of exogenous Gβγ induced the meiotic G2/M transition

as previously reported[48], whereas Gβγ-induced GVBD was inhibited by microinjection

of the anti-sfSGK-C antibody, but not by injection of the anti-sfAkt-C antibody (Fig.

6a). These findings indicate that signaling from exogenous Gβγ induces the meiotic

G2/M transition in a SGK-dependent manner, similar to 1-MeAde stimulation.

Consistently, immunoblot analysis showed that SGK was activated upon expression of

Gβγ (Fig. 6b). Furthermore, when Gβγ was expressed in the presence of the Cdk

inhibitor roscovitine, thereby avoiding any effect of cell cycle progression on signal

transduction, the level of A-loop phosphorylation of sfSGK was comparable to that

induced by 1-MeAde (Fig. 6c), suggesting that sfSGK is activated downstream of Gβγ

independently of Cdk activity. However, 1-MeAde- and Gβγ-induced SGK activation

was abolished upon inhibition of PI3K by wortmannin (Fig. 6d), suggesting that

activation of sfSGK by Gβγ requires PI3K.

Next, to determine whether PI3K activation solely induces SGK activation,

CA-PI3K was expressed in unstimulated oocytes. As previously reported[48], CA-PI3K

induced HM phosphorylation of Akt (Fig. 6e), but not the meiotic G2/M transition. In

these oocytes, A-loop phosphorylation of SGK was only detectable by immunoblotting

with a long time exposure for acquisition of the chemiluminescent image (Fig. 6e,

pA-loop with long exposure, open arrowhead), indicating that activation of PI3K barely

induces activation of SGK.

Taken together, Gβγ activates SGK in a PI3K-dependent manner, but activation of

PI3K alone is insufficient for SGK activation. These observations suggest that Gβγ

activates not only PI3K but also another pathway(s), which we call the atypical Gβγ

pathway, and that cooperation of the Gβγ-PI3K and atypical Gβγ pathways leads to

activation of SGK.

As to activation of Akt, CA-PI3K induced HM phosphorylation of Akt (Fig. 6e),

as previously reported[48]. In these oocytes, the anti-phospho-A-loop antibody detected a

weak band at the same mobility as Akt by immunoblotting after a long exposure (Fig.

6e, pA-loop with long exposure, closed arrowhead) even in whole-oocyte samples. This

band disappeared after immunodepletion of Akt from the oocyte extract (Fig. 6e),

showing that it corresponds to A-loop phosphorylation of Akt. Given that the level of

1-MeAde-induced A-loop phosphorylation of Akt was below the threshold of detection

in whole-oocyte samples (Fig. 1b–d), Akt is activated more strongly by CA-PI3K than

by 1-MeAde. This further indicates that activation of endogenous Akt is insufficient for

the meiotic G2/M transition. By contrast, in Gβγ-expressing oocytes, A-loop

phosphorylation of Akt was undetectable even in immunoprecipitated Akt samples (Fig.

6f, closed arrowhead), although HM phosphorylation was comparable to that in

1-MeAde-treated oocytes (Fig. 6c, Akt-pHM in Input). This suggests that Gβγ signaling

is insufficient for A-loop phosphorylation of Akt. This finding will be discussed later.

Discussion

This study showed that SGK is necessary and sufficient for the regulatory

phosphorylation of Cdc25 and Myt1 to trigger activation of cyclin B-Cdk1 in the

meiotic G2/M transition. Furthermore, we revealed that SGK is activated by

cooperation of the Gβγ-PI3K and atypical Gβγ pathways downstream of 1-MeAde

stimulation (Fig. 7). These findings clarify the molecular identity of the trigger kinase,

and offer important clues to elucidate the 1-MeAde signaling pathway. Moreover, they

provide new insights into the role and regulation of SGK.

In *Xenopus* oocytes, multiple pathways are proposed to reverse the balance

between the activities of Cdc25 and Myt1 (refs 23,30). Therefore, it is difficult to

determine the relative contribution of each pathway. In this context, SGK is the only

trigger kinase in starfish oocytes because it is necessary and sufficient for the regulatory

phosphorylation of Cdc25 and Myt1. Particularly, the phosphorylation of Cdc25 and

Myt1 was undetectable in SGK-inhibited oocytes (Fig. 2). In mouse oocytes, Cyclin

B-Cdk1 activation might be triggered by a phosphatase antagonistic to PKA, rather than

by kinases, and it is unclear whether the activity of this phosphatase is regulated at the

meiotic G2/M transition[29]. Thus, our present study is the first time that the trigger

kinase in oocyte meiosis has been clearly identified. To the best of our knowledge, this

is the first report that SGK is involved in M-phase entry.

In parallel, Hosoda et al. in the accompanying paper showed that sfSGK is

required for the increase in intracellular pH after 1-MeAde stimulation in starfish

oocytes[54]. They suggested that the increase in intracellular pH is not involved in cyclin

B-Cdk1 activation, but is required for progression of GVBD after full activation of

cyclin B-Cdk1 in ovarian oocytes. The functions of SGK have been mainly studied in

mammalian cells, which express three SGK isoforms: SGK1, SGK2, and SGK3[51].

Although mammalian SGKs are reportedly involved in various processes, such as cell

survival, stress responses, and tumorigenesis, in somatic cells[58,59], their functions in

germ cells remain unclear. To the best of our knowledge, studies presented here and in

the paper by Hosoda et al.[54] are the first indication of the functions of SGK in germ

cells.

Among the three mammalian SGK isoforms, only SGK3 contains a regulatory

domain, called the PX domain, N-terminal to the kinase domain[51,60]. sfSGK also has a

PX domain and is therefore likely an ortholog of mammalian SGK3 (ref. 54). The PX

domain of mammalian SGK3 binds to phosphatidylinositol 3-phosphate (PI-3P), which

is a phosphorylated derivative of phosphatidylinositol (PI) at the 3-position of the

inositol ring[56]. This interaction is essential for the localization and activation of SGK3

(refs 50,56). PI-3P is generated by dephosphorylation of PI-345P$_3$ at the 4- and

5-positions of the inositol ring[50]. PI3K, which produces PI-345P$_3$ from PI-45P$_2$, is

activated in signaling pathways downstream of receptor tyrosine kinases and GPCRs[45].

Thus, production of PI-3P from PI-345P$_3$ may occur downstream of these receptors.

Indeed, a recent report suggested that this mechanism is implicated in activation of

SGK3 after stimulation by IGF-1, which is an agonist for a receptor tyrosine kinase[61]. In

this context, SHIP2 and INPP4A/B were proposed to be phosphatases for PI-3P

production, although the regulatory mechanisms remain elusive[61]. Furthermore, it is

unclear whether PI-345P$_3$ is a source of PI-3P for SGK activation in GPCR-dependent

signaling. We previously reported that PI-345P$_3$ is generated on the plasma membrane

after 1-MeAde stimulation in starfish oocytes[48]. This PI-345P$_3$ may be metabolized to

PI-3P for SGK activation. Our present results suggest that the atypical Gβγ pathway

cooperates with PI3K to activate sfSGK (Fig. 6). Thus, we propose that the atypical

Gβγ pathway activates phosphatases that produce PI-3P from PI-345P$_3$ downstream of

GPCR-mediated signaling.

398   Another mechanism for PI-3P production is phosphorylation of PI at the

3-position, which occurs on early endosomes during intracellular vesicle trafficking and

endocytosis[45,50]. GPCRs are frequently endocytosed after agonist stimulation[62]. Thus,

1-MeAde may induce endocytosis, which increases the level of early endosomes. If this

is the case, the endocytic pathway may also contribute to activation of SGK in starfish

oocytes. Many aspects of these scenarios, such as the dynamics of PI-3P, the

localization of SGK, and the molecular identity of the atypical Gβγ pathway, remain

unclear. Nonetheless, these working models could be tested in future studies and may

help to elucidate the regulatory mechanisms of SGK3 activation in mammalian somatic

cells.

408   SGKs and Akt reportedly play redundant roles in some contexts[59]. In starfish

oocytes, Akt has been reported to have a potential to trigger cyclin B-Cdk1 activation[39].

Indeed, an excess amount of CA-Akt expression can induce the meiotic G2/M

transition[39,48]. Furthermore, regulatory effects of phosphorylation of the Akt/SGK

consensus motifs on Myt1 and Cdc25 activities were originally showed using Akt as a

kinase in *in vitro* experiments[39,48]. Nonetheless, our present results showed that SGK is

predominantly responsible for these phosphorylation *in ovo*. Contribution of Akt was, if

any, undetectable (Fig. 2) possibly due to its weak activation, as supported by low

levels of A-loop phosphorylation (Figs. 1) and by the kinase assay using oocyte extracts

(Fig. 5). The role of Akt may be other than the triggering of cyclin B-Cdk1 activation.

Even so, we cannot exclude the possibility that Akt makes a significant contribution to

the triggering as a redundant kinase with SGK in a limited number of unknown

circumstances in which Akt is more abundant than SGK in oocytes.

Contrary to our present conclusion, we previously reported that Akt is necessary

for the meiotic G2/M transition[39]. In our previous report, in addition to the anti-sfAkt

antibody, we also used competitive peptides derived from the A-loop or HM of Akt to

inhibit Akt activation. Akt and SGK have common upstream kinases; therefore, peptide

competition likely inhibited activation of both Akt and SGK. In the present study,

successful generation of anti-phospho-A-loop and anti-sfSGK-C antibodies enabled us

to evaluate the specificities of the neutralizing antibodies more precisely. In addition,

Akt was dispensable at least in all cases tested in the current study, which used oocytes

isolated from six starfish collected at three different areas in Japan. Thus, our current

conclusions are more convincing than our previous conclusions.

We made some interesting observations regarding activation of Akt. Distinct from

SGK, activation of Akt depends on binding of PI-345P$_3$ to its PH domain[59]. Expression

of CA-PI3K, but not of Gβγ, induced A-loop phosphorylation of Akt (Fig. 6e, f). We

previously reported that a comparable level of PI-345P$_3$ is produced upon 1-MeAde

stimulation and Gβγ expression, while CA-PI3K induces a higher level of PI-345P$_3$ (ref.

48). Thus, it is possible that a high level of PI-345P$_3$ alone can induce Akt activation,

whereas a low level of PI-345P$_3$ requires an additional mechanism to enhance A-loop

phosphorylation. In addition, it should be noted that the levels of HM phosphorylation

induced by CA-PI3K and exogenous Gβγ were comparable to that induced by 1-MeAde

stimulation (Fig. 6c, e). Thus, the threshold level of PI-345P$_3$ required for HM

phosphorylation may be lower than that required for A-loop phosphorylation.

In summary, we identified SGK as the trigger kinase for activation of cyclin

B-Cdk1 at M-phase entry and as the target of the collaborative actions of PI3K and the

atypical Gβγ pathway. These findings not only demonstrate the novel role of SGK in

the triggering of M-phase, but also provide insights into the regulatory mechanisms of

SGK activation, particularly with regard to PI-3P production.

Methods

Chemicals. Roscovitine (Calbiochem), wortmannin (LC Laboratories), and BX795 (Enzo Life Sciences) were dissolved in DMSO at concentrations of 50, 20, and 6 mM, respectively, as stock solutions and used at final concentrations of 45, 40, and 6 μM, respectively.

DNA constructs. For CA-SGK, an N-terminal fragment of sfSGK containing the PX domain (Met1−Asp132) was substituted with a myristoylation sequence (MGSSKSKPKDPSQR) by PCR. Thereafter, the insert was cloned into a modified pSP64-S vector, in which the SP6 promoter for *in vitro* transcription had been substituted by a T7 promoter sequence[48,63], using an In-Fusion kit (Takara Bio). Finally, Thr479 in the HM was substituted with Glu by PCR, and the point mutation was verified by sequencing. Constructs encoding untagged sfAkt, CA-PI3K in which Flag tag and plasma membrane-targeting sequence including a CAAX motif fused to the N- and C-terminus of the p110 catalytic subunit of sfPI3K, respectively, untagged starfish Gβ (sfGβ), and untagged starfish Gγ (sfGγ) for mRNA preparation were prepared as described previously[48].

Preparation of recombinant proteins. Recombinant sfMyt1 (N229A for the kinase-dead mutant or S75A mutant) cloned into the pGEX-4T-1 vector was expressed in *Escherichia coli* (*E. coli*) strain BL21 (DE3) (Invitrogen) as an N-terminal GST-tagged protein, purified using glutathione-Sepharose 4B beads (GE Healthcare), and dialyzed against storage buffer (20 mM PIPES pH 6.8, 200 mM sucrose, and 1 mM DTT). For His_6-Cdc25 (wild-type or S188A mutant), *E. coli* strain BL21 (DE3) was

transformed with the pET21a construct. Expressed His_6-Cdc25 were purified from

inclusion bodies under denaturing conditions (6 M urea) using Probond-Resin

(Invitrogen) and then refolded via stepwise reduction of the urea concentration by

dialysis using EasySep (TOMY) for 2 h against Tris-buffered saline (TBS; 50 mM Tris

and 150 mM NaCl, pH 7.5) containing 4 M urea, for 2 h against TBS containing 2 M

urea, and for 2 h against TBS lacking urea. To prepare the recombinant GST-fused AS

peptide (AGRPRAATFIESG), *E. coli* strain BL21-CodonPlus-RIL (Agilent

Technologies) was transformed with the pGEX-4T-1 construct. The expressed GST-AS

peptide was purified using glutathione-Sepharose 4B and dialyzed against

phosphate-buffered saline (137 mM NaCl, 2.68 KCl, 10 mM Na_2HPO_4, and 1.76 mM

KH_2PO_4, pH 7.4). A recombinant C-terminal 88-amino acid fragment of sfAkt was

prepared as previously described[39].

Antibody generation. Antibodies were generated by immunizing rabbits (Biologica)

and then purified using antigens. Antigens used for immunization and antibody

purification were a phospho-peptide of the A-loop of sfAkt (Thr304–Pro320,

phosphorylated at Thr315) for the anti-phospho-A-loop antibody, a C-terminal peptide

of sfSGK (Ser473–Asp489) for the anti-sfSGK-C antibody[54], and a C-terminal

88-amino acid fragment of sfAkt for the anti-sfAkt-C antibody. Control IgG for

microinjection or immunoprecipitation was purified from rabbit pre-immune serum

using protein A-Sepharose 4B (Sigma).

Oocyte preparation. Starfish *Asterina pectinifera* (renamed *Patiria pectinifera* in the

2007 National Center for Biotechnology Information Taxonomy Browser) was collected

during the breeding season and kept in laboratory aquaria supplied with circulating

seawater at 14°C. Fully grown unstimulated oocytes with follicle cells were manually

isolated from ovaries using forceps. Thereafter, the follicle cells were removed by

washing oocytes with calcium-free artificial seawater (476 mM NaCl, 10 mM KCl, 36

mM MgCl$_2$, 18 mM MgSO$_4$, and 20 mM H$_3$BO$_3$, pH 8.2). All treatments with 1-MeAde

or other chemicals, as well as microinjections, were performed in artificial seawater

(ASW; 462 mM NaCl, 9 mM CaCl$_2$, 10 mM KCl, 36 mM MgCl$_2$, 18 mM MgSO$_4$, and

20 mM H$_3$BO$_3$, pH 8.2) at 23°C. Oocytes were stimulated with 0.5 μM 1-MeAde.

Immunoblotting. Oocytes were lysed by vortexing in Laemmli sample buffer (LSB)

and then heated at 95°C for 5 min. Proteins were separated by SDS-PAGE[64] on an 8%

or 8.5% gel (separating gel: 8% or 8.5% acrylamide, 0.22% or 0.23%

N,N'-methylenebisacrylamide, 375 mM Tris, 0.1% SDS, 0.1% APS, and 0.1% TEMED,

pH 8.8; stacking gel: 3.6% acrylamide, 0.096% N,N'-methylenebisacrylamide, 120 mM

Tris, 0.1% SDS, 0.2% APS, and 0.1% TEMED, pH 6.8; and electrophoresis buffer: 25

mM Tris, 192 mM glycine, and 0.1% SDS) and transferred to a PVDF membrane

(Millipore)[65] using transfer buffer (100 mM Tris, 192 mM glycine, 0.1% SDS, and 20%

methanol). Phos-tag SDS-PAGE was performed in accordance with the manufacturer's

protocol using an 8% polyacrylamide gel containing 4 μM Phos-tag (FUJIFILM) and 8

μM MnCl$_2$. A modified 15% gel and buffer were used to detect Gγ (separating gel: 15%

acrylamide, 0.41% N,N'-methylenebisacrylamide, 750 mM Tris, 0.1% SDS, 0.1% APS,

and 0.1% TEMED, pH 8.8; stacking gel: 3.6% acrylamide, 0.096%

N,N'-methylenebisacrylamide, 240 mM Tris, 0.1% SDS, 0.2% APS, and 0.1% TEMED,

pH 6.8; and electrophoresis buffer: 50 mM Tris, 384 mM glycine, and 0.1% SDS). The

membranes were then blocked by incubation in blocking buffer (5% skimmed milk

prepared in TBS containing 0.1% Tween-20 (TBS-T)) for 1 h for the

anti-phospho-sfMyt1-Ser75, anti-sfMyt1, and anti-GST antibodies, but not for the other

antibodies. The primary antibodies were anti-phospho-A-loop (purified, 1:50 in TBS-T),

anti-sfAkt-C (purified, 1:1,000 in Can Get Signal Solution 1),

anti-phospho-sfAkt-Ser477 (ref. 47, purified, 1:1,000 in Can Get Signal Solution 1),

anti-sfSGK-C (purified, 1:2,000 in Can Get Signal Solution 1),

anti-phospho-sfCdc25-Ser188 (ref. 48, serum, 1:1,000 in Can Get Signal Solution 1),

anti-sfCdc25 (ref. 33, serum, 1:2,000 in Can Get Signal Solution 1),

anti-phospho-Cdk1-Tyr15 (Cell Signaling Technology, 1:1,000 in Can Get Signal

Solution 1), anti-PSTAIR[47] to detect Cdk1 (1:50,000 in Can Get Signal Solution 1),

anti-phospho-sfMyt1-Ser75 (ref. 48, purified, 1:100 in blocking buffer), anti-sfMyt1[39]

(purified, 1:200 in TBS-T), anti-GST (GE Healthcare, 1:500 in blocking buffer),

anti-pan-phospho-Akt/SGK substrate (Cell Signaling Technology, 1:1,000 in Can Get

Signal Solution 1), anti-sfp110β to detect sfPI3K[48] (purified, 1:1,000 in Can Get Signal

Solution 1), anti-sfGβ[48] (serum, 1:1,000 in Can Get Signal Solution 1), and anti-sfGγ[48]

(purified, 1:100 in Can Get Signal Solution 1). Horseradish peroxidase-conjugated

secondary antibodies were anti-goat IgG (Sigma, 1:500 in blocking buffer for anti-GST

antibody), anti-mouse IgG (Dako, 1:2,000 in Can Get Signal Solution 2 for

anti-PSTAIR antibody), and anti-rabbit IgG (GE Healthcare; 1:2,000 in TBS-T for

anti-phospho-A-loop, anti-sfSGK-C, anti-phospho-sfMyt1-Ser75, and

anti-pan-phospho-Akt/SGK substrate antibodies; in blocking buffer for the anti-sfMyt1

antibody; and in Can Get Signal Solution 2 for the other antibodies). When the signal

derived from IgG heavy chains hampered detection, TrueBlot (ROCKLAND, 1:2,000 in

Can Get Signal Solution 2) was used instead of the secondary antibody. Signals were

visualized with ECL Prime (GE Healthcare), and digital images were acquired on a

LAS4000 mini imager (Fujifilm). Quantitation was performed using ImageJ (National

Institutes of Health, USA). Graphs were generated using Microsoft Excel. Brightness

and contrast were adjusted using ImageJ.

Microinjection. Microinjection was performed as described previously[66]. mRNAs for

exogenous protein expression in starfish oocytes were transcribed from pSP64-S

constructs using a mMESSAGE mMACHINE kit (Ambion), dissolved in water, and

injected into unstimulated oocytes (20 pg for sfAkt, 20 pg for CA-SGK, and 40 pg for

CA-PI3K). To express starfish G$\beta\gamma$, 50 pg of an equimolar mixture of mRNAs encoding

sfGβ and sfGγ was injected. The incubation time for protein expression was determined

based on the amount of time taken to induce GVBD or to induce HM phosphorylation

of Akt to a comparable extent as that induced by 1-MeAde. For microinjection of

antibodies, concentration of antibodies and buffer exchange with phosphate-buffered

saline were performed using a 50 k cutoff Amicon Ultra filter (Millipore). Nonidet P-40

(NP-40) was added at a final concentration of 0.05%. Unstimulated oocytes were

injected with 230 pg of the anti-TOR antibody, 85 pg of the anti-sfAkt-C antibody, or

200 pg of the anti-sfSGK-C antibody. 1-MeAde was added after incubation for 1 h.

Immunoprecipitation and immunodepletion. To prepare antibody-bound beads,

antibodies (anti-sfAkt antibody, 2.5 μg/μl beads, 0.83 μg for 10 oocytes; and

anti-sfSGK antibody, 2 μg/μl beads, 2.7 μg for 10 oocytes) were incubated with Protein

A-Sepharose (Sigma) for more than 2 h on ice. Antibody-bound beads were washed

with TBS and lysis buffer (80 mM β-glycerophosphate, 20 mM EGTA, 10 mM MOPS

pH 7.0, 100 mM sucrose, 100 mM KCl, 1 mM DTT, 1× cOmplete EDTA-free (Merck),

0.5 mM sodium orthovanadate, 1 μM okadaic acid (Calbiochem), and 0.5% NP-40) and

used for immunoprecipitation. For cross-linking, antibody-bound beads were washed

with TBS and borate buffer (0.2 M NaCl and 0.2 M boric acid pH 9.0) and then

incubated with 3.75 mM disuccinimidyl suberate prepared in borate buffer at 24°C for

30 min. Thereafter, the supernatant was discarded and the beads were further incubated

in 1 M ethanolamine (pH 8.0) at 24°C for 15 min and finally washed with TBS and lysis

buffer. For immunoprecipitation, freeze-thawed oocytes in ASW (10 oocytes/μl of

ASW for immunoprecipitation of Akt and SGK) were lysed by adding 6 volumes of

lysis buffer and incubating samples on ice for 30 min, followed by gentle vortexing.

Lysates were centrifuged at 20 k × g at 4°C for 15 min. The supernatant represented the

oocyte extract. Antibody-bound beads were incubated with the oocyte extract (60

oocytes for immunoprecipitation of Akt, 30 oocytes for depletion of Akt, and 15

oocytes for immunoprecipitation and depletion of SGK) for 90 min on ice, separated

from flow-through extracts, and washed with lysis buffer. For immunoprecipitation of

Akt from oocytes that had been injected with the anti-sfSGK antibody, extracts were

pre-incubated with protein A-Sepharose 4B to remove injected IgG. LSB was added to

the input extracts, flow-through extracts, and beads for immunoblot analysis. All

samples were then heated at 95°C for 5 min and subjected to SDS-PAGE. To prepare

mock-, Akt-, and SGK-depleted extracts for the GST-AS peptide kinase assay,

freeze-thawed oocytes in ASW (5 oocytes/μl of ASW) were lysed by adding 6.5

volumes of lysis buffer. The extract was then prepared as described above and mixed

with beads bound to control IgG, the anti-sfAkt-C antibody, or the anti-sfSGK-C

antibody. After incubation at 4°C for 45 min, the supernatant was analyzed by

immunoblotting or used for the GST-AS peptide kinase assay.

***In vitro* phosphorylation of sfCdc25 and sfMyt1.** Recombinant sfCdc25 (final

concentration of 5 ng/μl) or sfMyt1 (final concentration of 200 ng/μl) was incubated

with N-terminal GST-tagged human SGK3 (final concentration of 10 ng/μl,

SignalChem) in reaction buffer (80 mM β-glycerophosphate, 20 mM EGTA, 1 mM

MOPS pH 7.0, 1 mM DTT, 1 mM ATP, 5 mM $MgCl_2$, and 0.3% NP-40) at 30°C for 50

min. The reaction was stopped by adding LSB followed by heating at 95°C for 5 min.

Samples were analyzed by immunoblotting with phospho-specific antibodies.

GST-AS peptide kinase assay. Freeze-thawed oocytes in ASW (5 oocytes/μl of ASW)

were lysed by adding 6.5 volumes of lysis buffer and then incubating samples on ice for

30 min, followed by gentle vortexing. Lysates were centrifuged at 20 k × g at 4°C for

15 min. The supernatant was used for the kinase assay. Mock, Akt-, and SGK-depleted

extracts were prepared as described above. For the kinase assay, a reaction mixture

containing the GST-AS peptide as a substrate (80 mM β-glycerophosphate, 15 mM

$MgCl_2$, 20 mM EGTA, 10 mM MOPS, pH 7.0, 1 mM DTT, 1× cOmplete EDTA-free, 2

mM ATP, 0.5 mM sodium orthovanadate, 1 μM okadaic acid, 0.1 mg/ml GST-AS

peptide, and 0.3% NP-40) was added to an equal volume of the extract. After incubation

at 30°C, the reaction solution was mixed with LSB, heated at 95°C for 5 min, and then

analyzed by immunoblotting.

Statistical analysis. Statistical significance was calculated using the unpaired one-tailed

Student's t-test in Microsoft Excel with StatPlus (AnalystSoft). Statistical significance

was defined as $p < 0.05$.

Accession numbers. Accession numbers are as follows: LC430700 for sfSGK,

AB076395 for sfCdc25, AB060280 for sfMyt1, AB060291 for sfAkt, LC017891 for

sfGβ, LC017892 for sfGγ, and LC017893 for the sfPI3K catalytic subunit p110β.

Data availability

Data supporting the findings of this study are available within the article or upon

reasonable requests to the corresponding authors.

Acknowledgements

We thank E. Okumura for helpful discussions, suggestions, and providing recombinant

Cdc25 and Myt1, as well as M. Terasaki for providing the pSP64-S vector. This work

was supported by grants-in-aid to T.K. from JSPS [grant numbers 25291043,

16H04783] and Takeda Science Foundation.

Author contributions

D.H. designed and performed all the experiments. E.H. and K.C. generated the

anti-sfSGK-C antibody. All authors discussed the results. D.H. and T.K. wrote the

manuscript. T.K. obtained funding and supervised the study.

Competing interests

The authors declare no competing interests.

Materials and correspondence

Correspondence and requests for materials should be addressed to D.H. and T.K.

References

References

1.   Dunphy, W. G. & Newport, J. W. Unraveling of mitotic control mechanisms. *Cell*

55, 925-928 (1988).

2.   Hunt, T. Maturation promoting factor, cyclin and the control of M-phase. *Curr.*

Opin. Cell Biol. **1,** 268-274 (1989).

3.   Nurse, P. Universal control mechanism regulating onset of M-phase. *Nature* **344,**

503-508 (1990).

4.   Lew, D.J. & Kornbluth, S. Regulatory roles of cyclin dependent kinase

phosphorylation in cell cycle control. *Curr. Opin. Cell Biol.* **8,**795-804 (1996).

5.   Lindqvist, A., Rodríguez-Bravo, V. & Medema, R. H. The decision to enter

mitosis: feedback and redundancy in the mitotic entry network. *J. Cell Biol.* **185,**

193-202 (2009).

6.   Qian, J., Winkler, C. & Bollen, M. 4D-networking by mitotic phosphatases. *Curr.*

Opin. Cell Biol. **25,** 697-703 (2013).

7.   Kishimoto, T. Entry into mitosis: a solution to the decades-long enigma of MPF.

Chromosoma **124,** 417-428 (2015).

8.   Hégarat, N., Rata, S. & Hochegger, H. Bistability of mitotic entry and exit switches

during open mitosis in mammalian cells. *Bioessays* 627-643 (2016).

9.   O'Farrell, P. H. Triggering the all-or-nothing switch into mitosis. *Trends Cell*

Biol. **11,** 512-519 (2001).

10.   Rieder, C. L. Mitosis in vertebrates: the G2/M and M/A transitions and their

associated checkpoints. *Chromosome Res.* **19,** 291-306 (2011).

11.   Thomas, Y. et al. Cdk1 Phosphorylates SPAT-1/Bora to promote Plk1 activation

in *C. elegans* and human cells. *Cell Rep.* **15,** 510-518 (2016).

12.   Gheghiani, L., Loew, D., Lombard, B., Mansfeld, J. & Gavet, O. PLK1 activation

in late G2 sets up commitment to mitosis. *Cell Rep.* **19,** 2060-2073 (2017).

13.   Vigneron, S. et al. Cyclin A-cdk1-dependent phosphorylation of Bora is the

triggering factor promoting mitotic entry. *Dev Cell.* **45,** 637-650 (2018).

14.   Okano-Uchida, T. et al. Distinct regulators for Plk1 activation in starfish meiotic

and early embryonic cycles. *EMBO J.* **22,** 5633-5642 (2003).

15.   Masui, Y. Meiotic arrest in animal oocytes. *Biology of Fertilization* **1,** 189–219

(Academic Press, San Diego, 1985).

16.   Kishimoto, T. MPF-based meiotic cell cycle control: Half a century of lessons

from starfish oocytes. *Proc. Jpn. Acad. Ser. B* **94,** 180-203 (2018).

17.   Kobayashi, H. et al. On the synthesis and destruction of A- and B-type cyclins

during oogenesis and meiotic maturation in *Xenopus laevis*. *J. Cell Biol*. **114,** 755-765

(1991).

18.   Minshull, J., Murray, A., Colman, A. & Hunt, T. *Xenopus* oocyte maturation does

not require new cyclin synthesis. *J. Cell Biol.* **114,** 767-772 (1991).

19.   Okano-Uchida, T. et al. *In vivo* regulation of cyclin A/Cdc2 and cyclin B/Cdc2

through meiotic and early cleavage cycles in starfish. *Dev. Biol.* **197,** 39-53 (1998).

20.   Nebreda, A. R. & Ferby, I. Regulation of the meiotic cell cycle in oocytes. *Curr.*

Opin. Cell Biol. **12,** 666-675 (2000).

21.   Kishimoto, T. Cell-cycle control during meiotic maturation. *Curr. Opin. Cell Biol.*

2003 **15,** 654-663 (2003).

22.   Pahlavan, G. et al. Characterization of polo-like kinase 1 during meiotic

maturation of the mouse oocyte. *Dev. Biol.* **220,** 392-400 (2000).

23.   Gaffré, M. et al. A critical balance between Cyclin B synthesis and Myt1 activity

controls meiosis entry in *Xenopus* oocytes. *Development* **138,** 3735-3744 (2011).

24. Maton, G. et al. Cdc2-cyclin B triggers H3 kinase activation of Aurora-A in

Xenopus oocytes. *J. Biol. Chem.* **278,** 21439-21449 (2003).

25. Abe, Y., Okumura, E., Hosoya, T., Hirota, T. & Kishimoto, T. A single starfish

Aurora kinase performs the combined functions of Aurora-A and Aurora-B in human

cells. *J. Cell Sci.* **123,** 3978-3988 (2010).

26. Komrskova, P. et al. Aurora kinase A is not involved in CPEB1 phosphorylation

and cyclin B1 mRNA polyadenylation during meiotic maturation of porcine oocytes.

PLoS One **9,** e101222 (2014). doi: 10.1371/journal.pone.0101222.

27. Maller, J. L. & Krebs, E. G. Progesterone-stimulated meiotic cell division in

Xenopus oocytes. Induction by regulatory subunit and inhibition by catalytic subunit of

adenosine 3':5'-monophosphate-dependent protein kinase. *J. Biol. Chem.* **252,**

1712-1718 (1977).

28. Jaffe, L. A. & Egbert, J. R. Regulation of mammalian oocyte meiosis by

intercellular communication within the ovarian follicle. *Annu. Rev. Physiol.* **79,** 237-260

(2017).

29. Adhikari, D. & Liu, K. The regulation of maturation promoting factor during

prophase I arrest and meiotic entry in mammalian oocytes. *Mol. Cell. Endocrinol.* **382,**

480-487 (2014).

30. Haccard, O. & Jessus, C. Redundant pathways for Cdc2 activation in *Xenopus*

oocyte: either cyclin B or Mos synthesis. *EMBO Rep.* **7,** 321-325 (2006).

31. Haccard, O. & Jessus, C. Oocyte maturation, Mos and cyclins--a matter of

synthesis: two functionally redundant ways to induce meiotic maturation. *Cell Cycle* **5,**

1152-1159 (2006).

32.   Dupré, A., Daldello, E. M., Nairn, A. C., Jessus, C. & Haccard, O.

Phosphorylation of ARPP19 by protein kinase A prevents meiosis resumption in

Xenopus oocytes. *Nat. Commun.* **5,** 3318 (2014). doi: 10.1038/ncomms4318.

33.   Okumura, E., Sekiai, T., Hisanaga, S., Tachibana, K. & Kishimoto, T., Initial

triggering of M-phase in starfish oocytes: a possible novel component of

maturation-promoting factor besides cdc2 kinase. *J. Cell Biol.* **132,** 125-135 (1996).

34.   Meijer, L. & Guerrier, P. Maturation and fertilization in starfish oocytes. *Int.*

Rev. Cytol. **86,** 129-196 (1984).

35.   Kanatani, H., Shirai, H., Nakanishi, K. & Kurokawa, T. Isolation and

identification on meiosis inducing substance in starfish *Asterias amurensis*. *Nature*. **221,**

273-274 (1969).

36.   Dorée, M. Protein synthesis is not involved in initiation or amplification of

the maturation-promoting factor (MPF) in starfish oocytes. *Exp. Cell Res.* **139,** 127-133

(1982).

37.   Tachibana, K., Tanaka, D., Isobe, T., Kishimoto, T. c-Mos forces the mitotic cell

cycle to undergo meiosis II to produce haploid gametes. *Proc. Natl. Acad. Sci. USA* **97,**

14301-14306 (2000).

38.   Ookata, K., Hisanaga, S., Okano, T., Tachibana, K. & Kishimoto, T. Relocation

and distinct subcellular localization of p34cdc2-cyclin B complex at meiosis reinitiation

in starfish oocytes. *EMBO J.* **11**, 1763–1772 (1992).

39.   Okumura, E. et al. Akt inhibits Myt1 in the signalling pathway that leads to

meiotic G2/M-phase transition. *Nat. Cell Biol.* **4,** 111-116 (2002).

40.   Kanatani, H. & Hiramoto, Y. Site of action of 1-methyladenine in inducing oocyte

maturation in starfish. *Exp. Cell Res.* **61,** 280-284 (1970).

41.   Chiba, K., Kontani, K., Tadenuma, H., Katada, T. & Hoshi, M. Induction of

starfish oocyte maturation by the βγ subunit of starfish G protein and possible existence

of the subsequent effector in cytoplasm. *Mol. Biol. Cell* **4,** 1027–1034 (1993).

42.   Jaffe, L. A., Gallo, C. J., Lee, R. H., Ho, Y. K. & Jones, T. L. Oocyte maturation

in starfish is mediated by the βγ-subunit complex of a G-protein. *J. Cell Biol.* **121,**

775–783 (1993).

43.   Shilling, F., Chiba, K., Hoshi, M., Kishimoto, T. & Jaffe, L. A. Pertussis toxin

inhibits 1-methyladenine-induced maturation in starfish oocytes. *Dev. Biol.* **133,**

605–608 (1989).

44.   Sadler, K. C. & Ruderman, J. V. Components of the signaling pathway linking the

1-methyladenine receptor to MPF activation and maturation in starfish oocytes. *Dev.*

Biol. **197,** 25–38 (1998).

45.   Vanhaesebroeck, B., Guillermet-Guibert, J., Graupera, M. & Bilanges B. The

emerging mechanisms of isoform-specific PI3K signalling. *Nat. Rev. Mol. Cell Biol.* **11,**

329-341 (2010).

46.   Hiraoka, D. et al. PDK1 is required for the hormonal signaling pathway leading to

meiotic resumption in starfish oocytes. *Dev. Biol.* **276,** 330-336 (2004).

47.   Hiraoka, D., Okumura, E., Kishimoto, T. Turn motif phosphorylation negatively

regulates activation loop phosphorylation in Akt. *Oncogene* **30,** 4487-4497 (2011).

48.   Hiraoka, D., Aono, R., Hanada, S., Okumura, E., & Kishimoto, T. Two new

competing pathways establish the threshold for cyclin-B-Cdk1 activation at the meiotic

G2/M transition. *J. Cell Sci.* **129,** 3153-3166 (2016).

49.   Webster, M. K., Goya, L., Ge, Y., Maiyar, A. C. & Firestone, G. L.

Characterization of sgk, a novel member of the serine/threonine protein kinase gene

family which is transcriptionally induced by glucocorticoids and serum. *Mol. Cell. Biol.*

13, 2031-2040 (1993).

50. Lien, E. C., Dibble, C. C. & Toker, A. PI3K signaling in cancer: beyond AKT.

Curr. Opin. Cell Biol. **45,** 62-71(2017).

51. Pearce, L. R., Komander, D. & Alessi, D. R. The nuts and bolts of AGC protein

kinases. *Nat. Rev. Mol. Cell Biol.* **11,** 9-22 (2010).

52. Kobayashi, T. & Cohen, P. Activation of serum- and glucocorticoid-regulated

protein kinase by agonists that activate phosphatidylinositide 3-kinase is

mediated by 3-phosphoinositide-dependent protein kinase-1 (PDK1) and PDK2.

Biochem. J. **339,** 319-328 (1999).

53. García-Martínez, J. M. & Alessi, D. R. mTOR complex 2 (mTORC2) controls

hydrophobic motif phosphorylation and activation of serum- and

glucocorticoid-induced protein kinase 1 (SGK1). *Biochem. J.* **416,** 375-385 (2008).

54. Hosoda, E. et al. SGK regulates meiotic resumption in ovarian starfish oocytes via

intracellular pH-increase and cyclin B-Cdk1 activation. Accompanying manuscript

(2018)

55. Andjelković, M. et al. Activation and phosphorylation of a pleckstrin homology

domain containing protein kinase (RAC-PK/PKB) promoted by serum and protein

phosphatase inhibitors. *Proc. Natl. Acad. Sci. USA* **93,** 5699-5704 (1996).

56. Virbasius, J. V. et al. Activation of the Akt-related cytokine-independent survival

kinase requires interaction of its phox domain with endosomal phosphatidylinositol

3-phosphate. *Proc. Natl. Acad. Sci. USA* **98,** 12908-12913 (2001).

57. Kohn, A. D., Takeuchi, F. & Roth, R. A. Akt, a pleckstrin homology domain

containing kinase, is activated primarily by phosphorylation. *J. Biol. Chem.* **271,**

21920-21926 (1996).

58.   Lang, F. et al. (Patho)physiological significance of the serum- and

glucocorticoid-inducible kinase isoforms. *Physiol. Rev.* **86,** 1151-1178 (2006).

59.   Bruhn, M. A., Pearson, R. B., Hannan, R. D. & Sheppard, K. E. AKT-independent

PI3-K signaling in cancer - emerging role for SGK3. *Cancer Manag. Res.* **5,** 281-292

(2013).

60.   Liu, D., Yang, X. & Songyang, Z. Identification of CISK, a new member of the

SGK kinase family that promotes IL-3-dependent survival. *Curr. Biol.* **10,** 1233-1236

(2000).

61.   Malik, N. et al. Mechanism of activation of SGK3 by growth factors via the Class

1 and Class 3 PI3Ks. *Biochem. J.* **475,** 117-135 (2018).

62.   Tsao, P. I. & von Zastrow, M. Diversity and specificity in the regulated endocytic

membrane trafficking of G-protein-coupled receptors. *Pharmacol. Ther.* **89,** 139-147

(2001).

63.   Terasaki, M., Okumura, E., Hinkle, B. & Kishimoto, T. Localization and

dynamics of Cdc2-cyclin B during meiotic reinitiation in starfish oocytes. *Mol. Biol.*

Cell. **14,** 4685-4694 (2003).

64.   Laemmli, U. K. Cleavage of structural proteins during the assembly of the head of

bacteriophage T4. *Nature* **227,** 680-685 (1970).

65.   Towbin, H., Staehelin, T. & Gordon, J. Electrophoretic transfer of proteins from

polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proc.*

Natl. Acad. Sci. USA **76,** 4350-4354 (1979).

66.   Kishimoto, T. Microinjection and cytoplasmic transfer in starfish oocytes.

Methods. Cell Biol. **27,** 379–394 (1986).

Fig. 1 SGK is activated after 1-MeAde stimulation in a PDK1- and TORC2-dependent manner. **a** The region of sfAkt used to raise the anti-phospho-A-loop antibody (upper, Lys304–Pro320 of sfAkt) is aligned with the corresponding region of sfSGK (Lys301–Pro317). The arrowhead indicates the residue phosphorylated by PDK1 (Thr315 in sfAkt and Thr312 in sfSGK). Dots indicate identical residues. **b** A-loop phosphorylation of endogenous sfAkt is undetectable level in whole oocyte extracts. Uninjected oocytes and oocytes injected with mRNA encoding untagged sfAkt were treated with 1-MeAde, and analyzed by immunoblotting with the anti-phospho-A-loop (pA-loop) and anti-sfAkt-C (Akt) antibodies. Closed and open arrowheads indicate the positions of Akt and A-loop-phosphorylated SGK (see **e** and **f**), respectively. **c, d**

A-loop phosphorylation of endogenous Akt is detectable in immunoprecipitates.
Oocytes were incubated with 1-MeAde in the presence or absence of a PDK1 inhibitor,
BX795, lysed, and subjected to immunoprecipitation with control IgG or the
anti-sfAkt-C antibody. Input and flow-through (FT) extracts (**c**), and precipitates (**d**)
were analyzed by immunoblotting (Akt-pHM represents phosphorylation of Ser477 in
the HM of Akt). A-loop phosphorylation of non-specifically included SGK was
detected in control and Akt immunoprecipitates (**d**, open arrowhead). **e** Unstimulated
and 1-MeAde-treated oocytes were subjected to immunoblotting with
anti-phospho-A-loop (left), anti-sfSGK-C (middle), and anti-sfAkt-C (right) antibodies.
f The anti-phospho-A-loop antibody against sfAkt cross-reacts with
A-loop-phosphorylated endogenous SGK. Unstimulated oocytes were treated with
1-MeAde. Immunoprecipitation was performed with control IgG or the anti-sfSGK-C
antibody. Input, flow-through, and precipitates were analyzed by immunoblotting. **g**
SGK is activated simultaneously with Akt activation. Oocytes were treated with
1-MeAde and collected at the indicated time points. Phosphorylation of Akt (Akt-pHM),
SGK (pA-loop), Cdc25 (pS188), and Cdk1 (pY15) were analyzed by immunoblotting.
GVBD occurred around 20 min. **h** SGK activation in starfish oocytes depends on TOR.
Unstimulated oocytes were injected with control IgG or the anti-TOR antibody, treated
with 1-MeAde, and analyzed by immunoblotting. The data are representative of two
independent experiments in **b**, **e**, and **f**; and three independent experiments in **c**, **d**, and **g**.
In all panels, closed and open arrowheads indicate the positions of Akt and
A-loop-phosphorylated sfSGK, respectively.

Fig. 2 SGK, but not Akt, is required for the meiotic G2/M transition. **a**

Immunoprecipitation with control IgG or the anti-sfAkt-C antibody was performed

using extracts prepared from unstimulated oocytes, followed by immunoblotting with

the anti-sfAkt-C antibody. **b, c** The anti-sfAkt-C antibody inhibits activation of Akt.

Unstimulated oocytes were injected with the anti-sfAkt-C antibody or control IgG and

then treated with 1-MeAde for 4 min. Immunoprecipitation with the anti-sfAkt-C

antibody was performed. Input extracts (**b**, Input) and precipitates (**c**, Akt-IP) were

analyzed by immunoblotting. **d, e** The anti-sfSGK-C antibody inhibits activation of
SGK. Unstimulated oocytes were injected with the anti-sfSGK-C antibody or control
IgG and then treated with 1-MeAde for 4 min. Extracts were prepared, and injected IgG
was removed by mixing samples with protein A-Sepharose beads. Thereafter, extracts
were subjected to immunoprecipitation with the anti-sfAkt-C antibody. Input extracts (**d**,
Input) and precipitates (**e**, Akt-IP) were analyzed by immunoblotting. **f** Akt is not
required for the meiotic G2/M transition. Unstimulated oocytes were injected with the
anti-sfAkt-C antibody or control IgG and then treated with 1-MeAde. GVBD was
monitored as a marker of the meiotic G2/M transition. The graph shows the percentage
of oocytes that had undergone GVBD by the indicated time points. "n" indicates the
number of oocytes observed. **g** SGK is required for the meiotic G2/M transition.
Unstimulated oocytes were injected with the indicated amount of the anti-sfSGK-C
antibody or control IgG and then treated with 1-MeAde. GVBD was monitored and
plotted as described in **f**. In all panels, open arrowheads indicate the position of
A-loop-phosphorylated SGK, closed arrowheads indicate the position of Akt, and
asterisks indicate non-specific bands. The data are representative of two independent
experiments in **a-c**, **f**, and **g**; and three independent experiments in **d** and **e**.

Fig. 3 SGK directly phosphorylates Cdc25 and Myt1. **a** SGK, but not Akt, is required

for Cdk-independent phosphorylation of Cdc25 and Myt1 upon 1-MeAde stimulation.

Unstimulated oocytes were injected with the indicated antibodies, treated with 1-MeAde

in the presence of roscovitine, and then analyzed by immunoblotting using a normal or

Phos-tag-containing SDS-PAGE gel. The open arrowhead indicates the position of

A-loop-phosphorylated SGK, while the closed arrowheads indicate the position of Akt.

b His-tagged recombinant wild-type (WT) sfCdc25 or the S188A mutant was incubated

with or without GST-tagged active human SGK3 (hSGK3) in the presence of ATP and

$MgCl_2$. Thereafter, immunoblotting was performed. **c** GST-tagged recombinant

kinase-dead (KD) or the S75A mutant of sfMyt1 was phosphorylated by human SGK3

as described in **b**. Asterisks indicate minor fragments of recombinant Myt1. The data in

all panels are representatives of each two independent experiments.

Fig. 4 SGK is a major kinase that phosphorylates the Akt/SGK consensus motif in starfish oocytes. **a, b** Kinase activity for the AS peptide is detectable in an extract of 1-MeAde-treated oocytes. Extracts were prepared from unstimulated or 1-MeAde-treated (3 min) oocytes and incubated with the GST-AS peptide for the indicated durations. Phosphorylation of the GST-AS peptide was detected by immunoblotting with an anti-pan-phospho-Akt/SGK substrate antibody (PAS), which recognizes phosphorylated Ser or Thr in the Akt/SGK consensus motif (**a**). Phosphorylation of the GST-AS peptide was quantitated from the immunoblots in **a** and normalized against the total amount of the GST-AS peptide. The graph shows the levels of phosphorylation relative to that at 60 min (**b**). **c–e** Akt is not responsible for the AS-peptide kinase activity in the extract. An extract was prepared from 1-MeAde-treated oocytes. Immunodepletion was performed with control IgG (mock) or

the anti-sfAkt-C antibody (ΔAkt). The crude and immunodepleted extracts were

analyzed by immunoblotting (**c**). The GST-AS peptide kinase assay was performed

using the extracts shown in **c** (**d**). Phosphorylation was quantitated as described in **a** and

b. The graph shows the levels of phosphorylation relative to that at 15 min in the crude

extract (**e**). **f–h** SGK is responsible for the AS-peptide kinase activity in the extract. An

extract was prepared from unstimulated or 1-MeAde-treated oocytes. Immunodepletion

was performed with the anti-sfSGK-C antibody (**f**), the GST-AS peptide kinase assay

was performed (**g**), and phosphorylation was quantitated (**h**) as described in **c**, **d**, and **e**.

The open arrowhead indicates the position of A-loop-phosphorylated SGK, while the

closed arrowheads indicate the position of Akt. The data in all panels are representatives

of each two independent experiments.

905
Fig. 5 SGK is sufficient for the meiotic G2/M transition. **a–c** CA-SGK expression
induces the meiotic G2/M transition. Unstimulated oocytes were injected with mRNA
encoding CA-SGK and incubated. GVBD in 14 oocytes was monitored every 15 min.
The graph shows the percentage of oocytes that had undergone GVBD by the indicated
time points (**a**). CA-SGK-expressing oocytes at GVBD, unstimulated oocytes, and
1-MeAde-treated oocytes were analyzed by immunoblotting (**b**). Phosphorylation levels
of the A-loop of endogenous SGK in 1-MeAde-treated oocytes and CA-SGK at GVBD
were quantitated from the immunoblots in **b**. The graph shows the relative
phosphorylation levels (**c**). Data represent mean values ± standard deviation from three
independent experiments. P = 0.06 (one-tailed t-test; not significant, n.s.). **d** CA-SGK
expression induces regulatory phosphorylation of Cdc25 and Myt1 to trigger cyclin
B-Cdk1 activation. Unstimulated oocytes were treated with 1-MeAde for 4 min in the
presence of roscovitine. For expression of CA-SGK, unstimulated oocytes were injected
with the mRNA, incubated with roscovitine, collected at the time point at which 50% of
CA-SGK-expressing oocytes exhibited GVBD in the absence of roscovitine (5 h 45 min
in **a**), and analyzed by immunoblotting. Note that, in the immunoblot of
phospho-A-loop after a long exposure (pA-loop, long exp.), weak non-specific bands
were detected at almost the same size as CA-SGK (asterisk). The data are representative
of three independent experiments in **a**, **b**, and **c**; and two independent experiments in **d**.

925
Fig. 6 SGK is activated by cooperation of the Gβγ-PI3K and atypical Gβγ pathways. **a,**
b Exogenous expression of Gβγ induces the meiotic G2/M transition in a
SGK-dependent manner. Unstimulated oocytes were injected with the indicated
antibodies and subsequently injected with mRNA encoding Gβγ. Thereafter, GVBD
was monitored (**a**). Oocytes were collected when GVBD occurred. Anti-sfSGK
antibody-injected oocytes, which did not undergo GVBD, were collected at 5 h after
mRNA injection. For 1-MeAde treatment, unstimulated oocytes were treated with
1-MeAde for 4 or 25 min (GVBD occurred at approximately 21 min) and then analyzed
by immunoblotting (**b**). **c** Exogenous Gβγ expression induces SGK activation.
Unstimulated oocytes were treated with 1-MeAde for 4 min in the presence of
roscovitine or injected with mRNA encoding Gβγ and then incubated with roscovitine
for 6 h. Thereafter, extracts were prepared from these oocytes and analyzed by
immunoblotting. These extracts were also used for immunoprecipitation of Akt in **f**. **d**
Gβγ-induced SGK activation requires PI3K. Unstimulated oocytes were treated with
1-MeAde or injected with mRNA encoding Gβγ, incubated in the presence of

roscovitine with or without wortmannin, and analyzed by immunoblotting. **e** PI3K is not
sufficient for activation of SGK. Unstimulated oocytes were treated with 1-MeAde for 4
min or injected with mRNA encoding CA-PI3K and incubated for 3 h. Extracts of
CA-PI3K-expressing oocytes were immunodepleted using control IgG (Mock) or the
anti-sfAkt-C antibody (ΔAkt). Thereafter, immunoblotting was performed. **f** Expression
of exogenous Gβγ fails to induce A-loop phosphorylation of Akt. Endogenous Akt was
immunoprecipitated from the extracts in **c** and analyzed by immunoblotting. In all
panels, open arrowheads indicate the position of A-loop-phosphorylated SGK, while
closed arrowheads indicate the position of Akt. The data are representative of two
independent experiments in **a**, **b**, **d**, and **e**; and three independent experiments in **c** and **f**.

951
Fig. 7 SGK triggers cyclin B-Cdk1 activation in 1-MeAde signaling. Downstream of
1-MeAde stimulation, Gβγ activates the PI3K pathway (green arrow) and the atypical
Gβγ pathway (cyan arrow). SGK is activated by cooperation of these pathways and
subsequently activates and inactivates Cdc25 and Myt1, respectively, via direct
phosphorylation. Consequently, reversal of the balance of Cdc25 and Myt1 activities
triggers cyclin B-Cdk1 activation. Akt is weakly activated after 1-MeAde stimulation,
and may phosphorylate Cdc25 and Myt1 at undetectable levels. However, Akt is
dispensable for cyclin B-Cdk1 activation and may have any other function in oocytes.