The bat influenza H17N10 can be neutralized by broadly-neutralizing monoclonal antibodies and its neuraminidase can facilitate viral egress.

George Carnell [a,b1], Efstathios S Giotis[c1], Keith Grehan [a,d1], Francesca Ferrara[a], Stuart Mather[a], Eleonora Molesti[a], Simon Scott[a], Martin Schwemmle[e,f], Antonello Pessi[g], Krzysztof Lacek[g,h], & Nigel Temperton[a]

[a] Viral Pseudotype Unit, Medway School of Pharmacy, University of Kent and University of Greenwich, Chatham, United Kingdom
[b] Laboratory of Viral Zoonotics, Department of Veterinary Medicine, University of Cambridge, United Kingdom
[c] Section of Virology, Faculty of Medicine, Imperial College London, St Mary's Campus, United Kingdom
[d] School of Molecular and Cellular Biology, University of Leeds, United Kingdom
[e] Institute of Virology, Medical Center University of Freiburg, Freiburg, Germany
[f] Faculty of Medicine, University of Freiburg, 79104 Freiburg, Germany
[g] Ceinge Biotecnologie Avanzate S.C.R.L., Napoli, Italy
[h] The Laboratory of Virus Molecular Biology, University of Gdansk, 80-822 Gdansk, Poland
[1] Authors contributed equally

Corresponding author: Nigel Temperton, n.temperton@kent.ac.uk, ORCID 0000-0002-7978-3815

Key words: Bat Influenza virus, broadly neutralizing antibody, Hemagglutinin, Neuraminidase

Abstract

The diversity of subtypes within the Influenza A virus genus has recently expanded with the identification of H17N10 and H18N11 from bats. In order to further study the tropism and zoonotic potential of these viruses, we have successfully produced lentiviral pseudotypes bearing both haemagglutinin H17 and neuraminidase N10. These pseudotypes were shown to be efficiently neutralized by the broadly-neutralizing monoclonal antibodies CR9114 and FI6. Our studies also confirm previous reports that H17 does not use sialic acid as its cellular receptor, as pseudotypes bearing the H17 envelope glycoprotein are released into the cell supernatant in the absence of NA. However, we demonstrate that N10 facilitates heterosubtypic (H5 and H7) influenza HA-bearing pseudotype release in the absence of another source of NA, significantly increasing luciferase pseudotype production titres. Despite this, N10 shows no activity in the enzyme-linked lectin assay used for traditional sialidases. These findings suggest that this protein plays an important role in viral egress, but is perhaps involved in further accessory roles in the bat influenza lifecycle that are yet to be discovered. Thus we show the lentiviral pseudotype system is a useful research tool, and amenable for investigation of bat influenza tropism, restriction and sero-epidemiology, without the constraints or safety issues with producing a replication-competent virus, to which the human population is naïve.

Significance statement

Influenza virus is responsible for mortality and morbidity across the globe; human populations are constantly at risk of newly emerging strains from the aquatic bird reservoir which harbors most of the subtypes of influenza A (H1-H16). Recently identified subtypes (H17N10, H18N11) from bats have broadened the reservoir from which potential pandemic strains of influenza can emerge. To evaluate the potential for these novel subtypes to cross over into human populations, their ability to establish an infection, in addition to the extent of cross-reactive immunity established by human seasonal strains needs to be investigated. This study highlights a novel platform for the study of the bat H17 and N10 envelope glycoproteins, using a lentiviral pseudotype system. Following the generation of this pseudotype it was employed in cell entry and microneutralization assays. These showed that two well-characterised monoclonal antibodies (mAb) which target avian and human influenza subtypes will also neutralize H17. Furthermore the data presented in this study show a novel aspect of the N10 glycoprotein in its ability to facilitate the budding of pseudotypes bearing different influenza HAs.

Introduction

Influenza A virus is the principal causative agent of influenza, which is a substantial burden to global economies and represents a significant public health risk worldwide (Ma et al. 2009; Reperant et al. 2012). The first stages of the viral life cycle are mediated by the influenza glycoprotein hemagglutinin (HA), which is located in the lipid outer membrane of the virus. This protein mediates both the attachment of the influenza virus to the target cell and the fusion process which allows the virus to infect its host cell. Through its activity, HA is a key determinant of virus tropism (Dos Reis et al. 2011; Sahini et al. 2010). Furthermore, HA is a highly variable protein and its features are often used as methods for distinguishing different viral strains (Ebrahimi et al. 2014; Wilson and Cox 1990).

While the association between influenza viruses and wild birds has long been established (Gelfond et al. 2009; Webster et al. 1992), the discovery of novel influenza-like viruses in New World bats (H17N10, H18N11) represents a possible challenge to the notion that avian species are its sole reservoir (Freidl et al. 2015; Sun et al. 2013; Tong et al. 2012, 2013). Recently, Egyptian *Rousettus aegyptiacus* bats have been reported to harbor a third bat influenza subtype (Kandeil et al. 2018). It is highly likely that further subtypes will be discovered in future years, warranting further research on existing strains of bat influenza.

The H17N10 and H18N11 subtypes of influenza A were discovered in bat species in Peru and Guatemala respectively. They are significantly diverged from other known influenza strains, in respect to all eight gene segments (Tong et al. 2012, 2013). The HA of these novel strains contain a number of apparently unique structural features and exhibit receptor-binding activities that differ from other influenza viruses (Sun et al. 2013). Notably, it has been shown that H17 and H18 hemagglutinins do not bind to sialic acid receptors, and the true receptors are currently still unknown (Tong et al. 2013).

The neuraminidase (NA) of these viruses is also divergent from other known NAs, but the overall structure is preserved, despite differences in the active site (Juozapaitis et al. 2014; Ma et al. 2015). Lack of sialidase activity has been reported previously for these NAs (García-Sastre 2012; Li et al. 2012; Tong et al. 2013; Zhu et al. 2012), suggesting they utilize a different substrate altogether. The combination of atypical HA binding profiles alongside a lack of NA activity suggests that this virus

functions differently to previously discovered influenza A subtypes, despite relative phylogenetic
relatedness.

Although the receptor remains unknown, H17 and H18 sequences have been extensively analyzed
and compared to those of other HA gene sequences, to investigate the potential for these bat
viruses becoming zoonotic (Freidl et al. 2015; Mänz et al. 2013).

At present, the degree to which the H17N10 and H18N11 isolates are capable of infecting non-bat
hosts is unknown, as attempts to isolate wildtype virus have not been successful. Sequence analysis
indicates that there is significant potential for a spillover occurrence, but further research is required
to assess the true potential of these viruses as pandemic threats (Freidl et al. 2015; Juozapaitis et al.
2014).

To study the zoonotic potential of H17N10 and H18N11 viruses, attempts to integrate bat HAs into
influenza A reverse genetics systems were made but none yielded infectious virus (Juozapaitis et al.
2014; Zhou et al. 2014). More recently, progress has been achieved by Moreira et al. 2016 to this
end, using a pseudotyped VSV platform.

Influenza pseudotyped viruses (PVs) are useful tools to study both viral entry mechanisms and the
antibody response directed against the influenza HA and NA. Furthermore, when use of PVs is
coupled with detailed sequence analysis and phylogenetic inference, they offer the potential to
establish safe and effective assays to inform epidemiological and public health models of viral spread
and risk. This is especially the case with later generation, single cycle lentiviral vectors, allowing
experimentation on functional glycoproteins with the use of a reporter incorporated in the lentivirus
genome.

In this study, we report generation of the first H17 and H17N10 pseudotyped lentiviruses, their use
in virus neutralization assays using broadly neutralizing monoclonal antibodies (bnmAbs) and
investigation into the debated substrate specificity of the putative N10 neuraminidase.

Results

Generation of H17- and N10- pseudotype viruses

Generation of H17- (A/little yellow shouldered bat/060/2010) and H5- (A/Vietnam/1194/2004)
lentiviral PVs was achieved after transfection of HEK293T/17 cells with lentiviral pseudotype
production plasmids (Figure 1), followed by transduction of target human U87 MG (glioblastoma)
cells with PV supernatant. Generation of H17- PVs was only possible in the presence of HAT or
TMPRSS2 proteases in HEK293T/17 cells (Figure 2), as has been seen with other subtypes of
influenza (Bertram et al. 2010; Böttcher et al. 2006). Subsequent experiments were carried out using
canine MDCK II and RIE1495 target cells which were previously reported to allow production of VSV-
H17 (Moreira et al. 2016). In initial experiments, attempts were made to generate functional H17
and H17N10 PV utilizing different protease-expressing plasmids to promote HA maturation. PV
luciferase-based titres increased significantly when HAT-, TMPRSS2- and TMPRSS4- expression
plasmids were used in the transfection mix when compared to controls, indicating activation of the
bat HA and fusion competence. Other proteases tested (furin, KLK5, and TMPRSS3) did not yield
significant titre increases. MDCK II and RIE1495 cells were notably transduced, whereas HEK293T/17
cells were not. In the absence of co-transfected, protease-encoding plasmids, no significant PV titre
was measurable (Figure 3).

H17 entry of target cells is inhibited by broadly neutralizing monoclonal antibodies

Treatment of RIE1495 cells with the acidifying agent ammonium chloride resulted in significantly lower levels of luciferase activity, demonstrating that H17 requires a low pH for membrane fusion similar to conventional influenza viruses, such as H5 (Figure 4). Several studies have demonstrated that the HA trimers of conventional influenza viruses must simultaneously coordinate their conformational changes to complete membrane fusion (Otterstrom et al. 2014). HA-binding broadly neutralising antibodies (bnmAbs) typically recognize loop regions surrounding the receptor binding site or conserved regions of the stem and their induced-inhibition is serotype-specific (Dreyfus et al. 2013; Ekiert et al. 2009; Sui et al. 2009). In order to test whether cross reactive antibody responses would affect the H17 glycoprotein, several characterised bnmAbs were employed. Neutralization potency was measured as IC_{50} (concentration or serum dilution required to neutralise 50% of input virus). Neutralization tests were conducted using both the permissible target cell lines identified above (MDCK II and RIE1495). The first bnmAb, CR9114, binds to a conserved epitope in the HA stalk of group 1 and 2 influenza A viruses, and has also been shown to protect against lethal influenza challenge in a mouse model against both lineages of influenza B (Dreyfus et al. 2012). The second, CR6261, has been shown to neutralize H1 and H5 subtypes (Friesen et al. 2010). The third, FI6-nt, can neutralize H1 to H16 subtypes of influenza A. A fourth, FI6-chol is a derivative of F16-nt, conjugated to cholesterol (Corti et al. 2011; Lacek et al. 2014). H17 is effectively neutralized by CR9114, at a concentration of 0.05 µg/ml. FI6-nt and FI6-chol also neutralize H17N10- PV, however, concentrations required exceed those of CR9114. In contrast, CR6261 was not able to neutralize H17 using the H17 PV assay (data not shown). Neutralization data is shown in Figure 5.

N10 facilitates production of conventional H5 and H7 bearing PV.

Pretreatment of target cells with sialidase, which removes cell surface sialic acids, did not affect the entry of H17 and H17N10 PVs, which supports previous studies that H17 does not bind sialic acids for infection (Maruyama et al. 2016), see Figure 6. N10-bearing PVs were titrated alongside H17 and H17N10 PV, in order to measure their ability to transduce cells. Results showed that N10 had no effect on H17 mediated viral entry and did not improve transduction when co-expressed with H17 (data not shown). However, addition of N10 in the generation of H5 or H7 PVs (H5N10, H7N10) produced high titre PV in the absence of any other neuraminidase source (Figure 7), indicating that N10 is facilitating release of viral particles bearing sialic acid-binding glycoproteins. H5N10 luciferase titres were one log lower than parental H5 PV produced with exogenous neuraminidase, but two logs higher than N10, ΔEnvelope (ΔEnv) or cell only controls (Figure 7). Despite this, the same H5N10 PV preparation did not show any detectable NA activity when titrated using the enzyme linked lectin assay (ELLA), confirming previous reports that this protein is not a sialidase (García-Sastre 2012; Li et al. 2012; Tong et al. 2013; Zhu et al. 2012). Similarly, multiple different preparations of H17N10 or of N10 bearing PV produced negative results in ELLA (data not shown).

Discussion

We have successfully produced H17 and H17N10 bearing lentiviral pseudotypes, and shown that N10 is not required for H17 pseudotype budding, but that it can mediate release of a heterologous, sialic acid-binding HA bearing PVs. These bat influenza PVs were neutralized by cross-reactive bnmAbs, suggesting that the stalk region of the H17 glycoprotein retains conserved epitopes present in group 1 and 2 influenza HAs (Sun et al. 2013).

Of the target cell lines tested, H17 and H17N10 lentiviral PV were able to transduce U87 MG, MDCK II and RIE1495 cells to varying degrees when activated by proteolytic cleavage. Reports differ on the permissibility of the MDCK I type cells (Hoffmann et al. 2016; Moreira et al. 2016). Due to the nature of the pseudotype based microneutralisation (pMN) assay used in this study, the addition of cells in suspension into PV containing supernatant may allow infection/transduction to occur before cell

adhesion and thus by a different route than *in vivo*. This may explain our results in the context of
Moriera and colleagues' findings that the bat influenza H18 VSV pseudotypes enter at the
basolateral membrane and inefficiently infected confluent cell monolayers (Moreira et al. 2016).
MDCK I and II differ in passage number (low and high, respectively), differences between them
include the mucin-type transmembrane protein podoplanin, which is expressed only in MDCK I, and
the Forssman glycosphingolipid, which is expressed only in MDCK II cells (Hansson et al. 1986;
Zimmer et al. 1997). The susceptibility of MDCK cells to bat influenza viruses is unclear, and is partly
compounded by the availability of various lineages with different characteristics (Dukes et al. 2011).
The widespread use of MDCK cells in influenza research is at odds with a previous study that showed
H17 failed to bind to MDCK cells (Sun et al. 2013). While the reasons for this are currently unclear, it
has been hypothesized that the level of HA binding is below the threshold of detection for the assays
used in this earlier study (Maruyama et al. 2016). Alternatively, only a subset of MDCK lineages may
be susceptible to these viruses (Table 2). RIE1495 cells are morphologically similar to MDCK II cells
and express the Forssman antigen, a possible factor in their susceptibility to H17 bearing PVs
(Moreira et al. 2016). The cell tropism data reported in the Moreira study raise some interesting
ideas concerning the H17 receptor, as only three out of eight bat cell lines were transduced by the
H17-VSV pseudotypes, originating from two *Miniopterus* and one *Pteropus* species. This suggests
that these viruses may be restricted to a particular set of closely related species belonging to the
Miniopteridae and *Pteropidae* families, which are closely related to the *Phyllostomidae* family, from
which the original H17 and H18 samples were isolated (Agnarsson et al. 2011; Tong et al. 2012).
However, this raises further questions, as the cell lines derived from *Pteropus dasymallus*
yayeyamae and *Rousettus leschenaultiii* were not permissive, indicating a complex pattern of
susceptibility of species to these new influenza viruses. Nevertheless, as these cell lines have only
recently been isolated and immortalised (Maeda et al. 2008; Maruyama et al. 2014), detailed
characterisation of their surface proteins and expressed proteases are not yet available, requiring
additional research before conclusions can be drawn. Further investigation is required to determine
the mechanisms involved for the transduction of canine cells but not HEK293T/17 cells, particularly
relating to the expression of putative receptors on permissive cell lines.

Our results, contrary to previous studies, indicate that N10 is performing a similar function to other
NAs in enabling release of new influenza virus particles, which indicate its ability to form VLPs rather
than releasing H17 from its cellular receptor (Yondola et al. 2011). PVs bearing H5/H7 and N10
envelope glycoproteins successfully budded from producer cells into the surrounding medium in the
absence of a sialic acid cleaving neuraminidase, resulting in significantly increased titres when
compared to the same glycoproteins generated without the addition of N10 (Figure 7). This increase
in budding PV may be due to the action of N10 on an unknown substrate, or perhaps action of the
protein itself in virus morphogenesis and budding (Barman et al. 2004; Yondola et al. 2011).
Sialidase activity was however not detected using the enzyme-linked lectin assay, suggesting either a
lack of sensitivity of this assay or another mechanism for the removal of surface sialic acids by N10
(Juozapaitis et al. 2014; Sun et al. 2013; Wu et al. 2014). Similarly, PVs bearing solely the N10
glycoprotein did not show sialidase activity via ELLA, or successfully transduce cells to give a
significant luciferase reading. Further investigation is required on the role of the bat influenza
neuraminidase in its lifecycle, the combination of the findings described in this article, with the fact
that the N10 enzymatic structure remains conserved and NA-like, suggests that it is involved in
accessory functions other than simply aiding in viral egress. Our results highlight the distinct
difference between bat and traditional influenza A viruses where a delicate balance is in place
between HA and NA. It may be the case that such a balance exists between bat HA and NAs which
will be made clear once the substrate(s) of the bat NA is discovered.

In a previous study, it was demonstrated that the TMPRSS2 protease was capable of inducing HA
maturation of H17 through cleavage from HA0 to HA1 and HA2 (Hoffmann et al. 2016). In our study,
we demonstrated that this maturation can also be facilitated by TMPRSS4 and HAT. This is of
particular interest as the expression of specific proteases is a known limiting factor in viral tropism
for a number of different viruses (Böttcher-Friebertshäuser et al. 2010; Ferrara et al. 2012; Millet
and Whittaker 2015). This, coupled with the observed susceptibility of a canine-derived cell line to
the H17 PV, will need to be factored into future analysis of the potential for zoonotic spillover from
bat origin influenza viruses.

The fact that bnmAbs were able to neutralize the virus particles via the H17 hemagglutinin, and their
requirement for proteolytic activation, reinforces that we have only scratched the surface in terms
of their characterization. Further exploration is required to establish whether these viruses are
endogenous or capable of forming infectious particles *in vivo*. Research has revealed that all eight
segments of the genome encode functional proteins (Juozapaitis et al. 2014; Moreira et al. 2016; Wu
et al. 2014; Zhu et al. 2012), despite the inability to isolate wildtype virus to date. Nevertheless, the
development of tools, such as those described here, which can be used in H17 receptor
identification studies may ultimately aid discovery of wildtype virus samples from the bat reservoir.

Methods

Plasmids:

The H17 HA and N10 NA genes from A/little yellow shouldered bat/Guatamala/060/2010 were
synthesised commercially by Genscript (Genscript, USA) and subcloned into vector pI.18 (Cox et al.
2002). Lentiviral packaging plasmids p8.91 (Zufferey et al. 1997) and pCSFLW (Demaison et al. 2002)
containing a firefly luciferase reporter were used to produce PV. Protease encoding plasmids
phCMV-Tag3 (TMPRSS4-myc) and pcDNA3.1-hTMPRSS3 were kindly provided by Prof. Stefan
Pöhlmann, Infection Biology Unit, German Primate Center, Germany. pCAGGS-TMPRSS2 and
pCAGGS-HAT were kindly provided by Eva Böttcher-Friebertshäuser, Philipps University of Marburg,
Germany. The plasmid bearing the Vesicular stomatitis virus envelope protein (VSV-G), pMD.G was
obtained from Dr Yasu Takeuchi, University College London, United Kingdom. All the plasmids that
were used in this study are listed in Table 3.

Cell lines:

Multiple cell lines were used for titration of PV containing supernatants. HEK293T/17 cells were
kindly provided by Dr Edward Wright (University of Westminster, UK). MDCK II and RIE1495 were
kindly provided by Dr. Gert Zimmer (Institute of Virology and Immunology, Switzerland). U-87 cells
were provided by Dr Simon Scott (University of Kent, UK). Madin-Darby Canine Kidney (MDCK) cells
were kindly provided by Prof. Sarah Gilbert (Jenner Institute, University of Oxford, UK). All cells were
cultured in Dulbecco's modified eagle medium supplemented with 10% fetal bovine serum.

Antibodies:

251 mAbs CR9114 and CR6261 were produced by Crucell (Janssen Vaccines AG, Bern, Switzerland). FI6-nt
252 and cholesterol conjugated FI6 (FI6-Chol) were produced by Alfredo Pesci and Krzysztof Lacek from
253 sequence information derived from Corti et al. 2011.

Production and quantification of H17N10 and H17 bearing lentiviral pseudotypes:

PV were produced as described previously (Ferrara et al. 2012; Temperton et al. 2007) and as shown
in Figure 1. Briefly, transfection of HEK293T/17 cells was performed using a variety of combinations

of plasmids pI.18-H17, pI.18-N10, p8.91 and pCSFLW using polyethylenimine transfection reagent
(Sigma Aldrich, UK). Protease-encoding expression plasmids were also included (Table 3). Medium
was replaced 12h post-transfection. Supernatants were harvested 48h post-transfection and passed
through a 0.45μm filter (Millipore, UK). PV-containing supernatants were titrated using the firefly
luciferase Bright-Glo™ system (Promega, UK). Serial dilution of 100μl of PV-containing supernatant
was performed across a white flat bottom 96-well Nunclon© plate (Thermo Fisher Scientific, UK).
Subsequently, approximately $1x10^4$ cells per well were added per well in 50μl of medium, plates
were incubated in a humidified incubator at 37°C 5% CO_2 for 48h, after which 50μl of Bright-Glo™
substrate was added and luciferase reading recorded in relative luminescence units (RLU) after a 5
minute incubation period. Further sets of N10-bearing PV were produced by transfection of
HEK293T/17 cells with 500ng p8.91, 750ng pCSFLW and various amounts of pI.18-N10 plasmid.
Medium was replenished after 12h; PV containing supernatants were collected 72h later and passed
through a 0.45μm filter. Transfections were carried out in 6-well Nunclon© plates (Thermo Fisher
Scientific, UK).

GFP-expressing pseudotypes were produced by substituting the pCSFLW firefly luciferase lentiviral
vector mentioned previously with the GFP expressing vector pCSGW. PV-containing supernatants
were titrated down clear 96-well Nunclon© plates (Thermo Fisher Scientific, UK) with the addition of
$1x10^4$ cells per well. Plates were incubated for 72h and visualized by Nikon Eclipse 50i
epiflourescence microscope with a charge-coupled QICAM Fast 1394 camera (QImaging) at x200
magnification.

Screening of cell lines

Two-fold serial dilutions of PV-containing supernatant were performed as previously described using
white 96-well Nunclon© plates (Thermo Fisher Scientific, UK). Subsequently, approximately $1x10^4$ of
each cell line was added in 50μl of medium per well. Plates were incubated in a humidified incubator
at 37°C and 5% CO_2 for 48h, after which 50μl of Bright-Glo™ substrate was added and luciferase
reading recorded in relative luminescence units (RLU), following a 5 minute incubation period.
Control wells were used to measure cell populations for each cell line.

Pseudotype based microneutralization assay (pMN) using H17N10 and H17 bearing lentiviral
pseudotypes:

Monoclonal antibodies (mAbs) were serially diluted 1:2 across white 96-well Nunclon© plates
(Thermo Fisher Scientific, UK) in 50 μl of DMEM. PV-containing supernatants were diluted and added
to each well to give an approximate RLU value of $1x10^6$ per well in 50 μl of DMEM. After 1h
incubation at 37°C and 5% CO_2, approximately $1x10^4$ cells were added per well in a volume of 50μl.
Plates were incubated for 48h at 37°C and 5% CO_2, then 50 μl of Bright-Glo™ was added and
luminescence read after a 5 minute incubation at room temperature. Results were analyzed with
Graphpad (Prism 7), using nonlinear regression on luminescence values normalized to cell only and
virus only control thresholds (100% and 0% neutralization equivalent, respectively). IC_{50} values
represent the concentration (ng/ml) required for each antibody to neutralize 50% of functional
pseudotyped virus, based on luciferase activity.

Cell treatment

Overnight treatment of RIE1495 cells with the endosomal acidification inhibitor
ammonium chloride (SIGMA) and pretreatment for 2h with neuraminidase (Roche) was done as
previously described (Maruyama et.al, 2016). Treated cells were washed three times with serum
free medium, and then incubated for 24h with H17 PV. PV titration and luciferase activity was
monitored with a luminometer as described previously.

Pseudotype based enzyme linked lectin assay (pELLA) utilizing N10:

ELLA was performed as described by Couzens et al. 2014, but adapted in order to allow use of
lentiviral pseudotypes as a source of NA (Biuso et al. 2018; Prevato et al. 2015). Briefly, PV
containing supernatant was serially diluted (1:2) across a standard clear microtitre plate in sample
diluent (PBS, 1% BSA, 0.5% Tween 20). 50 µl of the resulting dilution series was transferred in
duplicate to Nunclon© Maxisorp 96-well plates (Thermo Fisher Scientific, UK) previously coated with
Fetuin (Sigma Aldrich, UK) and containing 50 µl of sample diluent per well. Plates were then
incubated for 18h and washed, prior to addition of conjugate diluent (PBS, 1% BSA) containing a
1:500 concentration of peanut lectin conjugated to horseradish peroxidase (PNA-HRPO, Sigma
Aldrich, UK). Plates were then incubated for 2h in the dark, whereupon OPD-based substrate in
citrate buffer was added (Sigma Aldrich, UK). Reactions were stopped after 10 minutes using 50 µl
1M H_2SO_4, and readings recorded using a standard ELISA plate reader at 492nm. Exogenous
neuraminidase from *Clostridium perfringens* (Sigma Aldrich, UK) was used as a positive benchmark
control and samples were assayed alongside PV bearing neuraminidases from other influenza
subtypes.

Statistical analysis

Where possible, statistical analysis was carried out to determine whether differences in PV titre
were significant. One-way ANOVA t-tests were performed using fold change scores with a Tukey's
multiple comparisons test. P-values were set at 0.05 ($P≤0.05$) unless indicated otherwise.
Significantly different data are denoted with asteriks representing $p= <0.01$ (*), <0.001 (**), <0.0001
(***) and <0.00001 (****).

Figures and Tables

Figure 1. Cartoon showing the production of H17N10 PV via plasmid transfection.

Figure 2. Transduction of U87 MG cells by influenza GFP PV (x100 magnification). A) H17 produced
using pCAGGS-HAT. B) H17 PV produced using pCAGGS-TMPRSS2. C) Cell only control. D) H5 (A/Viet
nam/1194/2004) PV positive control.

Figure 3. Transduction of various target cell lines. A) HEK293T/17, B) RIE1495 and C) MDCK II with
H17 and H17N10 pseudotyped viruses carrying the luciferase reporter gene. Results given in Relative
Luminescence Units per ml; RLU/ml). Average cell only luminescence shown as a red line.

346 Figure 4. The effect of ammonium chloride based acidification on the transduction of RIE1495 cells
347 by H17 PV. Increases in acidification of RIE1495 cell cultures prior to transduction by H17 resulted in
348 lower luciferase based titres, indicating that the pH sensitivity of H17 is similar to that of H5.
349 Significant differences in transduction denoted by asteriks, ** represents p= <0.001 and *** p=
350 <0.0001 respectively.

IC50 (µg/ml)	MDCK II	RIE1495
CR9114	0.06	0.05
F16 nt	1.20	59.00
F16 chol	290.00	92.00

354 Figure 5. Neutralization curves and IC50 values for mAbs CR9114, FI6-nt and FI6-Chol against
355 H17N10 bearing lentiviral pseudotypes on cell lines RIE1495 and MDCK II. Nonlinear regression
356 carried out using Graphpad (Prism 7) in order to provide IC_{50} values for each graph. Each IC_{50} is the
357 concentration of mAb required for reduction of 50% of the virus input in terms of luciferase activity.

364 Figure 6. The effect of neuraminidase pre-treatment on the transduction of RIE1495 cells by H17 and
365 H5 PV. No relationship is seen between pre-treatment of cells with neuraminidase for H17 bearing
366 PV, but H5-based transduction is significantly reduced when sialic acids are stripped from target
367 cells. Significant difference in transduction denoted by asteriks, * represents p= <0.01 and ** p=
368 <0.001 respectively.

Figure 7. Release of H5 and H7 bearing PV by N10. PV bearing the glycoproteins H5 (A/Vietnam/1194/2004) and H7 (A/Shanghai/1/2013) were produced with exogenous neuraminidase (+NA), with no NA (-NA) or accompanied with an N10 glycoprotein (+N10). H5 and H7 PVs produced in the presence of N10 have significantly higher luficerase titres than those produced with no NA. Background luciferase titre control (Δ Env) control shown. Significant differences in transduction denoted by asteriks, **** represents p= <0.00001.

380   Table 1. Cell lines used in the production of Bat influenza VSV pseudotypes.

Cell Line	Species	Tissue:
Vero E6	*Chlorocebus sp.*	Kidney
HEK293	*Homo sapiens*	Kidney
MDCK	***Canis lupus familiaris***	**Kidney**
SK-L	*Sus scrofa*	Kidney
QT6	*Coturnix japonica*	Muscle
Cell Line:	**Bat species:**	**Tissue:**
BKT1	*Rhinolophus ferrumequinum*	Kidney
FBKT1	*Pteropus dasymallus yayeyamae*	Kidney
YubFKT1	***Miniopterus fuliginosus***	**Kidney**
IndFSPT1	***Pteropus giganteus***	**Spleen**
DemKT1	*Rousettus leschenaultii*	Kidney
ZFBK11-97	*Epomophorus gambianus*	Kidney
SuBK12-08	***Miniopterus schreibersii***	**Kidney**
ZFBS13-75A	*Eidolon helvum*	Spleen

List of cell lines used in VSV based pseudotyping of H17 and H18 viruses. Bold entries are those cell lines which were found to be permissive for viral transduction by Maruyama et al. 2016.

382   Table 2: MDCK cell lines.

NBL-2 (ATCC® cat # CCL-34™)
MDCK I (EEACC cat # 00062106)
MDCK II (EEACC cat# 00062107)
MDCK.1 (ATCC® cat# CRL-2935™)
MDCK.2 (ATCC® cat# CRL- CRL-2936™)
super dome (ATCC® cat# CRL-2286™)
super tube (ATCC® cat# CRL-2285™)
Different MDCK cell lines available commercially

391   Table 3: Genes, plasmids and sources.

Gene	Plasmid	Source
Transmembrane protease, serine 2	pCAGGS-TMPRSS2	(Böttcher et al. 2006)
Transmembrane protease, serine 3	pcDNA3.1-hTMPRSS3	(Bertram et al. 2010)
Transmembrane protease, serine 4	phCMV-Tag3 (TMPRSS4-myc)	(Jung et al. 2008)
Human Airway Trypsin	pCAGGS-HAT	(Böttcher et al. 2006)
Kallikrein-related peptidase 5	pI.18-KLK5	Synthesised, Sigma
Furin	pFurin	Kind gift from H. D. Klenk
H17 (A/little yellow shouldered bat/Guatamala/060/2010)	pI.18-H17	Synthesised, Genscript
N10 (A/little yellow shouldered bat/Guatamala/060/2010)	pI.18-N10	Synthesised, Genscript
H5 (A/Viet Nam/1194/2004)	pI.18-H5	(Temperton et al. 2007)
HIV-1 packaging plasmid	pCMVΔR8.91 (p8.91)	(Zufferey et al. 1997)
Lentiviral vector for expression of Firefly Luciferase	pHR-SIN-SE (pCSFLW)	(Demaison et al. 2002)
Vesicular Stomatitis Virus glycoprotein	pMD.G (VSV-G)	Yasu Takeuchi, University College London

393 **Acknowledgements**

394 EG is supported by the Biotechnology and Biological Sciences Research Council
395 (http://www.bbsrc.ac.uk) via Strategic LoLa grant BB/K002465/1 "Developing Rapid Responses to
396 Emerging Virus Infections of Poultry (DRREVIP)" and MS by the Deutsche Forschungsgemeinschaft
397 (DFG) (SCHW 632/17-1).
398

References

Agnarsson, I., Zambrana-Torrelio, C. M., Flores-Saldana, N. P., & May-Collado, L. J. (2011). A time-
calibrated species-level phylogeny of bats (chiroptera, Mammalia). *PLoS Currents*, *3*, RRN1212.
doi:10.1371/currents.RRN1212

Barman, S., Adhikary, L., Chakrabarti, A. K., Bernas, C., Kawaoka, Y., Nayak, D. P., et al. (2004). Role of
transmembrane domain and cytoplasmic tail amino acid sequences of influenza a virus
neuraminidase in raft association and virus budding. *Journal of virology*, *78*(10), 5258–5269.
doi:10.1128/JVI.78.10.5258

Bertram, S., Glowacka, I., Blazejewska, P., Soilleux, E. J., Allen, P., Danisch, S., et al. (2010). TMPRSS2
and TMPRSS4 facilitate trypsin-independent spread of influenza virus in Caco-2 cells. *J Virol*,
84(19), 10016–10025. doi:10.1128/JVI.00239-10

Biuso, F., Carnell, G. W., Montomoli, E., & Temperton, N. (2018). A Lentiviral Pseudotype ELLA for
the Measurement of Antibodies Against Influenza Neuraminidase. *Bio-Protocol*, *8*(14), 1–15.
doi:10.21769/BioProtoc.2936

Böttcher-Friebertshäuser, E., Freuer, C., Sielaff, F., Schmidt, S., Eickmann, M., Uhlendorff, J., et al.
(2010). Cleavage of influenza virus hemagglutinin by airway proteases TMPRSS2 and HAT
differs in subcellular localization and susceptibility to protease inhibitors. *Journal of virology*,
84(11), 5605–14. doi:10.1128/JVI.00140-10

Böttcher, E., Matrosovich, T. Y., Beyerle, M., Klenk, H.-D., Garten, W., & Matrosovich, M. N. (2006).
Proteolytic activation of influenza viruses by serine proteases TMPRSS2 and HAT from human
airway epithelium. *Journal of virology*, *80*(19), 9896–8. doi:10.1128/JVI.01118-06

Corti, D., Voss, J., Gamblin, S. J., Codoni, G., Macagno, A., Jarrossay, D., et al. (2011). A Neutralizing
Antibody Selected from Plasma Cells That Binds to Group 1 and Group 2 Influenza A
Hemagglutinins. *Science*, *333*(6044), 850–6. doi:10.1126/science.1205669

Couzens, L., Gao, J., Westgeest, K. B., Sandbulte, M. R., Lugovtsev, V. Y., Fouchier, R. A. M., &
Eichelberger, M. C. (2014). An optimized enzyme-linked lectin assay to measure influenza A
virus neuraminidase inhibition antibody titers in human sera. *Journal of Virological Methods*,
210, 7–14. doi:10.1016/j.jviromet.2014.09.003

Cox, R. J., Mykkeltvedt, E., Robertson, J. S., & Haaheim, L. R. (2002). Non-lethal viral challenge of
influenza haemagglutinin and nucleoprotein DNA vaccinated mice results in reduced viral
replication. *Scandinavian Journal of Immunology*, *55*(1), 14–23. doi:10.1046/j.1365-
3083.2002.01015.x

Demaison, C., Parsley, K., Brouns, G., Scherr, M., Battmer, K., Kinnon, C., et al. (2002). High-level
transduction and gene expression in hematopoietic repopulating cells using a human
immunodeficiency [correction of imunodeficiency] virus type 1-based lentiviral vector
containing an internal spleen focus forming virus promoter. *Human gene therapy*, *13*(7), 803–
813. doi:10.1089/10430340252898984

Dos Reis, M., Tamuri, A. U., Hay, A. J., & Goldstein, R. A. (2011). Charting the host adaptation of
influenza viruses. *Molecular Biology and Evolution*, *28*(6), 1755–1767.
doi:10.1093/molbev/msq317

Dreyfus, C., Ekiert, D. C., & Wilson, I. A. (2013). Structure of a Classical Broadly Neutralizing Stem
Antibody in Complex with a Pandemic H2 Influenza Virus Hemagglutinin. *Journal of Virology*,
87(12), 7149–7154. doi:10.1128/JVI.02975-12

Dreyfus, C., Laursen, N. S., Kwaks, T., Zuijdgeest, D., Khayat, R., Ekiert, D. C., et al. (2012). Highly
Conserved Protective Epitopes on Influenza B Viruses. *Science*, *337*(6100), 1343–1348.
doi:10.1126/science.1222908

Dukes, J. D., Whitley, P., Chalmers, A. D., Gaush, C., Hard, W., Smith, T., et al. (2011). The MDCK
variety pack: choosing the right strain. *BMC Cell Biology*, *12*(1), 43. doi:10.1186/1471-2121-12-
43

Ebrahimi, M., Aghagolzadeh, P., Shamabadi, N., Tahmasebi, A., Alsharifi, M., Adelson, D. L., et al.
(2014). Understanding the undelaying mechanism of HASubtyping in the level of physic-
chemical characteristics of protein. *PLoS ONE*, *9*(5), 1–14. doi:10.1371/journal.pone.0096984

Ekiert, D. C., Bhabha, G., Elsliger, M., Friesen, R. H. E., Jongeneelen, M., Throsby, M., et al. (2009).
Antibody recognition of a highly conserved influenza virus epitope◌: implications for universal
prevention and therapy. *Science*, *324*(5924), 246–251. doi:10.1126/science.1171491.Antibody

Ferrara, F., Molesti, E., Böttcher-Friebertshäuser, E., Cattoli, G., Corti, D., Scott, S. D., & Temperton,
N. J. (2012). The human Transmembrane Protease Serine 2 is necessary for the production of
Group 2 influenza A virus pseudotypes. *Journal of molecular and genetic medicine◌: an
international journal of biomedical research*, *7*, 309–14.

Freidl, G. S., Binger, T., Müller, M. A., De Bruin, E., Van Beek, J., Corman, V. M., et al. (2015).
Serological evidence of influenza a viruses in frugivorous bats from Africa. *PLoS ONE*, *10*(5), 1–
7. doi:10.1371/journal.pone.0127035

Friesen, R. H. E., Koudstaal, W., Koldijk, M. H., Weverling, G. J., Brakenhoff, J. P. J., Lenting, P. J., et al.
(2010). New class of monoclonal antibodies against severe influenza: Prophylactic and
therapeutic efficacy in ferrets. *PLoS ONE*, *5*(2), 1–7. doi:10.1371/journal.pone.0009106

García-Sastre, A. (2012). The neuraminidase of bat influenza viruses is not a neuraminidase. *Proc
Natl Acad Sci U S A*, *109*(46), 18635–18636. doi:10.1073/pnas.1215857109

Gelfond, J. A. L., Gupta, M., & Ibrahim, J. G. (2009). A bayesian hidden markov model for motif
discovery through joint modeling of genomic sequence and ChIP-chip data. *Biometrics*, *65*(4),
1087–1095. doi:10.1111/j.1541-0420.2008.01180.x

Hansson, G. C., Simons, K., & Meer, G. Van. (1986). Two strains of the Madin- Darby canine kidney (
NDCK) cell line have distinct glycosphingolipid compositions. *Proceedings of the National
Academy of Sciences*, *5*(3), 483–489.

Hoffmann, M., Krüger, N., Zmora, P., Wrensch, F., Herrler, G., & Pöhlmann, S. (2016). The
hemagglutinin of bat-associated influenza viruses is activated by TMPRSS2 for pH-dependent
entry into bat but not human cells. *PLoS ONE*, *11*(3), 1–18. doi:10.1371/journal.pone.0152134

Jung, H., Lee, K. P., Park, S. J., Park, J. H., Jang, Y. S., Choi, S. Y., et al. (2008). TMPRSS4 promotes
invasion, migration and metastasis of human tumor cells by facilitating an epithelial-
mesenchymal transition. *Oncogene*, *27*(18), 2635–2647. doi:10.1038/sj.onc.1210914

Juozapaitis, M., Aguiar Moreira, E., Mena, I., Giese, S., Riegger, D., Pohlmann, A., et al. (2014). An
infectious bat-derived chimeric influenza virus harbouring the entry machinery of an influenza
A virus. *Nat Commun*, *5*, 4448. doi:10.1038/ncomms5448

Kandeil, A., Gomaa, M. R., Shehata, M. M., El Taweel, A. N., Mahmoud, S. H., Bagato, O., et al.
(2018). Isolation and characterization of a distinct influenza A virus from Egyptian bats. *Journal
of virology*, JVI.01059-18. doi:10.1128/JVI.01059-18

Lacek, K., Urbanowicz, R. A., Troise, F., De Lorenzo, C., Severino, V., Di Maro, A., et al. (2014).

Dramatic potentiation of the antiviral activity of HIV antibodies by cholesterol conjugation.
Journal of Biological Chemistry, *289*(50), 35015–35028. doi:10.1074/jbc.M114.591826

Li, Q., Sun, X., Li, Z., Liu, Y., Vavricka, C. J., Qi, J., & Gao, G. F. (2012). Structural and functional
characterization of neuraminidase-like molecule N10 derived from bat influenza A virus.
Proceedings of the National Academy of Sciences of the United States of America, *109*(46),
18897–18902. doi:10.1073/pnas.1211037109

Ma, W., García-Sastre, A., & Schwemmle, M. (2015). Expected and Unexpected Features of the
Newly Discovered Bat Influenza A-like Viruses. *PLoS Pathog*, *11*(6), e1004819.
doi:10.1371/journal.ppat.1004819

Ma, W., Lager, K. M., Vincent, A. L., Janke, B. H., Gramer, M. R., & Richt, J. A. (2009). The role of
swine in the generation of novel influenza viruses. *Zoonoses and Public Health*, *56*(6–7), 326–
337. doi:10.1111/j.1863-2378.2008.01217.x

Maeda, K., Hondo, E., Terakawa, J., Kiso, Y., Nakaichi, N., Endoh, D., et al. (2008). Isolation of novel
adenovirus from fruit bat (Pteropus dasymallus yayeyamae) [7]. *Emerging Infectious Diseases*,
14(2), 347–349. doi:10.3201/eid1402.070932

Mänz, B., Schwemmle, M., & Brunotte, L. (2013). Adaptation of avian influenza A virus polymerase in
mammals to overcome the host species barrier. *Journal of virology*, *87*(April), 7200–9.
doi:10.1128/JVI.00980-13

Maruyama, J., Miyamoto, H., Kajihara, M., Ogawa, H., Maeda, K., Sakoda, Y., et al. (2014).
Characterization of the Envelope Glycoprotein of a Novel Filovirus, Lloviu Virus. *Journal of*
Virology, *88*(1), 99–109. doi:10.1128/JVI.02265-13

Maruyama, J., Nao, N., Miyamoto, H., Maeda, K., Ogawa, H., Yoshida, R., et al. (2016).
Characterization of the glycoproteins of bat-derived influenza viruses. *Virology*, *488*, 43–50.
doi:10.1016/j.virol.2015.11.002

Millet, J. K., & Whittaker, G. R. (2015). Host cell proteases: Critical determinants of coronavirus
tropism and pathogenesis. *Virus research*, *202*, 120–34. doi:10.1016/j.virusres.2014.11.021

Moreira, É. A., Locher, S., Kolesnikova, L., Bolte, H., Aydillo, T., García-Sastre, A., et al. (2016).
Synthetically derived bat influenza A-like viruses reveal a cell type- but not species-specific
tropism. *Proceedings of the National Academy of Sciences*, 201608821.
doi:10.1073/pnas.1608821113

Otterstrom, J. J., Brandenburg, B., Koldijk, M. H., Juraszek, J., Tang, C., Mashaghi, S., et al. (2014).
Relating influenza virus membrane fusion kinetics to stoichiometry of neutralizing antibodies at
the single-particle level. *Proceedings of the National Academy of Sciences*, *111*(48), E5143–
E5148. doi:10.1073/pnas.1411755111

Prevato, M., Cozzi, R., Pezzicoli, A., Taddei, A. R., Ferlenghi, I., Nandi, A., et al. (2015). An innovative
pseudotypes-based Enzyme-Linked Lectin Assay for the measurement of functional anti-
neuraminidase antibodies. *PLoS ONE*, *10*(8), e0135383. doi:10.1371/journal.pone.0135383

Reperant, L. A., Kuiken, T., & Osterhaus, A. D. (2012). Adaptive pathways of zoonotic influenza
viruses: From exposure to establishment in humans. *Vaccine*, *30*(30), 4419–4434.
doi:10.1016/j.vaccine.2012.04.049

Sahini, L., Tempczyk-Russell, A., & Agarwal, R. (2010). Large-scale sequence analysis of hemagglutinin
of influenza A virus identifies conserved regions suitable for targeting an anti-viral response.
PLoS ONE, *5*(2), 1–9. doi:10.1371/journal.pone.0009268

Sui, J., Hwang, W. C., Perez, S., Wei, G., Aird, D., Chen, L., et al. (2009). Structural and functional
bases for broad-spectrum neutralization of avian and human influenza A viruses. *Nature*
structural & molecular biology, *16*(3), 265–273. doi:10.1038/nsmb.1566

Sun, X., Shi, Y., Lu, X., He, J., Gao, F., Yan, J., et al. (2013). Bat-Derived Influenza Hemagglutinin H17
Does Not Bind Canonical Avian or Human Receptors and Most Likely Uses a Unique Entry
Mechanism. *Cell Reports*, *3*(3), 769–778. doi:10.1016/j.celrep.2013.01.025

Temperton, N. J., Hoschler, K., Major, D., Nicolson, C., Manvell, R., Hien, V. M., et al. (2007). A
sensitive retroviral pseudotype assay for influenza H5N1-neutralizing antibodies. *Influenza and*
other respiratory viruses, *1*(3), 105–112. doi:10.1111/j.1750-2659.2007.00016.x

Tong, S., Li, Y., Rivailler, P., Conrardy, C., Castillo, D. A. A., Chen, L.-M., et al. (2012). A distinct lineage
of influenza A virus from bats. *Proceedings of the National Academy of Sciences of the United*
States of America, *109*(11), 4269–4274. doi:10.1073/pnas.1116200109

Tong, S., Zhu, X., Li, Y., Shi, M., Zhang, J., Bourgeois, M., et al. (2013). New World Bats Harbor Diverse
Influenza A Viruses. *PLoS Pathogens*, *9*(10). doi:10.1371/journal.ppat.1003657

Webster, R. G., Bean, W. J., Gorman, O. T., Chambers, T. M., & Kawaoka, Y. (1992). *Evolution and*
Ecology of Influenza A Viruses. *MICROBIOLOGICAL REVIEWS* (Vol. 56).
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC372859/pdf/microrev00028-0170.pdf.
Accessed 27 August 2018

Wilson, I. a, & Cox, N. J. (1990). Structural basis of immune recognition of influenza virus
hemagglutinin. *Annual review of immunology*, *8*(1), 737–771.
doi:10.1146/annurev.iy.08.040190.003513

Wu, Y., Wu, Y., Tefsen, B., Shi, Y., & Gao, G. F. (2014). Bat-derived influenza-like viruses H17N10 and
H18N11. *Trends in Microbiology*, *22*(4), 183–191. doi:10.1016/j.tim.2014.01.010

Yondola, M. A., Fernandes, F., Belicha-Villanueva, A., Uccelini, M., Gao, Q., Carter, C., & Palese, P.
(2011). Budding capability of the influenza virus neuraminidase can be modulated by tetherin.
Journal of virology, *85*(6), 2480–91. doi:10.1128/JVI.02188-10

Zhou, B., Ma, J., Liu, Q., Bawa, B., Wang, W., Shabman, R. S., et al. (2014). Characterization of
Uncultivable Bat Influenza Virus Using a Replicative Synthetic Virus. *PLoS Pathogens*, *10*(10),
e1004420. doi:10.1371/journal.ppat.1004420

Zhu, X., Yang, H., Guo, Z., Yu, W., Carney, P. J., Li, Y., et al. (2012). Crystal structures of two subtype
N10 neuraminidase-like proteins from bat influenza A viruses reveal a diverged putative active
site. *Proceedings of the National Academy of Sciences of the United States of America*, *109*(46),
18903–18908. doi:10.1073/pnas.1212579109

Zimmer, G., Lottspeich, F., Maisner, A., Klenk, H. D., & Herrler, G. (1997). Molecular characterization
of gp40, a mucin-type glycoprotein from the apical plasma membrane of Madin-Darby canine
kidney cells (type I). *The Biochemical journal*, *326*, 99–108. doi:10.1042/bj3260099

Zufferey, R., Nagy, D., Mandel, R. J., Naldini, L., & Trono, D. (1997). Multiply attenuated lentiviral
vector achieves efficient gene delivery in vivo. *Nat Biotechnol*, *15*(9), 871–875.
doi:10.1038/nbt0997-871