Pandoravirus celtis illustrates the microevolution processes at work in the giant Pandoraviridae genomes

Matthieu Legendre[1], Jean-Marie Alempic[1], Nadège Philippe[1], Audrey Lartigue[1], Sandra Jeudy[1], Olivier Poirot[1], Ngan Thi Ta[1], Sébastien Nin[1], Yohann Couté[2], Chantal Abergel[1*], Jean-Michel Claverie[1*]

[1]Aix Marseille Univ, CNRS, IGS, Structural and Genomic Information Laboratory (UMR7256), Mediterranean Institute of Microbiology (FR3479), 163 Avenue de Luminy, F-13288 Marseille, France

[2]Univ. Grenoble Alpes, CEA, Inserm, BIG-BGE, 38000 Grenoble, France.

* Correspondence:
Jean-Michel Claverie
jean-michel.claverie@univ-amu.fr
Chantal Abergel
chantal.abergel@igs.cnrs-mrs.fr

Keywords: de novo gene creation, comparative genomics, Acanthamoeba, Giant viruses, Soil viruses, hAT transposase

Abstract

With genomes of up to 2.7 Mb propagated in µm-long oblong particles and initially predicted to encode more than 2000 proteins, members of the Pandoraviridae family display the most extreme features of the known viral world. The mere existence of such giant viruses raises fundamental questions about their origin and the processes governing their evolution. A previous analysis of six newly available isolates, independently confirmed by a study including 3 others, established that the Pandoraviridae pan-genome is open, meaning that each new strain exhibits protein-coding genes not previously identified in other family members. With an average increment of about 60 proteins, the gene repertoire shows no sign of reaching a limit and remains largely coding for proteins without recognizable homologs in other viruses or cells (ORFans). To explain these results, we proposed that most new protein-coding genes were created *de novo*, from pre-existing non-coding regions of the G+C rich pandoravirus genomes. The comparison of the gene content of a new isolate, *P. celtis*, closely related (96 % identical genome) to the previously described *P. quercus* is now used to test this hypothesis by studying genomic changes in a microevolution range. Our results confirm that the differences between these two similar gene contents mostly consist of protein-coding genes without known homologs (ORFans), with statistical signatures close to that of intergenic regions. These newborn proteins are under slight negative selection, perhaps to maintain stable folds and prevent protein aggregation pending the eventual emergence of fitness-increasing functions. Our study also unraveled several insertion events mediated by a transposase of the hAT family, 3 copies of which are found in *P. celtis* and are presumably active. Members of the Pandoraviridae are presently the first viruses known to encode this type of transposase.

1 Introduction

The Pandoraviridae is a proposed family of giant dsDNA viruses - not yet registered by the
International Committee on Taxonomy of Viruses (ICTV) - multiplying in various species of
Acanthamoeba through a lytic infectious cycle. Their linear genomes, flanked by large terminal
repeats, range from 1.9 to 2.7 Mb in size, and are propagated in elongated oblong particles
approximately 1.2 µm long and 0.6 µm in diameter (Fig. S1). The prototype strain (and the one
with the largest known genome) is *Pandoravirus salinus*, isolated from shallow marine sediments
off the coast of central Chile (Philippe et al., 2013). Other members were soon after isolated from
worldwide locations. Complete genome sequences have been determined for *P. dulcis*
(Melbourne, Australia) (Philippe et al., 2013), P. *inopinatum* (Germany) (Antwerpen et al.,
2015), *P. macleodensis* (Australia), *P. neocaledonia* (New Caledonia), and *P. quercus* (France)
(Legendre et al., 2018), and three isolates from Brazil (*P. braziliensis*, *P. pampulha*, and *P.*
massiliensis) (Aherfi et al., 2018). A standard phylogenetic analysis of the above strains
suggested that the Pandoraviridae family consists of two separate clades (Claverie et al., 2018;
Fig. 1). The average proportion of identical amino acids between Pandoraviridae orthologs within
each clade is above 70% while it is below 55% between members of the A and B clades.
Following a stringent reannotation of the predicted protein-coding genes using transcriptomic and
proteomic data, our comparative genomic analysis reached the main following conclusions
(Legendre et al., 2018):

1) the uniquely large proportion of predicted proteins without homologs outside of the
Pandoraviridae (ORFans) is real and not due to bioinformatic errors induced by the
above-average G+C content (>60%) of Pandoravirus genomes;
2) the Pandoraviridae pan genome appears "open" (i.e. unbounded);
3) as most of the genes are unique to each strain are ORFans, they were not horizontally
acquired from other (known) organisms;
4) they are neither predominantly the result of gene duplications.

The scenario of *de novo* and *in situ* gene creation, supported by the analysis of their sequence
statistical signatures, thus became our preferred explanation for the origin of strain specific genes.

In the present study, we take advantage of the high similarity (96.7% DNA sequence identity)
between *P. quercus* and a newly characterized isolate, *P. celtis*, to investigate the microevolution
processes initiating the divergence between Pandoraviruses. Our results further support *de novo*
gene creation as a main diversifying force of the Pandoraviridae family.

2 Materials and Methods

2.1 Virus Isolation, Production, and Purification

P. celtis and *P. quercus* were isolated in November 2014 from samples of surface soil taken at the
base of two trees (*Celtis autralis* and *Quercus ilex*) less than 50 meters apart in an urban green
space of Marseille city (GPS: 43°15'16.00"N, 5°25'4.00"E). Their particles were morphologically
identical to previously characterized pandoraviruses (Fig. S1). The viral populations were
amplified by co-cultivation with *A. castellanii*. They were then cloned, mass-produced and
purified as previously described (Philippe et al., 2013).

2.2 Genome and transcriptome sequencing, annotation

The *P. celtis* genome was fully assembled from one PacBio SMRT cell sequence data with the
HGAP 4 assembler (Chin et al., 2013) from the SMRT link package version 5.0.1.9585 with
default parameters and the "aggressive" option=true. Genome polishing was finally performed
using the SMRT package. Stringent gene annotation was performed as previously described
(Legendre et al., 2018). Briefly, data from proteomic characterization of the purified virions were
combined with stranded RNA-seq transcriptomic data, as well as protein homology among
previously characterized pandoraviruses. The transcriptomic data were generated from cells
collected every hour over an infectious cycle of 15 hours. They were pooled and RNAs were
extracted prior poly(A)$^+$ enrichment. The RNA were then sent for sequencing. Stranded RNA-seq
reads were then used to accurately annotate protein-coding as well as non-coding RNA genes. A
threshold of gene expression (median read coverage >5 over the whole transcript) larger than the
lowest one associated to proteins detected in proteomic analyses was required to validate all
predicted genes (including novel genes) (Table 1). Genomic regions exhibiting similar expression
levels but did not encompass predicted proteins or did overlap with protein-coding genes
expressed from opposite strands were annotated as "non-coding RNA" (ncRNA) after assembly
by Trinity (Grabherr et al., 2011). When only genomic data were available, namely for *P.*
pampulha, *P. massiliensi*s and *P. braziliensis* (Aherfi et al., 2018), we annotated protein-coding
genes using *ab initio* prediction coupled with sequence conservation information as previously
described (Legendre et al., 2018).

Gene clustering was performed on all available Pandoraviruses' protein-coding genes using
Orthofinder (Emms & Kelly, 2015) with defaults parameters except for the "msa" option for the
gene tree inference method.

Functional annotation of protein-coding genes was performed using a combination of protein
domains search with the CD-search tool (Marchler-Bauer & Bryant, 2004) and HMM-HMM
search against the Uniclust30 database with the HHblits tool (Remmert et al., 2011). In addition,
we used the same procedure to update the functional gene annotation of *P. salinus*, *P. dulcis*, *P.*
quercus, *P. macleodensis* and *P. neocaledonia* (Genbank IDs: KC977571, KC977570,
MG011689, MG011691 and MG011690).

2.3   Phylogenetic and Selection pressure analysis

The phylogenetic tree (Fig. 1) was computed from the concatenated multiple alignment of the
sequences of Pandoravirus core proteins corresponding to single-copy genes. The alignments of
orthologous genes peptide sequences were done using Mafft (Katoh et al., 2002). The tree was
computed using IQtree (Hoang et al., 2018) with the following options: -m MFP –bb 10000 –st
codon –bnni. The best model chosen was: GY+F+R5. Codon sequences were subsequently
mapped on these alignments. Ratios of non-synonymous (dN) over synonymous (dS) mutation
rates for pairs of orthologous genes were computed using the YN00 method from the PAML
package (Yang, 2007). Filters were applied so that dN/dS ratios were only considered if: dN > 0,
dS > 0, dS ≤ 2 and dN/dS ≤ 10. We also computed the Codon Adaptation Index (CAI) of *P. celtis*
genes using the cai tool from the Emboss package (Rice et al., 2000) as previously described
(Legendre et al., 2018).

2.4   Particle Proteomics

The *P. celtis* particle proteome was characterized by mass spectrometry-based proteomics from
purified viral particles as previously described (Legendre et al., 2018; Legendre et al., 2015).

3   Results

3.1   Main structural features of the *P. celtis* and *P. quercus* genomes

The *P. celtis* dsDNA genome sequence was assembled as a single 2,028,440 bp-long linear
contig, thus slightly shorter than the published 2,077,288-bp for *P. quercus*. Both contains 61%
of G+C. The two genomes exhibit a global collinearity well illustrated by a dotplot comparison of
their highly similar nucleotide sequences (Fig. 2). In particular, their 48.8 kb difference in
genome size does not obviously correspond to a large non-homologous region or a large-scale
duplication. Both genomes begin by a nearly perfect 19-kb long palindrome (labelled "P" in Fig.
2). As we did not observe this feature in the other published pandoravirus sequences, it may be
specific of *P. celtis* and *P. quercus*, or its absence in other genomes may result from flaws in the
assembly of terminal sequences due to insufficient read coverage or quality. Six kb downstream,
P. quercus exhibits a segment ([25,480-43,420]) nearly identical (20 indels) to the distal end of
the genome, inverted (labelled "T" in Fig. 2). Remnants of a similar feature appear blurred in *P.*
celtis, but are absent from the other pandoravirus genomes. As these regions are accurately
determined, we can infer that a duplication followed by an inversion/translocation of the distal
genome terminus occurred in the ancestor of *P. quercus* and *P. celtis* (between node 8 and 9 in
Fig. 1).

The next noticeable structural rearrangements consist of three segments denoted S_0, S_1, and S_2 in
Fig.2. Each of these segments are flanked by terminal inverted repeats (Fig. S2) and encode a
protein (respectively pclt_cds_98, pclt_cds_672, and pclt_cds_871) exhibiting both a BED zinc
finger domain and a C-terminal dimerisation region, typical of transposases of the
hobo/Activator/Tam3 (hAT) family. All these proteins exhibit the intact signature of hAT
transposases and are thus probably active (Atkinson, 2015). Using these sequences as queries, we
readily identified other well-conserved hAT transposase homologs in *P. pampulha*
(ppam_cds_67, cds_531, cds_663), *P. macleodensis* (pmac_cds_424, cds_799, cds 869) and *P.*
neocaledonia (pneo_cds_387, cds_113, cds_658, cds_798).

Besides their common transposase, the S_0 [10.3 kb, pos. 152,985-163,291], S_1 [10.3 kb, pos.
1,156,443-1,166,965], and S_2 [7.3 kb, pos. 1,452,773-1,460,053] transposons encode different
sets of proteins. S_0 encodes 9 proteins (pclt_cds_90-98). Except for the transposase, all of them
have no predicted function, and no recognizable homologs outside of the Pandoraviridae (i.e.
they are "family ORFans"). S_1 encodes 11 proteins (pclt_cds_672-682) all of which also have no
functional attributes and are family ORFans, except for the transposase. S_2 encodes 7 proteins
(pclt_cds_865-871), all of which have no recognizable signature (except for the transposase and a
F-box domain for pclt_cds_870). Besides the transposase, a single protein have paralogs in the
S_0, S_1, and S_2 transposons (pclt_cds_92, 681, 869), and one is only shared by S_0 and S_1
(pclt_cds_94, 678). These differences clearly suggest that S_0, S_1, S_2 are not the results of recent
duplication/transposition events from a common template.

A tentative scenario for the insertion of the *P. celtis* hAT transposons was inferred from their
presence/absence in *P. quercus* and the sequence similarity of the transposases. The S_1
transposon is the only one shared between the two strains (Fig. S2). Moreover, all the
orthologous proteins encoded in S_1 are 100% identical (pclt_cds_672-682 vs. pqer_cds_619-610),
including the transposases (pclt_cds_672 and pqer_cds_619). A first possibility is that the S_1
segment was already present in the ancestor of *P. celtis* and *P. quercus* (Fig. 1, node 9), then was
inverted and translocated about 70 kb downstream from its initial location in *P. celtis*. However,
since this transposon is absent from *P. inopinatum* (diverging after node 8, Fig. 1), it may have
been independently gained from the same source into *P. celtis* and *P. quercus* just after their
divergence as two variants within the local viral population. Interestingly, an unrelated sequence
of 30.5 kb was inserted at the homologous positions in *P. quercus* (pos. 1,177,379-1,207,935)
(Fig. S2). This insertion encodes 24 proteins (pqer_cds_659-682) 13 of which are anonymous and
ORFans (i.e. only homologous to other Pandoravirus proteins), the other exhibiting
uninformative motifs such as ankyrin repeats (pqer_cds_668, 669, 674-676, 680), F-box domains
(pqer_cds_681, 682), Morn repeat (pqer_cds_661) and Ring domain (pqer_cds_670). One protein
(pqer_cds_673) exhibits a low ($E<10^{-2}$) and partial similarity with a domain found at the N
terminus of structural maintenance of chromosomes (SMC) proteins. However, none of the
proteins encoded by this insertion bears any similarity with a hAT family transposase making the
mechanism and the origin of this insertion all the more puzzling.

Finally, the proposed scenario concerning S_0 and S_2 are simpler. As the two corresponding
transposases (pclt_cds_98 and pclt_cds_871) are only 91% identical to each other and less than
67% identical to pclt_cds_672, we propose that they resulted from two independent insertion
events that occurred after the *P. quercus*/*P. celtis* divergence, from distinct templates with little
overlap in their gene cargo (pclt_cds_92 and 869) besides active transposases. An alternative
scenario, would assume the presence of S_0 and S_2 in the *P. quercus*/*P. celtis* ancestor, and their
subsequent excision in *P. quercus*. This scenario is less likely as *P. inopinatum* (Fig. 1, node 8)
does not encode any hAT transposase.

3.2   Tandem duplications

Approximately 28 kb downstream from the S_2 transposon (absent in *P. quercus*), the *P. celtis vs.*
P. quercus dot plot indicates a cluster of closely interspersed direct repeats (Fig. S3) coding for
paralogous proteins that all contain a highly conserved N-terminal fascin-like domain
(CD00257). This 80-residue long domain is found in actin-bundling/crosslinking proteins. Many
copies of these proteins are found in each of the known pandoraviruses. There are 17 (full-length)
copies in *P. quercus*, and 14 in *P. celtis*, the paralogs sharing from 100% to 40% identical
residues. Using standard phylogenetic analysis each of the *P. celtis* paralogs are unambiguously
clustered with *P. quercus* counterparts in 14 orthologous pairs, indicating that the multiplication
of these proteins took place prior to their divergence (Fig. S3). As the 3 additional *P. quercus*
paralogs (pqer_cds_130, 131 and 870) with no obvious counterpart in *P. celtis* are not very
similar to other *P. quercus* paralogs (max is 72% sequence identity between pqer_cds_870 and
pqer_cds_869), it is unlikely that they arise from post-divergence duplications. The most
parsimonious scenario is thus that their counterparts have been recently lost by *P. celtis*. In a dot
plot, the pqer_cds_130 and pqer_cds_131 genes correspond to a 4 kb deletion at position 224,350
in *P. celtis*. The remaining pqer_cds_870 is located in a dense cluster of 6 contiguous paralogs
(pqer_cds_866-871) in *P. quercus*, clearly homologous to a similar cluster of 5 contiguous

paralogs (pclt_cds_884-888) *in P. celtis* (Fig. S3). The deletion of a short segment of the *P. celtis*
genome is visible at this exact position (1,487,300). Another member of this cluster
(pclt_cds_886) just entered a pseudogenization process through the several insertion/deletions
breaking its original reading frame (Fig. S3).

3.3   Core genes and pan genome update

We previously designed a stringent gene annotation scheme to minimize overpredictions in
G+C–rich genomes. In particular, genes predicted to encode proteins without database homolog
(i.e. ORFans) were required to overlap with detected sense transcripts to be validated (Legendre
et al., 2018). According to this protocol, *P. celtis* encodes at least 1184 protein-coding genes
(CDS) compared to 1185 protein-coding genes for *P. quercus*. Both encode a single tRNA(Pro).
It is worth noticing that despite a strong reduction compared to the number of proteins initially
predicted by standard methods (Philippe et al., 2013), the proportion of ORFans (i.e. w/o
homolog outside of the Pandoraviridae family) remains high at 68%. This confirms that the lack
of database homologs is not caused by bioinformatics overpredictions (Legendre et al., 2018). *P.*
celtis and *P. quercus* shared an average of 96% identical residues, as computed from 822
unambiguous orthologous proteins encoded by single copy genes (i.e. w/o paralogs).

We previously showed that the comparison of the first six available Pandoravirus genomes
rapidly converged toward a relatively small set of common genes, corresponding to 455 distinct
proteins (clusters) (Legendre et al., 2018). An even smaller estimate (352) was subsequently
proposed by another laboratory using a different clustering protocol and strains (Aherfi et al.,
2018). Now based on the ten fully sequenced Pandoraviruses and an optimized clustering
protocol (see Materials & Methods), the Pandoraviridae core gene set was found to contain 464
genes, close to the asymptotic limit suggested by Fig. 3. Including *P. celtis* in the analysis caused
the removal of two proteins from the list. Their homologs in *P. quercus* are pqer_cds_672, and
pqer_cds_370, two proteins w/o recognizable functional attribute or homolog outside of the
Pandoraviridae. Compared to the average number of distinct proteins encoded by individual
Pandoravirus genomes, the proportion of those presumably essential is thus less than half. On the
other hand, the predicted gene content of *P. celtis* further increased the Pandoraviridae pan-
genome. Despite its overall close similarity with *P. quercus*, *P. celtis* remains on a growth curve
whereby each new randomly isolated Pandoravirus is predicted to add about 60 distinct proteins
to the total number of protein clusters already identified in the family (Fig. 3). The process by
which such addition could arise was further investigated by a detailed comparison of the genomes
of the most similar relatives, *P. quercus* and *P. celtis*.

3.4   Protein-coding genes unique to *P. celtis* or *P. quercus*

Given the strong similarity between the two genomes, determining their difference in gene
content is straightforward. However, these differences can be due to various mechanisms: the
duplication of existing genes, their differential loss, their acquisition by horizontal transfer, or
their *de novo* creation. Here we wanted to focus on the later mechanism, previously proposed to
be prominent in the Pandoraviridae family (Legendre et al., 2018). The candidate genes most
likely resulting from *de novo* creation would be those unique to *P. celtis* or *P. quercus*. If there
are no homolog in any of the other Pandoraviridae, *de novo* creation (or acquisition from an
unknown source) becomes the most parsimonious evolutionary scenario, compared to a scenario

whereby such a gene, present in an ancestral Pandoraviridae would have been independently lost
multiple times. The 30 protein-coding genes unique to *P. celtis* (N=9) and *P. quercus* (N=21) are
listed in Table 1. As previously noticed (Legendre et al., 2018), these presumed novel genes are
associated with statistics that are significantly different from other pandoravirus protein-coding
genes. They exhibit a lower Codon Adaptation Index (CAI= 0.233 *vs* 0.351, Wilcoxon test, p <
10^{-14}), a lower G+C content (57.5% *vs* 64.4%, p < 10^{-16}), and encode smaller proteins (length (aa)
=126.6 *vs* 387.3, p < 10^{-15}). The two later characteristics make them similar to the random ORFs
found in intergenic regions (i.e. the so-called "protogenes") from which we proposed they
originated.

We also noticed that the proteins encoded by the novel genes exhibit an amino-acid composition
widely different from the rest of the predicted proteome (Chi-2, df=19, p<10^{-10}), with the largest
variations observed for lysine (increasing from 2% to 5%), phenylalanine (2.2% to 4%), and
arginine (8.6% to 11.3%). Owing to the anomalous proportions of these residues, 25 out of the 30
novel *P. celtis* and *P. quercus* unique proteins are predicted to be DNA-binding (Szilágyi &
Skolnick, 2006) (Table 1). Although such predictions are subject to caution (as the predictions
remains the same for shuffled sequences) binding novel proteins to the viral genome might help
segregate loosely folded (i.e. intrinsically disordered) proteins and diminish their potential
interference with the rest of the viral proteome. Over time, these proteins may also gain some
regulatory functions.

We investigated the putative intergenic origin of these 30 strain-specific protein-coding genes by
searching for remote sequence similarity in the genome of all other Pandoraviruses, using tblastn
(protein query against six reading frame translation, E<10^{-3}) (Sayers et al., 2018). For seven of
them, one unique to *P. celtis* and six unique to *P. quercus*, we found no similarity within the non-
coding moiety of other pandoraviruses. Our most parsimonious interpretation is that they
represent recent independent additions in each genome by *de novo* creation or transfer from an
unknown organism w/o known relative. A slightly less parsimonious scenario would be their
addition prior to node 9, followed by differential losses in *P. quercus* or *P. celtis.* Remote but
significant (E<10^{-3}) similarities were detected for the 25 other novel genes within non-coding
regions of other pandoraviruses, all of them members of Clade A (Fig. 1). These positive matches
were distributed as followed: five in strains that diverged since node 9, eight in all strains that
diverged since node 8, four in all strains that diverged since node 5, and six in all strains that
diverged since node 2. We did not detect significant matches in earlier diverging Clade B
members. The distribution of these traces in pandoraviruses at various evolutionary distances is
interpreted in the discussion section.

3.5   ncRNAs

We annotated *P. celtis* 161 ncRNAs (Table S1), a number close to the 157 ncRNAs predicted for
P. quercus using the same protocol. Such numerous transcripts without protein coding capability
were previously noticed for other pandoravirus strains (Legendre et al., 2018). The *P. celtis*
ncRNAs vary in length from 234 to 4,456 nucleotides (median= 1384, mean=1413±639). By
comparison, the *P. quercus* ncRNAs range from 273 to 5,176 (median=1,189, mean=1299±738).
The two distributions are not significantly different (T-test: p>0.13). The expression levels (in
median read coverage) of *P. celtis* ncRNAs vary from 92 to 1608 (median=283) compared to
3665 to 506,994 (median=690) for protein coding transcripts. A similar non-coding *versus* coding

expression median ratio (≈ 0.41) was previously found for *P. quercus* (ncRNAs median =228,
protein-coding transcript median = 551), in a distinct experiment. As previously noticed for other
pandoravirus strains, most of *P. celtis* ncRNAs (154/161=95.6%)) are overlapping by more than
half of their length with a protein-coding transcript expressed from the opposite strand, and only
7 are mostly intergenic.

Although the *P. celtis* and *P. quercus* genomes share a very large proportion of their protein-
coding genes (1146/1184=96.8%), this was not the case for ncRNAs. We found that only 87 of *P.*
celtis ncRNAs (87/161=54%) exhibit a homolog among *P. quercus* ncRNAs (Table S1). The 74
P. celtis ncRNAs without homologs thus mostly correspond to the lack of detectable transcription
in the cognate sequence of the *P. quercus* genome. Unexpectedly, the *P. celtis* and *P. quercus*
novel protein-coding genes discussed above (section 3.4) only rarely overlap with ncRNAs in
other strains. The single case is pclt_cds_308, the coding region of which overlaps with a *P.*
quercus ncRNA (pquer_ncRNA_47) (Table 1).

3.6   Selection pressure on new genes

Protein-coding genes shared by at least two different Pandoraviruses provide an opportunity to
estimate the selection pressure acting on them by computing the ratio of nonsynonymous (dN) to
synonymous (dS) substitutions per site. We computed the dN/dS ratio of genes shared by
increasingly close Pandoraviruses, dating their creation or acquisition by reference to the
corresponding node in the Pandoravirus phylogenetic tree (nodes 1 to 9 in Fig 1). All pairs of
orthologous gene sequences were thus analyzed using the YN00 algorithm from the PAM
package (Yang, 2007) and assigned to their most likely creation/acquisition node based on their
presence or absence in various clades of Pandoraviruses. As shown in Fig. 4 all genes are on
average under negative selection pressure (dN/dS < 1), including the presumably most recently
created/acquired genes as those only found in *P. quercus* and *P. celtis* (i.e. node 9 in Fig. 1). As
previously documented for short evolutionary distances and very close gene sequences (our case),
the computed dN/dS values are probably overestimated (i.e. closer to 1) given that a fraction of
the deleterious mutations might not yet be fixed (Rocha et al., 2006). This is consistent with the
observed negative correlation between the depth of creation/acquisition and the computed
purifying selection (Fig. 4). In other words recently created/acquired genes appear under selective
constraints weaker than that of "older" genes. The fact that dN/dS values tend to decrease with
longer divergence times was previously noticed (Rocha et al., 2006).

4   Discussion

Following our previous analysis of 6 available members of the Pandoraviridae family, we
proposed that the gigantism of their genomes as well as the large proportion of ORFans among
their encoded proteins was the result of unusual evolutionary mechanisms, including *de novo*
gene creation from previously non-coding sequences (Legendre et al., 2018). In this previous
study, we compared pandoravirus isolates exhibiting pairwise similarity ranging from 54 to 88%,
as computed from a super alignment of their orthologous proteins. At such evolutionary
distances, the observed differences are most often the result of multiple and overlapping
elementary variation processes the succession of which becomes impossible to retrace. The
subsequent isolation of *P. celtis*, a new pandoravirus strain very similar to *P. quercus* (with
orthologous proteins sharing 96% of identical residues in average), gave us the opportunity of

identifying and estimating the relative contributions of various types of genomic alterations at
work during their microevolution from their recent common ancestor.
335
The global analyses of the gene contents of the 10 members of the Pandoraviridae family
available today confirmed previous estimates of the number of core gene clusters (Aherfi et al.,
2018; Legendre et al., 2018) at 464. Such a small proportion (less than half) of presumably
"essential" genes compared to the total number of proteins encoded by each pandoravirus
genome (ranging from 1070 to 1430, using our stringent protocol) (Legendre et al., 2018) raises
the question of the origin and utility of so many "accessory" genes. Non-essential genes are
normally eliminated from the genomes of obligate intracellular parasites or symbionts through
reductive evolution (Floriano et al., 2018; Latorre & Manzano-Marín, 2017; McCutcheon &
Moran, 2011; Lopez-Madrigal et al., 2011; Corradi et al., 2010). The contrast is even more
puzzling when the size of the Pandoraviridae pan-genome shows no sign of leveling off after
reaching 1910 different protein-coding gene clusters, sustaining a trend predicting that each new
isolate will contribute 60 additional clusters. Understanding the mechanism by which new genes,
- most of which encode ORFan proteins -, appear in the genome of pandoraviruses was the main
goal of our study.
350
As we investigated the most visible alterations of the otherwise nearly perfect collinearity of the
P. celtis and *P. quercus* genomes, we identified 3 transposons of the hAT family (S_0, S_1 and S_2 in
Fig. 2). The cargo of these mobile elements was found to be variable in gene number (10 for S_0,
11 for S_1, and 7 for S_2) and with a single overlap (pclt_cds_92, 681, 869) in addition to the
transposases. These by-standing proteins exhibit no functional signature and have no homolog
outside of the Pandoraviridae family. Interestingly, hAT family transposases have recently been
identified in various Acanthamoeba species (Zhang et al., 2018). However, the gene contents of
the *P. celtis* and *P. quercus* hAT transposons indicates that these mobile elements are not
prominent vehicles of lateral gene transfers from the amoebal hosts to the pandoraviruses. With
the exception of pclt_cds_870 (encoded by S_2), newly inserted transposons do not contribute
genes unique to *P. celtis*. The hAT transposable elements only appear to ferry genes between
pandoravirus strains, generating non-local duplications. Such exchanges might occur within an
amoebal host undergoing multiple infections. To our knowledge, this is the first identification of
hAT family transposons in DNA viruses. The fact that hAT transposons are not present in other
well-documented families of large and giant virus infecting Acanthamoeba (Abergel et al., 2015;
Aherfi et al., 2016; Fabre et al., 2017; Zhang et al., 2018) suggests that its transfer from host to
virus is a rare event, or is specifically linked to Pandoravirus infections.
368
This work identified 30 genes unique to *P. celtis* or *P. quercus* (Table 1) that we interpreted as
encoding novel proteins that appeared after the recent divergence of these two strains from their
common ancestor (i.e. below node 9 in Fig. 1). These new genes are uniformly distributed along
the *P. celtis* and *P. quercus* genomes, and do not co-localize with large genomic insertions or
rearrangements, except for pqer_cds_665 and pqer_cds_673 encoded by the 30.5 kb segments
unique to *P. quercus* (section 3.1). As noticed in our previous study (Legendre et al., 2018) all
these proteins are strict ORFans (not detected in any other organism, virus or other pandoravirus
strains) and exhibit statistical features different from other pandoravirus genes, including a G+C
content closer to that of non-coding/intergenic regions. We thus proposed that new proteins could
be *de novo* created by the triggering of the transcription and translation of pre-existing non-
coding sequences (the so-called "protogenes"). Here we further investigated this hypothesis by

looking for eventual similarities of the new protein sequences with non-coding regions in other
pandoraviruses at increasing evolutionary distances (Fig. 1). For 8 of the 9 new *P. celtis* proteins
and 15 of the 21 new *P. quercus* proteins we detected significant non-coding matches in various
pandoravirus strains, in good agreement with their phylogenetic relationships. Trace of 5 new
genes were only detected at the level of node 9, 8 at the more ancient node 8, 4 at the node 5, and
6 at the root (node 2) of the clade A pandoraviruses. Although these numbers are small and
subject to fluctuations, such a distribution of hits suggests an evolutionary process combining a
continuous spontaneous generation of open reading frames, followed by their drift back as non-
coding sequences, unless they become transcriptionally active, and fixed as new proteins in a
given strain. The maximum value of detected traces at node 8 might thus result from a
compromise between the time interval required to generate *de novo* proteins, and the time during
which they could retain a detectable similarity with the drifting non-coding regions from which
they originated.
393
According to the above gene creation hypothesis, initially non-coding regions could act as
precursors for new proteins, by the random opening of a suitable reading frame, followed by its
transcriptional activation. Such a scenario is compatible with the distribution of new gene
matches in other pandoravirus strains discussed above, where 8 correspond to intergenic regions,
and 5 overlap with UTRs (Table 1). The flexibility of intergenic regions is further attested by the
fact that 11 of the new genes have no matches. As shown in Table 1, a single new gene
(pclt_cds_308) was found to co-localize with a ncRNA (pqer_ncRNA_47) despite the large
number of ncRNAs detected in the *P. celtis* and *P. quercus* genomes. ncRNAs thus do not appear
to be necessary intermediates in the emergence of new proteins. The low proportion of conserved
ncRNAs between the two close strains (54%) suggests that the on/off expression status of non-
coding genomic sequences is fluctuating very fast, and may even be variable at the viral
population level.
406
The fact that a negative selection pressure (dN/dS <1) is associated to the genes appeared since
the divergence of *P. celtis* and *P. quercus* decisively reinforces the hypothesis of their *de novo*
creation. Firstly, it provides the proof that these genes are truly expressed as *bona fide* proteins,
given that differences between synonymous *vs.* non-synonymous mutations cannot be generated
within non-translated nucleotide sequences. Any significant deviation from neutrality (dN/dS ≈1)
can only be due to a selection process exerted on certain amino acids at certain positions of a true
protein in order to improve, preserve, or eliminate its function.
414
The analysis of proteins whose creation appears most recent (node 9 in Fig. 4) indicates a
negative selection pressure whose (probably overestimated) average value (0.58) could
correspond to functions slightly increasing the pandoravirus fitness. However, the interpretation
of the selection pressure in the context of a function is a problem here, since we previously
pointed out that it is highly unlikely that a newly created small protein will have a function
(enzymatic, regulatory or structural) in the first place, even less a beneficial one (Legendre et al.,
2018).
Thus, we propose an alternative interpretation, based on the knowledge accumulated over the
years on the principles governing the stability of proteins. According to our scenario, most new
proteins are from the translation of pre-existing non-coding "protogenes" (see 3.4) and the rest
from DNA inserts of unknown origin (such as pqer_cds_665 and pqer_cds_673). It is then
unlikely that many of these new proteins lacking an evolutionary history will adopt a globular

fold (Watters et al. 2006; Monsellier & Chiti, 2007; Tanaka et al., 2010). Most will be toxic by
causing nonspecific aggregations within the host-virus proteome and be quickly eliminated
through the reversal of these newborn viral genes to an untranscribed/untranslated state. In
contrast, proteins whose folded 3-D structures do not prove to be toxic will enter an evolutionary
process involving a negative selection pressure promoting the conservation of the amino acids
responsible for their stability. Their proportion was estimated at about 34% for 100-200 residue
proteins (Miao et al., 2004). In absence of an initial function, the other positions/residues will
evolve in a neutral manner. Once "fixed", the new genes will exhibit an average selection
pressure lower than one, combining 1/3 of negative selection with 2/3 of neutrality. Following
many studies that have concluded that efficient folding (Watters et al., 2006) and prevention of
aggregation are important drivers of protein evolution (see Monsellier & Chiti, 2007), our
scenario predicts that many genes specific to pandoraviruses should encode proteins not
increasing their fitness, but whose stable 3-D structures may eventually serve as innovative
platforms for new functions.
441

In conclusion, the detailed comparative analysis of *P. celtis* with its very close relative *P. quercus*
enabled us to estimate the contributions of various microevolutionary processes to the steady
increase of the pan-genome of the proposed Pandoraviridae giant virus family. We first showed
that large-scale genomic rearrangements (segmental duplications, translocations) are associated
to transposable elements of the hAT family, widespread in metazoans, but until now unique to
this family of viruses. However, these mobile elements mostly appear to shuffle pandoravirus
genes between strains, without creating new ones or promoting host-to-virus horizontal gene
transfers. In contrast with the popular view that horizontal gene transfer plays an important role
in the evolution of large DNA viruses (Yutin & Koonin (2013); Koonin et al. 2015; Schulz et al.,
2017), this is definitely not the main cause of the inflation of the Pandoraviridae genomes, as we
previously argued (Claverie & Abergel, 2013; Philippe et al., 2013; Abergel et al., 2015;
Legendre et al., 2018). Finally, we also found that locally repeated regions are the siege of a
competition between tandem duplication and gene deletion, locally reshaping the genomes
without contributing to net genetic innovation (Fig. S3).

In continuity with our previous work on more distant pandoravirus strains, we found that the 30
protein-coding genes born since the divergence between the very close *P. quercus* and *P. celtis*
strains were derived from preexisting non-coding sequences or small DNA segments of unknown
origins inserted at randomly interspersed locations. We propose that random ORFs constantly
emerge in non-coding regions and that their transcription is turned on in some pandoravirus
strains, while they remain silent until they are deleted or diverge beyond recognition in others
strains. Our results add strong support to the constant *de novo* creation of proteins, few of which
are retained with a little initial impact on the virus fitness until eventually acquiring a selectable
function.

Such a scenario, particularly visible and active in the Pandoraviridae, might also apply to the
large proportion of ORFans encoded by other DNA viruses, from large eukaryotic viruses
(Abergel et al., 2015) to much smaller bacteriophages, in which they can be the target of global
functional studies (Berjon-Otero et al., 2017). Interestingly, the *de novo* gene creation scenario is
gaining more and more acceptance beyond the realm of virology, recently to explain the origin of
orphan protein even in mammals (Schmitz et al., 2018).

5   Author Contributions

CA, ML, JMC, and YC designed the experiments. JMA, AL, NP, SJ, and YC contributed to the
data and performed the experiments. ML, SN, NTT, OP, CA and JMC analyzed the data. ML,
CA, and JMC wrote the manuscript.

6   Funding

This work was partially supported by the French National Research Agency (ANR-14-CE14-
0023-01), France Genomique (ANR-10-INSB-01-01), Institut Français de Bioinformatique
(ANR–11–INSB–0013), the Fondation Bettencourt-Schueller (OTP51251), by a DGA-MRIS
scholarship, and by the Provence-Alpes-Côte-d'Azur région (2010 12125). Proteomic
experiments were partly supported by the Proteomics French Infrastructure (ANR-10-INBS-08-
01) and Labex GRAL (ANR-10-LABX-49-01). We thank the support of the discovery platform
and informatics group at EDyP and the PACA-Bioinfo platform.
483

7   Conflict of Interest Statement

The authors declare that the research was conducted in the absence of any commercial or
financial relationships that could be construed as a potential conflict of interest.

8   Acknowledgments

We kindly thank Hua-Hao Zhang for providing us with the hAT-transposon sequence he
identified in the *Acanthamoeba castellanii* genome.

9   Supplementary Material

The Supplementary Material for this article can be found online at:
https://www.frontiersin.org/articles/XXX/full#supplementary-material

Figure S1. TEM images of ultrathin sections of *P. celtis* and *P. quercus* viral particles.

Figure S2. Location of the hAT transposases in the S0, S1, and S2 segments and transposons
flanking sequences in *P. celtis*. The location of the tranposase gene is indicated in red**.**
498
499 **Figure S3.** Details of two tandem repeat clusters of fascin-domain containing genes.
500

10   References

Abergel, C., Legendre, M., Claverie, J.M. (2015) The rapidly expanding universe of giant
viruses: Mimivirus, Pandoravirus, Pithovirus and Mollivirus. FEMS Microbiol. Rev. 39(6), 779-
96. doi: 10.1093/femsre/fuv037.

Aherfi, S., Andreani, J., Baptiste, E., Oumessoum, A., Dornas, F.P., Andrade, A.C.D.S.P., et al.
(2018). A large open pangenome and a small core genome for giant pandoraviruses. Front.
Microbiol. 9, 1486. doi: 10.3389/fmicb.2018.01486.

Aherfi, S., Colson, P., La Scola, B., Raoult, D. (2016) Giant Viruses of Amoebas: An Update.
Front. Microbiol. 7, 349. doi: 10.3389/fmicb.2016.00349.

Antwerpen, M.H., Georgi, E., Zoeller, L., Woelfel, R., Stoecker, K., Scheid, P. (2015) Whole-
genome sequencing of a pandoravirus isolated from keratitis-inducing acanthamoeba. Genome
Announc. 3(2). pii: e00136-15. doi:10.1128/genomeA.00136-15.

Atkinson, P.W. (2015) hAT Transposable Elements. Microbiol. Spectr. 3(4). doi:
10.1128/microbiolspec. MDNA3-0054-2014.

Berjón-Otero M, Lechuga A, Mehla J, Uetz P, Salas M, Redrejo-Rodríguez M. (2017) Bam35
tectivirus intraviral interaction map unveils new function and localization of phage ORFan
proteins. J. Virol. 91, e00870-17. doi:10.1128/JVI.00870-17.

Chin, C.S., Alexander, D.H., Marks, P., Klammer, A.A., Drake, J., Heiner C., et al. (2013)
Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. Nat.
Methods 10(6), 563-9. doi: 10.1038/nmeth.2474.

Claverie, J.M., Abergel, C. (2013) Open questions about giant viruses. Adv. Virus Res. 85, 25-
56. doi: 10.1016/B978-0-12-408116-1.00002-1.

Claverie, J.M., Abergel, C., Legendre M. (2018) Giant viruses that create their own genes. Med.
Sci. (Paris) 34, 1087-1091. doi: 10.1051/medsci/2018300.

Corradi, N., Pombert, J.F., Farinelli, L., Didier, E.S., Keeling, P.J. (2010) The complete sequence
of the smallest known nuclear genome from the microsporidian Encephalitozoon intestinalis. Nat.
Commun. 1, 77. doi: 10.1038/ncomms1082.

Emms, D.M., Kelly, S. (2015) OrthoFinder: solving fundamental biases in whole genome
comparisons dramatically improves orthogroup inference accuracy. Genome Biol. 16, 157. doi:
10.1186/s13059-015-0721-2.

Fabre, E, Jeudy, S, Santini, S, Legendre, M, Trauchessec, M, Couté, Y, et al. (2017)
Noumeavirus replication relies on a transient remote control of the host nucleus. Nat. Commun.
8, 15087. doi: 10.1038/ncomms15087.

Floriano, A.M., Castelli, M., Krenek, S., Berendonk, T.U., Bazzocchi, C., Petroni, G., et al.
(2018) The Genome Sequence of "Candidatus Fokinia solitaria": Insights on Reductive Evolution
in Rickettsiales. Genome Biol. Evol. 10(4), 1120-1126. doi: 10.1093/gbe/evy072.

Grabherr, M.G., Haas, B.J., Yassour, M., Levin, J.Z., Thompson, D.A., Amit, I., et al. (2011)
Full-length transcriptome assembly from RNA-Seq data without a reference genome. Nat.
Biotechnol. 29, 644-652. doi: 10.1038/nbt.1883.

Hoang, D.T., Chernomor, O., von Haeseler, A., Minh, B.Q., Vinh, L.S. (2018) UFBoot2:
Improving the Ultrafast Bootstrap Approximation. Mol. Biol. Evol. 35, 518-522. doi:
10.1093/molbev/msx281.

Katoh, K., Misawa, K., Kuma, K., Miyata, T. (2002) MAFFT: a novel method for rapid multiple
sequence alignment based on fast Fourier transform. Nucleic Acids Res. 30, 3059-3066.

Koonin, E.V., Krupovic, M., Yutin, N. (2015) Evolution of double-stranded DNA viruses of
eukaryotes: from bacteriophages to transposons to giant viruses. Ann. N Y Acad. Sci. 1341, 10-
24. doi: 10.1111/nyas.12728.

Krumsiek, J., Arnold, R., Rattei, T. (2007) Gepard: a rapid and sensitive tool for creating dotplots
on genome scale. Bioinformatics 23, 1026-1028. doi: 10.1093/bioinformatics/btm039.

Latorre, A., Manzano-Marín, A. (2017) Dissecting genome reduction and trait loss in insect
endosymbionts. Ann. N Y Acad Sci. 1389, 52-75. doi: 10.1111/nyas.13222.

Lopez-Madrigal, S., Latorre, A., Porcar, M., Moya, A., Gil, R. (2011) Complete genome
sequence of 'Candidatus Tremblaya princeps' strain PCVAL, an intriguing translational machine
below the living-cell status. J. Bacteriol. 193, 5587–5588. doi: 10.1128/JB.05749-11.

Legendre, M., Fabre, E., Poirot, O., Jeudy, S., Lartigue, A., Alempic, J.M., et al. (2018).
Diversity and evolution of the emerging Pandoraviridae family. Nat. Commun. 9, 2285. doi:
10.1038/s41467-018-04698-4.

Legendre, M., Lartigue, A., Bertaux, L., Jeudy, S., Bartoli, J., Lescot, M., et al. (2015) In-depth
study of Mollivirus sibericum, a new 30,000-y-old giant virus infecting Acanthamoeba. Proc.
Natl Acad. Sci. U S A. 112, E5327-E5335. doi: 10.1073/pnas.1510795112.

McCutcheon JP, Moran NA (2011) Extreme genome reduction in symbiotic bacteria. Nat. Rev.
Microbiol. 10, 13-26. doi: 10.1038/nrmicro2670.

Marchler-Bauer, A., Bryant, S.H. (2004) CD-Search: protein domain annotations on the fly.
Nucleic Acids Res. 32, W327-331. doi : 10.1093/nar/gkh454.

Miao, J., Klein-Seetharaman, J., Meirovitch, H. (2004) The optimal fraction of hydrophobic
residues required to ensure protein collapse. J. Mol. Biol. 344, 797-811. doi:
10.1016/j.jmb.2004.09.061.

Monsellier, E., Chiti, F. (2007) Prevention of amyloid-like aggregation as a driving force of
protein evolution. EMBO Rep. 8, 737-742. doi: 10.1038/sj.embor.7401034.

Philippe, N., Legendre, M., Doutre, G., Couté, Y., Poirot, O., Lescot, M., et al. (2013)
Pandoraviruses: amoeba viruses with genomes up to 2.5 Mb reaching that of parasitic eukaryotes.
Science 341, 281-286. doi: 10.1126/science.1239181.

Remmert, M., Biegert, A., Hauser, A., Söding, J. (2011) HHblits: lightning-fast iterative protein
sequence searching by HMM-HMM alignment. Nat. Methods. 9, 173-175. doi:
10.1038/nmeth.1818.

Rice, P., Longden, I., Bleasby, A. (2000) EMBOSS: The European Molecular Biology Open
Software Suite. Trends Genet. 16, 276-277. doi:10.1016/S0168-9525(00)02024-2

Rocha, E.P.C., Smith, J.M., Hurst, L.D., Holden, M.T., Cooper, J.E., Smith, N.H., et al. (2006)
Comparisons of dN/dS are time dependent for closely related bacterial genomes. J. Theor. Biol.
239, 226-235. doi: 10.1016/j.jtbi.2005.08.037

Sayers, E.W., Agarwala, R., Bolton, E.E., Brister, J.R., Canese, K., Clark, K., et al. (2019)
Database resources of the National Center for Biotechnology Information. Nucleic Acids Res.
47(D1), D23-D28. doi: 10.1093/nar/gky1069.

Schulz, F., Yutin, N., Ivanova, N.N., Ortega, D.R., Lee, T.K., Vierheilig, J., et al. (2017) Giant
viruses with an expanded complement of translation system components. Science 356, 82-85.
doi: 10.1126/science.aal4657.

Schmitz JF, Ullrich KK, Bornberg-Bauer E. (2018) Incipient de novo genes can evolve from
frozen accidents that escaped rapid transcript turnover. Nat. Ecol. Evol. 2, 1626-1632. doi:
10.1038/s41559-018-0639-7.

Szilágyi, A, Skolnick, J. (2006) Efficient prediction of nucleic acid binding function from low-
resolution protein structures. J. Mol. Biol. 358, 922-933. doi: 10.1016/j.jmb.2006.02.053.

Tanaka, J., Doi, N., Takashima, H., Yanagawa, H. (2010) Comparative characterization of
random-sequence proteins consisting of 5, 12, and 20 kinds of amino acids. Protein Sci. 19, 786-
795. doi: 10.1002/pro.358.

Yang, Z. (2007) PAML 4: phylogenetic analysis by maximum likelihood. Mol. Biol. Evol. 24,
1586-1591. doi: 10.1093/molbev/msm088.

Watters AL, Deka P, Corrent C, Callender D, Varani G, Sosnick T, et al. (2007). The highly
cooperative folding of small naturally occurring proteins is likely the result of natural selection.
Cell 128, 613-624. doi: 10.1016/j.cell.2006.12.04.

Yutin, N., Koonin, E.V. (2013) Pandoraviruses are highly derived phycodnaviruses. Biol. Direct.
8, 25. doi: 10.1186/1745-6150-8-25.

Zhang, H.H., Zhou, Q.Z., Wang, P.L., Xiong, X.M., Luchetti, A., Raoult, D., et al. (2018)
Unexpected invasion of miniature inverted-repeat transposable elements in viral genomes. Mob.
DNA 9, 19. doi: 10.1186/s13100-018-0125-4.

11   Data Availability Statement

P. celtis annotated genome is deposited in GenBank under Accession n° MK174290.

12   Tables

609   **Table 1.** *P. celtis* and *P. quercus* unique protein-coding genes

Gene #	Size (aa)	Most ancestral detection	Homolog in *P. quercus*	Predicted DNA binding	Median RNAseq read coverage
pclt_cds_11	69	Since node 5	none	yes	704
pclt_cds_308	181	Since node 2	pqer_ncRNA_47	no	986
pclt_cds_350	121	Since node 9	intergenic	yes	57
pclt_cds_376	145	Since node 5	3'UTR pqer_cds_371	yes	454
pclt_cds_725	104	none	none	yes	46
pclt_cds_870	205	Since node 2	none	yes	2027
pclt_cds_995	149	Since node 5	5'UTR pqer_cds_981	yes	13
pclt_cds_1081	125	Since node 8	5'UTR pqer_cds_1061	yes	12,090
pclt_cds_1084	114	Since node 8	intergenic	yes	130
			Homolog in *P. celtis*		
pqer_cds_6	76	Since node 8	none	yes	311
pqer_cds_13	117	Since node 5	intergenic	yes	99
pqer_cds_17	82	Since node 2	intergenic	yes	482
pqer_cds_53	85	Since node 8	intergenic	yes	48
pqer_cds_143	93	Since node 8	anti 5'UTR pclt_cds_146	yes	177
pqer_cds_151	71	Since node 9	intergenic	yes	21
pqer_cds_203	101	Since node 8	antisense pclt_cds_206	yes	528
pqer_cds_350	94	none	none	no	160
pqer_cds_474	74	Since node 9	intergenic	yes	114
pqer_cds_486	146	Since node 2	3'UTR pclt_cds_499	yes	71
pqer_cds_665	114	none	none	yes	1,955
pqer_cds_673	383	none	none	yes	656
pqer_cds_685	124	Since node 8	alternative frame pclt_cds_685	no	117
pqer_cds_736	121	Since node 9	intergenic	yes	70
pqer_cds_875	136	none	none	no	2,050
pqer_cds_876	74	Since node 2	none	no	1,695
pqer_cds_877	78	none	none	yes	320
pqer_cds_878	224	none	none	yes	231
pqer_cds_1061	152	Since node 8	alternative frame pclt_cds_1081	yes	14,186
pqer_cds_1178	84	Since node 9	anti 5'UTR pclt_cds_1203	yes	170
pqer_cds_1183	158	Since node 2	none	yes	198

13   Figure legends

Figure 1.

Phylogenetic tree of known, fully characterized Pandoraviruses. The tree was computed from
the Pandoraviruses' core genes (see Materials and Methods for details). The estimated bootstrap
values were all equal to 1 and thus not reported. Internal nodes are labeled according to their
depth in the tree.

Figure 2.

Dot plot (nucleotide) comparison of the *P. celtis vs. P. quercus* genomes. Computations and
display were generated using Gepard (Krumsiek et al., 2007). Specific regions, labeled P, T, S_0,
S_1 and S_2 are described in the main text.

Figure 3.

Pandoraviridae pan-genome and core-genome. The boxplot represents the number of protein
clusters as a function of the number of sequenced genomes in all possible combinations. The
whiskers correspond to the extreme data points.

Figure 4.

Estimated selection pressure acting on Pandoravirus genes as a function of their ancestry.
Shown is the mean of dN/dS ratios (filled bars) for genes that are unique to the pandoravirus
strains below a given node in their phylogeny tree (see Fig. 1). Error bars correspond to standard
deviations. Bars are colored according to the depth of the node in the tree. As a control, we
calculated the dN/dS of Pandoravirus core genes (empty bars). For a given node, we considered
pairs of orthologous genes whose common ancestor correspond to that node.

Figure S1.

TEM image of an ultrathin section of *P. celtis* and *P. quercus* viral particles. The structures
of the particles of the other Pandoraviridae strains do not exhibit any noticeable difference.

Figure S2.

Details of the *P. celtis* hAT transposons. Details of the S_0, S_1, and S_2 segments highlighted in
Fig. 2. The genomic coordinates of the transposons in *P. celtis* (horizontal) and their homologous
locations in *P. quercus* (vertical) are indicated. The cognate *P. celtis* transposase gene name and
its approximate location is indicated for each transposon. Other genes are not represented.
Homologous segments are missing in *P. quercus* for S_0 and S_2 (See section 3.1). The sequences
of the transposon boundaries are shown above the dot-plots. The TIRs (Terminally Inverted
Repeats) are highlighted in black. Conserved TIRs and TSDs (Target Site Duplications) are
depicted using arrows.

Figure S3.

Details of two tandem repeat clusters of fascin-domain containing genes. See section 3.2 for
comments. The dot-plot (horizontal: *P. celtis*, vertical *P. quercus*) and the corresponding
phylogenetic tree of the genes illustrate the well-ordered dynamics of this highly repeated
regions, alternating expansion, deletions and pseudogenization. For instance, we noticed the
presence of two alleles of the pclt_886 proteins in our initial sequence data. The minor one (482
residues, used in this tree) is 96.7% identical to its pqer_868 homolog, while the major one
results into a smaller protein (363 residues) diverging after the first 125 residues due to the
presence of 4 indels in the rest of the gene. The evolutionary history of the fascin-domain
containing proteins was inferred using the Neighbor-Joining method as implemented in Mega
(Kumar S, et al. (2018) MEGA X: Molecular Evolutionary Genetics Analysis across computing
platforms. Mol. Biol. Evol. 35:1547-1549.). The evolutionary distances were computed using the
Poisson correction method. The analysis involved 31 amino acid sequences. All positions
containing gaps and missing data were eliminated. There were 323 positions in the final dataset.