Pandoravirus celtis illustrates the microevolution processes at work in the giant Pandoraviridae genomes

Matthieu Legendre[1], Jean-Marie Alempic[1], Nadège Philippe[1], Audrey Lartigue[1], Sandra Jeudy[1], Olivier Poirot[1], Ngan Thi Ta[1], Sébastien Nin[1], Yohann Couté[2], Chantal Abergel[1*], Jean-Michel Claverie[1*]

[1]Aix Marseille Univ, CNRS, IGS, Structural and Genomic Information Laboratory (UMR7256), Mediterranean Institute of Microbiology (FR3479), 163 Avenue de Luminy, F-13288 Marseille, France

[2]Univ. Grenoble Alpes, CEA, Inserm, BIG-BGE, 38000 Grenoble, France.

* Correspondence:
Jean-Michel Claverie
jean-michel.claverie@univ-amu.fr
Chantal Abergel
chantal.abergel@igs.cnrs-mrs.fr

Keywords: de novo gene creation, comparative genomics, Acanthamoeba, Giant viruses, Soil viruses, hAT transposase

Abstract

With genomes of up to 2.7 Mb propagated in µm-long oblong particles and initially predicted to encode more than 2000 proteins, members of the Pandoraviridae family display the most extreme features of the known viral world. The mere existence of such giant viruses raises fundamental questions about their origin and the processes governing their evolution. A previous analysis of six newly available isolates, independently confirmed by a study including 3 others, established that the Pandoraviridae pan-genome is open, meaning that each new strain exhibits protein-coding genes not previously identified in other family members. With an average increment of about 60 proteins, the gene repertoire shows no sign of reaching a limit and remains largely coding for proteins without recognizable homologs in other viruses or cells (ORFans). To explain these results, we proposed that most new protein-coding genes were created *de novo*, from pre-existing non-coding regions of the G+C rich pandoravirus genomes. The comparison of the gene content of a new isolate, *P. celtis*, closely related (96 % identical genome) to the previously described *P. quercus* is now used to test this hypothesis by studying genomic changes in a microevolution range. Our results confirm that the differences between these two similar gene contents mostly consist of protein-coding genes without known homologs (ORFans), with statistical signatures close to that of intergenic regions. These newborn proteins are under slight negative selection, perhaps to maintain stable folds and prevent protein aggregation pending the eventual emergence of fitness-increasing functions. Our study also unraveled several insertion events mediated by a transposase of the hAT family, 3 copies of which are found in *P. celtis* and are presumably active. Members of the Pandoraviridae are presently the first viruses known to encode this type of transposase.

1   Introduction

The Pandoraviridae is a proposed family of giant dsDNA viruses - not yet registered by the
International Committee on Taxonomy of Viruses (ICTV) - multiplying in various species of
Acanthamoeba through a lytic infectious cycle. Their linear genomes, flanked by large terminal
repeats, range from 1.9 to 2.7 Mb in size, and are propagated in elongated oblong particles
approximately 1.2 μm long and 0.6 μm in diameter (Fig. S1). The prototype strain (and the one with
the largest known genome) is *Pandoravirus salinus*, isolated from shallow marine sediments off the
coast of central Chile (Philippe et al., 2013). Other members were soon after isolated from worldwide
locations. Complete genome sequences have been determined for *P. dulcis* (Melbourne, Australia)
(Philippe et al., 2013), P. *inopinatum* (Germany) (Antwerpen et al., 2015), *P. macleodensis*
(Australia), *P. neocaledonia* (New Caledonia), and *P. quercus* (France) Legendre et al., 2018), and
three isolates from Brazil (*P. braziliensis*, *P. pampulha*, and *P. massiliensis*) (Aherfi et al., 2018). A
standard phylogenetic analysis of the above strains suggested that the Pandoraviridae family consists
of two separate clades (Claverie et al., 2018; Fig. 1). The average proportion of identical amino acids
between Pandoraviridae orthologs within each clade is above 70% while it is below 55% between
members of the A and B clades. Following a stringent reannotation of the predicted protein-coding
genes using transcriptomic and proteomic data, our comparative genomic analysis reached the main
following conclusions (Legendre et al., 2018):

1) the uniquely large proportion of predicted proteins without homologs outside of the
Pandoraviridae (ORFans) is real and not due to bioinformatic errors induced by the above-
average G+C content (>60%) of Pandoravirus genomes;
2) the Pandoraviridae pan genome appears "open" (i.e. unbounded);
3) as most of the genes are unique to each strain are ORFans, they were not horizontally
acquired from other (known) organisms;
4) they are neither predominantly the result of gene duplications,

The scenario of *de novo* and *in situ* gene creation, supported by the analysis of their sequence
statistical signatures, thus became our preferred explanation for the origin of strain specific genes.

In the present study, we take advantage of the high similarity (96.7% DNA sequence identity)
between *P. quercus* and a newly characterized isolate, *P. celtis*, to investigate the microevolution
processes initiating the divergence between Pandoraviruses. Our results further support *de novo* gene
creation as a main diversifying force of the Pandoraviridae family.

2   Materials and Methods

2.1   Virus Isolation, Production, and Purification

P. celtis and *P. quercus* were isolated in November 2014 from samples of surface soil taken at the
base of two trees (*Celtis autralis* and *Quercus ilex*) less than 50 meters apart in an urban green space
of Marseille city (GPS: 43°15'16.00"N, 5°25'4.00"E). Their particles were morphologically identical
to previously characterized pandoraviruses (Fig. S1). The viral populations were amplified by co-
cultivation with *A. castellanii*. They were then cloned, mass-produced and purified as previously
described (Philippe et al., 2013).

2.2   Genome and transcriptome sequencing, annotation

The *P. celtis* genome was fully assembled from one PacBio SMRT cell sequence data with the
HGAP 4 assembler (Chin et al., 2013) from the SMRT link package version 5.0.1.9585 with default

parameters and the "aggressive" option=true. Stringent gene annotation was performed as previously
described (Legendre et al., 2018). Briefly, data from proteomic characterization of the purified
virions were combined with stranded RNA-seq transcriptomic data, as well as protein homology
among previously characterized pandoraviruses. The transcriptomic data were generated from cells
collected every hour over an infectious cycle of 15 hours. They were pooled and RNAs were
extracted prior poly(A)$^+$ enrichment. The RNA were then sent for sequencing. Stranded RNA-seq
reads were then used to accurately annotate protein-coding as well as non-coding RNA genes. A
threshold of gene expression (median read coverage >5 over the whole transcript) larger than the
lowest one associated to proteins detected in proteomic analyses was required to validate all
predicted genes (including novel genes) (Table 1).When only genomic data were available, namely
for *P. pampulha*, *P. massiliensi*s and *P. braziliensis* (Aherfi et al., 2018), we annotated protein-
coding genes using *ab initio* prediction coupled with sequence conservation information as
previously described (Legendre et al., 2018).

Gene clustering was performed on all available Pandoraviruses' protein-coding genes using
Orthofinder (Emms & Kelly, 2015) with defaults parameters except for the "msa" option for the gene
tree inference method.

Functional annotation of protein-coding genes was performed using a combination of protein
domains search with the CD-search tool (Marchler-Bauer & Bryant, 2004) and HMM-HMM search
against the Uniclust30 database with the HHblits tool (Remmert et al., 2011). In addition, we used
the same procedure to update the functional gene annotation of *P. salinus*, *P. dulcis*, *P. quercus*, *P.*
macleodensis and *P. neocaledonia* (Genbank IDs: KC977571, KC977570, MG011689, MG011691
and MG011690).

2.3   Phylogenetic and Selection pressure analysis

The phylogenetic tree (Fig. 1) was computed from the concatenated multiple alignment of the
sequences of Pandoravirus core proteins corresponding to single-copy genes. The alignments of
orthologous genes peptide sequences were done using Mafft (Katoh et al., 2002). The tree was
computed using IQtree (Hoang et al., 2018) with the following options: -m MFP –bb 10000 –st codon
–bnni. The best model chosen was: GY+F+R5. Codon sequences were subsequently mapped on these
alignments. Ratios of non-synonymous (dN) over synonymous (dS) mutation rates for pairs of
orthologous genes were computed using the YN00 method from the PAML package (Yang, 2007).
Filters were applied so that dN/dS ratios were only considered if: dN > 0, dS > 0, dS ≤ 2 and dN/dS ≤
10. We also computed the Codon Adaptation Index (CAI) of *P. celtis* genes using the cai tool from the
Emboss package (Rice et al., 2000) as previously described (Legendre et al., 2018).

2.4   Particle Proteomics

The *P. celtis* particle proteome was characterized by mass spectrometry-based proteomics from
purified viral particles as previously described (Legendre et al., 2018; Legendre et al., 2015).

3   Results

3.1   Main structural features of the *P. celtis* and *P. quercus* genomes

The *P. celtis* dsDNA genome sequence was assembled as a single 2,028,579 bp-long linear contig,
thus slightly shorter than the published 2,077,288-bp for *P. quercus*. Both contains 61% of G+C. The
two genomes exhibit a global collinearity well illustrated by a dotplot comparison of their highly
similar nucleotide sequences (Fig. 2). In particular, their 48.7 kb difference in genome size does not
obviously correspond to a large non-homologous region or a large-scale duplication. Both genomes
begin by a nearly perfect 19-kb long palindrome (labelled "P" in Fig. 2). As we did not observe this
feature in the other published pandoravirus sequences, it may be specific of *P. celtis* and *P. quercus*,
or its absence in other genomes may result from flaws in the assembly of terminal sequences due to
insufficient read coverage or quality. Six kb downstream, *P. quercus* exhibits a segment ([25,480-
43,420]) nearly identical (20 indels) to the distal end of the genome, inverted (labelled "T" in Fig. 2).
Remnants of a similar feature appear blurred in *P. celtis*, but are absent from the other pandoravirus
genomes. As these regions are accurately determined, we can infer that a duplication followed by an
inversion/translocation of the distal genome terminus occurred in the ancestor of *P. quercus* and *P.
celtis* (between node 8 and 9 in Fig. 1).

The next noticeable structural rearrangements consist of three segments denoted S_0, S_1, and S_2 in
Fig.2. Each of these segments are flanked by terminal inverted repeats (Fig. S2) and encode a protein
(respectively pclt_cds_98, pclt_cds_672, and pclt_cds_871) exhibiting both a BED zinc finger
domain and a C-terminal dimerisation region, typical of transposases of the hobo/Activator/Tam3
(hAT) family. All these proteins exhibit the intact signature of hAT transposases and are thus
probably active (Atkinson, 2015). Using these sequences as queries, we readily identified other well-
conserved hAT transposase homologs in *P. pampulha* (ppam_cds_67, cds_531, cds_663), *P.
macleodensis* (pmac_cds_424, cds_799, cds 869) and *P. neocaledonia* (pneo_cds_387, cds_113,
cds_658, cds_798).

Besides their common transposase, the S_0 [10.3 kb, pos. 153,101-163,406], S_1 [10.3 kb, pos.
1,156,664-1,166,916], and S_2 [7.3 kb, pos. 1,452,969-1,460,249] transposons encode different sets of
proteins. S_0 encodes 9 proteins (pclt_cds_90-98). Except for the transposase, all of them have no
predicted function, and no recognizable homologs outside of the Pandoraviridae (i.e. they are "family
ORFans"). S_1 encodes 11 proteins (pclt_cds_672-682) all of which also have no functional attributes
and are family ORFans, except for the transposase. S_2 encodes 7 proteins (pclt_cds_865-871), all of
which have no recognizable signature (except for the transposase and a F-box domain for
pclt_cds_870). Besides the transposase, a single protein have paralogs in the S_0, S_1, and S_2
transposons (pclt_cds_92, 681, 869), and one is only shared by S_0 and S_1 (pclt_cds_94, 678). These
differences clearly suggest that S_0, S_1, S_2 are not the results of recent duplication/transposition events
from a common template.

A tentative scenario for the insertion of the *P. celtis* hAT transposons was inferred from their
presence/absence in *P. quercus* and the sequence similarity of the transposases. The S_1 transposon is
the only one shared between the two strains (Fig. S2). Moreover, all the orthologous proteins
encoded in S_1 are 100% identical (pclt_cds_672-682 vs. pqer_cds_619-610), including the
transposases (pclt_cds_672 and pqer_cds_619). A first possibility is that the S_1 segment was already
present in the ancestor of *P. celtis* and *P. quercus* (Fig. 1, node 9), then was inverted and translocated
about 70 kb downstream from its initial location in *P. celtis*. However, since this transposon is absent
from *P. inopinatum* (diverging after node 8, Fig. 1), it may have been independently gained from the
same source into *P. celtis* and *P. quercus* just after their divergence as two variants within the local
viral population. Interestingly, an unrelated sequence of 30.5 kb was inserted at the homologous
positions in *P. quercus* (pos. 1,177,379-1,207,935) (Fig. S2). This insertion encodes 24 proteins
(pqer_cds_659-682) 13 of which are anonymous and ORFans (i.e. only homologous to other

Pandoravirus proteins), the other exhibiting uninformative motifs such as ankyrin repeats
(pqer_cds_668, 669, 674-676, 680), F-box domains (pqer_cds_681, 682), Morn repeat
(pqer_cds_661) and Ring domain (pqer_cds_670). One protein (pqer_cds_673) exhibits a low (E<10^{-}
^{2}) and partial similarity with a domain found at the N terminus of structural maintenance of
chromosomes (SMC) proteins. However, none of the proteins encoded by this insertion bears any
similarity with a hAT family transposase making the mechanism and the origin of this insertion all
the more puzzling.

Finally, the proposed scenario concerning S_0 and S_2 are simpler. As the two corresponding
transposases (pclt_cds_98 and pclt_cds_871) are only 91% identical to each other and less than 67%
identical to pclt_cds_672, we propose that they resulted from two independent insertion events that
occurred after the *P. quercus*/*P. celtis* divergence, from distinct templates with little overlap in their
gene cargo (pclt_cds_92 and 869) besides active transposases. An alternative scenario, would assume
the presence of S_0 and S_2 in the *P. quercus*/*P. celtis* ancestor, and their subsequent excision in *P.*
quercus. This scenario is less likely as *P. inopinatum* (Fig. 1, node 8) does not encode any hAT
transposase.

3.2   Tandem duplications

Approximately 28 kb downstream from the S_2 transposon (absent in *P. quercus*), the *P. celtis vs*. *P.*
quercus dot plot indicates a cluster of closely interspersed direct repeats (Fig. S3) coding for
paralogous proteins that all contain a highly conserved N-terminal fascin-like domain (CD00257).
This 80-residue long domain is found in actin-bundling/crosslinking proteins. Many copies of these
proteins are found in each of the known pandoraviruses. There are 17 (full-length) copies in *P.*
quercus, and 14 in *P. celtis*, the paralogs sharing from 100% to 40% identical residues. Using
standard phylogenetic analysis each of the *P. celtis* paralogs are unambiguously clustered with *P.*
quercus counterparts in 14 orthologous pairs, indicating that the multiplication of these proteins took
place prior to their divergence (Fig. S3). As the 3 additional *P. quercus* paralogs (pqer_cds_130, 131
and 870) with no obvious counterpart in *P. celtis* are not very similar to other *P. quercus* paralogs
(max is 72% sequence identity between pqer_cds_870 and pqer_cds_869), it is unlikely that they
arise from post-divergence duplications. The most parsimonious scenario is thus that their
counterparts have been recently lost by *P. celtis*. In a dot plot, the pqer_cds_130 and pqer_cds_131
genes correspond to a 4 kb deletion at position 224,500 in *P. celtis*. The remaining pqer_cds_870 is
located in a dense cluster of 6 contiguous paralogs (pqer_cds_866-871) in *P. quercus*, clearly
homologous to a similar cluster of 5 contiguous paralogs (pclt_cds_884-888) *in P. celtis* (Fig. S3).
The deletion of a short segment of the *P. celtis* genome is visible at this exact position (1,487,500).

3.3   Core genes and pan genome update

We previously designed a stringent gene annotation scheme to minimize overpredictions in G+C–
rich genomes. In particular, genes predicted to encode proteins without database homolog (i.e.
ORFans) were required to overlap with detected sense transcripts to be validated (Legendre et al.,
2018). According to this protocol, *P. celtis* encodes at least 1207 protein-coding genes (CDS) and
173 non-coding RNAs (ncRNAs), compared to 1185 protein-coding genes and 157 ncRNAs for *P.*
quercus. Both encode a single tRNA(Pro). It is worth noticing that despite a strong reduction
compared to the number of proteins initially predicted by standard methods (Philippe et al., 2013),
the proportion of ORFans (i.e. w/o homolog outside of the Pandoraviridae family) remains high at
68%. This confirms that the lack of database homologs is not caused by bioinformatics
overpredictions (Legendre et al., 2018). *P. celtis* and *P. quercus* shared an average of 96.5% identical

residues, as computed from 815 unambiguous orthologous proteins encoded by single copy genes
(i.e. w/o paralogs).

We previously showed that the comparison of the first six available Pandoravirus genomes rapidly
converged toward a relatively small set of common genes, corresponding to 455 distinct proteins
(clusters) (Legendre et al., 2018). An even smaller estimate (352) was subsequently proposed by
another laboratory using a different clustering protocol and strains (Aherfi et al., 2018). Now based
on the ten fully sequenced Pandoraviruses and an optimized clustering protocol (see Materials &
Methods), the Pandoraviridae core gene set was found to contain 465 genes, close to the asymptotic
limit suggested by Fig. 3. Including *P. celtis* in the analysis caused the removal of a single protein
from the list. Its homolog in *P. quercus* is pqer_cds_672, a 220-residue protein w/o recognizable
functional attribute or homolog outside of the Pandoraviridae. Compared to the average number of
distinct proteins encoded by individual Pandoravirus genomes, the proportion of those presumably
essential is thus less than half. On the other hand, the predicted gene content of *P. celtis* further
increased the Pandoraviridae pan-genome. Despite its overall close similarity with *P. quercus*, *P.*
celtis remains on a growth curve whereby each new randomly isolated Pandoravirus is predicted to
add about 60 distinct proteins to the total number of protein clusters already identified in the family
(Fig. 3). The process by which such addition could arise was further investigated by a detailed
comparison of the genomes of the most similar relatives, *P. quercus* and *P. celtis*.

3.4   Protein-coding genes unique to *P. celtis* or *P. quercus*

Given the strong similarity between the two genomes, determining their difference in gene content is
straightforward. However, these differences can be due to various mechanisms: the duplication of
existing genes, their differential loss, their acquisition by horizontal transfer, or their *de novo*
creation. Here we wanted to focus on the later mechanism, previously proposed to be prominent in
the Pandoraviridae family (Legendre et al., 2018). The candidate genes most likely resulting from *de*
novo creation would be those unique to *P. celtis* or *P. quercus*. If there are no homolog in any of the
other Pandoraviridae, *de novo* creation (or acquisition from an unknown source) becomes the most
parsimonious evolutionary scenario, compared to a scenario whereby such a gene, present in an
ancestral Pandoraviridae would have been independently lost multiple times. The 33 protein-coding
genes unique to *P. celtis* (N=12) and *P. quercus* (N=21) are listed in Table 1. As previously noticed
(Legendre et al., 2018), these presumed novel genes are associated with statistics that are
significantly different from other pandoravirus protein-coding genes. They exhibit a lower Codon
Adaptation Index (CAI= 0.227 *vs* 0.352, p < 10^{-8}), a lower G+C content (56.9% *vs* 64.7%, p < 10^{-10}),
and encode smaller proteins (length (aa) =123.9 *vs* 387.3, p < 10^{-14}). The two later characteristics
make them similar to the random ORFs found in intergenic regions (i.e. the so-called "protogenes")
from which we proposed they originated.

We also noticed that the proteins encoded by the novel genes exhibit an amino-acid composition
widely different from the rest of the predicted proteome (Chi-2, df=19, p<10^{-5}), with the largest
variations observed for lysine (increasing from 2.1% to 4.6%), phenylalanine (2.3% to 4.2%), and
arginine (8.6% to 11.7%). Owing to the anomalous proportions of these residues, 28 out of the 33
novel *P. celtis* and *P. quercus* unique proteins are predicted to be DNA-binding (Szilágyi &
Skolnick, 2006) (Table 1). Although such predictions are subject to caution (as the predictions
remains the same for shuffled sequences) binding novel proteins to the viral genome might help in its
protection as well as in precluding loosely folded (i.e. intrinsically disordered) proteins to interfere
with the rest of the viral proteome. Over time, these proteins may also gain some regulatory
functions.

We investigated the putative intergenic origin of these 33 strain-specific protein-coding genes by
searching for remote sequence similarity in the genome of all other Pandoraviruses, using tblastn
(protein query against six reading frame translation, $E<10^{-3}$) (Sayers et al., 2018). For eight of them,
two unique to *P. celtis* and six unique to *P. quercus*, we found no similarity within the non-coding
moiety of other pandoraviruses. Our most parsimonious interpretation is that they represent recent
independent additions in each genome by *de novo* creation or transfer from an unknown organism
w/o known relative. A slightly less parsimonious scenario would be their addition prior to node 9,
followed by differential losses in *P. quercus* or *P. celtis.* Remote but significant ($E<10^{-3}$) similarities
were detected for the 25 other novel genes within non-coding regions of other pandoraviruses, all of
them members of Clade A (Fig. 1). These positive matches were distributed as followed: five in
strains that diverged since node 9, eleven in all strains that diverged since node 8, three in all strains
that diverged since node 5, and six in all strains that diverged since node 2. We did not detect
significant matches in earlier diverging Clade B members. The distribution of these traces in
pandoraviruses at various evolutionary distances is interpreted in the discussion section.

3.5   Selection pressure on new genes

Protein-coding genes shared by at least two different Pandoraviruses provide an opportunity to
estimate the selection pressure acting on them by computing the ratio of nonsynonymous (dN) to
synonymous (dS) substitutions per site. We computed the dN/dS ratio of genes shared by
increasingly close Pandoraviruses, dating their creation or acquisition by reference to the
corresponding node in the Pandoravirus phylogenetic tree (nodes 1 to 9 in Fig 1). All pairs of
orthologous gene sequences were thus analyzed using the YN00 algorithm from the PAM package
(Yang, 2007) and assigned to their most likely creation/acquisition node based on their presence or
absence in various clades of Pandoraviruses. As shown in Fig. 4 all genes are on average under
negative selection pressure (dN/dS < 1), including the presumably most recently created/acquired
genes as those only found in *P. quercus* and *P. celtis* (i.e. node 9 in Fig. 1). As previously
documented for short evolutionary distances and very close gene sequences (our case), the computed
dN/dS values are probably overestimated (i.e. closer to 1) given that a fraction of the deleterious
mutations might not yet be fixed (Rocha et al., 2006). This is consistent with the observed negative
correlation between the depth of creation/acquisition and the computed purifying selection (Fig. 4).
In other words recently created/acquired genes appear under selective constraints weaker than that of
"older" genes. The fact that dN/dS values tend to decrease with longer divergence times was
previously noticed (Rocha et al., 2006).

4   **Discussion**

Following our previous analysis of 6 available members of the Pandoraviridae family, we proposed
that the gigantism of their genomes as well as the large proportion of ORFans among their encoded
proteins was the result of unusual evolutionary mechanisms, including *de novo* gene creation from
previously non-coding sequences (Legendre et al., 2018). In this previous study, we compared
pandoravirus isolates exhibiting pairwise similarity ranging from 54 to 88%, as computed from a
super alignment of their orthologous proteins. At such evolutionary distances, the observed
differences are most often the result of multiple and overlapping elementary variation processes the
succession of which becomes impossible to retrace. The subsequent isolation of *P. celtis*, a new
pandoravirus strain very similar to *P. quercus* (with orthologous proteins sharing 96.5% of identical
residues in average), gave us the opportunity of identifying and estimating the relative contributions
of various types of genomic alterations at work during their microevolution from their recent
common ancestor.

301
The global analyses of the gene contents of the 10 members of the Pandoraviridae family available
today confirmed previous estimates of the number of core gene clusters (Aherfi et al., 2018;
Legendre et al., 2018) at 465. Such a small proportion (less than half) of presumably "essential"
genes compared to the total number of proteins encoded by each pandoravirus genome (ranging from
1070 to 1430, using our stringent protocol) (Legendre et al., 2018) raises the question of the origin
and utility of so many "accessory" genes. Non-essential genes are normally eliminated from the
genomes of obligate intracellular parasites or symbionts through reductive evolution (Floriano et al.,
2018; Latorre & Manzano-Marín, 2017; McCutcheon & Moran, 2011; Lopez-Madrigal et al., 2011;
Corradi et al., 2010). The contrast is even more puzzling when the size of the Pandoraviridae pan-
genome shows no sign of leveling off after reaching 1910 different protein-coding gene clusters,
sustaining a trend predicting that each new isolate will contribute 60 additional clusters.
Understanding the mechanism by which new genes, - most of which encode ORFan proteins -,
appear in the genome of pandoraviruses was the main goal of our study.
315
As we investigated the most visible alterations of the otherwise nearly perfect collinearity of the *P.*
celtis and *P. quercus* genomes, we identified 3 transposons of the hAT family (S_0, S_1 and S_2 in Fig.
2). The cargo of these mobile elements was found to be variable in gene number (10 for S_0, 11 for S_1,
and 7 for S_2) and with a single overlap (pclt_cds_92, 681, 869) in addition to the transposases. These
by-standing proteins exhibit no functional signature and have no homolog outside of the
Pandoraviridae family. Interestingly, hAT family transposases have recently been identified in
various Acanthamoeba species (Zhang et al., 2018). However, the gene contents of the *P. celtis* and
P. quercus hAT transposons indicates that these mobile elements are not prominent vehicles of
lateral gene transfers from the amoebal hosts to the pandoraviruses. With the exception of
pclt_cds_870 (encoded by S_2), newly inserted transposons do not contribute genes unique to *P. celtis*.
The hAT transposable elements only appear to ferry genes between pandoravirus strains, generating
non-local duplications. Such exchanges might occur within an amoebal host undergoing multiple
infections. To our knowledge, this is the first identification of hAT family transposons in DNA
viruses. The fact that hAT transposons are not present in other well-documented families of large and
giant virus infecting Acanthamoeba (Abergel et al., 2015; Aherfi et al., 2016; Fabre et al., 2017;
Zhang et al., 2018) suggests that its transfer from host to virus is a rare event, or is specifically linked
to Pandoravirus infections.
333
This work identified 33 genes unique to *P. celtis* or *P. quercus* (Table 1) that we interpreted as
encoding novel proteins that appeared after the recent divergence of these two strains from their
common ancestor (i.e. below node 9 in Fig. 1). These new genes are uniformly distributed along the
P. celtis and *P. quercus* genomes, and do not co-localize with large genomic insertions or
rearrangements, except for pqer_cds_665 and pqer_cds_673 encoded by the 30.5 kb segments unique
to *P. quercus* (section 3.1). As noticed in our previous study (Legendre et al., 2018) all these proteins
are strict ORFans (not detected in any other organism, virus or other pandoravirus strains) and exhibit
statistical features different from other pandoravirus genes, including a G+C content closer to that of
non-coding/intergenic regions. We thus proposed that new proteins could be *de novo* created by the
triggering of the transcription and translation of pre-existing non-coding sequences (the so-called
"protogenes"). Here we further investigated this hypothesis by looking for eventual similarities of the
new protein coding sequences with non-coding regions in other pandoraviruses at increasing
evolutionary distances (Fig. 1). For 10 of the 12 new *P. celtis* proteins and 15 of the 21 new *P.*
quercus proteins we detected significant non-coding matches in various pandoravirus strains, in good
agreement with their phylogenetic relationships. Trace of 5 new genes were only detected at the level
of node 9, 11 at the more ancient node 8, 3 at the node 5, and 6 at the root (node 2) of the clade A

pandoraviruses. Although these numbers are small and subject to fluctuations, such a distribution of
hits suggests an evolutionary process combining a continuous spontaneous generation of protogenes
followed by their drift as non-coding sequences, unless they are turned into a protein and fixed in a
given strain. The maximum value of detected traces at node 8 might thus result from a compromise
between the time interval required to generate enough non-coding protogenes, and the time during
which they retain a detectable similarity with the proteins they would eventually generate.
356
The fact that a negative selection pressure (dN/dS <1) is associated to the genes appeared since the
divergence of *P. celtis* and *P. quercus* decisively reinforces the hypothesis of their *de novo* creation.
Firstly, it provides the proof that these genes are truly expressed as *bona fide* proteins, given that
differences between synonymous *vs.* non-synonymous mutations cannot be generated within non-
translated nucleotide sequences. Any significant deviation from neutrality (dN/dS ≈1) can only be
due to a selection process exerted on certain amino acids at certain positions of a true protein in order
to improve, preserve, or eliminate its function.
364
The analysis of proteins whose creation appears most recent (node 9 in Fig. 4) indicates a negative
selection pressure whose (probably overestimated) average value (0.58) could correspond to
functions slightly increasing the pandoravirus fitness. However, the interpretation of the selection
pressure in the context of a function is a problem here, since we previously pointed out that it is
highly unlikely that a newly created small protein will have a function (enzymatic, regulatory or
structural) in the first place, even less a beneficial one (Legendre et al., 2018).
Thus, we propose an alternative interpretation, based on the knowledge accumulated over the years
on the principles governing the stability of proteins. According to our scenario, most new proteins are
from the translation of pre-existing non-coding "protogenes" (see 3.4) and the rest from DNA inserts
of unknown origin (such as pqer_cds_665 and pqer_cds_673). It is then unlikely that many of these
new proteins lacking an evolutionary history will adopt a globular fold (Watters et al. 2006;
Monsellier & Chiti, 2007; Tanaka et al., 2010). Most will be toxic by causing nonspecific
aggregations within the host-virus proteome and be quickly eliminated through the reversal of these
newborn viral genes to an untranscribed/untranslated state. In contrast, proteins whose folded 3-D
structures do not prove to be toxic will enter an evolutionary process involving a negative selection
pressure promoting the conservation of the amino acids responsible for their stability. Their
proportion has been estimated at about 34% for 100-200 residue proteins (Miao et al., 2004). In
absence of an initial function, the other positions/residues will evolve in a neutral manner. Once
"fixed", the new genes will exhibit an average selection pressure lower than one, combining 1/3 of
negative selection with 2/3 of neutrality. Following many studies that have concluded that efficient
folding (Watters et al., 2006) and prevention of aggregation are important drivers of protein evolution
(see Monsellier & Chiti, 2007), our scenario predicts that many genes specific to pandoraviruses
should encode proteins not increasing their fitness, but whose stable 3-D structures may eventually
serve as innovative platforms for new functions.
389
In conclusion, the detailed comparative analysis of *P. celtis* with its very close relative *P. quercus*
enabled us to estimate the contributions of various microevolutionary processes to the steady increase
of the pan-genome of the proposed Pandoraviridae giant virus family. We first showed that large-
scale genomic rearrangements (segmental duplications, translocations) are associated to transposable
elements of the hAT family, widespread in metazoans, but until now unique to this family of viruses.
However, these mobile elements mostly appear to shuffle pandoravirus genes between strains,
without creating new ones or promoting host-to-virus horizontal gene transfers. In contrast with the
popular view that horizontal gene transfer plays an important role in the evolution of large DNA

viruses (Yutin & Koonin (2013); Koonin et al. 2015; Schulz et al., 2017), this is definitely not the
main cause of the inflation of the Pandoraviridae genomes, as we previously argued (Claverie &
Abergel, 2013; Philippe et al., 2013; Abergel et al., 2015; Legendre et al., 2018). Finally, we also
found that locally repeated regions are the siege of a competition between tandem duplication and
gene deletion, locally reshaping the genomes without contributing to net genetic innovation (Fig. S3).

In continuity with our previous work on more distant pandoravirus strains, we found that the 33
protein-coding genes born since the divergence between the very close *P. quercus* and *P. celtis*
strains derived from preexisting non-coding sequences or small DNA segments of unknown origin
inserted at randomly interspersed locations. These non-coding regions, turned on in a given
pandoravirus strain, could remain silent or diverge beyond recognition in others. These results add
strong support to the constant *de novo* creation of proteins, few of which are retained with a little
impact on the virus fitness until eventually acquiring a selectable function.

5   Author Contributions

411 CA, ML, JMC, and YC designed the experiments. JMA, AL, NP, SJ, and YC contributed to the data
412 and performed the experiments. ML, SN, NTT, OP, CA and JMC analyzed the data. ML, CA, and
413 JMC wrote the manuscript.

6   Funding

This work was partially supported by the French National Research Agency (ANR-14-CE14-0023-
01), France Genomique (ANR-10-INSB-01-01), Institut Français de Bioinformatique
(ANR–11–INSB–0013), the Fondation Bettencourt-Schueller (OTP51251), by a DGA-MRIS
scholarship, and by the Provence-Alpes-Côte-d'Azur région (2010 12125). Proteomic experiments
were partly supported by the Proteomics French Infrastructure (ANR-10-INBS-08-01) and Labex
GRAL (ANR-10-LABX-49-01). We thank the support of the discovery platform and informatics
group at EDyP and the PACA-Bioinfo platform.
422

7   Conflict of Interest Statement

424 The authors declare that the research was conducted in the absence of any commercial or financial
425 relationships that could be construed as a potential conflict of interest.

8   Acknowledgments

427 We kindly thank Hua-Hao Zhang for providing us with the hAT-transposon sequence he identified in
428 the *Acanthamoeba castellanii* genome.

9   Supplementary Material

431 The Supplementary Material for this article is appended.

432 **Figure S1.** TEM images of ultrathin sections of *P. celtis* and *P. quercus* viral particles.

434   **Figure S2.** Location of the hAT transposases in the S0, S1, and S3 segments and transposons
435   flanking sequences in P. celtis. The location of the tranposase gene is indicated in red**.**
436
437   **Figure S3.** Details of a tandem repeat cluster of fascin-domain containing genes.
438

10   References

Abergel, C., Legendre, M., Claverie, J.M. (2015) The rapidly expanding universe of giant viruses:
Mimivirus, Pandoravirus, Pithovirus and Mollivirus. FEMS Microbiol Rev. 39(6), 779-96. doi:
10.1093/femsre/fuv037.

Aherfi, S., Andreani, J., Baptiste, E., Oumessoum, A., Dornas, F.P., Andrade, A.C.D.S.P., et al.
(2018). A large open pangenome and a small core genome for giant pandoraviruses. Front.
Microbiol. 9, 1486. doi: 10.3389/fmicb.2018.01486.

Aherfi, S., Colson, P., La Scola, B., Raoult, D. (2016) Giant Viruses of Amoebas: An Update. Front
Microbiol. 7, 349. doi: 10.3389/fmicb.2016.00349.

Antwerpen, M.H., Georgi, E., Zoeller, L., Woelfel, R., Stoecker, K., Scheid, P. (2015) Whole-
genome sequencing of a pandoravirus isolated from keratitis-inducing acanthamoeba. Genome
Announc. 3(2). pii: e00136-15. doi:10.1128/genomeA.00136-15.

Atkinson, P.W. (2015) hAT Transposable Elements. Microbiol Spectr. 3(4). doi:
10.1128/microbiolspec.MDNA3-0054-2014.

Chin, C.S., Alexander, D.H., Marks, P., Klammer, A.A., Drake, J., Heiner C., et al. (2013)
Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. Nat
Methods. 10(6), 563-9. doi: 10.1038/nmeth.2474.

Claverie, J.M., Abergel, C. (2013) Open questions about giant viruses. Adv Virus Res. 85, 25-56.
doi: 10.1016/B978-0-12-408116-1.00002-1.

Claverie, J.M., Abergel, C., Legendre M. (2018). Med Sci (Paris), in press.

Corradi, N., Pombert, J.F., Farinelli, L., Didier, E.S., Keeling, P.J. (2010) The complete sequence of
the smallest known nuclear genome from the microsporidian Encephalitozoon intestinalis. Nat
Commun. 1, 77. doi: 10.1038/ncomms1082.

Emms, D.M., Kelly, S. (2015) OrthoFinder: solving fundamental biases in whole genome
comparisons dramatically improves orthogroup inference accuracy. Genome Biol. 16, 157. doi:
10.1186/s13059-015-0721-2.

Fabre, E, Jeudy, S, Santini, S, Legendre, M, Trauchessec, M, Couté, Y, et al. (2017) Noumeavirus
replication relies on a transient remote control of the host nucleus. Nat Commun. 8, 15087. doi:
10.1038/ncomms15087.

Floriano, A.M., Castelli, M., Krenek, S., Berendonk, T.U., Bazzocchi, C., Petroni, G., et al. (2018)
The Genome Sequence of "Candidatus Fokinia solitaria": Insights on Reductive Evolution in
Rickettsiales. Genome Biol Evol. 10(4), 1120-1126. doi: 10.1093/gbe/evy072.

Hoang, D.T., Chernomor, O., von Haeseler, A., Minh, B.Q., Vinh, L.S. (2018) UFBoot2: Improving
the Ultrafast Bootstrap Approximation. Mol Biol Evol. 35(2), 518-522. doi:
10.1093/molbev/msx281.

Katoh, K., Misawa, K., Kuma, K., Miyata, T. (2002) MAFFT: a novel method for rapid multiple
sequence alignment based on fast Fourier transform. Nucleic Acids Res. 30(14), 3059-66.

Koonin, E.V., Krupovic, M., Yutin, N. (2015) Evolution of double-stranded DNA viruses of
eukaryotes: from bacteriophages to transposons to giant viruses. Ann N Y Acad Sci. 1341, 10-24.
doi: 10.1111/nyas.12728.

Krumsiek, J., Arnold, R., Rattei, T. (2007) Gepard: a rapid and sensitive tool for creating dotplots on
genome scale. Bioinformatics 23(8), 1026-8. doi: 10.1093/bioinformatics/btm039.

Latorre, A., Manzano-Marín, A. (2017) Dissecting genome reduction and trait loss in insect
endosymbionts. Ann N Y Acad Sci. 1389(1), 52-75. doi: 10.1111/nyas.13222.

Lopez-Madrigal, S., Latorre, A., Porcar, M., Moya, A., Gil, R. (2011) Complete genome sequence of
'Candidatus Tremblaya princeps' strain PCVAL, an intriguing translational machine below the
living-cell status. J; Bacteriol; 193, 5587–5588. doi: 10.1128/JB.05749-11.

Legendre, M., Fabre, E., Poirot, O., Jeudy, S., Lartigue, A., Alempic, J.M., et al. (2018). Diversity
and evolution of the emerging Pandoraviridae family. Nat. Commun. 9, 2285. doi: 10.1038/s41467-
018-04698-4.

Legendre, M., Lartigue, A., Bertaux, L., Jeudy, S., Bartoli, J., Lescot, M., et al. (2015) In-depth study
of Mollivirus sibericum, a new 30,000-y-old giant virus infecting Acanthamoeba. Proc Natl Acad Sci
U S A. 112, E5327-35. doi: 10.1073/pnas.1510795112.

McCutcheon JP, Moran NA (2011) Extreme genome reduction in symbiotic bacteria. Nat. Rev.
Microbiol. 10, 13–26. doi: 10.1038/nrmicro2670.

Marchler-Bauer, A., Bryant, S.H. (2004) CD-Search: protein domain annotations on the fly. Nucleic
Acids Res. 32, W327-31 doi : 10.1093/nar/gkh454.

Miao, J., Klein-Seetharaman, J., Meirovitch, H. (2004) The optimal fraction of hydrophobic residues
required to ensure protein collapse. J Mol Biol. 344(3), 797-811. doi: 10.1016/j.jmb.2004.09.061.

Monsellier, E., Chiti, F. (2007) Prevention of amyloid-like aggregation as a driving force of protein
evolution. EMBO Rep. 8(8), 737-42. doi: 10.1038/sj.embor.7401034.

Philippe, N., Legendre, M., Doutre, G., Couté, Y., Poirot, O., Lescot, M., et al. (2013)
Pandoraviruses: amoeba viruses with genomes up to 2.5 Mb reaching that of parasitic eukaryotes.
Science 341, 281-6. doi: 10.1126/science.1239181.

Remmert, M., Biegert, A., Hauser, A., Söding, J. (2011) HHblits: lightning-fast iterative protein
sequence searching by HMM-HMM alignment. Nat. Methods. 9(2), 173-5. doi: 10.1038/nmeth.1818.

Rice, P., Longden, I., Bleasby, A. (2000) EMBOSS: The European Molecular Biology Open
Software Suite. Trends Genet. 16(6), 276-7. doi:10.1016/S0168-9525(00)02024-2

Rocha, E.P.C., Smith, J.M., Hurst, L.D., Holden, M.T., Cooper, J.E., Smith, N.H., et al. (2006)
Comparisons of dN/dS are time dependent for closely related bacterial genomes. J Theor Biol. 239,
226-35.

Sayers, E.W., Agarwala, R., Bolton, E.E., Brister, J.R., Canese, K., Clark, K., et al. (2018) Database
resources of the National Center for Biotechnology Information. Nucleic Acids Res.
doi:10.1093/nar/gky1069.

Schulz, F., Yutin, N., Ivanova, N.N., Ortega, D.R., Lee, T.K., Vierheilig, J., et al. (2017) Giant
viruses with an expanded complement of translation system components. Science 356(6333), 82-85.
doi: 10.1126/science.aal4657.

Szilágyi, A, Skolnick, J. (2006) Efficient prediction of nucleic acid binding function from low-
resolution protein structures. J Mol Biol. 358(3), 922-33. doi: 10.1016/j.jmb.2006.02.053.

Tanaka, J., Doi, N., Takashima, H., Yanagawa, H. (2010) Comparative characterization of random-
sequence proteins consisting of 5, 12, and 20 kinds of amino acids. Protein Sci. 19(4), 786-95. doi:
10.1002/pro.358.

Yang, Z. (2007) PAML 4: phylogenetic analysis by maximum likelihood. Mol. Biol. Evol. 24(8),
1586-91. doi: 10.1093/molbev/msm088.

Watters AL, Deka P, Corrent C, Callender D, Varani G, Sosnick T, et al. (2007). The highly
cooperative folding of small naturally occurring proteins is likely the result of natural selection. Cell
128(3), 613-24. doi: 10.1016/j.cell.2006.12.04.

Yutin, N., Koonin, E.V. (2013) Pandoraviruses are highly derived phycodnaviruses. Biol Direct. 8,
25. doi: 10.1186/1745-6150-8-25..

Zhang, H.H., Zhou, Q.Z., Wang, P.L., Xiong, X.M., Luchetti, A., Raoult, D., et al. (2018)
Unexpected invasion of miniature inverted-repeat transposable elements in viral genomes. Mob DNA
9, 19. doi: 10.1186/s13100-018-0125-4.

11   Data Availability Statement

P. celtis annotated genome is deposited in GenBank under Accession n° MK174290.

12   Tables

537   **Table 1.** *P. celtis* and *P. quercus* unique protein-coding genes

Gene #	Size (aa)	Most ancestral sequence trace	Prediction	Median RNAseq read coverage
pclt_cds_11	69	Since node 5	DNA binding	473
pclt_cds_308	140	Since node 8	no	669
pclt_cds_350	121	Since node 9	DNA binding	32
pclt_cds_376	74	Since node 8	DNA binding	99
pclt_cds_649	77	Since node 8	DNA binding	221
pclt_cds_725	104	none	DNA binding	9
pclt_cds_764	188	none	DNA binding	11
pclt_cds_870	205	Since node 2	DNA binding	1,362
pclt_cds_898	69	Since node 5	no	1,302
pclt_cds_995	133	Since node 2	DNA binding	6
pclt_cds_1081	218	Since node 8	DNA binding	7,913
pclt_cds_1084	114	Since node 8	DNA binding	72
pqer_cds_6	76	Since node 8	DNA binding	303
pqer_cds_13	117	Since node 5	DNA binding	48
pqer_cds_17	82	Since node 2	DNA binding	403
pqer_cds_53	85	Since node 8	DNA binding	41
pqer_cds_143	93	Since node 8	DNA binding	160
pqer_cds_151	71	Since node 9	DNA binding	10
pqer_cds_203	101	Since node 8	DNA binding	528
pqer_cds_350	94	none	DNA binding	127
pqer_cds_474	74	Since node 9	DNA binding	111
pqer_cds_486	146	Since node 2	DNA binding	61
pqer_cds_665	114	none	DNA binding	1,929
pqer_cds_673	383	none	DNA binding	630
pqer_cds_685	124	Since node 8	no	95
pqer_cds_736	121	Since node 9	DNA binding	66
pqer_cds_875	136	none	no	2,005
pqer_cds_876	74	Since node 2	no	1,508
pqer_cds_877	78	none	DNA binding	305
pqer_cds_878	224	none	DNA binding	223
pqer_cds_1061	152	Since node 8	DNA binding	14,125
pqer_cds_1178	84	Since node 9	DNA binding	149
pqer_cds_1183	158	Since node 2	DNA binding	189

13 Figure legends

Figure 1.

Phylogenetic tree of known, fully characterized Pandoraviruses. The tree was computed from the
Pandoraviruses' core genes (see Materials and Methods for details). The estimated bootstrap values
were all equal to 1 and thus not reported. Internal nodes are labeled according to their depth in the tree.

Figure 2.

Dot plot (nucleotide) comparison of the *P. celtis vs. P. quercus* genomes. Computations and display
were generated using Gepard (Krumsiek et al., 2007). Specific regions, labeled P, T, S_0, S_1 and S_2 are
described in the main text.

Figure 3.

Pandoraviridae pan-genome and core-genome. The boxplot represents the number of protein
clusters as a function of the number of sequenced genomes in all possible combinations. The whiskers
correspond to the extreme data points.

Figure 4.

Estimated selection pressure acting on Pandoravirus genes as a function of their ancestry. Shown
is the mean of dN/dS ratios (filled bars) for genes that are unique to the pandoravirus strains below a
given node in their phylogeny tree (see Fig. 1). Error bars correspond to standard deviations. Bars are
colored according to the depth of the node in the tree. As a control, we calculated the dN/dS of
Pandoravirus core genes (empty bars). For a given node, we considered pairs of orthologous genes
whose common ancestor correspond to that node.

Figure S1.

TEM image of an ultrathin section of *P. celtis* and *P. quercus* viral particles. The structures of the
particles of the other Pandoraviridae strains do not exhibit any noticeable difference.

Figure S2.

Details of the P. celtis hAT transposons. Details of the S_0, S_1, and S_2 segments highlighted in Fig.
2. The genomic coordinates of the transposons in *P. celtis* (horizontal) and their homologous
locations in *P. quercus* (vertical) are indicated. The cognate *P. celtis* transposase gene name and its
approximate location is indicated for each transposon. Other genes are not represented. Homologous
segments are missing in *P. quercus* for S_0 and S_2 (See section 3.1). The sequences of the transposon
boundaries are shown above the dot-plots. The TIRs (Terminally Inverted Repeats) are highlighted
in black. Conserved TIRs and TSDs (Target Site Duplications) are depicted using arrows.

Figure S3.

Details of a tandem repeat cluster of fascin-domain containing genes. See section 3.2 for
comments. The dot-plot (horizontal: *P. celtis*, vertical *P. quercus*) and the corresponding
phylogenetic tree of the genes illustrate the well-ordered dynamics of this highly repeated regions,
alternating expansion and deletions. The evolutionary history of the fascin-domain containing

proteins was inferred using the Neighbor-Joining method as implemented in Mega (Kumar S, et al.
(2018) MEGA X: Molecular Evolutionary Genetics Analysis across computing platforms. Mol. Biol.
Evol. 35:1547-1549.). The evolutionary distances were computed using the Poisson correction
method. The analysis involved 31 amino acid sequences. All positions containing gaps and missing
data were eliminated. There were 323 positions in the final dataset.

Supplementary Figures S1-S3

Figure S1.

TEM image of an ultrathin section of *P. celtis* and *P. quercus* viral particles. The structures of the particles of the other Pandoraviridae strains do not exhibit any noticeable difference.

Figure S2.

Details of the P. celtis hAT transposons. Details of the S_0, S_1, and S_2 segments highlighted in Fig. 2. The genomic coordinates of the transposons in *P. celtis* (horizontal) and their homologous locations in *P. quercus* (vertical) are indicated. The cognate *P. celtis* transposase gene name and its approximate location is indicated for each transposon. Other genes are not represented. Homologous segments are missing in *P. quercus* for S_0 and S_2 (See section 3.1). The sequences of the transposon boundaries are shown above the dot-plots. The TIRs (Terminally Inverted Repeats) are highlighted in black. Conserved TIRs and TSDs (Target Site Duplications) are depicted using arrows.

S0_transposon `TAATCCT`GGTCAACGGCCAGCCGGCCCAGGGCCGGCCCATTTTGAAACGGGCC . . . CAAAATCTGCGGCCAGCCGGCCAGCCGCTAAGCCACGGGCCATTGGCAC`AGCATTA`

S1_transposon `TAACGCTGGGCAA`CGGCTAGCCGGCTCGGTCCTGGCTAAGGGTCGGACTAACC . . . AAAATGTGGCGGCTAGCCGGCTAGCCGGCTTACGCGTAGCCAT`T-GCCCAGCGTTA`

S2_transposon `TAACGCTGGGCAACGGCT`AGCCGGCTCGGTCCTGGCTAAGGGTCGGACTAACC . . . AACAGGTGGCGGCTAGCCGGTTAGCCGGCTAACGGTGA`GCCGTT-GCCCAGCGTTA`

Figure S3.

Details of a tandem repeat cluster of fascin-domain containing genes. See section 3.2 for comments. The dot-plot (horizontal: *P. celtis*, vertical *P. quercus*) and the corresponding phylogenetic tree of the genes illustrate the well-ordered dynamics of this highly repeated regions, alternating expansion and deletions. The evolutionary history of the fascin-domain containing proteins was inferred using the Neighbor-Joining method as implemented in Mega (Kumar S, et al. (2018) MEGA X: Molecular Evolutionary Genetics Analysis across computing platforms. Mol. Biol. Evol. 35:1547-1549.). The evolutionary distances were computed using the Poisson correction method. The analysis involved 31 amino acid sequences. All positions containing gaps and missing data were eliminated. There were 323 positions in the final dataset.