Motivation and Cognitive Control in Depression

Ivan Grahek[1][*], Amitai Shenhav[2], Sebastian Musslick[3], Ruth M. Krebs[4], & Ernst H.W. Koster[1]

1. *Department of Experimental Clinical and Health Psychology, Ghent University, Henri Dunantlaan 2, B-9000, Ghent, Belgium*

2. *Department of Cognitive, Linguistic, & Psychological Science and Brown Institute for Brain Science, Brown University, Providence, RI 02912, USA*

3. *Princeton Neuroscience Institute, Princeton University, Princeton, NJ 07001, USA*

4. *Department of Experimental Psychology, Ghent University, Henri Dunantlaan 2, B-9000, Ghent, Belgium*

* Corresponding author at: Department of Experimental Clinical and Health Psychology, Ghent University, Henri Dunantlaan 2, B-9000, Ghent, Belgium. Phone: +32 (0)9 264 94 46; Fax: +32 (0)9 264 64 89

E-mail address: ivan.grahek@ugent.be

Abstract

Depression is linked to deficits in cognitive control and a host of other cognitive impairments arise as a consequence of these deficits. Despite of their important role in depression, there are no mechanistic models of cognitive control deficits in depression. In this paper we propose how these deficits can emerge from the interaction between motivational and cognitive processes. We review depression-related impairments in key components of motivation along with new cognitive neuroscience models that focus on the role of motivation in the decision-making about cognitive control allocation. Based on this review we propose a unifying framework which connects motivational and cognitive impairments in depression. This framework is rooted in computational models of cognitive control and offers a mechanistic understanding of cognitive control deficits in depression.

Keywords: depression; cognitive control; anhedonia; motivation; reward anticipation; effort; controllability of environment.

Introduction

Depression[1] profoundly influences the way in which we process information and think about ourselves, others, and the world around us. An individual suffering from depression will take a longer time to disengage from the processing of negative information and will experience difficulties in suppressing irrelevant thoughts, or shifting attention from one task to another in order to reach a goal. Such issues will make it difficult for that individual to regulate emotions and adapt to the changing environment. This is why cognitive processes are a crucial target for understanding and treating depression (Clark and Beck, 2010; Kaser et al., 2017).

Depression is characterized by impairments in attention, memory, and cognitive control (Millan et al., 2012). Cognitive control deficits are related to central features of depression such as concentration and memory problems and a host of other cognitive impairments and biases arise as a consequence of these deficits (Disner et al., 2011; Gotlib and Joormann, 2010). Cognitive control is crucial in motivated, goal-directed behavior. It represents a set of processes that allow for the flexible adaptation of cognition and behavior in accordance with our current goals (Botvinick and Cohen, 2014; Friedman and Miyake, 2017; Shenhav et al., 2013). For example, cognitive control is necessary if we want to inhibit negative thoughts and shift our attention to a new task. Impairments in such control processes are found across a wide range of psychiatric disorders (Millan et al., 2012), and they have been consistently linked with depressive symptoms (Snyder, 2013). However, our understanding of cognitive control in depression is limited in several important ways.

Current understanding of cognitive control in depression is predominately descriptive and research is focused on detecting deficits in specific cognitive processes, such as the inhibition of

---

[1] Throughout this paper we use the term depression to denote the Major Depressive Disorder.

negative material (for a review see: Grahek et al., 2018). In spite of the important progress in charting cognitive control deficits related to depressive symptoms, the origin of these deficits remains poorly understood. It is not known why deficits in cognitive control develop, or how they are maintained. Most of the existing models view cognitive control deficits in depression as the inability to exert control and do not offer mechanisms through which these deficits emerge. Currently there is a strong need for the development of a more mechanistic account of cognitive control deficits in depression and recent progress in clinical and cognitive neuroscientific research provides building blocks for such an understanding. We believe that the recent developments in research on motivation and cognitive control, as well as the development of computational theories of cognitive control, can contribute to the mechanistic understanding of control deficits in depression.

In this paper we first review disparate literatures on motivation and cognitive control in depression. Further, we describe computational models of cognitive control and demonstrate how they can be used to link motivation and cognition in depression. On the basis of this review, we propose a framework in which cognitive control deficits in depression arise from alterations in crucial components of motivation: reward anticipation, effort costs, and estimates of environment controllability. This view offers the possibility for re-conceptualizing depression-related cognitive control deficits. Instead of an *inability* to employ control, we propose that control deficits can be viewed as changes in the decision-making process underlying cognitive control allocation. This decision-making process relies on crucial components of motivation: reward anticipation, effort, and estimates of the ability to control the environment. We use simulations of two cognitive tasks to demonstrate how this framework can be used to derive behavioral predictions about the impact of motivational impairments on cognitive control.

Cognitive Control in Depression

Deficits in cognitive control have not only been documented in clinically depressed individuals (Snyder, 2013), but also in patients in remission (Demeyer et al., 2012; Levens and Gotlib, 2015), and in at-risk populations (Derakshan et al., 2009; Owens et al., 2012). There is emerging evidence that cognitive remediation training aimed at improving cognitive control processes reduces depressive symptoms (for a review see: Koster, Hoorelbeke, Onraedt, Owens, & Derakshan, 2017). At the neural level, depressive symptoms have been linked to changes in the activity of the dorsolateral prefrontal cortex (dlPFC) and anterior cingulate cortex (ACC; Davidson, Pizzagalli, Nitschke, & Putnam, 2002; Gotlib & Hamilton, 2008; Pizzagalli, 2011). Multiple authors have proposed that the reduced activity in these regions is related to the diminished ability of depressed individuals to employ cognitive control (Disner et al., 2011; Joormann, Yoon, & Zetsche, 2007).

Cognitive control processes are considered to be an important vulnerability factor for depression. Impairments in attention, interpretation, and memory arise as a consequence of control deficits (Gotlib and Joormann, 2010; Millan et al., 2012; Siegle et al., 2007). Lowered levels of cognitive control increase and sustain depressive symptoms via their proximal links with emotion regulation strategies such as rumination (Everaert, Grahek, & Koster, 2017; Joormann & Vanderlind, 2014; Whitmer & Gotlib, 2013). Research on cognitive control in depression has been focused on charting deficits in different cognitive control processes. Specific deficits in processes such as inhibition, shifting, and updating have been documented (Joormann & Tanovic, 2015). These deficits are commonly thought of as the *inability* to inhibit certain thoughts or stimuli, shift attention away from them, or update the contents of working memory. However, it is important to note that not all accounts of cognitive impairments in depression postulate an

inability. For example, the cognitive-initiative account of memory in depression focuses on changes in initiative – a concept close to motivation – to explain memory impairments in depression (Hertel, 2000, 1994).

In a recent analysis of theoretical models of cognitive control in depression (Grahek et al., 2018) we identified three main conceptual problems in the field: (1) the use of descriptive models of cognitive control, (2) the reliance on describing the impairments instead of searching for mechanisms, and (3) the lack of integration between cognitive, motivational, and emotional impairments. These issues are hindering further progress in understanding how and why cognitive control is impaired in depression. In order to overcome some of the problems that we have outlined in our earlier work, in this paper we propose an integrated framework that links alterations in motivational processes with cognitive control deficits. This framework allows us to move away from the view that cognitive control deficits in depression stem from an inability to exert control. Instead, we will argue for a view in which the deficits arise as a result of altered expectations about the value of exerting control.

Components of Motivation in Depression

Cognitive impairments in depression are closely linked to impairments in emotional and motivational processes (Crocker et al., 2013). A wealth of depression research has focused on the relationship between cognitive and emotional processes. This approach has led to insights in key deficits related to the disengagement from emotionally negative material (Koster et al., 2011) and the ability to deploy cognitive control over emotional material (Joormann, 2010). While the processing of negative material and the presence of negative affect have been studied in relation to cognitive control impairments in depression, motivational deficits remain largely unexplored in this context. This is why we focus this paper on the link between motivation and cognition in

depression. The links between motivation and cognitive processing of emotional material our out of the scope of the current paper.

Motivation is goal-directed when effort is invested in order to bring about desired outcomes (Braver et al., 2014). Here we will focus on components of motivation that are relevant for goal-directed behavior because of their relevance for cognitive control processes. Motivated goal-directed behavior is flexible and sensitive to the current state of the individual and the environment. Two types of representations are crucial in driving this type of behavior: 1) action-outcome contingencies, and 2) the value of potential outcomes (Balleine and O'Doherty, 2010; Dickinson, 1985; Wood and Rünger, 2016). Action-outcome contingencies represent the probability that an action will result in a desired outcome. We will refer to these representations as *outcome controllability* (one's estimate of their ability to control outcomes in an environment) and *outcome value* (the expected reinforcement - total reward and/or punishment - for reaching an outcome). The third concept that we will consider is effort, a variable that is central to the study of motivation. Effort represents the intensification of physical or mental activity needed to reach a goal (Inzlicht et al., 2018; Kurzban et al., 2013). This intensification comes with a cost, and we will refer to the effort requirements for reaching an outcome as *effort costs*.

Goal-directed motivated behavior emerges with the integration of these three components. For example, imagine a person working long hours in order to get a promotion at work. This behavior is motivated and goal-directed because this person believes that working hard (high effort) will lead to the promotion (high outcome controllability), and the promotion is desired (high outcome value). As we review next, there is evidence that each of these three components of motivation can be impaired in depression (Barch et al., 2015; Griffiths et al., 2014).

Outcome Controllability

The classic paradigm for investigating the role of outcome controllability in response to stressors was developed by Seligman and Maier (1967). They demonstrated that animals who were subjected to uncontrollable stressors (inescapable shocks) exhibited passivity, a phenomenon they referred to as learned helplessness. They found that controllable stressors (escapable shocks) did not induce learned helplessness. Overall, uncontrollable stressors were found to induce responses that resemble some of the symptoms of depression (Maier, 1984; Maier and Watkins, 2005). Further work by Maier and colleagues (2006) revealed that animals were able to detect the possibility of control over their environment. While animals have a default reaction of passivity when experiencing stress, this default response can be overcome by learning that the stressors are controllable. The medial prefrontal cortex (mPFC) in rodents detects whether a stimulus is controllable and inhibits the default passivity in response to shocks. In this way, the ability to exert control over the environment serves as a protective factor against negative behavioral and physiological effects of stress (Maier and Seligman, 2016). Seligman and colleagues demonstrated that learned helplessness is also observed in humans, and suggested that it has significant relevance for understanding depression and related disorders (Abramson, Seligman, & Teasdale, 1978; Maier & Seligman, 2016; Seligman, 1972).

The idea that environmental controllability is crucial for one's health, security, and well-being is supported by several other lines of research (Leotti et al., 2010). For instance, Moscarello and Hartley (2017) have recently proposed that goal-directed behavior is strongly influenced by estimates of agency. The authors propose that animals and humans infer their ability to control their environment by ascertaining the relationship between their actions and motivationally significant outcomes. These estimates are generalized and determine the probability of being able to exert control in a novel environment. Estimates of agency are used to calibrate ongoing

behaviors. If the estimates of agency are high (i.e., high probability of being able to control the environment), goal-directed behavior is promoted. If the estimates are low, behavior is more likely to rely on an innate reactive repertoire. In this way, controllability of the environment, inferred from previous learning, is crucial for promoting either goal-directed or habitual behavior (Liljeholm, Tricomi, O'Doherty, & Balleine, 2011; Miller, Shenhav, & Ludvig, in press). Lowered estimates of outcome controllability, resulting in alterations of goal-directed behavior, could be an important factor in depression.

Recent theoretical frameworks of depression also emphasize the importance of outcome controllability. For example, Pizzagalli (2014) has proposed a model of anhedonia in depression in which stress plays a central role in the development of anhedonic symptoms. According to Pizzagalli, certain stressors, especially if they are uncontrollable, induce anhedonic behavior by causing dysfunctions in mesolimbic dopamine pathways crucial for motivated behavior. De Raedt and Hooley (2016) have argued that individual's expectancies about their ability to cope with future negative events play a crucial role in depression. These expectancies are proposed to be formed based on previous coping experiences and are one factor that determines the actual ability to cope with stressors when they occur. Notably, these authors suggest that expectancies influence the ability to cope with stressors by modulating the proactive allocation of cognitive control prior to encountering a stressful situation. In sum, altered estimates of outcome controllability seem to be an important factor contributing to the levels of depressive symptoms.

Outcome Value

Representations of outcome value are also strongly altered in depression. In recent years, motivational impairments in depression have received significant attention and a more fine-grained view of these impairments is emerging (Pizzagalli, 2014; Treadway and Zald, 2013).

Anhedonia is one of the two core symptoms of depression and is defined as a loss of pleasure in previously enjoyable activities or a loss of interest in pursuing them (DSM-5; American Psychiatric Association, 2013). Anhedonia is a good predictor of antidepressant treatment success (Uher et al., 2012), the course of depression (Spijker et al., 2001; Wardenaar et al., 2012), and the time to remission and number of depression-free days after antidepressant treatment (McMakin et al., 2012). In spite of the importance of anhedonia in depression, the conceptualization and measurement of anhedonia has been heterogeneous and inconsistent (for a discussion see: Rizvi, Pizzagalli, Sproule, & Kennedy, 2016). Traditionally, anhedonia has been primarily viewed as an impairment in consummatory pleasure (i.e. liking rewards when they are obtained). However, the evidence for impairments in consummatory pleasure in depression is mixed (Barch et al., 2015; Treadway and Zald, 2013).

Animal models suggest that the mesolimbic dopamine system is selectively involved in reward motivation, but not in hedonic responses when rewards are gained (Haber and Knutson, 2010; Salamone and Correa, 2012). These insights have stimulated research on anhedonia in depression and schizophrenia, revealing impairments in motivation that are not related to hedonic responses. As a result, an emerging, more nuanced account of motivation in anhedonia, emphasizes multiple reward processing deficits such as anticipation of rewards, reinforcement learning, effort expenditure, and value-based decision making (Barch and Dowd, 2010; Der-Avakian and Markou, 2012; Pizzagalli, 2014; Romer Thomsen et al., 2015; Strauss and Gold, 2012; Zald and Treadway, 2017). These deficits point toward changes in outcome value. They influence how outcome values are learned, anticipated, and translated into behavior.

Depression is related to a number of reward processing deficits (for reviews see: Barch et al., 2015; Pizzagalli, 2014; Treadway & Zald, 2011). Converging evidence from self-report

studies, behavioral tasks, physiological, and neuroimaging experiments suggests that depression, especially in the presence of anhedonia, is linked to reduced anticipation of rewards and impaired implicit reinforcement learning. For example, monetary incentive delay tasks (Knutson et al., 2000) were developed in order to decompose anticipatory (e.g., the period after notifying participants about a possible reward) and consummatory (e.g., receipt of a monetary reward) aspects of reward processing. In these tasks depressed individuals, relative to healthy controls, display reduced behavioral and neural responses in anticipation of rewards (Pizzagalli, 2014). The literature on motivational impairments in depression suggests that depressed individuals, especially in the presence of anhedonic symptoms, assign value outcomes in a manner that differs from healthy controls. This can be caused by anticipating lower payoffs and/or impaired reinforcement learning.

Effort costs

Another important factor influencing motivated behavior is effort that needs to be expended in order to reach a desired outcome. Effort involves the expected cost necessary to reach an outcome. This cost is weighed against expected benefits in order to choose which actions to pursue (Wallis and Rushworth, 2014). Reduced effort exertion is associated with multiple mental disorders (Culbreth et al., 2017; Salamone et al., 2016). Clinical studies have focused on the exertion of physical effort in order to obtain rewards, demonstrating that anhedonia in depression is related to reduced effort exertion (for an excellent review see: Zald & Treadway, 2017). In order to investigate this process, Treadway and colleagues developed the Effort Expenditure for Rewards Task (EEfRT; Treadway, Bossaller, Shelton, & Zald, 2012), which involves having participants choose how much effort they want to expend in order to gain varying amounts of reward. Using this task, the authors have demonstrated that depressed

individuals are less willing to exert effort than healthy controls (see also: Cléry-Melin et al., 2011; Yang et al., 2014). However, the research on effort exertion in anhedonia and depression has largely been focused on physical effort while ignoring cognitive effort.

Although cognitive effort has been an important topic of research for a long time (Kahneman, 1973), in recent years there has been an upsurge in cognitive and neuroscience research on this topic (Kool and Botvinick, 2018; Westbrook and Braver, 2015). More specifically, research on cognitive control has focused on the role of effort costs in decision-making about control allocation (for a recent review see: Shenhav et al., 2017). Cognitive control processes require more effort than automatic ones, and effort needs to be expended in order to override automatic processes in order to reach a goal. Research is starting to demonstrate that there are individual differences in the exertion of this type of effort (Westbrook et al., 2013). To date, there are not a lot of studies on cognitive effort in depression. However, a recent study has demonstrated the inverse relationship between depressive symptoms and a questionnaire-based measure of the willingness to exert cognitive effort (Marchetti et al., 2018). Also, the first experimental study of cognitive effort in depression revealed similar results to the results obtained with physical effort (Hershenberg et al., 2016).

Cognitive Control as a Process Reliant on Motivation

Controllability, value, and effort of a desired outcome depend on previous learning. They have been studied extensively in the context of animal and human learning at both behavioral and neural levels (for an overiew see: Daw and O'Doherty, 2014). The importance of these components of motivated behavior has also been recognized in other fields of psychology and neuroscience. For example, in social psychology the concepts of self-efficacy (Bandura, 2001, 1977) and locus of control (Rotter, 1966) are closely linked to controllability. Also, concepts of

feasibility and desirability of goals (Oettingen and Gollwitzer, 2001) correspond to the concepts

of controllability and outcome value. Similar concepts can be found in psychology of motivation

(Atkinson, 1957; Wigfield and Eccles, 2000) and work psychology (Bonner and Sprinkle, 2002).

The role of motivation is now increasingly recognized in cognitive psychology and cognitive

neuroscience.

Automatic or habitual responding can be suitable for a number of everyday situations, but

for tasks that are more novel, uncertain, or complex, individuals need to engage cognitive control.

This set of processes helps to overcome automatic response tendencies in favor of controlled

modes of information processing and behavior. This allows for the coordination of our thoughts

and actions in accordance with our goals. Recent research has started to emphasize the role of

motivation in the timing, intensity, and direction of cognitive control allocation. It is now

recognized that the allocation of cognitive control is driven by goals and therefore closely linked

to motivation (Braver et al., 2014).

Reward prospect enhances cognitive control processes (for reviews see: Botvinick and

Braver, 2015; Krebs and Woldorff, 2017). This effect has been observed on various tasks that tap

into different components of control such as: attentional control (Padmala and Pessoa, 2011),

response inhibition (Leotti and Wager, 2010), conflict adaptation (Braem et al., 2012), and task-

switching (Aarts et al., 2010). The effect is not restricted to situations in which reward is signaled

by advance cues augmenting preparatory control processes. Comparable results are found when

reward is signaled simultaneously with the target thereby promoting fast control adjustments

(Krebs et al., 2010). But why is there a need to adapt control in the first place? Is it not the most

optimal strategy to always exert maximal control over our thoughts and actions? The emerging

answer is that exerting cognitive control carries intrinsic effort-related costs (Shenhav et al.,

2017). It has been shown that exerting control is effortful and that individuals tend to avoid it (Kool et al., 2010). Both reward prospect associated with an outcome, and the effort needed to reach that outcome are important factors in determining the way in which cognitive control is allocated.

Another important factor in determining how cognitive control is allocated is the learned contingency between actions and outcomes. Although this factor has received less empirical attention, several computational models of cognitive control emphasize its importance. For example, the model of Alexander and Brown (2011) stresses the importance of predicting mappings between responses and outcomes in cognitive control. The model proposed by Shenhav and colleagues (2013) posits that the allocation of cognitive control relies, among other factors, on the probability of an outcome conditional on the type and intensity of control. In this way, allocation of cognitive control depends on predictions about probabilities of certain actions (e.g., intense control allocation or a certain response) producing desired outcomes (e.g., solving a task correctly).

Computational Models of Cognitive Control

The insight that cognitive control is closely related to motivation is formalized in a growing number of computational models of cognitive control. At present there are multiple theoretical and computational models which deal with the problem of how cognitive control is allocated (for recent reviews see: Botvinick & Cohen, 2014; Verguts, 2017). These models specify a set of components needed for optimal decision-making about control allocation and they are not mutually exclusive. Many of them include the representations of outcome value and/or outcome controllability (Brown and Alexander, 2017; Holroyd and McClure, 2015; Shenhav et al., 2013; Verguts et al., 2015). Also, many of these models assume that learning is

crucial in forming these representations and emphasize the importance of learning for cognitive control (Abrahamse et al., 2016; Bhandari et al., 2017). Our framework relies on the current formulation of the Expected Value of Control (EVC) theory (Shenhav et al., 2013). This theory includes representations of outcome controllability, value, and cost in the decision-making process about cognitive control allocation.

The EVC theory provides a normative account of cognitive control allocation as resulting from a decision-making process in which potential gains of control allocation are weighted against their costs (Shenhav et al., 2013). The theory postulates that these decisions determine how control is allocated across candidate control signals. These control signals vary on two dimensions: *signal identity*, which determines the processes which are engaged (e.g., paying attention to the ink color in a Stroop task), and *signal intensity* which determines the amount of control allocated (e.g., how intensely to pay attention to the ink color). The theory proposes that control signals are selected in such a way that maximizes the expected value of control at any given moment.

The EVC for a given control signal within a given state is determined by three components: efficacy, value, and cost. *Efficacy* is defined as the probability of a certain outcome given a signal of a particular identity and intensity, and the current state. In other words, the efficacy component can be described as the probability that certain outcomes will occur (i.e., that our actions lead to what we desire) or simply as action-outcome contingencies inferred from previous experience. In standard cognitive control paradigms, such as the Stroop task, an outcome can be a correct or an incorrect response on a particular trial. In this context, we can assign a probability to each of the two possible outcomes given each of the possible signal intensities and identities. *Value* is assigned to each of the possible outcomes and it represents the

value of an outcome in terms of possible rewards or punishments associated with the outcome. The rewards can be both intrinsic (i.e., motivation to do the task well) and extrinsic (i.e., monetary rewards for good performance). Punishments can also come from multiple sources such as monetary loss or poor task performance. Outcome values are also modulated by the time it takes for the outcomes to occur, as individuals try to maximize reward rates (rewards per unit of time) and not rewards per se. *Cost* is defined as the expected cost associated with the specified intensity of control allocation. Within the EVC theory, cost arises as the property of the neural system. This means that as the intensity of a signal to allocate control becomes stronger, so does the cost of allocating control. In sum, the EVC is determined by the probability of an outcome for a given control signal (efficacy), the value of that outcome, and the cost associated with control allocation.

At the neural level, the EVC theory was developed to explain the function of the dorsal anterior cingulate cortex (dACC). The theory assigns the critical role in decision-making about control allocation to this region. It is proposed that the dACC plays a key role in calculating the EVC and that the outcome of this process is then signaled to other regions which implement control, such as the dlPFC (Shenhav et al., 2016, 2013). The dACC is assigned a role in both monitoring for the changes in the environment relevant for control (rewards, punishments, errors, etc.), and, based on this, specifying which control signals maximize the EVC.

The EVC theory provides a detailed and formal account of how cognitive control is allocated. One of the main advantages of the theory is its ability to integrate a wide range of behavioral and neural findings. The computational implementation of the theory has been shown to account for various effects associated with the allocation of cognitive control such as the sequential adaptation effects and post-error slowing (Musslick et al., 2015), including how

individuals can learn about features of their environment that predict incentives for and demands of control allocation (Lieder et al., 2018).

Proposed Framework

We propose a framework that integrates disparate areas of research: cognitive control deficits and motivational impairments in depression. This framework encompasses the controllability, value, and effort of outcomes in order to provide a more mechanistic understanding of these deficits. Furthermore, the framework emphasizes the crucial role of learning in the process of allocating cognitive control. In this way the framework allows for the integration of motivational, learning, and cognitive control deficits in depression. We put forward direct links between components of motivation and reduced cognitive control in depression and propose how these links emerge through learning. In this way our framework offers a first step toward understanding how cognitive control deficits arise. In order to do so, we rely on the EVC theory which offers a computationally explicit model of cognitive control (Shenhav et al., 2013). This allows us to focus on the ways in which expected value of control might be reduced in depression, which results in allocating cognitive control with reduced intensity. We propose that cognitive control deficits in depression arise as a consequence of changes in the decision-making process about control allocation. These deficits in depression occur due to lowered expected value of control. The expected value of control relies on three components of motivation: controllability of outcomes, their value, and the effort needed to attain them.

We argue that research on motivation in depression provides insights into mechanisms leading to a reduction in the expected value of control. The efficacy component in the EVC theory is closely linked to what we have termed outcome controllability the estimates of the probability that actions will lead to desired outcomes. These representations are central in guiding

motivated behavior (Moscarello and Hartley, 2017). As we have reviewed, there is emerging evidence that estimates of outcome controllability and beliefs related to controllability of the environment, are changed in depression. The value component in the EVC theory is related to what we have termed outcome value - the expected reinforcers following an outcome. We have outlined how reward anticipation and reinforcement learning are impaired in depression, especially in the presence of anhedonic symptoms (Treadway and Zald, 2013). The cost component in the EVC theory is linked to the cost of effort needed to obtain rewards. We have reviewed how this process is altered in depression. All of these components of motivation rely on previous learning to estimate the probability of being able to exert control in a new environment, the value of outcomes in that environment, and the effort needed in order to reach them. These insights from the study of motivation in depression offer evidence for the reduction in efficacy, value, and cost of control. This leads to lower expected value of control which in turn results in reduced allocation of cognitive control (Figure 1).

$$EVC\ (signal,\ state) = \sum Pr(outcome_i\ |\ signal,\ state) * Value(outcome_i) - Cost(signal)$$

Figure 1. The schematic representation of the proposed framework. Depression-related changes in the outcome value (e.g. reduced reward anticipation), outcome controllability (e.g. lowered estimates of controllability), and effort costs (e.g. reduced effort exertion), lead to the reduced value of control. This leads to the lowered amounts of control being allocated. The expected value of control (*EVC*) for a signal of a given intensity is calculated as the sum of the values of each possible outcome weighted by the probability of reaching that outcome for the given signal. The cost of allocating control is subtracted from that sum. The figure was adapted from Shenhav et al., 2013 with permission from the authors.

Without a doubt, estimates of controllability, value, and effort of outcomes, are formed through prior learning. Estimates formed in one situation are generalized and used in novel similar environments. For example, imagine a person who developed low estimates of controllability in the work environment. Over the years that person may have learned that no matter what they do in the work environment, it is unlikely that desired outcomes occur. That person will tend to allocate less cognitive control in that environment and will develop beliefs about the inability to do the job well. This will lead to a reduction in motivated behavior and will further strengthen existing low controllability estimates and beliefs. Alternatively, another individual might estimate high controllability in the same work environment. However, that

individual can anticipate low amounts of rewards associated with achieving good results in work, or high amounts of effort needed to do so. That individual does not anticipate pleasure in work anymore and prefers to spend time doing other more pleasurable activities. That person will also tend to allocate less cognitive control during work hours.

Within our framework, depression-related deficits in cognitive control processes such as inhibition, task switching, or working memory updating (Joormann & Tanovic, 2015), can be conceptualized as products of changes in the expected value of control. In this way our framework integrates cognitive research on depression with impairments in other domains. Our framework outlines the links between cognitive control deficits and the study of motivation in depression. The framework can account for both the existing research findings and further develop the field by connecting these findings to previously unrelated literatures. Within our framework, cognitive deficits in depression (Gotlib and Joormann, 2010) can be studied in relation with reward processing impairments in depression (Admon and Pizzagalli, 2015) and changes in perceived controllability of an environment (Moscarello and Hartley, 2017).

This framework has several key implications. First, cognitive control deficits do not have a single cause and can be caused by impairments in different processes. Deficits can thus be present in depressed individuals with different clusters of symptoms (e.g., mainly depressive mood or mainly anhedonia). Second, cognitive deficits are causally related to reward processing impairments, effort costs, and estimates of controllability of an environment. These impairments influence allocation of cognitive control through learned estimates of reward probability, action-outcome contingencies, and needed effort. Third, cognitive control deficits are not a product of the simple inability to exert control. They are a result of lowered expectations about the value of exerting cognitive control.

Neural Level

Cognitive control deficits in depression are proposed to be related to functional changes in the dlPFC and the ACC (Disner et al., 2011). This view is based on the conceptualization of cognitive control functions as reliant purely on the prefrontal cortex. However, this view has been challenged by the discovery of the role of other regions, such as the basal ganglia and corticostriatal loops in cognitive control and goal-directed behavior (for reviews see: Balleine & O'Doherty, 2010; Chatham & Badre, 2015; Haber, 2016). By building on the EVC theory, our framework offers a more specific view on the role of these two regions in depression. We propose that depression-related changes in the decision-making process about control allocation are related to the activity of the dACC. These changes can further result in lower levels of dlPFC activation which implements control based on the signals from the dACC. In this way, depression is not related to an inability of the dlPFC to implement control, but related to changes in inputs to the dACC used in decision-making about control allocation. In line with the importance of the dACC in our framework, current research has pointed to the crucial role of the ACC in depression (for a review see: Holroyd and Umemoto, 2016). Our framework also points to the important role of efficacy and value representations which serve as inputs in the decision-making process. In this way, it can guide further research in connecting the role of the dACC and other regions and networks related to encoding value, reward processing, and efficacy (Haber and Knutson, 2010; Moscarello and Hartley, 2017; Treadway and Zald, 2011).

Computational Implementation

By relying on a formalized model of cognitive control (Shenhav et al., 2013), our framework provides the opportunity to make more precise predictions at both behavioral and neural level. The more recent work provides a computational implementation of the EVC theory

and provides a way to simulate the influence of changes in the efficacy, cost, and value

components on cognitive control allocation (Musslick et al., 2015). In this way testable

predictions of behavior and neural responses on cognitive control tasks can be generated.

To demonstrate how control-demanding behavior is affected by changes in the decision-

making process about control allocation, we simulated the behavior of an agent across two

paradigms while varying parameters of that agent's EVC-driven control valuation (Musslick et

al., 2015; for a detailed description of these simulations, see Supplementary Materials). We first

simulated performance in a Stroop task in which the simulated agent had to categorize a the ink

color (e.g. red or green) of a color word (e.g. "RED" or "GREEN") on each trial (Stroop, 1935).

We assessed the overall control allocation and the resulting performance cost associated with

responding to a color that is response-incongruent with the word (e.g. "RED" displayed in green)

compared to responding to a color that is response-congruent with a word (e.g. "RED" displayed

in red). We also simulated behavior in a cognitive effort discounting (COGED) task, in which the

agent must choose between performing a difficult, high-demand task for $2 and an easy, low-

demand (baseline) task for a variable amount of monetary reward on each trial. We assessed the

subjective value of performing a high-demand task by measuring amount of monetary reward

offered for the baseline task for which the EVC agent is indifferent between the two tasks, and by

normalizing this amount by the reward offered for the high-demand task ($2). We evaluated how

these complementary measures – of task performance vs. task preference – were influenced by

changes in the agent's (a) cost of cognitive control (b) sensitivity to reward, and (c) expected

efficacy of exerting cognitive control (control efficacy).

Consistent with observations in healthy human subjects, our simulated agents

demonstrated an incongruency cost, generating more errors when performing incongruent trials

compared to congruent trials (Stroop, 1935). Critically, agents chose to exert less control (Fig 1A-C), resulting in higher incongruency costs (Fig 1D-F) when they had (i) higher costs of control, (ii) lower reward sensitivity, and/or (iii) when they perceived their control as being less efficacious. The simulated agent also replicated the behavior of healthy human subjects in studies of cognitive effort selection, assigning lower subjective values to tasks that demanded higher amounts of cognitive control (Westbrook et al., 2013). However, the subjective value for a given task decreased as (i) the cost of cognitive control increased, (ii) reward sensitivity decreased, and/or (iii) the control efficacy decreases (Figure 2G-I). Interestingly, the influence of these changes in parameters are slightly magnified for high task difficulties compared to low task difficulties.

Figure 2. Effects of control cost, reward sensitivity and control efficacy on simulated behavior of the EVC model. Behavior of model was simulated in the Stroop paradigm and COGED paradigm (see supplementary Materials for details). (A-C) The amount of cognitive control allocated in a Stroop task is shown as a function of control cost, sensitivity to reward and expected control efficacy. (D-F) The performance impairment on incongruent compared to congruent trials is shown as a function of the three model parameters. (G-I) The subjective value of a task in the COGED paradigm is plotted as a function of task difficulty for different values of control costs, sensitivity to reward and control efficacy.

The computational implementation of our framework offers a possibility of studying cognitive impairments in depression within the developing framework of computational psychiatry (Huys et al., 2016; Montague et al., 2012). Several computational frameworks have already started to model behavioral control in depression (Huys et al., 2015; Huys and Dayan, 2009) and our framework offers the possibility of extending this research into the domain of cognitive control deficits in depression. We have described the advantages of the EVC theory

above, but we do recognize the importance of other related computational models of cognitive control. By focusing our framework on components which are shared across many models, we hope to provide a synthesis that is useful beyond the boundaries of the EVC theory and that can be broadly used to understand cognitive control deficits in depression.

Relationship with Other Constructs Relevant for Depression

Lowered levels of cognitive control have a reciprocal relationship with negative beliefs and attributions. Negative beliefs about the self, the others, and the world represent a crucial cognitive vulnerability factor to depression (Beck, 1972). Attributions that are global, internal, and stable are also an important vulnerability factor (Abramson et al., 1989). Cognitive control deficits can strengthen those beliefs, but are also strengthened by them. Finally, lowered levels of cognitive control are related to maladaptive use of emotion regulation strategies such as rumination (Nolen-Hoeksema et al., 2008) which contribute to the onset and maintenance of depressive symptoms (Joormann & Vanderlind, 2014). Impaired cognitive control also affects other cognitive processes thus producing cognitive biases in attention, interpretation, and memory, which further promote the maladaptive use of emotion regulation strategies (Everaert et al., 2012; Gotlib and Joormann, 2010).

An important research line has focused on the relationship between depressive symptoms and cognitive control over affectively negative material. These studies have demonstrated specific impairments in shifting attention away from negative material, removing negative material from the working memory, and inhibiting negative stimuli (Gotlib and Joormann, 2010; Joormann and Vanderlind, 2014; Koster et al., 2011). In its current form, our framework does not focus on these deficits. However, we hope to integrate this body of work in the framework in future studies. One of the interesting possibilities for such integration is the influence of negative

material on efficacy estimates through prior learning.

Future Directions and Open Questions

The proposed framework opens novel avenues for research, namely the links between cognitive control in depression and alterations in components of motivation. Several studies have demonstrated the importance of motivation for cognitive functioning in depression (Moritz et al., 2017; Scheurich et al., 2008). Recently, studies have focused on more specific components of motivation. For example, several studies have investigated the relationship between reward processing and cognitive control or attention in relation to depressive symptoms (Anderson et al., 2014; Ravizza and Delgado, 2014). Future studies should focus on precise distinctions between different types of impairments related to reward processing. In this context, we believe that reward anticipation and effort costs are the most interesting processes to investigate in relation to cognitive processes in depression. Depression research can rely on paradigms developed in cognitive science to study motivation and cognitive control (Botvinick and Braver, 2015). The study of effort expenditure is also gaining a great deal of attention and paradigms to study physical effort already exist (Treadway et al., 2012). For depression research, novel insights on cognitive effort (Shenhav et al., 2017) can be of particular relevance. Also, the relationship between anhedonic symptoms and cognitive impairments in depression will be a crucial next step in better understanding cognition in depression. In this domain research should be guided by recent developments in measuring anhedonia through behavioral tasks and questionnaires (for a review on measures of anhedonia see: Rizvi et al., 2016).

In the domain of efficacy, there are no readily available paradigms that could directly inform us about its relationship with cognitive control. However, developments in the field of learned helplessness suggest that the interactions between stress and controllability are crucial for

stress regulation (Maier and Seligman, 2016). Recent research is starting to provide paradigms to study stress controllability in humans (Bhanji et al., 2016; Hartley et al., 2014). The critical next step is to develop paradigms that would allow the investigation of cognitive control processes in relation to controllability. Recently, paradigms aimed to investigate different components of motivated action and learning have been developed in schizophrenia research. Using a novel paradigm, Morris and associates have demonstrated impairments in action-outcome learning in schizophrenia (Morris, Cyrzon, Green, Le Pelley, & Balleine, 2018; Morris, Quail, Griffiths, Green, & Balleine, 2015; see also: Liljeholm et al., 2011). The use of similar paradigms would allow for further work on computational models of motivation and cognition in depression. Recent computational work has demonstrated that generalization of learned action-outcome contingencies can account for a wide range of behavioral features associated with learned helplessness in animals and humans (Huys and Dayan, 2009; Lieder et al., 2013). However, there is a strong need for more empirical work which could precisely measure the importance of outcome controllability in depression.

Cognitive control deficits (as well as anhedonia) are transdiagnostic in nature and co-occur in many disorders (Goschke, 2014; Whitton et al., 2015). Although our model is focused on depression, the links between cognitive control and reward processing are relevant for other psychiatric disorders as well. Future studies in this domain hold the promise of not only detecting transdiagnostic constructs such as the positive valence and cognitive systems (Cuthbert and Insel, 2013; Insel et al., 2010), but starting to investigate the relations between these systems. In this context, the relationship between reward processing (positive valence system) and cognitive control (cognitive system) across different clinical populations can be of great interest.

In current research, depression is often regarded and measured as a single, homogeneous

disorder and we have treated it as such in this paper. Importantly, diagnosis of depression requires only presence of one out of the two core symptoms: prolonged negative affect and the loss of interest in previously pleasurable activities (DSM-5; American Psychiatric Association, 2013). A growing amount of evidence is pointing to the heterogeneous nature of depression and the need to analyze specific symptoms of depression (Fried and Nesse, 2015). Our framework emphasizes the possibility that cognitive impairments in depression emerge from different causes. This allows for a study of how potential different causes of cognitive deficits in depression can be related to different symptom clusters. For example, reward processing impairments, resulting in lowered levels of cognitive control allocation, can be related to anhedonic symptoms. In the same way, changes in efficacy estimates can produce control deficits, but they could be more related to the presence of negative affect.

The study of mechanisms through which cognitive control deficits in depression emerge is of great relevance for developing future cognitive treatments for depression. Currently, cognitive trainings are showing considerable promise (Koster et al., 2017; Siegle et al., 2007), but the mechanisms through which cognitive trainings improve depressive symptoms remain unknown. Understanding the components of motivation that are affected by these trainings will allow for more precise and individualized treatments.

Conclusions

Cognitive control deficits represent an important vulnerability factor for depression and play a central role in cognitive impairments in depression. We propose a framework of cognitive control in depression in which these deficits emerge as a consequence of changes in the decision-making process about control allocation. We argue that alterations in core components of motivated behavior – reward anticipation, effort costs, and estimates of environment

controllability – are crucial mechanisms contributing to dysfunctional cognitive control in depression. This view provides a more mechanistic understanding of cognitive control in depression, offers a better computational and neural understanding of the deficits, and connects cognitive research on depression with other fields of study such as motivation and agency. We believe that our framework can guide future research on mechanisms underlying depressive cognition which can result in improving treatments for depression.

Acknowledgements

This work was supported by the Special Research Fund (BOF) of Ghent University [grant number 01D02415 awarded to IG], by a Center of Biomedical Research Excellence grant from the National Institute of General Medical Sciences [grant number P20GM103645 awarded to AS], by the starting grant of the European Research Council (ERC) under the Horizon 2020 framework [grant number 636110 awarded to RK], and by the Concerted Research Action Grant of Ghent University [grant number BOF16/GOA/017 awarded to EHWK], and a grant from the John Templeton Foundation [awarded to SM; The opinions expressed in this publication are those of the authors and do not necessarily reflect the views of the John Templeton Foundation].

References

Aarts, E., Roelofs, A., Franke, B., Rijpkema, M., Fernández, G., Helmich, R.C., Cools, R., 2010. Striatal dopamine mediates the interface between motivational and cognitive control in humans: Evidence from genetic imaging. Neuropsychopharmacology 35, 1943–1951. https://doi.org/10.1038/npp.2010.68

Abrahamse, E., Braem, S., Notebaert, W., Verguts, T., 2016. Grounding cognitive control in associative learning. Psychol. Bull. 142. https://doi.org/10.1037/bul0000047

Abramson, L.Y., Metalsky, G.I., Alloy, L.B., 1989. Hopelessness depression: A theory-based subtype of depression. Psychol. Rev. 96, 358–372. https://doi.org/10.1037/0033-295X.96.2.358

Abramson, L.Y., Seligman, M.E.P., Teasdale, J.D., 1978. Learned helplessness in humans and reformulation. J. Abnorm. Psychol. 87, 49–74. https://doi.org/10.1037/0021-843X.87.1.49

Admon, R., Pizzagalli, D.A., 2015. Dysfunctional Reward Processing in Depression. Curr. Opin. Psychol. 4, 114–118. https://doi.org/10.1016/j.copsyc.2014.12.011

Alexander, W.H., Brown, J.W., 2011. Medial Prefrontal Cortex as an action-outcome predictor. Nat. Neurosci. 14, 1338–1344. https://doi.org/10.1038/nn.2921.Medial

American Psychiatric Association, 2013. Diagnostic and Statistical Manual of Mental Disorders, 5th ed. American Psychiatric Publishing, Washington DC.

Anderson, B.A., Leal, S.L., Hall, M.G., Yassa, M.A., Yantis, S., 2014. The attribution of value-based attentional priority in individuals with depressive symptoms. Cogn. Affect. Behav. Neurosci. 14, 1221–7. https://doi.org/10.3758/s13415-014-0301-z

Atkinson, J.W., 1957. Motivational determinants of risk taking behavior. Psychol. Rev. 64, 359–372. https://doi.org/10.1037/h0043445

Balleine, B.W., O'Doherty, J.P., 2010. Human and rodent homologies in action control: Corticostriatal determinants of goal-directed and habitual action. Neuropsychopharmacology 35, 48–69. https://doi.org/10.1038/npp.2009.131

Bandura, A., 2001. Social Cogntive Theory: An Agentic Perspective. Annu. Rev. Psychol. 52, 1–26. https://doi.org/10.1146/annurev.psych.52.1.1

Bandura, A., 1977. Self-efficacy: Toward a unifying theory of behavioral change. Psychol. Rev. 84, 191–215. https://doi.org/10.1037/0033-295x.84.2.191

Barch, D.M., Dowd, E.C., 2010. Goal representations and motivational drive in schizophrenia: The role of prefrontal-striatal interactions. Schizophr. Bull. 36, 919–934. https://doi.org/10.1093/schbul/sbq068

Barch, D.M., Pagliaccio, D., Luking, K.R., 2015. Mechanisms underlying motivational deficits in psychopathology: similarities and differences in depression and schizophrenia, in: Behavioral Neuroscience of Motivation. Springer International Publishing, pp. 411–449.

Beck, A.T., 1972. Depression: Causes and treatment. University of Pennsylvania Press.

Bhandari, A., Badre, D., Frank, M.J., 2017. Learning Cognitive Control, in: Egner, T. (Ed.), Wiley Handbook of Cognitive Control. Wiley-Blackwell, pp. 376–391.

Bhanji, J.P., Kim, E.S., Delgado, M.R., 2016. Perceived Control Alters the Effect of Acute Stress on Persistence 145, 356–365.

Bogacz, R., Brown, E., Moehlis, J., Holmes, P., Cohen, J.D., 2006. The physics of optimal decision making: A formal analysis of models of performance in two-alternative forced-choice tasks. Psychol. Rev. 113, 700–765. https://doi.org/10.1037/0033-295X.113.4.700

Bonner, S.E., Sprinkle, G.B., 2002. The effects of monetary incentives on effort and task performance: theories, evidence, and a framework for research. Accountine, Organ. Soc. 27, 303–345. https://doi.org/10.1016/S0361-3682(01)00052-6

Botvinick, M.M., Braver, T., 2015. Motivation and Cognitive Control: From Behavior to Neural Mechanism. Annu. Rev. Psychol. 66, 83–113. https://doi.org/10.1146/annurev-psych-010814-015044

Botvinick, M.M., Braver, T.S., 2015. Motivation and Cognitive Control : From Behavior to Neural Mechanism. Annu. Rev. Psychol. 1–31. https://doi.org/10.1146/annurev-psych-010814-015044

Botvinick, M.M., Cohen, J.D., 2014. The Computational and Neural Basis of Cognitive Control : Charted Territory and New Frontiers 38, 1249–1285. https://doi.org/10.1111/cogs.12126

Braem, S., Verguts, T., Roggeman, C., Notebaert, W., 2012. Reward modulates adaptations to conflict. Cognition 125, 324–332. https://doi.org/10.1016/j.cognition.2012.07.015

Braver, T.S., Krug, M.K., Chiew, K.S., Kool, W., Westbrook, J.A., Clement, N.J., Adcock, R.A., Barch, D.M., Botvinick, M.M., Carver, C.S., Cools, R., Custers, R., Dickinson, A., Dweck, C.S., Fishbach, A., Gollwitzer, P.M., Hess, T.M., Isaacowitz, D.M., Mather, M., Murayama, K., Pessoa, L., Samanez-Larkin, G.R., Somerville, L.H., 2014. Mechanisms of motivation-cognition interaction: challenges and opportunities. Cogn. Affect. Behav. Neurosci. 14, 443–72. https://doi.org/10.3758/s13415-014-0300-0

Brown, J.W., Alexander, W.H., 2017. Foraging Value, Risk Avoidance, and Multiple Control Signals: How the Anterior Cingulate Cortex Controls Value-based Decision-making. J. Cogn. Neurosci. 29, 1656–1673. https://doi.org/10.1162/jocn_a_01140

Chatham, C.H., Badre, D., 2015. Multiple gates on working memory. Curr. Opin. Behav. Sci. 1, 23–31. https://doi.org/10.1016/j.cobeha.2014.08.001

Clark, D.A., Beck, A.T., 2010. Cognitive theory and therapy of anxiety and depression: Convergence with neurobiological findings. Trends Cogn. Sci. 14, 418–424. https://doi.org/10.1016/j.tics.2010.06.007

Cléry-Melin, M.L., Schmidt, L., Lafargue, G., Baup, N., Fossati, P., Pessiglione, M., 2011. Why don't you try harder? an investigation of effort production in major depression. PLoS One 6, 1–8. https://doi.org/10.1371/journal.pone.0023178

Crocker, L.D., Heller, W., Warren, S.L., O'Hare, A.J., Infantolino, Z.P., Miller, G. a, 2013. Relationships among cognition, emotion, and motivation: implications for intervention and neuroplasticity in psychopathology. Front. Hum. Neurosci. 7, 261. https://doi.org/10.3389/fnhum.2013.00261

Culbreth, A.J., Moran, E.K., Barch, D.M., 2017. Effort-cost decision-making in psychosis and

depression: could a similar behavioral deficit arise from disparate psychological and neural mechanisms? Psychol. Med. 1–16. https://doi.org/10.1017/S0033291717002525

Cuthbert, B.N., Insel, T.R., 2013. Toward the future of psychiatric diagnosis: the seven pillars of RDoC. BMC Med. 11, 126. https://doi.org/10.1186/1741-7015-11-126

Davidson, R.J., Pizzagalli, D., Nitschke, J.B., Putnam, K., 2002. Depression: Perspectives from Affective Neuroscience. Annu. Rev. Psychol. 53, 545–574. https://doi.org/10.1146/annurev.psych.53.100901.135148

Daw, N.D., O'Doherty, J.P., 2014. Multiple Systems for Value Learning, in: Glimcher, P.W., Fehr, E. (Eds.), Neuroeconomics: Decision Making and the Brain: Second Edition. Academic Press, pp. 393–410. https://doi.org/10.1016/B978-0-12-416008-8.00021-8

De Raedt, R., Hooley, J.M., 2016. The role of expectancy and proactive control in stress regulation : A neurocognitive framework for regulation expectation. Clin. Psychol. Rev. 45, 45–55. https://doi.org/10.1016/j.cpr.2016.03.005

Demeyer, I., De Lissnyder, E., Koster, E.H.W., De Raedt, R., 2012. Rumination mediates the relationship between impaired cognitive control for emotional information and depressive symptoms: A prospective study in remitted depressed adults. Behav. Res. Ther. 50, 292–7. https://doi.org/10.1016/j.brat.2012.02.012

Der-Avakian, A., Markou, A., 2012. The neurobiology of anhedonia and other reward-related deficits. Trends Neurosci. 35, 68–77. https://doi.org/10.1016/j.tins.2011.11.005

Derakshan, N., Salt, M., Koster, E.H.W., 2009. Attentional control in dysphoria: An investigation using the antisaccade task. Biol. Psychol. 80, 251–255. https://doi.org/10.1016/j.biopsycho.2008.09.005

Dickinson, A., 1985. Actions and habits: the development of behavioural autonomy. Philos. Trans. R. Soc. Lond. B. Biol. Sci. 308, 67–78.

Disner, S.G.S., Beevers, C.G.C., Haigh, E.A.P., Beck, A.T., 2011. Neural mechanisms of the cognitive model of depression. Nat. Rev. Neurosci. 12, 467–77. https://doi.org/10.1038/nrn3027

Everaert, J., Grahek, I., Koster, E.H.W., 2017. Individual differences in cognitive control modulate cognitive biases linked to depressive. Cogn. Emot. 31, 736–746. https://doi.org/10.1080/02699931.2016.1144562

Everaert, J., Koster, E.H.W., Derakshan, N., 2012. The combined cognitive bias hypothesis in depression. Clin. Psychol. Rev. 32, 413–24. https://doi.org/10.1016/j.cpr.2012.04.003

Fried, E.I., Nesse, R.M., 2015. Depression sum-scores don't add up: why analyzing specific depression symptoms is essential. BMC Med. 13, 72. https://doi.org/10.1186/s12916-015-0325-4

Friedman, N.P., Miyake, A., 2017. Unity and diversity of executive functions: Individual differences as a window on cognitive structure. Cortex 86, 186–204. https://doi.org/10.1016/j.cortex.2016.04.023

Goschke, T., 2014. Dysfunctions of decision-making and cognitive control as transdiagnostic mechanisms of mental disorders: advances, gaps, and needs in current research. Int. J. Methods Psychiatr. Res. 23. https://doi.org/10.1002/mpr.1410

Gotlib, I.H., Hamilton, J.P., 2008. Neuroimaging and depression: Current status and unresolved issues. Curr. Dir. Psychol. Sci. 17, 159–163. https://doi.org/10.1111/j.1467-8721.2008.00567.x

Gotlib, I.H., Joormann, J., 2010. Cognition and depression: current status and future directions. Annu. Rev. Clin. Psychol. 6, 285–312. https://doi.org/10.1146/annurev.clinpsy.121208.131305

Grahek, I., Everaert, J., Krebs, R.M., Koster, E.H.W., 2018. Cognitive control in depression: Towards clinical models informed by cognitive neuroscience. Clin. Psychol. Sci. https://doi.org/10.1177/2167702618758969

Griffiths, K.R., Morris, R.W., Balleine, B.W., 2014. Translational studies of goal-directed action as a framework for classifying deficits across psychiatric disorders. Front. Syst. Neurosci. 8, 1–16. https://doi.org/10.3389/fnsys.2014.00101

Haber, S.N., 2016. Corticostriatal circuitry. Dialogues Clin. Neurosci. 18, 7–21. https://doi.org/10.1002/bit.260190307

Haber, S.N., Knutson, B., 2010. The reward circuit: Linking primate anatomy and human imaging. Neuropsychopharmacology 35, 4–26. https://doi.org/10.1038/npp.2009.129

Hartley, C.A., Gorun, A., Reddan, M.C., Ramirez, F., Phelps, E.A., 2014. Neurobiology of Learning and Memory Stressor controllability modulates fear extinction in humans. Neurobiol. Learn. Mem. 113, 149–156. https://doi.org/10.1016/j.nlm.2013.12.003

Hershenberg, R., Satterthwaite, T.D., Daldal, A., Katchmar, N., Moore, T.M., Kable, J.W., Wolf, D.H., 2016. Diminished effort on a progressive ratio task in both unipolar and bipolar depression. J. Affect. Disord. 196, 97–100. https://doi.org/10.1016/j.jad.2016.02.003

Hertel, P., 2000. The cognitive-initiative account of depression-related impairments in memory. Psychol. Learn. Motiv. 39, 47–71. https://doi.org/10.1016/S0079-7421(00)80030-4

Hertel, P.T., 1994. Depression and Memory: Are Impairments Remediable Through Attentional Control? Curr. Dir. Psychol. Sci. 3, 190–193. https://doi.org/10.1111/1467-8721.ep10770707

Holroyd, C.B., McClure, S.M., 2015. Hierarchical control over effortful behavior by rodent medial frontal cortex: A computational model. Psychol. Rev. 122, 54–83.

https://doi.org/10.1037/a0038339

Holroyd, C.B., Umemoto, A., 2016. The Research Domain Criteria Framework: The Case for
    Anterior Cingulate Cortex. Neurosci. Biobehav. Rev. 71, 418–443.
    https://doi.org/10.1016/j.neubiorev.2016.09.021

Huys, Q.J.M., Daw, N.D., Dayan, P., 2015. Depression: A Decision-Theoretic Analysis. Annu.
    Rev. Neurosci. 38, 1–23. https://doi.org/10.1146/annurev-neuro-071714-033928

Huys, Q.J.M., Dayan, P., 2009. A Bayesian formulation of behavioral control. Cognition 113,
    314–328. https://doi.org/10.1016/j.cognition.2009.01.008

Huys, Q.J.M., Maia, T. V, Frank, M.J., 2016. Computational psychiatry as a bridge from
    neuroscience to clinical applications. Nat. Neurosci. 19, 404–13.
    https://doi.org/10.1038/nn.4238

Insel, T., Cuthbert, B., Garvie, M., Heinssen, R., Pine, D.S., Quinn, K., Sanislow, C., Wang, P.,
    2010. Research Domain Criteria (RDoC): Toward a New Classification Framework for
    Research on Mental Disorders. Am. J. Psychiatry 167, 748–751.
    https://doi.org/10.1176/appi.ajp.2010.09091379

Inzlicht, M., Shenhav, A., Olivola, C.Y., 2018. The Effort Paradox: Effort Is Both Costly and
    Valued. Trends Cogn. Sci. 22, 337–349. https://doi.org/10.1016/j.tics.2018.01.007

Joormann, J., 2010. Cognitive Inhibition and Emotion Regulation in Depression. Curr. Dir.
    Psychol. Sci. 19, 161–166. https://doi.org/10.1177/0963721410370293

Joormann, J., Tanovic, E., 2015. Cognitive vulnerability to depression: examining cognitive
    control and emotion regulation. Curr. Opin. Psychol. 4, 86–92.
    https://doi.org/10.1016/j.copsyc.2014.12.006

Joormann, J., Vanderlind, W.M., 2014. Emotion Regulation in Depression: The Role of Biased
    Cognition and Reduced Cognitive Control. Clin. Psychol. Sci. 2, 402–421.
    https://doi.org/10.1177/2167702614536163

Joormann, J., Yoon, K.L., Zetsche, U., 2007. Cognitive inhibition in depression. Appl. Prev.
    Psychol. 12, 128–139. https://doi.org/10.1016/j.appsy.2007.09.002

Kahneman, D., 1973. Attention and Effort. Prentice-Hall, Englewood Cliffs, NJ.

Kaser, M., Zaman, R., Sahakian, B.J., 2017. Cognition as a treatment target in depression.
    Psychol. Med. 47, 987–989. https://doi.org/10.1017/S0033291716003123

Knutson, B., Westdorp, A., Kaiser, E., Hommer, D., 2000. FMRI visualization of brain activity
    during a monetary incentive delay task. Neuroimage 12, 20–27.
    https://doi.org/10.1006/nimg.2000.0593

Kool, W., Botvinick, M., 2018. Mental labour. Nat. Hum. Behav. https://doi.org/10.1038/s41562-

018-0401-9

Kool, W., McGuire, J.T., Rosen, Z.B., Botvinick, M.M., 2010. Decision making and the avoidance of cognitive demand. J. Exp. Psychol. Gen. 139, 665–682. https://doi.org/10.1037/a0020198

Koster, E.H.W., De Lissnyder, E., Derakshan, N., De Raedt, R., 2011. Understanding depressive rumination from a cognitive science perspective: the impaired disengagement hypothesis. Clin. Psychol. Rev. 31, 138–45. https://doi.org/10.1016/j.cpr.2010.08.005

Koster, E.H.W., Hoorelbeke, K., Onraedt, T., Owens, M., Derakshan, N., 2017. Cognitive control interventions for depression: A systematic review of findings from training studies. Clin. Psychol. Rev. 53, 79–92. https://doi.org/10.1016/j.cpr.2017.02.002

Krebs, R.M., Boehler, C.N., Woldorff, M.G., 2010. The influence of reward associations on conflict processing in the Stroop task. Cognition 117, 341–347. https://doi.org/10.1016/j.cognition.2010.08.018

Krebs, R.M., Woldorff, M.G., 2017. Cognitive control and reward., in: Egner, T. (Ed.), Wiley Handbook of Cognitive Control. Wiley-Blackwell, pp. 422–439.

Kurzban, R., Duckworth, A., Kable, J.W., Myers, J., 2013. An opportunity cost model of subjective effort and task performance. Behav. Brain Sci. 36, 661–679. https://doi.org/10.1017/S0140525X12003196

Leotti, L. a, Wager, T.D., 2010. Motivational influences on response inhibition measures. J. Exp. Psychol. Hum. Percept. Perform. 36, 430–47. https://doi.org/10.1037/a0016802

Leotti, L.A., Iyengar, S.S., Ochsner, K.N., 2010. Born to choose: The origins and value of the need for control. Trends Cogn. Sci. 14, 457–463. https://doi.org/10.1016/j.tics.2010.08.001

Levens, S.M., Gotlib, I.H., 2015. Updating emotional content in recovered depressed individuals: Evaluating deficits in emotion processing following a depressive episode. J. Behav. Ther. Exp. Psychiatry 48, 156–163. https://doi.org/10.1016/j.jbtep.2015.03.009

Lieder, F., Goodman, N.D., Huys, Q.J., 2013. Learned helplessness and generalization. Proc. Annu. Meet. Cogn. Sci. Soc. 35.

Lieder, F., Shenhav, A., Musslick, S., Griffiths, T.L., 2018. Rational metareasoning and the plasticity of cognitive control. Plos Comput. Biol. 14. https://doi.org/10.13140/RG.2.2.24500.14721

Liljeholm, M., Tricomi, E., O'Doherty, J.P., Balleine, B.W., 2011. Neural Correlates of Instrumental Contingency Learning: Differential Effects of Action-Reward Conjunction and Disjunction. J. Neurosci. 31, 2474–2480. https://doi.org/10.1523/JNEUROSCI.3354-10.2011

Maier, S.F., 1984. Learned helplessness and animal models of depression. Prog. Neuro-Psychopharmacology Biol. Psychiatry 8, 435–446. https://doi.org/10.1016/S0278-5846(84)80032-9

Maier, S.F., Amat, J., Baratta, M. V., Paul, E., Watkins, L.R., 2006. Behavioral control, the medial prefrontal cortex, and resilience. Dialogues Clin. Neurosci. 8, 397–406. https://doi.org/10.1016/j.ynstr.2014.09.003

Maier, S.F., Seligman, M.E.P., 2016. Learned helplessness at fifty: Insights from neuroscience. Psychol. Rev. 123, 1–19. https://doi.org/10.1037/rev0000033

Maier, S.F., Watkins, L.R., 2005. Stressor controllability and learned helplessness: The roles of the dorsal raphe nucleus, serotonin, and corticotropin-releasing factor. Neurosci. Biobehav. Rev. 29, 829–841. https://doi.org/10.1016/j.neubiorev.2005.03.021

Marchetti, I., Shumake, J., Grahek, I., Koster, E.H.W., 2018. Temperamental factors in remitted depression: The role of effortful control and attentional mechanisms. J. Affect. Disord. 235. https://doi.org/10.1016/j.jad.2018.04.064

McMakin, D.L., Olino, T.M., Porta, G., Dietz, L.J., Emslie, G., Clarke, G., Wagner, K.D., Asarnow, J.R., Ryan, N.D., Birmaher, B., Shamseddeen, W., Mayes, T., Kennard, B., Spirito, A., Keller, M., Lynch, F.L., Dickerson, J.F., Brent, D.A., 2012. Anhedonia predicts poorer recovery among youth with selective serotonin reuptake inhibitor treatmentresistant depression. J. Am. Acad. Child Adolesc. Psychiatry 51, 404–411. https://doi.org/10.1016/j.jaac.2012.01.011

Millan, M.J., Agid, Y., Brüne, M., Bullmore, E.T., Carter, C.S., Clayton, N.S., Connor, R., Davis, S., Deakin, B., DeRubeis, R.J., Dubois, B., Geyer, M.A., Goodwin, G.M., Gorwood, P., Jay, T.M., Joëls, M., Mansuy, I.M., Meyer-Lindenberg, A., Murphy, D., Rolls, E., Saletu, B., Spedding, M., Sweeney, J., Whittington, M., Young, L.J., 2012. Cognitive dysfunction in psychiatric disorders: characteristics, causes and the quest for improved therapy. Nat. Rev. Drug Discov. 11, 141–168. https://doi.org/10.1038/nrd3628

Miller, K.J., Shenhav, A., Ludvig, E.A., n.d. Habits without values. Psychol. Rev.

Montague, P.R., Dolan, R.J., Friston, K.J., Dayan, P., 2012. Computational psychiatry. Trends Cogn. Sci. 16. https://doi.org/10.1016/j.tics.2011.11.018

Moritz, S., Stöckert, K., Hauschildt, M., Lill, H., Jelinek, L., Beblo, T., Diedrich, S., Arlt, S., 2017. Are we exaggerating neuropsychological impairment in depression? Reopening a closed chapter. Expert Rev. Neurother. 17, 839–846. https://doi.org/10.1080/14737175.2017.1347040

Morris, R.W., Cyrzon, C., Green, M.J., Le Pelley, M.E., Balleine, B.W., 2018. Impairments in action–outcome learning in schizophrenia. Transl. Psychiatry 8, 54. https://doi.org/10.1038/s41398-018-0103-0

Morris, R.W., Quail, S., Griffiths, K.R., Green, M.J., Balleine, B.W., 2015. Corticostriatal control of goal-directed action is impaired in schizophrenia. Biol. Psychiatry 77, 187–195. https://doi.org/10.1016/j.biopsych.2014.06.005

Moscarello, J.M., Hartley, C.A., 2017. Agency and the Calibration of Motivated Behavior. Trends Cogn. Sci. 21, 725–735. https://doi.org/10.1016/j.tics.2017.06.008

Musslick, S., Shenhav, A., Botvinick, M.M., Cohen, J.D., 2015. A computational model of control allocation based on the Expected Value of Control. Reinf. Learn. Decis. Mak. Conf. 59, 2014.

Navarro, D.J., Fuss, I.G., 2009. Fast and accurate calculations for first-passage times in Wiener diffusion models. J. Math. Psychol. 53, 222–230. https://doi.org/10.1016/j.jmp.2009.02.003

Nolen-Hoeksema, S., Wisco, B.E., Lyubomirsky, S., 2008. Rethinking Rumination. Perspect. Psychol. Sci. 3, 400–424. https://doi.org/10.1111/j.1745-6924.2008.00088.x

Oettingen, G., Gollwitzer, P.M., 2001. Goal setting and goal striving, in: Tesser, A., Schwarz, N. (Eds.), Blackwell Handbook of Social Psychology: Intraindividual Processes. Blackwell Publishers, pp. 329–348.

Owens, M., Koster, E.H.W., Derakshan, N., 2012. Impaired filtering of irrelevant information in dysphoria: An ERP study. Soc. Cogn. Affect. Neurosci. 7, 752–763. https://doi.org/10.1093/scan/nsr050

Padmala, S., Pessoa, L., 2011. Reward reduces conflict by enhancing attentional control and biasing visual cortical processing. J. Cogn. Neurosci. 23, 3419–3432. https://doi.org/10.1162/jocn_a_00011

Pizzagalli, D.A., 2014. Depression, Stress, and Anhedonia: Toward a Synthesis and Integrated Model. Annu Rev Clin Psychol 393–423. https://doi.org/10.1146/annurev-clinpsy-050212-185606.Depression

Pizzagalli, D.A., 2011. Frontocingulate dysfunction in depression: Toward biomarkers of treatment response. Neuropsychopharmacology 36, 183–206. https://doi.org/10.1038/npp.2010.166

Ratcliff, R., 1978. A theory of memory retrieval. Psychol. Rev. 85, 59–108. https://doi.org/10.1037/0033-295X.85.2.59

Ravizza, S.M., Delgado, M.R., 2014. Motivational enhancement of cognitive control depends on depressive symptoms. Emotion 14, 646–50. https://doi.org/10.1037/a0036754

Rizvi, S.J., Pizzagalli, D.A., Sproule, B.A., Kennedy, S.H., 2016. Assessing anhedonia in depression: Potentials and pitfalls. Neurosci. Biobehav. Rev. 65, 21–35. https://doi.org/10.1016/j.neubiorev.2016.03.004

Romer Thomsen, K., Whybrow, P.C., Kringelbach, M.L., 2015. Reconceptualizing anhedonia: novel perspectives on balancing the pleasure networks in the human brain. Front. Behav. Neurosci. 9, 1–23. https://doi.org/10.3389/fnbeh.2015.00049

Rotter, J.B., 1966. Generalized expectancies for internal versus external control of reinforcement. Psychol. Monogr. 80, 1–28. https://doi.org/10.1037/h0092976

Salamone, J.D., Correa, M., 2012. The Mysterious Motivational Functions of Mesolimbic Dopamine. Neuron 76, 470–485. https://doi.org/10.1016/j.neuron.2012.10.021

Salamone, J.D., Yohn, S.E., Lopez-Cruz, L., San Miguel, N., Correa, M., 2016. Activational and effort-related aspects of motivation: Neural mechanisms and implications for psychopathology. Brain 139, 1325–1347. https://doi.org/10.1093/brain/aww050

Scheurich, A., Fellgiebel, A., Schermuly, I., Bauer, S., Wölfges, R., Müller, M.J., 2008. Experimental evidence for a motivational origin of cognitive impairment in major depression. Psychol. Med. 38, 237–246. https://doi.org/10.1017/S0033291707002206

Seligman, M.E., 1972. Learned helplessness. Annu. Rev. Med. 23, 407–412. https://doi.org/10.1146/annurev.me.23.020172.002203

Seligman, M.E.P., Maier, S.F., 1967. Failure To Escape Traumatic Shock. J. Exp. Psychol. 74, 1–9. https://doi.org/10.1037/h0024514

Shenhav, A., Botvinick, M., Cohen, J., 2013. The expected value of control: An integrative theory of anterior cingulate cortex function. Neuron 79, 217–240. https://doi.org/10.1016/j.neuron.2013.07.007

Shenhav, A., Cohen, J.D., Botvinick, M.M., 2016. Dorsal anterior cingulate cortex and the value of control. Nat. Neurosci. 19, 1286–1291. https://doi.org/1038/nn.4382

Shenhav, A., Musslick, S., Lieder, F., Kool, W., Griffiths, T.L., Cohen, J.D., Botvinick, M.M., 2017. Toward a Rational and Mechanistic Account of Mental Effort. Annu. Rev. Neurosci. 40, 99–124. https://doi.org/10.1146/annurev-neuro-072116-031526

Siegle, G.J., Ghinassi, F., Thase, M.E., 2007. Neurobehavioral therapies in the 21st century: Summary of an emerging field and an extended example of cognitive control training for depression. Cognit. Ther. Res. 31, 235–262. https://doi.org/10.1007/s10608-006-9118-6

Snyder, H.R., 2013. Major depressive disorder is associated with broad impairments on neuropsychological measures of executive function: a meta-analysis and review. Psychol. Bull. 139, 81–132. https://doi.org/10.1037/a0028727

Spijker, J., Bijl, R. V, de Graaf, R., Nolen, W. a, 2001. Determinants of poor 1-year outcome of DSM-III-R major depression in the general population: results of the Netherlands Mental Health Survey and Incidence Study (NEMESIS). Acta Psychiatr. Scand. 103, 122–30. https://doi.org/acp016 [pii]

Strauss, G.P., Gold, J.M., 2012. A new perspective on anhedonia in schizophrenia. Am. J. Psychiatry 169, 364–373. https://doi.org/10.1176/appi.ajp.2011.11030447

Stroop, J.R., 1935. Studies of interference in serial verbal reactions. J. Exp. Psychol. 18, 643–662. https://doi.org/10.1037/h0054651

Treadway, M.T., Bossaller, N.A., Shelton, R.C., Zald, D.H., 2012. Effort-based decision-making in major depressive disorder: A translational model of motivational anhedonia. J. Abnorm. Psychol. 121, 553–558. https://doi.org/10.1037/a0028813

Treadway, M.T., Zald, D.H., 2013. Parsing Anhedonia: Translational Models of Reward-Processing Deficits in Psychopathology. Curr. Dir. Psychol. Sci. 22, 244–249. https://doi.org/10.1177/0963721412474460

Treadway, M.T., Zald, D.H., 2011. Reconsidering anhedonia in depression: Lessons from translational neuroscience. Neurosci. Biobehav. Rev. 35, 537–555. https://doi.org/10.1016/j.neubiorev.2010.06.006

Uher, R., Perlis, R.H., Henigsberg, N., Zobel, A., Rietschel, M., Mors, O., Hauser, J., Dernovsek, M.Z., Souery, D., Bajs, M., Maier, W., Aitchison, K.J., Farmer, A., McGuffin, P., 2012. Depression symptom dimensions as predictors of antidepressant treatment outcome: Replicable evidence for interest-activity symptoms. Psychol. Med. 42, 967–980. https://doi.org/10.1017/S0033291711001905

Verguts, T., 2017. Computational models of cognitive control, in: Egner, T. (Ed.), The Wiley Handbook of Cognitive Control. Wiley-Blackwell, Hoboken, NJ, pp. 376–391.

Verguts, T., Vassena, E., Silvetti, M., 2015. Adaptive effort investment in cognitive and physical tasks: a neurocomputational model. Front. Behav. Neurosci. 9. https://doi.org/10.3389/fnbeh.2015.00057

Wallis, J.D., Rushworth, M.F.S., 2014. Integrating benefits and costs in decision making, in: Neuroeconomics (Second Edition). pp. 411–433.

Wardenaar, K.J., Giltay, E.J., Van Veen, T., Zitman, F.G., Penninx, B.W.J.H., 2012. Symptom dimensions as predictors of the two-year course of depressive and anxiety disorders. J. Affect. Disord. 136, 1198–1203. https://doi.org/10.1016/j.jad.2011.11.037

Westbrook, A., Braver, T.S., 2015. Cognitive effort: A neuroeconomic approach. Cogn. Affect. Behav. Neurosci. 15, 395–415. https://doi.org/10.3758/s13415-015-0334-y

Westbrook, A., Kester, D., Braver, T.S., 2013. What Is the Subjective Cost of Cognitive Effort? Load, Trait, and Aging Effects Revealed by Economic Preference. PLoS One 8, 1–8. https://doi.org/10.1371/journal.pone.0068210

Whitmer, A.J., Gotlib, I.H., 2013. An attentional scope model of rumination. Psychol. Bull. 139, 1036–61. https://doi.org/10.1037/a0030923

Whitton, A.E., Treadway, M.T., Pizzagalli, D.A., 2015. Reward processing dysfunction in major depression, bipolar disorder and schizophrenia. Curr. Opin. Psychiatry 28, 7–12.

Wigfield, A., Eccles, J.S., 2000. Expectancy–Value Theory of Achievement Motivation. Contemp. Educ. Psychol. 25, 68–81. https://doi.org/10.1006/ceps.1999.1015

Wood, W., Rünger, D., 2016. Psychology of Habit. Annu. Rev. Psychol. 67, 289–314. https://doi.org/10.1146/annurev-psych-122414-033417

Yang, X. hua, Huang, J., Zhu, C. ying, Wang, Y. fei, Cheung, E.F.C., Chan, R.C.K., Xie, G. rong, 2014. Motivational deficits in effort-based decision making in individuals with subsyndromal depression, first-episode and remitted depression patients. Psychiatry Res. 220, 874–882. https://doi.org/10.1016/j.psychres.2014.08.056

Zald, D.H., Treadway, M.T., 2017. Reward Processing, Neuroeconomics, and Psychopathology. Annu. Rev. Clin. Psychol 133, 1–325. https://doi.org/10.1146/annurev-clinpsy-032816-044957

Supplementary Materials

Simulations of control allocation

In order to simulate the effects of motivational variables on control allocation we generated behavior from an agent that performs cognitive control tasks[2] (Musslick et al., 2015). In these simulations, each trial is described as an interaction between the control system and the task environment. The agent specifies the optimal control signal based on an internal representation that is an estimate of the next trial (inferred state $\hat{S}$). The agent implements the specified control signal while performing the corresponding task in the actual task environment (actual state S). After each trial, the agent updates the inferred state for the next trial based on its observation of the current trial.

To generate response probabilities and reaction times we used a drift diffusion model (DDM, Bogacz et al., 2006; Ratcliff, 1978) that accumulates evidence towards one of two responses in a

---

[2] The source code for all simulations is available at https://github.com/musslick/EVCDepression

Stroop task (e.g. one response indicating the color green and the other response indicating the color red). We assume that the rate of accumulation toward one of the two response boundaries is governed by an automatic component and a controlled component

$$d = d_{CONTROL} + d_{AUOTMATIC}$$

where the automatic component reflects automatic processing of each of the color and word of the stimulus that is unaffected by control:

$$d_{AUTOMATIC} = a_{COLOR} + a_{WORD}$$

The absolute magnitude of the color-response association a_{COLOR}, as well as the magnitude of the word-response association a_{WORD} depends on the strength of the association of each stimulus feature with a given response and its sign depends on the response (e.g. $a_{COLOR} < 0$ if the response is associated with the left button, $a_{COLOR} > 0$ if response is associated with the right button). Thus, for congruent trials a_{COLOR}, and a_{WORD} have the same sign, whereas the opposite sign for incongruent (conflict) trials. The controlled component of the drift rate is the sum of the two stimulus values, each weighted by the efficacy ε with which cognitive control is implemented, as well as the intensity of the corresponding control signal, one for processing the color dimension of the stimulus u_{COLOR} and one for processing the word dimension of the stimulus u_{WORD}:

$$d_{CONTROL} = a_{COLOR} \cdot u_{COLOR} \cdot \varepsilon + a_{WORD} \cdot u_{WORD} \cdot \varepsilon$$

Thus, each control signal biases processing towards one of the stimulus dimensions. As a result, higher control signal intensity for processing the color dimension of the stimulus improves performance – speeds responses and lowers error rates – for the Stroop task. The simulations reported in this paper varied the inferred efficacy of control $\hat{\varepsilon}$ (thereby influencing decisions

about control allocation) but held the actual efficacy of control constant ($\varepsilon = 1$). Mean reaction

times (RTs) and response probabilities for a given parameterization of drift rate on trial t are

derived from an analytical solution to the DDM (Navarro and Fuss, 2009).

In order to specify the optimal set of control signals $\boldsymbol{U} = \{u_{COLOR}, u_{WORD}\}$ on a given trial, the

model estimates the expected value for each configuration of control signal intensities based on

its internal model of the next trial $\hat{S} = \{\hat{a}_{COLOR}, a_{WORD}\}$. This is done by weighting the expected

reward for an outcome against the cost associated with the chosen control signal configuration:

$$EVC(\boldsymbol{U}, \hat{S}) = P(Correct|\boldsymbol{U}, \hat{S}) \cdot V(R) - Cost(\boldsymbol{U})$$

where $P(Correct|\boldsymbol{U}, \hat{S})$ corresponds to the probability of reaching the decision threshold for the

correct response and $V(R)$ corresponds to the subjective value of responding correctly. Here, the

subjective value $V(R) = v \cdot R$ corresponds to the amount of reward offered for a correct response

R weighted by the model's sensitivity to the reward v. The total cost of cognitive control

$Cost(\boldsymbol{C})$ is computed as the sum of the costs for each control signal,

$$Cost(\boldsymbol{C}) = e^{c \cdot u_{COLOR}} + e^{c \cdot u_{WORD}}$$

where the cost for each control signal is an exponential function of the intensity of the control

signal, scaled by the cost parameter c. The model selects the control signal configuration with the

maximum EVC within the inferred next trial $\hat{S}$, out of all the configurations under consideration:

$$U^* \leftarrow argmax_i[EVC(\boldsymbol{U}, \hat{S})]$$

Performance in the actual state S is determined by the influence of the chosen control signals on

the true parameters of the DDM (e.g., by adjusting the drift rate).

Task environments

Stroop Paradigm. To illustrate effects of distractor interference in the EVC model, we simulated behavior on a Stroop task. In this task, the model is presented with a two-dimensional stimulus, one dimension representing an ink color and another dimension representing a color word. On each trial, the EVC model is required to indicate the response associated with the ink color. The trial sequence encompassed 200 trials, half of which were response congruent and half of which were response incongruent. To simulate congruent trials, we set $a_{COLOR} = 0.1, a_{WORD} = 0.33$ such that both stimuli dimension promote the same response. On incongruent trials, we set $a_{COLOR} = 0.1, a_{WORD} = -0.33$ such that the word dimension is associated with a different response than the color dimension. Note that the word-response association has a higher absolute magnitude than the color-response association, reflecting the assumption that word reading is a more automatic process than color naming. To simulate an expected mixture of congruent and incongruent trials, we parameterized the expected state as the average of the two trial conditions, $\hat{S} = \{\hat{a}_{COLOR} = 0.1, a_{WORD} = 0\}$. Control was implemented in the form of two control signals, one for processing the color dimension and one for processing the word dimension. The range of control signal intensities as varied from 0 to 10 in steps of 0.2 and the reward received for a correct response was set to $R = 40$. DDM parameters were set as follows: starting point = 0.0, noise coefficient = 0.5, non-decision time = 0.25s and threshold = 0.36. To demonstrate how control-demanding behavior is affected by changes in the decision-making process about control allocation we systematically varied the control cost parameter c from 0.2 to 0.3 in steps of 0.1, the reward sensitivity v from 0.5 to 1.5 in steps of 0.1 and the expected control efficacy $\hat{\varepsilon}$ from 0.5 to 1.5 in steps of 0.1 across simulations. Note that we varied only one parameter at a time while holding the other parameters constant at $c = 0.25, v = 1, \hat{\varepsilon} = 1$. For each parameter setting, we assessed the mean amount of control that the model allocated for each

trial type, as well as the mean error rate.

COGED Paradigm. To demonstrate the effects of changing model parameters on

demand-avoidance, we simulated behavior in the cognitive effort discounting (COGED)

experiment described by Westbrook & Braver (2015). In this paradigm, subjects can choose on

each trial whether they want to perform a baseline low-demand task for a low reward or a higher-

demand alternative task for a higher reward. The amount of reward offered for the baseline task is

adjusted to identify the point of indifference, that is, the reward at which subjects are indifferent

between performing the low-demand baseline task and performing the high-demand task. To

simulate this paradigm, we modeled both tasks as different types of trials that the model can

choose between. Each trial encompassed a stimulus with a color dimension that mapped to one of

two responses with $a_{COLOR} > 0$. However, unlike in the Stroop task described above there was no

word dimension, $a_{WORD} = 0$. The difficulty of the high-demand task was manipulated across

experiment blocks by varying the color-response association a_{COLOR} from 0.2 to 0.4 in steps of

0.0667 and the difficulty of the baseline task was fixed to $a_{COLOR} = 1$ (higher color-response

associations may reflect higher saturation values for a color patch). For each set of simulations,

we fixed the reward for the high-demand task to ($R = 200$) while steadily increasing the amount

of reward offered for the low-demand task in steps of 1, beginning from an initial reward value of

$R = 1$. On each trial, the EVC agent determined the highest EVC separately for each task and

chose the task with the highest predicted EVC. We then assessed the amount of reward offered

for the low-demand task for which the model would be indifferent between performing the low-

demand task and the (more rewarding) high-demand task and normalized this value by the

amount of reward offered for the high-demand task. Following the notation by Westbrook &

Braver (2015), we refer to this reward value as the subjective value of completing the high-

demand task. For instance, if the model would switch to performing the low-demand task at an

offered reward of 120 then the (discounted) subjective value of the high-demand task would be

120. The range of control signal intensities was varied from 0 to 4 in steps of 0.05 and DDM

parameters were set as follows: starting point = 0.0, noise coefficient = 0.8, non-decision time =

0.25s and threshold = 1. To demonstrate how control-demanding behavior is affected by changes

in the decision-making process about control allocation, we systematically varied the control cost

parameter c from 1 to 1.5 to 2, the reward sensitivity v from 0.5 to 1.0 to 2.0 and the expected

control efficacy $\hat{\varepsilon}$ from 1 to 3 to 5 across simulations. Note that we varied only one parameter at a

time while holding the other parameters constant at $c = 1.5, v = 1, \hat{\varepsilon} = 1$. For each parameter

setting, we assessed the subjective value as a function of the task difficulty of the high-demand

task.