Medial prefrontal cortex activity reflects ordinal retrieval modes and hippocampal activity reflects temporal context retrieval modes

Running Head: SEQUENCE MEMORY RETRIEVAL MODES IN THE HUMAN BRAIN

Puck C. Reeders[1], Amanda G. Hamm[1], Timothy A. Allen[1,2,3], Aaron T. Mattfeld[1,3]

[1]Cognitive Neuroscience Program, Department of Psychology, Florida International University, Miami, FL, 33199

[2]Department of Environmental Health Sciences, Florida International University, Miami, FL, 33199

[3]Center for Children and Families, Florida International University, Miami, FL, 33199

Title: 118 characters of 120 characters

Number of text pages: 28

Number of figures: 4

Word/Character count in:

Title: 118 Characters (120 characters max)
Abstract: 147 words (150 words max)
Significance Statement: 127 words
Introduction: 747 words (incl citations)
Discussion: 16653 words (incl citations)
Conclusion: 82 words

Corresponding Author:

Aaron T. Mattfeld, PhD
Department of Psychology
Florida International University
11200 SW 8th Street
Miami, FL 33199
email: amattfel@fiu.edu

Keywords: medial prefrontal cortex, hippocampus, declarative memory, sequence memory, ordinal, episodic memory, retrieval

1   Conflict of Interest: The authors declare no competing financial interests.

1   **Acknowledgments.** The research was conducted with funds provided by FIU to TAA and ATM, and

2   the Feinberg Foundation to TAA. We thank Adam Kimbler for his help with coding, and Dr. Leila Allen

3   for useful feedback on the manuscript.

ABSTRACT

Remembering sequences of events defines episodic memory, but retrieval can be driven by both ordinality and temporal contexts. Whether these modes of retrieval operate at the same time or not remains unclear. Theoretically, medial prefrontal cortex (mPFC) confers ordinality, while the hippocampus (HC) associates events in gradually changing temporal contexts. Here, we looked for evidence of each with BOLD fMRI in a sequence task that taxes both retrieval modes. To test ordinal modes, items were transferred between sequences but retained their position (e.g., AB3). Ordinal modes activated mPFC, but not HC. To test temporal contexts, we examined items that skipped ahead across lag distances (e.g., ABD). HC, but not mPFC, tracked temporal contexts. There was a mPFC and HC by retrieval mode interaction. These current results suggest that the mPFC and HC are concurrently engaged in different retrieval modes in support of remembering *when* an event occurred.

Significance Statement

Memory for sequences of events is a defining aspect of everyday episodic memory allowing our brain to separate unique experiences that otherwise have overlapping sensory and spatial content. Sequence memory is impaired in typical aging and in disorders such as Alzheimer's disease. The results of the current study provide new evidence that two retrieval modes concurrently arise during sequence memory, and they have distinct neural correlates. The medial prefrontal cortex contributes to an ordinal retrieval mode, while at the same time, the hippocampus contributes a gradually-changing temporal context mode of retrieval. These data shed new light on why typical episodic memory requires both the medial prefrontal cortex and the hippocampus, and suggests a functional dissociation between the medial prefrontal cortex and hippocampus across these modes of retrieval.

Introduction

Memory for sequences of events is a fundamental component of episodic memory (Tulving

1984; 2002; Allen and Fortin, 2013; Howard and Eichenbaum, 2013; Eichenbaum, 2017). While

different experiences share overlapping elements, the sequence of events is unique. Remembering the

order of events allows us to disambiguate episodes with similar content and make detailed predictions

supporting decision making.

At least two complementary memory processes contribute to the retrieval of events in the

correct sequence: ordinal (Orlov et al., 2002) and temporal context (Howard and Kahana, 2002)

retrieval modes. Whether these disparate retrieval modes operate coincidently or not remains an open

question with consequences for understanding basic mechanisms of how we remember the events that

unfold throughout our day. According to an ordinal retrieval mode, items are remembered by their

position within an event sequence (Dubrow and Davachi, 2013; Allen et al., 2014; Long and Kahana,

2019) providing for sequential memory through well-established semantic or abstracted relationships

(1st then 2nd, etc.). While for a temporal context retrieval mode, events are remembered through a

gradually changing temporal context within which specific items have been associated. According to

temporal context, when an element of a sequence is presented or retrieved (e.g., 'C' in AB**C**DEF),

items that are close in the sequence (e.g. the 'D' in the sequence) have a higher retrieval rate

compared to items that are further away (e.g. the 'F' in the sequence). These temporal contexts result

from item associations that are dependent on time varying neural activity (e.g., Eichenbaum, 2014), and

contribute to sequence memory through the reactivation of neighboring items during retrieval (Dubrow

and Davachi, 2013; Long and Kahana, 2019).

The medial prefrontal cortex (mPFC) and hippocampus (HC) are thought to contribute to

sequence memory through ordinal representations and temporal contexts, respectively (Agster et al.,

2002; Fortin et al., 2002; Kesner et al., 2002; DeVito and Eichenbaum, 2011; Allen et al., 2016; Jenkins

and Ranganath, 2016). In rodents, mPFC disruptions impair sequence memory (DeVito et al., 2011;

Jayachandran et al., 2019), mPFC 'time cells' are evident (Tiganj et al., 2017), and positions within a

sequence can be the main determinant of differential activity in mPFC neurons during spatial

sequences (Euston and McNaughton, 2006). In humans, mPFC activation is sensitive to temporal order

memory (Preston and Eichenbaum, 2013), and codes for information about temporal positions within

image sequences irrespective of the image itself (Hsieh and Ranganath, 2015). HC activations are also

generally associated with temporal order memory (Kumaran and Maguire, 2006; Ekstrom and

Bookheimer, 2007; Lehn et al., 2009; Ross et al., 2009; Jenkins and Ranganath, 2010; Tubridy and

Davachi, 2011; Kalm et al., 2013; Hsieh et al., 2014; Goyal et al., 2018). The HC binds events within

temporal contexts (Eichenbaum et al., 2007; Dubrow and Davachi, 2013; Bladon et al., 2019) through a

gradually changing neural context (Manns et al., 2007; Mankin et al., 2012). Similarly, medial temporal

lobe neuronal and BOLD activations in humans have demonstrated evidence for gradually evolving

temporal contexts (Howard et al., 2012; Kalm et al., 2013; Kragel et al., 2015).

Here we tested the contributions of the mPFC and HC during a visual sequence memory task

that provides behavioral evidence of both ordinal and temporal context retrieval modes (Allen et al.,

2014). In the task, these two retrieval modes are parsed using probe trials that place conflicting

demands on ordinal (Allen et al., 2014; 2015; Orlov et al., 2000) and temporal context modes

(Jayachandran et al., 2019). We first evaluated ordinal retrieval modes using items that were

transferred from one sequence to another while retaining their ordinal position. Evidence for an ordinal-

based retrieval mode occurs when these probes are identified as in sequence because they occur in

the same ordinal position of their original sequence. mPFC activations (but not HC) was strongest for

these ordinal retrievals, and did not reflect general task accuracy. Second, we evaluated a temporal

context retrieval mode using items that skipped ahead with shorter lag distances (ABC**F**EF) to larger

lag distances away from their position (A**F**CDEF). HC activations (but not mPFC) tracked with lag

distance, providing evidence the HC reflects a temporal context-based retrieval mode. In fact, a

significant interaction was observed such that mPFC and HC differentially activated for ordinal and

temporal context retrievals. Our data shows that sequence memory involves both retrieval modes. In

line with these results, we suggest that understanding episodic memory requires more insight into the

neurobiology of ordinal processing in mPFC, in addition to the more often studied temporal contexts in

HC.

Materials and Methods

Participants

Thirty-nine right-handed volunteers were recruited from Florida International University (FIU)

and University of Miami to perform a magnetic resonance imaging (MRI) study that included a

sequence memory task designed to investigate the ability of humans to learn and remember arbitrary

sequences of items, and a temporal reward discounting task. Task order was counterbalanced across

all participants. Here we report results from the sequence memory task. All participants provided written

consent in compliance with the local Institutional Review Board. Five participants were excluded from

the final analysis due to failure to complete the task (n = 1) or because of poor performance (n = 4;

sequence memory index score $\leq$ 0; see Sequence Memory Analysis section below). Participants

excluded for poor performance had d-prime scores 2 standard deviations below the mean (M = 2.030,

SD = 0.807). The final sample consisted of 34 individuals (19 females; mean age = 21 years, SD = 2).

Task Apparatus

18   The sequence task was run on a Dell computer using Matlab (R2015b) with custom scripts that

19 included functions from Psychtoolbox (Psychtoolbox-3 distribution; http://www.psychtoolbox.org).

20 Images were back-projected and viewed by participants with an angled mirror mounted on the head

21 coil. Responses were recorded using a Current Designs MR-compatible 4-button inline response

22 device (https://www.curdes.com).

Experimental Design and Statistical Analysis

Prescan Training

All participants began with a practice session to become acquainted with the structure of the

task and the method of responding. During the practice session, participants viewed four low-memory

demand sequence sets, each comprising six unique images: (1) individual arrows at 0°, 60°, 120°,

180°, 240°, and 300°, presented in a clockwise fashion (Fig1A, Seq1); (2) a dot moving from the upper

left to the lower right corner (Fig. 1A, Seq2); (3) bars of different colors moving from left to right (not

shown); and (4) letters A, B, C, D, E, and F positioned in the center of the screen (not shown).

Participants were asked to memorize sequences after a single "study" presentation during which the

items were passively viewed. Later the practice sequences were tested with 15 unique memory probes

per sequence. Testing was self-paced; each sequence was preceded by a screen with the words

"Press the button to begin". To initiate an image in the sequence, participants were required to press

and hold a button. If the image was in sequence (InSeq), participants were instructed to hold down the

button until the image disappeared on its own at 1 s (the decision threshold), after which they could

release the button. If the item was out of sequence (OutSeq), participants were instructed to release the

button prior to the decision threshold (<1 s), at which point the image would disappear upon button

release. Prescan training was conducted on a Dell desktop computer.

Sequence Memory Task

During the Sequence Memory Task, participants were presented with two sequences from

prescan training (*low-memory demand;* Fig. 1A, Seq 1-2), and four novel sequences consisting of six

unique fractal images (*high-memory demand;* Fig. 1A, Seq 3-6). The exact composition of the novel

fractal sequence sets was different for each participant. Sets were selected randomly, without

replacement, from a bank of 240 unique fractal images. Similar to the prescan training, participants

were asked to view and memorize the sequences after a single presentation in a passive viewing

phase. Following the passive viewing phase, sequences were presented in a pseudorandom order,

participants made judgments as to whether each item in a sequence was presented InSeq or OutSeq.

Testing was self-paced and

followed the same structure as

the prescan training. The words

"Press the button to begin"

preceded the beginning of each

sequence. A button depression

with the right index finger

initiated the presentation of

each image in a sequence. If

the participants judged the

current image to be InSeq, they

would hold the button until the

decision threshold (1 s, when

the image disappeared), after

which they could release the

button. If the image was

OutSeq, they were instructed to

release the button before the 1

s decision threshold. Sequence

order was determined by the

Figure 1*. Sequence memory task and overall performance levels*. Participants were tested on a sequence memory task that pressures different retrieval modes using different out of sequence probe trial types. **A,** participants were presented sequences, one item at a time, to study before being tested on whether an item was in sequence or out of sequence. The total sequence set included 6 sequences, from which two sequences were *low memory demand* sequences (Seq 1-2), which were easy, predictable sequences that could be solved using a rule-based strategy. The other four sequence were *high memory demand* sequences (Seq 3-6), that required memorization of the items during the study phase. **B,** there were three out of sequence probe trials: Items that were repeated in the sequence (Repeats), items that were presented too early in the sequence (Skips), and items that transferred from one sequence to another, while remaining in their ordinal position (Ordinal Transfers). Repeats and Skips occurred throughout the whole task, whereas Ordinal Transfers occurred during the second half to avoid participants relying solely on ordinal strategies. **C,** participants performed best on identifying Repeats, then Skips and then Ordinal Transfers. For all three probe trial types, learning was rapid (asymptotic within a few trials), and performance was steady throughout the experiment. Thus, performance in the task and the accompanying brain imaging data primarily reflect memory and the associated retrieval modes.

following rules: (1) each sequence was presented first with all images in the correct order. (2) In the first

half of testing (first 120 sequences), OutSeq items were either the same image appearing twice

(Repeats) or an item appearing too early (Skips; see probe trial description below). In the second half

of testing (second 120 sequences), OutSeq items could now include images appearing from a different

sequence, but in the correct ordinal position (Ordinal Transfers). Ordinal Transfers were introduced

later so participants would not adopt an explicit ordinality strategy at the outset of the study given the

dominance of ordinal retrieval modes early in sequence memory (Orlov et al., 2000). The six

sequences were presented 40 times each, for a total of 240 sequence presentations. Half of the total

sequences contained one OutSeq and five InSeq images; the remaining consisted only of correctly

sequenced items. InSeq and OutSeq sequence sets were randomly presented throughout testing. The

Sequence Memory Task consisted of 15 min blocks of continuous performance separated by a brief

break (<1 min) to provide participants a rest. The number of blocks was dependent on the pace of the

participant. Out of all the analyzed participants, most participants completed the task in four blocks (n =

29), while the remainder finished in either three (n=1) or five (n=4).

OutSeq Probe Trials

Three distinct types of OutSeq probe trials were used during the test phase: Ordinal Transfers,

Repeats (reverse lags) and Skips (forward lags). OutSeq items were counterbalanced across sequence

sets, never presented in first position (Pos1), and each OutSeq instance was unique. Due to the nature

of the OutSeq trials, there could not be equal distribution of probe trials across all possible positions,

and occurred as follows: Pos1, 0%; Pos2, 7.5%; Pos3, 17,5%; Pos4, 50%; Pos5, 17.5%; Pos6, 7.5%.

Ordinal Transfers. OutSeq trials where an item from one sequence was 'transferred' to a

different sequence while retaining its correct ordinal position were considered Ordinal Transfers (Fig.

1B, *Ordinal Transfer*). For example, consider two sequences consisting of ABCDEF and UVWXYZ. An

ordinal transfer into the first sequence might have Y at the 5th position ABCD$\underline{Y}$F. While Y occupies its

original 5th position, it otherwise does not belong to the current sequence (Y does not normally follow

D). Distribution of Ordinal Transfers across all possible positions: Pos1, 0%; Pos2, 7.5%; Pos3, 17,5%;

Pos4, 50%; Pos5, 17.5%; Pos6, 7.5%.

Repeats (reverse lags). An OutSeq item was considered a Repeat if the image previously

appeared in the current sequence and was repeated (Fig. 1B, *Repeat*). Repeats occurred at multiple

lag distances, represented as negative values from repeat presentation to the original presentation,

ranging from -2 (e.g. ABC$\underline{B}$EF) to -5 (e.g. ABCDE$\underline{A}$). A lag distance of -1 was not used due to the lack

of an intervening item. Distribution of Repeats across all possible positions: Pos1, 0%; Pos2, 0%; Pos3, 7.5%; Pos4, 50%; Pos5, 27.5%; Pos6, 15%.

Skips (forward lags). OutSeq images were considered Skips when presented too early in the sequence (e.g. AB<u>E</u>DEF; Fig. 1B, *Skip*). Skips occurred at all lag distances, represented as positive values from early presentation to original presentation, ranging from +1 (e.g. ABCD<u>F</u>F) to +4 (e.g. A<u>F</u>CDEF). Distribution of Skips across all possible positions: Pos1, 0%; Pos2, 15%; Pos3, 27.5%; Pos4, 50%; Pos5, 7.5%; Pos6, 0%.

Sequence Memory Analysis

To evaluate whether participants demonstrated sequence memory, we compared the observed and expected frequencies of InSeq and OutSeq responses using *G*-tests (Sokal and Rohlf 1995). Responses to each item were sorted into a 2 x 2 matrix based on accuracy (correct/incorrect) and sequence condition (InSeq/OutSeq), as done previously (Allen et al., 2014; Jayachandran et al., 2019). For the sequence memory analysis, only "high memory" sequence sets were included. Responses to the first item of each sequence were excluded from analysis. Ordinal Transfers were excluded from this analysis because they were used to parse retrieval modes rather than measure performance. For comparison we evaluated sequence memory using d-prime, a measure of memory specificity derived from signal detection theory. Hits were defined as InSeq trials that were correctly identified as InSeq, misses were InSeq trials that were incorrectly identified as OutSeq, correct rejections were OutSeq probe trials that were correctly identified as OutSeq, and false alarms were defined as OutSeq probe trials that were incorrectly judged to be InSeq. The same conclusions were drawn with both approaches, thus, here we only report the results from the G-tests given its robustness to response biases as the current task has an overall greater number of InSeq responses.

Additionally, we examined overall sequence memory performance, using a summary statistic called the Sequence Memory Index (SMI; Equation 1). SMI normalizes the proportion of InSeq and OutSeq items across different conditions and represents sequence memory performance as a single

value ranging from -1 to 1. A SMI of "1" represents perfect sequence performance (response time > 1 s

for all InSeq items and < 1 s for all OutSeq items), and 0 represents chance performance. Note that an

SMI score of -1 would indicate an incorrect response to every single item (response times < 1 s for all

InSeq items and > 1 s for all OutSeq items). Negative SMIs rarely occurred, poor performance was

typically captured by SMI scores close to zero. SMI was calculated for both low and high memory

demand sequences and has been used in previous human studies to facilitate comparisons (Allen et

al., 2014, 2015).

Equation 1. $SMI = \dfrac{(0.9\text{Incor})(0.1\text{Outcor}) - (0.9\text{Ininc})(0.1\text{Outinc})}{\sqrt{(0.9\text{Incor} + 0.9\text{Ininc})(0.1\text{Outcor} + 0.1\text{Outinc})(0.9\text{Incor} + 0.1\text{Outinc})(0.9\text{Ininc} + 0.1\text{Outcor})}}$

Detailed Position and Lag Analysis

To evaluate strategies or mechanisms of sequence memory performance, a detailed analysis

based on ordinal position and lag distance was performed for each OutSeq probe type (Ordinal

Transfers, Skips, and Repeats). For Ordinal Transfers, we evaluated performance across positions

(Pos2 thru Pos6). For Skips, we evaluated performance across n-forward lag distances (Fig. 3A; n-

forward lags: +1, +2, +3, and +4). Smaller n-forward lags occurred more often because more

combinations were available. For Repeats, we evaluated performance across the n-reverse lags (Fig.

3A; n-reverse lags: -2, -3, -4, and -5). Smaller n-reverse lags occurred more often because more

combinations were available. We compared each position or lag distance performance (accuracy and

SMI) using repeated-measures ANOVAs followed by one-sample t-tests. To account for the potential of

a response bias (the assumption by participants of each item being in sequence) in the accuracy

measures, we calculated an adjusted chance level as follows: (1) we calculated the observed response

bias for holding the button for >1 sec on Pos 2 through Pos 6, irrespective of the InSeq or OutSeq

status, which was 90.466%, and (2) we calculated the complement of that response bias and set that

as the chance response for probe trials, which was 9.534%. To assess differences in variability

amongst probe trial types trial specific performance was converted to a z-score across positions

(Ordinal Transfers) and lags (Skips and Repeats) for each participant. Three mixed general linear

models were performed with subject as a random effect and position or lag as a fixed effect. The

resulting residuals were subsequently squared and averaged across positions or lags resulting in probe

trial type specific residuals for each participant. We conducted a repeated measures ANOVA with

Greenhouse-Geisser corrections and LSD posthoc testing to compare the averaged squared residuals

across probe trial types (Ordinal Transfers, Skips and Repeats).

Neuroimaging Acquisition

Neuroimaging data were collected on a General Electric Discovery MR750 3.0T scanner using a

32-channel head coil at the University of Miami Neuroimaging Facility. Structural T1-weighted images

were collected (186 slices, flip angle = 12°; TE = 3.68 ms, TR = 9.184 ms, TI = 650 ms, matrix = 256 x

256 mm; FOV = 256 mm, slice thickness = 1.0 mm). Whole-brain T2*-weighted, blood oxygen level

dependent (BOLD) echo-planar imaging data (42 slices, interleaved, bottom-up; flip angle = 75°; TE =

25 ms; TR = 2000 ms; matrix = 96x96 mm; FOV = 240 mm, slice thickness = 3.0 mm, voxel size = 2.5

mm^2) were collected during the task.

Neuroimaging Preprocessing

Preprocessing was performed using a pipeline developed in Nipype v0.1 (Gorgolewski et al.,

2011), wrapping tools from Analysis of Functional Neuroimages (AFNI v16.3.18; Cox, 1996), FSL

(v5.0.10), FreeSurfer (v5.1.0; Dale et al., 1999), Advanced Normalization Tools (v2.2.0; Avants et al.,

2008), and Artifact Detection Tools (ART; www.nitrc.org). Following DICOM conversion, cortical surface

reconstruction and cortical/subcortical segmentation was performed on all T1-weighted structural

scans. Functional data were first 'despiked' to remove and replace intensity outliers in the functional

time series. The data were simultaneously slice-time and motion corrected (Roche, 2011). An affine

transformation matrix was calculated during coregistration of the mean of each participant's functional

scans to their structural scan using Freesurfer's boundary-based registration algorithm (BBregister).

Brain masks were created by binarizing the aparc+aseg file and dilating by one voxel. The resulting

binary masks were subsequently coregistered to the functional data by applying the inverse of the

affine coregistration matrix. Motion and intensity outliers were then identified using the rapid art artifact

detection tool as implemented in nipype. Time-points at which intensity either exceeded three standard

deviations or composite frame-wise displacement was greater than 1 mm were flagged as outliers to

serve as subsequent regressors of no interest in the first-level general linear models. Finally, functional

data were spatially filtered with a 5 mm FWHM maximum Gaussian kernel using the *SUSAN* algorithm

(FSL).

Neuroimaging Normalization

A study-specific template was generated using Advanced Normalization Tools (ANTs). Each

structural scan was skull-stripped by multiplying the T1-weighted structural scan by the binarized and

dilated aparc+aseg file in structural space. Each skull-stripped brain was then rigid-body transformed

(no scaling or shearing) to Montreal Neurological Institute space using FSL's *FLIRT* algorithm. This first

pass was used to minimize large spatial shifts between participants and generate a template close to a

commonly used reference. Following visual inspection, a study template was created using the

buildtemplateparallel.sh script from ANTs. After template generation, each participant's skull-stripped

brain was normalized using non-linear symmetric diffeomorphic mapping implemented by ANTs. The

resulting warps were applied to contrast parameter estimates following fixed-effects modeling for

subsequent group-level tests.

Neuroimaging Analysis

Functional data were analyzed according to a general linear model approach using FMRIB's

Software Library (www.fmrib.ox.ac.uk/fsl). Two separate univariate general linear models were used for

first-level analyses: (1) a performance (correct/incorrect) model; and (2) a lag model.

All first-level models included event and nuisance regressors. The performance analysis

contained eight event regressors of interest: correct and incorrect InSeq, Repeats, Skips, and Ordinal

Transfers. Event regressors were convolved with FSL's double gamma hemodynamic response

function with an onset beginning at the first stimulus item of a sequence and a duration equal to the

length of time required to evaluate all the items in that sequence (mean = 9.917 s per subject). The lag

analysis comprised six event regressors of interest: Repeats at different n-reverse lags (n-reverse lags:

-2, -3, and -4), and Skips at different n-forward lags (n-forward lags: +1, +2, and +3). InSeq and Ordinal

Transfers irrespective of behavioral performance were also included. We evaluated linear changes in

activation across lags with the following contrast weights: -1, 0, 1 and 1, 0, -1 for Repeats (n-reverse

lags: -2, -3, and -4) and Skips (n-forward lags: +1, +2, and +3). We did not include a lag of -5 or +4, as

the numbers of trials with these lags per participant was not sufficient (1 trial). Regressors for baseline

or low memory sequences were also included in both models. Nuisance regressors included motion

parameters (x, y, z translations; pitch, roll, yaw rotations), first and second derivatives of the motion

parameters, normalized motion, first, second, and third order Lagrange polynomials, as well as each

outlier time-point that exceeded the artifact detection thresholds identified during preprocessing.

Following the first-level analysis, a fixed effects analysis across experimental runs was

performed for each participant for the respective contrasts of interest (e.g., performance analysis:

correct versus incorrect probe trials; lag analysis: linear contrasts). Contrast parameter estimates from

the fixed effects analysis were normalized to the study-specific template and group-level analyses were

performed using FSL's *randomise* threshold-free cluster enhancement (tfce) one sample t-test. To test

a priori hypotheses with respect to the functional contributions of the mPFC and HC during sequence

memory retrieval, we constrained our voxel-wise analyses at the group level to the bilateral mPFC and

medial temporal lobe using masks of these regions. Whole-brain exploratory analyses were used to

follow up our anatomically directed tests. In the performance analysis, some participants were missing

key events of interest (e.g., correct Ordinal Transfers or incorrect Repeat probes). These participants

were not included in the relevant analyses, reducing the sample size for the correct versus incorrect

Ordinal Transfer contrast to n = 26, and for the correct versus incorrect repeat probe trial contrast to n =

31.

To directly examine the relation between sequence memory retrieval mode and regional

contribution, as a post hoc follow up to our voxelwise analyses, we used an anatomical region of

interest analysis to perform a 2x2 repeated measures factorial ANOVA with brain region (right anterior

HC vs. mPFC) and strategy (ordinal retrieval mode [Ordinal Transfer incorrect > Ordinal Transfer

correct] vs. temporal context retrieval mode [TCM; Positive Linear Skip contrast]) as the within subjects

factors and participants as the repeated measure. Anterior HC region of interest was manually drawn in

coronal slices. The anterior most boundary of the hippocampus was defined by the white matter

separating the hippocampus from the amygdala, the lateral and medial boundaries were defined by the

cerebral spinal fluid of the lateral ventricle, the inferior boundary was white matter of the

parahippocampal gyrus, while the superior boundary was the wavelike contour of the pes

digitations/alveus/horizontal line connecting the middle of the medial border of the lateral ventricle to

the surface of the uncus. The anterior HC was delineated from the posterior HC by the presence of a

small anatomical protrusion of medial HC into the lateral ventricle that was absent in the posterior HC –

uncal apex. The mPFC region of interest was created by binarizing the medial orbitofrontal, superior

frontal, rostral anterior cingulate, and the caudal anterior cingulate Freesurfer labels.

Results

Sequence Memory

We first looked at overall sequence memory in the task by comparing performance on low and

high memory demand sequence sets. We calculated an SMI separately for the low and high memory

demand sequences (Fig. 1A). As expected, participants performed significantly better than chance (SMI

= 0) on both low memory sequences (SMI_{low}: 0.751 ± 0.173; $SMI_{low\ vs.\ chance}$: $t_{(33)}$ = 37.065, p = 3.487 x

10^{-23}) and high memory sequences (SMI_{high}: 0.582 ± 0.170; $SMI_{low\ vs.\ chance}$: $t_{(33)}$ = 19.690, p = 5.391 x 10^{-20}

^{20}), but performance was significantly better with low memory sequences ($SMI_{low\ vs.\ high}$: $t_{(33)}$ = -4.839, p =

2.958 x 10^{-5}). High memory sequences were used for all subsequent analyses because they optimally

taxed the different sequence retrieval modes. Importantly, there was no testing order effect (1st or 2nd

experimental block) on sequence memory ($SMI_{1st\ block}$ = 0.571 ± 0.190, $SMI_{2nd\ block}$ = 0.594 ± 0.152;

$SMI_{1st\ vs.\ 2nd}$: $t_{(32)}$ = -0.386, p = 0.702), nor did we observe any sex differences (SMI_{female} = 0.566 ± 0.197,

SMI_{male} = 0.602 ± 0.133; $SMI_{female\ vs.\ male}$ = $t_{(32)}$ = -0.612, p = 0.545). Thus, we pooled these groups.

Next, we evaluated overall performance across each of the three memory probes individually

using percent correct for ease of interpretation. As in previously published studies, we found that

Ordinal Transfers (Accuracy = 34.927 ± 4.120%) were the most difficult, followed by Skips (Accuracy =

49.205 ± 2.686%), with Repeats being the easiest (Accuracy = 78.584 ± 2.853%). There was a

significant difference in performance across probe types ($F_{(2,66)}$ = 85.714, P = 3.433 x 10^{-14}) suggesting

differences in the available retrieval modes driven by the different conditions. Post hoc pairwise

comparisons showed that the performance on Repeats was significantly higher compared to Skips

(mean difference = 0.294 ± 0.020, p = 8.137 x 10^{-16}) and Ordinal Transfers (mean difference = 0.437 ±

0.039, p = 1.078 x 10^{-12}) and performance on Skips was significantly higher compared to Ordinal

Transfers (mean difference = 0.143 ± 0.039, p = 0.001). The same pattern of results was evident when

using SMI which controls for idiosyncratic response patterns. Importantly, participants performed each

of the three probe types significantly better than chance (Ordinal Transfers: $t_{(33)}$ = 6.164, p = 5.951 x 10^{-7}

^{7}, Repeats: $t_{(33)}$ = 24.201, p = 1.403 x 10^{-22}; Skips: $t_{(33)}$ = 14.771, p = 4.203 x 10^{-16}). For all probe types,

learning was rapid (asymptotic within a few trials), and performance was steady throughout the duration

of the experiment (Fig. 1C). Thus, performance in the task and the accompanying brain imaging data

primarily reflect distinct memory retrieval modes driven by specific probe conditions.

Ordinal Memory Retrievals

An ordinal retrieval mode is known to contribute to memory for sequences of events which we

tested for here using Ordinal Transfer probe trials (e.g., A**2**CDEF, Fig. 2A). If participants *exclusively*

used an ordinal retrieval mode on these trials (e.g., A goes in the 1^{st} position, B goes in the 2^{nd} position,

etc.), then these probes would be remembered as being InSeq (since "2" is in the same position as B).

An ordinal retrieval mode would be expected to drive performance on these trials to very low levels,

possibly to chance, if no other retrieval mode was engaged. Moreover, if an ordinal retrieval mode was

used, we would not expect differences in accuracy as a function of transfer position. Conversely, if

participants relied on another process, such as a temporal context, then Ordinal Transfers would be

easily identified as OutSeq (since "2" does not follow A, and in fact it doesn't go with any of the items in

that sequence). Non-ordinal retrieval modes would drive performance to very high levels on Ordinal

Transfer probe trials, probably to the same levels as at Repeats (as a good empirical benchmark for

asymptotic performance).

We found Ordinal Transfers were performed significantly better than chance across all positions

(Fig. 2B, red stars; Pos2: 0.314 ± 0.357; Pos2$_{vs. chance}$: $t_{(33)} = 3.566$, $p = 0.001$; Pos3: 0.338 ± 0.313;

Pos3$_{vs. chance}$: $t_{(33)} = 4.512$, $p = 7.712 \times 10^{-5}$; Pos4: 0.347 ± 0.258; Pos4$_{vs. chance}$: $t_{(33)} = 5.670$, $p = 2.549 \times$

10^{-6}; Pos5: 0.354 ± 0.283; Pos5$_{vs. chance}$: $t_{(33)} = 5.334$, $p = 6.892 \times 10^{-6}$; Pos6: 0.373 ± 0.336; Pos6$_{vs. chance}$:

$t_{(33)} = 4.811$, $p = 3.212 \times 10^{-5}$) suggesting participants utilized non-ordinal retrieval modes for these

trials. However, performance was much lower than Repeats (Fig. 2B, purple stars; Pos2$_{vs. Repeats}$: $t_{(33)} = -$

7.711, $p = 7.019 \times 10^{-9}$; Pos3$_{vs. Repeats}$: $t_{(33)} = -8.338$, $p = 1.224 \times 10^{-9}$; Pos4$_{vs. Repeats}$: $t_{(33)} = -9.917$, $p =$

1.996×10^{-11}; Pos5$_{vs. Repeats}$: $t_{(33)} = -8.919$, $p = 2.621 \times 10^{-10}$; Pos6$_{vs. Repeats}$: $t_{(33)} = -7.175$, $p = 3.185 \times 10^{-8}$)

and lower than Skips (Fig. 2B, blue stars; Pos2$_{vs. Skips}$: $t_{(33)} = -2.912$, $p = 6.390 \times 10^{-3}$; Pos3$_{vs. Skips}$: $t_{(33)} = -$

2.870, p = 7.104 x 10^{-3}; $Pos4_{vs.\ Skips}$: $t_{(33)}$ = -3.284, p = 2.426 x 10^{-3}; $Pos5_{vs.\ Skips}$: $t_{(33)}$ = -2.854, p = 7.400 x

10^{-3}; $Pos6_{vs.\ Skips}$: $t_{(33)}$ = -2.075, p = 4.589 x 10^{-2}) suggesting a heavy reliance on an ordinal retrieval

mode. We found no significant difference in performance across positions (Fig. 2B; $F_{(4,\ 132)}$ = 0.331, p =

0.786). The observed behavioral performance suggests that participants were disproportionally using

an ordinal retrieval mode (e.g., lower performance when compared to Repeats and Skips) in

combination with other non-ordinal retrieval modes (i.e., still at performance better than chance) to a

lesser extent. A prediction from the these conflicting retrieval modes is that performance variability

would be high between participants and across positions. In line with this hypothesis we observed

higher Ordinal Transfer variability when compared to Repeats, and much higher when compared to

Skips (Fig. 2C-D; $M_{residual}$ = 0.795 ± 0.074). A repeated-measure ANOVA with Greenhouse-Geisser

correction showed significant main effect among regression residuals of Ordinal Transfers, Skips and

Repeats ($F_{(2,\ 54)}$ = 23.866, p = 1.843 x 10^{-8}). Regression residuals of Ordinal Transfers were significantly

higher compared to Skips (mean difference = 0.530 ± 0.089, p = 2.523 x10^{-6}) and Repeats (mean

difference = 0.158 ± 0.028, p = 5.627x10^{-6}). Regression residuals of Repeats were significantly higher

compared to Skips (mean difference = 0.373 ± 0.099, p = 0.001).

mPFC Activations and Ordinal Retrievals

We hypothesized that using an ordinal retrieval mode would be related to activations in the mPFC (e.g.,

Hsieh and Ranganath, 2015). Algorithmically, this would be evident when contrasting undetected (i.e.,

using an ordinal retrieval mode) versus detected (i.e., using a non-ordinal retrieval mode) during

Ordinal Transfer trials. We also reasoned that the use of an ordinal retrieval mode would facilitate the

detection of Skips and Repeats (because of the ordinal position mismatch), thus, when assuming an

ordinal retrieval mode for Skips and Repeats we evaluated activations contrasting detected versus

undetected trials. We used an anatomical region of interest (ROI) analysis to evaluate mPFC activity

across the three probe trial types. In support of our hypothesis the mPFC was significantly more active

during identification of Ordinal Transfers relative to both Skips and Repeats for contrasts that assumed

ordinal retrieval modes (Fig. 2E; $F_{(2, 56)}$ = 4.038, p = 0.034). Post hoc tests (LSD) further supported the

conclusion that the mPFC was most active in the comparison of undetected versus detected Ordinal

Transfers (M = 0.708 ± 1.998) relative to detected versus undetected Skips (M = -0.649 ± 1.445, p =

0.051) and Repeats (M = -0.431 ± 1.728, p = 0.005). No significant difference was identified in mPFC

activation between detected versus undetected Skips and Repeats (mean difference = 0.061 ± 0.437, p

= 0.890). As a follow-up and to explore contributions of distinct regions within our anatomical mPFC

ROI we evaluated the same contrasts at the voxel-wise level. We observed activations throughout the

mPFC bilaterally (prelimbic cortex, anterior cingulate cortex, medial superior frontal gyrus) following the

comparison of undetected versus detected Ordinal Transfers (Fig. 2F; FWE-tfce p < 0.05), while no

mPFC clusters survived corrections for multiple comparisons when comparing detected versus

undetected Skips (data not shown) and Repeats (Fig. 2I; FWE-tfce p > 0.05).

One potential confound to this interpretation is that mPFC activations might reflect relative

performance levels (incorrect > correct trials) rather than the utilization of ordinal retrieval modes per

se. If true, we would expect similar mPFC activation clusters when examining incorrect compared to

correct Skips and Repeats if performance was the main contributor to activation. When evaluating

performance related activations for Repeats and Skips, undetected (i.e., incorrect) compared to

detected (i.e., correct) Skips exhibited activations predominantly in rostrolateral prefrontal cortex (Fig.

2G) while a cluster for detected greater than undetected Repeats was identified in the right fusiform

gyrus (Fig. 2H). These results further support the view that the mPFC contributes to ordinal retrieval

modes, and do not generally reflect performance accuracy. Notably, the observed activations in the

rostrolateral prefrontal cortex during undetected Skips, and the fusiform gyrus for detected Repeats

may reflect prospective memory (Umeda et al., 2011; Volle et al., 2011; Benoit et al., 2012) and object

processing (Grill-Spector et al., 1998), respectively.

Figure 2. *mPFC activations and ordinal retrievals.* **A,**

Ordinal retrieval modes were probed using Ordinal

Transfer trials. Ordinal Transfers occurred when an

item from one sequence (e.g., UVWXYZ) was

transferred to another sequence (e.g., ABCDEF)

while retaining in its original ordinal position (e.g.

ABCDYF). Theoretically, Ordinal Transfers are

identified as InSeq when an ordinal association is

used, and identified as OutSeq when a sequential

item-item association is used. **B,** Ordinal Transfers

occurred in every position in each sequence except

the first. For each position (2 through 6), mean

performance was significantly higher than chance

(red dashed line; one minus the calculated response

bias), and significantly lower than the mean

performance on Repeats (purple dashed line) and

Skips (blue dashed line). Performance did not

significantly differ among positions. **C,** the

normalized performance slope did not significantly

differ across positions, and variability, measured by

the averaged squared residuals per participant of

the linear regression, was high overall. High

performance variability suggests conflicting

strategies (Repeats and Skips variability levels

plotted for comparison in **D**). For the subsequent

BOLD analysis, we focused on contrasts assuming

participants were using an ordinal retrieval mode.

When an ordinal retrieval mode is engaged,

participants would identify Ordinal Transfers as

InSeq, but would identify Repeats and Skips as

OutSeq. **E**, a BOLD fMRI mPFC ROI analysis

showed that the Ordinal Transfers Undetected –

Detected contrast activation was significantly higher

compared with Repeats and Skips (Detected – Undetected Contrasts). **F,** a voxel-wise BOLD fMRI analysis using mPFC as a

mask (outlined in green) resulted in significant activity in a large area of the mPFC. To investigate whether there was a

difference between these probe trials, we compared Ordinal Transfers Undetected – Detected, Repeats Detected –

Undetected, and Skips Detected – Undetected trials. **G,** mPFC activation was also evident in whole brain analysis for Ordinal

Transfers Undetected – Detected. **H,** additionally, we found activation when comparing Skips Undetected – Detected in the

rostrolateral prefrontal cortex. No significant activation was found when comparing Repeats Undetected – Detected (not

shown). **I,** when comparing Repeats Detected – Undetected we found activation in the right fusiform gyrus. No significant

activation was found when comparing Skips Detected – Undetected (not shown). Notably, mPFC activation did not simply

reflect inaccuracy on OutSeq trials. Symbol(s): ns, not significant; *, $p < 0.05$, **, $p < 0.01$, ***, $p < 0.001$.

Sequence memory as a function of lag direction and distance

Memory for sequences of events at different lags can be supported by a variety of cognitive

processes including working memory (WM) and temporal context memory (TCM) retrieval modes. The

use of WM and TCM retrieval modes predict different patterns in behavioral performance across n-

forward and n-reverse lags (see Fig. 3B; also see Jayachandran et al., 2019), which can be exploited

here to test the use of different retrieval modes (e.g., WM versus TCM).

Skips (forward lags). OutSeq probe trials that skipped ahead in the sequence (n-forward, e.g.,

the "D" in ABD̲DEF) afford the opportunity to evaluate predictions of the use of a TCM retrieval mode

during rapid sequence memory decisions. Specifically, successful performance on Skips relies on a

TCM retrieval mode as it requires participants to have precise expectations for the subsequent items in

the sequence. Accordingly, the likelihood of a memory retrieval is highest for the very next item in the

forward direction (lag distance = +1) and drops off (in a graded fashion) for more distal items (from +2

to +4). In the context of this task, the use of a TCM retrieval mode predicts the inverse in performance

compared with free recall tasks because of interference. Specifically, performance on Skip OutSeq

probe trials should be most difficult to detect for the shortest forward lag distance (+1) and improve at

longer distances (+2, +3, or +4; Fig. 3B, right side) precisely because proximal items in a sequence

(i.e., Skips with a short forward lag) are more likely to be retrieved (Howard and Kahana 2002; Kragel

et al. 2015) and then falsely match up with an out of sequence probe image (thus be judged as InSeq;

undetected).

We tested this prediction by examining performance across n-forward lag distances. First, all

forward lags were performed better than chance (Fig. 3C, blue bars; +1: 0.420 ± 0.178; $+1_{vs.\ chance}$: $t_{(33)} =$

10.609, $p = 3.608 \times 10^{-12}$; +2: 0.665 ± 0.156; $+2_{vs.\ chance}$: $t_{(33)} = 21.299$, $p = 7.417 \times 10^{-21}$; +3: $0.747 \pm$

0.248; $+3_{vs.\ chance}$: $t_{(33)} = 15.343$, $p = 1.402 \times 10^{-16}$; +4: 0.765 ± 0.431; $+4_{vs.\ chance}$: $t_{(33)} = 9.066$, $p = 1.780 \times$

10^{-10}), suggesting that the use of a TCM retrieval mode does not completely interfere with OutSeq

detection at the different forward lag distances. Second, subjects exhibited graded performance

improvement as the skip distance increased ($F_{(3,\ 99)} = 13.790$, $p = 1.379 \times 10^{-4}$). We found that the

identification of Skips with a +1 lag were the most difficult to detect compared with all other forward lags

(*post hoc* LSD: +2: mean difference = 0.245 ± 0.029, p = 1.218 x 10^{-9}, +3: mean difference = 0.327 ±

0.049, p = 1.248 x 10^{-7}, +4: mean difference = 0.345 ± 0.079, p = 1.186 x 10^{-4}). These results are

consistent with TCM retrieval mode predictions (Fig. 3B, 3C). Detection of Skips with a +2 lag was

significantly lower than Skips with a +3 lag (*post hoc* LSD: mean difference = 0.082 ± 0.038, p = 0.039).

No significant difference in detection was observed, however, when comparing Skips with lags of +2

and +4 (*post hoc* LSD: mean difference = 0.100 ± 0.074, p = 0.187), or +3 and +4 (*post hoc* LSD:

mean difference = 0.018 ± 0.074, p = 0.814), suggesting that performance approached asymptote.

These behavioral results suggest that the participants' sequence memory was driven by the use of a

TCM retrieval mode on Skips.

11   *Repeats (reverse lags).* OutSeq trials that repeated an item from earlier in the sequence (*n*-

reverse; e.g., the second "A" in AB<u>A</u>DEF) can be solved by either the use of a TCM retrieval mode, a

WM retrieval mode, or a mixture of the two (Fig. 3B, left side). Notably, the pattern of OutSeq

detections across *n*-reverse lag distances should differentiate the two competing processes

(Jayachandran et al., 2019). For Repeats, a TCM retrieval mode predicts that short backward lag

distances (e.g., -2) would be the most difficult to detect due to heightened interference, while further *n*-

reverse lag distances (e.g. -5) would be readily detectable as OutSeq. By contrast, a WM retrieval

mode predicts the opposite pattern because the most recently experienced items would be the most

accessible and therefore easiest to detect.

20   We found that performance was better than chance for all *n*-reverse lag distances (Fig. 3C,

purple bars, -2: 0.859 ± 0.116; -2$_{vs. chance}$: $t_{(33)}$ = 38.333, p = 6.058 x 10^{-29}; -3: 0.805 ± 0.171; -3$_{vs. chance}$:

$t_{(33)}$ = 24.138, p = 1.522 x 10^{-22}; -4: 0.782 ± 0.233; -4$_{vs. chance}$: $t_{(33)}$ = 17.204, p = 4.821 x 10^{-18}; -5: 0.529 ±

0.507; -5$_{vs. chance}$: $t_{(33)}$ = 4.996, p = 1.865 x 10^{-5}). As the *n*-reverse lag distance increased, detection as

OutSeq decreased ($F_{(3, 99)}$ = 9.760, p = 0.001; Fig. 3C, purple bars), suggestive of the dominant use of a

WM retrieval mode. Items with an *n*-reverse lag closest to their original position (e.g., ABC<u>B</u>EF; lag = -

2) were easiest to detect as OutSeq compared with all other *n*-reverse lag positions (*post hoc* LSD lag -

2 compared with: lag -3: mean difference = 0.055 ± 0.022, p = 0.020; lag -4: mean difference = 0.077

± 0.033, p = 0.025; or lag -5: mean difference = 0.330 ± 0.084, p = 4.262 x 10^{-4}). By contrast, items

with an *n*-reverse lag farthest from their original position (e.g. ABCDE<u>A</u>; lag = -5) were the most difficult

to detect as OutSeq compared with all other *n*-reverse lag positions (*post hoc* LSD lag -5 compared

with: lag -4: mean difference = 0.253 ± 0.091, p = 0.009; lag -3: mean difference = 0.275 ± 0.091, p =

0.005; or lag -2: mean difference = 0.330 ± 0.084, p = 4.262 x 10^{-4}). No significant differences in

detection between -3 and -4 *n*-reverse lags were observed (*post hoc* LSD: mean difference = 0.022 ±

0.035, p = 0.532). Taken together, performance at the *n*-reverse lag extremes (-2 and -5) supports the

notion that the use of a WM retrieval mode plays an important role in identifying Repeats. The absence

of a graded performance across *n*-reverse lags of -3 and -4 supports the idea that a combination of

cognitive processes is being used, but overall these analyses do not support the use of a TCM retrieval

mode as an isolated process driving the identification of Repeats in this task.

TCM retrievals for Skips, but a mixture of retrieval modes for Repeats.

Differences in the overall ability to detect OutSeq probe trials, and the residuals from a lag-

based linear regression model, helped to further elucidate the contributions of either a TCM retrieval

mode, a WM retrieval mode, or their combination. First, we predicted that the ability to detect OutSeq

probes would be greater for Repeats than Skips because TCM and WM retrieval modes can both

contribute to the evaluation of Repeats but not Skips (Fig. 3B, left, purple dashed line). Second, we

predicted that a linear regression of Skip detections would positively increase across *n*-forward lags,

whereas a similar analysis across *n*-reverse lags for Repeats would be essentially flat. Third, we

predicted that the residuals from the linear regressions would be, highest on Repeats compared with

Skips, reflecting the use of multiple retrieval modes, whereas Skips involve a single mode (i.e., TCM-

based retrieval). To test these predictions, we compared the overall ability to detect the two probe trial

types, calculated a linear regression based on the z-scores (accounting for individual baseline detection

levels) across different lag positions, and averaged the squared residuals as a measure of detection

Figure 3. *BOLD fMRI analysis indicates that*
the right anterior hippocampus (raHC)
temporal context across forward lags. **A,** Skips
and Repeats could occur at different lags away
from their InSeq position. Skip lags ranged from
+1 (e.g. ABCDEFF) to +4 (e.g. AFCDEF). Repeat
lags ranged from -2 (e.g. ABCBEF) to -5 (e.g.
ABCDEA). **B,** theoretical predictions based on
temporal context memory (TCM) and working
memory (WM) retrieval modes for different lag
distances on Repeats (e.g., ABA) and Skips
(e.g., ABD). The TCM (blue lines) predicts that
item retrieval in sequences is graded by lag
distance (for both Repeats and Skips).
According to the TCM, OutSeq probe trials with
small lags would be more difficult to detect than
those with larger lags reflecting the fact that
they were more temporally contiguous in the
original sequence and likely to be judged InSeq.
Conversely, the WM (red line) predicts that
items with small lags would be easier to detect
than those with larger lags, but WM can only
contribute to performance on Repeats. A
mixture of TCM and WM modes is shown by
the dashed purple line. **C,** the mean
performance accuracy for all participants on
Repeats and Skips at different lags. **D,** we
predicted and found that overall performance
on Repeats would be higher than for Skips
because TCM and WM retrieval modes
complement each other. **E,** we also predicted
and found that performance would show a
positive slope on Skips defining of TCM
retrieval modes, **F,** and would be flat on

Repeats reflecting the use of a mixture of TCM and WM retrieval modes. Specifically, a linear regression in Skips performance
positively increased across forward lags, but did not have a significant slope on Repeats. **G,** finally, we predicted and found
significantly higher variability, measured by the averaged squared residuals per participant of the linear regression, in Repeats
compared with Skips, which is thought to reflect the integration of variance from using both TCM and WM retrieval modes.
Symbol(s): ns, not significant; *, $p < 0.05$, **, $p < 0.01$, *** $p < 0.001$. **H,** BOLD fMRI analysis of Skips with a linear contrast of -
1, 0, 1 for lags of +1, +2, +3 respectively using the bilateral medial temporal lobe as a revealed significant activation in the
raHC (right anterior Hippocampus). **I,** BOLD fMRI analysis of Repeats with a linear contrast of 1, 0, -1 for lags of -4, -3, -2
respectively using bilateral medial temporal lobe as a mask showed no significantly activated clusters.

variability across lags. Participants were significantly better at detecting Repeats ($M_{accuracy}$ = 0.786 ±

0.166) than Skips ($M_{accuracy}$ = 0.492 ± 0.157; $t_{(33)}$ = 14.435, p = 8.137 x 10^{-16}; Fig. 3D).

Consistent with our hypothesis, the linear regression in Skips positively increased across

forward lags accounting for a large effect on performance (Fig. 3E; R^2 = 0.572, β = 0.757, p = 3.711 x

10^{-20}), whereas a linear regression across reverse lags for Repeats had no significant slope (Fig. 3F; R^2

= 0.024, β = 0.154, p = 0.139). When quantifying variability, Repeats ($M_{residual}$ = 0.593 ± 0.243) were

significantly more variable compared with Skips ($M_{residual}$ = 0.285 ± 0.478) across different lag positions

(Fig. 3G; $t_{(30)}$ = 3.051, p = 0.004). The patterns in OutSeq detection and variability further support the

conclusion that multiple retrieval modes likely contribute to identifying Repeats as OutSeq, while Skip

detection is mediated more exclusively by a TCM retrieval mode.

HC activations and TCM retrievals

Converging evidence indicates that the medial temporal lobe, specifically the HC formation,

plays a disproportionate role in the use of a TCM retrieval mode in the brain (Manns et al., 2007; Hsieh

et al., 2014; Bladon et al., 2019). To test whether regions of the medial temporal lobe contribute to a

TCM retrieval mode during Skip and Repeat probe trials, we looked for linear changes in activation

across the different lag distances. We observed a significant activation cluster in the right anterior HC

following small volume corrections (bilateral medial temporal lobe, FWE-tfce p < 0.05) that increased its

activation across n-forward lags (Fig. 3H). The statistical significance of similar patterns of activation

across n-reverse lags did not survive corrections for multiple comparisons (Fig. 3I). These results

suggest that the right anterior HC contributes to a TCM retrieval mode during Skips.

Brain region by retrieval mode interaction effect

After finding individual evidence supporting the mPFC contributes to sequence memory by the

use of an ordinal retrieval mode and the right anterior HC by a TCM retrieval mode, we wanted to

directly compare the mode-related activations (ordinal vs. TCM) across the regions that exhibited

voxelwise activations (right anterior HC vs. mPFC). A repeated-measures factorial ANOVA was

conducted with brain region (anatomically defined right anterior HC vs. mPFC) and strategy (ordinal

retrieval vs. TCM retrieval modes) as within-subjects factors, and BOLD activations as the dependent

measure. We observed greater activations in the mPFC in relation to the use of an ordinal retrieval

relative to a TCM retrieval mode, and the opposite pattern in the right anterior HC, evidenced by a

significant interaction effect (Fig. 4A. $F_{(1,31)}$ = 4.782, p = 0.036).

Figure 4. *mPFC and HC retrieval mode interactions.* **A,** there was a significant interaction effect between mode and brain

region ($F_{(1,31)}$ = 4.782, p = 0.036), where we observed greater activations in the mPFC for ordinal retrievals relative to TCM

retrievals, and the opposite pattern in the right anterior HC. **B,** based on the results of the current experiment, **1)** the mPFC

engages in ordinal-based retrieval calling on associations between items and their ordinal position (which could be

represented elsewhere) and **2)** the HC engages in temporal context-based retrieval.

Discussion

The current study used different out of sequence probe trials during a memory task to test the

concurrent use of distinct retrieval modes, adding to a growing literature on sequence memory as a

fundamental component of episodic memory (Tulving, 1984; 2002; Allen and Fortin, 2013; Howard and

Eichenbaum, 2013; Eichenbaum, 2017). Complementary behavioral results across the different out of

sequence probe trials in the current task support the conclusion that memory for sequences of events is

supported by both ordinal and temporal context retrieval modes. The BOLD fMRI evidence showed that

mPFC activations reflect the use of an ordinal retrieval mode, while activations in the HC reflected a

temporal context memory retrieval mode, thus dissociating the neurobiological substrates of two distinct

processes contributing to sequence memory. While considerable evidence indicates that the mPFC and

HC are involved in sequence memory, the current study provides new results that the mPFC and HC

are concurrently engaged in different retrieval modes in support of remembering *when* an event

occurred. This evidence provides an important baseline for further investigation of how sequence

memory is impaired in typical aging and diseases such as Alzheimer's Disease on the neurobiological

level, especially since evidence suggests the relative dependence on an ordinal retrieval mode

increases while the use of a temporal context retrieval mode declines with age (Bastin and Van der

Linden, 2010; Allen et al., 2015).

Ordinal retrieval modes in mPFC

Our results suggest that the positional information associated with retrieving memories in

sequence is represented in the mPFC (e.g., Hsieh and Ranganath, 2015), and/or that mPFC

activations help engage these representations elsewhere such as in HC neurons (e.g., Allen et al.,

2016). Ordinal Transfers were detected less often than both Repeats and Skips (Fig 2B, see also Allen

et al., 2014, 2015), but better than chance, suggestive of the use of multiple retrieval mode perhaps

concurrent ordinal and temporal context retrieval modes. These results are not totally surprising

because mPFC has been shown to be generally important to temporal order memory (Milner et al.,

1985; Shimamura et al., 1990; DeVito and Eichenbaum, 2011; Hsieh et al., 2015) and other semantic

representations (Preston and Eichenbaum, 2013; Hyman et al., 2012). Our study shows that when an

ordinal retrieval mode is strongly engaged, it interferes with the ability to detect Ordinal Transfers,

supported by the fact that Transfers went undetected at a very high rate, and mPFC activation was

similarly high in such a way that could not be attributed to undetected probes more globally.

Ordinal retrieval modes have been demonstrated in rats (Allen et al., 2014), monkeys (Orlov et

al., 2000; 2006) and humans (Allen et al., 2015; Hsieh et al., 2015). In fact, monkeys natural and

dominant strategy is to categorize items within a sequence by their ordinal position, a strategy that

occurred first in Orlov et al. (2000), and only later in trials did monkeys employ other strategies such as

sequential associations for adjacent items and working memory (Orlov et al., 2000; 2006). Together,

this evidence strongly suggests that an ordinal retrieval mode dominates early in sequence memory,

and is subsequently bolstered by other retrieval modes such as a TCM mode. Likewise, in humans it

has been shown that multi-voxel patterns from the mPFC are significantly higher for objects that share

the same position information, compared to objects in different positions (Hsieh et al., 2015),

suggesting convergent patterns of activation reflect shared ordinal representations within mPFC (Tiganj

et al., 2017) despite the difference in object identity. Taken together, prior research and the current

study show compelling evidence that the mPFC helps remember *when* events occurred by engaging an

ordinal retrieval mode.

Theoretically, an ordinal retrieval mode generated by mPFC would be input to the HC during

episodic memory through indirect cortical or thalamic pathways (for review see Dolleman-van der Weel

et al., 2019). The engagement of an ordinal retrieval mode in HC might then allow for the rapid

formation of conjunctive item-position representations (e.g., 1^{st} – A, 2^{nd} – B, etc.; Fig. 4B), and provide

sequential structure without an explicit need to represent the elapsing time between items (a useful

form of neural compression for temporal information). There is indirect evidence that item-position

representations are reflected in rodent CA1 neurons during spatial sequence tasks (Euston and

McNaughton, 2006), and direct evidence for conjunctive item-position representations during an

analogous odor sequence task (Allen et al., 2016). Importantly, item-position representations are

learned and retrieved *before* sequential item-item associations in non-human primates (Orlov et al.,

2000). This begs the question whether episodic memories typically rely on an ordinal retrieval mode for

recalling events with timelines. Subsequent replay events or other consolidation processes could either

strengthen temporal context modes and/or weaken ordinal modes, although we suspect the former.

Interestingly, in monkeys ordinal transfers show graded interference over lag distances, suggesting

they share similar properties with temporal context retrievals when studied in this way (Orlov et al.,

2006), indicating two temporal dimensions may be complementary and normally integrated. Future

neuroimaging studies in humans using ordinal transfers distributed across lag distances will be useful

for examine the neural activity when these processes are both contributing to retrieval patterns.

Temporal context retrieval modes in HC

Consistent with the literature, our results show that the HC contributes to sequence memory

through the use of a temporal context retrieval mode. The HC has been shown to be important for

sequence memory (Hsieh et al., 2014; Goyal et al., 2018) but its precise contribution has remained an

open question. According to TCM, the HC associates items in sequences through a drifting contextual

representation (Howard et al., 2005; Polyn and Kahana, 2008). As such, the presentation or retrieval of

an item from a sequence elicits the retrieval of neighboring items that share a temporal context,

decreasing in likelihood or strength for more distal items (Howard and Kahana, 2002). According to this

framework, a temporal context retrieval mode accounts for the increased likelihood that adjacent items

in word lists are recalled (Howard and Kahana, 2002). This pattern in free-recall performance predicted

by temporal context memory has been validated in computational models (Howard and Kahana, 2002),

behavioral studies (Kahana, 1996; Sederberg, 2010; Morton and Polyn, 2016), and neurobiological

studies (Polyn and Kahana, 2008; Jenkins and Ranganath, 2010; Hsieh et al., 2014; Bladon et al.,

2019). In the current sequence memory task, we reasoned that the presentation of InSeq items would

elicit the retrieval of neighboring items that shared temporal contexts and lead to graded impairments in

the detection of Skip probes. This is because, as the forward lag increased, the upcoming

representations were less likely to be retrieved and thus less likely to interfere with an out of sequence

determination. Similar patterns in performance are observed in other tasks probing temporal memory

(Allen et al., 2014; 2015; DuBrow and Davachi, 2014).

The HC has shown activations consistent with the use of a TCM retrieval mode. Activation in the

HC is elevated during the processing of overlapping compared to nonoverlapping sequences (Kumaran

and Maguire, 2006; Brown et al., 2010; Brown and Stern, 2013). Population activity in CA1 drifts across

both small- and large-time scales (Manns et al., 2007; Mankin et al., 2012; Ziv et al., 2013; Rubin et al.,

2015; Mau et al., 2018). Additionally, HC lesions impair the discrimination of overlapping odor

sequences (Agster et al., 2002). A recent MRI study showed that HC multi-voxel pattern similarity was

higher for pairs of adjacent trials that belonged to the same temporal context within a sequence

compared to pairs of sequence items that bridged between sequences, even when the temporal

distance between the pairs of items was similar (Hsieh et al., 2014). The same study observed that the

HC carries information about the temporal context between items within a sequence, rather than

information about the objects themselves (Hsieh et al., 2014). Specifically, Hsieh et al. (2014)

demonstrated that when the same sequence item is repeated, hippocampal voxel patterns were

dissimilar, unless the temporal context was reinstated. Our results add to this by showing that as Skips

lag further away from their InSeq location, HC activity also increases closely matching predictions of

TCM.

Limitations and theoretical considerations

While the use of specialized out of sequence probe trials provided important insight regarding

different retrieval modes, and their related neurobiological substrates, several limitations of the current

study remain. First, the timing of the task and self-paced design of the experiment preclude detailed

item-based analyses at the neurobiological level. At the expense of repetitions of trials, future studies

should include temporal jitter between sequence items to isolate signals at the different item positions.

Second, similar to multivariate approaches, univariate approaches are subject to interpretational

ambiguities (Hebart and Baker, 2018; Ritchie et al., 2017). However, to avoid overfitting issues,

especially with time-varying representations, we chose to use a univariate approach over a multivariate

approach as it is better suited to our task design to differentiate retrieval modes using different out-of-

sequence probe trials.

Key questions remain concerning the interactions between ordinal and temporal context

retrieval modes, and their related neurobiological constituents (mPFC and HC). While the current study

did not include probe trials that investigated ordinal transfer and transpositions, future studies should

examine these critical probe trial types and expect to see increased interactions between the mPFC

and HC. mPFC-HC coupling may lead to conjunctive representations of temporal contexts and

positional coding. As shown here, mPFC activation reflects ordinal retrieval modes (Fig. 4B) which

interacts with temporal contexts in the HC (Fig. 4B). In theory, this information could be merged

allowing the formation of conjunctive temporal context and item-position association in HC neurons.

Direct evidence for this latter possibility was provided by recent studies that showed HC neurons

encode item-context and item-position conjunctions (Komorowski et al., 2009; Allen et al., 2016),

suggesting that these conjunctive representations provide the neuronal basis for time and place

integrations in episodic memory. However, future studies are required for examining mPFC-HC

interdependent interactions directly in humans.

Conclusion

The results from our study highlight novel evidence that ordinal and temporal context retrieval

modes both contribute to remembering items with a timeline. In particular we showed that mPFC

activity reflected an ordinal retrieval mode and HC activity reflected a TCM retrieval mode. Further

experiments in both animals and humans are necessary to delineate the precise mechanisms by which

the mPFC engages ordinal retrievals, and how this process interacts with HC to integrate ordinality and

temporal contexts in support everyday episodic memory.

References

Agster KL, Fortin NJ, Eichenbaum H. 2002. The hippocampus and disambiguation of overlapping sequences. J Neurosci 22(13):5760-5768.

Allen TA, Fortin NJ. 2013. The evolution of episodic memory. Proc Natl Acad Sci U S A 110 (suppl 2):10379-10386.

Allen TA, Morris AM, Mattfeld AT, Stark CEL, Fortin NJ. 2014. A sequence of events model of episodic memory shows parallels in rats and humans. Hippocampus 24:1178-1188.

Allen TA, Morris AM, Fortin NJ, Stark CEL. 2015. Memory for sequences of events impaired in typical aging. Learn Mem 22:138-148.

Allen TA, Salz DM, McKenzie SA, Fortin NJ. 2016. Nonspatial sequence coding in CA1 neurons. J Neurosci 36(5):1547-1563.

Avants BB, Epstein CL, Grossman M, Gee JC. 2008. Symmetric diffeomorphic image registration with cross-correlation: Evaluating automated labeling of elderly and neurodegenerative brain. Med Image Anal 12(1):26-41.

Bastin C, Van der Linden M. 2005. Memory for temporal context: Effects of ageing, encoding instructions, and retrieval strategies. Memory, 13:1.

Benoit RG, Gilbert SJ, Frith CD, Burgess PW. 2012. Rostral prefrontal cortex and the focus of attention in prospective memory. Cerebral Cortex 22(8):1876-1886.

Bladon JH, Sheehan DJ, De Freitas CS, Howard MW. 2019. In a temporally segmented experience hippocampal neurons represent temporally drifting context, but not discrete segments. J Neurosci 39(35):6936-6952.

Brown TI, Stern CE. 2013. Contributions of medial temporal lobe and striatal memory systems to learning and retrieving overlapping spatial memories. Cereb Cortex 24(7):1906-22.

Brown TI, Ross RS, Keller JB, Hasselmo ME, Stern CE. 2010. Which way was I going? Contextual

retrieval supports the disambiguation of well learned overlapping navigational routes. J Neurosci

30:7414 –7422.

Cox RW, 1996. AFNI: software for analysis and visualization of functional magnetic resonance

Neuroimages. Comput Biomed Res 29(3):162-173.

Dale AM, Fischl B, Sereno MI. 1999. Cortical surface-based analysis. I. Segmentation and surface

reconstruction. Neuroimage 9:179-194.

DeVito LM, Eichenbaum H. 2011. Memory for the order of events in specific sequences: Contributions

of the hippocampus and medial prefrontal cortex. J Neurosci 31(9):3169-3175.

Dolleman-van der Weel, Griffin AL, Ito HT, Shapiro ML, Witter MP, Vertes RP, Allen TA. 2019. The

nucleus reuniens of the thalamus sits at the nexus of a hippocampus and medial prefrontal cortex

circuit enabling memory and behavior. Learn Mem 26:191-205.

Dubrow S, Davachi L. 2014. Temporal memory is shaped by encoding stability and intervening item

reactivation. J Neurosci 34(42):13998-14005.

Dubrow S, Davachi L. 2013. The influence of context boundaries on memory for the sequential order of

events. J Exp Psychol Gen 142(4):1277-1286.

Eichenbaum H. 2014. Time cells in the hippocampus: a new dimension for mapping memories. Nat

Neurosci 15:732-744.

Eichenbaum H. 2017. On the integration of space, time and memory. Neuron. 95(5):1007-1018.

Eichenbaum H, Yonelinas A, Ranganath C. 2007. The medial temporal lobe and recognition memory.

Annu Rev Neurosci 30:123-152.

Ekstrom AD, Bookheimer SY, 2007. Spatial and temporal episodic memory retrieval recruit dissociable

functional networks in the human brain. Learn Mem 14(10):645-654.

Euston DR, McNaughton BL. 2006. Apparent encoding of sequential context in rat medial prefrontal

cortex is accounted for by behavioral variability. J Neurosci 26(51):13143-13155.

Fortin NJ, Agster KL, Eichenbaum HB. 2002. Critical role of the hippocampus in memory for sequences

of events. Nat Neurosci 5:458-462.

Gorgolewski K, Burns CD, Madison C, Clark D, Halchenko YO, Waskom ML, Ghosh SS. 2011. Nipype:

a flexible, lightweight extensible neuroimaging data processing framework in python. Front

Neuroinform 5:13.

Goyal A, Miller J, Watrous AJ, Lee SA, Coffey T, Sperling MR, Sharan A, Worrell G, Berry B, Lega B,

Jobst BC, Davis KA, Inmann C, Sheth SA, Wanda PA, Azzyat Y, Das SR, Stein J, Gorniak R,

Jacobs J. 2018. Electrical stimulation in hippocampus and entorhinal cortex impairs spatial and

temporal memory. J Neurosci 38(19):4471-4481.

Grill-Spector K, Kushnir T, Hendler T, Edelman S, Itzchak Y, Malach R. 1998. A sequence of object-

processing stages revealed by fMRI in the human occipital lobe. Hum. Brain Mapp. 6:316-328.

Hebart MN, Baker CI. 2018. Deconstructing multivariate decoding for the study of brain function.

Neuroimage. 180:4-18.

Howard MW, Eichenbaum H. 2013. The hippocampus, time and memory across scaled. J Exp Psychol

142(4):1211-1230.

Howard MW, Kahana MJ. 2002. A distributed representation of temporal context. J Math Psychol 46:

269-299.

Howard MW, Fotedar MS, Datey AV, Hasselmo ME. 2005. The temporal context model in spatial

navigation and relational learning: toward a common explanation of medial temporal lobe function

across domains. Psych Rev 112(1):75-116.

Howard MW, Viskontas IV, Shankar KH, Fried I. 2012. Ensembles of human MTL neurons "jump back

in time" in response to a repeated stimulus. Hippocampus 22:1833-1847.

Hsieh LT, Gruber MJ, Jenkins LJ, Ranganath C. 2014. Hippocampal activity patterns carry information

about objects in temporal context. Neuron 81:1165-1178.

Hsieh LT, Ranganath C. 2015. Cortical and subcortical contributions to sequence retrieval: Schematic

coding of temporal context in the neocortical recollection network. Neuroimage 121:78-90.

Hyman JM, Ma L, Balaquer-Ballester e, Durstewitz D, Seamans JK. 2012. Contextual encoding by

ensembles of medial prefrontal cortex neurons. Proc Natl Acad Sci U S A. 109(13): 5086-5091.

Jayachandran M, Linley SB, Schlecht M, Mahler SV, Vertes RP, Allen TA. 2019. Prefrontal pathways

provide top-down control of memory for sequences of events. Cell Rep 28(3):640-645.

Jenkins LJ, Ranganath C. 2010. Prefrontal and medial temporal lobe activity at encoding predicts

temporal context memory. J Neurosci 30(46):15558-15565.

Jenkins LJ, Ranganath C. 2016. Distinct neural mechanisms for remembering when an event occurred.

Hippocampus 26(5):554-559.

Kahana MJ. 1996. Associative retrieval processes in free recall. Mem and Cogn. 24(1): 103-109.

Kalm K, Davis MH, Norris D. 2013. Individual sequence representations in the medial temporal lobe. J

Cogn Neurosci 25:1111-1121.

Kesner RP, Gilbert PE, Barua LA. 2002. The role of the hippocampus in memory for the temporal order

of a sequence of odors. Behav Neurosci 116(2):286-290.

Komorowski RW, Manns JR, Eichenbaum H. 2009. Robust conjunctive item-place coding by

hippocampal neuros parallels learning what happens where. J Neurosci 29(31):9918-9929.

Kragel JE, Morton NW, Polyn SM. 2015. Neural activity in the medial temporal lobe reveals the fidelity

of mental time travel. J Neurosci 35(7):2914-2926.

Kumaran D, Maguire EA. 2006. An unexpected sequence of events: Mismatch detection in the human

hippocampus. PLoS Biol 4(12):2372-2382.

Lehn H, Steffenach HA, Strien NM, Veltman DJ, Witter MP, Haberg AK. 2009. A specific role of the

human hippocampus in recall of temporal sequences. J Neurosci 29(11):3475-3484.

Long HMK, Kahana NM. 2019. Hippocampal contributions to serial-order memory. Hippocampus doi:

10.1002/hipo.23025.

Mankin EA, Sparks FT, Slayyeh B, Sutherland RJ, Leutgeb S, Leutgeb JK. 2012. Neuronal code for

extended time in the hippocampus. Proc Natl Acad Sci U S A 109(47):19462-19467.

Manns JR, Howard MW, Eichenbaum H. 2007. Gradual changes in hippocampal activity support

remembering the order of events. Neuron 56(3):530-540.

Mau W, Sullivan DW, Kinsky NR, Hasselmo ME, Howard MW, Eichenbaum H. 2018. The same

hippocampal CA1 population simultaneously codes temporal information over multiple timescales.

Curr Biol 28(10):1499-1508.

Milner B, Petrides M, Smith ML. 1985. Frontal lobes and temporal organization of memory. Hum

Neurobiol 4:137-142.

Morton NW, Polyn SM. 2016. A predictive framework for evaluating models of semantic organization in

free recall. Journal of memory and language. 86: 119-140.

Orlov T, Amit DJ, Yakovlev V, Zohary E, Hochstein S. 2006. Memory of ordinal number categories in

macaque monkeys. J Cogn Neurosci 18(3):399-417.

Orlov T, Yakovlev V, Amit D, Hochstein S, Zohary E. 2002. Serial memory strategies in macaque

monkeys: Behavioral and theoretical aspects. Cereb Cortex 12(3):306-317.

Orlov T, Yakovlev V, Hochstein S, Zohary E. 2000. Macaque monkeys categorize images by their

ordinal number. Nature 404:77-80.

Preston AR, Eichenbaum H. 2013. Interplay of hippocampus and prefrontal cortex in memory. Curr Biol

23(17):764-773.

Polyn SM, Kahana MJ. 2008. Memory search and the neural representation of context. Trends Cogn

Sci 12(1):24-30.

Ritchie JB, Kaplan DM, Klein C. 2017. Decoding the brain: Neural representation and the limits of

multivariate pattern analysis in cognitive neuroscience. Brit J Philos Sci.70(2):581-607.

Roche A. 2011. A four-dimensional registration algorithm with application to joint correction of motion

and slice timing in fMRI. IEE Trans Med Imaging 30(8):1546-1554.

Ross RS, Brown TI, Stern CE. 2009. The retrieval of learned sequences engages the hippocampus:

Evidence from fMRI. Hippocampus 19:790-799.

Rubin A, Geva N, Sheintuch L, Ziv Yaniv. 2015. Hippocampal ensemble dynamics timestamp events in

long-term memory. eLife 4:e12247.

Shimamura AP, Janowsky JS, Squire LR. 1990. Memory for the temporal order of events in patients

with frontal lobe lesions and amnesic patients. Neuropsychologia 28:803-814.

Sederberg PB, Miller JF, Howard MW, Kahana MJ. 2010. The temporal contiguity effect predicts

episodic memory performance. Mem Cognit. 38(6):689-699.

Sokal RR, Rohlf FJ. 1995. Biometry: The Principles and Practices of Statistics in Biological Research.

New York (NY): W.H. Freeman and Company.

Tiganj Z, Kim J, Jung MW, Howard MW. 2017. Sequential firing codes for time in rodent medial

prefrontal cortex. Cereb Cortex 27(12):5663-5671.

Tubridy S, Davachi L. 2011. Medial temporal lobe contributions to episodic sequence encoding.

Cerebral Cortex 21:272-280.

Tulving E. 1984. Precis of elements of episodic memory. Behav Brain Sci 7:223-268.

Tulving E. 2002. Episodic memory: from mind to brain. Annu Rev Psychol 53:1-25.

Umeda S, Kurosaki Y, Terasawa Y. Kato M, Miyahara Y. 2011. Deficits in prospective memory

following damage to the prefrontal cortex. Neuropsychologia 49(8):2178-2184.

Volle E, Gonen-Yaacovi G, Costello AL, Burgess PW. 2011. The role of rostral prefrontal cortex in

prospective memory: a voxel-based lesion study. Neuropsychologia 49(8):2185-2198.

Ziv Y, Burns LD, Cocker ED, Hamel EO, Ghosh KK, Kitch LJ, El Gamal A, Schnitzer MJ. 2013. Long-

term dynamics of CA1 hippocampal place codes. Nat Neurosci 16(3):264-266.