M1 disruption delays motor processes but not deliberation about action choices

Gerard Derosiere[1], David Thura[2], Paul Cisek[3], Julie Duque[1]

[1] Institute of Neuroscience, Laboratory of Neurophysiology, Université catholique de Louvain, Brussels, Belgium

[2] Lyon Neuroscience Research Center - Impact team, Inserm U1028 - CNRS UMR 5292, Bron, France

[3] Department of Neuroscience, Université de Montréal, Montréal, QC H3T 1J4, Canada

Corresponding author contact details:
Gerard Derosiere
CoActions Lab
Institute of Neuroscience
Université catholique Louvain
Av. Mounier, 53 - Bte B1.53.04
1200 Bruxelles, Belgium
Tel: + 32 (0)2 764 54 20
Email address: gerard.derosiere@uclouvain.be

Running title: "M1 disruption and motor decisions"

Abstract

Decisions about actions typically involve a period of deliberation that ends with the commitment to a choice and the motor processes overtly expressing that choice. Previous studies have shown that neural activity in sensorimotor areas, including the primary motor cortex (M1), correlates with deliberation features during action selection. Yet, the causal contribution of these areas to the decision process remains unclear. Here, we investigated whether M1 determines choice commitment, or whether it simply reflects decision signals coming from upstream structures and instead mainly contributes to the motor processes that follow commitment. To do so, we tested the impact of a disruption of M1 activity, induced by continuous theta burst stimulation (cTBS), on the behavior of human subjects in *(1)* a simple reaction time (SRT) task allowing us to estimate the duration of the motor processes and *(2)* a modified version of the tokens task (Cisek et al., 2009), which allowed us to estimate subjects' time of commitment as well as accuracy criterion. The efficiency of cTBS was attested by a reduction in motor evoked potential amplitudes following M1 disruption, as compared to those following a sham stimulation. Furthermore, M1 cTBS lengthened SRTs, indicating that motor processes were perturbed by the intervention. Critically, all of the behavioral results in the tokens task were similar following M1 disruption and sham stimulation, suggesting that the contribution of M1 to the deliberation process is negligible. Taken together, these findings indicate that M1 contribution is downstream of the decision process.

News and noteworthy

Decisions between actions are ubiquitous in the animal realm. Previous work indicates that

deliberation during action choices entails changes in the activity of the sensorimotor areas

controlling those actions, but the causal role of these areas has yet to be established. Using

continuous theta burst stimulation, we show that disrupting the primary motor cortex delays the

motor processes that follow instructed commitment but does not alter volitional deliberation,

suggesting that the contribution of M1 is downstream of the decision process.

INTRODUCTION

The physical world provides animals with a variety of action opportunities, constantly requiring them to make decisions, some of which are critical for survival. For instance, the choice of a car driver to turn left or right in front of a sudden obstacle may have dramatic consequences on her/his life and on that of the pedestrians around. The driver will have to quickly deliberate and commit to one action.

Classical models of decision-making postulate that commitment to an action occurs when a neural signal related to that action reaches a critical decision threshold (Laming, 1968; Ratcliff, 1978; Stone, 1960). During deliberation, the signal grows from a starting point, at a rate that depends on the amount of sensory evidence favoring the chosen action. Evidence may be strong or weak, bringing the signal either quickly or slowly to the threshold (Alamia et al., 2018; Derosiere et al., 2018; Donner et al., 2009; Gould et al., 2012; Tosoni et al., 2014; Wyart et al., 2012). In the car driver example above, the presence of pedestrians on the right side of the street will strengthen the neural signal coding for the movement of rotating the wheel towards the left, and possibly weaken the signal favoring the opposite rightward rotation movement. Ultimately, the driver will commit to turning left and execute the related action, to avoid hitting the group of people.

Importantly, making decisions often requires balancing the desire to take time to deliberate (*i.e.*, to accumulate sensory evidence and make the best choice) with the urge to act (Churchland et al., 2008; Forstmann et al., 2008; Hanks et al., 2011; Thura and Cisek, 2014a). During a speeded decision, the urge to act increases as time passes (Cisek et al., 2009, Ditterich, 2006, Drugowitsch et al., 2012, Seideman et al., 2018; Thura et al., 2012) but the overall level of urgency also varies depending on the context (Murphy et al., 2016, Thura et al., 2014; Thura and Cisek, 2016, 2017). In the situation described earlier, the driver's level of urgency will be higher if the obstacle suddenly appears close to the car than if it appears far away. As evident

in this example, adjustments in urgency alter the balance between decision speed and accuracy

– *i.e.*, the so-called speed-accuracy tradeoff: choices are more likely to be incorrect when time

pressure is elevated, while accuracy improves when temporal demands permit long deliberation

(*e.g.*, Hanks et al., 2011; Seideman et al., 2018).

At the neural level, several lines of evidence indicate that higher levels of urgency during

deliberation about action choices amplify activity in the sensorimotor regions controlling these

actions. For reach movements, this concerns the dorsal premotor (PMd) and primary motor

cortex (M1; Steinemann et al., 2018, Thura and Cisek, 2014a, 2016). In these areas, neural

activity is globally increased at baseline and then builds-up at a faster rate when urgency is high

compared to when it is low, reducing the time needed to reach decision threshold but at the cost

of accuracy (Thura and Cisek, 2016). Recent findings suggest that the basal ganglia (Thura and

Cisek, 2017; van Maanen et al., 2016) and the locus coereleus (Hauser et al., 2018; Murphy et

al., 2016) may contribute to generate such a modulation of motor cortical activity. To date,

however, the specific contribution of motor cortical areas to the decision process remains

unclear.

Here, we investigated whether M1 itself contributes to deliberation about action choices,

or whether it simply reflects decision signals coming from upstream areas, such as PMd, the

basal ganglia, or the locus coereleus. It has previously been shown that disrupting M1 activity

by means of continuous theta burst stimulation (cTBS) causes finger responses to slow down

(Huang et al., 2005; Lakhani et al., 2014; McAllister et al., 2013). However, it is unclear

whether this effect should be interpreted as a slowing down of processes involved in deciding

which action to perform (*i.e.*, in the deliberation process) or as a slowing down of the motor

processes that follow commitment (*i.e.*, of movement initiation and execution). If M1 is

causally involved in the decision process, then M1 disruption should lengthen deliberation in a

manner consistent with reduced urgency, and therefore should increase the accuracy of

subjects' choices as compared to a sham cTBS session. Conversely, if M1 is mainly involved

in initiating and executing movements, then its disruption should have no effect on the

deliberation portion of response time or the accuracy of choices, and should only slow down

the motor processes that follow commitment to an action. These hypotheses were tested by

characterizing action choices in a modified version of the tokens task (Cisek et al., 2009), which

allows one to estimate subjects' time of commitment, their accuracy criterion and infer from

those variables their urgency functions (Thura et al., 2014).

MATERIALS AND METHODS

Participants

19 healthy right-handed subjects participated in this study (10 women; 24 ± 3.5 years old).

Participants were financially compensated for their participation and earned additional money

depending on their performance in a decision-making task (see *Task* section below). The

protocol was approved by the institutional review board of the catholic University of Louvain

(UCLouvain), Brussels, Belgium, and required written informed consent, in compliance with

the principles of the Declaration of Helsinki.

Experimental design

Experiments were conducted in a quiet and dimly-lit room. Subjects were seated at a table

in front of a 21-inch cathode ray tube computer screen. The display was gamma-corrected and

its refresh rate was set at 100 Hz. The computer screen was positioned at a distance of 70 cm

from the subject's eyes and was used to display stimuli during the decision-making task. Left

and right forearms were rested upon the surface of the table with the palms facing the table. A

computer keyboard was positioned upside-down under the dominant (*i.e.*, right) hand with the

response keys F9 and F8 under the index and middle fingers, respectively (see Figure 1).

Task

The task used in the current study is a variant of the "tokens task" (Cisek et al., 2009) and

was implemented by means of LabView 8.2 (National Instruments, Austin, TX). The sequence

of events in each trial is depicted in Figure 1. In between trials, subjects well always presented

with a default screen consisting of three circles (4.5 cm diameter), displayed for 2000 ms on a

white background. Fifteen randomly arranged tokens (0.3 cm diameter) then appeared in the

central circle. After a delay of 800 ms, the tokens began to jump, one-by-one every 200 ms

from the center to one of the two lateral circles (*i.e.*, $Jump_1$ to $Jump_{15}$). The subjects' task was

to indicate by a right index or right middle finger key-press which lateral circle they thought

would ultimately receive the majority of the tokens (*i.e.*, F9 and F8 key-presses to choose left

and right circles, respectively). They could provide their response as soon as they felt

sufficiently confident, but between $Jump_1$ and $Jump_{15}$. Once the response was provided, the

tokens kept on jumping every 200 ms until the central circle was empty. At this time, the

selected circle was highlighted either in green or in red depending on whether the response was

correct or incorrect, respectively, and a score was displayed above the central circle to provide

the subjects with further feedback of their performance. In correct trials, subjects received a

positive score (*i.e.*, a monetary reward) which was equal to the number of tokens remaining in

the central circle at the time of the response (in € cents). Conversely, incorrect responses led to

a fixed penalty of 7 cents, regardless of the RT. Thus, the longer the subjects waited to provide

a response, the lower was the reward/penalty ratio, generating an increasing sense of urgency

as time passed within each trial. In the absence of any response before $Jump_{15}$, the central circle

was highlighted in red and a "Time Out" (TO) message appeared on the top of the screen. The

subjects were neither rewarded nor penalized in these trials. The feedback cue remained on the

screen for 1000 ms and then disappeared at the same time as the tokens did, denoting the end

of the trial. Subjects were told that they would receive a monetary reward at the end of the

experiment corresponding to their final score. Each trial lasted 6600 ms.

Figure 1: Schematic of the tokens task. In each trial, 15 tokens jumped one-by-one every 200
ms from the central circle to one of the two lateral circles (*i.e.*, $Jump_1$ to $Jump_{15}$). The subjects
had to indicate by a right index or right middle finger key-press (*i.e.*, F9 and F8 keys,
respectively) which lateral circle they thought would receive more tokens (*i.e.*, left or right
circle, respectively) at the end of the trial. They could provide their response whenever they
wanted between $Jump_1$ and $Jump_{15}$. For a correct response, the subjects earned, in € cents, the
number of tokens remaining in the central circle at the time of the response. Hence, the reward
received for a correct response decreased over time, as depicted on the upper right side of the
figure (green trace). The example presented on the lower left side of the figure represents a
correct response provided between $Jump_9$ and $Jump_{10}$ (*i.e.*, the score indicates that 6 tokens
remained in the central circle at the moment the right circle was chosen). In contrast, if subjects
chose the incorrect lateral circle, they lost 7 € cents, regardless of their RT. As such, the penalty
score was fixed, as shown in red on the upper right side of the figure: the lower middle example
represents an incorrect choice of the left circle. Thus, the reward/penalty ratio decreased over
time, producing an increasing sense of urgency over the course of a trial. In the absence of
response ("Time Out" trial, lower right side example), subjects were neither rewarded, nor
penalized (score = 0). For representative purposes, the "Time Out" message is depicted below
the circles in this example, while it was presented above them in the experiment.

Blocks and sessions

The study included 3 sessions, conducted on separate days at a 24-hour interval. Testing

always occurred at the same time of the day for a given subject, to avoid variations that could

be due to changes in chronobiologic states (Derosiere et al., 2015a; Schmidt et al., 2006). Each

session comprised 4 blocks of 50 trials, with each block lasting about 5.5 minutes. Subjects also

performed 4 blocks of 5 trials of a simple reaction time (SRT) task, two at the beginning and

two at the end of each session. In the SRT task, subjects were presented with the same display

as in the tokens task described above. However, after 50 ms in the central circle, the 15 tokens

all jumped together into one of the two lateral circles at the same time. Subjects were instructed

to respond to this "GO signal" by pressing the corresponding key as fast as possible.

Importantly, the 15 tokens always jumped into the same lateral circle in all trials of a given SRT

block and subjects were told which lateral circle this would be in advance. This SRT task

allowed us to estimate the sum of the delays attributable to the motor processes ensuing from

commitment to an action in the absence of a choice (see Cisek et al., 2009; Thura et al., 2014).

Day 1 served as a training session. Day 2 and 3 corresponded to the actual experimental

sessions with the cTBS intervention (Figure 2). cTBS was applied before subjects engaged in

the blocks of trials, either over the left M1 hand area (M1-Disruption session) or over the right

primary somatosensory cortex (S1), 2 cm behind the right M1 area (Sham session; Derosiere et

al., 2014; Alexandre et al., 2015; Torta et al., 2013), in a randomized order. The Sham session

allowed us to ensure that the putative behavioral effects observed following M1 cTBS were not

due to the tactile and auditory sensations elicited by the TMS pulses (Derosiere et al., 2017a,

2017b).

TMS procedure

TMS was delivered through a 2x75 mm figure-of-eight coil connected to a Magpro X100

Stimulator (Magventure Company, Farum, Denmark). The coil was placed tangentially on the

scalp with the handle oriented towards the back of the head and laterally at a 45° angle away

from the midline. At the beginning of each session, M1 was localized by identifying the optimal

spot for eliciting MEPs in the first dorsal interosseous (FDI) muscle of the right hand (M1-

Disruption session) or the left hand (Sham session). This site (called the "hotspot") was marked

on an electroencephalography cap fitted on the participant's head to provide a reference point

throughout the session (Duque et al., 2010, 2014; Vandermeeren et al., 2009). We then

determined the resting motor threshold (rMT) at the hotspot. The rMT was defined as the

minimal intensity required to evoke MEPs of 50 µV peak-to-peak in the targeted muscle on 5

out of 10 consecutive trials (Grandjean et al., 2018; Vassiliadis et al., 2018).

The cTBS procedure consisted of a series of short TMS trains (three pulses at 50 Hz)

repeated every 200 ms for a total duration of 40 s (600 pulses) at an intensity of 80 % of rMT

(Derosiere et al., 2017a, 2017b; Huang et al., 2005; Solopchuk et al., 2017). Such an

intervention has been shown to inhibit the stimulated cortical area, producing a temporary

"virtual lesion", often effective as soon as the train is over (Derosiere et al 2017a, 2017b; Do et

al., 2018; Sasaki et al. 2018) and lasting for between 20 (Clerget et al., 2011; Oberman et al.,

2011; Zénon et al., 2015) and 45 minutes (Huang et al., 2005).

In order to monitor the inhibitory effect of cTBS on motor activity, single TMS pulses were

applied at the M1 hotspot at 115 % of the rMT to elicit MEPs at different time points in the

M1-Disruption and Sham sessions (Klein et al., 2014; Labruna et al., 2014; Quoilin et al., 2016,

2017). In the M1-Disruption session, MEPs were recorded in the right FDI following TMS over

left M1, to evaluate the impact of left M1 cTBS on left motor excitability. In the Sham session,

MEPs were obtained from the left FDI following TMS over right M1, to control for the absence

of effect of right S1 cTBS on right M1 excitability.

The time points at which MEPs were elicited in the M1-Disruption and Sham sessions were

comparable (see Figure 2). In both sessions, 20 MEPs were elicited at the beginning of the

session (*i.e.*, just before cTBS; $TMS_{Baseline}$). Then, 15 MEPs were elicited just following cTBS

(TMS_1), after the two initial SRT blocks (TMS_2), and after each block of trials of the tokens

task (TMS_{3-6}). Finally, 20 additional MEPs were evoked following the two last SRT blocks

(TMS_7). These seven timings (TMS_{1-7}) fell 1 min, 3 min, 11 min, 19 min, 27 min, and

37 min after the cTBS intervention, respectively.

Figure 2: Experimental protocol. Subjects came to the lab for 3 consecutive days (Day 1, 2
and 3). On each day, they performed the tokens task during 4 blocks of 50 trials (Block$_{1-4}$; light
grey rectangles). They also performed Simple Reaction Time blocks at the beginning and at the
end of each session (SRT$_1$ and SRT$_2$, respectively; dark grey rectangles). Day 1 served as a
training session and did not involve any continuous Theta Burst Stimulation (cTBS). A cTBS
train was applied for 40 s at the beginning of Day 2 and 3 (red rectangles), either over the left
(L) primary motor cortex (M1-Disruption session, magenta coil) or over the right (R) primary
somatosensory cortex (Sham session, cyan coil), in a randomized order. Motor Evoked
Potentials (MEPs) were elicited at different time points (TMS$_{EPOCHS}$) throughout the sessions
(TMS$_{Baseline}$ and TMS$_{1-7}$, yellow rectangles), either in the R first dorsal interosseous (FDI)
muscle (M1-Disruption session) or in the L FDI (Sham session), by applying single-pulse TMS
over the L or R M1, respectively. Note that in the Sham session, this implied targeting different
sites for the cTBS intervention (R S1) and the MEP assessments (R M1; coil position not shown
on the figure).

Data collection

Electromyography (EMG) was used to measure the peak-to-peak amplitude of FDI MEPs

elicited by single TMS pulses over the contralateral M1. EMG activity was recorded from

surface electrodes placed over the right FDI or the left FDI (M1-Disruption or Sham sessions,

respectively). EMG data were collected for 1000 ms on each trial, starting 300 ms before the

TMS pulse. EMG signals were amplified, bandpass filtered on-line (10-500 Hz) and digitized

at 2000 Hz for off-line analysis.

Data analysis

Motor evoked potential data

MEP data were collected with Signal (Signal 3.0, Cambridge, UK) and analyzed with

custom Signal scripts. MEP amplitudes were measured for each TMS pulse. Trials with

background EMG activity greater than 100 μV in the 200-ms window preceding the TMS

artifact were excluded from the analysis (Derosiere et al., 2015b; Labruna et al. 2014). 3.63 ±

5.43 % of trials were discarded based on this criterion. Moreover, the data of three subjects had

to be discarded because they presented outlying MEP values – *i.e.*, values exceeding 2.5 SD

above the mean of the group, reflecting a large increase in motor excitability following M1

cTBS. This indicates that cTBS was not effective in inducing a virtual lesion in these subjects,

consistent with previous reports of inter-individual differences in the efficiency of the

intervention (Do et al., 2018; Jannati et al., 2017; Rocchi et al., 2018). The analyses of both the

MEP and the behavioral data were performed on the remaining pool of subjects (n = 16). The

amplitude of MEPs elicited at each time point was averaged to obtain a measure of motor

excitability at $TMS_{Baseline}$ and at TMS_{1-7} in the M1-Disruption and Sham sessions (Figure 2).

For each session, we then expressed MEP amplitudes obtained at TMS_{1-7} (*i.e.*, after the cTBS

intervention) in percentage of the amplitudes measured at $TMS_{Baseline}$ (*i.e.*, before the cTBS

intervention).

Behavioral data

Behavioral data were collected with LabVIEW 8.2 (National Instruments, Austin, TX),

stored in a database (Microsoft SQL Server 2005, Redmond, WA), and analyzed with custom

MATLAB scripts (MathWorks, Natick, MA). Because Day 1 served as a training session (see

section *Blocks and sessions*), the behavioral analyses focused on the data acquired on Day 2

and 3 (*i.e.*, in the M1-Disruption and Sham sessions).

SRT task

For each subject, we computed the mean SRT for both fingers (right index and right middle

fingers), defined as the difference between the time at which subjects pressed the key and the

time at which the 15 tokens appeared simultaneously in the lateral circle, obtained at the

beginning (SRT_1) and at the end (SRT_2) of each session. This SRT allowed us to quantify the

impact of M1 cTBS on the motor processes that follow commitment to an action in the absence

of a choice.

Tokens task

Methods to analyze behavior in the tokens task have been described previously (Cisek et

al., 2009). Briefly, the task allows us to calculate, at each moment in time, the "success

probability" $p_i(t)$ associated with choosing each lateral circle *i*. For instance, assuming a total

of 15 tokens, if at a particular moment in time the right (R) circle contains N_R tokens, the left

(L) circle contains N_L tokens, and the central (C) circle contains N_C tokens, then the probability

that the circle on the right will ultimately be the correct one is described as follows:

$$p(\text{R}|N_R, N_L, N_C) = \text{N} \frac{N_C!}{2^{N_C}} \sum_{k=0}^{\min(N_C, 7-N_L)} \frac{1}{k!\,(N_C - k)!}$$

322   *(1)*

For some of the analyses, we pooled trials according to a classification determined based

on the temporal profile of $p_i(t)$. That is, although the side of each token jump was completely

random in each trial, we could classify some trials as belonging to one of two specific groups a

posteriori. Trials were categorized as "obvious" when the $p_i(t)$ was above 0.6 after Jump_2 and

above 0.75 after five; that is, the initial token jumps consistently favored the correct circle.

Other trials were categorized as "ambiguous" when the initial jumps were balanced between

the lateral circles, keeping the $p_i(t)$ close to 0.5 until late in the trial: $p_i(t)$ remained between 0.5

and 0.66 up to Jump_7 in these trials. For each trial type (obvious and ambiguous; trials that were

neither obvious nor ambiguous were not considered here) and each session (M1-Disruption and

Sham), we analyzed the following behavioral variables: the decision time (DT), the percentage

of correct choices (%Correct) and the percentage of time out trials (%TO). To estimate the DT,

we first calculated the RT during the tokens task by computing the difference between the time

at which subjects pressed the key and Jump_1. We then subtracted the mean RT (SRT_1 and SRT_2

pooled together) estimated in the SRT task on the same day from this RT for each subject to

yield an estimate of DT – *i.e.*, reflecting the duration of the deliberation process.

One key feature of the tokens task is that it allows us to calculate the available sensory

evidence for the chosen circle at the time when commitment is made (*i.e.*, at decision time

[DT]). Although we believe that subjects cannot calculate equation 1 to compute sensory

evidence after each jump, we can expect that they can make a reasonable estimate. For instance,

a simple "first-order" approximation of sensory evidence is the sum of log-likelihood ratios

(SumLogLR) of individual token movements, calculated as follows:

344
$$SumLogLR(n) = \sum_{k=1}^{n} log \frac{p(e_k|S)}{p(e_k|U)}$$

345   *(2)*

Where $p(e_k/S)$ is the likelihood of a token event e_k (a token jumping into either the selected

or unselected lateral circle) during trials in which the selected target S is correct and (e_k/U) is

the likelihood of e_k during trials in which the unselected circle U is correct. The SumLogLR is

proportional to the difference between the number of tokens that moved in each direction before

the decision. To characterize the decision policy of the subjects in the Sham and M1-Disruption

sessions, we binned trials as a function of the total number of tokens that moved before the

decision, and calculated the average SumLogLR for each bin. Seven bins were defined, with

the first bin (Bin_1) including responses provided between $Jump_5$ and $Jump_6$, the second bin

(Bin_2) including responses provided between $Jump_6$ and $Jump_7$ and so on, until the last bin

(Bin_7) covering the period between $Jump_{11}$ and $Jump_{12}$. SumLogLR at DT preceding $Jump_5$ or

following $Jump_{12}$ were not considered for this analysis because part of the subjects did not

respond at these timings. Importantly, the SumLogLR at DT was computed based on every trial

where a response was provided (*i.e.*, for correct and incorrect responses, in obvious and

ambiguous trials, as well as in other trials with different $p_i(t)$).

According to recent models of decision-making, action choices result from the combination

of signals that track the available sensory evidence and the level of urgency that grows over

time (Cisek et al., 2009, Ditterich, 2006, Drugowitsch et al., 2012). A direct prediction of such

urgency-based models is that decisions made with lower levels of evidence should be associated

with higher levels of urgency. Hence, we exploited the SumLogLR at DT values to estimate an

urgency function for each subject and each block condition. To do so, we rectified the

SumLogLR at DT (*i.e.*, by multiplying it by -1 and adding a constant of 3) and then fitted a

linear regression over the rectified values. We then extracted the intercept and the slope of these

urgency functions, which we used as estimates of the initial level and the growth rate of the

urgency signal, respectively. Second-order polynomial regressions were also performed on the

rectified SumLogLR data but did not yield significantly better fits (*i.e.*, group-level Bayesian

Information Criterion [BIC] values for linear and polynomial fits were 5.75 ± 1.61 and $5.74 \pm$

1.79 a.u., respectively; $t_{14} = 0.02$, $p = .984$).

Statistical analysis

All statistical analyses were performed with custom R Scripts (R Version 3.4.1, Car and

BayesFactor packages). MEP data (expressed in percentage of $TMS_{Baseline}$) were analyzed using

a two-way repeated-measure ANOVA ($ANOVA_{RM}$) with SESSION (M1-Disruption, Sham)

and TMS_{EPOCH} (TMS_1, TMS_2, TMS_3, TMS_4, TMS_5, TMS_6, TMS_7) as within-subject factors.

Similarly, the SRT data were analyzed using a two-way $ANOVA_{RM}$ with SESSION (M1-

Disruption, Sham) and SRT_{EPOCH} (SRT_1, SRT_2) as within-subject factors. The DT, the

%Correct and the %TO data were analyzed using two-way $ANOVA_{RM}$ with SESSION (M1-

Disruption, Sham) and TRIAL (obvious, ambiguous) as within-subject factors. The SumLogLR

at DT was analyzed using a two-way $ANOVA_{RM}$ with SESSION (M1-Disruption, Sham) and

BIN (Bin_1 to Bin_7) as within-subject factors. When appropriate, Fisher's LSD post-hoc tests

were used to detect paired differences in these ANOVAs. Furthermore, the intercept and the

slope of the urgency functions were compared between each session using Student's t-tests. The

significance level for these tests was set at $p < .05$ Results are expressed as mean $\pm$ SE.

RESULTS

Motor evoked potentials

As mentioned above (see section *TMS procedure*), during M1-Disruption sessions, MEPs were recorded in the right FDI following single-pulse TMS over left M1, in order to evaluate the impact of left M1 cTBS on left M1 excitability. Conversely, in the Sham sessions, MEPs were obtained from the left FDI following TMS over right M1, to control for the absence of effect of right S1 cTBS on right M1 excitability. The MEPs elicited following cTBS (TMS$_{1-7}$) were expressed in percentage of MEPs elicited before (TMS$_{Baseline}$), allowing us to directly compare potential MEP changes elicited by the intervention in the M1-Disruption and Sham sessions.

The ANOVA$_{RM}$ revealed a main effect of the factor SESSION on these MEP changes ($F_{1,15}$ = 15.41; p = .001; see Figure 3). As such, percentage MEP values were lower following cTBS in the M1-Disruption session (78.78 ± 3.04 %) than in the Sham session (105.53 ± 5.32 %; TMS$_{1-7}$ timings pooled together). This effect on percentage MEP values did not depend on time as the ANOVA$_{RM}$ did not reveal any effect of the factor TMS$_{EPOCH}$ ($F_{6,90}$ = 1.99, p = .075) nor interaction with the factor SESSION ($F_{6,90}$ = 0.72, p = .637). Hence, percentage MEP values remained stable following the cTBS intervention; they were consistently lower in the M1-Disruption than in the Sham session, regardless of the time at which MEPs were considered during the course of the experiment.

Additional Student's t-tests against 100 % (run for each TMS$_{EPOCHS}$) showed that, as expected, the MEP difference between the two sessions reported above was due to a selective suppression of MEPs in the M1-Disruption but not in the Sham session. As such, percentage MEP values were significantly lower than 100 % at almost all timings in the M1-Disruption session, except for the last one (*i.e.*, at TMS$_{1-6}$; all p-values = [.000008 .017]). Conversely, amplitudes were not significantly different from 100 % (*i.e.,* from TMS$_{Baseline}$) in the Sham session (all p-values = [.163 .826]).

Figure 3: Motor Evoked Potentials (MEPs) amplitudes. A. Mean value of MEPs (in percentage of MEPs at $TMS_{Baseline}$) elicited after the cTBS intervention in the First Dorsal Interosseous (FDI) muscle at each of the TMS_{EPOCHS} (TMS_{1-7}) in the M1-Disruption (magenta traces) and Sham (cyan traces) sessions. Note the significant disruption of MEPs with respect to baseline (*i.e.*, dashed horizontal line) in the M1-Disruption session (#: significantly different from 100 at p< .05). **B. Cumulative percentage of subjects.** MEPs obtained at TMS_{1-7} are pooled together. Note the larger percentage of subjects showing MEPs smaller than 100% in the M1-disruption than in the sham session, as also evident from the inset representing the group-level average with the effect of SESSION (*: significantly different at p < .05). Error bars represent SE.

Behavioral data

Simple Reaction Time (SRT)

The $ANOVA_{RM}$ revealed a main effect of the factor SESSION on the SRT data ($F_{1,15} = 5.34$, p = .035; see Figure 4.A). Indeed, SRTs were significantly prolonged in the M1-Disruption session (237.31 ± 7.44 ms) compared to the Sham session (220.83 ± 5.62 ms; SRT_1 and SRT_2 pooled together). The impact of M1 disruption on SRTs did not vary over the course of the session. As such, the $ANOVA_{RM}$ did not reveal any significant effect of the factor SRT_{EPOCH} ($F_{1,15} = 0.02$, p = .892) nor of its interaction with the factor SESSION ($F_{1,15} = 1.78$,

p = .202). These findings indicated that M1-disruption altered the motor processes underlying

initiation and/or execution of the cued movements (Huang et al., 2005; Lakhani et al., 2014;

McAllister et al., 2013).

Tokens task

Decision time (DT)

As expected, the ANOVA$_{RM}$ revealed a main effect of the factor TRIAL on the DT data

($F_{1,15}$ = 119.30, p < .00001; see Figure 4.B). Indeed, DTs were significantly shorter in obvious

trials (871.77 ± 57.14 ms) than in ambiguous ones (1426.57 ± 88.59 ms; M1-Disruption and

Sham sessions pooled together), consistent with the literature (*e.g.,* Thura et al., 2014; Thura

and Cisek, 2014a, 2016, 2017). Critically, the ANOVA$_{RM}$ did not reveal any significant effect

of the factor SESSION ($F_{1,15}$ = 0.24, p = .631) nor of its interaction with the factor TRIAL ($F_{1,15}$

= 0.02, p = .888). Hence, the time taken by the subjects to deliberate depended on the trial type

they encountered (*i.e.*, obvious vs ambiguous) but was not affected by M1 disruption.

Percentage of correct choices (%Correct)

Also consistent with the literature is the significant main effect of the factor TRIAL for the

%Correct data ($F_{1,15}$ = 31.727, p = .00005; see Figure 4.C; *e.g.,* Thura et al., 2014; Thura and

Cisek, 2014a, 2016, 2017). Indeed, %Correct was significantly higher in obvious trials (99.80

± 0.13 %) than in ambiguous ones (88.61 ± 1.80 %; M1-Disruption and Sham sessions pooled

together). However, neither the factor SESSION ($F_{1,15}$ = 0.05, p = .497) nor its interaction with

the factor TRIAL were significant ($F_{1,15}$ = 0.44, p = .517). Hence, M1 disruption did not alter

the accuracy of the decision process.

<u>Percentage of Time Out trials (%TO)</u>

The %TO data revealed a similar pattern as the variables described above. Indeed, the %TO

was significantly lower in obvious (0.20 ± 0.13 %) than in ambiguous trials (4.43 ± 1.42 %;

M1-Disruption and Sham sessions pooled together), as confirmed by the ANOVA$_{RM}$ (factor

TRIAL: $F_{1,15} = 7.47$, p = .015; see Figure 4.D). Moreover, there was no effect of the factor

SESSION ($F_{1,15} = 0.01$, p = .920) or interaction with the factor TRIAL ($F_{1,15} = 0.01$, p = .922).

Hence, the proportion of trials in which subjects refrained from responding was unaffected by

M1-Disruption.

<u>Sensory evidence at decision time (SumLogLR at DT) and urgency</u>

The amount of sensory evidence based on which subjects made their decision was

estimated using the SumLogLR (computed at DT): the higher the SumlogLR, the higher the

evidence at DT (Cisek, 2009; Thura et al., 2012, 2014). SumLogLR values are presented for

each Bin$_n$ on Figure 4.E (see *Methods*), separately for the M1-Disruption and the Sham sessions.

Note that two subjects were excluded from this analysis as they responded too early on most

trials, resulting in a lack of SumLogLR values after Jump$_9$ in these participants. Hence,

SumLogLR analyses were run on 14 subjects.

Overall, fast decisions were made based on more sensory evidence than slow decisions, as

confirmed by the ANOVA$_{RM}$ showing a main effect of the factor BIN on the SumLogLR at DT

($F_{6,78} = 12.86$, p < .00001; see Figure 4.E; BF value above 100). Indeed, the SumLogLR at DT

was significantly higher at Bin$_1$ (1.18 ± 0.06 a.u.) than for any other bin after Bin$_4$ (all

SumLogLR at DT $\leq 0.84 \pm 0.05$ a.u.; M1-Disruption and Sham sessions pooled together).

Hence, the amount of sensory evidence based on which subjects made their choices decreased

as a function of time, in agreement with previous studies in which changing sensory evidence

was used to guide subjects' choices (*e.g.,* Cisek et al., 2009; Gluth et al., 2012, Murphy et al.,

2016; Thura et al., 2012, 2014; Thura and Cisek, 2014a, 2016, 2017). Here again, the

ANOVA$_{RM}$ did not reveal any significant effect of the factor SESSION ($F_{6,78} = 0.04$, p = .852)

nor of its interaction with the factor BIN ($F_{6,78} = 0.41$, p = .868). Hence, subjects made their

decisions based on a similar amount of sensory evidence in both sessions, suggesting a

preservation of the urgency drive during deliberation with M1 disruption.

To further confirm this finding, we obtained a simple approximation of the urgency signal

underlying the subjects' decisions by fitting a linear regression over the rectified version of

SumLogLR at DT for each session (M1-Disruption, Sham) and extracted the intercept and the

slope of the regression functions (see section *Methods* and Figure 4. F, G and H). Again, student

t-tests did not reveal any significant impact of the session on the intercept ($t_{14} = 0.48$, p =.798)

or the slope ($t_{14} = -0.22$, p =.832) of the urgency functions.

Figure 4: Behavioral results. A. Mean Simple Reaction Time (SRT) (obtained in the SRT task) measured at each of the SRT_{EPOCHS} (SRT_{1-2}) in the M1-Disruption (magenta bars) and Sham (cyan bars) sessions. **B. Mean Decision Time (DT)** measured in each TRIAL (obvious, ambiguous) in the M1-Disruption (magenta bars) and Sham (cyan bars) sessions. **C** and **D** same as B for the percentages of correct choices (%Correct) and of Time Out trials (%TO),

respectively. **E. SumLogLR at DT** measured in each Bin_n of decision time (*i.e.*, between $Jump_5$
and $Jump_{12}$, see main text) in the M1-Disruption (magenta traces) and Sham (cyan traces)
sessions. **F. Urgency functions** computed based on the rectified SumLogLR at DT for the M1-
Disruption (magenta traces) and Sham (cyan traces) sessions. The small bar graphs on the
bottom right represent the group-level average intercept and slope of the functions. **G.**
Individual intercept values, represented for the M1-Disruption Session (y-axis) as a function
of the values for the Sham session (x-axis). Points above the diagonal (n = 7/14) represent the
subjects showing a higher intercept in the M1-Disruption than following Sham session, while
points below the diagonal (n = 7/14) represent the subjects showing a lower intercept in the
M1-Disruption than following Sham session. **H**. Same as G. for the slope values. *: significant
difference (p < .05). Error bars represent SE.

<u>Verifying that M1 disruption does not impact deliberation using Bayesian analyses</u>

The $ANOVA_{RM}$ and the t-tests revealed that M1 disruption did not significantly alter the

behavioral data measured in the tokens task, suggesting that M1 is not functionally involved in

the deliberation process underlying action choices. In order to confirm this result, a Bayes

Factor (BF) was computed for each analysis involving the factor SESSION (10 BF values

obtained in total), providing us with a ratio of the likelihood probability of the null hypothesis

(*i.e.*, H0: the probability that data do not exhibit an effect of SESSION) over the alternative

hypothesis (*i.e.*, H1: the probability that data exhibit the effect; Morey and Rouder 2011). A BF

value of 1 would reflect an equal probability that H0 and H1 are correct, while a BF value above

1 would reflect a higher probability that H0 is correct. In accordance with conventional

interpretation of BF values (Jeffreys, 1961), a BF value ranging between 1 and 3 is interpreted

as indicating *anecdotal* evidence in favor of H0, a value between 3 and 10 as indicating

substantial evidence for H0 and a value between 10 and 30 a *strong* evidence for H0.

Table 1 summarizes the BF values obtained for each factor tested. The average BF value

was of 5.66 ± 1.74 (all BFs ranged between 3.46 and 21.28) indicating substantial to strong

evidence for an absence of difference in subjects' behavior between the M1-Disruption and the

Sham sessions. Hence, Bayesian analyses further reinforce the conclusion that M1 disruption

did not influence the deliberation process underlying decision-making, but solely altered the motor processes that ensue commitment to an action.

Factor tested	DT	%Correct	%TO	SumLogLR	Urgency Intercept	Urgency Slope
SESSION	3.87	3.77	3.91	5.23	3.46	3.48
SESSION*TRIAL	3.91	3.77	3.91	---	---	---
SESSION*TIME	---	---	---	21.28	---	---

Table 1: Bayes factor (BF) values. The first column specifies the factors tested for which a BF was computed. Other columns represent the BFs obtained for each behavioral measure in the tokens task. Overall, BFs ranged between 3.46 and 21.28 indicating substantial to strong evidence for a lack of effect of the SESSION on subjects' behavior.

DISCUSSION

Previous studies have suggested that urgency amplifies motor activity during decisions about actions (Steinemann et al., 2018, Thura and Cisek, 2014a, 2016). To date, however, the specific contribution of motor cortical areas to the decision process remains unclear. Here, we investigated whether M1 itself contributes to deliberation during action choices, or whether it simply reflects decision signals coming from upstream structures and contributes exclusively to the motor processes overtly expressing commitment. To do so, we tested the impact of a disruption of M1 activity, induced by continuous theta burst stimulation (cTBS), on the behavior of human subjects in *(1)* a simple reaction time (SRT) task allowing us to estimate the duration of the motor processes and *(2)* a modified version of the tokens task (Cisek et al., 2009), which allowed us to estimate subjects' time of commitment as well as accuracy criterion.

Subjects were generally faster and more accurate in obvious than in ambiguous trials,

suggesting that late decisions relied on weaker sensory evidence compared to early decisions.

This is confirmed by the systematic analysis of the sensory evidence available at DT, which

indicates that subjects committed to a choice based on less sensory evidence as time elapsed

during the course of a trial. This dropping of the accuracy criterion is consistent with previous

studies in which similar tasks were used (*e.g.,* Cisek et al., 2009; Gluth et al., 2012; Murphy et

al., 2016; Thura et al., 2012, 2014;) and supports recent models postulating that urgency grows

over time during speeded decisions (Churchland et al., 2008; Ditterich, 2006, Drugowitsch et

al., 2012 Hanks et al., 2011).

The efficiency of the cTBS intervention was attested by a reduction in MEP amplitudes in

the entire M1 disruption session, compared to amplitudes obtained following a sham

stimulation. Moreover, M1 cTBS lengthened the SRTs (*i.e.*, compared to when sham cTBS was

performed), indicating that motor processes that are known to involve M1 were successfully

perturbed by the intervention (Huang et al., 2005; Lakhani et al., 2014; McAllister et al., 2013).

Notably, in the present study, motor responses were recorded through key-presses. Hence, RTs

involved both a premotor and a motor time – *i.e.*, times occurring before and after movement

initiation, respectively (Spieser et al., 2017). As a consequence, the lengthening of SRT

observed here might reflect an increase of the time needed for both initiating and executing the

required motor response.

Critically, all the behavioral data collected in the tokens task were similar in the two cTBS

sessions, whether M1 was disrupted or not. This finding suggests that the contribution of M1

to decision-making is negligible and that past reports of decision-related changes in M1 are

likely to reflect the influence of signals coming from upstream structures rather than an actual

involvement in the deliberation process itself (Thura and Cisek, 2017; van Maanen et al., 2016).

In a similar vein, our recent work shows that M1 disruption negatively alters value-based

choices when action values are freshly acquired but does not following consolidation,

suggesting that M1 contribution to value-based decision-making vanishes as subjects become

more proficient at using the value information (Derosiere et al., 2017a, 2017b). As such, the

causal involvement of M1 may be restricted to the motor processes that follow commitment to

an action.

If M1 does not influence action selection, where in the brain are decisions about actions

determined? Among many possible areas, PMd emerges as a promising candidate. First, single-

cell recordings in behaving monkeys have shown that during deliberation, activity of some PMd

neurons tuned for a particular action reflects the unfolding sensory evidence favoring that

action. This observation also makes it possible that this decision-related activity influences M1

neurons through cortico-cortical projections (Duque et al., 2012; Martinez-Gracia et al., 2015).

Second, the same studies found that PMd activity related to the selected target reaches a peak

about 280 ms before movement initiation regardless of decision difficulty, while a peak of M1

activity occurs about 140 ms later (Thura and Cisek, 2014a). Third, neurons in the globus

pallidus internus, which are insensitive to sensory information during deliberation, become

directionally tuned around the time of the PMd activity peak (Thura and Cisek, 2017).

Altogether, these results suggest that PMd might be the primary site where decisions are

computed and commitment determined.

In agreement with this hypothesis, a recent study found that microstimulation of PMd

neurons alters the deliberation duration, especially if current is applied shortly before

commitment time (Thura and Cisek, 2014b). Crucially, stimulation has much less influence on

decision duration if it is applied long before commitment or between commitment and

movement onset. Relevant for the present work, this microstimulation study also shows similar

time-dependent effect of M1 microstimulation on decision durations but the effect size is

smaller when M1 is stimulated compared to PMd. Finally, other non-primary motor areas may

be causally involved in decision-making, including the pre-supplementary motor area (pre-SMA; Tosun et al., 2017). Investigating their precise contribution represents an interesting issue for future investigations.

Of note, we cannot completely rule out the possibility that the cTBS intervention led to some fast reorganization of the decision network following M1 disruption. According to this idea, M1 might still be part of the network involved in the deliberation process but some compensatory mechanisms may have occurred in this network following M1 disruption, leading to similar decision behaviors after M1 and sham cTBS (Bestmann et al., 2004; Briend et al., 2017; Cash et al., 2017; Derosiere et al., 2017a; Rastogi et al., 2017). One solution to tackle this issue in the future would be to exploit online rTMS techniques (Duque et al., 2010, 2013), which could allow to perturb neural activity at a specific moment during the decision process, leaving less time for compensatory mechanisms to occur. As such, previous work has shown that while online microstimulation of a decision-related area alters behavior during perceptual decision-making (*i.e.*, the lateral intraparietal area; Hanks et al., 2006), (offline) inactivation of the same area does not (Katz et al., 2016).

In conclusion, we show that the offline disruption of M1 activity delays motor processes that follow commitment to an action, but does not alter volitional decision behavior. Taken together, these findings suggest that the contribution of M1 is downstream of the decision process. Future studies should use online disruption protocols and broaden their investigation to the role of non-primary motor areas in deliberation, especially the PMd and the pre-SMA.

REFERENCES

Alamia A, **Zénon A**, **VanRullen R**, **Duque J**, **Derosiere G**. Implicit visual cues tune oscillatory motor activity during decision-making. *Neuroimage* 186: 424–436, 2019.

Alexandre F, **Derosiere G**, **Papaiordanidou M**, **Billot M**, **Varray A**. Cortical motor output decreases after neuromuscular fatigue induced by electrical stimulation of the plantar flexor

muscles. *Acta Physiol* 214: 124–134, 2015.

Bestmann S, **Baudewig J**, **Siebner HR**, **Rothwell JC**, **Frahm J**. Functional MRI of the immediate impact of transcranial magnetic stimulation on cortical and subcortical motor circuits. *Eur J Neurosci* 19: 1950–1962, 2004.

Briend F, **Leroux E**, **Delcroix N**, **Razafimandimby A**, **Etard O**, **Dollfus S**. Impact of rTMS on functional connectivity within the language network in schizophrenia patients with auditory hallucinations. *Schizophr Res* 189: 142–145, 2017.

Cisek P, **Puskas GA**, **El-Murr S**. Decisions in changing conditions: the urgency-gating model. *J Neurosci* 29: 11560–71, 2009.

Clerget E, **Andres M**, **Olivier E**. Deficit in Complex Sequence Processing after a Virtual Lesion of Left BA45. *PLoS One* 8: e63722, 2013.

Derosière G, **Alexandre F**, **Bourdillon N**, **Mandrick K**, **Ward TE**, **Perrey S**. Similar scaling of contralateral and ipsilateral cortical responses during graded unimanual force generation. *Neuroimage* 85: 471–477, 2014.

Derosiere G, **Farrugia N**, **Perrey S**, **Ward T**, **Torre K**. Expectations induced by natural-like temporal fluctuations are independent of attention decrement: evidence from behavior and early visual evoked potentials. *Neuroimage* 104: 278–86, 2015a.

Derosiere G, **Billot M**, **Ward ET**, **Perrey S**. Adaptations of Motor Neural Structures' Activity to Lapses in Attention. *Cereb Cortex* 25: 66–74, 2015b.

Derosiere G, **Klein P-A**, **Nozaradan S**, **Zénon A**, **Mouraux A**, **Duque J**. Visuomotor Correlates of Conflict Expectation in the Context of Motor Decisions. *J Neurosci* 38: 9486–9504, 2018.

Derosiere G, **Vassiliadis P**, **Demaret S**, **Zénon A**, **Duque J**. Learning stage-dependent effect of M1 disruption on value-based motor decisions. *Neuroimage* 162: 173–185, 2017a.

Derosiere G, **Zénon A**, **Alamia A**, **Duque J**. Primary motor cortex contributes to the implementation of implicit value-based rules during motor decisions. *Neuroimage* 146: 1115–1127, 2017b.

Ditterich J. Stochastic models of decisions about motion direction: Behavior and physiology. *Neural Network*. 19: 981–1012, 2006.

Do M, **Kirkovski M**, **Davies CB**, **Bekkali S**, **Byrne LK**, **Enticott PG**. Intra- and Inter-Regional Priming of Ipsilateral Human Primary Motor Cortex With Continuous Theta Burst Stimulation Does Not Induce Consistent Neuroplastic Effects. *Front Hum Neurosci* 12: 123, 2018.

Donner TH, **Siegel M**, **Fries P**, **Engel AK**. Buildup of Choice-Predictive Activity in Human Motor Cortex during Perceptual Decision Making. *Curr Biol* 19: 1581–1585, 2009.

Drugowitsch J, Moreno-Bote R, Churchland AK, Shadlen MN, Pouget A. The Cost of Accumulating Evidence in Perceptual Decision Making. *J Neurosci*, 32:3612–3628. 2012.

Duque J, **Labruna L**, **Cazares C**, **Ivry RB**. Dissociating the influence of response selection and task anticipation on corticospinal suppression during response preparation. *Neuropsychologia* 65: 287–296, 2014.

Duque J, **Labruna L**, **Verset S**, **Olivier E**, **Ivry RB**. Dissociating the role of prefrontal and

premotor cortices in controlling inhibitory mechanisms during motor preparation. *J Neurosci* 32: 806–16, 2012.

Duque J, **Lew D**, **Mazzocchio R**, **Olivier E**, **Ivry RB**. Evidence for two concurrent inhibitory mechanisms during response preparation. *J Neurosci* 30: 3793–802, 2010.

Duque J, **Olivier E**, **Rushworth M**. Top–Down Inhibitory Control Exerted by the Medial Frontal Cortex during Action Selection under Conflict. *J Cogn Neurosci* 25: 1634–1648, 2013.

Forstmann BU, **Dutilh G**, **Brown S**, **Neumann J**, **von Cramon DY**, **Ridderinkhof KR**, **Wagenmakers E-J**. Striatum and pre-SMA facilitate decision-making under time pressure. *Proc Natl Acad Sci* 105: 17538–17542, 2008.

Gluth S, Rieskamp J, Büchel C. Deciding not to decide: computational and neural evidence for hidden behavior in sequential choice. *PLoS comp biol* 9:e1003309, 2012.

Grandjean J, **Derosiere G**, **Vassiliadis P**, **De Wilde Y**, **Quemener L**, **Duque J**. Towards assessing corticospinal excitability bilaterally: Validation of a double-coil TMS method. *J Neurosci Methods* 293: 162–168, 2018.

Hanks TD, **Ditterich J**, **Shadlen MN**. Microstimulation of macaque area LIP affects decision-making in a motion discrimination task. *Nat Neurosci* 9: 682–689, 2006.

Hanks TD, **Mazurek ME**, **Kiani R**, **Hopp E**, **Shadlen MN**. Elapsed Decision Time Affects the Weighting of Prior Probability in a Perceptual Decision Task. J Neurosci 31:6339-6352, 2011.

Hauser TU, **Moutoussis M**, **Purg N**, **Dayan P**, **Dolan RJ**. Beta-Blocker Propranolol Modulates Decision Urgency During Sequential Information Gathering. *J Neurosci* 38: 7170–7178, 2018.

Huang Y-Z, **Edwards MJ**, **Rounis E**, **Bhatia KP**, **Rothwell JC**. Theta Burst Stimulation of the Human Motor Cortex. *Neuron* 45: 201–206, 2005.

Jannati A, **Block G**, **Oberman LM**, **Rotenberg A**, **Pascual-Leone A**. Interindividual variability in response to continuous theta-burst stimulation in healthy adults. *Clin Neurophysiol* 128: 2268–2278, 2017.

Jeffreys H. Theory of probability [Online]. Clarendon Press. https://global.oup.com/academic/product/the-theory-of-probability-9780198503682?cc=be&lang=en& [6 Dec. 2018].

Katz LN, **Yates JL**, **Pillow JW**, **Huk AC**. Dissociated functional significance of decision-related activity in the primate dorsal stream. *Nature* 535: 285–288, 2016.

Klein PA, **Petitjean C**, **Olivier E**, **Duque J**. Top-down suppression of incompatible motor activations during response selection under conflict. *Neuroimage* 86: 138–149, 2014.

Labruna L, **Lebon F**, **Duque J**, **Klein P-A**, **Cazares C**, **Ivry RB**. Generic inhibition of the selected movement and constrained inhibition of nonselected movements during response preparation. *J Cogn Neurosci* 26: 269–78, 2014.

Lakhani B, **Bolton DAE**, **Miyasike-daSilva V**, **Vette AH**, **McIlroy WE**. Speed of processing in the primary motor cortex: A continuous theta burst stimulation study. *Behav Brain Res* 261: 177–184, 2014.

Laming DRJ. Information theory of choice reaction time. Wiley; New York: 1968.

van Maanen L, **Fontanesi L**, **Hawkins GE**, **Forstmann BU**. Striatal activation reflects urgency in perceptual decision making. *Neuroimage* 139: 294–303, 2016.

Martinez-Garcia M, **Insabato A**, **Pannunzi M**, **Pardo-Vazquez JL**, **Acuña C**, **Deco G**. The Encoding of Decision Difficulty and Movement Time in the Primate Premotor Cortex. *PLOS Comput Biol* 11: e1004502, 2015.

Mcallister CJ, **Rönnqvist KC**, **Stanford IM**, **Woodhall GL**, **Furlong PL**, **Hall SD**. Oscillatory Beta Activity Mediates Neuroplastic Effects of Motor Cortex Stimulation in Humans. *J Neurosci* 33: 7919-7927, 2013.

Morey RD, **Rouder JN**. Bayes factor approaches for testing interval null hypotheses. *Psychol Methods* 16: 406–419, 2011.

Murphy PR, **Boonstra E**, **Nieuwenhuis S**. Global gain modulation generates time-dependent urgency during perceptual choice in humans. *Nat Commun* 7: 13526, 2016.

Oberman L, **Edwards D**, **Eldaief M**, **Pascual-Leone A**. Safety of Theta Burst Transcranial Magnetic Stimulation: A Systematic Review of the Literature. *J Clin Neurophysiol* 28: 67–74, 2011.

Quoilin C, **Lambert J**, **Jacob B**, **Klein P-A**, **Duque J**. Comparison of Motor Inhibition in Variants of the Instructed-Delay Choice Reaction Time Task. *PLoS One* 11: e0161964, 2016.

Quoilin C, Wilhelm E, Maurage P, de Timary P, Duque J. Deficient inhibition in alcohol-dependence: let's consider the role of the motor system! *Neuropsychopharm*, 2018.

Rastogi A, **Cash R**, **Dunlop K**, **Vesia M**, **Kucyi A**, **Ghahremani A**, **Downar J**, **Chen J**, **Chen R**. Modulation of cognitive cerebello-cerebral functional connectivity by lateral cerebellar continuous theta burst stimulation. *Neuroimage* 158: 48–57, 2017.

Rocchi L, **Ibáñez J**, **Benussi A**, **Hannah R**, **Rawji V**, **Casula E**, **Rothwell J**. Variability and Predictors of Response to Continuous Theta Burst Stimulation: A TMS-EEG Study. *Front Neurosci* 12: 400, 2018.

Sasaki T, **Kodama S**, **Togashi N**, **Shirota Y**, **Sugiyama Y**, **Tokushige S**, **Inomata-Terada S**, **Terao Y**, **Ugawa Y**, **Hamada M**. The intensity of continuous theta burst stimulation, but not the waveform used to elicit motor evoked potentials, influences its outcome in the human motor cortex. *Brain Stimul* 11: 400–410, 2018.

Schmidt C, **Peigneux P**, **Muto V**, **Schenkel M**, **Knoblauch V**, **Münch M**, **de Quervain DJ-F**, **Wirz-Justice A**, **Cajochen C**. Encoding Difficulty Promotes Postlearning Changes in Sleep Spindle Activity during Napping. *J Neurosci* 26: 8976–8982, 2006.

Seideman JA, **Stanford TR**, **Salinas E**. Saccade metrics reflect decision-making dynamics during urgent choices. *Nat Commun* 9: 2907, 2018.

Solopchuk O, **Alamia A**, **Dricot L**, **Duque J**, **Zénon A**. cTBS disruption of the supplementary motor area perturbs cortical sequence representation but not behavioural performance. *Neuroimage* 163: 34–40, 2017.

Spieser L, **Servant M**, **Hasbroucq T**, **Burle B**. Beyond decision! Motor contribution to speed–accuracy trade-off in decision-making. *Psychon Bull Rev* 24: 950–956, 2017.

Steinemann NA, O'Connell RG, Kelly SP. Decisions are expedited through multiple neural adjustments spanning the sensorimotor hierarchy. *Nat. Commun* 9(1):3627, 2018.

Thura D, **Beauregard-Racine J**, **Fradet C-W**, **Cisek P**. Decision making by urgency gating: theory and experimental support. *J Neurophysiol* 108: 2912–2930, 2012.

Thura D, **Cisek P**. Deliberation and Commitment in the Premotor and Primary Motor Cortex during Dynamic Decision Making. *Neuron* 81: 1401–1416, 2014a.

Thura D, **Cisek P**. Micro-stimulation of premotor and motor cortex delays the commitment to an action choice. Program No. 651.04. 2014 Neuroscience Meeting Planner. Washington, DC: Society for Neuroscience, 2014b.

Thura D, Cisek P. Modulation of premotor and primary motor cortical activity during volitional adjustments of speed-accuracy trade-offs. *J Neurosci* 36: 938-956. 2016

Thura D, **Cisek P**. The Basal Ganglia Do Not Select Reach Targets but Control the Urgency of Commitment. *Neuron* 95: 1160–1170.e5, 2017.

Thura D, **Cos I**, **Trung J**, **Cisek P**. Context-dependent urgency influences speed-accuracy trade-offs in decision-making and movement execution. *J Neurosci* 34: 16442–54, 2014.

Torta D, **Legrain V**, **Algoet M**, **Olivier E**, **Duque J**, **Mouraux A**. Theta Burst Stimulation Applied over Primary Motor and Somatosensory Cortices Produces Analgesia Unrelated to the Changes in Nociceptive Event-Related Potentials. *PloS One* 8: e73263, 2013.

Tosoni A, **Corbetta M**, **Calluso C**, **Committeri G**, **Pezzulo G**, **Romani GL**, **Galati G**. Decision and action planning signals in human posterior parietal cortex during delayed perceptual choices. *Eur J Neurosci* 39: 1370–1383, 2014.

Tosun T, **Berkay D**, **Sack AT**, **Çakmak YÖ**, **Balcı F**. Inhibition of Pre–Supplementary Motor Area by Continuous Theta Burst Stimulation Leads to More Cautious Decision-making and More Efficient Sensory Evidence Integration. *J Cogn Neurosci* 29: 1433–1444, 2017.

Ratcliff R. A theory of memory retrieval. *Psych rev* 85:59, 1978.

Stone M. Models for choice-reaction time. *Psychometrika* 25:251-260, 1960.

Vandermeeren Y, **Davare M**, **Duque J**, **Olivier E**. Reorganization of cortical hand representation in congenital hemiplegia. *Eur J Neurosci* 29: 845–854, 2009.

Vassiliadis P, **Grandjean J**, **Derosiere G**, **de Wilde Y**, **Quemener L**, **Duque J**. Using a Double-Coil TMS Protocol to Assess Preparatory Inhibition Bilaterally. *Front Neurosci* 12: 139, 2018.

Wyart V, **de Gardelle V**, **Scholl J**, **Summerfield C**. Rhythmic Fluctuations in Evidence Accumulation during Decision Making in the Human Brain. *Neuron* 76: 847–858, 2012.

Zénon A, **Klein P-A**, **Alamia A**, **Boursoit F**, **Wilhelm E**, **Duque J**. Increased Reliance on Value-based Decision Processes Following Motor Cortex Disruption. *Brain Stimul* 8: 957–964, 2015.

GRANTS

This work was supported by grants from the "Fonds Spéciaux de Recherche" (FSR) of the Université Catholique de Louvain, the Belgian National Funds for Scientific Research (FRS-

790 FNRS: MIS F.4512.14) and the "Fondation Médicale Reine Elisabeth" (FMRE). GD was a
791 postdoctoral fellow supported by the FNRS.

AUTHOR CONTRIBUTIONS

795 G.D., D.T., P.C. and J.D. conceived and designed research; G.D. performed experiments; G.D.
796 analyzed data; G.D., D.T., P.C. and J.D. interpreted results of experiments; G.D. prepared
797 figures; G.D. and J.D. drafted manuscript; G.D., D.T., P.C. and J.D. edited and revised
798 manuscript; G.D., D.T., P.C. and J.D. approved final version of manuscript.

DISCLOSURES

801 No conflicts of interest, financial or otherwise, are declared by the authors.