More than a flying syringe: Using functional traits in vector-borne disease research

Lauren J. Cator*[1], Leah R. Johnson[2], Erin A. Mordecai[3], Fadoua El Moustaid[4], Thomas R.C. Smallwood[1], Shannon L. LaDeau[5], Michael A. Johansson[6], Peter J. Hudson[7], Michael Boots[8], Matthew B. Thomas[9], Alison G. Power[10] & Samraat Pawar[1]

[1] Department of Life Sciences, Imperial College London, Silwood Park, Ascot, SL5 7PY, UK

[2] Department of Statistics, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061, USA

[3] Department of Biology, Stanford University, Stanford, CA 94305, USA

[4] Department of Biological Sciences, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061, USA

[5] The Cary Institute of Ecosystem Studies, Millbrook, NY 12545, USA

[6] CDC Dengue Branch and Harvard T.H. Chan School of Public Health, Boston, MA 02115, USA

[7] Huck Institutes of Life Sciences, Pennsylvania State University, University Park, PA 16802, USA

[8] Department of Integrative Biology, University of California Berkeley, Berkeley, CA 94720, USA

[9] Department of Entomology, Pennsylvania State University, University Park, PA 16802, USA

[10] Department of Ecology and Evolution, Cornell University, Ithaca, NY 14853, USA

*Corresponding Author: Department of Life Sciences, Imperial College London, Silwood Park Campus, Ascot SL57PY United Kingdom, Phone: 02075941785, l.cator@imperial.ac.uk

Abstract

Many important endemic and emerging diseases are vector-borne. The functional traits of vectors

affect not just pathogen transmission rates, but also the fitness and population dynamics of these

animals themselves. Increasing empirical evidence suggests that vector traits vary significantly at time

scales relevant to transmission dynamics. Currently, an understanding of how this variation in key

traits impacts transmission is hindered by a lack of both empirical data and theoretical methods for

mechanistically incorporating traits into transmission models. Here, we present a framework for

incorporating both intrinsic and environment-driven variation in vector traits into empirical and

theoretical vector-borne disease research. This framework mechanistically captures the effect of trait

variation on vector fitness, the correlation between vector traits, and how these together determine

transmission dynamics. We illustrate how trait-based vector-borne disease modelling can make novel

predictions, and identify key steps and challenges in the construction, empirical parameterisation and

validation of such models, as well as the organization and prioritization of data collection efforts.

Keywords: Vector-borne Disease Modelling, Traits, Population Dynamics, Transmission, Fitness,

Reproductive number

27 **Contents**

I. INTRODUCTION

Vector-borne diseases (VBDs) remain a serious threat to health and economic welfare worldwide. At least one-third of the human population is thought to be impacted by mosquito-borne diseases alone (WHO, 2018). Brazil loses an estimated 1.3 billion USD in productivity each year to Chagas disease, which is transmitted by triatomine bugs (Lee *et al.*, 2013). VBDs include re-emerging infections, such as dengue (San Martín *et al.*, 2010; Dick *et al.*, 2012), and new threats, such as Chikungunya, Zika, and Lyme disease (Mead, 2015; CDC, 2016; Faria *et al.*, 2016). The impacts of VBDs are not only medical. Many important and emerging diseases of plants (e.g., Tomato yellow leaf curl virus, cassava mosaic virus, and citrus greening virus (Taylor *et al.*, 2016)) and livestock (e.g., bluetongue (Wilson & Mellor, 2009)) are transmitted by vectors.

VBD transmission dynamics are driven by a system of interconnected population abundances that vary over time and, usually, also over space. Evidence from a wealth of empirical and theoretical studies indicates that the behaviour and life history of vectors are key determinants of these VBD dynamics (reviewed below). The importance of vector abundance and life history in transmission dynamics is reflected by the fact that almost all VBD models incorporate vector biting rate, competence, adult survival, and density (Jeger *et al.*, 2004; Reiner *et al.*, 2013). These aspects of vector biology can be described as traits.

A trait is any measurable feature of an individual organism. Specifically, a trait is functional when it directly (e.g., mortality rate/probability) or indirectly (e.g., development time) determines individual fitness (McGill *et al.*, 2006; Gibert *et al.*, 2015) (henceforth we use "functional trait" and "trait" synonymously). In natural populations, trait variation between individuals, and within individuals over time is ubiquitous and has been shown to alter population and community level processes (Imura, Toquenaga, & Fujii, 2003; McGill *et al.*, 2006; Agashe, 2009; Gibert *et al.*, 2015). It is widely accepted that vector traits are also important for transmission and they are likely to be temporally and spatially variable (Smith *et al.*, 2014). While some models incorporate these traits, most do not accommodate trait variation. Mechanistically incorporating trait variation into transmission models should therefore improve the ability of models to predict transmission through time and space.

Trait variation can change transmission dynamics, either by changing vector abundances on

similar time scales as pathogen transmission, or by directly changing transmission potential itself

(e.g., vector biting rate or competence). This is partly due to the fact, that the majority of animal and

plant disease vectors are arthropods, and therefore have significantly shorter generation times (and

thus, operate at faster timescales) than their host (May & Anderson 1979). Furthermore, arthropods

are small ectotherms, which means their abundances vary with fluctuations in environmental

variables. Work from other fields indicates that both the mean and variation of a trait's distribution

can have strong effects on the dynamics of populations (Norberg *et al.*, 2001; Bolnick *et al.*, 2011;

Gibert *et al.*, 2015). Trait variation has now begun to be identified as essential variables that are key

for understanding the state and future trajectories of biodiversity and ecosystems (Kissling *et al.*,

2018). Disease dynamics are strongly nonlinear, which means that variation in traits over time and

across individuals can have compounding effects on transmission (Lloyd-Smith *et al.*, 2005; Martin *et*

al., 2019). Until we incorporate trait variation into VBD transmission models our ability to predict

transmission dynamics at longer temporal and larger spatial scales will remain limited, like any

ecological model that lacks a mechanistic basis (Getz *et al.*, 2018). To incorporate trait variation

robustly, the mechanistic links between biological parameters, and how their effects on vector

population fitness can change population dynamics and transmission rates need to be considered.

There are two related challenges to understanding the parameter space in which trait variation

could affect transmission. First, for most vector species, we lack basic data on traits underlying these

parameters, and how they vary. Researchers are frequently forced to infer parameter values (for

example, using the time it takes a mosquito to produce a clutch of eggs to infer biting rate) or use data

from related species to parameterise models (for example, Johnson *et al.* 2015, Mordecai *et al.* 2013).

Second, while there are many statistical, correlative methods, we lack a theoretical framework to

mechanistically incorporate parameter variation (through trait variation) into transmission models.

Here, we propose a trait-based framework as the way forward on both empirical and

theoretical fronts for making more accurate VBD transmission dynamic predictions. We begin by

reviewing empirical evidence for trait variation in vectors and the degree which it has (or in some

cases has not) been incorporated into previous transmission models. We then describe the trait-based

modelling framework and provide an example of how this approach can produce novel predictions and allows for investigation into the degree to which different traits drive transmission dynamics and underlying fitness effects. Next we outline how this can be operationalized, both in terms of incorporating traits into transmission models, and data collection efforts. Finally, we discuss further challenges in the way of implementing a trait-based approach in VBD research.

II.   TRAIT VARIATION IN VECTORS

Variation in vector traits can be grouped into three primary types: 1) variation across the lifespan of an individual; 2) variation within a population; and 3) environmentally driven variation (Fig. 1).

(1) Variation across the lifespan of an individual

For transmission to occur a vector must survive the time it takes from acquiring the parasite to the parasite becoming infectious (extrinsic incubation period or EIP), which can be a large proportion of the vector life span. Therefore, older vector individuals are; 1) more likely to become infected because they are more likely to have been exposed, 2) more likely to be infectious because they are more likely to have survived EIP, and 3) are more likely to transmit the pathogen onwards because they are more likely to bite subsequent hosts after becoming infectious. Therefore, trait values at the end of life disproportionately contribute to transmission. Vector behaviour and life history may vary over the lifetime of an individual also due to intrinsic processes such as aging. There is evidence for age-specific vector competence (Soliman *et al.*, 1993), immune function (Hillyer *et al.*, 2005; Christensen *et al.*, 2005; Laughton, Fan, & Gerardo, 2014), flight performance (Nayar & Sauerman, 1973) and feeding behaviour (Alto, Lounibos, & Juliano, 2003; Den Otter, Tchicaya, & Schutte, 2008; Bohbot *et al.*, 2013). When multiple life stages of the vector are involved in transmission, transmission efficiency may vary with stage (Caraco *et al.*, 2002; Coletta-Filho *et al.*, 2014). All these time-dependent changes in vector traits could lead to significant variation in the number of infectious vectors and their contact rates with hosts.

(2) Variation within a population

Variation within a population (for example due to intrinsic genetic variation) can lead to

subgroups of the vector population having disproportionate effects on the average population fitness

and transmission potential. There are multiple examples of this kind of within-population variation in

vectors. For example, the nutritional status of vectors can alter both their vector competence and

several behavioural parameters linked to VBD transmission (Takken *et al.*, 2013; Shapiro *et al.*,

2016). Body size varies within populations (Renshaw, Service, & Birley, 1994; De Xue, Edman, &

Scott, 1995; Kindlmann & Dixon, 2003) and can drive significant variation in feeding, assimilation,

and respiration, and therefore development and mortality rates (Brown *et al.*, 2004; Savage, 2004;

Amarasekare & Savage, 2012). Inter-individual variation in age-specific mortality is important for

transmission (Clements & Paterson, 1981; Harrington *et al.*, 2001, 2008; Styer *et al.*, 2007). Recent

evidence from several different systems shows that vectors infected with a variety of pathogens

exhibit altered foraging behaviours (reviewed in (Murdock, Luckhart, & Cator, 2017)). These types of

variation between individuals in key traits such as biting rate, host preference, and longevity

ultimately impact VBD pathogen transmission rate.

(3) Environmentally driven variation

The majority of vectors are small ectotherms, so their behaviour and life history are sensitive to

their environment. Variation due to environmental drivers, may have both short- or long-term effects

on vector traits. At present most of the data on this kind of variation comes from investigations of

temperature. Many studies have measured an effect of variation in environmental temperature on

vector life history (Kersting, Satar, & Uygun, 1999; Bayoh & Lindsay, 2003; Delatte *et al.*, 2009;

Ciota *et al.*, 2014) and competence (Kramer, Hardy, & Presser, 1983; Murral *et al.*, 1996; Dohm,

O'Guinn, & Turell, 2002; Paweska, Venter, & Mellor, 2002; Wittmann, Mellor, & Baylis, 2002).

Other environmental variables, such as humidity, can also directly affect vector life history (Wittmann

et al., 2002; Costa *et al.*, 2010).

(4) Traits are mechanistically linked

Not only do vector traits vary but multiple traits also co-vary because they are mechanistically

linked. For example, mosquitoes infected with bird malaria parasites exhibit reduced fecundity, which

in turn increases longevity (Vézilier *et al.*, 2012). These kinds of trait covariances, often appearing in

the form of life-history trade-offs (Charnov 1993), have implications for both vector population

fitness (and therefore abundance) and transmission rate. Therefore, any theoretical framework that

incorporates vector traits and their variation must also account for the mechanistic relationships

between traits. There is evidence that this type of incorporation of multiple, mechanistically-linked

traits can yield new insights in transmission of VBDs as well as non-VBDs (Mordecai *et al.*, 2013,

2017; Molnár *et al.*, 2017). For example, recent work has incorporated metabolic traits into micro-

and macro-parasite disease transmission (Molnár *et al.*, 2017).

In summary, any theoretical framework for VBD should be able to capture changes in

transmission dynamics emerging from trait variation (including for those not directly related to

transmission), correlations between traits, and the resulting effects on vector population dynamics. We

now consider past approaches towards incorporating traits into mathematical VBD models.

III.   CURRENT   APPROACHES   TO   INCORPORATING   TRAITS   INTO

TRANSMISSION DYNAMICS

(1) Classical compartment models

Classical compartment models focus on the frequency of different (Susceptible, Exposed,

Infected, Recovered) sub-populations of the host and vector, often ignoring fluctuations in absolute

abundances of the two species. For example, the Ross-Macdonald type model for malaria

transmission by a mosquito (Fig. 2, SI Section 2), on which many current VBD models (including

non-mosquito vectors) are based (MacDonald, 1957; May & Anderson, 1979; Jeger *et al.*, 2004;

Smith *et al.*, 2012; Reiner *et al.*, 2013) focuses exclusively on the parameters governing transmission

rate of the pathogen between susceptible and infected vector and host subpopulations, most of which

are mosquito traits. It yields a relatively simple equation for the basic reproduction number of the

disease (R_0)—the number of new infectious cases that would arise from a single infectious case

introduced into a fully susceptible host population—which quantifies its transmission potential or risk

(MacDonald, 1957; Smith *et al.*, 2012) (see SI section 2 for derivation):

$$R_0 = \left(\frac{V a^2 b c \mathrm{e}^{-\mu P}}{H d \mu} \right)^{\frac{1}{2}} \qquad\qquad 1$$

Here, V is vector density, a is per-vector biting rate, b is the proportion of the bites by infective

mosquitoes that produce infection in susceptible humans, c is the proportion of bites by susceptible

mosquitoes on infectious humans that infect mosquitoes (thus, bc is vector competence), μ is adult

vector mortality rate, P is the extrinsic incubation period of the pathogen within the vector, H is host

density, and d is the rate at which infected hosts recover and acquire immunity (Fig 2).

Compartment models do implicitly incorporate traits. The vector (a, b, c, P, μ) and host (d)

parameters used in this approach are all essentially traits that are either directly-measurable properties

of the vector itself (a, μ) or which can be derived from underlying traits of the vector and parasite (b,

c, P). However, in these models, it is assumed that vector (and host) traits do not affect total vector or

host population size and that these traits do not vary across individuals or over time (Fig. 1).

Furthermore, all of the vector's biology and ecology are represented by its adult (often female) biting

rate, survival, extrinsic incubation period, competence, and abundance. All of these traits are assumed

to be independent of each other despite the fact that they are known to trade-off and feedback on each

other with potentially compounding effects on transmission. For example, biting rate (a) is assumed to

have no effect on mortality (μ) or vector population density (V) and does not vary across individuals

or over time. All individual hosts and vectors are modelled as having an identical risk of infection that

remains constant over time and vectors interact exclusively with the target host. This effectively

reduces the vector to a homogeneous vehicle to transmit pathogens– a "flying syringe".

(2) Compartment models with vector population dynamics

Classical compartment models have been extended to incorporate vector population

dynamics, by adding vector life-stage compartments (May & Anderson, 1979; Anderson & May,

1979; Hoshi, Higa, & Chaves, 2014; Johnson *et al.*, 2018; Ng *et al.*, 2018) (Fig. 3). This introduces

additional parameters for the vector's life history, all of which are directly measurable vector traits

(also called demographic rates, e.g., mortality, fecundity), or parameters which can be derived from

underlying traits (e.g. the transition probabilities (p_i's) in Fig. 3 can be derived from stage-specific

survivorship and development time). These models do relax the assumption that vector population

dynamics are constant, but do not incorporate trait variation, mechanistic links between traits, or allow

trait variation to drive vector population dynamics.

(3) Adding in trait variation

Subsequent studies have incorporated trait variation into the compartment as well as

Anderson-May models (Fig. 4). Reiner et al (2013) found that while extensions of the Ross-

MacDonald model to incorporate parasite latency in malaria vectors are commonplace, efforts to

include variation in vector traits such as heterogeneity in host-vector contact or variation other vector

(and host) attributes are rare. Several studies have shown that variation in single traits such as age-

specific vector mortality drives changes in the predicted sensitivity of R_0 to vector control (Styer *et*

al., 2007; Bellan, 2010; Novoseltsev *et al.*, 2012). Incorporating variation in single life history traits

associated with infection (McElhany, Real, & Power, 1995; Koella, 2005; Lefèvre & Thomas, 2008)

or nutrition (Shapiro *et al.*, 2016) has also been found to affect key VBD parameters such as biting

rate. In some cases, components of vector ecology have also been added to Ross-MacDonald type

models, including environmental drivers (Beck-Johnson *et al.*, 2013) or species interactions (Depinay

et al., 2004; Nakazawa, Yamanaka, & Urano, 2012). While these approaches incorporate trait

variation, they don't necessarily link this to vector population dynamics or link between traits.

Other studies have derived transmission parameters as a function of traits without

mechanistically linking traits. For example, Brand, Rock, & Keeling (2016) determined the number of

infectious bites delivered by midges by combining the probability of survival through bluetongue

virus EIP, age specific-biting rate, and mortality. They demonstrate that determining model

parameters with traits can dramatically change the interpretation of R_0 as well as the estimated impact

of vector control for bluetongue. However, while the traits drive a common transmission parameter,

they are not mechanistically linked to one another. In a similar example, Brady *et al.* (2016)

incorporated adult female mosquito blood feeding, egg laying, larval ecology and how these affect

vectorial capacity. They found that this increased the relative importance of vector control methods

relative to those that specifically targeted adults.

Most recently, trait variation has been incorporated directly into compartment models by

incorporating environment driven trait variation. In this case multiple traits are determined by the

same driver. For example, Parham & Michael (2010) derived an equation for mean, asymptotic vector

population size as a function of traits that are then allowed to be functions of environmental

conditions. Mordecai et al. (2013, 2017) built upon this approach to include empirically derived

unimodal thermal responses for life-history and transmission related traits. Brand et al. (2017) took a

similar approach for allowing biting and EIP parameters to depend on temperature, but modelled the

vector population using a statistical approach. All these studies have provided interesting insights into

possible effects of trait variation, but they do not mechanistically link traits. We have not found

examples of studies that have systematically incorporated trait variation, mechanistic linkages

between traits (e.g., co-variation in biting rate and survival), and allowed traits to drive population and

transmission dynamics.

IV.　A TRAIT-BASED FRAMEWORK FOR VBD RESEARCH

Here, we present a general trait-based research framework that unifies these previous

approaches, and can provides a foundation for systematically tackling the challenge of incorporating

trait variation into VBD dynamics. An overview of the framework is given in Fig. 5 (SI section 1,

contains a more detailed framework description). In general, a fully trait-based VBD system's

specification must include:

1. **Transmission compartments for each focal host and vector species:** For example, the SIR

compartments as presented in the Ross-MacDonald type models (e.g., H_S, H_R) and may have

additional host sub-compartments (H_j, where $j \geq 0$) tailored to a particular disease.

2. **Vector life history compartments:** These would include the commonly used infected susceptible

vector sub populations (V_S, V_I), but additionally include the vectors' juvenile life stage

subpopulations, starting at birth (V_0) and followed by immature stages (V_l, where $l > 0$). In adult

stages, we include the potential for additional stages (V_k) leading to infectious adults (V_I).

3. **Species interaction compartments:** These will depend on the VBD system, but typically at least

one consumer-resource interaction would be necessarily incorporated, because all vectors

experience significant mortality from predation at one or more life stages. This would require

compartments for a resource species consumed by the vector (R_1) and predator species that

consume the vector (C_1). In more complex systems, additional consumer and resource

compartments (C_n, R_m) could be added.

4. **Trait Variation**: A suite of trait (e.g., vector mortality, fecundity and biting rates) to parameter

mappings that determine fitness effects, and at least a single type of trait variation, such as with an

environmental factor (dz/dE; e.g., temperature) (Fig. 1, 5). As we explain earlier, under this

framework "fitness" has a more general meaning in its evolutionary connotation: it is the change

in abundance value of any compartment (population) between time steps or spatial locations.

Finally, note that a fitness effect in the transmission component (Fig. 5) changes abundances of

the host's infected sub-population through changes in transmission rate (e.g., through biting rate)

and reflects a change in fitness of the pathogen's (population) fitness. While many features of

individuals and populations are measureable and may be variable, this framework specifies that

these features are only important for transmission when they affect demographic, interaction, or

transmission rates that drive population and transmission dynamics. We incorporate three key

classes of functional traits; life history, interaction, and transmission traits (Table 1).

5. **Mechanistic Links Between Traits**: Traits should be mechanistically linked. This may be

accomplished by explicitly modelling how multiple traits together affect a fitness parameter

through shared bio-mechanical and metabolic constraints (see Section VI.1.a).

We do not show explicit linkages between trait variation, consumer-resource, and life history

sub-compartments in Fig. 5 because these will vary with VBD system. For example, in the case of

most aphid-transmitted diseases, the resource (R) and host (H) are often the same. For other vectors,

such as *Anopheles* mosquitoes, the transmission relevant hosts (H) may make up only a proportion of

the resources (R) that regulate growth and reproduction (e.g. (LoGiudice *et al.*, 2003; Donnelly *et al.*,

2015)).

V.   AN EXAMPLE: A TRAIT-BASED MODEL OF MOSQUITO ABUNDANCE AND

DISEASE TRANSMISSION

We now illustrate how a trait-based approach (Fig. 5) can lead to novel predictions about

vector population dynamics and therefore transmission using the effect of temperature on

transmission as an example. Temperature is a major source of environment-driven trait variation (Fig.

2), and affects variation in both adult and juvenile traits in vectors. To incorporate trait variation into

transmission, we model the effects of temperature-driven life-stage specific trait variation on vector

population density, V, through the population's intrinsic growth rate, r_m. Full details of this worked

example and the model are provided in SI section 3. Briefly, r_m is a function of adult peak fecundity

(b_{pk}), age-related fecundity decline rate (κ), adult mortality rate (μ) and juvenile development time (α)

and juvenile mortality (μ_J). Variation in each of these traits across temperature is characterized by the

thermal performance curve of each trait. By incorporating such environment-driven trait variation into

a vector population abundance model we can derive the transmission dynamics over time (Fig. 6).

We contrast the trait-based approach with a phenomenological one that has been used in

previous studies (Hoshi *et al.*, 2014; Johnson *et al.*, 2018; Ng *et al.*, 2018), where abundance (*V*) is

directly associated with temperature by fitting a time-series model or where abundance is assumed, a

priori, to follow a sinusoidal function (Bacaër & Guernaoui, 2006; Bacaër 2007; Bacaër & Ouifki,

2007; Bacaër & Ait Dads, 2012). This results in a transmission dynamic where R_0 tracks temperature

variation (Fig. 6), potentially with a time-lag. This result contrasts strongly with that from the trait-

based approach that maps traits through parameters to vector population size. Specifically, we observe

key differences in predictions of both abundance dynamics and the knock-on effects on transmission:

the trait-based model predicts that vector populations will emerge earlier in the year and persist later

into the cooler late summer season with a dip in the warmest period of the summer. These differences

in *V* feed through to a longer period of annual transmission with an early and late summer peak. The

trait-based model predicts a longer transmission season than the phenomenological model, and reveals

a decrease in transmission risk in the warmest period. The latter result in particular is surprising given

the general "warmer is better" view, but is consistent with the results of Mordecai et al (2013) which

incorporated variation (SI section 2.2). Interestingly, when metabolic theory was used to

mechanistically model of infection of an endothermic host with an arctic nematode parasite, the

continuous spring-to-fall transmission season morphed into two distinct transmission seasons as

climate warmed was also observed (Molnár, 2013). The similarity in predicted transmission dynamics

across these two very different systems suggests that mechanistically incorporating trait variation can

reveal general constraints on VBD systems — in this case, the effect of temperature on VBD

dynamics through life-history traits. Additionally, a trait-based approach enables us to determine

which traits matter the most for determining thermal sensitivity of transmission dynamics (through

trait sensitivity analyses; see next section).

The example we have developed here also illustrates a key theoretical point we raised at the

start. If vector traits (eqn 1; assumption 2) change at the same or shorter timescales (here, driven by

within-year temperature change) than the rate of pathogen transmission, the classical approach will

fail to capture important aspects of contemporary transmission dynamics (Anderson & May, 1981;

Heesterbeek & Roberts, 1995; Bacaër, 2007). One approach towards relaxing the constant vector

population assumption of classical transmission models has been to build phenomenological models for time varying vector populations, for example by assuming that the vector population oscillates sinusoidally (Heesterbeek & Roberts, 1995; Bacaër, 2007). Although this approach is relatively simple, it cannot quantify or predict how variation in key vector traits or parameters (e.g., *a, b, c, μ*) interact to drive transmission dynamics over time if the relationship between abundance and temperature is non-linear. A trait-based approach provides new mechanistic insights and hypotheses, and predicts non-trivial dynamics that would likely inform public health and control decisions.

(1) The importance of trait sensitivity analyses

In addition to producing novel predictions, a trait-based approach allows investigation of the degree to which different traits drive transmission dynamics and underlying fitness effects. For example, a local trait sensitivity analysis of the population fitness component of the above trait-based model allows us to investigate the relative importance of juvenile versus adult traits in determining effects of temperature on abundance (and therefore transmission) (Fig. 7).

This leads to a key insight: juvenile traits are expected to play a major role in determining vector fitness across temperatures, and therefore abundance, and ultimately transmission. In particular, the trait sensitivity analysis adds further insight to *why* the trait-based approach yields very different predictions for population abundance and R_0 compared to a phenomenological approach. The thermal sensitivity of abundance (V) and the underlying population fitness (r_m) is driven by the variation of temperature-driven variation in larval stage traits. These predictions and insights provide quantitative targets for validation using field data. Sensitivity analyses of transmission measures with respect to traits also allow key traits to be identified, guiding further empirical and theoretical work on the contributions of traits to VBD system dynamics.

VI.   A TRAIT-BASED FRAMEWORK ROADMAP

Our relatively simple worked example in the previous section gives an idea of the steps one would

need to take in order to use a trait-based approach. A full trait-based study would involve making the

following links:

1. Trait→Parameter

2. Trait-Variation→Fitness Variation

3. Fitness→Population Dynamics

4. Population Dynamics→Transmission Dynamics

Research programs may focus on all or a portion of the following four sequential components or

steps. In our worked example, we only attempt to tackle step 2 and 3. Each step requires a relationship

or mapping (→) to be quantified through empirical studies coupled with mathematical modelling. To

be clear, useful progress and insights can be gained at each step. We now explain each of these steps,

followed by an example.

(1) Trait→Parameter

A key component of any trait-based framework is the mapping of trait values onto

mathematical VBD model parameters (Fig. 5). In this framework a parameter represents a process that

would have a fitness effect within the model (e.g., it can change the abundance of one of the model

compartments). For example, mortality rate decreases population abundance (V_S) and fecundity

increases abundance (V_S). The trait/parameter delineation is not exclusive. Mortality rate and

fecundity could also be interpreted as traits because they are directly measurable. In contrast, vectorial

capacity is not a trait as it is a derived measure and cannot be directly measured. The distinction

between "trait" and "parameter" is a matter of context; when a quantity is being used to drive another

parameter it is a trait and when is directly affecting fitness it is a parameter. Deconstruction of

parameters into their underlying traits bounds the parameter's feasible range (parameter space), and

also reveals how different parameters are linked, and therefore interact or trade-off with one another

to impact compartment fitness (Charnov, 1993) and ultimately, transmission dynamics.

Biomechanical and metabolic approaches that link physical and performance traits could be

very useful for tackling this challenge (Brown *et al.*, 2004; McGill *et al.*, 2006; Amarasekare &

Savage, 2012; Pawar, Dell, & Savage, 2015a). These approaches use a common currency (energy) to

map parameters onto traits. This inevitably establishes mechanistic links between traits. For example,

body size drives not just adult vector biting rate, but also its fecundity and mortality rates. Recent

advances in metabolic modelling offer an opportunity to determine encounter rate parameters between

vectors and hosts (Dell, Pawar, & Savage, 2011; Pawar, Dell, & Savage, 2012; Dell, Pawar, &

Savage, 2014; Gilbert *et al.*, 2014; Pawar *et al.*, 2015a; Rizzuto, Carbone, & Pawar, 2018) (see

example below) and even capture within-host parasite dynamics (Kirk *et al.*, 2018). Empirical studies

that validate trait-fitness models for specific vectors and VBD systems are of equal importance.

Ideally, such studies should measure multiple traits within individuals (for example, vector

competence, fecundity, and survival) and how they relate to individual fitness (Ohm *et al.*, 2016).

(a) A Worked Example: Deconstruction of Biting Rate

To illustrate Trait→Parameter mapping, we use recent advances in mechanistic, metabolic

trait-based theory for species interactions (Pawar *et al.*, 2012; Gibert *et al.*, 2015; Pawar *et al.*, 2015a)

to develop a mathematical model that deconstructs biting rate (a parameter in the model) into its

underlying functional traits and show how it is ultimately linked to the vector's adult body size.

Heterogeneity in biting behaviour is considered important for determining transmission dynamics

(Dye & Hasibeder, 1986; Smith *et al.*, 2014). This key transmission parameter in all VBD systems,

that also underpins vector fitness, can be derived from body size (a physical trait) (Fig. 8, SI section

1.2)(De Leo & Dobson, 1996). Adult vector size is a key physical trait as it determines many

functional traits, and also itself responds strongly and predictably to external factors such as

environmental temperature and resource availability (e.g., larval nutrition).

Consider a flying vector such as a mosquito, fly, or midge, seeking larger, warm-blooded

hosts (e.g., a terrestrial mammal, which is also the energy resource for the vector). A general field

encounter rate model for such vector-host interactions in two-dimensional Euclidean space (2*D*) is

(Pawar *et al.*, 2012),

$$a = 2v_r d f(.)H$$

Here, a is biting rate (bites $\times$ time^{-1}) for a given host density H (individuals $\times$ area^{-1}), v_r is relative

velocity between the vector and host, d is the reaction distance (minimum distance at which the vector

detects and reacts to host), and $f(.)$ is the host "risk" function that determines the (biting) functional

response of the vector. That is, $f(.)$ encapsulates the constraints on the vector's handling time (h) (the

post-encounter phase in Fig. 8), which would be determined by its ability to land on and complete the

biting action successfully. Assuming each encounter between vector and host results in one bite, we

can derive biting rate as a function of body size (as physical trait) two key functional traits (vector's

flight velocity and reaction distance) (details in SI section 1.2):

$$a = a_0 m_V^{p_v + 2p_d} k^{p_d} f(.)H \qquad\qquad 2$$

where a_0 is a normalization constant that also includes the effect of environmental temperature, m_V is

the vector's body mass, p_v and p_d are scaling exponents for the size-dependence of vector flight

velocity and reaction distance (respectively) k is the vector-host body mass-ratio (a measure of their

relative size difference). In the case where hosts are relatively rare, $f(.) = 1$, but in the case where hosts

are common (e.g., mosquito populations are often localized near human villages), it can be replaced

with a more complex function that includes the effect of host handling time (physical limits on

vector's biting & ingestion rate; Fig. 2), which itself is likely to be vector size and temperature

dependent (Dell *et al.*, 2011; Pawar *et al.*, 2012).

Breaking down biting rate into its component traits recovers known patterns: that smaller

flying vectors will have a lower mean or maximum biting rate relative to larger ones (e.g., midges vs.

mosquitoes), that biting rates will be lower for walking vectors compared to flying ones (e.g., ticks vs.

mosquitoes), and explains how temperature (through the scaling constants a_0) affects biting rate (e.g.,

(Dell *et al.*, 2011; Mordecai *et al.*, 2013; Dell *et al.*, 2014), providing a mechanistic basis for

determining the effect of temperature on vector fitness as well as vectorial capacity. This

Trait→Parameter mapping approach can be modified to accommodate vectors with different moving

or sensory modes (e.g., ticks vs. mosquitoes), and for the same host density (say, humans).

(2) Trait-Variation→Fitness

A population's fitness is essentially a distribution of fitness values across its individuals.

Therefore, the second key step in a trait-based VBD framework is to use Trait-Parameter mappings to

quantify how variation in functional traits generates variation in vector fitness and transmission. That

is, after identifying key functional traits and building trait–parameter models, the effect of trait

variation on the parameter, and eventually, fitness distribution in each class of compartment needs to

be quantified. The same trait can affect the fitness of more than one compartment. For example,

variation in biting rate or intrinsic mortality would affect both vector population fitness (e.g., r_{max}) and

transmission fitness (e.g., R_0). Incorporating or mapping any of the above three types of trait variation

(Fig. 1) onto vector population or transmission parameter requires the re-definition of the parameters

as functions (e.g., $p(z)$, dz/dt, dz/dE; Fig. 3), which are better constructed mechanistically using Trait-

Parameter mappings. In our worked example, we explicitly derive population fitness using

environment-driven traits.

(3) Fitness→Population Dynamics

The third step is to quantify how trait variation determines vector population abundance or

dynamics over time through effects on fitness. This requires the construction of stage-structured

population dynamic models (Fig. 5). There are two key challenges here. The first is quantification of

the impact of trait variation (Fig. 3) on the vector's stage-structured population dynamics. One

approach for doing so is trait-driver theory, which provides methods inherited from quantitative

genetics to study how trait variation drives abundance dynamics (Norberg *et al.*, 2001; Webb *et al.*,

2010; Enquist *et al.*, 2015). This challenge may be made more difficult because most vectors have

stage-structured life cycles underlying population dynamics, which may have transgenerational effects

(e.g., maternal effects) or carryover effects across life stages (Lorenz & Koella, 2011). Initial progress

can be made by simply measuring trait variation and "plugging" it into population dynamics models

(through Trait-Parameter models) with variation to allow deviations from the standard exponential

assumptions (for example see (Brand *et al.*, 2016)) to determine how sensitive abundance is to trait

variation. In the worked example, we mapped trait-based fitness effects onto vector abundance using

one type of trait variation (environment-driven, specifically temperature). Because body-size is a key

physical trait that also changes with life stage (over time), integral projection models may be another

way to incorporate traits into VBDs (Coulson, 2012; Rees, Childs, & Ellner, 2014; Metcalf *et al.*,

2016).

There is increasing interest in incorporating species interactions into VBD transmission dynamics

(Keesing *et al.*, 2010). This is an area of ongoing investigation not just in VBD research, but in

ecology in general. Species interactions impact life history traits, especially fecundity and mortality,

by shifting them from the baseline, interaction-independent values (Roux *et al.*, 2015). For example,

in consumer-resource interactions, fecundity increases with availability of the vector's resources

(vector-resource or vector-host interaction), and mortality with the vector's consumers (vector-

predator interaction). Tackling this challenge will require mathematical models (and complementary

empirical studies) that tractably include the impacts of species interactions on baseline (or idealized)

life history (demographic) parameters (typically measured in the laboratory), especially fecundity and

mortality. One relatively simple way to make progress in this direction is to re-define the otherwise

idealized life history parameter (e.g., development rate, fecundity, and mortality) functions to include

losses or gains due to species interactions. This approach could circumvent the additional complexity

of explicitly adding consumer-resource dynamics to vector population and transmission dynamics.

(4) Population Dynamics→Transmission Dynamics

The final step is to quantify transmission dynamics using trait-based vector population

dynamics and resulting vector-host interaction rates (Fig. 5). To achieve this, two key challenges on

the theoretical front need to be tackled. First, how trait variation determines the timescale of

fluctuations in vector population sizes and vector-host interaction traits relative to the timescale of

transmission dynamics needs to be modeled and validated. The trait-based approach, by deriving the

timescales of population fluctuations mechanistically, would "naturally" determine whether and when

the separation of the timescales of population and transmission dynamics, implicit in classical

(compartment-type) VBD models, is valid (see example below). Second, the appropriate level of

biological complexity (essentially, number of parameters or traits) needed to capture the effect of trait

variation through population dynamics on transmission dynamics needs to be determined. For

example, trait variation in both juveniles and adults may need to be incorporated simultaneously into

transmission models (Fig 7). In addition to vector traits that affect its population dynamics, models

and data on traits that determine the ability of the vector to transmit a pathogen (e.g., P and bc in eqn.

1) and interact with the host (e.g., a in eqn. 1) will be needed. In many cases, these transmission-

relevant traits will be the same as those determining fitness. For example, both, encounter rate with

host and with the vector's resources (or predators) are determined by body size and velocity (Fig. 8).

Indeed, the host is the primary or sole resource in many vectors (e.g., aphids) which links

transmission parameters directly to the vector's fitness though biting and feeding rate.

VII.   KEY CHALLENGES

The four components for building trait-based models that we have outlined above will each

face four challenges to differing degrees; *data*: how to prioritize experiments and report data;

parameterisation: how to link model components to empirical data; *validation*: how to test the trait –

fitness, trait variation – fitness, population dynamics, and transmission models; *selection*: how to pick

the most parsimonious model.

(1) Data

Experiments to quantify vector traits involve taking individual measures on small insects that

are often logistically difficult to work with due to ethical and biosafety concerns. As we have

described above, theoretical work can help to prioritize which traits we should expect to have the

greatest impact on fitness and transmission. New data collection efforts are currently underway in

several disease systems, but the limited accessibility of data that have already been collected is a

major issue that needs to be addressed. It is critical to get as much information as possible from these

time- and labor-intensive experiments by reporting data at the most disaggregated level possible.

Many studies report point estimates of trait means/demographic rates across populations, across

lifespans, and over environmental gradients rather than presenting distributions or individual values.

Individual-level data are critical for accurately estimating within-population variation in trait values.

Beyond individual studies, consolidating datasets with individual measurements into common formats

would allow for the identification of gaps and coordinated data collection efforts to specifically target

the traits and conditions that are data-poor. We have recently launched a hub for storing and accessing

vector trait data (www.vectorbyte.org) and a platform for coordinating data collection efforts

(www.vectorbite.org). Current data can be used to parameterise mechanistic models, including trait

distributions that accurately represent uncertainty and variation within populations. This would allow

us to identify which traits and types of variation in these traits are not well characterized and may play

an important role in transmission.

(2) Parameterisation

Accurately quantifying the distribution of trait values within populations is a major barrier to

developing and refining Trait → Parameter and Trait Variation → Population Fitness models. First,

better identification of how individual measurements link to modeled quantities needs to be more

consistently addressed. For example, many models assume a single fixed value of the extrinsic

incubation period (EIP – the time it takes for an exposed vector to become infectious). However,

studies may report any number of variations in EIP – such as EIP_{50} (time to 50% of vectors being

infectious) of EIP_{90} (time to 90% infectious) (Ohm *et al.*, 2018). Correctly identifying and

incorporating empirical measurements from the literature to match with the meaning of the trait or

parameters or transformations is necessary. Then the variation and uncertainty in the traits must be

quantified. This involves the use of statistical methods for separating true biological variation from

measurement error, and thereby quantifying uncertainty in estimates of trait values (e.g., the mean or

variance of a trait). Using sophisticated approaches that allow quantification and propagation of

uncertainty (e.g., from Trait→Parameter to population dynamics modelling), such as Bayesian

inference, with or without prior information, will be key here (Clark, 2007; Johnson *et al.*, 2015). In

addition, parameter sensitivity analyses in trait-based models (Fig. 7) are crucial, and can provide

biological insights into the traits driving variation in transmission (Johnson *et al.*, 2015).

(3) Model Validation and Selection

A fundamental goal of trait-based VBD research should be to determine the conditions under

which vector traits drive significant variation in realized transmission rate. This requires validation of

models at each level or compartment of the framework (ideally) with data on the spatial or temporal

distribution of traits/drivers as predictors. In contrast to inference or calibration of a model, validation

is the process of assessing how well a parameterised model can replicate data that was not used for

parameter inference/calibration (i.e., out-of-sample prediction) (Hooten & Hobbs, 2015). For

example, Trait→Parameter models need to be validated with data on population growth rates, the

subsequent population dynamics models need to be validated with data on abundance variation over

space or time, and the transmission models need to be validated using disease incidence data over

space and time. However, when the data available for validation are very different in nature from the

data used for model fitting or model outputs do not correspond to easily measured quantities, it can be

difficult to determine when a model captures sufficient detail to adequately represent reality. Efforts

to develop hierarchical model validation methods for the types of dynamic, trait-based models

described here are ongoing (e.g., (LaDeau *et al.*, 2011; Johnson, Pecquerie, & Nisbet, 2013; Johnson

et al., 2014; Sun, Lee, & Hoeting, 2015). These will include Bayesian methods, which allow

quantification of uncertainty and the incorporation of prior data (Clark, 2007; Hooten & Hobbs,

2015). Not only do the statistical methods for (VBD) dynamical systems need to be refined, but as

they are developed it is important that these methods are made accessible for non-statisticians doing

research in this area. This requires training a new generation of researchers in both the new modeling

techniques (so they can develop models that include details such as behaviour) as well as statistical

techniques appropriate for parameterizing and validating the models as they are developed.

It is inevitable and useful that multiple models will be built to address the same question

within any of the compartments of a trait-based framework (Johnson & Omland, 2004). For example,

there are multiple ways to predict fitness from underlying life history traits (Amarasekare & Coutinho,

2013). Comparing the predictions from multiple models allows us to identify which models are most

should make validation data sets publically available and accessible, and standardized metrics of

goodness-of-fit or similar should be reported for all models against validation sets. Second, publishing

the code used to generate model outputs as a standard practice would facilitate comparing

transmission models of all types—not just those incorporating trait variation. These steps would

enable model comparison and multi-model ensembles to be used for future predictions.

VIII.   WHY NOW?

This is the ideal time to undertake the development of fully trait-based approaches in VBD

research. Recent public health crises have spurred government agencies to support the collection of

large amounts of data on these systems. This, combined with innovations in empirical data collection

and sharing, means that the necessary data for parameterizing and validating trait-based models are

now becoming available. At the same time, the movement towards reproducible research has resulted

in open models and computational approaches that allow more direct comparison of models and easier

expansion of existing methods.

At the same time, the broader field of trait-based research is maturing across ecological

systems, with both the theory and experimental methods growing apace (McGill *et al.*, 2006; Pawar,

Woodward, & Dell, 2015b). This provides VBD trait-based approaches with a solid foundation. There

are many other areas of ecology that are currently striving to mechanistically incorporate trait

variation to understand emergent (especially community and ecosystem level) processes. Methods

developed for VBDs would have utility in solving problems in these other systems, such as predicting changes in ecosystem services like primary production (Díaz *et al.*, 2007; Blanchard *et al.*, 2012).

Building a fully trait-based approach to modeling VBD dynamics is not the "quick and easy path" (Kershner & Lucas, 1980). It is data-hungry and requires extensive efforts to building models that integrate knowledge about processes from the individual to populations and beyond. However, we argue that in comparison to phenomenological approaches (e.g., regressions or other correlative approaches) taking a more mechanistic approach, in general, provides a better way to extrapolate dynamics across time or space (Bayarri *et al.*, 2009). This kind of predictive power is becoming increasingly critical as these pathogens continue to emerge and expand their ranges in the context of a globalized and rapidly changing environment. Beyond that, incorporating traits with this bottom up approach should allow us to better understand *why* we see the patterns of transmission that we do, and potentially inform efforts at control. By explicitly modelling the fitness of a given trait and its effect on population dynamics and fitness, trait-based approaches could be used to incorporate trait evolution into transmission models. For instance, although we have focused on traits that directly affect vector population fitness in idealized conditions, the addition of other traits that are mediated by human intervention, such as insecticide resistance, is also possible within this framework. The evolution of insecticide resistance is arguably the largest challenge to sustainable management of vector-borne diseases. A trait-based approach has the potential to better understand the implication of both current (e.g., chemical pesticides) and future control measures (e.g., genetically altered vectors) that inherently alter traits, while suggesting innovative and nuanced ways to apply control in a way that to anticipates changes driven by the inherent complexities of these systems.

IX.   **CONCLUSION**

1. Mounting evidence shows that vector functional traits vary over lifespan, within populations, and across environmental gradients. This variation in key parameters is expected to alter disease transmission dynamics.

2. Mathematical models of Vector-Borne Disease (VBD) dynamics to date have not mechanistically incorporated trait variation, and are therefore unable to capture changes in transmission dynamics that may emerge from different types and magnitudes of trait variation, the correlations between traits, and resulting vector population dynamics.

3. A fully trait-based VBD approach would involve explicitly modelling trait variation and its effect on transmission, vector life history, species interactions and also allowing feedback between traits. This requires mechanistic Trait-Parameter mapping, and the incorporation of trait variation (and co-variation) into the fitness of vector population and disease transmission compartments.

4. Using a trait-based VBD approach can lead to novel predictions for transmission dynamics, and allows analyses of the relative importance of different traits in transmission dynamics. Incorporating trait variation into VBD transmission models will enhance our ability to predict transmission dynamics at longer temporal and larger spatial scales.

5. Increasing research efforts in the development of trait-based theory and data, combined with increasing investment in vector population surveillance makes this an ideal time for the community to systematically and strategically incorporate trait variation into VBD research.

Acknowledgements

We would like to thank the members of the VectorBiTE RCN (www.vectorbite.org) for their energy and stimulating discussions around these ideas. This work was funded by NIH grant 1R01AI122284-01 and BBSRC grant BB/N013573/1 as part of the joint [NIH-NSF-USDA-BBSRC] Ecology and Evolution of Infectious Diseases program. EAM and LRJ were funded by NSF grant DEB-1518681.

Author Contributions: LJC, SP, LRJ, EM and PJH conceived the study. LJC and SP wrote the manuscript with inputs from MBT, AGP, MB, SLL, MAJ, LRJ, EM, and PJH. SP, LRJ, TS, and FEM developed the mathematical models and worked examples.

References

AGASHE, D. (2009) The stabilizing effect of intraspecific genetic variation on population dynamics in

novel and ancestral habitats. *American Naturalist* **174**, 255–267.

ALTO, B.W., LOUNIBOS, L.P. & JULIANO, S.A. (2003) Age-dependent bloodfeeding of *Aedes aegypti*

and Aedes albopictus on artificial and living hosts. *Journal of the American Mosquito Control*

Association **19**, 347–352.

AMARASEKARE, P. & SAVAGE, V. (2012) A framework for elucidating the temperature dependence of

fitness. *The American Naturalist* **179**, 178–191.

AMARASEKARE, P. & COUTINHO, R.M. (2013) The intrinsic growth rate as a predictor of population

viability under climate warming. *Journal of Animal Ecology* **82**, 1240–1253.

ANDERSON, R.M. & MAY, R.M. (1979) Population biology of infectious diseases: Part I. *Nature* **280**,

361–367.

ANDERSON, R.M. & MAY, R.M. (1981) The population dynamics of microparasites and their

invertebrate hosts. *Philosophical Transactions of the Royal Society of London B* **291**, 451–

524.

BACAËR, N. (2007) Approximation of the basic reproduction number R0 for vector-borne diseases

with a periodic vector population. *Bulletin of Mathematical Biology* **69**, 1067–1091.

BACAËR, N. & AIT DADS, E. (2012) On the biological interpretation of a definition for the parameter

R0 in periodic population models. *Journal of Mathematical Biology* **65**, 601–621.

BACAËR, N. & GUERNAOUI, S. (2006) The epidemic threshold of vector-borne diseases with

seasonality. *Journal of Mathematical Biology* **53**, 421–436.

BACAËR, N. & OUIFKI, R. (2007) Growth rate and basic reproduction number for population models

with a simple periodic factor. *Mathematical Biosciences* **210**, 647–658.

BAYARRI, M., CALDER, E., LUNAGOMEZ, S., PITMAN, E., BERGER, J., DALBEY, K., PATRA, A.,

SPILLER, E. & WOLPERT, R. (2009) Using statistical and computer models to quantify

volcanic hazards. *Technometrics* **51**, 401–413.

BAYOH, M.N. & LINDSAY, S.W. (2003) Effect of temperature on the development of the aquatic

stages of *Anopheles gambiae* sensu stricto (Diptera: Culicidae). *Bulletin of Entomological*

Research **93**, 375–381.

BECK-JOHNSON, L.M., NELSON, W.A., PAAIJMANS, K.P., READ, A.F., THOMAS, M.B. & BJØRNSTAD,

O.N. (2013) The effect of temperature on *Anopheles* mosquito population dynamics and the

potential for malaria transmission. *PLoS ONE* **8**, e79276.

BELLAN, S. (2010) The importance of age dependent mortality and the extrinsic incubation period in

models of mosquito-borne disease transmission and control. *PLoS One* **5**, e10165.

BLANCHARD, J.L., JENNINGS, S., HOLMES, R., HARLE, J., MERINO, G., ALLEN, J.I., HOLT, J., DULVY,

N.K. & BARANGE, M. (2012) Potential consequences of climate change for primary

production and fish production in large marine ecosystems. *Philosophical Transactions of the*

Royal Society of London B **367**, 2979–2989.

BOHBOT, J., DURAND, N., VINYARD, B. & DICKENS, J. (2013) Functional development of the octenol

response in *Aedes aegypti*. *Frontiers in Physiology*.

BOLNICK, D.I., AMARASEKARE, P., ARAÚJO, M.S., BÜRGER, R., LEVINE, J.M., NOVAK, M., RUDOLF,

V.H.W., SCHREIBER, S.J., URBAN, M.C. & VASSEUR, D. (2011) Why intraspecific trait

variation matters in community ecology. *Trends in ecology & evolution* **26**, 183–192.

BRADY, O.J., GODFRAY, H.C.J., TATEM, A.J., GETHING, P.W., COHEN, J.M., MCKENZIE, F.E.,

PERKINS, T.A., REINER, R.C., TUSTING, L.S., SINKA, M.E., MOYES, C.L., ECKHOFF, P.A.,

SCOTT, T.W., LINDSAY, S.W., HAY, S.I., ET AL. (2016) Vectorial capacity and vector control:

reconsidering sensitivity to parameters for malaria elimination. *Transactions of the Royal*

Society of Tropical Medicine and Hygiene **110**, 107–117.

BRAND, S., ROCK, K. & KEELING, M. (2016) The interaction between vector life history and short

vector life in vector-borne disease control. *PLoS computational biology* **12**, e1004837.

BROWN, J.H., GILLOOLY, J.F., ALLEN, A.P., SAVAGE, V.M. & WEST, G.B. (2004) Toward a

metabolic theory of ecology. *Ecology* **85**, 1771–1789.

CARACO, T., GLAVANAKOV, S., CHEN, G., FLAHERTY, J.E., OHSUMI, T.K. & SZYMANSKI, B.K.

742   (2002) Stage-structured infection transmission and a spatial epidemic: a model for Lyme

743   disease. *The American Naturalist* **160**, 348–359.

744   CDC (2016) Lyme disease: data and statistics. www.cdc.gov/lyme/stats/ [accessed 31 May 2016].

745   CHARNOV, E.L. (1993) *Life History Invariants: Some Explorations of Symmetry in Evolutionary*

746   *Ecology*. Oxford University Press.

747   CHRISTENSEN, B.M., LI, J., CHEN, C.-C. & NAPPI, A.J. (2005) Melanization immune responses in

748   mosquito vectors. *Trends in Parasitology* **21**, 192–199.

749   CIOTA, A.T., MATACCHIERO, A.C., KILPATRICK, A.M. & KRAMER, L.D. (2014) The effect of

750   temperature on life history traits of *Culex* mosquitoes. *Journal of medical entomology* **51**, 55–

751   62.

752   CLARK, J. (2007) *Models for Ecological Data: An Introduction*. Princeton University Press, Princeton

753   N.J.

754   CLEMENTS, A.N. & PATERSON, G.D. (1981) The analysis of mortality and survival rates in wild

755   populations of mosquitoes. *Journal of Applied Ecology* **18**, 373–399.

756   COLETTA-FILHO, H.D., DAUGHERTY, M.P., FERREIRA, C. & LOPES, J.R. (2014) Temporal progression

757   of *Candidatus Liberibacter asiaticus* infection in citrus and acquisition efficiency by

758   *Diaphorina citri*. *Phytopathology* **104**, 416–421.

759   COSTA, E.A.P. DE A., SANTOS, E.M. DE M., CORREIA, J.C. & ALBUQUERQUE, C.M.R. DE (2010)

760   Impact of small variations in temperature and humidity on the reproductive activity and

761   survival of *Aedes aegypti* (Diptera, Culicidae). *Revista Brasileira de Entomologia* **54**, 488–

762   493.

763   COULSON, T. (2012) Integral projections models, their construction and use in posing hypotheses in

764   ecology. *Oikos* **121**, 1337–1350.

765   CRUTSINGER, G., COLLINS, M., FORDYCE, J., GOMPERT, Z., NICE, C. & SANDERS, N. (2006) Plant

766   genotypic diversity predicts community structure and governs an ecosystem process. *Science*

767   **313**, 966–968.

DE LEO, G.A. & DOBSON, A.P. (1996) Allometry and simple epidemic models for microparasites. *Nature* **379**, 720–722.

DE XUE, R., EDMAN, J.D. & SCOTT, T.W. (1995) Age and body size effects on blood meal size and multiple blood feeding by *Aedes aegypti* (Diptera: Culicidae). *Journal of Medical Entomology* **32**, 471–474.

DELATTE, H., GIMONNEAU, G., TRIBOIRE, A. & FONTENILLE, D. (2009) Influence of temperature on immature development, survival, longevity, fecundity, and gonotrophic cycles of *Aedes albopictus*, vector of chikungunya and dengue in the Indian Ocean. *Journal of Medical Entomology* **46**, 33–41.

DELL, A.I., PAWAR, S. & SAVAGE, V.M. (2011) Systematic variation in the temperature dependence of physiological and ecological traits. *Proceedings of the National Academy of Sciences* **108**, 10591–10596.

DELL, A.I., PAWAR, S. & SAVAGE, V.M. (2014) Temperature dependence of trophic interactions are driven by asymmetry of species responses and foraging strategy. *Journal of Animal Ecology* **83**, 70–84.

DEN OTTER, C., TCHICAYA, T. & SCHUTTE, M. (2008) Effect of age, sex and hunger on the antennal olfatory sensitivity of tsetse flies. *Physiological Entomology* **16**, 173–182.

DEPINAY, J.M.O., MBOGO, C.M., KILLEEN, G., KNOLS, B., BEIER, J., CARLSON, J., DUSHOFF, J., BILLINGSLEY, P., MWAMBI, H., GITHURE, J. & OTHERS (2004) A simulation model of African *Anopheles ecology* and population dynamics for the analysis of malaria transmission. *Malaria Journal* **3**, 29.

DÍAZ, S., LAVOREL, S., DE BELLO, F., QUÉTIER, F., GRIGULIS, K. & ROBSON, T.M. (2007) Incorporating plant functional diversity effects in ecosystem service assessments. *Proceedings of the National Academy of Sciences* **104**, 20684–20689.

DICK, O.B., SAN MARTÍN, J.L., MONTOYA, R.H., DEL DIEGO, J., ZAMBRANO, B. & DAYAN, G.H. (2012) The history of dengue outbreaks in the Americas. *The American Journal of Tropical Medicine and Hygiene* **87**, 584–593.

DOHM, D.J., O'GUINN, M.L. & TURELL, M.J. (2002) Effect of environmental temperature on the

ability of *Culex pipiens* (Diptera: Culicidae) to transmit West Nile Virus. *Journal of Medical*

Entomology **39**, 221–225.

DONNELLY, B., BERRANG-FORD, L., ROSS, N.A. & MICHEL, P. (2015) A systematic, realist review of

zooprophylaxis for malaria control. *Malaria Journal* **14**, 313.

DYE, C. & HASIBEDER, G. (1986) Population dynamics of mosquito-borne disease: effects of flies

which bite some people more frequently than others. *Transactions of the Royal Society of*

Tropical Medicine and Hygiene **80**, 69–77.

ENQUIST, B.J., NORBERG, J., BONSER, S.P., VIOLLE, C., WEBB, C.T., HENDERSON, A., SLOAT, L.L. &

SAVAGE, V.M. (2015) Scaling from traits to ecosystems: developing a general trait driver

theory via integrating trait-based and metabolic scaling theories. In *Advances in Ecological*

Research pp. 249–318. Elsevier.

FARIA, N.R., AZEVEDO, R. DO S. DA S., KRAEMER, M.U.G., SOUZA, R., CUNHA, M.S., HILL, S.C.,

THÉZÉ, J., BONSALL, M.B., BOWDEN, T.A., RISSANEN, I., ROCCO, I.M., NOGUEIRA, J.S.,

MAEDA, A.Y., VASAMI, F.G. DA S., MACEDO, F.L. DE L., ET AL. (2016) Zika virus in the

Americas: Early epidemiological and genetic findings. *Science* **352**, 345–349.

GETZ, W.M., MARSHALL, C.R., CARLSON, C.J., GIUGGIOLI, L., RYAN, S.J., ROMAÑACH, S.S.,

BOETTIGER, C., CHAMBERLAIN, S.D., LARSEN, L., D'ODORICO, P. & O'SULLIVAN, D. (2018)

Making ecological models adequate. *Ecology Letters* **21**, 153–166.

GIBERT, J.P., DELL, A.I., DELONG, J.P. & PAWAR, S. (2015) Scaling-up trait variation from

individuals to ecosystems. In *Advances in Ecological Research* (ed G.W. AND A.I.D.

SAMRAAT PAWAR), pp. 1–17. Academic Press.

GILBERT, B., TUNNEY, T.D., MCCANN, K.S., DELONG, J.P., VASSEUR, D.A., SAVAGE, V., SHURIN,

J.B., DELL, A.I., BARTON, B.T. & HARLEY, C.D. (2014) A bioenergetic framework for the

temperature dependence of trophic interactions. *Ecology Letters* **17**, 902–914.

HARRINGTON, L.C., BUONACCORSI, J.P., EDMAN, J.D., COSTERO, A., KITTAYAPONG, P., CLARK,

G.G. & SCOTT, T.W. (2001) Analysis of survival of young and old *Aedes aegypti* (Diptera:

Culicidae) from Puerto Rico and Thailand. *Journal of medical entomology* **38**, 537–547.

HARRINGTON, L.C., FRANÇOISEVERMEYLEN, null, JONES, J.J., KITTHAWEE, S., SITHIPRASASNA, R.,

EDMAN, J.D. & SCOTT, T.W. (2008) Age-dependent survival of the dengue vector *Aedes*

aegypti (Diptera: Culicidae) demonstrated by simultaneous release-recapture of different age

cohorts. *Journal of Medical Entomology* **45**, 307–313.

HEESTERBEEK, J. & ROBERTS, M. (1995) Threshold quantities for infectious diseases in periodic

environments. *Journal of biological systems* **3**, 779–787.

HILLYER, J.F., SCHMIDT, S.L., FUCHS, J.F., BOYLE, J.P. & CHRISTENSEN, B.M. (2005) Age-

associated mortality in immune challenged mosquitoes (*Aedes aegypti*) correlates with a

decrease in haemocyte numbers. *Cellular Microbiology* **7**, 39–51.

HOOTEN, M. & HOBBS, N. (2015) A guide to Bayesian model selection for ecologists. *Ecological*

Monographs **85**, 3–28.

HOSHI, T., HIGA, Y. & CHAVES, L.F. (2014) *Uranotaenia novobscura ryukyuana* (Diptera: Culicidae)

population dynamics are denso-dependent and autonomous from weather fluctuations. *Annals*

of the Entomological Society of America **107**, 136–142.

IMURA, D., TOQUENAGA, Y. & FUJII, K. (2003) Genetic variation can promote system persistence in

an experimental host-parasitoid system. *Population Ecology* **45**, 205–212.

JEGER, M.J., HOLT, J., VAN DEN BOSCH, F. & MADDEN, L.V. (2004) Epidemiology of

insect-transmitted plant viruses: modelling disease dynamics and control interventions.

Physiological Entomology **29**, 291–304.

JOHNSON, J. & OMLAND, K. (2004) Model selection in ecology and evolution. *Trends in Ecology &*

Evolution **19**, 101–108.

JOHNSON, L., PECQUERIE, L. & NISBET, R. (2013) Bayesain inference for bioenergetic models.

Ecology **94**, 882–889.

JOHNSON, L., LAFFERTY, K., MCNALLY, E., MORDECAI, E., PAAIJMANS, K., PAWAR, S. & RYAN, S.

(2014) Mapping the distribution of malaria: current approaches and future directions. Pages

191–211 in D. Chen, B. Moulin, and J. Wu, editors. In *Analyzing and modeling spatial and*

temporal dynamics of infectious diseases. John Wiley & Sons, Hoboken, New Jersey, USA.

pp. 191–211. John Wiley & Sons, Hoboken, USA.

JOHNSON, L.R., BEN-HORIN, T., LAFFERTY, K.D., MCNALLY, A., MORDECAI, E., PAAIJMANS, K.P.,

PAWAR, S. & RYAN, S.J. (2015) Understanding uncertainty in temperature effects on vector-

borne disease: a Bayesian approach. *Ecology* **96**, 203–213.

JOHNSON, L.R., GRAMACY, R.B., COHEN, J., MORDECAI, E., MURDOCK, C., ROHR, J., RYAN, S.J.,

STEWART-IBARRA, A.M. & WEIKEL, D. (2018) Phenomenological forecasting of disease

incidence using heteroskedastic Gaussian processes: A dengue case study. *The Annals of*

Applied Statistics **12**, 27–66.

KEESING, F., BELDEN, L.K., DASZAK, P., DOBSON, A., HARVELL, C.D., HOLT, R.D., HUDSON, P.,

JOLLES, A., JONES, K.E. & MITCHELL, C.E. (2010) Impacts of biodiversity on the emergence

and transmission of infectious diseases. *Nature* **468**, 647.

KERSHNER, I. & LUCAS, G. (1980) The Empire strikes back. Twentieth Century-Fox Film

Corporation.

KERSTING, U., SATAR, S. & UYGUN, N. (1999) Effect of temperature on development rate and

fecundity of apterous *Aphis gossypii* Glover (Hom., Aphididae) reared on *Gossypium*

hirsutum L. *Journal of Applied Entomology* **123**, 23–27.

KINDLMANN, P. & DIXON, A.F.G. (2003) Insect predator–prey dynamics and the biological control of

aphids by ladybirds. In *First international symposium on biological control of arthropods.*

USDA Forest Service, USA pp. 118–124.

KIRK, D., JONES, N., PEACOCK, S., PHILLIPS, J., MOLNÁR, P.K., KRKOŠEK, M. & LUIJCKX, P. (2018)

Empirical evidence that metabolic theory describes the temperature dependency of within-

host parasite dynamics. *PLOS Biology* **16**, e2004608.

KISSLING, W.D., WALLS, R., BOWSER, A., JONES, M.O., KATTGE, J., AGOSTI, D., AMENGUAL, J.,

BASSET, A., VAN BODEGOM, P.M. & CORNELISSEN, J.H. (2018) Towards global data

products of Essential Biodiversity Variables on species traits. *Nature ecology & evolution* **2**,

1531–1540.

KOELLA, J.C. (2005) Malaria as a manipulator. *Behavioural processes* **68**, 271-273.

KRAMER, L.D., HARDY, J.L. & PRESSER, S.B. (1983) Effect of temperature of extrinsic incubation on

the vector competence of *Culex tarsalis* for Western Equine Encephalomyelitis Virus. *The*

American Journal of Tropical Medicine and Hygiene **32**, 1130–1139.

LADEAU, S.L., GLASS, G.E., HOBBS, N.T., LATIMER, A. & OSTFELD, R.S. (2011) Data-model fusion

to better understand emerging pathogens and improve infectious disease forecasting.

Ecological Applications: A Publication of the Ecological Society of America **21**, 1443–1460.

LAUGHTON, A.M., FAN, M.H. & GERARDO, N.M. (2014) The combined effects of bacterial symbionts

and aging on life history traits in the pea aphid, *Acyrthosiphon pisum. Applied and*

environmental microbiology **80**, 470–477.

LLOYD-SMITH, J.O., SCHREIBER, S.J., KOPP, P.E. & GETZ, W.M. (2005) Superspreading and the

effect of individual variation on disease emergence. *Nature* **438**, 355–359.

LEE, B.Y., BACON, K.M., BOTTAZZI, M.E. & HOTEZ, P.J. (2013) Global economic burden of Chagas

disease: a computational simulation model. *The Lancet Infectious Diseases* **13**, 342–348.

LEFÈVRE, T. & THOMAS, F. (2008) Behind the scene, something else is pulling the strings:

emphasizing parasitic manipulation in vector-borne diseases. *Infection, genetics and evolution*

8, 504–519.

LLOYD-SMITH, J.O., SCHREIBER, S.J., KOPP, P.E. & GETZ, W.M. (2005) Superspreading and the

effect of individual variation on disease emergence. *Nature* **438**, 355–359.

LOGIUDICE, K., OSTFELD, R.S., SCHMIDT, K.A. & KEESING, F. (2003) The ecology of infectious

disease: effects of host diversity and community composition on Lyme disease risk.

Proceedings of the National Academy of Sciences **100**, 567–571.

LORENZ, L. & KOELLA, J. (2011) Maternal environment shapes the life history and susceptibility to

malaria of *Anopheles gambiae* mosquitoes. *Malaria Journal* **10**, 1–8.

MACDONALD, G. (1957) *The epidemiology and control of malaria*. Oxford University Press, London.

MARTIN, L.B., ADDISON, B., BEAN, A.G., BUCHANAN, K.L., CRINO, O.L., EASTWOOD, J.R., FLIES,

A.S., HAMEDE, R., HILL, G.E. & KLAASSEN, M. (2019) Extreme Competence: Keystone

Hosts of Infections. *Trends in Ecology & Evolution*.

MAY, R.M. & ANDERSON, R.M. (1979) Population biology of infectious diseases: Part II. *Nature* **280**,

455–461.

MCELHANY, P., REAL, L.A. & POWER, A.G. (1995) Vector preference and disease dynamics: a study

of barley yellow dwarf virus. *Ecology*, 444–457.

MCGILL, B.J., ENQUIST, B.J., WEIHER, E. & WESTOBY, M. (2006) Rebuilding community ecology

from functional traits. *Trends in Ecology & Evolution* **21**, 178–185.

MEAD, P.S. (2015) Epidemiology of Lyme disease. *Infectious Disease Clinics of North America* **29**,

187–210.

METCALF, C.J.E., GRAHAM, A.L., MARTINEZ-BAKKER, M. & CHILDS, D.Z. (2016) Opportunities and

challenges of Integral Projection Models for modelling host–parasite dynamics. *Journal of*

Animal Ecology **85**, 343–355.

MOLNÁR, P. (2013) Metabolic approaches to understanding climate change impacts on seasonal host-

macroparasite dynamics. *Ecology Letters* **16**, 9–21.

MOLNÁR, P.K., SCKRABULIS, J.P., ALTMAN, K.A. & RAFFEL, T.R. (2017) Thermal performance

curves and the metabolic theory of ecology—a practical guide to models and experiments for

parasitologists. *Journal of Parasitology* **103**, 423–439.

MORDECAI, E.A., COHEN, J.M., EVANS, M.V., GUDAPATI, P., JOHNSON, L.R., LIPPI, C.A.,

MIAZGOWICZ, K., MURDOCK, C.C., ROHR, J.R., RYAN, S.J., SAVAGE, V., SHOCKET, M.S.,

STEWART IBARRA, A., THOMAS, M.B. & WEIKEL, D.P. (2017) Detecting the impact of

temperature on transmission of Zika, dengue, and chikungunya using mechanistic models.

PLoS neglected tropical diseases **11**, e0005568.

MORDECAI, E.A., PAAIJMANS, K.P., JOHNSON, L.R., BALZER, C., BEN-HORIN, T., DE MOOR, E.,

MCNALLY, A., PAWAR, S., RYAN, S.J., SMITH, T.C. & LAFFERTY, K.D. (2013) Optimal

temperature for malaria transmission is dramatically lower than previously predicted. *Ecology*

Letters **16**, 22–30.

MURDOCK, C.C., LUCKHART, S. & CATOR, L.J. (2017) Immunity, host physiology, and behaviour in

infected vectors. *Current Opinion in Insect Science* **20**, 28–33.

MURRAL, D.J., NAULT, L.R., HOY, C.W., MADDEN, L.V. & MILLER, S.A. (1996) Effects of

temperature and vector age on transmission of two Ohio strains of aster yellows phytoplasma

by the aster leafhopper (Homoptera: Cicadellidae). *Journal of Economic Entomology* **89**,

1223–1232.

NAKAZAWA, T., YAMANAKA, T. & URANO, S. (2012) Model analysis for plant disease dynamics co-

mediated by herbivory and herbivore-borne phytopathogens. *Biology Letters* **8**, 685–688.

NAYAR, J.K. & SAUERMAN, D.M. (1973) A comparative study of flight performance and fuel

utilization as a function of age in females of Florida mosquitoes. *Journal of insect physiology*

19, 1977–1988.

NG, K.C., CHAVES, L.F., TSAI, K.H. & CHUANG, T.W. (2018) Increased adult *Aedes aegypti* and

Culex quinquefasciatus (Diptera: Culicidae) abundance in a dengue transmission hotspot,

compared to a coldspot, within Kaohsiung City, Taiwan. *Insects* **9, 1-16**.

NORBERG, J., SWANEY, D.P., DUSHOFF, J., LIN, J., CASAGRANDI, R. & LEVIN, S. (2001) Phenotyptic

diversity and ecosystem functioning in chaning environments: a theoretical framework. *PNAS*

98, 11376–11381.

NOVOSELTSEV, V.N., MICHALSKI, A.I., NOVOSELTSEVA, J.A., YASHIN, A.I., CAREY, J.R. & ELLIS,

A.M. (2012) An age-structured extension to the vectorial capacity model. *PLOS ONE* **7**,

e39479.

OHM, J.R., BALDINI, F., BARREAUX, P., LEFEVRE, T., LYNCH, P.A., SUH, E., WHITEHEAD, S.A. &

THOMAS, M.B. (2018) Rethinking the extrinsic incubation period of malaria parasites.

Parasites & Vectors **11**, 178.

OHM, J.R., TEEPLE, J., NELSON, W.A., THOMAS, M.B., READ, A.F. & CATOR, L.J. (2016) Fitness

consequences of altered feeding behaviour in immune-challenged mosquitoes. *Parasites &*

Vectors **9**, 113.

PARHAM, P.E. & MICHAEL, E. (2010) Modeling the effects of weather and climate change on malaria

transmission. *Environmental Health Perspectives* **118**, 620–626.

957 PAWAR, S., DELL, A.I. & SAVAGE, V. (2015a) From Metabolic Constraints on Individuals to the
958  Dynamics of Ecosystems. *Aquatic Functional Biodiversity: An Ecological and Evolutionary*
959  *Perspective*, 3–36.

960 PAWAR, S., DELL, A.I. & SAVAGE, V.M. (2012) Dimensionality of consumer search space drives
961  trophic interaction strengths. *Nature* **486**, 485.

962 PAWAR, S., WOODWARD, G. & DELL, A. (2015b) *Trait-Based Ecology- from structure to function*, 1st
963  edition. Elsevier Academic Press, Oxford, UK.

964 PAWESKA, J.T., VENTER, G.J. & MELLOR, P.S. (2002) Vector competence of South African
965  *Culicoides* species for bluetongue virus serotype 1 (BTV-1) with special reference to the
966  effect of temperature on the rate of virus replication in *C. imicola* and *C. bolitinos*. *Medical*
967  *and Veterinary Entomology* **16**, 10–21.

968 REINER, R.J., PERKINS, T., BARKER, C., NIU, T., CHAVES, L., ELLIS, A., GEORGE, D., LE MANACH, A.,
969  PULLIAM, J., BISANZIO, D., BUCKEE, C., CHIYAKA, C., CUMMINGS, D., GARCIA, A., GATTON,
970  M., ET AL. (2013) A systemic review of mathematical models of mosquito-bourne pathogen
971  transmission: 1970:2010. *J R Soc Interface* **10**, 20120921.

972 RENSHAW, M., SERVICE, M.W. & BIRLEY, M.H. (1994) Size variation and reproductive success in the
973  mosquito *Aedes cantans*. *Medical and Veterinary Entomology* **8**, 179–186.

974 RIZZUTO, M., CARBONE, C. & PAWAR, S. (2018) Foraging constraints reverse the scaling of activity
975  time in carnivores. *Nature ecology & evolution* **2**, 247.

976 ROUX, O., VANTAUX, A., ROCHE, B., YAMEOGO, K.B., DABIRÉ, K.R., DIABATÉ, A., SIMARD, F. &
977  LEFÈVRE, T. (2015) Evidence for carry-over effects of predator exposure on pathogen
978  transmission potential. *Proceedings of the Royal Society B* **282**, 20152430.

979 SAN MARTÍN, J.L., BRATHWAITE, O., ZAMBRANO, B., SOLÓRZANO, J.O., BOUCKENOOGHE, A.,
980  DAYAN, G.H. & GUZMÁN, M.G. (2010) The epidemiology of dengue in the Americas over
981  the last three decades: a worrisome reality. *The American Journal of Tropical Medicine and*
982  *Hygiene* **82**, 128–135.

SAVAGE, V.M. (2004) Improved approximations to scaling relationships for species, populations, and

ecosystems across latitudinal and elevational gradients. *Journal of Theoretical Biology* **227**,

525–534.

SHAPIRO, L.L.M., MURDOCK, C.C., JACOBS, G.R., THOMAS, R.J. & THOMAS, M.B. (2016) Larval

food quantity affects the capacity of adult mosquitoes to transmit human malaria. *Proceedings*

of the Royal Society B **283**, 20160298.

SMITH, D.L., BATTLE, K.E., HAY, S.I., BARKER, C.M., SCOTT, T.W. & MCKENZIE, F.E. (2012) Ross,

Macdonald, and a theory for the dynamics and control of mosquito-transmitted pathogens.

PLoS Pathog **8**, e1002588.

SMITH, D.L., PERKINS, T.A., REINER, R.C., BARKER, C.M., NIU, T., CHAVES, L.F., ELLIS, A.M.,

GEORGE, D.B., MENACH, A.L., PULLIAM, J.R.C., BISANZIO, D., BUCKEE, C., CHIYAKA, C.,

CUMMINGS, D.A.T., GARCIA, A.J., ET AL. (2014) Recasting the theory of mosquito-borne

pathogen transmission dynamics and control. *Transactions of The Royal Society of Tropical*

Medicine and Hygiene **108**, 185–197.

SOLIMAN, B., ABO GHALIA, A., SHOUKRY, A. & MERDAN, A. (1993) Mosquito age as a factor

influencing the transmission of *Wuchereria bancrofti*. *Journal of the Egyptian Society of*

Parasitology **23**, 717–722.

STYER, L.M., CAREY, J.R., WANG, J.-L. & SCOTT, T.W. (2007) Mosquitoes do senesce: departure

from the paradigm of constant mortality. *The American Journal of Tropical Medicine and*

Hygiene **76**, 111–117.

SUN, L., LEE, C. & HOETING, J.A. (2015) Parameter inference and model selection in deterministic

and stochastic models via approximate Bayesian computation: modelling a wildlife epidemic.

Environmentrics **26**, 451-462.

TAKKEN, W., SMALLEGANGE, R.C., VIGNEAU, A.J., JOHNSTON, V., BROWN, M., MORDUE-LUNTZ, A.

& BILLINGSLEY, P.F. (2013) Larval nutrition differentially affects adult fitness and

Plasmodium development in the malaria vectors *Anopheles gambiae* and *Anopheles stephensi*.

Parasites & Vectors **6**, 345.

TAYLOR, R., MORDECAI, E., GILLIGAN, C., ROHR, J. & JOHNSON, L. (2016) Mathematical models are

a powerful method to understand and control the spread of Huanglongbing. *PeerJ* **4**, e2642.

VÉZILIER, J., NICOT, A., GANDON, S. & RIVERO, A. (2012) *Plasmodium* infection decreases fecundity

and increases survival of mosquitoes. *Proceedings of the Royal Society B* **279**, 4033–4041.

WEBB, C.T., HOETING, J.A., AMES, G.M., PYNE, M.I. & LEROY POFF, N. (2010) A structured and

dynamic framework to advance traits-based theory and prediction in ecology. *Ecology Letters*

13, 267–283.

WILSON, A.J. & MELLOR, P.S. (2009) Bluetongue in Europe: past, present and future. *Philosophical*

Transactions of the Royal Society B: Biological Sciences **364**, 2669–2681.

WITTMANN, E.J., MELLOR, P.S. & BAYLIS, M. (2002) Effect of temperature on the transmission of

orbiviruses by the biting midge, *Culicoides sonorensis*. *Medical and Veterinary Entomology*

16, 147–156.

WORLD HEALTH ORGANIZATION (2017) Vector-borne diseases factsheet.

www.who.int/mediacentre/factsheets/fs387/en/ [accessed 14 April 2017].

1024

 FIGURE LEGENDS

 Figure 1. Types of trait variation found in all VBD systems. A: Across-individuals: for example,

 variation in a trait (z) within a population at a fixed point in time. The distribution (p(z) around the

 mean ($\bar{z}$), represents amount individual variation not measurement error. For example, we could

 measure the probability of biting of individuals in a population at a particular age (based on data

 reviewed in Murdock et al. 2017) . **B**: Within-individual, over time: for example, biting probability

 may vary over the vector's lifespan (based on data reviewed in Murdock et al. 2017). Such variation is

 quantifiable as a continuous time-dependent function dz/dt, where dz is the (differential) change in

 trait variation change with time (dt). **C:** Environment-driven: For example, biting rate varies

 unimodally with temperature (based on Mordecai et al. 2013). Such variation is quantifiable as a

 continuous environment state-dependent function, dz/dE. **D:** Combined variation: The three types of

 trait variation may appear in combination. For example, across-individual trait variation may change

 over time. The upper line represents change in mean and variance of the trait, while the lower line

 represents change only in the mean. We use derivatives for quantifying over-time, with-environment

 and combined types of trait variation to represent the idealized scenario that the trait value function

 varies continuously and smoothly with respect to another variable. In reality, it may not always be

 possible to express these as smooth functions for empirical reasons.

 Figure 2. **Traits in classical compartment models.** As an illustration, we show here a Ross-

 Macdonald type model for malaria transmission by a mosquito vector. Most of the vector parameters

 in this model — a, b, c, P, μ — are directly measurable vector traits (a, μ), or can be derived from

 underlying vector and parasite traits (b, c, P). The corresponding equations and parameter definitions

 are shown in SI section 2. H is the total host density which is made up of susceptible (H_S), Infected

 (H_I) and Recovered (H_S) sub-groups. Total vector density (V) is made up for Infected (V_I), Exposed

 (V_E) and Susceptible (V_S) sub-groups.

 Figure 3. Traits in the classical compartment model with vector population dynamics included

 (Anderson-May framework). In addition to the host and vector transmission compartments, the

Anderson-May framework includes the vector life-history compartments. As an illustration, we

extend the classical Ross-MacDonald type models model for malaria (Fig. 2) to include mosquito life

history. This introduces additional parameters (vector life stage-specific μ's, fecundity f, and stage

transition probabilities p_i's), all of which are directly measurable vector traits (the μ's, f), or which

can be derived from underlying traits (the p_i's). The corresponding equations and parameter

definitions are shown in SI section 2.

1059

1060   **Figure 4. Trait variation in compartment models.** In some cases traits and trait variation are used

1061   to drive specific components of the classic compartment model (Fig. 2) directly. In some cases traits

1062   are individually driven by a common driver (in most examples temperature). Traits are still not related

1063   to each other mechanistically, and do not drive the transmission process dynamically through changes

1064   in population abundances (e.g., as would happen in the Anderson-May model framework).

1065

Figure 5. The trait-based framework for vector-borne disease systems. For illustration, we have

used a mosquito-borne disease, but this framework can be applied to any VBD with distinct stage or

age classes (further details in SI). Arrows between panels represent parameters (potentially with

underlying traits) that determine population or transmission dynamics through (absolute) fitness

effects. *Transmission dynamics compartments*: Number of Susceptible (S), Infected (I), and

Recovered (R) hosts (H_S, H_I, H_R, respectively; additional compartments, H_j, can be added) and

number of Susceptible, Infected, and Exposed (E) vectors (V_S, V_I , V_E respectively). *Vector population*

dynamics compartments: number of vector individuals at egg, larval pupal and adult stages (V_0, V_1, V_2,

V_S respectively; additional compartments V_k can be added); *Species interaction compartments*:

Abundance of a single resource species (R_1) that is the primary energy source of the vector population

(may actually be the host itself, so $R_1 = H_S$), and a single consumer species (C_1) that is the primary

source of mortality for the vector population; *Trait variation*: A suite of trait to parameter mappings

that determine fitness (e.g., vector mortality, fecundity and biting rates), and a single type of trait

variation, such as variation with an environmental factor (dz/dE; e.g., temperature; see Fig. 3). For

developing a mathematical model of such a system, the most common tool would be ordinary

differential equation (ODE) systems, as illustrated in SI Section 1.

1082

1083 **Figure 6. An example trait-based model for malaria transmission.** We illustrate here the contrast

1084 in models and resulting dynamics produced from phenomenological vs. trait-based approaches. Both

1085 models cover a time scale of one year and seek to predict the fluctuation in transmission risk or rate

1086 (R_0) during that period. Full details of both models can be found in SI section 3.

1087

Figure 7. An example trait sensitivity analysis. Analyzing the influence of individual fitness traits

on the thermal sensitivity of population fitness of mosquito vectors reveals that juvenile traits have a

stronger influence than adult traits in shaping population fitness (r_m) across temperatures- (b_{pk}), age-

related fecundity decline rate (κ), adult mortality rate (μ) and juvenile development time (α) and

juvenile mortality (μ_J). The vertical dashed line marks the thermal optimum of fitness. The sensitivity

of r_m to the thermal dependence can be assessed by deviation of dr_m/dT from zero. When dr_m/dT is

positive this means that population fitness increases at this temperature. **A.** The sensitivity of r_m to the

thermal dependence of each parameter can be assessed by deviation of each evaluated partial

derivative with respect to each parameter (the differently coloured lines) from zero and compared to

the full model (black).   **B.** Traits can be combined by life stage, with development rate and juvenile

mortality categorised as juvenile traits and adult mortality, fecundity, and rate of loss of fecundity as

adult traits. This reveals the relative contribution of the thermal performance curve of each life stage

to the thermal performance curve of r_m, again with greater deviation from zero indicating a greater

contribution. Full details of the trait sensitivity analysis are in SI Section 4.

1102

Figure 8. Illustration of a Trait-Parameter mapping. We develop a mechanistic, trait-based model

for biting rate (a key transmission parameter; eqn. 1) of vectors by deconstructing it into component

functional traits (shown above). Biting rate (a) can be decomposed into relative velocity (v_r), distance

between vectors and hosts (d), risk ($f(.)$) and host density (H). These parameters can be further

decomposed so that f(.) is determined by handling time (h) and the mass-specific velocities of the

vector and host (V_V, V_R) as in eq. 2. The model derivation is in SI section 1.2.

1109

$\square \rightarrow$ Population or sub-population (compartment); $\vdots\vdots \rightarrow$ Optional compartments; H_S, V_I, V_0, R_1 etc.: Abundance; $\vdash \rightarrow$ Fitness (abundance) loss from compartment; $\wedge \rightarrow$ Fitness effect between compartments; $z \rightarrow$ functional trait

A. Across-individuals

Biting probability

$\bar{z}$

$p(z)$

Trait value (z)

B. Over time

Biting probability

$\frac{d\bar{z}}{dt}$

Time

C. With environment

Biting rate

$\frac{d\bar{z}}{dE}$

Environmental value

D. Combined

Trait value (z)

$\frac{dp(z)}{dt}$

z

Time