Partial-resistance against aphids in wild barley involves phloem and

mesophyll-based defences

Authors:

1: Daniel J Leybourne [1,2,3]

2: Tracy A Valentine [3]

3: Jean AH Robertson [4]

4: Estefania Pérez-Fernández [4]

5: Angela M Main [4]

6: Alison J Karley [3]

7: Jorunn IB Bos [1,2*]

Affiliations:

[1] Division of Plant Sciences, School of Life Sciences, University of Dundee, Dundee, DD2 5DA

[2] Cell and Molecular Sciences, the James Hutton Institute, Invergowrie, Dundee, DD2 5DA,

[3] Ecological Sciences, the James Hutton Institute, Invergowrie, Dundee, DD2 5DA

[4] Environmental and Biochemical Sciences, the James Hutton Institute, Cragiebuckler,

Aberdeen, AB15 8QH

[*] Corresponding author

Tables:

Partial-resistance against aphids in wild barley involves phloem- and mesophyll-based defences

Highlight:

Partial-resistance against aphids in wild barley is based in the mesophyll and vascular tissue and is potentially associated with higher basal defence gene expression and altered phloem amino acid composition.

Abstract:

Aphids, including the bird cherry-oat aphid (*Rhopalosiphum padi*), are significant agricultural pests. Aphid populations are typically controlled using insecticides, but there is increasing demand for more sustainable pest management practices. The wild relative of barley, *Hordeum spontaneum* 5 (Hsp5) has been described as partially-resistant to *R. padi.* Partial-resistance is proposed to involve higher thionin and lipoxygenase gene expression. However, the underlying mechanistic processes are unknown. In this study we compared Hsp5 with a susceptible cultivar of barley (Concerto) to test the extent to which partial-resistance affects aphid fitness. We used the electrical penetration graph technique to monitor *R. padi* feeding patterns to elucidate the tissue location of partial-resistance factors alongside molecular and biochemical analyses to identify potential mechanisms. We show that partial-resistance in Hsp5 extends to three aphid species and is mediated by phloem/mesophyll-based factors, leading to a three-fold increase in the time aphids take to establish sustained phloem ingestion. Partial-resistance likely involves elevated expression of defence and phytohormone genes alongside altered phloem amino acid composition. Further work is required to establish the function of these traits, however this study highlights plant tissues which are important in conferring broad-spectrum partial-resistance against aphids in barley.

Keywords

Amino Acid Composition, Defence gene expression, Electrical Penetration Graph (EPG), *Hordeum spontaneum,* Partial-resistance, *Rhopalosiphum padi, Sitobion avenae*, *Utamphorophora humboldti*

Abbreviations

ABA: Abscisic Acid

Ala: Alanine

Arg: Arginine

Asn: Asparagine

Asp: Aspartic Acid

Concerto: *Hordeum vulgare* cv. Concerto

EPG: Electrical penetration graph

ET: Ethylene

FTIR: Fourier Transform Infrared

Gln: Glutamine

Glu: Glutamic Acid

Gly: Glycine

His: Histidine

Hsp5: *Hordeum spontaneum* 5

Ile: Isoleucine

JA: Jasmonic Acid

Leu: Leucine

Lys: Lysine:

Met: Methionine

Phe: Phenylalanine

r_m: the intrinsic rate of population increase

SA: Salicylic Acid

Ser: Serine

Thr: Threonine

Trp: Tryptophan

Tyr: Tyrosine

Val: Valine

Introduction

Barley, *Hordeum vulgare,* is the fourth most agriculturally important cereal by production quantity (Newton et al., 2011), with 65% of barley yield being used as feed for livestock, 30% in malting and distilling processes, and 5% contributing towards other uses (Newman and Newman, 2006). By 2050 it is estimated that crop yields will need to increase by 60% in order to meet the demands of global population growth (Alexandratos and Bruinsma, 2012). One way to help achieve this is to reduce yield losses to abiotic and biotic stress factors. Aphids such as the bird cherry-oat aphid, *Rhopalosiphum padi*, and the English grain aphid, *Sitobion avenae*, are major pests of cereals. Aphids feed directly from plant phloem using specialised mouthparts known as stylets (Auclair, 1963). Aphids cause direct plant damage by feeding and ingesting phloem sap and act as vectors of economically important plant viruses. Many cereal aphids are vectors for viruses such as *Barley Yellow Dwarf Virus* (*BYDV*), which are transmitted into plant tissues during stylet probing (Powell, 2005). Disease resulting from virus infection can cause yield losses of up to 30% in barley and up to 80% in wheat (Smith and Sward, 1982; Perry et al., 2000; Murray and Brennan, 2010).

Aphids are primarily controlled through the application of insecticides, with the amount of insecticide used per annum increasing (Jess et al., 2018). However the use of these has led to the emergence of insecticide-resistant aphid populations (Chen et al., 2007; Foster et al., 2014), caused negative impacts on non-target organism (Unal and Jepson, 1991; James et al., 2016), and potentially resulted in an increase in *BYDV* prevalence (Dewar and Foster, 2017), leading to legislative restrictions on insecticide use and promotion of integrated pest management (Directive, 2009/128/EC). As a consequence, there is a growing requirement for alternative pest management solutions which are more sustainable, and improving crop resistance to aphids is one avenue which could be explored to achieve this.

To date, only partial-resistance against aphids has been reported for cereal crops. Plant resistance traits which are effective against insect pests can be allocated to one of three categories: 1) chemical deterrence of insect settling and feeding; 2) physical barriers to insect feeding; and 3) reduction in palatability (Mitchell et al., 2016). Partial-resistance against aphids has been associated with each of these categories (respectively, Gibson and Pickett (1983), Tsumuki et al. (1989) and Greenslade et al. (2016). Inherent molecular defences also

play a key role in conferring resistance against aphids (Delp et al., 2009; Guan et al., 2015;

Escudero-Martinez et al., 2017). Molecular defences shown to be involved in aphid resistance

in *Hordeum* spp. include increased expression of thionin (antimicrobial peptides) genes (Delp

et al., 2009; Mehrabi et al., 2014; Escudero-Martinez et al., 2017), increased chitinase and β-

1,3-glucanase activity (Forslund et al., 2000), and the presence of plant secondary metabolites

(Gianoli and Niemeyer, 1998). Plant phythormone signalling pathways, including Abscisic Acid

(ABA), Salicylic Acid (SA), Jasmonic Acid (JA) and Ethylene (ET) signalling processes, mediate

co-ordinated molecular responses to herbivory via the regulation of defence signalling genes

and the biosynthesis of defensive allelochemicals (Smith and Boyko, 2007; Bari and Jones,

2009; Morkunas et al., 2011; Foyer et al., 2016); higher constitutive expression of

phytohormone signalling genes can lead to improved resistance against aphids in cereals

(Losvik et al., 2017).

Constitutive and induced plant defences are frequently lost during plant breeding and

domestication (Moreira et al., 2018). However, wild relatives of modern crops possess traits

which confer improved resistance against herbivorous pests (Dempewolf et al., 2014).

Screening crop wild relatives for (partial-)resistance against aphids provides an opportunity

to identify potentially beneficial traits (Zhang et al., 2017) which can be introduced into

agricultural cultivars to improve resistance (Xu et al., 2015; Li et al., 2018). Screening of wild

relatives in wheat (Xu et al., 2015; Aradottir et al., 2017), barley (Nieto-Lopez and Blake, 1994;

Åhman et al., 2000) and maize (Maag et al., 2015) has resulted in successful identification of

partial-resistance traits in wild relatives of these species (Tsumuki et al., 1989; Barria et al.,

1992; Gianoli and Niemeyer, 1998; Ahmad et al., 2011; Greenslade et al., 2016;

Chandrasekhar et al., 2018; Li et al., 2018).

Partial-resistance against *R. padi* was identified in the barley wild relative *H. spontaneum* 5

(Hsp5) (Åhman et al., 2000; Delp et al., 2009). Although Hsp5 features higher tissue

concentrations of indole alkaloid gramine, a plant defensive compound, the presence of

gramine did not correlate with increased partial-resistance (Åhman et al., 2000). Instead,

partial-resistance in Hsp5 is thought to involve increased expression of plant defence genes

(Delp et al., 2009; Mehrabi et al., 2014), including higher constitutive expression of thionin

genes (Delp et al., 2009) and higher expression of a proteinase inhibitor (Mehrabi et al., 2014).

Initial attempts have been made to characterise the molecular processes contributing to

partial-resistance in Hsp5 (Delp et al., 2009), although the underlying mechanism(s) of
resistance and the tissue location of these resistance factors remains to be elucidated.

The primary aim of this study was to characterise the plant traits and mechanisms
contributing to partial-resistance in Hsp5 by comparison with a susceptible commercial barley
cultivar, Concerto, and explore whether this partial-resistance is effective against several
aphid species, including two agricultural pests (*R. padi* and *Sitobion avenae*) and an invasive
species (*Utamphorophora humboldti*). The factors contributing towards partial-resistance in
Hsp5, and their tissue location, were elucidated using the electrical penetration graph (EPG)
technique to monitor *R. padi* feeding processes. Subsequently, we assessed differences
between the two barley species in leaf surface architecture, defence gene expression, and
phloem amino acid composition to gain insight into the potential mechanisms underlying
partial resistance in Hsp5.

Materials and methods

Plant growth and insect rearing conditions

Hordeum vulgare Linnaeus cv. Concerto (Concerto) and *H. spontaneum* 5 Linnaeus (Hsp5)
seeds were surface-sterilised by washing in 2% (v/v) hypochlorite and rinsing with d.H_2O.
Seeds were kept moist in the dark: Hsp5 seeds were incubated at $4^{o}C$ for 14 days and Concerto
seeds were kept at room temperature for 48h. Germinated seedlings were planted into a
bulrush compost mix (Bulrush, Northern Ireland) and grown under glasshouse conditions with
a 16:8h day:night length and a $20:15^{o}C$ day:night temperature. Plants were grown until the
first true leaf stage (1.1 – 1.2 on the Zadoks et al. (1974) growth staging key).

Asexual laboratory cultures of the bird cherry-oat aphid, *Rhopalosiphum padi* (Linnaeus)
(genotype B; Leybourne et al. (2018))*,* the English grain aphid, *Sitobion avenae* Fabricius and
the American grass leaf aphid, *Utamphorophora humboldti* Essig were established from
individual apterous adults collected from Dundee, UK. Molecular barcoding of the
cytochrome oxidase subunit I gene (Folmer et al., 1994) was used to confirm identity of aphid
species. Aphid cultures were reared on one week old barley seedlings (cv. Optic) contained in
ventilated cups at $20^{o}C$ and 16:8h (L:D).

Insect fitness measurements

The performance of all aphid species was assessed in glasshouse conditions (described above) on Hsp5 and Concerto (*n* = 12) in a randomised block design (each block contained one replicate of each treatment combination). Plants were infested with a single apterous aphid which was allowed to reproduce overnight, then a total of three nymphs were retained on each plant. Mean nymph mass and number of surviving nymphs were recorded at 72h and 168h, after which a random single nymph was returned to the plant; for this nymph, data were collected on the length of the pre-reproductive period (d) and the intrinsic rate of population increase (r_m). Aphids were caged onto the first fully expanded true leaf using Perspex clip-cages of 25mm internal diameter (MacGillivray and Anderson, 1957). Nymph mass gain was calculated as the change in mass between 168 and 72h and aphid r_m was calculated using the equation of Wyatt and White (1977): $rm = 0.74 \frac{\ln(Fd)}{d}$, where d is the time period between birth and production of first progeny, and Fd is the total progeny produced over a time period equal to d.

Electrical penetration graph (EPG) monitoring of aphid feeding

The DC-EPG technique (Tjallingii, 1978; Tjallingii, 1988; Tjallingii, 1991; Tjallingii, 2001) was employed to monitor the probing and feeding behaviour of apterous *R. padi* over a 6h period using a Giga-4 DC-EPG device (EPG Systems, The Netherlands) on plants at the true-leaf stage. A plant probe (copper rod approximately 50 mm long x 5 mm diameter), was soldered to electrical wire extending from the plant voltage output of the Giga-4 device and inserted into the plant soil. An aphid probe was made by soldering a piece of copper wire (30 mm long x 2 mm diameter) to a brass pin (tip diameter 2 mm). Approximately 30 mm of gold wire (20 μm diameter; EPG Systems, The Netherlands) was adhered to the copper end of the aphid probe using water-based silver glue (EPG Systems, The Netherlands) and aphids were connected by adhering the free end of the gold wire onto the aphid dorsum using the same water-based adhesive. Wired aphids were connected to the Giga-4 device by placing the end of the brass pin into the EPG probes with a 1 GΩ input resistance and a 50x gain (Tjallingii, 1988). The order in which *R. padi* – plant combinations were tested and allocated to an EPG probe was randomised. Data were acquired using Stylet+D software (EPG Systems, The Netherlands). A

total of 18 and 16 successful recordings were made for aphids feeding on Concerto and Hsp5, respectively. All EPG recordings were obtained within a grounded Faraday cage.

EPG waveforms were annotated using Stylet+A software (EPG Systems, The Netherlands). Waveforms were annotated by assigning waveforms to np (non-probing), C (stylet penetration/pathway), pd (intercellular punctures), E1 (saliva secretion into phloem), E2 (saliva secretion and passive phloem ingestion), F (penetration difficulty) or G (xylem ingestion) phases (Tjallingii, 1988; Alvarez et al., 2006). No E1e (extracellular saliva secretion) phases were detected. Annotated waveforms were converted into time-series data using the excel macro developed by Dr Schliephake (Julius Kühn-Institut, Germany).

Determination of non-glandular trichome density

Non-glandular trichome densities of Hsp5 and Concerto (n = 9) were determined using polarised light microscopy following a procedure adapted from Pomeranz et al. (2013). Briefly, the first true leaf was excised from plant stems and treated to two incubation steps to clear the leaf epidermis. Firstly, leaf area was measured and leaves were soaked in 96% Ethanol (Sigma-Aldrich, UK) for 48h before being treated with 1.25 M NaOH:EtOh (1:1, v:v) at 70°C for 2h. Leaves were stored in 50% glycerol prior to trichome visualisation; if NaOH:EtOH treatment did not clear the leaf epidermis leaves were further treated with 80% lactic acid for 6h. To analyse trichome density samples were placed adaxial side up on microscope slides (75 x 25 mm; Corning, UK). A polarising light microscope was created using two 50 mm^2 polarising filters (Sigma-Aldrich, UK), one placed below the stage of a stereo-microscope but above the light source and the second attached below the objective lens. Non-glandular trichomes appeared illuminated under the polarised microscope setup and the number of trichomes per unit area were counted. To allow correlation of trichome density with aphid performance, an additional *R. padi* performance experiment (as described above) was carried out on these plants prior to trichome density analysis.

Epicuticular wax analysis

A Fourier Transform Infrared (FTIR) spectrometer (Bruker Vertex 70 FTIR spectrometer, Bruker, Ettlingen, Germany), incorporating a Diamond Attenuated Total Reflection (DATR) sampling accessory, was used to identify the functional groups present in chemical extracts from the surfaces of Hsp5 and Concerto leaves. Briefly, the waxes from upper and lower leaf

surfaces of the first true leaf (n = 4) were extracted with dichloromethane (DCM) by running approximately 1-2 ml of DCM along the leaf surface directly onto the diamond window of the ATR sampling accessory; after evaporation of the DCM, FTIR spectra were recorded of the films deposited on the DATR. Signal-to-noise ratio was enhanced by taking 200 scans for each sample and averaging to obtain a single spectrum. The spectral range scanned was 4000 cm^{-1} to 400 cm^{-1} and a background reading was taken before each sample was analysed.

Analysis of phloem amino acid composition

Hsp5 and Concerto plants were grown under glasshouse conditions (as described above) in a temporally-split randomised block design: three temporal blocks, one initial temporal block comprising two replicate sub-blocks, and two further temporal blocks each comprising four replicate sub-blocks. Each sub-block contained a single replicate of every treatment combination. Plants at the first true-leaf stage were infested with either ten apterous *R. padi* adults caged onto the plants with microperforated bags (Polybags, UK) or were aphid-free bagged controls. Samples were collected at 0 and 24h post-aphid infestation. Phloem sap was collected using the method of King and Zeevaart (1974); leaves (*n = 10*) were excised from the stem, placed in a dish containing 1 mM EDTA (Sigma-Aldrich, UK) solution and immediately re-cut. The cut surface of the leaf was placed into 200μl of filter-sterilised 1mM EDTA solution (pH 7.5 with NaOH) in a 2 mL Eppendorf™ microcentrifuge tube and incubated for 1.5h at room temperature in a darkened exudation chamber (a polystyrene box containing a dish of 6 M K_2HPO_4 (Sigma-Aldrich, UK), and moistened paper towels to maximise humidity and decrease leaf transpiration). After incubation, leaves were removed and the samples of EDTA solution containing exuded phloem sap were stored at -80ºC until analysis.

Reverse-phase HPLC was used to separate phloem amino acids Asp, Glu, Asn, His, Ser, Gln, Arg, Gly, Thr, Tyr, Ala, Trp, Met, Val, Phe, Ile, Leu and Lys using an Agilent HP1100 series auto-sampling LC system equipped with a ZORBAX™ Eclipse AAA column and fluorescence detector. Amino acids were derivatised using *o*-phthaldialdehyde (Sigma-Aldrich, UK) following the method developed by Jones et al. (1981); all protein amino acids except proline and cysteine could be detected by this method, with a detection limit of approximately 0.5 pmol. Samples were processed in their experimental blocks along with blanks (1 mM EDTA solution that was not exposed to an excised leaf) as controls. Amino acids were identified and

quantified by comparison with known concentrations of amino acid standards (AA-S-18

(Sigma, UK), supplemented with Asp, Glu and Trp).

qPCR analysis of plant defence genes

Five sub-blocks over the three temporal blocks (one from the first block and two from each

subsequent temporal block) were chosen at random from the phloem exudation experiment

described above and leaf tissue was sampled for gene expression analysis. Approximately

25mm length of leaf tissue was cut from the true leaf and flash-frozen in liquid nitrogen

immediately prior to the excision of the leaf for phloem exudation. Plant material was stored

at -80°C until RNA extraction.

For RNA extraction leaf samples were ground to a fine powder under liquid nitrogen using a

mortar and pestle and total RNA was extracted with the Norgen Plant/Fungi RNA Extraction

Kit (Sigma-Aldrich, UK) following the manufacturer's protocol with additional DNAse I

treatment (Qiagen, UK); RNA quality and quantity was assessed with a NanoDrop® ND-1000

Spectrophotometer (ThermoFischer, UK). cDNA synthesis was carried out with approximately

1000 ng RNA using the SuperScript® III cDNA synthesis kit (Sigma-Aldrich, UK), following the

manufacturer's protocol. qPCR primers were designed using the Roche Universal Probe

Library Assay Design Centre. Reference gene primers were described by Hua et al. (2015). All

primer sequences are shown in Supplementary Table 1 and were supplied by Sigma-Aldrich,

UK. Primers were validated for PCR efficiency prior to use, and reference gene primers were

tested for stability across all treatments following the geNorm procedure (Vandesompele et

al., 2002). Gene expression analysis used SYBR® Green chemistry with GoTaq® qPCR Master

Mix (Promega, UK) on a StepOne™ Real-Time PCR Machine (Applied Biosystems, UK).

Reactions were carried out in 12.5 µl reactions with a final concentration of 1x GoTaq® qPCR

Master Mix, 1µM of each primer, 1.4 mM $MgCl_2$, 2.4 µM CXR reference dye and a cDNA

quantity of approx. 12.5 ng (assuming 1:1 RNA:cDNA conversion). The qPCR conditions were

as follows: 95°C for 15 mins followed by 40 cycles of denaturing for 15s at 95°C, annealing for

30s at 60°C, and 30s at 72°C for DNA extension. Fluorescence was recorded at the end of each

annealing cycle and a melting curve was incorporated into the end of the qPCR programme.

Data were normalised to the mean expression of two reference genes, *HvCYP* (AK253120.1) and

HvUBC (AK248472.1)*,* and $2^{-\Delta\Delta Ct}$ methodology (Livak and Schmittgen, 2001) was used to

determine differential expression, with Concerto at 0h aphid infestation used as the

experimental control treatment. Expression levels at 24h were further normalised to uninfested control plants. Samples were processed in experimental blocks with three technical replicates at the PCR level.

Statistical analysis

All statistical analyses were carried out using R Studio v.1.0.143 running R v.3.4.3 (R Core Team, 2014) using packages: car v.2.1-4 (Fox and Weisberg, 2011), coxme v.2.2-7 (Therneau, 2018), dunn.test v.1.3.5 (Dinno, 2017)**,** ggplot2 v.2.2.1 (Wickham, 2009), ggpubr v. 0.1.2 (Kassambara, 2017), ggfortify v.0.4.5 (Tang et al., 2016), lawstat v.3.2 (Hui et al., 2008), lme4 v.1.1-13 (Bates et al., 2015), lmerTest v.2.0-33 (Kuznetsova et al., 2017), lsmeans v.2.27-62 (Lenth, 2016), survival v.2.41-3 (Therneau and Grambsch, 2000)**,** survminer v.0.4.2 (Kassambara and Kosinski, 2017), vegan v.2.5-3 (Oksanen et al., 2013).

Aphid juvenile mass gain and adult r_m, were modelled using linear mixed effects models with experimental block incorporated as a random factor. ANOVA with type III Satterthwaite approximation for degrees of freedom was used to analyse the final models with calculation of Least Squares Means used for *post-hoc* testing. A Cox proportional hazards regression model was used for nymph survival analysis (Therneau and Grambsch, 2000; Therneau, 2018), incorporating experimental block as a random factor; a χ^2 test was used on the final model. A log-rank test with Benjamini-Hochberg correction was used to carry out pairwise comparisons between aphid-plant combinations. Aphid feeding behaviour was assessed globally by fitting a permutated MANOVA to the dataset. Response variables with normal data distribution were analysed by either ANOVA or general linear models, while non-normally distributed data were analysed with Kruskal-Wallis tests.

Levene's test (Hui et al., 2008) was used to analyse differences in trichome density; a Kendall's rank correlation tau was then used to test for correlations between trichome density and *R. padi* performance. Differences in leaf surface chemistry were analysed using a Welch two sample *t*-test by comparing the total number of identified functional groups in the two plant species.

Phloem amino acid concentrations (pmol/µl) were expressed as relative amino acid composition by converting into mole % of total amino acid content and subjected to principal component analysis with a correlation matrix. Three principal components explained 69% of

observed variation and were analysed by fitting a linear mixed effects model incorporating aphid treatment, plant, time-point, plant x time-point interaction, and leaf subsampling for RNA (to test effect on amino acid composition) as explanatory variables with temporal block, randomised block and HPLC batch as random factors. Final models were analysed with a χ^2 test. Differential gene expression ($2^{-\Delta\Delta Ct}$) data were analysed using Kruskall-Wallis rank-sum tests, with subsequent Dunn's test *post-hoc* analysis of the plant x infestation interaction.

Results

A wild relative of barley exhibits partial-resistance against multiple aphid species

Aphid performance experiments were undertaken to assess whether partial-resistance in Hsp5 against *R. padi* (Delp et al., 2009; Leybourne et al., 2018) extends to other aphid species. We assessed the fitness of *R. padi, S. avenae,* and *U. humboldti* on Hsp5 in comparison with a commercial barley cultivar (Concerto) and found evidence for partial-resistance against all three aphid species. Specifically, nymph survival was significantly lower on Hsp5 (χ^2_1 = 10.65; p = 0.001; Fig. 1A; Table 1); pairwise comparison showed that nymph survival was reduced on Hsp5 for *U. humboldti* (p = <0.001), but not for *R. padi* (p = 0.057) or *S. avenae* (p = 0.973). Furthermore, nymph mass gain ($F_{56,1}$ = 9.14, p = 0.003; Fig. 1B; Table 1) and the rate of population increase (r_m) ($F_{54,1}$ = 27.43, p = <0.001; Fig. 1C; Table 1) were reduced for all aphid species when feeding from Hsp5. Differences were also detected between aphid species, with *R. padi* exhibiting the highest r_m ($F_{54,2}$ = 99.10, p = <0.001; Fig. 1B; Table 1) and *S. avenae* the largest mass gain ($F_{56,2}$ = 49.82, p = <0.001; Fig. 1C; Table 1). On average *U. humboldti* was the least fit species (Fig. 1A-C).

Partial-resistance against R. padi involves mesophyll and phloem resistance factors

To elucidate the potential underlying partial-resistance mechanism(s) against aphids in Hsp5 *R. padi* feeding behaviour was monitored using the EPG technique. Overall, the feeding behaviour of *R. padi* differed significantly when feeding from Hsp5 compared with Concerto ($F_{1,33}$ = 2.61; p = 0.022; Fig. 2; Table 2). Aphids showed significant differences in feeding behaviour at the leaf epidermis and within mesophyll tissue (Fig. 3; Table 2). We observed an approximate three-fold decrease in the time to the first epidermal probe when feeding on Hsp5 ($F_{1,33}$ = 7.99; p = 0.008), alongside a shorter duration of the first probe ($F_{1,33}$ = 6.94; p = 0.013) and an increased time to the first sieve element puncture ($F_{1,33}$= 6.33; p = 0.017). In

addition, aphids feeding on Hsp5 showed a two-fold delay in initiating passive phloem

ingestion ($\chi^2_{1,33}$ = 4.29; p = 0.038) and the number of intracellular punctures observed during

the first aphid probe of plant tissue was lower, with an average of ten intracellular probes,

compared with 24 on Concerto (F $_{1,33}$ = 3.95; p = 0.047; Table 2).

Feeding patterns within the vascular tissue also differed (Fig. 4; Table 2). Specifically, aphids

feeding on Hsp5 showed delayed initiation of sustained phloem ingestion (ingestion for >10

mins) after the first sieve element puncture ($\chi^2_{1,33}$ = 4.29; p = 0.038). The total length of time

aphids spent in the sustained feeding phase was also 2.5x shorter ($\chi^2_{1,33}$ = 7.49; p = 0.006),

and the ratio of time aphids spent ingesting phloem relative to the length of time probing

plant tissue (ingestion:pathway ratio) was four-fold lower ($\chi^2_{1,33}$ = 7.43; p = 0.006).

Furthermore, aphids feeding on Hsp5 spent three times longer ingesting xylem ($\chi^2_{1,33}$ = 5.28;

p = 0.022). Table 2 reports additional feeding parameters which highlight further mesophyll

and vascular-mediated resistance against *R. padi* in Hsp5.

Leaf surface architecture differs between susceptible and partially-resistant plants.

The leaf surface represents a key interface between plants and insects, with many factors

reported to modulate plant-insect interactions, including leaf trichomes and epicuticular

waxes (Agrawal et al., 2009; Glas et al., 2012; Karley et al., 2016). Leaf trichome counts

revealed Hsp5 had a significantly higher abundance of non-glandular leaf trichomes ($F_{1,18}$ =

6.24; p = 0.022; Fig. 5). Non-glandular trichome abundance showed a negative correlation

with *R. padi* fecundity (Z = -3.60; p = <0.001; T = -0.56; Supplementary Fig. 1A) but there was

no relation with nymph mass gain (Z = -0.29; p = 0.770; T = -0.04; Supplementary Fig. 1B). In

addition, we found differences in the epicuticular wax composition between the two plants.

The complexity of the epicuticular wax differed in that aliphatic hydrocarbons were detected

in surface extracts of Hsp5 leaves, two Hsp5 replicates also contained weak ester bands

(Supplementary Table 3). In Concerto the chemical mixture contained a more consistent

mixture of aliphatic hydrocarbons and more pronounced ester bands, alongside a more

complex, but variable, mixture of chemical groups, including carboxylates and amides (*t* = -

4.89; df = 6; p = 0.002). Overall, the results were consistent within Hsp5 but more variable in

Concerto (Supplementary Table 3). However, the detection of some Nitrogen-containing

functional groups in Concerto leaf extracts could be a result of extracting some components

within the upper tissue layers, this potentially highlights differences in the thickness of the
wax between Hsp5 and Concerto.

Basal expression levels of several plant defence genes are elevated in Hsp5

To investigate the level of defence gene expression in Concerto versus Hsp5 we selected
marker genes for defensive thionins (*HvTHIO1, HvTHIO2, HvβTHIO*) alongside JA (HvLOXA,
HvLOX2 and *HvJAZ*), SA (*HvNPR1*), ET (*HvERF*), and ABA (*HvA1*) signalling pathways. We
assessed expression of marker genes in both plant types constitutively and in response to 24h
of aphid infestation. We observed 10-(*HvTHIO1)*, 13-(*HvTHIO)* and 7-fold (*HvβTHIO)* higher
expression of the three thionin genes in Hsp5 (Fig 6; Table 3). Thionin gene expression levels
remained higher in Hsp5 24h after aphid infestation but were not differentially regulated in
either plant in response to aphid infestation (Fig 6; Table 3). In addition, *HvLOXA*, *HvLOX2,*
HvA1, and *HvERF1*, were more highly expressed in Hsp5, with 3.5-, 6-, 10- and 14-fold higher
expression levels, respectively (Fig 6; Table 3).

No genes were differentially expressed in response to aphid infestation. However, several of
the phytohormone signalling genes were differentially expressed in response to the plant x
aphid infestation interaction; namely *HvLOXA*, *HvLOX2, HvNPR1,* and *HvERF1* (Table 3).
HvLOXA was significantly down-regulated in Hsp5 after 24h aphid infestation to levels similar
to those observed in Concerto (Fig. 6; Table 4). For *HvLOX2, HvNPR1,* and *HvERF,* expression
levels were higher in Hsp5 after 24h of aphid infestation compared with levels in Concerto
(Fig. 6; Table 4). *HvJAZ* was not differentially regulated in response to any treatment factor.

Phloem amino acid composition in Hsp5 is characterised by a reduction in the proportion of
essential amino acids and an increased abundance of asparagine

To investigate phloem factors contributing to partial-resistance against aphids (Fig. 4; Table
2) we analysed the amino acid composition and the proportion of essential and non-essential
amino acids in phloem exudates (Fig. 7). Principal component analysis of amino acid
composition (mol% of Asp, Glu, Asn, His, Ser, Gln, Arg, Gly, Thr, Tyr, Ala, Trp, Met, Val, Phe,
Ile, Leu, and Lys) of phloem exudates revealed that three principal components explained 69%
of the observed variation in overall amino acid composition, with all three principle
components showing a degree of separation between the two plant types (Table 5).
Furthermore, >50% of the variation was explained by the first two principal components (Fig.

7C; Table 5) with separation of Hsp5 from Concerto largely occurring along PC1 due to differences in the relative amounts of Asn and His vs. most other essential amino acids (Fig. 7C). The total percentage of essential amino acids (Arg, His, Ile, Leu, Lys, Met, Phe, Thr, Trp, Val; described in Morris (1991)) was higher in Concerto compared with Hsp5 (18.97% vs 11.45%; χ^2_1 = 25.43; p = <0.001; Fig. 7D; Table 6).

The difference in composition of non-essential amino acids was most pronounced between the two plant types, with a higher proportion of Asn and lower proportions of Glu and Gly in Hsp5 (42.75%, 7.34% and 12.69%, respectively) compared with Concerto (7.55%, 15.18% and 22.43%, respectively) (Fig 7A-B). Amino acid composition and percentage of essential amino acids changed over the 24h of the experiment, but not in response to aphid treatment (Table 5; Table 6). Harvesting of leaf material for qPCR analysis had no effect on the phloem amino acid composition (Table 5; Table 6).

Discussion

In this study we provide evidence for mesophyll and phloem mediated partial-resistance against aphids in a wild barley relative, which is effective against three aphid species, including two agricultural pests and an invasive species. Our data suggest the partial-resistance in this progenitor of modern barley is complex and may be associated with elevated defence gene expression and reduced phloem quality.

Mesophyll and phloem-based factors contribute to partial-resistance in Hsp5

Our observation that *R. padi* feeding on Hsp5 spends more time in the pathway phase (probing into plant tissue) and takes longer to reach the phloem and initiate salivation/phloem ingestion is representative of feeding patterns associated with mesophyll-based resistance (Alvarez et al., 2006). Factors associated with mesophyll resistance are thought to antagonise and impede stylet penetration (Montllor and Tjallingii, 1989; Alvarez et al., 2006). In addition, *R. padi* spends less time ingesting phloem sap and salivating into the phloem, in line with resistance factors residing in the phloem (Alvarez et al., 2006). Phloem based resistance factors generally alter the ingestion efficiency of aphids and can involve feeding antagonists, alterations to the phloem metabolic profile, and the plugging of sieve-tube elements (Klingler et al., 1998; Klingler et al., 2005; Pegadaraju et al., 2007; Medina-Ortega and Walker, 2015; Greenslade et al., 2016; Peng and Walker, 2018).

Partial-resistance against aphids in other plant-aphid systems have similarly pointed to the

importance of mesophyll and phloem-based resistance factors (Montllor and Tjallingii, 1989;

Caillaud et al., 1995; Pegadaraju et al., 2007; Guo et al., 2012; Greenslade et al., 2016). For

example, partial resistance in *Triticum monococcum* lines (a wild relative of wheat) against *S.*

avenae and *R. padi* involved resistance at the phloem level, as determined by EPG (Caillaud

et al., 1995; Greenslade et al., 2016). Interestingly, metabolic profiling of resistant and

susceptible lines in the latter study indicated different metabolite profiles between

susceptible and partially-resistant plants, including lower levels of primary metabolites in the

partially-resistant lines and elevated Asn (Greenslade et al., 2016). Resistance against *D. noxia*

in five cereal species was shown to involve phloem and mesophyll resistance factors, with

elevated 2,4-dihydroxy-7-methoxy-1,4-benzoxazin-3-one (DIMBOA) levels hypothesised to be

a contributing factor towards resistance (Mayoral et al., 1996). Resistance against *D. noxia* in

wheat also involved individual resistance genes; the most recent of these to be characterised

is *Dn10* (Li et al., 2018), although the tissue types involved in conferring this resistance are

unknown.

In addition, *Arabidopsis thaliana* defence against *Myzus persicae* involved *phytoalexin*

deficient4 (PAD4), and reduced susceptibility in *PAD4* overexpressing lines was associated

with reduced phloem ingestion (Pegadaraju et al., 2007). Furthermore, gene-for-gene type

resistance against the pea aphid, *Acyrthosiphon pisum,* in *Medicago truncatula* line Jemalong

A17 (A17) is conferred by a single gene, *RAP1,* which maps to chromosome 3 (Stewart et al.,

2009). Further characterisation showed that A17 also conferred resistance against the

bluegreen aphid, *A. kondoi,* as well as *A. pisum,* and that this resistance was phloem-mediated

and linked to a QTL on chromosome 3 (Guo et al., 2012). The inorganic chemistry of leaf tissue

can also play a role in plant defence (Lane, 2002; Boyd, 2012), with oxalates highlighted as

important contributors towards plant defence against pathogens in cereals (Lane, 2002). The

inorganic chemistry of Hsp5 and Concerto leaves was not assessed in this study, but a

comparison of leaf chemistry could highlight further chemical groups which could be involved

in mediating plant-aphid interactions.

Differences in leaf surface characteristics contribute little to partial-resistance against

aphids in Hsp5

Leaf surface architectural differences were detected between Concerto and Hsp5 and
included a higher abundance of non-glandular leaf trichomes in Hsp5 and a more complex
chemistry in the epicuticular extracts of Concerto. Non-glandular trichomes have been
reported to decrease *R. padi* fitness in wheat (Roberts and Foster, 1983), but have contrasting
effects on herbivore fitness in other plants. For example, the density of non-glandular
trichomes in raspberry negatively affected spider mite performance but had minimal impact
on aphid fitness (Karley et al., 2016).

Although we detected a positive correlation between trichome abundance and *R. padi* r_m,
there was no correlation between trichome abundance and juvenile *R. padi* mass gain,
indicating that either non-glandular trichomes have differential effects on aphids at different
life-stages or that the higher abundance of non-glandular trichomes on the leaf surface of
Hsp5 are not a primary cause of partial-resistance to aphids. Indeed, EPG analysis indicated
that leaf surface traits contributed little to partial-resistance in Hsp5. Aphids feeding from
Hsp5 did not show increased time spent in the non-probing phase (a key indicator of
epidermal-mediated resistance: Alvarez et al. (2006). Moreover, the time taken for aphids to
penetrate the leaf surface was shorter on Hsp5, indicating aphids experienced fewer barriers
to probing the leaf epidermis on Hsp5. Less complex chemistry at the leaf epicuticle in Hsp5
may have played a role in promoting aphid probing of plant tissue and previous studies have
shown that epicuticular chemical compounds can either promote or deter aphid probing of
plant tissue (Powell et al., 1999). Barley epicuticular waxes can contribute towards resistance
against *R. padi* (Tsumuki et al., 1989), although the influence of specific chemical functional
groups on aphid behaviour is not well characterised.

Basal defence gene expression is elevated in Hsp5

In line with Delp et al. (2009) and Mehrabi et al. (2014), basal expression levels of thionin
genes, which contribute to barley defences against aphids (Escudero-Martinez et al., 2017),
and *LOX2*, a JA-signalling marker, were elevated in Hsp5. The feeding parameters exhibited
by *R. padi* while feeding on Hsp5 indicated that mesophyll-based mechanisms conferred
short-term (6h) partial-resistance against aphids. Thionins are antimicrobial peptides present
in plant cell walls and within the intracellular space (Reimann-Philipp et al., 1989; Stec, 2006).
It is therefore likely that aphids are exposed to thionins when probing mesophyll tissue, and
these could contribute to reduced aphid performance on Hsp5.

In addition, several plant defence signalling genes, *LOX* (JA)*, ERF1* (ET) and *A1* (ABA) showed

higher basal expression in Hsp5, suggesting that Hsp5 may be able to respond more efficiently

to biotic stimuli. In line with our data, *HvLOX2* genes and other components of the JA (*HvAOS,*

HvAOC), ET (*HvACCO, HvACCS*), and Auxin (*HvTDS, HvTSM*) signalling pathways showed higher

basal expression in the barley cultivar Stoneham, which is partially-resistant against *D. noxia,*

relative to susceptible cultivar Otis (Marimuthu and Smith, 2012). Furthermore,

overexpression of *HvLOX2* in barley resulted in decreased *R. padi* fecundity five days after

infestation (Losvik et al., 2017)*,* pointing to a functional role of this gene and JA signalling in

short-term barley defence against aphids. Expression levels of *CmERF1* in early responses (6h)

to aphid infestation were around ten-fold higher in a resistant variety of melon compared

with a susceptible variety (Anstead et al., 2010), and higher ET levels have been reported to

contribute to *R. maidis* resistance in maize (Louis et al., 2015). ABA mediated-processes have

been implicated in increasing both plant resistance (Zhu-Salzman et al., 2004; Park et al.,

2006) and susceptibility (Kerchev et al., 2013; Hillwig et al., 2016) to aphids. Therefore the

consequence of elevated *HvA1* in the context of partial aphid resistance in Hsp5 is not clear.

Guo et al. (2016) reported that elevated ABA levels in response to drought conditions can lead

to increased aphid xylem ingestion. This finding suggests that elevated expression of

components of the ABA signalling pathway could indirectly contribute to partial-resistance by

encouraging xylem ingestion, thereby reducing phloem ingestion. Indeed, longer periods of

xylem ingestion were detected in *R. padi* feeding from Hsp5, indicating that there may be

potential fitness consequences for aphids as a result of higher basal expression levels.

In contrast, basal expression levels of *HvNPR1*, a SA-signalling marker, were not significantly

different between Hsp5 and Concerto. However, *HvNPR1* levels in Concerto were down-

regulated 24h after aphid infestation to a level that was significantly lower than *HvNPR1* levels

in Hsp5 after 24h of aphid infestation. Therefore, SA-mediated defences in Concerto, but not

Hsp5, may be repressed upon aphid infestation leading to increased susceptibility of this

cultivar. Indeed, NPR1 is required for initiation of SA-mediated defence against aphids in *A.*

thaliana (Moran and Thompson, 2001; Wu et al., 2012).

Essential amino acid availability is reduced in Hsp5

The observation that the relative concentration of essential amino acids was lower in the

phloem of Hsp5 compared with Concerto could be associated with lower aphid performance.

Alterations to amino acid composition can reduce plant palatability and nutritional quality
and contribute to increased resistance against aphids (Sandström and Pettersson, 1994;
Ponder et al., 2000; Karley et al., 2002). For example, decreased survival and fecundity of *M.*
persicae and *Macrosiphum euphorbiae* feeding from older "tuber-filling" potato plants
compared with younger "pre-tuber-filling" plants was linked to changes in amino acid
composition (Karley et al., 2002). When presented with chemically-defined diets
representative of the phloem sap composition of young and mature potato plants, *M.*
persicae and *Ma. euphorbiae* showed decreased feeding rate on the "tuber-filling" diets
(Karley et al., 2002). In addition, *R. padi* feeding from barley plants grown under nitrogen-
limited conditions exhibited reduced r_m compared with *R. padi* feeding on barley grown under
nitrogen-rich conditions (Ponder et al., 2000). Examination of phloem amino acid composition
indicated that composition was altered under nitrogen-limited conditions. Interestingly,
plants under nitrogen-limited conditions contained a higher percentage of Asn, a lower
percentage of Gly and a small reduction in the concentration of essential amino acids
compared with nitrogen-fertilised plants (Ponder et al., 2000).

Changes in the abundance of specific amino acids cannot be readily associated with increased
insect resistance/susceptibility in a given plant species. It is most likely that any observed
effect on aphid fitness is due primarily to the ratio of essential:non-essential amino acids as a
result of compositional changes. Indeed, essential amino acids are generally elevated in
susceptible plants compared with varieties showing increased resistance against aphids
(Auclair, 1976). Vogel and Moran (2011) found that the mass of multiple *A. pisum* biotypes
was reduced when essential amino acids were removed from aphid artificial diets; this was
observed even though aphids have the capacity to synthesis essential amino acids via their
essential endosymbiont, *Buchnera aphidicola* (Douglas and Prosser, 1992). Furthermore,
some aphid species actively remobilise plant nutrients to increase the abundance of essential
amino acids in susceptible plant hosts (Telang et al., 1999; Sandström et al., 2000). Enhanced
mobilisation of plant nutrients is thought to be the mechanism explaining resistance
breakdown in wheat by a virulent biotype of the greenbug, *Schizaphis graminum* (Dorschner
et al., 1987). In a study by Sandström et al. (2000), *R. padi* feeding did not lead to increased
phloem mobilisation of amino acids, similar to observations made in our experimental system

where *R. padi* infestation did not alter the phloem amino acid composition in Hsp5 or
Concerto.

Conclusion

This study shows that partial-resistance against aphids in wild barley can be wide-ranging and
effective against multiple aphid species. Broad partial-resistance traits are agriculturally and
economically important as crops are likely to be infested by multiple insect species under field
conditions (Stam et al., 2014). However, compared with gene-for-gene resistance traits,
comparatively little is known about the genetic determinants of partial-resistance. The
underlying factors contributing towards partial-resistance are complex and likely to involve
multiple, independent resistance pathways (Mayoral et al., 1996; Moharramipour et al., 1997;
Seah et al., 1998). We show that partial-resistance against aphids in wild barley involves
resistance factors that reside in both the mesophyll and vascular tissues. This partial-
resistance is potentially caused by heightened basal expression of defence and
phytohormone signalling genes, alongside altered phloem amino acid quality. However, the
functional molecular role of these differences, and how they contribute towards partial-
resistance, remains to be established.

Supplementary Data

612 Supplementary Table 1: primers used in RT-qPCR analysis.

613 Supplementary Table 2: Additional EPG parameters.

614 Supplementary Table 3: FTIR results for leaf surface extracts

615 Supplementary Fig. 1: Correlation of insect fitness with non-glandular trichome density.

Acknowledgements

D.J.L was funded by the James Hutton Institute and the Universities of Aberdeen and Dundee
through a Scottish Food Security Alliance (Crops) PhD studentship. A.J.K, T.A.V, A.H.J.R,
A.M.M, and E.P-F were supported by the strategic research programme funded by the

Scottish Government's Rural and Environment Science and Analytical Services Division. J.I.B.B.

was supported by the European Research Council (310190-APHIDHOST). We would like to thank

Dr Freddy Tjallingii (EPG Systems, The Netherlands), Professor Alberto Fereres (CSIC, Spain)

and Professor Gregory Walker (University of California, Riverside, USA) for providing EPG

training and Gaynor Malloch (James Hutton Institute) for kindly providing COI barcoding

primers.

629 Author Contributions

630 JIBB, AJK, and DJL conceived and designed the experiments. DJL, EP-F, and AMM performed

631 the experiments. JIBB, AJK, DJL, and JAHR analysed the data. DJL wrote the manuscript with

632 input from JIBB, AJK, TAV, and JAHR. All authors read and approved the final manuscript.

References

Agrawal, A. A., Fishbein, M., Jetter, R., Salminen, J. P., Goldstein, J. B., Freitag, A. E. and Sparks, J. P. (2009). Phylogenetic ecology of leaf surface traits in the milkweeds (*Asclepias spp.*): chemistry, ecophysiology, and insect behavior. *New Phytologist* **183**, 848-67.

Ahmad, S., Veyrat, N., Gordon-Weeks, R., Zhang, Y., Martin, J., Smart, L., Glauser, G., Erb, M., Flors, V., Frey, M. et al. (2011). Benzoxazinoid Metabolites Regulate Innate Immunity against Aphids and Fungi in Maize. *Plant Physiology* **157**, 317-327.

Åhman, I., Tuvesson, S. and Johansson, M. (2000). Does Indole Alkaloid Gramine Confer Resistance in Barley to Aphid *Rhopalosiphum Padi*? *Journal of Chemical Ecology* **26**, 233-255.

Alexandratos, N. and Bruinsma, J. (2012). World Agriculture Towards 2030/2050: the 2012 Revision. ESA Working Paper No. 12-03. Rome: FAO.

Alvarez, A. E., Tjallingii, W. F., Garzo, E., Vleeshouwers, V., Dicke, M. and Vosman, B. (2006). Location of resistance factors in the leaves of potato and wild tuber-bearing *Solanum* species to the aphid *Myzus persicae*. *Entomologia Experimentalis et Applicata* **121**, 145-157.

Anstead, J., Samuel, P., Song, N., Wu, C., Thompson, G. A. and Goggin, F. (2010). Activation of ethylene-related genes in response to aphid feeding on resistant and susceptible melon and tomato plants. *Entomologia Experimentalis et Applicata* **134**, 170-181.

Aradottir, G. I., Martin, J. L., Clark, S. J., Pickett, J. A. and Smart, L. E. (2017). Searching for wheat resistance to aphids and wheat bulb fly in the historical Watkins and Gediflux wheat collections. *Annals of Applied Biology* **170**, 179-188.

Auclair, J. L. (1963). Aphid Feeding and Nutrition. *Annual Review of Entomology* **8**, 439-490.

Auclair, J. L. (1976). Feeding and Nutrition of the Pea Aphid, *Acyrthosiphon Pisum* (Harris), with Special Reference to Amino Acids. In *The Host-Plant in Relation to Insect Behaviour and Reproduction*, (ed. T. Jermy), pp. 29-34. Boston, MA: Springer US.

Bari, R. and Jones, J. D. G. (2009). Role of plant hormones in plant defence responses. *Plant Mol Biol* **69**, 473-488.

Barria, B. N., Copaja, S. V. and Niemeyer, H. M. (1992). Occurrence of DIBOA in wild *Hordeum* species and its relation to aphid resistance. *Phytochemistry* **31**, 89-91.

Boyd, R. S. (2012). Plant defense using toxic inorganic ions: Conceptual models of the defensive enhancement and joint effects hypotheses. *Plant Science* **195**, 88-95.

Caillaud, C. M., Pierre, J. S., Chaubet, B. and Pietro, J. P. (1995). Analysis of wheat resistance to the cereal aphid *Sitobion avenae* using electrical penetration graphs and flow charts combined with correspondence analysis. *Entomologia Experimentalis et Applicata* **75**, 9-18.

Chandrasekhar, K., Shavit, R., Distelfeld, A., Christensen, S. A. and Tzin, V. (2018). Exploring the metabolic variation between domesticated and wild tetraploid wheat genotypes in response to corn leaf aphid infestation. *Plant Signaling & Behavior* **https://doi.10.1080/15592324.2018.1486148**.

Chen, M.-h., Han, Z.-j., Qiao, X.-f. and Qu, M.-j. (2007). Mutations in acetylcholinesterase genes of *Rhopalosiphum padi* resistant to organophosphate and carbamate insecticides. *Genome* **50**, 172-179.

Delp, G., Gradin, T., Åhman, I. and Jonsson, L. M. (2009). Microarray analysis of the interaction between the aphid *Rhopalosiphum padi* and host plants reveals both differences and similarities between susceptible and partially resistant barley lines. *Molecular Genetics and Genomics* **281**, 233-248.

Dempewolf, H., Eastwood, R. J., Guarino, L., Khoury, C. K., Müller, J. V. and Toll, J. (2014). Adapting Agriculture to Climate Change: A Global Initiative to Collect, Conserve, and Use Crop Wild Relatives. *Agroecology and Sustainable Food Systems* **38**, 369-377.

Dewar, A. M. and Foster, S. P. (2017). Overuse of Pyrethroids may be implicated in the Recent *BYDV* Epidemics in Cereals. *Outlooks on Pest Management* **28**, 7-12.

Dinno, A. (2017). dunn.test: Dunn's Test of Multiple Comparisons Using Rank Sums. *CRAN repository* **available: https://CRAN.R-project.org/package=dunn.test**.

Directive. (2009/128/EC). Directive 2009/128/EC of the European Parliament and of the Council of 21 October 2009 establishing a framework for Community action to achieve the sustainable use of pesticides.

Dorschner, K. W., Ryan, J. D., Johnson, R. C. and Eikenbary, R. D. (1987). Modification of Host Nitrogen Levels by the Greenbug (Homoptera: Aphididae): Its Role in Resistance of Winter Wheat to Aphids. *Environmental Entomology* **16**, 1007-1011.

Douglas, A. E. and Prosser, W. A. (1992). Synthesis of the essential amino acid tryptophan in the pea aphid (*Acyrthosiphon pisum*) symbiosis. *Journal of Insect Physiology* **38**, 565-568.

Escudero-Martinez, C. M., Morris, J. A., Hedley, P. E. and Bos, J. I. B. (2017). Barley transcriptome analyses upon interaction with different aphid species identify thionins contributing to resistance. *Plant, Cell & Environment* **40**, 2628-2643.

Folmer, O., Black, M., Hoeh, W., Lutz, R. and Vrijenhoek, R. (1994). DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology Biotechnology* **3**, 294-299.

Forslund, K., Pettersson, J., Bryngelsson, T. and Jonsson, L. (2000). Aphid infestation induces PR-proteins differently in barley susceptible or resistant to the bird cherry-oat aphid (*Rhopalosiphum padi*). *Physiologia Plantarum* **110**, 496-502.

Foster, S. P., Paul, V. L., Slater, R., Warren, A., Denholm, I., Field, L. M. and Williamson, M. S. (2014). A mutation (L1014F) in the voltage-gated sodium channel of the grain aphid, *Sitobion avenae*, is associated with resistance to pyrethroid insecticides. *Pest Management Science* **70**, 1249-1253.

Foyer, C. H., Rasool, B., Davey, J. W. and Hancock, R. D. (2016). Cross-tolerance to biotic and abiotic stresses in plants: a focus on resistance to aphid infestation. *Journal of experimental botany* **67**, 2025-2037.

Gianoli, E. and Niemeyer, H. M. (1998). DIBOA in wild Poaceae: Sources of resistance to the Russian wheat aphid (*Diuraphis noxia*) and the greenbug (*Schizaphis graminum*). *Euphytica* **102**, 317-321.

Gibson, R. W. and Pickett, J. A. (1983). Wild potato repels aphids by release of aphid alarm pheromone. *Nature* **302**, 608-609.

Glas, J. J., Schimmel, B. C., Alba, J. M., Escobar-Bravo, R., Schuurink, R. C. and Kant, M. R. (2012). Plant glandular trichomes as targets for breeding or engineering of resistance to herbivores. *International Journal of Molecular Sciences* **13**, 17077-17103.

Greenslade, A. F. C., Ward, J. L., Martin, J. L., Corol, D. I., Clark, S. J., Smart, L. E. and Aradottir, G. I. (2016). *Triticum monococcum* lines with distinct metabolic phenotypes and phloem-based partial resistance to the bird cherry–oat aphid *Rhopalosiphum padi*. *Annals of Applied Biology* **168**, 435-449.

Guan, W., Ferry, N., Edwards, M. G., Bell, H. A., Othman, H., Gatehouse, J. A. and Gatehouse, A. M. R. (2015). Proteomic analysis shows that stress response proteins are significantly up-regulated in resistant diploid wheat (*Triticum monococcum*) in response to attack by the grain aphid (*Sitobion avenae*). *Molecular Breeding* **https://doi.org/10.1007/s11032-015-0220-x**.

Guo, H., Sun, Y., Peng, X., Wang, Q., Harris, M. and Ge, F. (2016). Up-regulation of abscisic acid signaling pathway facilitates aphid xylem absorption and osmoregulation under drought stress. *Journal of experimental botany* **67**, 681-693.

Guo, S.-M., Kamphuis, L. G., Gao, L.-L., Klingler, J. P., Lichtenzveig, J., Edwards, O. and Singh, K. B. (2012). Identification of distinct quantitative trait loci associated with defence against the closely related aphids *Acyrthosiphon pisum* and *A. kondoi* in *Medicago truncatula*. *Journal of experimental botany* **63**, 3913-3922.

Hillwig, M. S., Chiozza, M., Casteel, C. L., Lau, S. T., Hohenstein, J., Hernández, E., Jander, G. and MacIntosh, G. C. (2016). Abscisic acid deficiency increases defence responses against *Myzus persicae* in Arabidopsis. *Molecular Plant Pathology* **17**, 225-235.

Hua, W., Zhu, J., Shang, Y., Wang, J., Jia, Q. and Yang, J. (2015). Identification of Suitable Reference Genes for Barley Gene Expression Under Abiotic Stresses and Hormonal Treatments. *Plant Molecular Biology Reporter* **33**, 1002-1012.

Hui, W., Gel, Y. R. and Gastwirth, J. L. (2008). lawstat: An R Package for Law, Public Policy and Biostatistics. *Journal of Statistical Software* **28**, 1-26.

James, K. L., Randall, N. P., Walters, K. F. A., Haddaway, N. R. and Land, M. (2016). Evidence for the effects of neonicotinoids used in arable crop production on non-target organisms and concentrations of residues in relevant matrices: a systematic map protocol. *Environmental Evidence* **https://doi.org/10.1186/s13750-016-0072-9**.

Jess, S., Matthews, D., Murchie, A. and Lavery, M. (2018). Pesticide Use in Northern Ireland's Arable Crops from 1992–2016 and Implications for Future Policy Development. *Agriculture* **https://doi.org/10.3390/agriculture8080123**.

Jones, B. N., Pääbo, S. and Stein, S. (1981). Amino Acid Analysis and Enzymatic Sequence Determination of Peptides by an Improved o-Phthaldialdehyde Precolumn Labeling Procedure. *Journal of Liquid Chromatography* **4**, 565-586.

Karley, A. J., Douglas, A. E. and Parker, W. E. (2002). Amino acid composition and nutritional quality of potato leaf phloem sap for aphids. *Journal of Experimental Biology* **205**, 3009-3018.

Karley, A. J., Mitchell, C., Brookes, C., McNicol, J., O'Neill, T., Roberts, H., Graham, J. and Johnson, S. N. (2016). Exploiting physical defence traits for crop protection: leaf trichomes of *Rubus idaeus* have deterrent effects on spider mites but not aphids. *Annals of Applied Biology* **168**, 159-172.

Kassambara, A. and Kosinski, M. (2017). survminer: Drawing Survival Curves using "ggplot". *CRAN repository* **available: https://cran.r-project.org/web/packages/survminer/index.html**.

Kerchev, P. I., Karpińska, B., Morris, J. A., Hussain, A., Verrall, S. R., Hedley, P. E., Fenton, B., Foyer, C. H. and Hancock, R. D. (2013). Vitamin C and the Abscisic Acid-Insensitive 4 Transcription Factor Are Important Determinants of Aphid Resistance in *Arabidopsis*. *Antioxidants & Redox Signaling* **18**, 2091-2105.

King, R. W. and Zeevaart, J. A. D. (1974). Enhancement of Phloem Exudation from Cut Petioles by Chelating Agents. *Plant Physiology* **53**, 96-103.

Klingler, J., Creasy, R., Gao, L., Nair, R. M., Calix, A. S., Jacob, H. S., Edwards, O. R. and Singh, K. B. (2005). Aphid Resistance in *Medicago truncatula* Involves Antixenosis and Phloem-Specific, Inducible Antibiosis, and Maps to a Single Locus Flanked by NBS-LRR Resistance Gene Analogs. *Plant Physiology* **137**, 1445-1455.

Klingler, J., Powell, G., Thompson, G. A. and Isaacs, R. (1998). Phloem specific aphid resistance in Cucumis melo line AR 5: effects on feeding behaviour and performance of *Aphis gossypii*. *Entomologia Experimentalis et Applicata* **86**, 79-88.

Kuznetsova, A., Brockhoff, P. B. and Christensen, R. H. B. (2017). lmerTest package: tests in linear mixed effects models. *Journal of Statistical Software* **82**, 1-26.

Lane, B. G. (2002). Oxalate, Germins, and Higher-Plant Pathogens. *IUBMB Life* **53**, 67-75.

Lenth, R. V. (2016). Least-Squares Means: The R Package lsmeans. *Journal of Statistical Software* **69**, 1-33.

Leybourne, D. J., Bos, J. I. B., Valentine, T. A. and Karley, A. J. (2018). The price of protection: a defensive endosymbiont impairs nymph growth in the bird cherry-oat aphid, *Rhopalosiphum padi*. *Insect Science* **https://doi.org/10.1111/1744-7917.12606**

Li, G., Xu, X., Carver, B. F., Guo, P. and Puterka, G. (2018). *Dn10*, a New Gene Conferring Resistance to Russian Wheat Aphid Biotype 2 in Iranian Wheat Landrace PI 682675. *Crop Science* **https://doi.org/10.2135/cropsci2017.10.0649**.

Livak, K. J. and Schmittgen, T. D. (2001). Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the 2−ΔΔCT Method. *Methods* **25**, 402-408.

Losvik, A., Beste, L., Glinwood, R., Ivarson, E., Stephens, J., Zhu, L.-H. and Jonsson, L. (2017). Overexpression and Down-Regulation of Barley Lipoxygenase *LOX2.2* Affects Jasmonate-Regulated Genes and Aphid Fecundity. *International Journal of Molecular Sciences* **https://doi:10.3390/ijms18122765**.

Louis, J., Basu, S., Varsani, S., Castano-Duque, L., Jiang, V., Williams, W. P., Felton, G. W. and Luthe, D. S. (2015). Ethylene Contributes to *maize insect resistance1*-Mediated Maize Defense against the Phloem Sap-Sucking Corn Leaf Aphid. *Plant Physiology* **169**, 313-324.

Maag, D., Erb, M., Kollner, T. G. and Gershenzon, J. (2015). Defensive weapons and defense signals in plants: some metabolites serve both roles. *Bioessays* **37**, 167-74.

MacGillivray, M. and Anderson, G. (1957). Three useful insect cages. *The Canadian Entomologist* **89**, 43-46.

Marimuthu, M. and Smith, C. M. (2012). Barley tolerance of Russian wheat aphid (Hemiptera: Aphididae) biotype 2 herbivory involves expression of defense response and developmental genes. *Plant Signaling & Behavior* **7**, 382-391.

Mayoral, A. M., Tjallingii, W. F. and Castañera, P. (1996). Probing behaviour of *Diuraphis noxia* on five cereal species with different hydroxamic acid levels. *Entomologia Experimentalis et Applicata* **78**, 341-348.

Medina-Ortega, K. J. and Walker, G. P. (2015). Faba bean forisomes can function in defence against generalist aphids. *Plant, Cell & Environment* **38**, 1167-1177.

Mehrabi, S., Åhman, I. and Jonsson, L. M. V. (2014). Transcript abundance of resistance- and susceptibility-related genes in a barley breeding pedigree with partial resistance to the bird cherry-oat aphid (Rhopalosiphum padi L.). *Euphytica* **198**, 211-222.

Mitchell, C., Brennan, R. M., Graham, J. and Karley, A. J. (2016). Plant Defense against Herbivorous Pests: Exploiting Resistance and Tolerance Traits for Sustainable Crop Protection. *Frontiers in Plant Science* **7**, https://doi.org/10.3389/fpls.2016.01132.

Moharramipour, S., Tsumuki, H., Sato, K. and Yoshida, H. (1997). Mapping resistance to cereal aphids in barley. *Theoretical and Applied Genetics* **94**, 592-596.

Montllor, C. B. and Tjallingii, W. F. (1989). Stylet penetration by two aphid species on susceptible and resistant lettuce. *Entomologia Experimentalis et Applicata* **52**, 103-111.

Moran, P. J. and Thompson, G. A. (2001). Molecular Responses to Aphid Feeding in Arabidopsis in Relation to Plant Defense Pathways. *Plant Physiology* **125**, 1074-1085.

Moreira, X., Abdala-Roberts, L., Gols, R. and Francisco, M. (2018). Plant domestication decreases both constitutive and induced chemical defences by direct selection against defensive traits. *Scientific Reports* **https://doi.org/10.1038/s41598-018-31041-0**.

Morkunas, I., Mai, V. C. and Gabryś, B. (2011). Phytohormonal signaling in plant responses to aphid feeding. *Acta Physiologiae Plantarum* **33**, 2057-2073.

Morris, J. G. (1991). Nutrition. In *Environmental and Metabolic Animal Physiology*, (ed. C. L. Prosser), pp. 231 - 276. New York, USA: John Wiley & Sons.

Murray, G. M. and Brennan, J. P. (2010). Estimating disease losses to the Australian barley industry. *Australasian Plant Pathology* **39**, 85-96.

Newman, C. and Newman, R. (2006). A brief history of barley foods. *Cereal foods world* **51**, 4-7.

Newton, A. C., Flavell, A. J., George, T. S., Leat, P., Mullholland, B., Ramsay, L., Revoredo-Giha, C., Russell, J., Steffenson, B. J., Swanston, J. S. et al. (2011). Crops that feed the world 4. Barley: a resilient crop? Strengths and weaknesses in the context of food security. *Food Security* **https://doi.org/10.1007/s12571-011-0126-3**.

Nieto-Lopez, R. M. and Blake, T. K. (1994). Russian Wheat Aphid Resistance in Barley: Inheritance and Linked Molecular Markers. *Crop Science* **34**, 655-659.

Oksanen, J., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'hara, R., Simpson, G. L., Solymos, P., Stevens, M. H. H. and Wagner, H. (2013). R Package vegan: Community Ecology Package. *CRAN repository* **available: https://CRAN.R-project.org/package=vegan**.

Park, S.-J., Huang, Y. and Ayoubi, P. (2006). Identification of expression profiles of sorghum genes in response to greenbug phloem-feeding using cDNA subtraction and microarray analysis. *Planta* **223**, 932-947.

Pegadaraju, V., Louis, J., Singh, V., Reese, J. C., Bautor, J., Feys, B. J., Cook, G., Parker, J. E. and Shah, J. (2007). Phloem-based resistance to green peach aphid is controlled by Arabidopsis

PHYTOALEXIN DEFICIENT4 without its signaling partner ENHANCED DISEASE SUSCEPTIBILITY1. *Plant J* **52**, 332-41.

Peng , H. C. and Walker, G. P. (2018). Sieve element occlusion provides resistance against *Aphis gossypii* in TGR-1551 melons. *Insect Science* **https://doi.org/10.1111/1744-7917.12610**.

Perry, K. L., Kolb, F. L., Sammons, B., Lawson, C., Cisar, G. and Ohm, H. (2000). Yield Effects of *Barley yellow dwarf virus* in Soft Red Winter Wheat. *Phytopathology* **90**, 1043-1048.

Pomeranz, M., Campbell, J., Siegal-Gaskins, D., Engelmeier, J., Wilson, T., Fernandez, V., Brkljacic, J. and Grotewold, E. (2013). High-resolution computational imaging of leaf hair patterning using polarized light microscopy. *The Plant Journal* **73**, 701-708.

Ponder, K. L., Pritchard, J., Harrington, R. and Bale, J. S. (2000). Difficulties in location and acceptance of phloem sap combined with reduced concentration of phloem amino acids explain lowered performance of the aphid *Rhopalosiphum padi* on nitrogen deficient barley (*Hordeum vulgare*) seedlings. *Entomologia Experimentalis et Applicata* **97**, 203-210.

Powell, G. (2005). Intracellular salivation is the aphid activity associated with inoculation of non-persistently transmitted viruses. *Journal of General Virology* **86**, 469-72.

Powell, G., Maniar, S. P., Pickett, J. A. and Hardie, J. (1999). Aphid responses to non-host epicuticular lipids. *Entomologia Experimentalis et Applicata* **91**, 115-123.

Reimann-Philipp, U., Schrader, G., Martinoia, E., Barkholt, V. and Apel, K. (1989). Intracellular thionins of barley. A second group of leaf thionins closely related to but distinct from cell wall-bound thionins. *Journal of Biological Chemistry* **264**, 8978-8984.

Roberts, J. J. and Foster, J. E. (1983). Effect of Leaf Pubescence in Wheat on the Bird Cherry Oat Aphid (Homoptera: Aphidae). *Journal of Economic Entomology* **76**, 1320-1322.

Sandström, J. and Pettersson, J. (1994). Amino acid composition of phloem sap and the relation to intraspecific variation in pea aphid (*Acyrthosiphon pisum*) performance. *Journal of Insect Physiology* **40**, 947-955.

Sandström, J., Telang, A. and Moran, N. A. (2000). Nutritional enhancement of host plants by aphids — a comparison of three aphid species on grasses. *Journal of Insect Physiology* **46**, 33-40.

Seah, S., Sivasithamparam, K., Karakousis, A. and Lagudah, E. S. (1998). Cloning and characterisation of a family of disease resistance gene analogs from wheat and barley. *Theoretical and Applied Genetics* **97**, 937-945.

Smith, C. M. and Boyko, E. V. (2007). The molecular bases of plant resistance and defense responses to aphid feeding: current status. *Entomologia Experimentalis et Applicata* **122**, 1-16.

Smith, P. and Sward, R. (1982). Crop loss assessment studies on the effects of *barley yellow dwarf virus* in wheat in Victoria. *Australian Journal of Agricultural Research* **33**, 179-185.

Stam, J. M., Kroes, A., Li, Y., Gols, R., Loon, J. J. A. v., Poelman, E. H. and Dicke, M. (2014). Plant Interactions with Multiple Insect Herbivores: From Community to Genes. *Annual Review of Plant Biology* **65**, 689-713.

Stec, B. (2006). Plant thionins – the structural perspective. *Cellular and Molecular Life Sciences* **63**, 1370-1385.

Stewart, S. A., Hodge, S., Ismail, N., Mansfield, J. W., Feys, B. J., Prospéri, J.-M., Huguet, T., Ben, C., Gentzbittel, L. and Powell, G. (2009). The *RAP1* Gene Confers Effective, Race-Specific Resistance to the Pea Aphid in *Medicago truncatula* Independent of the Hypersensitive Reaction. *Molecular Plant-Microbe Interactions* **22**, 1645-1655.

Tang, Y., Horikoshi, M. and Li, W. (2016). ggfortify: Unified Interface to Visualize Statistical Result of Popular R Packages. *The R Journal* **8**, 478 - 489.

Telang, A., Sandström, J., Dyreson, E. and Moran, N. A. (1999). Feeding damage by *Diuraphis noxia* results in a nutritionally enhanced phloem diet. *Entomologia Experimentalis et Applicata* **91**, 403-412.

Therneau, T. (2018). coxme: Mixed effects Cox models. *CRAN repository* **available: https://cran.r-project.org/web/packages/coxme/index.html**.

Therneau, T. M. and Grambsch, P. M. (2000). Modeling Survival Data: Extending the Cox Model. New York: Springer.

Tjallingii, W. (2001). Comparison of AC and DC systems for electronic monitoring of stylet penetration activities by homopterans. In *Principles and applications of electronic monitoring and other techniques in the study of homopteran feeding behavior*, eds. G. P. Walker and E. A. Backus), pp. 41. Beijing, China.

Tjallingii, W. (1991). Continuous recording of stylet penetration activities by aphids. In *Aphid-Plant Genotype Interactions*, eds. C. RK and E. RD), pp. 89 - 101. New York, USA: Elsevier.

Tjallingii, W. (1988). Electrical Recording of Stylet Penetration Activities. In *Aphids Their Biology, Natural Enemies and Control*, eds. M. AK and H. P), pp. 95 - 108. Amsterdam, The Netherlands: Elsevier.

Tjallingii, W. F. (1978). Electronic recording of penetration behaviour by aphids. *Entomologia Experimentalis et Applicata* **24**, 721-730.

Tsumuki, H., Kanehisa, K. and Kawada, K. (1989). Leaf surface wax as a possible resistance factor of barley to cereal aphids. *Applied Entomology and Zoology* **24**, 295-301.

Unal, G. and Jepson, P. C. (1991). The toxicity of aphicide residues to beneficial invertebrates in cereal crops. *Annals of Applied Biology* **118**, 493-502.

Vandesompele, J., De Preter, K., Pattyn, F., Poppe, B., Van Roy, N., De Paepe, A. and Speleman, F. (2002). Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biology and Evolution* **https://doi.org/10.1186/gb-2002-3-7-research0034**.

Vogel, K. J. and Moran, N. A. (2011). Sources of variation in dietary requirements in an obligate nutritional symbiosis. *Proceedings of the Royal Society B: Biological Sciences* **278**, 115-121.

Wu, Y., Zhang, D., Chu, Jee Y., Boyle, P., Wang, Y., Brindle, Ian D., De Luca, V. and Després, C. (2012). The *Arabidopsis* NPR1 Protein Is a Receptor for the Plant Defense Hormone Salicylic Acid. *Cell Reports* **1**, 639-647.

Xu, X., Bai, G., Carver, B. F., Zhan, K., Huang, Y. and Mornhinweg, D. (2015). Evaluation and Reselection of Wheat Resistance to Russian Wheat Aphid Biotype 2. *Crop Science* **55**, 695-701.

Zadoks, J. C., Chang, T. T. and Konzak, C. F. (1974). A decimal code for the growth stages of cereals. *Weed Research* **14**, 415-421.

Zhang, H., Mittal, N., Leamy, L. J., Barazani, O. and Song, B.-H. (2017). Back into the wild—Apply untapped genetic diversity of wild relatives for crop improvement. *Evolutionary Applications* **10**, 5-24.

Zhu-Salzman, K., Salzman, R. A., Ahn, J.-E. and Koiwa, H. (2004). Transcriptional Regulation of Sorghum Defense Determinants against a Phloem-Feeding Aphid. *Plant Physiology* **134**, 420-431.

Tables

Table 1: Statistical results of aphid performance experiments (significant values are shown in bold).

Response Variable	Explanatory Variable	Model Basis	Error Distribution or Statistical Method	Statistical Test	Test Statistic	P value
Nymph Survival	Plant	Cox Proportional Hazards Regression	N/A	Type II Wald χ^2 Analysis of Deviance	$X^2_1 = 10.65$	**0.001**
	Aphid				$X^2_2 = 13.94$	**<0.001**
	Plant x Aphid				$X^2_2 = 5.85$	0.054
Nymph mass gain (mg)	Plant	Linear Mixed Effects Model	Restricted Maximum Likelihood	Type III Analysis of Variance with Satterthwaite approximation for degrees of freedom	$F_{1,56} = 9.14$	**0.004**
	Aphid				$F_{2,56} = 49.83$	**<0.001**
	Plant x Aphid				$F_{2,46} = 3.12$	0.054
Rate of population increase (r_m)	Plant				$F_{1,54} = 27.43$	**<0.001**
	Aphid				$F_{2,54} = 99.10$	**<0.001**
	Plant x Aphid				$F_{2,52} = 1.49$	0.234

Table 2: Statistical results of the significant EPG parameters. The mean value for each plant host and the standard error of the mean are also displayed; in the test statistic column χ² values are italicised and F values are underlined.

| EPG Parameter | Hypothesised location of resistance factor | Transformation | Mean Value | | Standard Error of the Mean | | Statistical Test | Test Statistic | Degrees of Freedom (Residuals) | P Value |
			cv. Concerto	HsP5	cv. Concerto	HsP5				
All parameters (Global analysis)	All tissue	-	-	-	-	-	Permutated Multiple Analysis of Variance	2.61	1(33)	**0.022**
Time to the first aphid probe of plant tissue	Epidermis	Log[2]	1211 s	421 s	354 s	85 s	Analysis of Variance (ANOVA)	7.99	1(33)	**0.008**
Duration of the first aphid probe	Mesophyll	Log[10]	7717 s	2147 s	1737 s	1161 s	ANOVA	6.94	1(33)	**0.013**
Number of potential drops (intracellular punctures) in first probe	Mesophyll	Sqrt	24	10	5.36	2.82	ANOVA on glm	3.95	1(33)	**0.047**
Time to the first E1 phase (sieve element puncture and salivation into phloem)	Mesophyll	Log[2]	2856 s	7452 s	660 s	1683 s	ANOVA	6.33	1(33)	**0.017**
Time to the first phloem phase containing E1 and/or E2	Mesophyll	-	3761 s	9864 s	675 s	1884 s	Kruskall-Wallis Test (KW)	5.50	1(33)	**0.019**
Time to the first phloem ingestion (E2) phase	Mesophyll	-	4395 s	9946 s	846 s	1880 s	ANOVA	5.32	1(33)	**0.028**
Time to first sustained phloem ingestion (>10 minutes)	Mesophyll	-	8210 s	13660 s	1345 s	1954 s	KW	4.29	1(33)	**0.038**

Number of probes before first E1	Mesophyll	-	0.66	2.38	0.34	0.76	KW	*9.36*	1(33)	**0.002**
Number of probes before first E2	Mesophyll	-	1	2.5	0.38	0.79	KW	*4.14*	1(33)	**0.042**
Total time ingesting xylem	Xylem	-	2197 s	6859 s	721 s	1408 s	KW	*5.28*	1(33)	**0.022**
Total number of xylem phases	Xylem	-	0.72	1.56	0.13	0.30	KW	*4.77*	1(33)	**0.029**
Total time ingesting phloem (E2)	Phloem	-	7965 s	3299 s	1306 s	1188 s	KW	*8.00*	1(33)	**0.004**
Average time of phloem ingestion (E2)	Phloem	-	5317 s	1717 s	1533 s	702 s	KW	*6.52*	1(33)	**0.011**
Total time of phloem phases containing E1 and/or E2	Phloem	-	8824 s	3620 s	1348 s	1241 s	KW	*9.22*	1(33)	**0.002**
Maximum time of a single phloem phase with E1 and/or E2	Phloem	-	7913 s	3225 s	1442 s	1172 s	KW	*9.01*	1(33)	**0.002**
Maximum length of a single E2 period	Phloem	-	7215 s	2884 s	1371 s	1134 s	KW	*8.40*	1(33)	**0.004**
Average time of phloem phase containing E1 and/or E2	Phloem	-	5911 s	1913 s	1653 s	722 s	KW	*6.88*	1(33)	**0.008**
Time from first E1 phase to sustained E2 phase	Phloem	-	8210 s	13660 s	1345 s	1954 s	KW	*4.29*	1(33)	**0.038**
Total number of sustained E2 phases	Phloem	-	1.39	0.75	0.16	0.17	KW	*6.06*	1(33)	**0.014**
Mean duration of sustained E2	Phloem	-	6500 s	2779 s	1424 s	1141 s	KW	*7.12*	1(33)	**0.008**
Median duration of sustained E2	Phloem	-	6425 s	2779 s	1440 s	1141 s	KW	*6.40*	1(33)	**0.011**

Total time of sustained E2	Phloem	-	7694 s	3060 s	1342 s	1162 s	KW	*7.49*	1(33)	**0.006**
Ingestion:Pathway Ratio (%)	Phloem	-	3.01	0.67	1.40	0.33	KW	*7.43*	1(33)	**0.006**

Table 3: Statistical results of plant defence gene expression assessed by qPCR showing χ^2 and p values (significant values are shown in bold)

Gene	Explanatory variable			
	Plant	Aphid infestation	Plant x infestation	Experimental Block
HvTHIO1	$X^2 = 11.06$ p = **<0.001**	$X^2 = 0.57$ p = 0.450	$X^2 = 11.67$ p = **0.009**	$X^2 = 1.10$ p = 0.894
HvTHIO2	$X^2 = 13.62$ p = **<0.001**	$X^2 = 0.57$ p = 0.449	$X^2 = 14.31$ p = **0.002**	$X^2 = 0.47$ p = 0.976
HvβTHIO	$X^2 = 14.29$ p = **<0.001**	$X^2 = 1.12$ p = 0.290	$X^2 = 16.10$ p = **0.001**	$X^2 = 0.15$ p = 0.997
HvA1	$X^2 = 5.77$ p = **0.016**	$X^2 = 1.46$ p = 0.226	$X^2 = 4.57$ p = 0.210	$X^2 = 4.58$ p = 0.206
HvNPR1	$X^2 = 1.12$ p = 0.290	$X^2 = 1.85$ p = 0.174	$X^2 = 8.46$ p = **0.037**	$X^2 = 3.64$ p = 0.457
HvERF1	$X^2 = 7.00$ p = **0.008**	$X^2 = 4.81$ p = **0.028**	$X^2 = 12.26$ p = **0.006**	$X^2 = 2.24$ p = 0.691
HvLOXA	$X^2 = 4.81$ p = **0.028**	$X^2 = 4.48$ p = **0.034**	$X^2 = 8.05$ p = **0.045**	$X^2 = 3.11$ p = 0.539
HvLOX2	$X^2 = 5.49$ p = **0.019**	$X^2 = 2.76$ p = 0.096	$X^2 = 8.46$ p = **0.037**	$X^2 = 3.97$ p = 0.410
HvJAZ	$X^2 = 2.85$ p = 0.131	$X^2 = 0.57$ p = 0.450	$X^2 = 3.55$ p = 0.315	$X^2 = 9.42$ p = 0.055

Table 4: Dunn's test *post-hoc* analysis of qPCR results significant for plant x time-point interaction, showing *t* and p values for each pairwise comparison; C = Concerto, H = Hsp5, 0 = 0 h infestation, 24 = 24 h aphid infestation (significant values are shown in bold).

Gene	Pairwise comparison											
	C0:C24		C0:H0		C0:H24		C24:H0		C24:H24		H0:H24	
HvTHIO1	t = 0.69	P = 0.244	t = -2.19	P = **0.014**	t = -1.82	P = **0.034**	t = -2.89	P = **0.001**	t = -2.51	P = **0.006**	t = 0.37	P = 0.354
HvTHIO2	t = 1.28	P = **0.099**	t = -1.76	P = **0.038**	t = -1.97	P = **0.024**	t = -3.05	P = **0.001**	t = -*3.26*	P = **<0.001**	t = -0.21	P = 0.415
HvβটHIO	t = 1.33	P = **0.09**	t = -2.08	P = **0.018**	t = -1.92	P = **0.022**	t = -3.42	P = **<0.001**	t =-3.26	P = **<0.001**	t = 0.16	P = 0.436
HvNPR1	t = 2.62	P = **0.004**	t = 0.91	P = 0.182	t = 0.21	P = 0.415	t = -1.71	P = **0.043**	t = -2.41	P = **0.008**	t = -0.69	P = 0.244
HvERF1	t = 2.03	P = **0.021**	t = -1.38	P = 0.082	t = -0.32	P = 0.374	t = -3.42	P = **<0.001**	t = -2.35	P = **<0.001**	t = 1.06	P = 0.143
HvLOXA	t = 1.12	P = 0.132	t = -1.66	P = **0.048**	t = 0.21	P = 0.415	t = -2.77	P = **0.002**	t = -0.91	P = 0.182	t = 1.87	P = **0.030**
HvLOX2	t = 1.49	P = 0.067	t = -1.33	P = 0.091	t = -0.48	P = 0.315	t = -2.83	P = **0.002**	t = -1.97	P = **0.024**	t = 0.85	P = 0.196

Table 5: Analysis of deviance table for the scores on each principal component (% variation explained is indicated in brackets) derived from Principal Component Analysis of phloem amino acid composition (significant values are shown in bold)

Explanatory Variable	PC1 (34.38%)		PC2 (17.80%)		PC3 (10.89%)	
	X^2 (df)	p-value	X^2 (df)	p-value	X^2 (df)	p-value
Plant	40.11 (1)	**<0.001**	5.40 (1)	**0.020**	11.98 (1)	**<0.001**
Aphid infestation	0.29 (1)	0.585	0.24 (1)	0.626	2.26 (1)	0.131
Time-point	5.42 (1)	**0.019**	3.05 (1)	0.080	0.08 (1)	0.765
Plant x time-point	0.05 (1)	0.816	2.17 (1)	0.140	1.97 (1)	0.159
RNA Harvest	0.17 (1)	0.671	1.92 (1)	0.166	0.67 (1)	0.413

Table 6: Analysis of deviance table for essential amino acids as a percentage of total amino acids in phloem sap of Hsp5 and Concerto leaves (significant values are shown in bold).

Explanatory Variable	Essential Amino acid (%)	
	X^2 (df)	p-value
Plant	23.35 (1)	**<0.001**
Aphid infestation	0.13 (1)	0.716
Time-point	8.82 (1)	**0.002**
Plant x time-point	0.39 (1)	0.530
RNA Harvest	2.01 (1)	0.155

Figure Legends

Fig 1. Aphid performance on a susceptible barley cultivar (Concerto) and partially-resistant wild relative (HsP5). A) Survival of aphid nymphs over seven days. Number of model observations = 216. C = Concerto; H5 = Hsp5; Rp = *R. padi;* Sa = *S. avenae;* Uh = *U. humboldti.* B) Nymph mass gain and C) intrinsic rate of population increase (rm) of the three aphid species while feeding from the two plant types. Values represent means ± SE. Number of model observations = 72. Letters indicate which groups are similar to each other based on least squares means analysis with Tukey correction.

Fig. 2: Representative examples of feeding patterns derived from EPG waveforms of *R. padi* feeding from susceptible Concerto (A) and partially-resistant Hsp5 (B). Diagrams show the frequency and length of each feeding parameter over a 6h (21600s) period. Feeding patterns are split into the five main waveforms observed: non-probing of plant tissue (np), probing into epidermal and mesophyll tissue (the pathway phase (C), saliva secretion into phloem (E1), phloem ingestion (E2), and xylem ingestion (G).

Fig 3: EPG parameters indicative of epidermal, mesophyll and mesophyll/phloem resistance factors. A) Time (s) from start of EPG recording to first stylet puncture of leaf tissue. B) Duration (s) of first stylet probe into plant tissue. C) Time (s) since start of EPG recording until stylet puncture of a sieve tube element. D) Time (s) from start of EPG recording until sustained phloem ingestion. A, B and C values represent means ± SE. Number of model observations = 34.

Fig 4: EPG parameters indicative of vascular (phloem and xylem) resistance factors. A) Time (s) from first stylet penetration of a sieve tube element to first sustained phloem ingestion. B) Total time (s) of sustained phloem feeding. C) Ratio of time spent in phloem phase relative to the pathway phase; values represent means ± SE. D) Total time (s) ingesting xylem. Number of model observations = 34.

Fig. 5: Trichome densities in the two plant types. A) Median non-glandular trichome densities (No. cm^{-2}) and confidence intervals for Concerto and Hsp5. B) Polarised light micrograph showing the contrasting leaf hair densities between HsP5 (top) and Concerto (bottom). Number of model observations = 30.

Fig 6. Expression patterns of thionin and phytohormone signalling genes 0 h and 24 h after aphid infestation. All gene expression values are relative to the mean expression of two reference genes, *HvUBC* and *HvCYP,* and normalised to uninfested control plants. Fold-change, 2-ΔΔCt, is relative to the expression of cv. Concerto at 0 h. Boxplots show median and confidence intervals; total number of model observations for each gene = 20.

Fig. 7: Amino acid composition of leaf phloem exudates of Hsp5 and Concerto. A and B) phloem amino acid composition in uninfested leaves of Concerto and Hsp5, respectively, at time zero. C) Biplot of scores on the first two principal components explaining >50% of the detected variation in amino acid composition. D) % proportion of essential and non-essential amino acids in phloem exudates of uninfested Concerto and Hsp5leaves at time zero.

Figures

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7